

- transplantation (HCT). The 49th American Society of Hematology Annual Meeting, Atlanta, December 8-11, 2007. (abstr. # 5330).
11. Yokoyama H, Fukuda T, Kobayashi Y, Kim SW, Maruyama D, Mori S, Watanabe T, Tanosaki R, Tajima K, Tobinai K, Takaue Y: Hematopoietic cell transplantation (HCT) for chemoradiotherapy-related myelodysplastic syndrome and acute leukemia: A single-center analysis of 47 patients. The 49th American Society of Hematology Annual Meeting, Atlanta, December 8-11, 2007.
  12. Kato S, Atsuta Y, Morishima Y, Nagamura T, Takaue Y, Sakamaki H, Okamoto S, Takahashi S, Taniguchi S, Kato K, Isoyama K, Takashi M, Kai S, Azuma H, Tsuchida M, Kawa K, Kodera Y for the Japan Marrow Donor Program (JMDP) and the Japan Cord Blood Bank Network (JCBBN): Comparison of 6586 unrelated bone marrow transplants (UBMT) and 2713 unrelated cord blood transplants (UCBT) in Japan. The 49th American Society of Hematology Annual Meeting, Atlanta, December 8-11, 2007.
  13. Kim SW, Lee JW, Kim HP, Fukuda T, Kanda Y, Kim HJ, Min WS, Mori S, Tanosaki R, Lee JH, Lee KH, Yamashita T, Sakamaki H, Sohn SK, Chae YS, Ohshima K, Ichinohe T, Kanda J, Kasai M, Kobayashi N, Ando K, Onizuka M, Taniguchi S, Seo S, Yang DH, Kim YK, Shin HJ, Takaue Y, Kim CC: Similar background of graft-versus-host disease (GVHD) following allogeneic blood and marrow transplantation (Tx) in Korean and Japanese populations allows a uniform foundation to perform large-scale clinical studies in the region. The 49th American Society of Hematology Annual Meeting, Atlanta, December 8-11, 2007.
  14. Fukuda T, Takahashi T, Iisaka Y, Ohara-Waki F, Kim SW, Mori S, Tanosaki R, Makimoto A, Tobinai K, Takaue Y: Primary treatment for acute graft-versus-host disease (GVHD): A single-center analysis of 458 Japanese patients. The 49th American Society of Hematology Annual Meeting, Atlanta, December 8-11, 2007.
  15. Yokoyama H, Fukuda T, Kobayashi Y, Kim SW, Maruyama D, Mori S, Watanabe T, Tanosaki R, Tajima K, Tobinai K, Takaue Y: Hematopoietic cell transplantation for chemoradiotherapy-related myelodysplastic syndrome and acute leukemia: a single-center analysis of 47 patients. American Society for Blood and Marrow Transplantation Annual Meeting, San Diego, February 13-17, 2008.
  16. 岡本康裕, 河野嘉文 他 6 名. 地方施設における再生不良性貧血に対する同種骨髄移植の後方視的検討. 第 69 回日本血液学会・第 49 回日本臨床血液学会 (横浜 2007.10.11-13)
  17. 岡本康裕, 河野嘉文 他 7 名. 造血幹細胞移植の医師チーム制による管理. 第 30 回日本造血細胞移植学会 (大阪 2008.3.1)
  18. 川森有里子, 松井利充 他 6 名. 非血縁者間同種骨髄移植後、閉塞性障害を伴う放射線肺臓炎を来たしアジスロマイシンが奏功した一例. 近畿血液学地方会 (大阪 2007.6.)
  19. 薬師神公和, 松井利充 他 11 名. 類洞閉塞症候群(SOS)発症予測因子としての血小板凝集能検査の有用性. 第 69 回日本血液学会・第 49 回日本臨床血液学会 (横浜 2007.10.11-13)
  20. 岡村篤夫, 松井利充 他 6 名. 薬物血中濃度解析(PK)に基づく移植後至適ミコフェノール酸モチフェル(MMF)投与方法の確立に関する検討. 第 69 回日本血液学会・第 49 回日本臨床血液学会 (横浜 2007.10.11-13)
  21. 岡村篤夫, 松井利充 他 2 名. 急性 GVHD 予防薬ミコフェノール酸モチフェル(MMF)の移植後 Day30 以降における漸減投与の必要性. 第 30 回日本造血細胞移植学会 (大阪 2008.2.28-3.1)
  22. 高橋義行, 松井利充 他 5 名. 後天性再生不良性貧血 31 例に対する非血縁者間臍帯血移植の治療成績. 第 30 回日本造血細胞移植学会 (大阪 2008.2.28-3.1)

23. 井上潤一郎、松井利充 他 5 名. 同種造血幹細胞移植患者の身体活動量が身体機能に与える影響. 第 30 回日本造血細胞移植学会 (大阪 2008.2.28-3.1)
24. 増田委加、松井利充 他 4 名. 造血幹細胞移植療法 (HSCT) における皮下埋め込み型中心静脈ポートの安全性および有用性. 第 30 回日本造血細胞移植学会 (大阪 2008.2.28-3.1)
25. 安永和世、松井利充 他 5 名. 造血幹細胞移植時生野菜を含む普通食提供の安全性および有用性に関する検討. 第 30 回日本造血細胞移植学会 (大阪 2008.2.28-3.1)
26. 宇都宮興 他 10 名. ヒト化抗 CCR4 抗体 KW-0761 の再発 ATLL および PTCL 患者での第 1 相試験-予備解析結果. 第 69 回日本血液学会・第 49 回日本臨床血液学会 (横浜 2007.10.11-13)
27. 中野伸亮、宇都宮興 他 4 名. 成人 T 細胞白血病リンパ腫(ATLL)に対する同種造血幹細胞移植後の再発例の検討. 第 69 回日本血液学会・第 49 回日本臨床血液学会 (横浜 2007.10.11-13)
28. 高塚祥芝、宇都宮興 他 4 名. ATLL における同種造血細胞移植後の長期生存例の検討. 第 30 回日本造血細胞移植学会 (大阪 2008.2.28-3.1)
29. 糸山貴浩, 中野伸亮, 窪田歩, 竹内昇吾, 高塚祥芝, 宇都宮興: 同種骨髓移植後, 第 XIII 因子欠乏を背景とする消化管出血を繰り返した ATLL の 1 例. 第 30 回日本造血細胞移植学会総会 2008 年 2 月 29 日-3 月 1 日, 大阪
30. 窪田歩, 高塚祥芝, 中野伸亮, 糸山貴浩, 竹内昇吾, 宇都宮興: RIST 後モノクローナル T リンパ球増殖症を起こした ATLL の 1 症例. 第 30 回日本造血細胞移植学会総会 2008 年 2 月 29 日-3 月 1 日, 大阪
31. Utsunomiya A, Tanosaki R, Uike N, Masuda M, Saburi Y, Tomonaga M, Etoh T, Kikuchi H, Harada M, Kawano F, Choi I, Matsuoka M, Kannagi M, Sonoda S, Okamura J: Kinetics of HTLV-1 proviral loads after allogeneic hematopoietic stem cell transplantation using reduced intensity conditioning adult T-cell leukemia/lymphoma. 13th International Conference on Human Retrovirology HTLV AND RELATED VIRUSES in Japan, May 22-25, 2007
32. Yonekura K, Utsunomiya A, Nakano N, Takeuchi S, Itoyama T, Takatsuka Y, Tashiro Y, Kanzaki T, Kanekura T: Graft-versus-adult T-cell leukemia/lymphoma (Gv-ATL) effect in ATL patients after allogeneic stem cell transplantation. 13th International Conference on Human Retrovirology HTLV AND RELATED VIRUSES in Japan, May 22-25, 2007
33. Tsukasaki K, Utsunomiya A, Fukuda H, Masuda M, Yamada Y, Tobinai K, Tomonaga M, Shimoyama M, Hotta T: Clinical trials for the treatment of adult T-cell leukemia/lymphoma (ATLL) by the Japan Clinical Oncology Group (JCOG). 13th International Conference on Human Retrovirology HTLV AND RELATED VIRUSES in Japan, May 22-25, 2007
34. Uike N, Utsunomiya A, Tanosaki R, Masuda M, Saburi Y, Tomonaga M, Etoh T, Kikuchi H, Harada M, Hidaka M, Choi I, Yufu Y, Matsuoka M, Kannagi M, Okamura J: Comparable outcome of reduced-intensity conditioning with or without antithymocyte globulin in allogeneic stem cell transplantation for adult T-cell leukemia: The results of two prospective studies. 13th International Conference on Human Retrovirology HTLV AND RELATED VIRUSES in Japan, May 22-25, 2007
35. Harashima N, Shimizu Y, Takamori A, Kurihara K, Utsunomiya A, Tanosaki R, Masuda M, Okamura J, Kannagi M: Positive conversion of Tax-specific CTL response in ATL patients after hematopoietic stem cell transplantation. 13th International Conference on Human Retrovirology HTLV AND RELATED VIRUSES in Japan, May 22-25, 2007

## **Unrelated-Donor Bone Marrow Transplantation with a Conditioning Regimen Including Fludarabine, Busulfan, and 4 Gy Total Body Irradiation**

Yasushi Onishi, Shin-ichiro Mori, Shigeru Kusumoto, Kyoko Sugimoto, Daigo Akahane, Yuriko Morita-Hoshi, Sung-Won Kim, Takahiro Fukuda, Yuji Heike, Ryuji Tanosaki, Kensei Tobinai, Yoichi Takaue

*Division of Hematology and Hematopoietic Stem Cell Transplantation, National Cancer Center Hospital, Tokyo, Japan*

Received October 10, 2006; received in revised form December 26, 2006; accepted January 9, 2007

### **Abstract**

We investigated the feasibility of reduced-intensity conditioning with 4 Gy total body irradiation, fludarabine (30 mg/m<sup>2</sup> for 6 days), and busulfan (4 mg/kg for 2 days) for bone marrow transplantation from a serologically HLA-matched unrelated donor. Seventeen adult patients (median age, 55 years; range, 27-67 years) with various hematologic malignancies (6 in remission, 11 not in remission) were treated. Successful engraftment was achieved in all patients at a median of day 18 (range, day 14-35) after transplantation, although subsequent secondary graft failure was observed in 2 patients. The cumulative incidence of acute graft-versus-host disease (GVHD) of grades II to IV at day 100 was 48%. With a median follow-up of 286 days (range, 56-687 days), the rates of 1-year overall survival, 100-day nonrelapse mortality, and 1-year nonrelapse mortality were 41%, 14%, and 46%, respectively. Eleven patients died, and the causes of death were relapse (n = 4), pulmonary complications (n = 4), acute GVHD (n = 2), and sepsis (n = 1). The remaining 6 patients (at transplantation, 2 were in remission, and 4 were not in remission) are currently still in remission. These results suggest that this regimen reduces the risk of graft failure, but further studies are needed to ameliorate transplantation-related toxicities, primarily GVHD and/or pulmonary complications.

*Int J Hematol.* 2007;85:256-263. doi: 10.1532/IJH97.06199

© 2007 The Japanese Society of Hematology

**Key words:** Unrelated donor bone marrow transplantation; Fludarabine; Busulfan, TBI

### **1. Introduction**

Although allogeneic hematopoietic stem cell transplantation (HSCT) is a possible curative approach for patients with various hematologic malignancies, only 30% to 40% of patients in Japan have an appropriate family donor available [1]. Hence, the application of unrelated-donor transplantation using bone marrow or cord blood cells has been expanding. Another area of current interest is the application of reduced-intensity conditioning regimens, mostly incorporating fludarabine as a primary agent, because conventional allogeneic HSCT using a conditioning regimen

with high doses of systemic chemotherapy/radiation is associated with significant toxicities. In contrast, HSCT with a reduced-intensity conditioning regimen allows older patients and those who have contraindicating comorbidities to undergo HSCT [2-7].

Nevertheless, special consideration should be paid to developing reduced-intensity conditioning protocols for the unrelated-donor HSCT setting, because the incidences of both graft rejection and graft-versus-host disease (GVHD) are greater than in related-donor transplantation. In addition, the intensity of the reduced-intensity conditioning regimen influences transplantation-related toxicities and the relapse rate, and the stem cell source (ie, peripheral blood stem cells or bone marrow cells) influences engraftment [8]. Accordingly, several reduced-intensity conditioning protocols have been tested to address a variety of problems [8-17]. In this study, we investigated the feasibility of bone marrow transplantation (BMT) from a serologically HLA-matched unrelated donor with a regimen containing

Correspondence and reprint requests: Yoichi Takaue, National Cancer Center Hospital, 5-1-1, Tsukiji, Chuo-ku, Tokyo 104-0045, Japan; 81-3-3542-2511; fax: 81-3-3542-3815 (e-mail: ytakaue@ncc.go.jp).

4 Gy of total body irradiation (TBI), fludarabine (Flu), and busulfan (BU).

## 2. Patients and Methods

### 2.1. Patients and Donors

The data for adult patients with hematologic malignancies who underwent unrelated-donor BMT through the Japan Marrow Donor Program between June 2002 and December 2003 at the National Cancer Center Hospital were analyzed retrospectively. This protocol was approved by the Ethics Committee, and written informed consent was obtained from each patient. The patients who were enrolled in this study were ineligible for conventional allogeneic HSCT because of age (older than 50 years) and/or concomitant diseases or preceding intensive therapies, such as autologous HSCT or multiple chemotherapies. Donor-recipient pairs were selected on the basis of serologic matching for HLA-A and HLA-B and molecular matching for HLA-DRB1. HLA allele typing was performed by intermediate-resolution polymerase chain reaction (PCR) analysis. The stem cell source, which was determined by the Japan Marrow Donor Program donor center, was bone marrow in all cases.

### 2.2. Treatment Plan and Evaluations

The conditioning regimen consisted of 30 mg/m<sup>2</sup> Flu intravenously daily for 6 days (day -8 to day -3), 4 mg/kg BU orally daily for 2 days (days -6 and -5, without BU dose adjustment), and 4 Gy TBI without lung shielding (day -9 or day -1, single dose or 2 divided doses). Non-T-cell-depleted bone marrow was infused on day 0. The time of neutrophil engraftment was defined as the first of 3 consecutive days with an absolute neutrophil count  $\geq 0.5 \times 10^9/L$ , and the time of platelet engraftment was defined as the first of 7 consecutive days with a platelet count  $\geq 20 \times 10^9/L$  without transfusion support. Granulocyte colony-stimulating factor (G-CSF) was administered at 300  $\mu\text{g}/\text{m}^2$  from day 6 and continued until neutrophil engraftment. The degree of donor chimerism among peripheral blood mononucleated cells was evaluated by PCR analysis of short tandem repeat polymorphisms with fluorescently labeled primers. Secondary graft failure was defined as cytopenia with an absolute neutrophil count  $< 0.1 \times 10^9/L$  or decreasing chimerism not associated with relapsing disease in patients who had recovered in the early posttransplantation period.

GVHD prophylaxis consisted of cyclosporin A (CsA) from day -1 (daily administration of 3 mg/kg by continuous intravenous infusion or 6 mg/kg orally in 2 divided doses) and methotrexate (10 mg/m<sup>2</sup> intravenously on day 1 and 7 mg/m<sup>2</sup> on days 3, 6, and 11). The CsA dosage was adjusted according to the patient's renal function and to maintain therapeutic levels (250-350 ng/mL) with continuous infusion or trough levels (150-250 ng/mL) with oral administration. In patients without GVHD, CsA was tapered from day 100 over a 3- to 6-month period. Standard criteria were used to grade acute and chronic GVHD [18,19]. Chronic GVHD was evaluated in patients who survived at least 100 days and was classified as limited or extensive. Patients who developed acute

GVHD  $\geq$  grade II were treated with methylprednisolone at 1 to 2 mg/kg per day.

### 2.3. Supportive Care

Antimicrobial prophylaxis consisted of ciprofloxacin, fluconazole, acyclovir, and trimethoprim/sulfamethoxazole according to our institutional protocol. All patients were nursed in a room equipped with high-efficiency air filtration of particulates. Monitoring for cytomegalovirus (CMV) antigenemia was performed once a week after neutrophil engraftment by means of the horseradish peroxidase-C7 method. Patients positive for CMV antigenemia were started preemptively on ganciclovir therapy.

### 2.4. Statistical Analysis

Overall survival was calculated from the time of transplantation until death from any cause. Progression-free survival was measured from transplantation until disease progression or death from any cause. Nonrelapse death was defined as death due to any cause other than relapse. Survival curves for overall survival and progression-free survival were estimated by the Kaplan-Meier method.

## 3. Results

### 3.1. Patients

The median age of the 17 patients was 55 years (range, 27-67 years; Table 1). The diagnoses were acute myeloid leukemia (AML) (n = 7), myelodysplastic syndrome (MDS) (n = 4), chronic myelogenous leukemia (n = 1), non-Hodgkin's lymphoma (n = 4), and multiple myeloma (n = 1). Six patients were in remission at transplantation, and the remaining 11 were not in remission. Three patients with MDS or AML following MDS underwent unrelated-donor BMT as a primary treatment. Seven donor-recipient pairs were fully matched for HLA-A, HLA-B, and HLA-DRB1 at the allele level, 4 donor-recipient pairs had an allele-level mismatch at the HLA-A locus, and 5 pairs had an allele-level mismatch at the HLA-DRB1 locus. One patient was mismatched with the donor at 3 HLA alleles.

### 3.2. Engraftment and Chimerism

The median number of infused nucleated cells was  $2.7 \times 10^8/\text{kg}$  (range,  $0.65\text{-}5.5 \times 10^8/\text{kg}$ ). All patients achieved neutrophil recovery, but 5 patients did not become independent of platelet transfusion during their follow-up period (Table 2). The median times until neutrophil and platelet recoveries were 18 days (range, 14-35 days) and 26 days (range, 15-112 days), respectively (Figure 1). Late graft failure was observed in 2 patients, one of whom had secondary graft failure due to myelosuppression caused by ganciclovir treatment for CMV colitis. In this patient, donor chimerism was not assessed after day 30 when complete donor chimerism was confirmed. In the other case, donor chimerism decreased from 89% on day 30 to 33% on day 60, despite the tapering of CsA from day 30. Chimerism was

**Table 1.**  
Patient and Disease Characteristics\*

Patient No.	Age, y/Sex	Disease	Status	Time from Dx to HSCT, mo	HLA Allelic Mismatch	GVH Vector	HVG Vector	Contraindications to Conventional HSCT	Pretransplantation Comorbidities
1	55/F	AML	CR3	117				Age	No
2	52/F	AML	Primary Ref	13	DRB1	1	1	Age + comorbidity	Pneumonia
3	57/F	AML	Rel2	28				Age	Atrial fibrillation
4	55/M	MDS	Primary Ref	3				Age	Atrial fibrillation
5	57/M	MDS	CR1	8				Age	No
6	59/M	CML	CP2	8				Age	No
7	55/M	PTCL	PR	16	DRB1	1	1	Age	Gastric ulcer
8	58/M	AML	Untreated	10	DRB1	1	1	Age	Bronchial asthma, FEV <sub>1</sub> 75%
9	59/M	AML	Untreated	33	DRB1	1	1	Age	Bilirubin 1.5 mg/dL
10	52/M	AML	CR1	11	A	1	1	Age	FEV <sub>1</sub> 67%
11	57/M	MDS	CR1	13				Age	Prior gastric cancer
12	61/M	AML	CR2	58	A, both DRB1	3	3	Age	No
13	67/F	FL	Primary Ref	58	A	1	1	Age + comorbidity	Dyspnea requiring oxygen
14	27/M	DLBCL	Rel3	38	A	1	0	Prior autologous HSCT	No
15	48/F	MM	Primary Ref	80				Comorbidity	Ventricular septal defect
16	52/F	MDS	Untreated	130	A	1	1	Age	No
17	49/M	FL	Rel1	28	DRB1	1	1	Prior multiple chemotherapies	No

\*Dx indicates diagnosis; HSCT, hematopoietic stem cell transplantation; GVH, graft-versus-host; HVG, host-versus-graft; AML, acute myeloid leukemia; CR3, third complete remission; Ref, refractory; Rel2, second relapse; MDS, myelodysplastic syndrome; CML, chronic myelogenous leukemia; CP2, second chronic phase; PTCL, peripheral T-cell lymphoma; PR, partial remission; FEV<sub>1</sub>, forced expiratory volume in 1 second; FL, follicular lymphoma; DLBCL, diffuse large B-cell lymphoma; MM, multiple myeloma.

evaluated by analysis of short tandem repeats in 14 patients, and complete donor chimerism was confirmed in 12 of these patients. One patient who relapsed on day 32 had exhibited 54% donor chimerism on day 30. In the remaining 3 patients who relapsed after transplantation, complete donor chimerism had been achieved by day 30. In the patient who relapsed on day 78, donor chimerism decreased from 100% on day 30 to 64% on day 60. Mixed chimerism was not confirmed in the other 2 patients before disease progression or relapse. The patients without graft failure or relapse did not have mixed chimerism during their follow-up periods.

### 3.3. Regimen-Related Toxicities and Infections

Regimen-related toxicity was graded according to the National Cancer Institute Common Toxicity Criteria, version 2.0, and maximum toxicities are shown in Table 3. Fifteen of the 17 patients had grade III oral/pharyngeal mucositis that required morphine as an analgesic. Reversible elevation (grades III-IV) in transaminase and bilirubin levels occurred in 35% and 12% of the cases, respectively. No veno-occlusive disease was observed. Four patients developed transient grade III hyponatremia within 28 days after transplantation. Four patients developed transient pulmonary infiltration or congestive heart failure due to hypercytokinemia at engraftment, and 2 of these patients developed grade II acute GVHD after engraftment. No histologic findings of acute GVHD were seen in the other 2 patients. One patient developed reversible paroxysmal

supraventricular tachycardia. One patient developed bloody diarrhea and abdominal pain even after improvement of acute GVHD of the skin, and we diagnosed intestinal thrombotic microangiopathy from the results of a gut biopsy. This patient was successfully managed by diminishing immunosuppressive treatment. Four patients who had blood cultures positive for bacterial infection (*Pseudomonas aeruginosa*, *Acinetobacter lwoffii*, *Corynebacterium* sp, and *Staphylococcus* sp) within 28 days after transplantation were successfully treated with antibiotics. Invasive aspergillosis was encountered in 2 patients (1 proven and 1 possible case). In the proven case, the patient had bronchiolitis obliterans, which was the ultimate cause of death. Of the 17 patients, CMV antigenemia was detected in 12 patients, 2 of whom had CMV colitis.

### 3.4. Graft-versus-Host Disease


Acute GVHD of grades II to IV was diagnosed in 8 patients (48%; 95% confidence interval [CI], 36%-59%); the GVHD was grade II in 3 patients and grade IV in 5. The median time to the onset of acute GVHD was 32 days (range, 20-81 days) after transplantation (Figure 2A). Two of 4 patients who skipped methotrexate treatment on day 11 because of severe mucositis developed grade IV acute GVHD. Two of the 5 patients with grade IV acute GVHD subsequently died. One of these patients had acute GVHD after the withdrawal of CsA treatment at the time of leukemia relapse, and the other patient had received bone

**Table 2.**  
Transplantation Outcomes\*

Patient No.	Time to		Acute GVHD					mPSL, mg/kg	Response	Chronic GVHD (Involved Organs)	Follow-up, d	Current Disease Status	Cause of Death
	ANC >0.5 × 10 <sup>9</sup> /L, d	Platelets >20 × 10 <sup>9</sup> /L, d	Grade	Skin	Liver	Gut							
1	16	26	IV	3	4	4	2	PG	NE	121	Dead	Acute GVHD	
2	35	30	0	0	0	0	—	—	NE	133	Dead	Relapse	
3	17	—	0	0	0	0	—	—	NE	56	Dead	Relapse	
4	14	15	IV	4	0	0	2	CR	Ext (skin, mouth, eyes, liver, lung)	439	Dead	BO	
5	15	22	I	1	0	0	—	—	Ext (skin, mouth, liver)	286	Dead	IP	
6	21	38	II	3	0	0	—	—	NE	260	Dead	Relapse	
7	14	25	I	2	0	0	—	—	Ext (mouth, liver)	687+	CR, alive		
8	20	30	II	3	1	0	1	PR	Ext (skin)	667+	CR, alive		
9	22	—	II	3	0	0	—	—	Ext (skin, mouth, eyes)	336	Dead	Organizing pneumonia	
10	18†	—	0	0	0	0	—	—	NE	94	Dead	Secondary graft failure	
11	16	23	0	0	0	0	—	—	Ext (skin, mouth)	564+	CR, alive		
12	16†	—	IV	2	4	3	2	PG	NE	69	Dead	Acute GVHD	
13	18	23	I	1	0	0	1	CR	Ext (mouth, eyes, liver)	525+	CR, alive		
14	18	—	IV	3	4	2	1	UE	NE	64	Dead	Relapse	
15	14	16	0	0	0	0	—	—	Ext (mouth, eyes)	511+	CR, alive		
16	26	112	0	0	0	0	—	—	Lim (mouth)	463+	CR, alive		
17	22	38	IV	4	0	0	2	CR	Ext (skin, mouth, eyes, liver, lung)	276	Dead	BO + aspergillosis	

\*ANC indicates absolute neutrophil count; GVHD, graft-versus-host disease; mPSL, methylprednisolone; PG, progressive response; NE, not evaluable; CR, complete response; Ext, extensive disease; BO, bronchiolitis obliterans; IP, interstitial pneumonitis; PR, partial response; UE, unevaluated; Lim, limited disease.

†Secondary graft failure occurred after neutrophil recovery.



**Figure 1.** Engraftment after unrelated-donor bone marrow transplantation following reduced-intensity conditioning expressed as the cumulative probability of a neutrophil count  $>0.5 \times 10^9/L$  (A) and a platelet count  $>20 \times 10^9/L$  (B). All patients achieved neutrophil recovery, but 5 patients did not achieve platelet recovery. The median times until neutrophil and platelet recoveries were 18 days (range, 14-35 days) and 26 days (15-112 days), respectively. Late graft failure was observed in 2 patients.

marrow from a donor with allele-level mismatches at 3 HLA loci. Two patients with grade IV acute GVHD involving only the skin were successfully treated with methylprednisolone. Grade II acute GVHD involving only the skin was treated solely with CsA in 2 patients (Table 2). In 7 patients without relapse or secondary graft failure, CsA was tapered from a median of day 120 (range, day 96-169). Only 2 of the 7 patients were able to discontinue CsA (at days 203 and 288). Chronic GVHD was documented in all patients who

survived beyond day 100 (1 with limited GVHD, 9 with extensive disease). There was no significant correlation between HLA disparity at the allele level and the incidence of GVHD, although it was difficult to analyze the data statistically because of the small number of patients in this study.

### 3.5. Survival and Causes of Death

The median follow-up period was 286 days (range, 56-687 days). Overall, 11 patients died, but 6 patients are currently in remission (2 in remission and 4 not in remission at the time of transplantation). The estimated 100-day and 1-year nonrelapse mortality rates were 14% (95% CI, 12%-17%) and 46% (95% CI, 33%-57%), respectively (Figure 2B). Estimated 1-year overall survival and progression-free survival rates were both 41% (95% CI, 32%-51%; Figure 3). There were 4 deaths due to recurrent or progressive disease at a median time of 55 days (range, 32-93 days). The causes of the 7 treatment-related deaths included acute GVHD ( $n = 2$ ), secondary graft failure with sepsis ( $n = 1$ ), interstitial pneumonitis ( $n = 1$ ), organizing pneumonia ( $n = 1$ ), bronchiolitis obliterans ( $n = 1$ ), and bronchiolitis obliterans with invasive aspergillosis ( $n = 1$ ).

## 4. Discussion


In our previous study in an unrelated-donor BMT setting, 5 patients underwent conditioning with a combination of Flu ( $30 \text{ mg/m}^2$  for 6 days) or cladribine ( $0.11 \text{ mg/kg}$  for 6 days), BU ( $4 \text{ mg/kg}$  for 2 days), and antithymocyte globulin ( $2.5 \text{ mg/kg}$  for 4 days) without TBI, but secondary graft failure in 2 of these patients alerted us to a possible higher risk of graft rejection when we used bone marrow instead of peripheral blood cells as the stem cell source. In this study, we demonstrated that the addition of 4 Gy of TBI to the widely applied combination of Flu ( $30 \text{ mg/m}^2$  for 6 days) and BU ( $4 \text{ mg/kg}$  for 2 days) reduces the risk of graft failure and enables the rapid achievement of full donor chimerism without donor lymphocyte infusion (DLI) and that the regimen-related toxicity was acceptable. Nevertheless, a relatively high incidence of nonrelapse mortality was observed. We lost 4 patients who developed extensive chronic GVHD and subsequent pulmonary complications in the later phase, more than 6 months after transplantation. Because many patients develop extensive GVHD, we assume that the pulmonary complications were primarily due to GVHD and not the consequence of our reduced-intensity stem cell transplantation (RIST) regimen incorporating 4 Gy of TBI. However, Deeg et al reported that more pulmonary compli-

**Table 3.**

Maximum Toxicities (N = 17)\*

Grade	Cardiac, n	Mucositis, n	GI, n	Hepatic, n	CNS, n	Hyponatremia, n	Pulmonary, n	Renal, n
0	12	0	9	1	16	6	11	15
I	4	0	3	2	0	7	2	0
II	0	2	4	7	0	0	0	2
III	1	15	1	5	1	4	4	0
IV	0	0	0	2	0	0	0	0

\*GI indicates gastrointestinal tract; CNS, central nervous system.




**Figure 2.** Cumulative incidence of acute GVHD (grades II-IV) (A) and nonrelapse mortality (B) after unrelated bone marrow transplantation following reduced-intensity conditioning. Acute GVHD (grades II-IV) was diagnosed in 8 patients (48%) (grade II in 3 patients and grade IV in 5) at a median of day 32 (range, day 20-81). The estimated 100-day and 1-year nonrelapse mortality rates were 14% and 46%, respectively.

cations developed in patients with aplastic anemia who received 4 to 6 Gy of TBI in combination with cyclophosphamide/antithymocyte globulin for unrelated-donor BMT than in patients who received 2 Gy TBI [20]. These investigators recommended that a 2-Gy TBI dose is sufficient to allow stable engraftment without increased toxicities, and this proposal should be evaluated in future studies. On the other hand, Maris et al described a nonmyeloablative conditioning regimen consisting of 2 Gy TBI and Flu (90 mg/m<sup>2</sup>) for unrelated-donor HSCT [8]. In their study, the use of bone marrow rather than G-CSF-mobilized peripheral blood cells as the source of hematopoietic stem cells led to a lower engraftment rate (56% versus 85%), as well as lower rates of overall survival (33% versus 57%) and progression-free survival (17% versus 44%). Because bone marrow is currently the only stem cell source available from volunteer donors in Japan, we may need a more intensified regimen than the combination of 2 Gy TBI and 90 mg/m<sup>2</sup> Flu.

In this study, the rates of acute GVHD of grades II to IV and extensive chronic GVHD in patients who survived for more than 100 days were 48% and 90%, respectively. Grade IV acute GVHD was the primary cause of death in 2

patients. Moreover, the quality of life of patients who develop extensive chronic GVHD rapidly deteriorates, particularly in elderly patients. Although CsA was tapered from a median of day 120 in this series, it might be better to delay the start of CsA tapering in elderly patients, who are associated with higher GVHD rates. Studies have incorporated in vivo T-cell depletion through the addition of antithymocyte globulin or alemtuzumab in order to reduce the risk of GVHD [21-26]. In the study reported by Chakraverty et al, severe GVHD following RIST from an unrelated donor was decreased with in vivo use of alemtuzumab in the preparative regimen [23]. In their study, the rates of acute GVHD (grades II to IV) and chronic GVHD were 21% and 8%, respectively. The long half-life of alemtuzumab (15-21 days) may disturb the induction of full donor chimerism, however. If patients cannot achieve full donor chimerism, the usual option is DLI, which carries a risk of GVHD [26]. Moreover, lymphocytes for DLI are not always available for every patient, particularly in unrelated-donor transplantation settings. In this regard, we think that a regimen that routinely involves DLI after transplantation cannot be considered a universal strategy. In the present study, 2 patients who had



**Figure 3.** Kaplan-Meier actuarial probability of overall survival (OS) (A) and progression-free survival (PFS) (B) after unrelated-donor bone marrow transplantation following reduced-intensity conditioning. The median follow-up was 286 days (range, 56-687 days). The 1-year OS and PFS rates were both 41%. All 6 of the surviving patients (2 in remission and 4 not in remission at transplantation) remain in remission.



secondary graft failure did not receive DLI, because of grade IV acute GVHD in 1 patient and a reduced performance status in the other. Another approach to preventing severe GVHD is the use of novel immunosuppressive regimens. Several combinations of agents for GVHD prophylaxis, including CsA/mycophenolate mofetil [8,14,16] and tacrolimus/methotrexate [10,15,27], have been reported previously, and their value should be tested in prospective trials.

The induction of adequate antileukemic activity is another primary concern with a RIST procedure, particularly for patients with refractory diseases. de Lima et al reported a promising regimen that consisted of once-daily intravenous BU (130 mg/m<sup>2</sup> for 4 days) and Flu (40 mg/m<sup>2</sup> for 4 days) for patients with AML or MDS [27]. Replacement of oral BU with an intravenous preparation may result in an improved toxicity/survival profile. In our series, 4 patients achieved remission after RIST, although they were not in remission at the time of transplantation. Hence, it is likely that the antileukemic effect exerted by 4 Gy TBI in combination with Flu and BU is valuable even for the immediate control of leukemic blasts, although this possibility needs to be confirmed in further studies. The use of DLI has allowed the rescue of relapsed patients after allogeneic HSCT. In this study, however, we did not give DLI to 4 patients with progressive or relapsed diseases after transplantation because the relevance of the graft-versus-leukemia effect in rapidly proliferating diseases was not fully established and 2 of the patients had developed acute GVHD.

In conclusion, our regimen of 4 Gy TBI, Flu (180 mg/m<sup>2</sup>), and BU (8 mg/kg) was effective in reducing the risk of graft failure following unrelated-donor transplantation. We confirmed, however, that a high incidence of nonrelapse mortality, primarily due to GVHD and/or pulmonary complications, still remains a major obstacle for the wider application of this procedure to elderly or medically infirm patients. Further studies to identify ways to ameliorate transplantation-related toxicities are urgently required.

## References

- Kanda Y, Chiba S, Hirai H, et al. Allogeneic hematopoietic stem cell transplantation from family members other than HLA-identical siblings over the last decade (1991-2000). *Blood*. 2003;102:1541-1547.
- Giralt S, Estey E, Albitar M, et al. Engraftment of allogeneic hematopoietic progenitor cells with purine analog-containing chemotherapy: harnessing graft-versus-leukemia without myeloablative therapy. *Blood*. 1997;89:4531-4536.
- Khouri IF, Keating M, Korbling M, et al. Transplant-lite: induction of graft-versus-malignancy using fludarabine-based nonablative chemotherapy and allogeneic blood progenitor-cell transplantation as treatment for lymphoid malignancies. *J Clin Oncol*. 1998;16:2817-2824.
- Slavin S, Nagler A, Naparstek E, et al. Nonmyeloablative stem cell transplantation and cell therapy as an alternative to conventional bone marrow transplantation with lethal cytoreduction for the treatment of malignant and nonmalignant hematologic diseases. *Blood*. 1998;91:756-763.
- Childs R, Clave E, Contentin N, et al. Engraftment kinetics after nonmyeloablative allogeneic peripheral blood stem cell transplantation: full donor T-cell chimerism precedes alloimmune responses. *Blood*. 1999;94:3234-3241.
- Bornhauser M, Thiede C, Schuler U, et al. Dose-reduced conditioning for allogeneic blood stem cell transplantation: durable engraftment without antithymocyte globulin. *Bone Marrow Transplant*. 2000;26:119-125.
- McSweeney PA, Niederwieser D, Shizuru JA, et al. Hematopoietic cell transplantation in older patients with hematologic malignancies: replacing high-dose cytotoxic therapy with graft-versus-tumor effects. *Blood*. 2001;97:3390-3400.
- Maris MB, Niederwieser D, Sandmaier BM, et al. HLA-matched unrelated donor hematopoietic cell transplantation after nonmyeloablative conditioning for patients with hematologic malignancies. *Blood*. 2003;102:2021-2030.
- Bornhauser M, Thiede C, Platzbecker U, et al. Dose-reduced conditioning and allogeneic hematopoietic stem cell transplantation from unrelated donors in 42 patients. *Clin Cancer Res*. 2001;7:2254-2262.
- Giralt S, Thall PF, Khouri I, et al. Melphalan and purine analog-containing preparative regimens: reduced-intensity conditioning for patients with hematologic malignancies undergoing allogeneic progenitor cell transplantation. *Blood*. 2001;97:631-637.
- Nagler A, Aker M, Or R, et al. Low-intensity conditioning is sufficient to ensure engraftment in matched unrelated bone marrow transplantation. *Exp Hematol*. 2001;29:362-370.
- Kroger N, Sayer HG, Schwerdfeger R, et al. Unrelated stem cell transplantation in multiple myeloma after a reduced-intensity conditioning with pretransplantation antithymocyte globulin is highly effective with low transplantation-related mortality. *Blood*. 2002;100:3919-3924.
- Bornhauser M, Storer B, Slattery JT, et al. Conditioning with fludarabine and targeted busulfan for transplantation of allogeneic hematopoietic stem cells. *Blood*. 2003;102:820-826.
- Niederwieser D, Maris M, Shizuru JA, et al. Low-dose total body irradiation (TBI) and fludarabine followed by hematopoietic cell transplantation (HCT) from HLA-matched or mismatched unrelated donors and postgrafting immunosuppression with cyclosporine and mycophenolate mofetil (MMF) can induce durable complete chimerism and sustained remissions in patients with hematologic diseases. *Blood*. 2003;101:1620-1629.
- Wong R, Giralt SA, Martin T, et al. Reduced-intensity conditioning for unrelated donor hematopoietic stem cell transplantation as treatment for myeloid malignancies in patients older than 55 years. *Blood*. 2003;102:3052-3059.
- Rodriguez R, Parker P, Nademanee A, et al. Cyclosporine and mycophenolate mofetil prophylaxis with fludarabine and melphalan conditioning for unrelated donor transplantation: a prospective study of 22 patients with hematologic malignancies. *Bone Marrow Transplant*. 2004;33:1123-1129.
- Inamoto Y, Oba T, Miyamura K, et al. Stable engraftment after a conditioning regimen with fludarabine and melphalan for bone marrow transplantation from an unrelated donor. *Int J Hematol*. 2006;83:356-362.
- Sullivan KM, Agura E, Anasetti C, et al. Chronic graft-versus-host disease and other late complications of bone marrow transplantation. *Semin Hematol*. 1991;28:250-259.
- Przepiorka D, Weisdorf D, Martin P, et al. 1994 Consensus Conference on Acute GVHD Grading. *Bone Marrow Transplant*. 1995;15:825-828.
- Deeg HJ, Amylon ID, Harris RE, et al. Marrow transplants from unrelated donors for patients with aplastic anemia: minimum effective dose of total body irradiation. *Biol Blood Marrow Transplant*. 2001;7:208-215.
- Bacigalupo A, Lamparelli T, Bruzzi P, et al. Antithymocyte globulin for graft-versus-host disease prophylaxis in transplants from unrelated donors: 2 randomized studies from Gruppo Italiano Trapianti Midollo Osseo (GITMO). *Blood*. 2001;98:2942-2947.
- Hale G, Jacobs P, Wood L, et al. CD52 antibodies for prevention of graft-versus-host disease and graft rejection following transplantation of allogeneic peripheral blood stem cells. *Bone Marrow Transplant*. 2000;26:69-76.
- Chakraverty R, Peggs K, Chopra R, et al. Limiting transplantation-related mortality following unrelated donor stem cell transplantation by using a nonmyeloablative conditioning regimen. *Blood*. 2002;99:1071-1078.

24. Ho AY, Pagliuca A, Kenyon M, et al. Reduced-intensity allogeneic hematopoietic stem cell transplantation for myelodysplastic syndrome and acute myeloid leukemia with multilineage dysplasia using fludarabine, busulphan, and alemtuzumab (FBC) conditioning. *Blood*. 2004;104:1616-1623.
25. van Besien K, Artz A, Smith S, et al. Fludarabine, melphalan, and alemtuzumab conditioning in adults with standard-risk advanced acute myeloid leukemia and myelodysplastic syndrome. *J Clin Oncol*. 2005; 23: 5728-5738.
26. Lim ZY, Ho AY, Ingram W, et al. Outcomes of alemtuzumab-based reduced intensity conditioning stem cell transplantation using unrelated donors for myelodysplastic syndromes. *Br J Haematol*. 2006;135:201-209.
27. de Lima M, Couriel D, Thall PF, et al. Once-daily intravenous busulfan and fludarabine: clinical and pharmacokinetic results of a myeloablative, reduced-toxicity conditioning regimen for allogeneic stem cell transplantation in AML and MDS. *Blood*. 2004;104:857-864.

# Prospective phase II trial to evaluate the complications and kinetics of chimerism induction following allogeneic hematopoietic stem cell transplantation with fludarabine and busulfan

Akiko M. Saito,<sup>1,2\*</sup> Masahiro Kami,<sup>3</sup> Shin-Ichiro Mori,<sup>4</sup> Yoshinobu Kanda,<sup>5</sup> Ritsuro Suzuki,<sup>6</sup> Shin Mineishi,<sup>7</sup> Akiyoshi Takami,<sup>8</sup> Shuichi Taniguchi,<sup>9</sup> Yoshinobu Takemoto,<sup>10</sup> Masamichi Hara,<sup>11</sup> Masaki Yamaguchi,<sup>12</sup> Masayuki Hino,<sup>13</sup> Takashi Yoshida,<sup>14</sup> Sung-Won Kim,<sup>4</sup> Akiko Hori,<sup>4</sup> Yasuo Ohashi,<sup>2</sup> and Yoichi Takaue<sup>4</sup>

<sup>1</sup> Department of Hematology and Oncology, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

<sup>2</sup> Department of Biostatistics/Epidemiology and Preventive Health Sciences, School of Health Sciences and Nursing, University of Tokyo, Tokyo, Japan

<sup>3</sup> Division of Exploratory Research, Institution of Medical Science, The University of Tokyo, Tokyo, Japan

<sup>4</sup> Hematopoietic Stem Cell Transplantation Unit, National Cancer Center Hospital, Tokyo, Japan

<sup>5</sup> Department of Cell Therapy and Transplantation Medicine, University of Tokyo Hospital, Tokyo, Japan

<sup>6</sup> Department of HSCT Data Management, Nagoya University School of Medicine, Nagoya, Japan

<sup>7</sup> Division of Hematology/Oncology, University of Michigan, Ann Arbor, Michigan

<sup>8</sup> Department of Cellular Transplantation Biology, Kanazawa University Graduate School of Medicine, Ishikawa, Japan

<sup>9</sup> Department of Hematology, Toranomon Hospital, Tokyo, Japan

<sup>10</sup> Medical Corporation Kouryokai CPC Clinic, Kagoshima, Japan

<sup>11</sup> Division of Hematology, Ehime Prefectural Central Hospital, Ehime, Japan

<sup>12</sup> Department of Hematology and Immunology, Ishikawa Prefectural Central Hospital, Ishikawa, Japan

<sup>13</sup> Department of Clinical Hematology and Clinical Diagnostics, Graduate School of Medicine, Osaka City University, Osaka, Japan

<sup>14</sup> Department of Internal Medicine, Toyama Prefectural Central Hospital, Toyama, Japan

**This prospective trial assessed the safety and efficacy of allogeneic hematopoietic stem cell transplantation from a HLA-matched donor with a reduced-intensity regimen (RIST) consisting of iv fludarabine 30 mg/m<sup>2</sup> for 6 days and oral busulfan 4 mg/kg/day for 2 days in patients older than 50 years with hematological malignancies. Cyclosporine alone or cyclosporine with short-term methotrexate was randomized for graft-versus-host disease prophylaxis. After 30 patients had been enrolled, an interim analysis was performed, and this report focuses on a precise evaluation of the toxicity profile and chimerism kinetics. Sustained engraftment in all patients, no severe regimen-related toxicity (RRT) within 20 days, and no transplant-related mortality through Day 100 were observed. T-cell (CD3+) full-donor (over 90%) chimerism was observed in 22 of the 30 patients, while the remaining eight had mixed-donor chimerism over 77% on Day 90. Thereafter, five subsequently converted to full-donor chimerism without donor lymphocyte infusion by day 120 ( $n = 4$ ) or Day 180 ( $n = 1$ ). Two showed persistent mixed chimerism without relapse through Day 180. Grade III–IV acute graft-versus-host disease and extensive chronic graft-versus-host disease occurred in 10% and 73%, respectively. With a median follow-up of 1.5 years, overall survival and disease-free survival at 1 year was 83% and 62%, respectively. Seven patients hematologically relapsed overall, and five of them had myelodysplastic syndrome with poor prognostic factors. In older patients, RIST with fludarabine and busulfan was associated with acceptable toxicities and a satisfactory antileukemia effect, regardless of the early chimerism status. *Am. J. Hematol.* 00:000–000, 2007. © 2007 Wiley-Liss, Inc.**

## Introduction

Allogeneic hematopoietic stem cell transplantation (HSCT) is a potentially curative treatment of choice for hematological malignancies. However, many centers limit HSCT to younger patients because of the threat of a higher risk of treatment-related toxicities including graft-versus-host disease (GvHD), nonrelapse mortality, and lower disease-free survival (DFS) in the older population, although the median age of onset of chronic myeloid leukemia (CML) is in the sixth decade of life, and the peak incidence of acute myeloid leukemia (AML) and myelodysplastic syndrome (MDS) is in the seventh decade. To overcome this obstacle, allogeneic HSCT with a reduced-intensity (RIST) or nonmyeloablative conditioning regimen has recently been explored for patients who are ineligible to receive conventional myeloablative HSCT (CIST) due to age limits or comorbidities. Many studies suggested that RIST is a reasonable option for older patients or

patients with comorbidities with acceptable treatment-related complications or morbidity, while preserving adequate anti-tumor effects [1–12]. However, these studies mostly pursued different variables including disease types, stages [1,4–6,8,12], donor type [1,2,5,10], graft source [1,2], conditioning regimens [4,5,7,9], and/or GvHD prophylaxis [1,4,9]. This

Contract grant sponsor: Health and Labour Sciences Research Grant from the Ministry of Health, Labour and Welfare.

\*Correspondence to: Akiko M. Saito, MD, PhD, Dana-Farber Cancer Institute, 44 Binney Street, SM 275 Boston, MA 02115.  
E-mail: akiko\_saito@dfci.harvard.edu

Received for publication 11 January 2007; Revised 5 March 2007; Accepted 30 March 2007

*Am. J. Hematol.* 00:000–000, 2007.

Published online in Wiley InterScience (www.interscience.wiley.com).

DOI: 10.1002/ajh.20977

**TABLE I. Patient and Donor Characteristics**

UPN	Patient age/sex	Donor age/sex	Stage and diagnosis	IPSS/cytogenetic risk in MDS patients	CD34+ cells (10 <sup>6</sup> )	Blood type patient/donor
1	60/M	46/F	MDS (RA)	Intermediate-2/Poor	3.92	A/A
2	61/M	54/F	MDS (RAEB)	Intermediate-2/Poor	5.84	O/O
3	67/M	60/F	AML (M4) in 2CR		2.74	B/O
4	60/M	55/F	AML (M2) in 2CR		4.58	B/B
5	63/M	60/M	CML in 2CP		12.59	O/O
6	54/M	59/F	MDS (RAEB)	High/Intermediate	5.4	O/A
7	52/M	55/F	AML (M2) in 1CR		6.77	A/A
8	61/M	54/M	AML (M1) in 1CR		3.29	B/A
9	58/F	64/F	CML in 1CP		2.9	A/AB
10	64/F	59/F	ALL (L2) in 1CR		5.54	A/A
11	55/M	44/M	AML (M1) in 1CR		3.13	A/A
12	55/F	51/F	CML in 1CP		4.94	A/O
13	52/F	42/M	AML (M4) in 1CR		3.59	A/A
14	59/M	64/M	MDS (RAEB)	Intermediate-2/intermediate	3.58	A/AB
15	59/M	56/M	MDS (RA)	Intermediate-1/Good	3.58	AB/A
16	53/F	55/F	MDS (RA)	Intermediate-2/Poor	2.2	O/O
17	55/F	68/M	AML (M3) in 2CR		2.63	A/A
18	54/M	50/M	MDS (RA)	Intermediate-1/Poor	3.74	O/B
19	51/M	44/F	AML (M1) in 1CR		4.86	AB/A
20	64/F	66/M	CML in 2CP		3.59	O/A
21	68/F	64/M	MDS (RAEB)	Intermediate-1/Good	3.56	B/B
22	53/M	44/M	MDS (RAEB)	High/Intermediate	7.2	B/B
23	60/F	53/M	AML (M2) in 1CR		2.83	A/B
24	59/M	62/M	AML (M4) in 2CR		5.47	A/O
25	51/F	47/F	MDS (RAEB)	Intermediate-2/Poor	5.93	A/A
26	59/M	62/F	MDS (RA)	Intermediate-2/Poor	4.02	B/O
27	59/M	48/M	AML (M2) in 2CR		4.94	B/A
28	56/M	62/F	MDS (RAEB-t)	High/Good	4.38	AB/A
29	53/F	62/F	AML (M2) in 1CR		3.06	O/O
30	54/F	63/M	AML (M2) in 1CR		6.47	A/O

M, male; F, female; MDS, myelodysplastic syndrome; AML, acute myeloid leukemia; CML, chronic myeloid leukemia; ALL, acute lymphoblastic leukemia; RA, refractory anemia; RAEB, refractory anemia with excess blasts; RAEB-t, refractory anemia with excess blasts in transformation; CR, complete remission; CP, chronic phase. All donors were HLA-matched siblings.

makes overall interpretation of studies difficult. Additionally, there has been no study to prospectively assess whether RIST consisting of 180 mg fludarabine plus 8 mg/kg busulfan without antithymocyte globulin actually produces less significant organ toxicities and treatment-related toxicities in an older patient population. Information regarding the impact of the speed and degree of lineage-specific donor chimerism on clinical outcomes after RIST in older patients has been limited [3,8,13–17]. Moreover, even studies evaluated with more homogeneous patient population, type of GvHD prophylaxis and/or tempo of withdrawal of immunosuppressive agents varied depending on transplant centers and a feasible prophylaxis regimen for acute GvHD has not been well evaluated in RIST, which is considered to require a sophisticated balance between GvHD and a graft-versus-leukemia (GvL) effect.

To address these points, we conducted a prospective randomized clinical trial to evaluate the safety and efficacy of RIST with fludarabine and oral busulfan in patients aged over 50 years and with appropriate GvHD prophylaxis. In this report, the results of an interim analysis, including clinical outcomes, complications, and chimerism kinetics, were compared with those previously published in the literature.

**Patients and Methods**


**Patient eligibility and accrual**

Eligible patients ranged in age from 50 to 69 years (median 58.5, range 51–68 years) and had a hematological malignancy, including

AML or acute lymphoblastic leukemia (ALL) in 1st or 2nd complete remission (CR), CML in 1st or 2nd chronic phase (CP), and MDS. They were required to have an HLA-identical related donor. The study protocol was reviewed and approved by the institutional review boards of the participating transplantation centers (Appendix). Eligible patients and their donors gave written informed consent before enrollment. The enrollment criteria included a performance status (PS) of the Eastern Cooperative Oncology Group (ECOG) of less than two, a serum creatinine concentration of less than 2.0 mg/dl, a cardiac ejection fraction of more than 50%, arterial oxygen saturation without supplemental oxygen of more than 93%, liver function tests less than fourfold the upper limit of normal, total bilirubin less than 2.0 mg/dl, no active infection, and no previous allergy for drugs used for conditioning or GvHD prophylaxis. Donors were required to have a normal physical examination, and normal values in the serum chemistry and blood counts, and negative results of serologic testing for human immunodeficiency virus and hepatitis B. The patient and donor characteristics are shown in Table I. Those with AML/ALL in 1st CR, CML in 1st CP, or MDS in refractory anemia were defined as low risk, and the others were defined as high risk. All 12 patients with MDS except one (UPN 22) were transfusion dependent, and all those were grouped according to the International Prognostic Scoring System (IPSS) into intermediate or high risk at the time of transplantation: Intermediate-1, n = 3; intermediate-2, n = 6; high risk, n = 3. By IPSS criteria, 3 patients had good-risk, 3 had intermediate-risk, and 6 had poor-risk cytogenetics.

**Donor selection and blood stem cell harvest**

Related donors were selected based on compatibility of HLA-A, B and DRB1 by intermediate- or high-resolution DNA typing. After G-CSF treatment, apheresis procedures were performed daily until at least 2.0 × 10<sup>6</sup> CD34+ cells per kilogram of the recipient's body weight, up to three times, and all of the collected cells were cryopreserved until stem cell infusion.



**Figure 1. Treatment schedule.** CyA; cyclosporine, MTX; methotrexate. \*1: When acute GvHD was not observed, CyA was tapered by 10% a week starting at Day 28, and was eliminated by Day 100. When mixed chimerism was seen without active acute GvHD over Day 60, CyA was tapered and discontinued within 2 weeks. Patients who did not convert to complete chimerism after CyA withdrawal received donor lymphocyte infusion.

### Treatment schedule

The treatment schedule is shown in Fig. 1. The conditioning regimen consisted of fludarabine (30 mg/m<sup>2</sup>/day) infused over 30 min once a day on Days 8, 7, 6, 5, 4, and 3, and oral busulfan (4 mg/kg/day) on Days 6 and 5. To prevent seizures, the patients received oral valproate sodium, at a dose of 600 mg divided into 3 doses 2 days before busulfan administration, and this was continued until 24 hr after the last dose of busulfan.

Patients were randomized to receive either cyclosporine (CyA) alone or CyA plus short-term methotrexate (MTX) for GvHD prophylaxis. Randomization was performed by stratifying according to disease (AML, ALL, CML or MDS), transplant center, age (less than 60 years or more than or equal to 60 years), and sex (male or female). All patients received 3 mg/kg/day CyA by continuous iv infusion daily from Day 1 to maintain a therapeutic trough level of 250–400 ng/ml, and thereafter orally in an attempt to maintain a therapeutic trough level of 150–250 ng/ml. The patients who were assigned to CyA plus short-term MTX received a dose of 10 mg/m<sup>2</sup> iv MTX on Day +1, and 7 mg/m<sup>2</sup> on Days +3 and +6 after stem cell infusion. CyA was tapered starting at Day 28 in the absence of acute GvHD and was discontinued by Day 100 after transplantation. When a patient did not achieve complete donor chimerism by Day 60, CyA was tapered rapidly and discontinued within 2 weeks if clinically feasible, since anti-leukemic effect was presumed to occur after development of complete donor chimerism [14]. Cases of Grade II–IV acute GvHD were treated with 2 mg/kg/day of methylprednisolone in addition to CyA.

### Supportive care

The following infection prophylaxis was recommended: prophylactic antibiotics (fluoroquinolones) were given during cytopenia, fluconazole (200 mg/day) was given at the start of conditioning and continued until the discontinuation of immunosuppressant, and oral acyclovir (1,000 mg/day) or iv acyclovir (750 mg/day) was given for prophylaxis of herpes simplex virus (HSV) and varicella zoster virus (VZV) from Day -7 to Day 35. Prophylaxis against *Pneumocystis carinii* was consisted of trimethoprim-sulfamethoxazole after neutrophil engraftment ( $\geq 0.5 \times 10^9 \text{ L}^{-1}$ ) and was continued until the discontinuation of immunosuppressant. During the first 100 days after transplantation, cytomegalovirus antigenemia assay with HRP-C7 or C10/C11 monoclonal antibody was performed weekly after neutrophil engraftment until Day 100 after transplantation. Pre-emptive therapy with ganciclovir was recommended upon the detection of positive antigenemia and was continued until it became negative. Patients were treated with G-CSF from Day +6 to neutrophil engraftment.

### Chimerism analysis

Hematopoietic chimerism was evaluated with regard to peripheral T cell (CD3+) fraction by an analysis of DNA microsatellite polymorphisms by polymerase chain reaction (PCR) with D18S51, D20S471, and D22S684 fluorescence-labeled primers, which identified differences

between patient and donor (on the basis of polymorphisms found in pretransplant patient/donor samples) using an BECKMAN COULTER CEQ8000 GENETIC ANALYSIS SYSTEM. T cell (CD3+) chimerism studies post HSCT were performed on Days 30, 60, 90, 120, and thereafter every other month through 1 year.

### Assessment of response

Day 0 was defined as the day of stem cell infusion day. The day of neutrophil engraftment was defined as the first of two consecutive days on which the patient's absolute neutrophil count was above  $0.5 \times 10^9 \text{ L}^{-1}$ . The day of platelet engraftment was defined as the first of seven consecutive days on which the platelet count was above  $20 \times 10^9 \text{ L}^{-1}$  without platelet transfusion.

Regimen-related toxicity (RRT) was graded using the Seattle criteria [18] on the day before the initiation of conditioning regimens and at least 3 days a week until Day 20 after transplantation. All other observed adverse events were graded according to the National Cancer Institute Common Toxicity Criteria version 2.0 (NCI-CTC ver. 2.0) until Day 100 after transplantation. Infectious diseases were diagnosed based on any positive blood culture or histologic evidence of tissue invasion.

To evaluate the general condition of patients associated with the toxicity profile, PS, and dietary oral intake were also reported at least three times a week during the initial hospitalization and once a week afterwards up to Day 100 post-transplant.

The diagnosis and grading of acute and chronic GvHD was made based on the date of onset (within or beyond 100 days) and clinical findings in conjunction with biopsy of the skin and digestive tract using the published criteria [19,20]. Patients who survived 100 days or longer were evaluable for the assessment of chronic GvHD.

### Pharmacokinetic studies of fludarabine phosphate and busulfan

Blood sampling for pharmacokinetic studies was done on Day -5 to investigate the effect of concomitant busulfan administration on the pharmacokinetics of 2-fluoro-ara A (2F-ara-A), which is the major metabolite of fludarabine phosphate. Blood samples for determining the 2F-ara-A plasma level were collected at 0, 0.5, 1, 2, 5, and 23.5 hr after the 4th infusion of fludarabine. We also obtained blood samples for determining the busulfan plasma level at 0, 0.5, 1, 1.5, 2, 3, and 6 hr after the sixth administration of busulfan (1 mg/kg/dose for 8 times). Blood samples were taken in tubes containing heparin and erythro-9-(2-hydroxy-3-nonyl)adenine. Plasma was obtained by centrifugation, and then transported to the laboratory and were stored at -20°C until analysis. Plasma levels of 2F-ara-A and busulfan were determined using high-performance liquid chromatography with fluorescence and UV detection, respectively. The accuracy and precision of the assays for 2F-ara-A and busulfan were confirmed by measuring QC samples of both before this study. The maximum concentration of drug in plasma after drug administration ( $C_{max}$ ,  $C_{peak}$ ) and the time to reach

the maximum concentration following drug administration ( $T_{max}$ ) were observed. The area under the plasma concentration-versus-time curve (AUC) for 2F-ara-A or busulfan was calculated by dividing the administered dose by the final plasma clearance estimate, whereas the plasma clearance was determined by modeling all plasma concentration versus time data. Terminal half-lives ( $T_{1/2}$ ) were calculated from the primary parameters.

### Statistical analysis

The primary endpoint of this study was to determine the percentage of patients who were alive at 100 days after transplantation with complete donor chimerism (over 90%) achieved by Day 90. Secondary endpoints included the time to engraftment of neutrophils and platelets, the incidence and severity of RRT, the incidence and severity of acute and chronic GvHD, the anti-leukemia effect, DFS, and overall survival (OS). A descriptive statistical analysis was performed to assess patient baseline characteristics and disease. Time to engraftment, complete chimerism, acute or chronic GvHD, OS, and DFS were calculated using the Kaplan-Meier method. OS was defined as the time between stem cell infusion to death from any cause. DFS was defined as the time between stem cell infusion to relapse and death from any cause, whichever occurred first. After 30 patients had been enrolled in the study, a data and safety monitoring committee undertook an interim analysis. This analysis, completed in October 2004, included data for the primary endpoint, i.e. survival at Day 100 and chimerism status at Day 90, and data on acute and chronic GvHD, survival, chimerism status, and anti-tumor effect through Day 180. Neither of the predefined criteria for stopping the study was met; however, a review of available safety data including incidence and severity of RRT and Day 100 mortality indicated that this conditioning regimen was adequately safe for older patients. According to the recommendation of the committee, we decided to continue the study and published an interim report when 30 patients were enrolled and evaluated without comparing the two different GvHD prophylaxis procedures. This report includes data on these 30 patients with all available follow-up data through December 2005, and does not include the results of a comparison of the two different GvHD prophylaxis procedures.

## Results

### Engraftment and chimerism analysis

The results are summarized in Table II. One and four patients were not evaluated for neutrophil and platelet engraftment, respectively, because they did not show a nadir. The remaining patients achieved sustained engraftment and none experienced graft failure. The median number of days to achieve a neutrophil count  $\geq 0.5 \times 10^9 L^{-1}$  was 13 (range, 10–25 days), and this was 18 (range, 11–24 days) for a platelet count  $\geq 20 \times 10^9 L^{-1}$  without transfusion. Full-donor (over 90%) T-cell (CD3+) chimerism was observed in 2 and 9 of the 30 patients on Day 30 and Day 60, respectively (median [range], Day 30:71 [40 to  $\geq 90$ ] %, day 60:81 [41 to  $\geq 90$ ] %). Twenty-two patients achieved full-donor chimerism, while the remaining eight patients had mixed chimerism ranging from 78% to 88% on Day 90. Among those with mixed chimerism on Day 90, five subsequently converted to full-donor chimerism without early CyA withdrawal because of the severe acute GvHD ( $n = 2$ : UPN 1 and 15) and/or donor lymphocyte infusion (DLI) by day 120 ( $n = 4$ ) or day 180 ( $n = 1$ ). One achieved full-donor chimerism on Day 120 after DLI since the patient did not respond to the discontinuation of immunosuppressive drugs, and two had persistent mixed chimerism without relapse through 180 days after transplantation (71% and 75% donor-type chimerism on Day 180). The diagnoses of two patients with persistent mixed chimerism through Day 180 were CML and MDS, and they had not received preceding cytotoxic chemotherapy; the patient with CML (UPN 12) received immunomodulators, imatinib mesylate and hydroxyurea, and the patient with MDS (UPN 21) received low-dose cytarabine and aclarubicin in combination with granulocyte colony stimulating factor before RIST.

### Regimen-related toxicities, complications, and general condition

The frequencies of Grade I–IV organ toxicities within 20 days after transplantation are listed in Table III. Although non-fatal toxicities including Grade I/II were seen in all 30 patients, all of the observed episodes were reversible and in no case required suspension of fludarabine. Stomatitis was the most frequently observed organ toxicity (57%, 17/30), with 47% of them (8/17) had Grade II events. None of the patients experienced veno-occlusive disease of the liver (VOD). Twenty patients had at least one episode of infectious complications within the first 100 days, with a total of 44 documented episodes (median, 2; range, 1–7 episodes) within the first 100 days after transplantation. These included proven bacterial infection (1 episode), suspected bacterial infection (1), suspected fungal infection (2), cytomegalovirus antigenemia (6), HSV infection (1), suspected viral infection (1), and uncertain causes (33). All infectious complications were recovered with or without appropriate antibiotic therapy.

The median PS for the first 28 days was 0 (range, 0–3). The worst PS of 2 ( $n = 5$ ) or 3 ( $n = 2$ ) within the first 28 days was experienced temporarily due to infection ( $n = 2$ ), Grade III GvHD ( $n = 1$ ), and nausea/vomiting ( $n = 4$ ). Those ( $n = 6$ ) observed from Day 29 to Day 100 were all caused by Grade II or III acute GvHD. A one-thirds reduction in dietary oral intake was temporarily seen in 20 and 11 patients within the first 28 days and from 29 days to 100 days post HSCT, respectively, which resulted from nausea/vomiting ( $n = 18$ ) and treatment-related mucositis ( $n = 2$ ) within Day 28, and Grade II–III acute GvHD ( $n = 9$ ), prolonged infection with Grade II acute GvHD ( $n = 1$ ) and gastroesophageal reflux disease ( $n = 1$ ) between Day 29 and Day 100.

### GvHD

Grade I–IV acute GvHD at 100 days was documented, respectively, in 5 (17%), 15 (50%), 3 (10%), and 0 (0%) patients. The median time to the occurrence of Grade II–IV acute GvHD was 74 days (range, 18–100 days). All 30 patients survived beyond Day 100 and were evaluated for chronic GvHD. Twenty-six of the 30 patients (87%) developed chronic GvHD (limited type in four cases and extensive type in 22 cases) with the onset at a median of 123 days after transplantation (range, 116–217 days).

### Disease response, survival, and cause of death

No patient died within the first 100 days, and the median follow-up period was 555 days (149–1114 days) after transplantation. Twenty-nine of the 30 patients achieved CR within 100 days after transplantation, but two of them with MDS, who had poor-risk cytogenetics and were classified into intermediate-2, subsequently relapsed on Day 141 (UPN 26) and Day 156 (UPN 25). One was treated with DLI (UPN 25) and showed a temporary response, but died because of the disease progression on Day 401. The other patient (UPN 26) did not respond to DLI and died of progressive disease on Day 412. One patient (UPN 22) with MDS with high risk IPSS achieved full-donor chimerism on Day 90, but could not achieve CR on Day 98 and died with progressive disease on Day 306. This patient showed full-donor chimerism through Day 180. Five other patients died between 100 days and 1 year after transplantation (149, 151, 169, 187, and 354 days). In six patients who died within the first year, two patients were over 60 years and four patients were classified into high risk disease group. Causes of death included progressive disease of MDS with poor IPSS in 1, GvHD and/or its complications in 4, and recurrence of interstitial pneumonia in 1. In four patients, who died of GvHD and/or its complications, all had experienced

**TABLE II. Summary of Clinical Outcomes**

UPN	Chimerism analysis			Post transplant DLI (reason)	GvHD		Infection until day 100 (etiological agent)	Relapse	Outcome (Cause of death)	Follow up
	Day 90(%)				Acute	Chronic				
	Day 90(%)	Day 120(%)	Day 180(%)		Gr II (S, G)	Gr II (S)				
1	88.40	>90	>90	Yes (d662, relapse)	Gr II (S, G)	Extensive	-	-	Alive	1,114
2	85	>90	>90		Gr II (S)	Extensive	Yes (unknown)	Yes (d402)	Dead (recurrent disease and its complication)	652
3	>90	>90	>90		Gr II (S)	Extensive	Yes (S. maltophilia, unknown)	-	Alive	735
4	>90	>90	D		-	-	-	-	Dead (IP)	169
5	>90	>90	>90		Gr II (L)	Extensive	-	-	Alive	731
6	>90	>90	>90		Gr II (S, G)	Extensive	-	-	Alive	716
7	>90	>90	>90		Gr II (S, G)	Extensive	Yes (CMV antigenemia)	-	Dead (GvHD)	354
8	>90	>90	>90		Gr III (S, G, L)	Extensive	Yes (bacteremia susp., fungal susp., CMV antigenemia, unknown)	-	Alive	431
9	>90	>90	>90		-	Extensive	Yes (unknown)	-	Alive	592
10	>90	>90	>90		Gr II (S, G)	Extensive	Yes (unknown)	-	Dead (GvHD)	757
11	>90	>90	>90		Gr II (S)	Extensive	-	-	Alive	360
12	88	79	71		Gr III (S, L)	Extensive	Yes (unknown)	-	Alive	720
13	>90	>90	>90		Gr I (S)	Extensive	Yes (HSV, unknown)	-	Dead (GvHD)	517
14	>90	>90	>90		Gr II (S, G, L)	Limited	Yes (fungal susp., unknown)	-	Dead (GvHD and its complication)	187
15	85	88	>90		Gr III (S, G)	-	-	-	Alive	702
16	84	88	>90		-	Limited <sup>a</sup>	Yes (CMV antigenemia)	-	Alive	642
17	80	>90	D	Yes (d98, mixed chimerism)	Gr II (S, G) <sup>b</sup>	-	-	-	Dead (GvHD and its complication)	149
18	>90	88	>90		Gr II (S)	Extensive	Yes (CMV antigenemia)	-	Alive	729
19	>90	>90	>90		Gr I (S)	Limited	-	-	Alive	737
20	>90	>90	>90		Gr II (G)	-	Yes (CMV antigenemia)	Yes (d147) <sup>c</sup>	Alive	688
21	78	77	75		Gr II (S, L)	Extensive	-	Yes (d364)	Dead (BOOF)	593
22	>90	>90	>90		Gr II (S, G)	Extensive	Yes (unknown)	Yes (d98)	Dead (progressive disease)	306
23	>90	>90	D		-	Extensive	Yes (unknown)	-	Dead (GvHD and its complication)	151
24	>90	>90	>90		Gr II (G)	Extensive	-	Yes (>d365) <sup>d</sup>	Dead (recurrent disease)	825
25	>90	87	>90	Yes (d186, d238, relapse)	Gr II (S)	Extensive	Yes (CMV antigenemia)	Yes (d156)	Dead (recurrent disease)	401
26	>90	>90	>90	Yes (d204, relapse)	Gr I (S)	Extensive	Yes (unknown)	Yes (d141)	Dead (recurrent disease and its complication)	412
27	84	>90	>90		-	Extensive	Yes (unknown)	-	Alive	371
28	>90	>90	>90		Gr II (S)	Extensive	Yes (viral susp., unknown)	-	Alive	365
29	>90	>90	>90		Gr I (S)	Extensive	Yes (unknown)	-	Alive	366
30	>90	>90	>90		Gr I (S)	Limited	Yes (unknown)	Yes (d370)	Alive	370

ND, not done; D, dead; DLI, donor lymphocyte infusion; Gr, grade; GvHD, graft-versus-host disease; GvHD site codes: S-skin, G-gut, L-liver; CMV, cytomegalovirus; susp., suspected; unknown, no microbiological evidence despite symptoms; IP, interstitial pneumonia.

<sup>a</sup>This patient developed a GvHD starting on day 112 after receiving DLI for mixed chimerism.

<sup>b</sup>This patient developed gut GvHD starting on day 92.

<sup>c</sup>CNS relapse without hematological relapse.

<sup>d</sup>This patient relapsed after day 365, but the exact date of relapse is unknown.

**TABLE III. Regimen-Related Toxicities Within 20 Days After HSCT According to the Seattle Criteria in 30 Patients**

Toxicity	Grade			
	1	2	3	4
Heart	1	0	0	0
Bladder	0	1	0	0
Kidney	5	1	0	0
Lung	2	0	0	0
Liver	8	0	0	0
CNS	1	0	0	0
Stomatitis	9	8	0	0
GI toxicity	4	1	0	0

HSCT, hematopoietic stem cell transplantation; CNS, central nervous system; GI, gastro-intestinal.

gut GvHD, three of those developed extensive chronic GvHD and all were treated with corticosteroid.

The Kaplan-Meier estimated probability of OS and DFS at 1 year was, respectively, 83% and 62% (Fig. 2). Both patients age ( $\leq 55$  years versus  $>55$  years) and CD34+ cell dose ( $>5.0 \times 10^6 \text{ kg}^{-1}$  versus  $\leq 5.0 \times 10^6 \text{ kg}^{-1}$ ) were not associated with better outcomes by a stratified analysis (data not shown).


#### Pharmacokinetic results for fludarabine and busulfan

2F-ara-A and busulfan PK parameters were calculated from data obtained from blood samples from six consenting patients (UPN 1, 3–7). After the start of the 4th infusion of fludarabine phosphate (30 mg/m<sup>2</sup>/dose), the maximum plasma level of 2F-ara-A was  $3.12 \pm 1.08 \text{ nmol/ml}$ , with a subsequent decline to  $T_{1/2}$  of  $8.59 \pm 1.57 \text{ h}$ . The AUC (0–24 hr) and CL were  $17.7 \pm 2.82 \text{ nmol hr/ml}$  and  $78.9 \pm 13.1 \text{ ml/min/m}^2$ , respectively. After the 6th administration of busulfan (1 mg/kg/dose for eight times), the maximum plasma level of busulfan was  $1.37 \pm 0.34 \text{ nmol/ml}$ , with a subsequent decline to a  $T_{1/2}$  of  $2.88 \pm 0.65 \text{ hr}$ . The AUC (0–6 hr) and CL were  $4.85 \pm 1.07 \text{ nmol hr/ml}$  and  $3.60 \pm 0.88 \text{ ml/min/m}^2$ , respectively. Since these parameters are similar to those in a previous study with the repeated administration of fludarabine phosphate alone at 15, 20, and 25 mg/m<sup>2</sup>/dose (data not shown), combination with busulfan seemed to have no effect on the pharmacokinetics of 2F-ara-A. The steady-state plasma level of busulfan ( $808 \pm 178 \text{ ng/ml}$ ) was observed to remain within a therapeutic level (600–900 ng/ml) in adults [21].

#### Discussion

In this prospective study, we showed that a combination of fludarabine (180 mg/m<sup>2</sup>) and oral busulfan (8 mg/kg), despite the omission of antithymocyte globulin from the original regimen by Slavin et al. [6], can be successfully used to help prepare patients older than 50 years with hematological malignancies for HSCT from an HLA-matched related donor: All patients achieved sustained engraftment without graft failure, only an insignificant occurrence of RRT and treatment-related complications were seen, and PS and dietary intake were well maintained, which agrees with published observational studies on RIST with fludarabine and busulfan [16,22,23].

The rapid induction of complete donor-type chimerism was considered as an essential part of the RIST procedure. Although all of our patients rapidly developed conventional neutrophil and platelet engraftment, two of the 30 patients without preceding cytotoxic chemotherapy remained in mixed T-cell chimerism during the first 6 months after transplantation. A more rapid induction of T-cell chimerism has



**Figure 2. Kaplan-Meier product estimates of overall survival and disease-free survival.**

been observed in other studies of RIST in patients who had been previously treated with chemotherapy for diseases other than CML or MDS [24]. Although a close association between the occurrence of acute GvHD and the induction of higher levels of donor T-cell chimerism has been reported [14], in our experience over 50% of patients did not achieve complete chimerism at the onset of acute GvHD, demonstrating that mixed chimerism status did not provide absolute protection from GvHD, which is in agreement with data published by Baron et al. [15]. We speculate that differences in the conditioning regimen and GvHD prophylaxis may result in different observations.

While our less intensive regimen was associated with less toxicity, this strategy will only work if modifications to the conditioning regimen intensity that allow early clinical benefits do not also lead to reduced induction of GvL effect or other complications that increase relapse rate or result in worse survival in later time period [25]. A recent observational study from European Group of Blood and Marrow Transplantation Registry compared treatment-related mortality (TRM) and other outcomes between 315 RIST recipients and 407 CIST recipients, who were over 50 years and transplanted from a HLA matched sibling donor [26], and suggested that lower TRM but higher relapse rate were seen in RIST recipients. Given the fact that all three patients, who relapsed within 6 months after transplantation, were MDS with poor prognostic factors, the incidence of relapse in our study seems to be no higher than that in published data for CIST [27–30]. Taussig et al. evaluated the feasibility and safety of the fludarabine based RIST regimen in 16 patients with standard risk diseases [31]. In this study, TRM rate within 100 days was 0%, however, OS and DFS at 1 year read from Fig. 2 were 69% and 56%, respectively, where most of the patients included in this study had early stage diseases and over 30% of patients were aged less than 50 years. Despite the older patient population, our data showing no treatment-related mortality (TRM) within the first 100 days after transplantation and OS and DFS at 1 year of 83% and 62%, respectively, was encouraging.

In a previous report, we suggested that the development of GvHD is not essential for the control of low-risk myeloid malignancies, and that GvHD and infection, rather than relapse, are more important problems to be addressed in these patients [25]. Although our data showed favorable outcomes, six patients with four low risk disease and three patients aged less than 55 years died of GvHD or its complication within the first year should be interpreted with care. The incidence of Grade II–IV acute GvHD in this



study was somewhat higher than that in published literature and our own observational data with elder patients and high risk diseases [25]. However, Grade III–IV acute GvHD was infrequent and none died from acute GvHD. The incidence of chronic GvHD was higher than that in our previous experience (56%) [32] or in other reports [31,33] even after considering inevitable differences in the ethnicity, GvHD prophylaxis and matching practice of HLA, or disease risk. G-CSF mobilized peripheral blood stem cells may have been associated with an increased incidence of GvHD, particularly in its chronic form [34,35]. Conditioning regimen excluded antithymocyte globulin was also a possible explanation of this finding [23]. Most importantly, patients undergoing RIST are usually older than those undergoing CIST, which leads to a higher risk for GvHD [36,37]. Early CyA withdrawal regulation to get speedy achievement of complete donor chimerism after RIST in our protocol might have influenced the increased incidence of Grade II–IV acute GvHD, which might have affected the rate of chronic GvHD [33,35,38,39]. Although severe GvHD will be unavoidable for some patients including MDS with poor prognostic factors [40,41], the balance between GvHD and GvL is a significant concern in RIST and we should seriously evaluate the type and tapering speed of immunosuppressive agents after RIST. Current findings suggested GvHD control might be improved simply by extending the duration of CyA administration. Additionally, we noticed that the clinical features of GvHD are different in RIST than in CIST, i.e. a syndrome compatible with acute GvHD occurs well after Day 100. Hence, the current grading system for GvHD, which was developed on the basis of experience in ablative settings, may not be an optimal tool for assessing GvHD after RIST. We observed a late onset of acute GvHD and an early onset of chronic GvHD, and therefore believe that a significant number of late-onset acute GvHD may have been judged as chronic GvHD in this study simply because the onset of GvHD was over 100 days after transplantation. Our results support the current proposition by Mielcarek and Storb concerning the abandonment of the traditional Day 100 cutoff for separating acute from chronic GvHD [35].

In this prospective study, we confirmed the short-term safety and efficacy of our RIST procedure for hematological malignancies in the elderly. Long-term follow-up of patients to evaluate disease control and the consequence of therapy is mandatory, and the development of optimal GvHD prophylaxis, with the use of novel assessment criteria, will be of primary importance for the wider application of the RIST procedure. RIST may also be beneficial in young patients, since organ damage, including infertility, might be milder and less frequent in RIST than in CIST, which should be confirmed by further prospective clinical trials. Although the number of patients studied was limited, the analysis of fludarabine pharmacokinetics has for the first time provided reliable information on the interaction of key drugs, and we found no evidence to suggest that synergic or specific toxicities were associated with increased exposure to the concomitant use of busulfan, or vice versa. This information should be useful in future studies in which different drugs are combined with fludarabine.

## Acknowledgments

This study was supported solely by a Health and Labour Sciences Research Grant from the Ministry of Health, Labour and Welfare, with fludarabine phosphate kindly supported by Nihon Schering K. K. We thank the medical and nursing staff of all of the transplant centers for their important contributions. We are indebted to the members of the

Japan Clinical Research Support Unit for data management. We are also grateful to the members of the University of Tokyo, the Statcom, and the EPS for the statistical analyses.

## Appendix

The following institutions contributed data to this study: National Cancer Center Hospital, Kanazawa University Hospital, Toranomon Hospital, Imamura Bun-in Hospital, Ehime Prefectural Central Hospital, Ishikawa Prefectural Central Hospital, Osaka City University Hospital, and Toyama Prefectural Central Hospital.

## References

1. Alyea EP, Kim HT, Ho V, et al. Comparative outcome of nonmyeloablative and myeloablative allogeneic hematopoietic cell transplantation for patients older than 50 years of age. *Blood* 2005;105:1810–1814.
2. Bertz H, Potthoff K, Finke J. Allogeneic stem-cell transplantation from related and unrelated donors in older patients with myeloid leukemia. *J Clin Oncol* 2003;21:1480–1484.
3. McSweeney PA, Niederwieser D, Shizuru JA, et al. Hematopoietic cell transplantation in older patients with hematologic malignancies: Replacing high-dose cytotoxic therapy with graft-versus-tumor effects. *Blood* 2001;97:3390–3400.
4. Shimoni A, Kroger N, Zabelina T, et al. Hematopoietic stem-cell transplantation from unrelated donors in elderly patients (age >55 years) with hematologic malignancies: Older age is no longer a contraindication when using reduced intensity conditioning. *Leukemia* 2005;19:7–12.
5. Shapira MY, Resnick IB, Bitan M, et al. Low transplant-related mortality with allogeneic stem cell transplantation in elderly patients. *Bone Marrow Transplant* 2004;34:155–159.
6. Slavin S, Nagler A, Naparstek E, et al. Nonmyeloablative stem cell transplantation and cell therapy as an alternative to conventional bone marrow transplantation with lethal cytoreduction for the treatment of malignant and nonmalignant hematologic diseases. *Blood* 1998;91:756–763.
7. Wong R, Giralt SA, Martin T, et al. Reduced-intensity conditioning for unrelated donor hematopoietic stem cell transplantation as treatment for myeloid malignancies in patients older than 55 years. *Blood* 2003;102:3052–3059.
8. Tsiangotis P, Bitan RO, Resnick IB, et al. A non-myeloablative conditioning regimen in allogeneic stem cell transplantation from related and unrelated donors in elderly patients. *Haematologica* 2006;91:852–855.
9. Giralt S, Estey E, Albitar M, et al. Engraftment of allogeneic hematopoietic progenitor cells with purine analog-containing chemotherapy: Harnessing graft-versus-leukemia without myeloablative therapy. *Blood* 1997;89:4531–4536.
10. Kroger N, Shimoni A, Zabelina T, et al. Reduced-toxicity conditioning with treosulfan, fludarabine and ATG as preparative regimen for allogeneic stem cell transplantation (alloSCT) in elderly patients with secondary acute myeloid leukemia (sAML) or myelodysplastic syndrome (MDS). *Bone Marrow Transplant* 2006;37:339–344.
11. Gupta V, Daly A, Lipton JH, et al. Nonmyeloablative stem cell transplantation for myelodysplastic syndrome or acute myeloid leukemia in patients 60 years or older. *Biol Blood Marrow Transplant* 2005;11:764–772.
12. Khouri IF, Keating M, Korbling M, et al. Transplant-free: Induction of graft-versus-malignancy using fludarabine-based nonablative chemotherapy and allogeneic blood progenitor-cell transplantation as treatment for lymphoid malignancies. *J Clin Oncol* 1998;16:2817–2824.
13. Niederwieser D, Maris M, Shizuru JA, et al. Low-dose total body irradiation (TBI) and fludarabine followed by hematopoietic cell transplantation (HCT) from HLA-matched or mismatched unrelated donors and postgrafting immunosuppression with cyclosporine and mycophenolate mofetil (MMF) can induce durable complete chimerism and sustained remissions in patients with hematological diseases. *Blood* 2003;101:1620–1629.
14. Childs R, Clave E, Contentin N, et al. Engraftment kinetics after nonmyeloablative allogeneic peripheral blood stem cell transplantation: Full donor T-cell chimerism precedes alloimmune responses. *Blood* 1999;94:3234–3241.
15. Baron F, Baker JE, Storb R, et al. Kinetics of engraftment in patients with hematologic malignancies given allogeneic hematopoietic cell transplantation after nonmyeloablative conditioning. *Blood* 2004;104:2254–2262.
16. Bornhauser M, Storer B, Slattery JT, et al. Conditioning with fludarabine and targeted busulfan for transplantation of allogeneic hematopoietic stem cells. *Blood* 2003;102:820–826.
17. Keil F, Prinz E, Moser K, et al. Rapid establishment of long-term culture-initiating cells of donor origin after nonmyeloablative allogeneic hematopoietic stem-cell transplantation, and significant prognostic impact of donor T-cell chimerism on stable engraftment and progression-free survival. *Transplantation* 2003;76:230–236.
18. Bearman SI, Appelbaum FR, Buckner CD, et al. Regimen-related toxicity in patients undergoing bone marrow transplantation. *J Clin Oncol* 1988;6:1562–1568.
19. Przepiorcka D, Weisdorf D, Martin P, et al. 1994 Consensus conference on acute GVHD grading. *Bone Marrow Transplant* 1995;15:825–828.

20. Shulman HM, Sullivan KM, Weiden PL, et al. Chronic graft-versus-host syndrome in man. A long-term clinicopathologic study of 20 Seattle patients. *Am J Med* 1980;69:204–217.
21. Slattery JT, Risler LJ. Therapeutic monitoring of busulfan in hematopoietic stem cell transplantation. *Ther Drug Monit* 1998;20:543–549.
22. Bornhauser M, Thiede C, Schuler U, et al. Dose-reduced conditioning for allogeneic blood stem cell transplantation: Durable engraftment without antithymocyte globulin. *Bone Marrow Transplant* 2000;26:119–125.
23. Schetelig J, Bornhauser M, Kiehl M, et al. Reduced-intensity conditioning with busulfan and fludarabine with or without antithymocyte globulin in HLA-identical sibling transplantation—A retrospective analysis. *Bone Marrow Transplant* 2004;33:483–490.
24. Maris MB, Niederwieser D, Sandmaier BM, et al. HLA-matched unrelated donor hematopoietic cell transplantation after nonmyeloablative conditioning for patients with hematologic malignancies. *Blood* 2003;102:2021–2030.
25. Hamaki T, Kami M, Kim SW, et al. Reduced-intensity stem cell transplantation from an HLA-identical sibling donor in patients with myeloid malignancies. *Bone Marrow Transplant* 2004;33:891–900.
26. Aoudjane M, Labopin M, Gorin NC, et al. Comparative outcome of reduced intensity and myeloablative conditioning regimen in HLA identical sibling allogeneic haematopoietic stem cell transplantation for patients older than 50 years of age with acute myeloblastic leukaemia: A retrospective survey from the Acute Leukemia Working Party (ALWP) of the European group for blood and marrow transplantation (EBMT). *Leukemia* 2005;19:2304–2312.
27. Schattenberg A, Schaap N, Preijers F, van der Maazen R, de Witte T. Outcome of T cell-depleted transplantation after conditioning with an intensified regimen in patients aged 50 years or more is comparable with that in younger patients. *Bone Marrow Transplant* 2000;26:17–22.
28. Du W, Dansey R, Abella EM, et al. Successful allogeneic bone marrow transplantation in selected patients over 50 years of age—A single institution's experience. *Bone Marrow Transplant* 1998;21:1043–1047.
29. Ringden O, Horowitz MM, Gale RP, et al. Outcome after allogeneic bone marrow transplant for leukemia in older adults. *JAMA* 1993;270:57–60.
30. Farag SS, Elder PJ, Marcucci G, et al. Radiation-free regimens result in similar outcomes of allogeneic hematopoietic progenitor cell transplantation in patients aged  $\geq 50$  years compared to younger adults with low-risk disease. *Bone Marrow Transplant* 2003;31:87–93.
31. Tauszig DC, Davies AJ, Cavenagh JD, et al. Durable remissions of myelodysplastic syndrome and acute myeloid leukemia after reduced-intensity allografting. *J Clin Oncol* 2003;21:3060–3065.
32. Mineishi S, Kanda Y, Saito T, et al. Impact of graft-versus-host disease in reduced-intensity stem cell transplantation (RIST) for patients with haematological malignancies. *Br J Haematol* 2003;121:296–303.
33. Mielcarek M, Martin PJ, Leisenring W, et al. Graft-versus-host disease after nonmyeloablative versus conventional hematopoietic stem cell transplantation. *Blood* 2003;102:756–762.
34. Cutler C, Giri S, Jeyapalan S, Paniagua D, Viswanathan A, Antin JH. Acute and chronic graft-versus-host disease after allogeneic peripheral-blood stem-cell and bone marrow transplantation: A meta-analysis. *J Clin Oncol* 2001;19:3685–3691.
35. Mielcarek M, Storb R. Graft-vs-host disease after non-myeloablative hematopoietic cell transplantation. *Leuk Lymphoma* 2005;46:1251–1260.
36. Nash RA, Pepe MS, Storb R, et al. Acute graft-versus-host disease: Analysis of risk factors after allogeneic marrow transplantation and prophylaxis with cyclosporine and methotrexate. *Blood* 1992;80:1838–1845.
37. Weisdorf D, Hakke R, Blazar B, et al. Risk factors for acute graft-versus-host disease in histocompatible donor bone marrow transplantation. *Transplantation* 1991;51:1197–1203.
38. Burroughs L, Mielcarek M, Leisenring W, et al. Extending postgrafting cyclosporine decreases the risk of severe graft-versus-host disease after nonmyeloablative hematopoietic cell transplantation. *Transplantation* 2006;81:818–825.
39. Ochs LA, Miller WJ, Filipovich AH, et al. Predictive factors for chronic graft-versus-host disease after histocompatible sibling donor bone marrow transplantation. *Bone Marrow Transplant* 1994;13:455–460.
40. Mohty M, Bay JO, Faucher C, et al. Graft-versus-host disease following allogeneic transplantation from HLA-identical sibling with antithymocyte globulin-based reduced-intensity preparative regimen. *Blood* 2003;102:470–476.
41. Castro-Malaspina H, Harris RE, Gajewski J, et al. Unrelated donor marrow transplantation for myelodysplastic syndromes: Outcome analysis in 510 transplants facilitated by the National Marrow Donor Program. *Blood* 2002;99:1943–1951.



# Comparable Antileukemia/Lymphoma Effects in Nonremission Patients Undergoing Allogeneic Hematopoietic Cell Transplantation with a Conventional Cytoablative or Reduced-Intensity Regimen

*Dai Maruyama, Takahiro Fukuda, Ruri Kato, Satoshi Yamasaki, Eiji Usui, Yuriko Morita-Hoshi, Sung-Won Kim, Shin-ichiro Mori, Yuji Heike, Atsushi Makimoto, Kinuko Tajima, Ryuji Tanosaki, Kensei Tobinai, Yoichi Takaue*

Hematology and Stem Cell Transplantation Division, National Cancer Center Hospital, Tokyo, Japan

Correspondence and reprint requests: Takahiro Fukuda, MD, PhD, Hematology and Stem Cell Transplantation Division, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo, 104-0045, Japan (e-mail: tafukuda@ncc.go.jp).

Received February 1, 2007; accepted April 12, 2007

## ABSTRACT

To evaluate the potential of allogeneic hematopoietic cell transplantation (HCT) with a reduced-intensity conditioning regimen (RIST) for the treatment of patients with hematologic malignancies not in remission, we retrospectively reviewed the medical records of 132 patients (89 leukemia or myelodysplastic syndrome, 40 malignant lymphoma, and 3 others) who received conventional myeloablative HCT (CST,  $n = 52$ ) or RIST ( $n = 80$ ). The median age of the RIST group was significantly higher than that of the CST group (53 years versus 40 years,  $P < .01$ ). The RIST group also included a higher proportion of patients with an HCT-specific comorbidity index (HCT-CI) of 1 or more than the CST group (65% versus 37%,  $P = .03$ ). The probabilities of achieving complete remission and the incidences of grades II-IV and III-IV acute graft-versus-host disease (aGVHD) in the CST and RIST groups were, respectively, 77% and 64%, 50% and 50%, and 23% and 28%, with no significant differences. Similarly, there was no difference in the 2-year probabilities of nonrelapse mortality (NRM, 36% and 38%), progressive disease or relapse (PD 51% and 49%), overall survival (OS, 31% and 38%), and progression-free survival (PFS, 28% and 29%). Multivariate analyses revealed that a higher HCT-CI score and transplant from donors other than HLA-matched relatives were associated with increased risks of NRM and poor OS, and patients who received chemotherapy within 2 months before HCT were associated with increased risks of PD, poor OS, and PFS after transplantation. After adjusting for these variables, the risks of NRM, PD, OS, and PFS in the RIST group were not significantly different from those in the CST group. In conclusion, these results suggest that the antileukemia/lymphoma effect associated with RIST is comparable to that associated with CST. RIST appears to be feasible for the treatment of hematologic malignancies not in remission.

© 2007 American Society for Blood and Marrow Transplantation

## KEY WORDS

Transplantation • Leukemia • Lymphoma

## INTRODUCTION

Allogeneic hematopoietic cell transplantation (HCT) has the potential to achieve long-term cure of hematologic malignancies by pretransplant conditioning and a graft-versus-leukemia/lymphoma (GVL) effect. It has been well established that the disease status

at the time of transplantation is the most important prognostic factor, and the rates of relapse and nonrelapse mortality (NRM) significantly increase in patients with hematologic malignancies who were not in remission. Therefore, conventional stem cell transplantation (CST) using a myeloablative conditioning

regimen has been universally used in the hope of maximally reducing the tumor burden before HCT in patients not in remission. However, CST may not be an option for many patients because of their older age or associated comorbidities. Alternatively, over the past few years, nonmyeloablative and reduced-intensity conditioning stem cell transplantation (RIST) have been offered to these patients undergoing HCT, on the assumption that RIST would be better tolerated [1-4].

There have been several reports that the outcome of older patients who underwent RIST while in remission was comparable to that of patients who received CST [5-9], which suggests that the GVL effect associated with RIST might be adequate for controlling chemosensitive or slowly progressing disease. On the other hand, it still remains controversial whether RIST is feasible for patients not in remission, although small pilot studies have shown that RIST was unsuccessful for advanced hematologic malignancies [3,10-14]. To address this issue, we retrospectively analyzed 132 patients who were not in remission at the time of CST or RIST.

## PATIENTS AND METHODS

### Study Patients

We retrospectively reviewed the medical records of 132 patients with various hematologic malignancies who underwent allogeneic HCT (CST,  $n = 52$ ; RIST,  $n = 80$ ) while not in remission at our institution from January 2000 to December 2004. Patients with chronic myelogenous leukemia (CML) in the chronic phase, myelodysplastic syndrome (MDS)-refractory anemia, and those with lymphoma in partial remission (PR) were not included because the response to treatment and the outcome of these patients is generally considered to be similar to those in patients who are in complete remission (CR). Bone marrow or granulocyte colony-stimulating factor (G-CSF)-mobilized peripheral blood stem cells (PBSC) were harvested from donors according to protocols approved by the guidelines of the Japan Marrow Donor Program, the Japanese Society for Hematopoietic Cell Transplantation, and the Japanese Society of Blood Transfusion. Informed consent was obtained according to the Declaration of Helsinki.

### Transplantation Procedures

The conditioning regimens used in CST included the combination of cyclophosphamide (CY; 60 mg/kg i.v. daily for 2 days) and fractionated total body irradiation (TBI; 12 Gy in 6 fractions over 3 days) in 34 patients, CY and oral busulfan (BU; 16 mg/kg divided over 4 days) in 13 patients, and other combinations in 5 patients (Table 1). Targeted dose adjustment of BU

was not performed. Patients who underwent RIST were older than 50 years of age or those who had comorbidities or prior transplantation. The conditioning regimens for RIST consisted of fludarabine (30 mg/m<sup>2</sup> i.v. daily for 6 days) or cladribine (0.11 mg/kg i.v. daily for 6 days) plus 8 mg/kg of oral BU [15] with ( $n = 27$ ) or without ( $n = 53$ ) 4 Gy TBI. In Japan, only bone marrow is permitted as a stem cell source in transplantation from an unrelated healthy volunteer donor. In the setting of nonmyeloablative stem cell transplantation from an unrelated donor, the sustained engraftment rate has been reported to be lower for recipients of bone marrow than for those given PBSC [13]. Therefore, low-dose TBI was also added to the conditioning regimen for RIST from an unrelated donor to facilitate engraftment.

Day 0 was defined as the day of stem cell infusion. G-CSF was administered after transplantation in all patients until neutrophil engraftment. Most patients who underwent CST were given cyclosporine (CSP) with methotrexate (MTX) [16], and all patients who underwent RIST were given CSP with or without MTX for graft-versus-host disease (GVHD) prophylaxis (Table 1). GVHD was treated with 1 to 2 mg/kg/day prednisolone equivalents, resumption of full-dose CSP administration if applicable, or both. Initial doses of corticosteroids and tapering schedules of immunosuppressive medications were modified at the discretion of the attending physicians according to the presence or absence of malignant cells and the severity of GVHD. Treatment for relapse after transplantation was left to the discretion of the attending physicians.

All patients received ciprofloxacin (200 mg orally 3 times daily) for bacterial prophylaxis until neutrophil engraftment. Fluconazole (100 mg once daily) was administered for fungal prophylaxis. Patients who had positive serologic test results for herpes simplex virus or varicella zoster virus received prophylactic low-dose acyclovir until the cessation of immunosuppressive agents [17]. Prophylaxis against *Pneumocystis jiroveci* infection was provided with trimethoprim-sulfamethoxazole from the first day of conditioning to day -3 of transplantation, and from day 28 until day 180 or the cessation of immunosuppressive agents. Patients were monitored with weekly cytomegalovirus (CMV) pp65 antigenemia testing, and positive antigenemia was treated with ganciclovir as described previously [18,19].

### Definitions

Chemotherapy within 2 months before HCT was defined as chemotherapy to control the disease except for rituximab alone for lymphoma and imatinib mesylate alone for CML. Pretransplantation comorbidities were determined by the HCT-specific comorbidity index (HCT-CI) [20] with a minor modification [21]. Neutrophil engraftment was defined as the first