

Figure 9. *Wfs1* mRNA signals and protein immunoreactivity in the normal mouse brain.




Figure 10. Wfs1 protein immunoreactivity in the normal mouse suprachiasmatic nucleus (SCN), optic chiasm (OX), and optic tract (OT).

TABLE 1. Primary Antibodies Used in This Study

Antibody	Manufacturer	Catalog No.	Lot No.	Antibody isotype	Clone	Dilution	Antigen	Specificity
Mouse Wfs1	This Study			Rabbit serum		1:5000- 1:10000	Amino acids 1-179 representing N-terminus Wfs1 protein of mouse origin	A single band of ~100 kDa in extracts from the mouse brain and optic nerve. ~100 kDa and ~70 kDa bands in extracts from the mouse retina. Antigen block of immunolabeling. Figs. 1 and 2D
Calbindin-D-28K	Sigma-Aldrich, St. Louis, MO	C9848	113K4867	Mouse IgG ₁	Clone CB-955	1:500	Purified bovine kidney calbindin-D-28K	A single 28-kDa band*1. Staining pattern identical to published data*2
Gou ₁	Chemicon, Temecula, CA	MAB3073	24040782	Mouse IgG ₁	Clone 2A	1:250	Purified bovine brain Gou ₁	A single protein band of 39-42 kDa*1. Staining pattern identical to published data*2
Choline Acetyltransferase	Chemicon	AB144P	24030848	Goat serum		1:100	Choline acetyltransferase from human placenta	Staining pattern identical to published data*3
Tubulin, β III isoform	Chemicon	MAB1637	25010305	Mouse IgG ₁	Clone TU-20	1:50	A synthetic peptide corresponding to amino acids 443-450 (ESESQGPK) of human class III β -tubulin conjugated to keyhole limpet hemocyanin (KLH), a carrier for haptens	A single protein band of 50 kDa in extracts from the porcine brain*4
Bm-3a	Santa Cruz Biotechnology, Santa Cruz, CA	sc-8429	L1106	Mouse IgG _{2b}	Clone 14A6	1:50	Amino acids 1-109 representing N-terminus Bm-3a protein of mouse origin	Staining pattern identical to published data*5
Glutamine Synthetase	BD Biosciences Pharmingen, San Diego, CA	610517	01915	Mouse IgG _{2a}	Clone 6	1:500	Sheep glutamine synthetase (1-373: Full Length)	A single protein band of 45 kDa in extracts from the rat brain*6
Glial Fibrillary Acidic Protein (GFAP)	Chemicon	AB5804	23080114	Rabbit serum		1:1000	Purified bovine GFAP	A single protein band of 51 kDa in extracts from the mouse brain*7
GFAP	Chemicon	MAB360	25040157	Mouse IgG ₁	Clone GA5	1:1000	Purified GFAP from porcine spinal cord	A single protein band of 51 kDa in extracts from a human glioma cell line*8. Staining pattern (Fig. 8B) identical to that of the anti-GFAP rabbit serum (Fig. 6A)
Oligodendrocytes [RIP]	Chemicon	MAB1580	25040064	Mouse IgG ₁	Clone NS-1	1:100000	Rat olfactory bulb	Staining pattern identical to published data*9
Mouse CD11b (Mac-1 α chain)	BD Biosciences Pharmingen	550282	01651	Rat (DA) IgG _{2b,1}	Clone M1/70	1:50	C57BL/10 mouse splenic T cells and concanavalin A-activated C57BL/10 splenocytes	Staining pattern identical to published data*10

*1: Manufacturer's technical information; Gargini et al., 2007

*2: Haverkamp and Wässle, 2000

*3: Jeon et al., 1998; Haverkamp and Wässle, 2000; Kang et al., 2004

*4: Manufacturer's technical information; Dráberová et al., 1998

*5: Xiang et al., 1995

*6: Manufacturer's technical information

*7: Jalil et al., 2005

*8: Manufacturer's technical information; Debus et al., 1983

*9: Saari et al., 1997; Morcos and Chan-Ling, 2000

*10: Retina: Dräger, 1983; Optic nerve: Reichert and Rotshenker, 1996; Neufeld, 1999.

TABLE 2. Distribution of Wfs1 Immunoreactivity in the Normal Mouse Retina

Retinal cells	
Photoreceptor	+
Outer plexiform layer	+
Horizontal cells	+
Bipolar cells	+
Amacrine cells	+
Inner plexiform layer	+
Ganglion cells	+
Displaced amacrine cells	+
Müller cells	+
Astrocytes	-
Microglia	-

+, present; -, not present.

TABLE 3. Distribution of Wfs1 Immunoreactivity in the Normal Mouse Optic Nerve

Optic nerve cells	
Astrocytes	+
Oligodendrocytes	-
Microglia	-

+, present; -, not present.

TABLE 4. Distribution of Wfs1 Immunoreactivity in the Normal Mouse Vision-Related Brain Structures

Vision-related brain structures	
Optic chiasm	-
Optic tract	-
Suprachiasmatic nucleus	+
Lateral geniculate nucleus	-
Pretectum	-
Superior colliculus	+
Medial terminal nucleus	-
Primary visual cortex	+
Secondary visual cortex	+

+, present; -, not present.

Identification of Glypican3 as a novel GLUT4 binding protein

Akihiko Taguchi^a, Masahiro Emoto^a, Shigeru Okuya^{a*}, Naofumi Fukuda^a, Yoshitaka Nakamori^a, Mutsuko Miyazaki^a, Sachiko Miyamoto^a, Katsuya Tanabe^a, Hiroyuki Aburatani^b, Yoshitomo Oka^c, Yukio Tanizawa^a

^aDivision of Endocrinology, Metabolism, Hematological Sciences and Therapeutics, Department of Bio-Signal Analysis Yamaguchi University Graduate School of Medicine

^bGenome Science Division, Research Center for Advanced Science and Technology, University of Tokyo, Tokyo, Japan

^cDivision of Molecular Metabolism and Diabetes, Tohoku University Graduate School of Medicine, Sendai 980-8575, Japan

***Corresponding author.** Fax: +81-836-22-2342.

E-mail address: okuya@yamaguchi-u.ac.jp (S. Okuya)

Abstract

Insulin stimulates glucose uptake in fat and muscle primarily by stimulating the translocation of vesicles containing facilitative glucose transporters, GLUT4, from intracellular compartments to the plasma membrane. Although cell surface externalization of GLUT4 is critical for glucose transport, the mechanism regulating cell surface GLUT4 remains unknown. Using a yeast two-hybrid screening system, we have screened GLUT4-binding proteins, and identified a novel glycosyl phosphatidyl inositol (GPI)-linked proteoglycan, Glypican3 (GPC3). We confirmed their interaction using immunoprecipitation and a GST pull-down assay. We also revealed that GPC3 and GLUT4 to co-localized at the plasma membrane, using immunofluorescent microscopy. Furthermore, we observed that glucose uptake in GPC3-overexpressing adipocytes was increased by 30% as compared to control cells. These findings suggest that GPC3 may play roles in glucose transport through GLUT4.

Keywords: GLUT4; 3T3-L1 adipocytes; GPC3

Insulin stimulation of glucose uptake into skeletal muscle and adipose tissue is achieved via the translocation of intracellular-sequestered GLUT4 protein to the cell surface membrane [1; 2]. On the plasma membrane, GLUT4 proteins, responding to insulin stimulation, remain externalized for a certain period of time and facilitate glucose transport. Although cell surface externalization of GLUT4 is critical for glucose transport, the mechanism regulating cell surface GLUT4 remains largely unknown. We speculated that it would require a protein capable of interacting with the glucose transporter.

Over the past decade, several GLUT4-binding proteins, such as mUbc9 [3], TUG [4], DAXX [5], L-3-hydroxyacyl-CoA dehydrogenase [6] and carboxyl esterase [7], have been identified using the C-terminus region of GLUT4 as bait in either a two-hybrid system or an immobilized GST fusion protein pull-down experiment. For example, mUbc9 was demonstrated to regulate transporter degradation [3], whereas TUG was shown to modulate GLUT4 distribution [4]. However, the functions of other GLUT4 binding proteins are not fully understood. We speculated that a change in the three-dimensional structure of the GLUT4 partial sequence has made it difficult to identify and analyze functional GLUT4 binding proteins. Thus, we used full length GLUT4 for screening, and thereby identified GPC3 as a GLUT4 binding protein.

GPC3 is one of the heparan sulfate proteoglycans that are anchored to the cell membrane by a glycosyl-phosphatidylinositol protein [8]. This family of proteins was shown to be related to morphogenesis and GPC3 was originally reported to be a negative regulator of cell proliferation as well as the progression

Taguchi et al. Identification of Glypican3 as a novel GLUT4 binding protein.
Biochem Biophys Res Comm. In press

of malignant tumors [9; 10; 11; 12; 13]. Herein, we report GPC3 as a newly identified GLUT4 binding protein. This is the first report describing a protein that acts directly on GLUT4 molecules at the plasma membrane.

MATERIALS AND METHODS

Antibodies. Mouse monoclonal GPC3 antibody was provided by Dr. Hiroyuki Aburatani (University of Tokyo, Japan). The following antibodies were used: anti-GLUT4 rabbit and goat, anti-GST (Santa Cruz Biotechnology, CA); anti-FLAG (M2) (Sigma); anti-Myc (9E10) (Convance, CA) and fluorescent-conjugated and Horseradish peroxidase-conjugated secondary antibodies (Jackson Immuno-Research Laboratories).

Constructs. Mouse GPC3 cDNA was purchased from Open Biosystems (Huntsville, AL). Wild-type GPC3 was subcloned into a pGEX-6P1 (GE Healthcare Biosciences) vector. FLAG-tagged GPC3 and 4×Myc-tagged GLUT4-eGFP was subcloned into a pcDNA3 (Invitrogen) vector. All chemically synthesized and PCR-derived DNA sequences were verified by DNA sequencing.

Preparation of recombinant adenovirus vectors. Recombinant adenovirus encoding eGFP or FLAG-tagged GPC3 was constructed using the AdEasy adenovirus vector system according to the manufacturer's instructions (Stratagene). All amplified viruses were stored at -80°C. 3T3-L1 adipocytes were infected with recombinant adenovirus vectors encoding eGFP and FLAG-tagged GPC3 at a multiplicity of infection (m.o.i.) of 50.

Taguchi et al. Identification of Glypican3 as a novel GLUT4 binding protein.
Biochem Biophys Res Comm. In press

Cell culture. 3T3-L1 fibroblasts were grown in DMEM with 10% fetal bovine serum (FBS) at 37°C. The cells (3-5 days post-confluent) were differentiated into adipocytes by incubation in the same DMEM containing 0.5 mM isobutylmethylxanthine, 0.25 µM dexamethasone, and 4 µg/ml insulin for 3 days, and then grown in DMEM with 10% FBS for an additional 3-6 days. Human hepatocellular carcinoma cell line HepG2 cells were cultured in Eagle's Minimum Essential Medium (EMEM) with 10% FBS in at 37°C.

Primary culture of mouse hepatocytes. Mouse hepatocytes were prepared from 5-month male C57BL/6J mice as described previously [14]. Isolated cells were seeded onto coverslips and allowed to recover for 24h.

Yeast two-hybrid screening. The MATCHMAKER LexA Two-hybrid System (Clontech) was used for identification of GLUT4 binding proteins. As bait for screening, the vector pLexA-GLUT4 expressing a fusion protein composed of full-length rat GLUT4 linked to the DNA-binding domain was constructed. A rat adipocyte cDNA library already cloned into the pB42AD vector was obtained from Origene Technologies (Rockville, MD). Approximately 300 colonies showed activation of the yeast reporter gene, and 15 colonies showing dependence on the LexA-GLUT4 fusion protein for activation of the reporter gene were selected. Plasmids from positive clones were subsequently isolated from the yeast, transferred to *E. coli*, and sequenced. Full-length cDNA was obtained by 5'RACE (Rapid Amplification of cDNA End) using a kit (Clontech) and the GenBank/NCBI databases were screened for similar sequences using BLAST Search.

In vitro GST pull-down assay. GST fusion proteins of full-length GPC3 and GST alone were purified according to the manufacturer's instructions. GLUT4 protein was generated from Myc-tagged GLUT4-eGFP (four-Myc epitope-tags in the first exofacial loop and eGFP in the C-terminus) transfected 293 cells and further purified using anti-Myc antibody. These GST fusion proteins and purified 4×Myc-tagged GLUT4-eGFP were mixed in PBS and incubated at 4°C for 4h. The proteins were then pulled down with glutathione-sepharose beads (GE Healthcare Biosciences).

Immunoprecipitation and immunoblotting. Cells were lysed in lysis buffer [20 mM HEPES (pH 7.2), 100 mM NaCl, 1mM EDTA, 25 mM NaF, 1 mM sodium vanadate, 1 mM benzamidine, 5 µg/ml leupeptin, 5 µg/ml aprotinin, 1 mM PMSF, 1mM DTT] and the protein concentration was measured with BCA protein assay reagent (Pierce, Rockford, IL). For immunoprecipitation, the cell lysate was pre-incubated with protein-G Sepharose at 4°C for 30 min to remove nonspecific bound proteins. Then, samples were incubated with primary antibody at 4°C for 8-12h followed by incubation with protein-G Sepharose. Lysates and immunoprecipitates were resolved by SDS-PAGE and transferred to a polyvinylidene difluoride (PVDF) membrane (GE Healthcare Biosciences). The membranes were incubated with appropriate antibodies.

Immunofluorescence microscopy. 3T3-L1 adipocytes, HepG2 cells, and primary hepatocytes were seeded onto coverslips and allowed to recover for 24-48h. 3T3-L1 adipocytes were serum-starved for 4h in DMEM, followed by incubation with or without 100 nM of insulin for 15 min at 37°C. Then, all of the

cells were fixed with 3.7% formaldehyde in PBS and permeabilized with buffer A (0.5% Triton X-100, 1% FBS in PBS) for 15min, and finally incubated for 2h with primary antibodies at room temperature. The cells were washed and incubated with an appropriate secondary antibody, or rhodamine conjugated wheat germ agglutinin (molecular probes, inc.) as a counter staining of cell membrane and Golgi system. The coverslips were washed thoroughly and mounted on glass slides. Immunostained cells were observed at room temperature with a laser-scanning confocal microscope (LSM5 PASCAL; Carl Zeiss Inc).

Plasma membrane sheet assay. Cell surface protein was assayed using plasma membrane lawns as described previously [15]. The cells were subsequently swollen using hypotonic buffer and sonicated to generate a lawn of plasma membrane fragments. The membranes were immunostained with anti-GPC3 and anti-GLUT4 antibody. These cells were observed by laser confocal microscopy.

2-deoxy-glucose uptake. Differentiated adipocytes were prepared in 24-well plates. Cells were infected with the recombinant adenoviruses. Two days thereafter, the cells were serum-starved for 2 h at 37°C in Krebs-Ringer phosphate buffer (130 mM NaCl, 5 mM KCl, 1.3 mM CaCl₂, 1.3 mM MgSO₄, and 10 mM Na₂HPO₄, pH 7.4). The cells were then stimulated with or without 100 nM of insulin for 15 min, and deoxy-glucose uptake was determined by 2-deoxy-D-[2,6 ³H] glucose incorporation.

RESULTS AND DISCUSSION

Identification of GLUT4 binding proteins

Regulation of glucose uptake in muscle and adipose tissues by insulin is important for proper maintenance of blood glucose. This hormone stimulates translocation of the GLUT4 glucose transporter from the intracellular membrane to the cell surface. After translocation to the plasma membrane, GLUT4 remains on the cell surface temporarily and facilitates glucose transport. However, the mechanism regulating cell surface GLUT4 is still largely unknown. We hypothesized that it requires protein-protein interactions at the plasma membrane. We used the yeast two-hybrid screening system to identify proteins that interact physically with full length GLUT4. As bait, we used full-length rat GLUT4 cDNA and, as prey, a rat adipose tissue cDNA library. After the first screening, we obtained more than 300 colonies. After a second screening, 15 colonies remained positive. Ultimately, nine colonies were left. Full length cDNAs were obtained by 5'RACE. One of them was 100% identical to GPC3 [16], which was originally cloned as OCI-5, a GPI anchored membrane protein [17]. Since we also obtained mUbc9 [3], which was previously identified using the same method, our experimental procedure was thought to have worked correctly. The GPC3 sequence obtained with the yeast two-hybrid system was comprised of the residues from 521 to 597. In order to confirm the interaction between GLUT4 and GPC3, we employed the yeast two-hybrid system again; using full-length GLUT4 cDNA and full-length GPC3 cDNA as the bait and prey, respectively. We

confirmed the LacZ signal indicating a direct interaction between GPC3 and GLUT4 (Fig. 1A). Next, we determined GPC3 protein expression using Western blot analysis. As shown in Fig. 1B, we detected high molecular weight form corresponding to the glycosylated GPC3 and bands corresponding to the non-glycosylated GPC3 core protein and its cleavage product of in 3T3-L1 adipocytes as in HepG2 cells. These data were consistent with the previous reports [11; 13].

Intracellular localization of GPC3

The intracellular localization of GPC3 was observed in adipocytes by laser confocal microscopy. First, we determined GPC3 protein by immunofluorescent microscopy using our specific antibody against GPC3 in HepG2 cells and primary hepatocytes as a positive or negative control, respectively (Fig. 2A). As shown in Fig. 2B, most of the GPC3 was at the plasma membrane, as reported for other cell lines [17]. As expected, we observed co-localization of GLUT4 and GPC3 after insulin stimulation (Fig 2B). Next, in order to confirm plasma membrane integrity, we prepared plasma membrane sheets from 3T3-L1 adipocytes, stained with anti-GPC3 and anti-GLUT4 antibodies, and then observed GPC3 expression by confocal microscopy (Fig. 2C). We observed that GLUT4 translocated to the plasma membrane in response to insulin stimulation. GPC3 localized at the plasma membrane on under both conditions, and merged more clearly with GLUT4 after insulin stimulation.

The results shown in Fig. 1A and Fig. 2 suggest that GPC3 interacted with GLUT4 at the plasma membrane.

Interaction between GLUT4 and GPC3

In order to reconfirm the interaction between GPC3 and GLUT4, we conducted two other experiments; immunoprecipitation and a GST pull-down assay. First, we examined endogenous protein-protein interactions employing immunoprecipitation experiments. We prepared the total lysates of insulin-treated 3T3-L1 adipocytes. These lysates were immunoprecipitated with anti-GLUT4 antibody, and blotted with anti-GPC3 antibody. In adipocytes, we detected the interaction between GLUT4 and non-glycosylated GPC3 core protein (Fig 3A). We also confirmed these interactions by immunoprecipitation with anti-GPC3 antibody (Fig. 3B). Second, we assessed the direct interaction between GLUT4 and GPC3 using the GST pull-down assay. We generated GST-GPC3 protein and GST alone using a bacterial system, and then purified these proteins with glutathione sepharose beads. GLUT4 protein was purified from 293 cells transfected with 4×Myc-tagged-GLUT4-eGFP. These fusion proteins were mixed in PBS, and incubated for several hours. The proteins were then pulled down with glutathione sepharose beads. We confirmed the GLUT4 signal in the GST-GPC3 precipitate lane, indicating a direct GLUT4 and GPC3 interaction in vitro (Fig. 3C).

Overexpression of GPC3 increased insulin stimulated glucose uptake

Since GPC3 was proved to GLUT4 at the plasma membrane, we examined whether GPC3 might regulate glucose uptake through GLUT4. We measured

basal and insulin stimulated glucose uptake in 3T3-L1 adipocytes which transiently overexpressed GPC3 following transfection with adenovirus vectors. Overexpression of GPC3 increased insulin induced glucose uptake (Fig. 4). These data suggested that GPC3 overexpression to enhance insulin-stimulated glucose uptake via cell surface GLUT4.

Simpson-Golabi-Behmel syndrome (SGBS) is a rare, complex congenital syndrome with affected individuals having loss of function mutations in the GPC3 gene. Although many researchers reported that the patients have general feature of overgrowth and cancer development, little is known about the clinical characteristics on glucose metabolism [18; 19; 20; 21]. It was reported that some infants had hypoglycemia due to hyperplasia of islets of Langerhans rather than modification of insulin action in peripheral tissues [22]. More detailed analysis for the peripheral insulin action, especially on the adipose tissue would clarify the physiological role of GPC3 on glucose metabolism.

In conclusion, we identified a novel GLUT4 binding protein GPC3, using a yeast two-hybrid screening system. We also demonstrated GPC3 and GLUT4 to co-localized at the plasma membrane in response to insulin. Furthermore, overexpression of GPC3 increased insulin stimulated glucose uptake in cultured adipocytes. GPC3 may play a role in insulin stimulated glucose uptake.

ACKNOWLEDGEMENTS

We are also very grateful to Dr. Teruo Nishida for generously supporting us in the using a laser confocal microscope experiments. The authors thank Y Kora

Taguchi et al. Identification of Glypican3 as a novel GLUT4 binding protein.
Biochem Biophys Res Comm. In press

and M Kaneko for technical assistance. This study was supported in part by Grants-in Aid for Scientific Research (17590934 to M Emoto, 15590939 and 17590935 to S Okuya) from the Ministry of Education, Culture, Sports, Science and Technology of Japan and a grant from Takeda Science Foundation to S Okuya.

REFERENCES

- [1] J.C. Hou, and J.E. Pessin, Ins (endocytosis) and outs (exocytosis) of GLUT4 trafficking. *Curr Opin Cell Biol* 19 (2007) 466-73.
- [2] S. Huang, and M.P. Czech, The GLUT4 glucose transporter. *Cell Metab* 5 (2007) 237-52.
- [3] F. Giorgino, O. de Robertis, L. Laviola, C. Montrone, S. Perrini, K.C. McCowen, and R.J. Smith, The sentrin-conjugating enzyme mUbc9 interacts with GLUT4 and GLUT1 glucose transporters and regulates transporter levels in skeletal muscle cells. *Proc Natl Acad Sci U S A* 97 (2000) 1125-30.
- [4] J.S. Bogan, N. Hendon, A.E. McKee, T.S. Tsao, and H.F. Lodish, Functional cloning of TUG as a regulator of GLUT4 glucose transporter trafficking. *Nature* 425 (2003) 727-33.
- [5] V.S. Lalioti, S. Vergarajauregui, D. Pulido, and I.V. Sandoval, The insulin-sensitive glucose transporter, GLUT4, interacts physically with Daxx. Two proteins with capacity to bind Ubc9 and conjugated to SUMO1. *J Biol Chem* 277 (2002) 19783-91.
- [6] Y. Shi, S.J. Samuel, W. Lee, C. Yu, W. Zhang, M. Lachaal, and C.Y. Jung, Cloning of an L-3-hydroxyacyl-CoA dehydrogenase that interacts with the GLUT4 C-terminus. *Arch Biochem Biophys* 363 (1999) 323-32.
- [7] W. Lee, J. Ryu, J. Hah, T. Tsujita, and C.Y. Jung, Association of carboxyl esterase with facilitative glucose transporter isoform 4 (GLUT4) intracellular compartments in rat adipocytes and its possible role in insulin-induced GLUT4 recruitment. *J Biol Chem* 275 (2000) 10041-6.
- [8] J. Filmus, Glypicans in growth control and cancer. *Glycobiology* 11 (2001) 19R-23R.
- [9] D.F. Cano-Gauci, H.H. Song, H. Yang, C. McKerlie, B. Choo, W. Shi, R. Pullano, T.D. Piscione, S. Grisaru, S. Soon, L. Sedlackova, A.K. Tanswell, T.W. Mak, H. Yeger, G.A. Lockwood, N.D. Rosenblum, and J. Filmus, Glypican-3-deficient mice exhibit developmental overgrowth and some of the abnormalities typical of Simpson-Golabi-Behmel syndrome. *J Cell Biol* 146 (1999) 255-64.
- [10] H. Lin, R. Huber, D. Schlessinger, and P.J. Morin, Frequent silencing of the GPC3 gene in ovarian cancer cell lines. *Cancer Res* 59 (1999) 807-10.
- [11] Y. Midorikawa, S. Ishikawa, H. Iwanari, T. Imamura, H. Sakamoto, K. Miyazono, T. Kodama, M. Makuuchi, and H. Aburatani, Glypican-3, overexpressed in hepatocellular carcinoma, modulates FGF2 and BMP-7 signaling. *Int J Cancer* 103 (2003) 455-65.
- [12] H.H. Song, W. Shi, Y.Y. Xiang, and J. Filmus, The loss of glypican-3 induces alterations in Wnt signaling. *J Biol Chem* 280 (2005) 2116-25.
- [13] M.I. Capurro, W. Shi, S. Sandal, and J. Filmus, Processing by convertases is not required for glypican-3-induced stimulation of hepatocellular carcinoma growth. *J Biol Chem* 280 (2005) 41201-6.
- [14] J.D. Horton, H. Shimano, R.L. Hamilton, M.S. Brown, and J.L. Goldstein, Disruption of LDL receptor gene in transgenic SREBP-1a mice unmask hyperlipidemia resulting from production of lipid-rich VLDL. *J Clin Invest* 103 (1999) 1067-76.
- [15] M. Emoto, S.E. Langille, and M.P. Czech, A role for kinesin in insulin-stimulated GLUT4 glucose transporter translocation in 3T3-L1 adipocytes. *J Biol Chem* 276 (2001) 10677-82.
- [16] J. Filmus, J.G. Church, and R.N. Buick, Isolation of a cDNA corresponding to a developmentally regulated transcript in rat intestine. *Mol Cell Biol* 8 (1988) 4243-9.

Taguchi et al. Identification of Glypican3 as a novel GLUT4 binding protein.
Biochem Biophys Res Comm. In press

- [17] K. Watanabe, H. Yamada, and Y. Yamaguchi, K-glypican: a novel GPI-anchored heparan sulfate proteoglycan that is highly expressed in developing brain and kidney. *J Cell Biol* 130 (1995) 1207-18.
- [18] G. Pilia, R.M. Hughes-Benzie, A. MacKenzie, P. Baybayan, E.Y. Chen, R. Huber, G. Neri, A. Cao, A. Forabosco, and D. Schlessinger, Mutations in GPC3, a glypican gene, cause the Simpson-Golabi-Behmel overgrowth syndrome. *Nat Genet* 12 (1996) 241-7.
- [19] N. Okamoto, M. Yagi, K. Imura, and Y. Wada, A clinical and molecular study of a patient with Simpson-Golabi-Behmel syndrome. *J Hum Genet* 44 (1999) 327-9.
- [20] S. Lindsay, M. Ireland, O. O'Brien, J. Clayton-Smith, J.A. Hurst, J. Mann, T. Cole, J. Sampson, S. Slaney, D. Schlessinger, J. Burn, and G. Pilia, Large scale deletions in the GPC3 gene may account for a minority of cases of Simpson-Golabi-Behmel syndrome. *J Med Genet* 34 (1997) 480-3.
- [21] J.Y. Xuan, A. Besner, M. Ireland, R.M. Hughes-Benzie, and A.E. MacKenzie, Mapping of Simpson-Golabi-Behmel syndrome to Xq25-q27. *Hum Mol Genet* 3 (1994) 133-7.
- [22] M.R. DeBaun, J. Ess, and S. Saunders, Simpson Golabi Behmel syndrome: progress toward understanding the molecular basis for overgrowth, malformation, and cancer predisposition. *Mol Genet Metab* 72 (2001) 279-86.