

図2 生理活性蛋白質の高分子バイオコンジュゲーションの特徴  
 生体に投与された生理活性蛋白質は、各種プロテアーゼにより分解されてしまうだけでなく、分子量数万以下のものは速やかに腎排泄されてしまうため、一般にその生体内安定性・滞留性はあまりにも乏しい。一方で高分子バイオコンジュゲーションは、高分子化による腎排泄速度の現象をもたらすだけでなく、蛋白質分子の表面を覆う修飾高分子により立体配位的にプロテアーゼからの攻撃をブロックするために、体内滞留時間の延長をもたらす。しかし、高分子レセプターとの結合を要するサイトカインなどのバイオコンジュゲーションは、同時に分子量増大に伴う組織移行性の低下や、蛋白質に結合した修飾高分子に由来する立体障害のために、レセプター親和性の低下(比活性の低下)を招いてしまう。




図3 最適バイオコンジュゲーションによるIL-6の血小板産生促進作用の選択増強  
 最適条件下でバイオコンジュゲーションしたPEG化IL-6のin vivoにおける血小板産生促進作用は、未修飾IL-6の500倍にも増強されていた。また未修飾IL-6では、発熱や非特異的抗体産生の誘導といった強い副作用が観察されたが、PEG化IL-6投与群ではこれらの副作用はほとんど認めなかった。

これらリジン残基への高分子導入により必然的に著しい比活性低下を招いてしまう。また、ランダムに修飾高分子が導入されるため、得られたバイオコンジュゲート体は蛋白質のさまざまな部位に種々個数の修飾高分子が結合した分子的に不均一な混合物となる。したがって、プロテオーム創薬を推進するためには部位特異的に効率よく高分子導入でき、高い比活性を有するバイオコンジュゲート体を創製できる方法の確立が待望されている。

この点著者らは、前述したファージ表面提示法を駆使した“医薬価値に優れた機能性人工蛋白質を迅速創製できるネオ・ダーウィニズム的分子進化戦略”との融合アプローチにより、完全に活性を保持したリジン欠損機能性人工蛋白質を創製することによって、“N末端アミノ基だけを標的とした部位特異的バイオコンジュゲーション”にはじめて成功した<sup>1)</sup>。このリジン欠損機能性人工蛋白質に対する部位特異的バイオコンジュゲーションは

N末端アミノ基にのみ高分子導入されるため、分子的均一性に優れたバイオコンジュゲート体がほぼ100%の収率で得られる。たとえば、TNF- $\alpha$ の場合、上述したように、全6個のリジン残基のうちLys11やLys65・Lys90はその立体構造(三量体)形成やレセプター結合に必須と考えられており<sup>12-14)</sup>、アミノ基に対するランダムなバイオコンジュゲーション法ではこれら活性発現や構造形成に関与するリジン残基までもが修飾されてしまうため、活性低下を避け得なかった。事実、wTNF- $\alpha$ のアミノ基に対するランダムPEGylationでは多様な修飾率(PEG導入率)のPEG化wTNF- $\alpha$ が得られてしまうが、そのなかから1分子のPEG導入体(ランダムモノPEG化wTNF- $\alpha$ :ran-PEG-TNF- $\alpha$ )の収率は20%程度、このran-PEG-TNF- $\alpha$ の残存活性はwTNF- $\alpha$ の約10%にまで減少していた(図1)。一方で、N末端側の8個のアミノ酸を欠損させてもTNF- $\alpha$ の活性は損なわれないことから、活性発現にN末端側は重要でないものと考えられている<sup>15)</sup>。そのため、N末端アミノ基に対する部位特異的モノPEG化リジン欠損TNF- $\alpha$ (sp-PEG-mTNF- $\alpha$ )は80%以上の活性を保持しているなど、圧倒的な利点を有していることが判明した(図1)。

この分子的均一性や比活性、収率に優れた部位特異的PEG化リジン欠損TNF- $\alpha$ は、血中滞留性や抗腫瘍作用の選択的発現能に優れているうえ、従来法で作製したランダムPEG化TNF- $\alpha$ よりも著しく強い*in vivo*抗腫瘍効果を有していることも見出しており、現在臨床応用に向けた研究を推進中である。一方、N末端領域が活性発現に必須である蛋白質の場合でも機能性リジン欠損体を創製したうえで、活性発現とは無関係な領域にあらたなりジン残基を挿入することにより、 $\alpha$ アミノ基と $\epsilon$ アミノ基との反応性の違いを利用した部位特異的バイオコンジュゲーションが可能となることも判明している。

以上の革新的な部位特異的バイオコンジュゲーション法は、著者らが確立した機能性人工蛋白質の分子進化戦略との融合により機能性リジン欠損体を創製することによってはじめて可能となる。現在、種々の蛋白質に関して活性を十二分に保持

したりジン欠損体創出を進めており、今後N末端アミノ基への部位特異的バイオコンジュゲーションの有用性をさらに追求していく予定である。

## DDS機能を有した機能化高分子キャリアの設計

著者らは従来よりバイオコンジュゲート化蛋白質の生体内挙動や*in vivo*薬効発現強度が蛋白質表面を覆う修飾高分子の諸性質によって運命づけられることに着目し、バイオコンジュゲーション法のさらなるグレードアップを目的に、薬物徐放化能や標的指向能といったDDS機能を有する高分子キャリアの分子設計をはかってきた。たとえば、血中滞留性の向上を目的としたバイオコンジュゲーションにはPEGよりもポリビニルピロリドン(PVP)が圧倒的に優れた修飾高分子であること、新規合成したマレイン酸導入PVPやラウリル酸導入PVPがそれぞれIFN- $\gamma$ 誘導能(抗腫瘍免疫誘導能)や高度な脾指向能を有していることなどを明らかにしてきた<sup>6)</sup>。これら新規修飾高分子を用いたバイオコンジュゲーションは単に蛋白質の生体内安定性を高めるだけでなく、高度な組織ターゲティング能やあらたな薬理活性を導入することにより、生理活性蛋白質の目的とする治療作用の選択的発現をさらに保証することを認めている。

このような一連の研究を通じて最近、著者らは腎への高度な薬物送達能とpH応答性薬物徐放化能を合わせもった高分子キャリア(Poly(vinylpyrrolidone-co-dimethyl maleic anhydride):PVD)を新規合成することに成功した<sup>3)</sup>。このPVDはpH8以上で蛋白質のアミノ基と結合し、pH7以下で結合蛋白質を徐々に解離する。一般に炎症組織や癌組織では正常組織よりも低pHであることからPVDを薬物キャリアとして適用した場合、病態組織でのみ効果的に蛋白質がpH応答的に徐放されることを意味している。このPVDをマウスに尾静脈内投与したところ、数時間後に投与量の約80%が腎へ選択的に集積し、4日後には40%に減少していた(図4)。このPVDは腎尿管上皮細胞へのみ選択的に取り込まれるが、細胞毒性をまったく示さないうえ、大量投与しても腎を含め他の組織




図4 PVDの静脈内投与後の組織分布

PVDをマウスに尾静脈内投与したところ、わずか数分で投与量の約30%が腎に集積し、投与後数時間~24時間では投与量の約80%もが腎へ選択的に滞留した。また徐々に尿中排泄され、投与後96時間後には約40%が腎に滞留していた。

に何ら傷害を及ぼさない。さらに、PVDでバイオコンジュゲーションした抗炎症蛋白質(SOD)は体内安定性に優れ、かつ静脈内投与後、選択的に腎へ高集積し、著しい腎炎治療作用を発揮することを見出した。

高齢化社会を迎え、腎不全をはじめとする腎疾患が世界的に深刻な社会問題となっている<sup>16)</sup>。しかし、慢性腎疾患に対する治療は腎移植と透析に頼らざるをえないのが現状であり、患者のQOL (quality of life)の観点からも安全かつ有効な薬物療法の確立が待望されている<sup>17)</sup>。本観点から現在、上述した“医薬価値に優れた機能性人工蛋白質を迅速創製できる分子進化戦略”による機能性人工蛋白質の創製や部位特異的バイオコンジュゲーションシステムとの融合により、あらたな腎疾患治療戦略の確立をさらに推進している。

#### おわりに

本稿で紹介した3段階の“プロテオーム創業にかなうDDS基盤テクノロジー”は、疾患プロテオ

ミクス情報を有効活用したプロテオーム創業の実現と安全かつ有効な蛋白療法の確立に向けて相乗的に機能するものと期待している。また、プロテオーム創業を推進するためにはまず、多種多様な蛋白質とその構造変異体を網羅的に作製し、これらのレセプター・リガンド結合の様式・強度などをも含めた機能情報をハイスループットに評価可能な方法論の構築とその立体構造との連関を網羅的に評価することが必須となる。そのうえで、ゲノムシーケンス情報をもとにあらたに見出された蛋白性シーズなどの機能と構造を予測しうるバイオインフォマティクスが構築されて、ようやく真の意味でプロテオーム創業が可能となってくる。この点、ファージ表面提示法を駆使した“医薬価値に優れた機能性人工蛋白質を迅速創製できる分子進化戦略”は、膨大な多様性をもった構造変異体を創出し、その機能解析を迅速に大量解析しうる最適の基盤テクノロジーとなりうる。

著者らの研究成果は、得られた数多くの機能性人工蛋白質の立体構造と機能特性との連関評価を通じて“機能(医薬価値)→構造”に関する知見の集積が可能となり、将来的に機能性人工蛋白質を合理的設計しうるファーマコ・バイオインフォマティクスの構築にも貢献しうるものと期待している。

#### 文献/URL

- 1) Yamamoto, Y. et al.: Site-specific PEGylation of a lysine-deficient TNF- $\alpha$  with full bioactivity. *Nat. Biotechnol.*, 21: 546-552, 2003.
- 2) Tsubsumi, Y. et al.: Site-specific chemical modification with polyethylene glycol of recombinant immunotoxin anti-Tac(Fv)-PE38(LMB-2) improves antitumor activity and reduces animal toxicity and immunogenicity. *Proc. Natl. Acad. Sci. USA*, 97: 8548-8553, 2000.
- 3) Kamada, H. et al.: Synthesis of a poly(vinylpyrrolidone-co-dimethyl maleic anhydride)co-polymer and its application as renal targeting carrier. *Nat. Biotechnol.*, 21: 399-404, 2003.
- 4) Onda, M. et al.: Lowering the isoelectric point of the Fv portion of recombinant immunotoxins leads to decreased nonspecific animal toxicity without affecting antitumor activity. *Cancer Res.*, 61: 5070-5077, 2001.
- 5) Smith, G. P.: Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science*, 228: 1315-1317, 1985.
- 6) Kamada, H. et al.: Antitumor activity of tumor

- necrosis factor- $\alpha$  conjugated with polyvinylpyrrolidone on solid tumors in mice. *Cancer Res.*, 60 : 6416-6420, 2000.
- 7) Kaneda, Y. et al. : Antitumor activity of tumor necrosis factor- $\alpha$  conjugated with divinyl ether and maleic anhydride copolymer on solid tumors in mice. *Cancer Res.*, 58(2) : 290-295, 1998.
  - 8) Tsutsumi, Y. et al. : PEGylation of interleukin-6 effectively increases its thrombopoietic potency. *Thromb. Haemost.*, 77 : 168-173, 1997.
  - 9) Tsutsumi, Y. et al. : Molecular design of hybrid tumour necrosis factor alpha with polyethylene glycol increases its anti-tumour potency. *Br. J. Cancer*, 71 : 963-968, 1995.
  - 10) Tsutsumi, Y. et al. : Molecular design of hybrid tumour necrosis factor- $\alpha$  II : The molecular size of polyethylene glycol-modified tumour necrosis factor- $\alpha$  affect its anti-tumour potency. *Br. J. Cancer*, 74 : 1090-1095, 1996.
  - 11) Tsutsumi, Y. et al. : Molecular design of hybrid tumor necrosis factor- $\alpha$  III : Polyethylene glycol-modified tumor necrosis factor- $\alpha$  has markedly enhanced anti-tumor potency due to longer plasma half-life and higher tumor-accumulation. *J. Pharmacol. Exp. Ther.*, 278 : 1006-1011, 1996.
  - 12) Yamagishi, J. et al. : Mutational analysis of structure-activity relationships in human tumor necrosis factor- $\alpha$ . *Protein Eng.*, 3 : 713-719, 1990.
  - 13) Van Ostade, X. et al. : Localization of the active site of human tumour necrosis factor(hTNF) by mutational analysis. *Embo J.*, 10 : 827-836, 1991.
  - 14) Loetscher, H. et al. : Human tumor necrosis factor alpha(TNF $\alpha$ ) mutants with exclusive specificity for the 55 kDa or 75 kDa TNF receptors. *J. Biol. Chem.*, 268 : 26350-26357, 1993.
  - 15) Jones, E. Y. et al. : Structure of tumour necrosis factor. *Nature*, 338 : 225-228, 1989.
  - 16) Jones, C. A. : Serum creatinine levels in the US population : third National Health and Nutrition Examination Survey. *Am. J. Kidney Dis.*, 32 : 992-999, 1998.
  - 17) Progress and Priorities : Renal disease research plan. Report of the strategic planning conferences- Renal research properties-sponsored by National Institute of Diabetes, and Digestive and Kidney Disease, Council of American Kidney Societies (December 5-6, 1998 and February 4-5, 1999). (<http://www.niddk.nih.gov/federal/planning.htm#g>)

●お知らせ●

第14回運動処方講習会

テーマ：セミアドバンスコース  
 日 時：2004年12月4日(土)午後1時～6時30分  
 会 場：東京体育館第一研修室  
 受講資格・対象者：病院、スポーツ施設等の運動療法に携わっている医師、看護師、臨床検査技師、理学療法士、健康運動指導士、トレーナー等  
 受講料：医師 18,000円、医師以外 15,000円  
 受講定員：約120名(先着順)  
 申込要領：  
 HP <http://www.senmon-i.ne.jp/cepp-koushuukai/>から「参加申込書」をプリントアウトし、参加希望回にはっきりと○をつけ、必要事項をご記入の上、FAXまたは郵便でお申し込み下さい。申込確認後に、受講料、受講料振込用紙等をご送付いたします。当日

は必ず受講票および振込用紙の控えをお持ち下さい。

申込先：ミナト医科学株式会社東京支店 運動処方講習会担当 松岡  
 住所：〒113-0033 東京都文京区本郷3-40-3  
 TEL：03-3815-3710 FAX：03-3818-0940  
 申し込み期間：  
 第14回：平成16年8月2日(月)～11月19日(金)(定員になり次第申し込み終了)  
 問い合わせ先：運動処方研究会事務局  
 FAX：03-3401-0472 E-mail：cepp@cepp.ne.jp  
 ミナト医科学(担当 染谷)  
 FAX：03-3818-0940 E-mail：m-someya@minato-med.co.jp  
 事務局：(財)心臓血管研究所内(E-mail：cepp@cepp.ne.jp)

《若手研究者紹介》



蛋白療法の最適化に叶う DDS の開発を目指して

堤 康 央 Yasuo Tsutsumi

大阪大学薬学研究科薬剤学分野

1. はじめに

若手研究者紹介コラムへの執筆機会を賜り、そのタイトルを思い巡らせ、最終的に上記とさせて頂いた。筆者は、学部学生として大阪大学薬学部薬剤学講座に配属してからの14年間、一貫してこのタイトルに取り組んできた。この間、国立循環器病センター研究所や米国の国立衛生研究所 (NIH) への留学などにより、多くの先生方から最新の医用高分子化学、遺伝子工学、蛋白質工学をご教授賜り、研究の質的グレード・アップを図ってきたものの、その研究対象は遺伝子そのものではなく、常にその翻訳産物である蛋白質にあった。今改めて周りを見渡すと、蛋白質の医薬品化を目指した DDS 研究、特に PEGylation に代表される蛋白質の高分子バイオコンジュゲーションにしがみついているのは筆者ぐらいなもので、恩師 真弓忠範先生からの叱咤激励と情熱、愛情が無ければ、途中で挫折していたかもしれない。その真弓忠範先生が、この2004年3月31日付けで大阪大学をご停年退官される。本稿では、これまでの想いを込めて、上記タイトルに関して、

筆者紹介：1991年3月大阪大学薬学部薬学科卒業、同年4月国立循環器病センター研究所生体工学部研究員(松田武久先生)、1993年3月大阪大学大学院薬学研究科応用薬学専攻修士課程修了、1994年8月大阪大学大学院薬学研究科応用薬学専攻博士課程中退後、直ちに大阪大学薬学部助手、1997年8月薬学博士(大阪大学)、現在に至る。この間、1999年4月～2001年1月米国 National Cancer Institute (NCI) / National Institutes of Health (NIH) 博士研究員 (Laboratory of Molecular Biology [Dr. Ira PASTAN]) に留学。平成9年度日本薬学会近畿支部奨励賞、平成16年度日本薬学会奨励賞、平成16年度日本薬剤学会奨励賞などを受賞。

我々がこれまで歩んできた道のりの一端を、紙面の許す限り、紹介させて頂きたい。

2. プロテオーム創薬と21世紀型蛋白療法

ヒトゲノム解読が完了し、ポストゲノム研究は蛋白質の時空間的、質的、量的な機能発現様式と立体構造との連関を包括的に解明しようとするプロテオミクスや構造ゲノミクスへと集約されつつある。これは、ゲノムが単なる情報(遺伝型)に過ぎず、そのままでは何ら機能を発揮し得ないこと、転写した RNA が機能する極一部の例外を除き、このゲノム情報に基づき、多様な蛋白質群(プロテオーム; 表現型)が翻訳され、その機能を発揮することで初めて生命現象が営めるようになることを考えると当然のことと言えよう。この研究の流れは、癌や種々感染症、自己免疫疾患などに対する抗体療法やサイトカイン療法といった21世紀医療が台頭してきたことも相俟って、疾病治療に有効な蛋白質を探索・創製しようとするプロテオーム創薬と、これらを適用した新規蛋白療法への期待を加速度的に高めている。

しかし過去の多くの事例が示しているように、蛋白質は一般に、体内安定性に極めて乏しいため、臨床応用の際には大量頻回投与を余儀なくされ、往々にして重篤な副作用を招いてしまう。なかでもサイトカインなどは、多彩な細胞上の複数種類のレセプターを介して、多様な *in vivo* 生理活性を示すため、目的とする治療作用のみならず副作用の原因となる他の作用までもを同時に発揮してしまう。そのため周知のように、医薬品化された蛋白質は極一部にすぎないのが現状である。従ってプロテオーム創薬や

21世紀型蛋白療法を推進するためには、このような蛋白質固有の問題点を克服できるDDSの確立が依然として必須となっている。本観点から我々は、(1)レセプター親和性・特異性などが高く、医薬価値に優れた機能性人工蛋白質を迅速創出できるネオ・ダーウィニズムの分子進化戦略(プロテオーム創薬システム)の構築<sup>1)</sup>、(2)蛋白質の生体内安定性を向上させ、かつ目的治療作用の選択的発現能を付与できる部位特異的高分子バイオコンジュゲーション法の確立<sup>1~4)</sup>、(3)DDS機能(標的指向能・薬物徐放化能など)を有した高機能化修飾高分子の設計<sup>5)</sup>など、上記三者を融合させた「蛋白療法の最適化に叶うDDSの確立」に関する研究を進めている。

### 3. プロテオーム創薬システム

蛋白療法の最適化に向け、従来から産官学の多くのバイオ研究機関が、特定レセプターへの親和性や選択性に優れた機能性人工蛋白質などを創製するため、Kunkel法といった点突然変異法を用いた構造変異蛋白質(アミノ酸置換体)の作製を精力的に試みている。しかし点突然変異法では、まず構造変異蛋白質の立体構造や機能をシミュレーションし、トライ・アンド・エラーで生理活性蛋白質の構成アミノ酸を一つずつ別の特定アミノ酸に改変することにより、個々の構造変異蛋白質を作製せねばならない。そのうえで目的とする機能性人工蛋白質を探索・同定するため、作製した構造変異蛋白質の諸機能を個別に評価する必要がある。そのため従来法では、時間ばかりが消費され、かつ作製し得る構造変異構造変異蛋白質の多様性(種類)にも限界があるなど、期待通りの成果は得られていない。そのため、より迅速かつ効率よく目的作用を有する機能性人工蛋白質を創製できるテクノロジーの開発が望まれている。

この点我々は最近、ファージ表面提示法を独自に改良することにより、蛋白質中の多数のアミノ酸を各々20種類のアミノ酸へ一挙に置換することで、 $10^8$ 種類以上もの多様性を有した構造変異蛋白質をCombinatorial Biosynthesisし、この中からレセプター親和性・特異性などの高い「医薬価値に優れた機能性人工蛋白質」を迅速かつ効率良く同定できる基盤テクノロジー(プロテオーム創薬システム)を確立した。例えばアラニン・スキャンといった従来

の点突然変異法を用いた構造-活性相関研究により、腫瘍壊死因子(TNF- $\alpha$ )のLys 11やLys 65, Lys 90はその立体構造(三量体)形成やレセプター結合に必須の役割を担っているものと考えられていた。これはTNF- $\alpha$ に限らず、一般にリジン残基は多くの場合、生理活性蛋白質の高次構造形成やリガンド-レセプター結合などに必須の役割を担っているため、他のアミノ酸への置換は致命的な活性低下を招いてしまうことが、従来までの点突然変異解析によって常識となっていた。事実これまで、活性を完全に保持させたまま、蛋白質中のリジン残基全てを欠損させ得た例(機能性リジン欠損体)は皆無であった。しかし我々はこの概念を覆す知見、即ちTNF- $\alpha$ 中の全6個のリジン残基を一挙に他のアミノ酸へ置換しても、wild型TNF- $\alpha$ (wTNF- $\alpha$ )と同等から10倍以上ものレセプター親和性や生物活性を有する機能性リジン欠損TNF- $\alpha$ を創製することに初めて成功した。このwTNF- $\alpha$ と同等以上の生物活性を有する種々の機能性リジン欠損TNF- $\alpha$ は、BIAcoreを用いたTNFレセプター1やTNFレセプター2への結合性評価により、wTNF- $\alpha$ と同等以上のレセプター親和性を有していること、ゲル濾過解析などから三量体を形成していることも確認している。またこれらの機能性リジン欠損TNF- $\alpha$ の体内挙動を評価したところ、wTNF- $\alpha$ よりも血中滞留性に優れていることを認めた。この血中滞留性の向上は、カチオン性のリジン残基を他のアミノ酸へ置換したことによるLowering pI効果に起因したものと考えられる。さらにこれらの機能性リジン欠損TNF- $\alpha$ の*in vivo*における抗腫瘍効果は、Lowering pI効果により、wTNF- $\alpha$ と比較して数倍から10倍にまで向上しているうえ、その毒性は顕著に低下しており、その治療域はwTNF- $\alpha$ の十数倍以上にも向上していることが明らかとなった。これらの知見は、TNF- $\alpha$ 分子中の全6個のリジン残基を他のアミノ酸へ一挙かつ網羅的に置換した $20^6$ (6,400万)種類もの構造変異TNF- $\alpha$ (TNF- $\alpha$ のアミノ酸置換体)を表面提示したファージライブラリを作製したうえで、TNFレセプター1や抗TNF中和抗体に対するアフィニティー・バイオパンニングを行い、選択・濃縮された構造変異TNF- $\alpha$ 群の諸機能を高速解析することによって得られたものである。即ち以上の事実は、ファージ表面提示法を駆使した基盤テクノ

ロジを適用することで、今まで全く創造し得なかった医薬価値に優れた機能性人工蛋白質を新たに産み出し得ることを強く示しており、我々が確立した「プロテオーム創薬システム」は、他を圧倒する“プロテオーム創薬のための競争力(DDS基盤テクノロジー)”を提供するだけでなく、従来までの点突然変異法(アラニン・スキャン)で得られた知見では想像もでき得なかった“蛋白改変の概念”や“蛋白質の構造—活性相関概念”をも新たに提唱するものと期待している。

#### 4. プロテオーム創薬のための バイオインフォマティクス

プロテオーム創薬は、プロテオミクスおよび構造ゲノミクスの進展と、これらの知見を統括したバイオインフォマティクスが駆動力となり、近い将来、上記の「プロテオーム創薬システム」との融合により加速度的に推進されるものと期待される。即ちこのようなプロテオーム創薬を指向したバイオインフォマティクスの進展は、蛋白質のアミノ酸配列(一次構造)と立体構造(三次構造)、機能との連関を理解可能とするため、近未来的には蛋白質をコードした塩基配列や蛋白質のアミノ酸配列さえ判明すれば、その立体構造と機能が予測できることになる。これは逆に目的とする機能や立体構造を有した機能性蛋白質の新規デザインを可能とするだけでなく、蛋白質の立体構造やその機能を模倣した有機化合物の合理的設計をも可能にするものと期待される。このようなバイオインフォマティクスをシステムアップするためには、未知蛋白質の機能解明や立体構造解析に加え、種々の蛋白質について膨大な多様性を有する構造変異体を網羅的に作製し、レセプター・リガンド結合の様式、生物活性等をも含めた機能情報を集積し、立体構造との連関を追求しなければならない。この点我々が開発した「プロテオーム創薬システム」は、視点を変えれば僅か1週間で $10^8$ 種類以上もの多様性を有する構造変異蛋白質のライブラリを作製し、その機能情報を高速集積できる基盤技術と言える。本観点から現在、機能性人工TNF- $\alpha$ を含む様々な構造変異蛋白質の機能評価と共に、そのX線結晶構造解析を進めており、近未来的なバイオインフォマティクスへの研究展開を予定している。

一方で最近、Gene shufflingや人工遺伝暗号システムなどを用いた機能性人工蛋白質の創出に注目が集まっている。これら興味深いアプローチは天然に存在しない新たなアミノ酸配列を有した蛋白質を人工的に作製しようとするものであるが、残念ながら臨床応用可能な非天然型生理活性蛋白質の創製には至っていない。当然のことながら、上述した我々の基盤テクノロジーはこれら非天然型生理活性蛋白質の探索や創製、安定化や高機能化にも適用可能であり、現在Gene shuffling法とファージ表面提示法を融合した新たな機能性人工蛋白質の創製システムの構築を進めている。

#### 5. 部位特異的高分子バイオコンジュゲーション法

前述した(2)および(3)について我々は、主として酵素蛋白質への適用に限定されていたバイオコンジュゲーション法を、生理活性蛋白質一般に適用するためのグレードアップを図ってきた。その結果、1. 蛋白質の作用機構を考慮し最適の修飾高分子を選択し、2. 比活性—修飾率—分子サイズ等の相関をもとに最適条件を見出すことにより、①蛋白質の生体内安定性を飛躍的に向上させ得ること、②多様な*in vivo*作用の中から、目的治療作用のみを数百倍にも高め得ることを明らかにしている。この②の生理活性蛋白質への作用の選択性付与は、投与量の削減や副作用発現組織への移行性低下に因ることを明らかにし、Interleukin-6やTNF- $\alpha$ のバイオコンジュゲーションの場合、副作用を増幅することなく目的とする血小板産生促進効果や抗腫瘍効果が各々500倍および100倍にも選択増強されることを認めている。このようにバイオコンジュゲーションは蛋白質の医薬品開発において、最適DDS技術の地位を確立しつつあるが、その成功例は極めて少ない。この最大の原因は、水溶性高分子を蛋白質に結合させる際に、水溶性高分子がリジン $\epsilon$ アミノ基やN末端 $\alpha$ アミノ基ヘランダムに結合してしまうことにある。そのため、生理活性蛋白質の活性発現部位への結合は致命的な比活性低下をきたし、結合分子数・部位の違いはバイオコンジュゲート体の分子的・機能的不均一性をもたらす結果となる。我々はこれらの問題を解決するため、前述した①を応用して、生理活性蛋白質の活性を完全に保持したリジン欠損体を創製し、「N末端アミノ基のみを標的とした部位

特異的バイオコンジュゲーション法」を確立した。このリジン欠損体に対する部位特異的バイオコンジュゲーションは、分子的均一性に優れたバイオコンジュゲート体が最大 100% の収率で得られる。またリコンビナントの野生型 TNF- $\alpha$  の場合、従来までのランダムなバイオコンジュゲーションでは僅か 1 分子の高分子導入により 10% 以下まで活性低下してしまうが、リジン欠損 TNF- $\alpha$  に対する部位特異的バイオコンジュゲーションは殆ど活性低下を招かないなど、圧倒的な利点を有していた (図 1)。この分子的均一性や比活性に優れた部位特異的 PEG 化リジン欠損 TNF- $\alpha$  は、血中滞留性や抗腫瘍作用の選択的発現能にも優れているうえ、従来法で作製したランダム PEG 化 TNF- $\alpha$  よりも著しく強い *in vivo* 抗腫瘍効果を有していた。一方で我々は、従来よりバイオコンジュゲート体の生体内挙動や薬効発現強度が、蛋白質表面を覆う修飾高分子の諸性質によって運命付けられることに着目し、バイオコンジュゲーション法のさらなるグレードアップを目的に、多数の機能化修飾高分子の設計を図ってきた。例えば最近、我々は腎臓への高度な薬物送達能と pH 応答性薬物徐放化能を併せ持った PVD を新規合成することに成功した (図 2)。現在、上述した「プロテオーム創薬システム」によるリジン欠損機能性人工蛋白質の創製や部位特異的バイオコンジュゲーションシステムとの融合により、新たな治療戦略の確立を目指した研究を推進している。

## 6. おわりに

以上、足早ではあるが、真弓忠範先生と共に走り続けてきた「蛋白療法の最適化に叶う DDS の開発を目指して」という道のりの極一端を紹介させて頂いた。前述したように蛋白質の臨床応用はまだまだ著しく制限されており、そのほとんどは医薬品化されていない。従って、プロテオーム創薬を推進し、種々の蛋白質を有効な医薬品として適用した 21 世紀型蛋白療法を実現するためには、蛋白質固有の種々問題点を克服できる戦略を早急に構築せねばならない。即ち、特定レセプターへの親和性や指向性に優れたスーパーアゴニストやアンタゴニスト、体内安定性や血中滞留性に優れたミュータント等、高い安全性と有効性を保証し得る機能性人工蛋白質を、疾病や治療目的に応じて迅速かつ合理的に創出




図 1 リジン欠損 TNF- $\alpha$  およびその部位特異的バイオコンジュゲート体の *in vitro* 活性評価  
○, wTNF- $\alpha$ ; △, ran-PEG-TNF- $\alpha$  (ランダム・モノ PEG 化体); ●, リジン欠損 TNF- $\alpha$ ; ▲, sp-PEG-mTNF- $\alpha$  (部位特異的モノ PEG 化体)。




図 2 PVD の静脈内投与後の組織分布

できる創薬テクノロジーの確立が不可欠である。この創薬テクノロジーは、薬剂的観点から言えば、蛋白性薬物による疾病治療の最適化を目指した DDS であり、ポストゲノム基礎研究と 21 世紀医療との架け橋となり得るものと位置づけられるものと考えている。



本研究は、恩師・真弓忠範先生にご指導賜り、遂行されたものであり、また本稿の執筆に際しても格段のご配慮を賜りました。心より御礼を申し上げます。また種々ご教授頂いた共同研究者の先生方、種々ご協力頂いた大阪大学薬学研究科薬剤学分野助教授 中川晋作先生ならびに学生諸氏に深謝申し上げます。

#### 引用文献

- 1) Y. Yamamoto, Y. Tsutsumi, Y. Yoshioka, T. Nishibata, K. Kobayashi, T. Okamoto, Y. Mukai, T. Shimizu, S. Nakagawa, S. Nagata, T. Mayumi, Site-specific PEGylation of a lysine-deficient TNF-alpha with full bioactivity, *Nat. Biotechnol.*, **21**, 546-552 (2003).
- 2) Y. Kaneda, Y. Yamamoto, H. Kamada, S. Tsunoda, Y. Tsutsumi, T. Hirano, T. Mayumi, Antitumor activity of tumor necrosis factor-alpha conjugated with divinyl ether and maleic anhydride copolymer on solid tumors in mice, *Cancer Res.*, **58** (2), 290-295 (1998).
- 3) H. Kamada, Y. Tsutsumi, Y. Yamamoto, T. Kihira, Y. Kaneda, Y. Mu, H. Kodaira, S. Tsunoda, S. Nakagawa, T. Mayumi, Antitumor activity of tumor necrosis factor-alpha conjugated with polyvinylpyrrolidone on solid tumors in mice, *Cancer Res.*, **60**, 6416-6420 (2000).
- 4) Y. Tsutsumi, M. Onda, S. Nagata, B. Lee, R.J. Kreitman, I. Pastan, Site-specific chemical modification with polyethylene glycol of recombinant immunotoxin anti-Tac (Fv)-PE38 (LMB-2) improves antitumor activity and reduces animal toxicity and immunogenicity, *Proc. Natl. Acad. Sci. U.S.A.*, **97**, 8548-8553 (2000).
- 5) H. Kamada, Y. Tsutsumi, K. Sato-Kamada, Y. Yamamoto, Y. Yoshioka, T. Okamoto, S. Nakagawa, S. Nagata, T. Mayumi, Synthesis of a poly(vinylpyrrolidone-co-dimethyl maleic anhydride) o-polymer and its application as renal targeting carrier, *Nat. Biotechnol.*, **21**, 399-404 (2003).



## Poly(vinylpyrrolidone-co-dimethyl maleic acid) as a novel renal targeting carrier

Yoko Yamamoto<sup>1</sup>, Yasuo Tsutsumi<sup>\*1</sup>, Yasuo Yoshioka<sup>1</sup>, Haruhiko Kamada, Keiko Sato-Kamada, Takayuki Okamoto, Yohei Mukai, Hiroko Shibata, Shinsaku Nakagawa, Tadanori Mayumi

*Department of Biopharmaceutics, Graduate School of Pharmaceutical Sciences, Osaka University, 1-6 Yamadaoka, Suita, Osaka 565-0871, Japan*

Received 18 August 2003; accepted 25 November 2003

### Abstract

Poly(vinylpyrrolidone-co-dimethyl maleic acid) (PVD) was found to have high renal-targeting capability and safety as a drug carrier. To optimize the renal drug delivery system using PVD, the relationship between the molecular weight of PVD and its renal accumulation were evaluated in mice by their intravenous injection. It was found that the molecular size of 6–8 kDa was associated with the highest renal accumulation. The specific bioactivity of PVD-conjugated superoxide dismutase (SOD) relative to that of unmodified SOD gradually decreased with an increase in the degree of modification to SOD with PVD6K. The conjugated SOD (L-PVD-SOD) with the molecular size of 73 kDa, which had comparable specific bioactivity with native SOD, showed longer plasma half-life than native SOD. About sixfold more L-PVD-SOD was distributed to the kidneys than native SOD 3 h after intravenous injection, whereas extensive PVD modification did not enhance the renal accumulation of SOD. This L-PVD-SOD effectively accelerated recovery from mercuric chloride-induced acute renal failure in vivo. These results suggest that L-PVD-SOD may be the optimal derivative as a potential therapeutic agent to various renal diseases. © 2004 Elsevier B.V. All rights reserved.

**Keywords:** Renal targeting; Vinylpyrrolidone; Dimethyl maleic acid; Polymeric carrier; Acute renal failure; Drug delivery system

### 1. Introduction

In recent years, clinical application of bioactive proteins, such as cytokines and growth factors, have

been studied. However, most of these proteins are limited in their clinical application because of their various side effects [1,2]. In general, the plasma half-lives of bioactive proteins in vivo are very short [3–5], so that their frequent administration at a high dosage are needed to obtain sufficient therapeutic effects. This administration markedly destroys homeostasis, resulting in unexpected side effects. In addition, since bioactive proteins exhibit diverse pharmacological actions in various tissues, it is difficult to obtain only the favorable actions (therapeutic effects)

*Abbreviations:* DMMA, dimethyl maleic anhydride; PVP, polyvinylpyrrolidone; VP, vinylpyrrolidone; SOD, superoxide dismutase; PVD-SOD, PVD-modified SOD; PEG, Polyethylene glycol.

\* Corresponding author. Tel./fax: +81-6-6879-8178.

E-mail address: [tsutsumi@phs.osaka-u.ac.jp](mailto:tsutsumi@phs.osaka-u.ac.jp) (Y. Tsutsumi).

<sup>1</sup> These authors contributed equally to the work.

selectively. To overcome their problems, bioactive proteins have been conjugated with water-soluble polymeric carriers. We have already reported that polymer conjugation of cytokines typified with tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ), interleukin-6 (IL-6), and immunotoxin (IT), with polyethylene glycol (PEG), and polyvinylpyrrolidone (PVP) improved its resistance to proteinase and plasma half-lives and resulted in greater therapeutic potency [6–10]. We have also shown that conjugation with polymeric carriers regulate the tissue distribution of cytokine, resulted in a selective increase its desirable therapeutic effects, and decrease its undesirable side effects. However, for further enhancement of the therapeutic potency and safety of conjugated bioactive protein, more precise control of the *in vivo* behavior of each protein for selective expression of their therapeutic effect is necessary. Thus, it is expected to develop novel polymeric carriers with targeting capability to specific tissue, whereas PEG and PVP are useful and powerful polymeric carriers for improving the plasma half-lives of proteins.

Renal disease is a serious problem on the increase in the world. There is no cure for renal disease, and few strategies are available for prevention [11,12]. Bioactive proteins, such as superoxide dismutase (SOD) and interleukin-10 (IL-10), have been expected to prevent the progression of renal disease, but their therapeutic potency were too low because they were poorly distributed to the kidney. The development of a renal delivery system that selectively carries drugs to the kidneys is a promising approach for limiting tissue distribution and controlling toxicity. Several renal drug delivery systems have been previously described. One approach involves prodrugs that are cleaved by kidney-associated enzymes to release the drugs in the kidney [13]. However, these prodrugs generally do not accumulate in the kidneys as a result of plasma protein binding and limited transport to the kidney. Low-molecular-weight proteins, such as lysozyme, have been used as carriers because they are reabsorbed by the kidneys. Unfortunately, they also produced strong renal toxicity and cardiovascular side effects [14]. Streptavidin carriers bind to biotin in the kidney, but they are immunogenic and have limited renal accumulation due to their large molecular size [15,16]. Thus, it is important to develop an effective renal drug delivery

system that not only targets the kidney but also has excellent safety.

Recently, we synthesized poly(vinylpyrrolidone-co-dimethyl maleic acid) (PVD) as a new renal targeting carrier [17]. About 80% of the dose of PVD selectively distributed to kidneys after intravenous injection and then gradually excreted to urinary and approximately 40% remained in the kidneys 4 days after intravenous injection. No side effects occurred in the kidney and other tissues by administration of excessively high dose of PVD. In this study, to assess the usefulness of PVD as a renal targeting polymeric carrier of drugs, we first evaluated the relationship between PVD molecular weight and renal accumulation. We then prepared a conjugated SOD with PVD and evaluated the pharmacokinetic characteristics and therapeutic effects on HgCl<sub>2</sub>-induced acute renal failure (ARF). This study will provide fundamental information enabling us to create a powerful drugs against renal disease.

## 2. Materials and methods

### 2.1. Synthesis and purification of PVD

Chemicals were obtained from Wako (Osaka, Japan) and 2,3-dimethyl maleic anhydride (DMMA) was purchased from Acros Organic (New Jersey, USA). PVD was synthesized by the radical polymerization method using 4,4'-azobis-4-cyanovaleric acid (ACVA) as a radical initiator. Briefly, DMMA and *N*-vinyl-2-pyrrolidone (VP) were mixed in a ratio of 1:5 in a glass tube containing dimethyl formamide (DMF) and incubated at 60 °C for 96 h. The resulting copolymer was precipitated in dry diethyl ether, collected immediately after filtration, and dried under vacuum. The molecular weight was determined by GFC (TSKgel G4000PW, Tosoh, Tokyo). PVD was separated into four fractions of different number-average molecular weight values. In addition, the number-average molecular weight ( $M_n$ ) of each fraction was calculated by comparing it with PEG standards (Fr.1; 14 kDa, Fr.2; 8 kDa, Fr.3; 6 kDa, Fr.4; 3 kDa). The polydispersity [weight-average molecular weight ( $M_w$ )/ $M_n$ ] of these PVDs were about 1.1.




Fig. 1. Plasma clearance and tissue distribution of PVDs with various molecular weight after intravenous injection. PVD with various molecular weight were  $^{125}\text{I}$ -labeled by chloramine-T method. Mice were intravenously injected with  $^{125}\text{I}$ -labeled polymers. (A) After administration, blood was collected from the tail at indicated time, and the radioactivity was measured by a  $\gamma$ -counter. (B) At 3 h after intravenous injection, mice were sacrificed, and each organ was collected. The radioactivity was measured by a  $\gamma$ -counter. Data represent mean  $\pm$  S.D. ( $n=5$ ).

## 2.2. Preparation of $^{125}\text{I}$ -labeled PVDs

To study the pharmacokinetics of PVDs with various  $M_n$ , radiolabeled PVDs were prepared by the chloramine-T method. PVDs were dissolved in DMF and activated with dicyclohexyl carbodiimide and *N*-hydroxysuccinimide for 24 h at room temperature. The PVDs were reacted with tyramine hydrochloride for 24 h at 4 °C, dialyzed in water, and lyophilized. PVD-tyramine conjugates were radiolabeled by the chloramine-T method.  $^{125}\text{I}$ -labeled PVDs were purified by GFC. The specific activities of  $^{25}\text{I}$ -labeled PVDs were about 4.44  $\mu\text{Ci}/\text{mg}$  polymer.

## 2.3. Measurement of plasma clearance and tissue distribution

All experimental protocols for animal studies were in accordance with the "Guide for Laboratory Animal Facilities and Care" (NIH Publication 85-23, revised 1985). These protocols have been approved by the committee in the Pharmaceutical School, Osaka University. The clearance of PVD in male BALB/c mice (5 weeks old; SLC, Hamamatsu, Japan) were studied after intravenous injection of 10  $\mu\text{g}$  of polymer per mouse. Blood was collected from the tail vein at intervals, and radioactivity was measured in each sample using a  $\gamma$ -counter. After 3 h, most plasma

$^{125}\text{I}$ -radioactivity was present as  $^{125}\text{I}$ -labeled polymers and not as free  $^{125}\text{I}$ . To evaluate tissue distribution, mice were housed in metabolic cages to collect urine and sacrificed 3 h after treatment. Clearance value was pharmacokinetically evaluated based on the clearance concept as described [18,19].

## 2.4. Preparation of PVD-conjugated SOD

Bovine erythrocyte Cu-Zn SOD was reacted with a 22-fold molar excess of activated PVD at room

Table 1  
Clearance of PVDs with various molecular weight

	Clearance ( $\mu\text{l}/\text{h}$ )			
	Kidney	Liver	Lung	Spleen
PVD $M_n=14$ kDa	2786.4 $\pm$ 52.9	195.4 $\pm$ 13.9	10.6 $\pm$ 3.0	9.8 $\pm$ 0.4
PVD $M_n=8$ kDa	6614.0 $\pm$ 199.5	119.3 $\pm$ 3.2	14.3 $\pm$ 4.4	5.7 $\pm$ 0.3
PVD $M_n=6$ kDa	8888.4 $\pm$ 214.5	63.6 $\pm$ 1.4	35.2 $\pm$ 8.3	4.6 $\pm$ 0.2
PVD $M_n=3$ kDa	3871.7 $\pm$ 216.8	45.4 $\pm$ 6.1	15.7 $\pm$ 7.0	3.2 $\pm$ 0.3

Mice were used in groups of five. Each value is the mean  $\pm$  S.E.




Fig. 2. Renal accumulation and urinary excretion of PVD with various molecular weight after intravenous injection. Mice were intravenously injected with  $^{125}\text{I}$ -labeled PVDs. (A) At indicated time after intravenous injection, mice were sacrificed, and kidneys were collected. The radioactivity was counted. (B) Mice were housed in metabolic cages to collect urine for 24 h after intravenous injection. The radioactivity of urine was counted. The radioactivity was measured by a  $\gamma$ -counter. Data represent mean  $\pm$  S.D. ( $n=5$ ).

temperature for 90 min in pH 8.5 borate buffer. The resultant PVD-SOD (PVD-SOD) was separated into three fraction of different molecular size ( $M_s$ ) by gel filtration chromatography (Superose 12 HR10/3; Amersham Pharmacia Biotech, Uppsala, Sweden) in 0.05 M phosphate buffer (pH 7.2). The  $M_s$  of each fraction was estimated by gel filtration chromatography with protein standard (gel filtration calibration kit; Amersham Pharmacia Biotech). The specific activity of PVD-SODs was determined by the cytochrome *c* method. The bioactivity of  $^{125}\text{I}$ -native SOD and PVD-SODs prepared by the chloramine-T method were indistinguishable from their non-radiolabeled forms.

### 2.5. Therapeutic effects of PVD-SOD on ARF

We used a mouse model of  $\text{HgCl}_2$ -induced ARF to assess the therapeutic effects of PVD-SOD (L-PVD-SOD) with  $M_s$  of 73 kDa. ARF was induced by subcutaneous injection of 8 mg/kg  $\text{HgCl}_2$ . Then, 12 h later, these mice were given 4 mg protein/kg native SOD or L-PVD-SOD intravenously. Therapeutic efficacy was qualitatively assessed 48 h after treatment with  $\text{HgCl}_2$  by measuring the level of urinary alkaline phosphatase (ALP), urinary  $\gamma$ -glutamyltransferase (GTP), urinary *N*-acetyl- $\beta$ -glucosaminidase (NAG), urinary glucose, urinary keton, urinary protein, uri-

nary hemoglobin, and serum creatinine as ARF markers.

## 3. Results

### 3.1. Pharmacokinetics of PVD with various molecular weight

To evaluate the influence of molecular weight on renal accumulation of PVD, we estimated the plasma clearance and tissue distribution of PVD with various  $M_n$  after intravenous injection (Fig. 1). Radioactivity

Table 2  
Characterization of PVD-conjugated SOD

	Number average <sup>a</sup> molecular weight	SOD enzymatic activity <sup>b</sup> ( $\times 10^3$ U/mg)	Remaining <sup>c</sup> activity (%)
H-PVD-SOD	220,000	$1.47 \pm 0.11$	63.1
M-PVD-SOD	120,000	$1.97 \pm 0.31$	84.5
L-PVD-SOD	73,000	$2.19 \pm 0.05$	93.9
Native SOD	32,000	$2.33 \pm 0.03$	100.0

<sup>a</sup> Number-average molecular weight was determined by gel filtration HPLC (protein standard).

<sup>b</sup> Specific activity was measured by cytochrome *c* method. Each value represents the mean  $\pm$  S.D.

<sup>c</sup> Remaining activity (%) was calculated base on native SOD.

- [22] P.M. Tiidus, Radical species in inflammation and overtraining. *Can. J. Physiol. Pharmacol.* 76 (1998) 533–538.
- [23] M.J. Finnen, C.M. Lawrence, S. Shuster, Inhibition of dithranol inflammation by free-radical scavengers. *Lancet* 2 (1984) 1129–1130.
- [24] A. Guglielmotti, A. Capezzone De Joannon, N. Cazzolla, M. Marchetti, L. Soldo, G. Cavallo, M. Pinza, Radical scavenger activity of bendazac, an anticataract non-steroidal anti-inflammatory agent. *Pharmacol. Res.* 32 (1995) 369–373.
- [25] A. Garcia-Gonzalez, J.L. Ochoa, Anti-inflammatory activity of *Debaryomyces hansenii* Cu, Zn-SOD. *Arch. Med. Res.* 30 (1999) 69–73.
- [26] C. Canavese, P. Stratta, A. Vercellone, Oxygen free radicals in nephrology. *Int. J. Artif. Organs* 10 (1987) 379–389.
- [27] T. Fujita, H. Furitsu, M. Nishikawa, Y. Takakura, H. Sezaki, M. Hashida, Therapeutic effects of superoxide dismutase derivatives modified with mono- or polysaccharides on hepatic injury induced by ischemia/reperfusion. *Biochem. Biophys. Res. Commun.* 189 (1992) 191–196.
- [28] K. Mihara, Y. Oka, K. Sawai, Y. Takakura, M. Hashida, Improvement of therapeutic effect of human recombinant superoxide dismutase on ischemic acute renal failure in the rat via cationization and conjugation with polyethylene glycol. *J. Drug Target.* 2 (1994) 317–321.



## Augmentation of antigen-specific immune responses using DNA-fusogenic liposome vaccine

Tomoaki Yoshikawa<sup>a,b</sup>, Susumu Imazu<sup>a</sup>, Jian-Qing Gao<sup>a</sup>, Kazuyuki Hayashi<sup>a</sup>, Yasuhiro Tsuda<sup>a</sup>, Mariko Shimokawa<sup>b</sup>, Toshiki Sugita<sup>a,b</sup>, Takako Niwa<sup>a,b</sup>, Atushi Oda<sup>a,b</sup>, Mitsuru Akashi<sup>b,c</sup>, Yasuo Tsutsumi<sup>b,d</sup>, Tadanori Mayumi<sup>e</sup>, Shinsaku Nakagawa<sup>a,b,\*</sup>

<sup>a</sup> Department of Biopharmaceutics, Graduate School of Pharmaceutical Sciences, Osaka University, Japan

<sup>b</sup> Creation of Bio-devices and Bio-systems with Chemical and Biological Molecules for Medical Use, CREST, Japan Science and Technology Corporation (JST), Japan

<sup>c</sup> Department of Molecular Chemistry, Graduate School of Engineering, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

<sup>d</sup> National Institute of Health Science, Osaka Branch Fundamental Research Laboratories for Development of Medicine, 1-1-43 Hoenzaka, Chuo-ku, Osaka 540-0006, Japan

<sup>e</sup> Department of Cell Therapeutics, Graduate School of Pharmaceutical Sciences, Kobe-Gakuin University, 518 Arise, Igawadani, Nishiku, Kobe 651-2180, Japan

Received 27 September 2004

### Abstract

In an attempt to enhance the immunological efficacy of genetic immunization, we investigated a new biological means for delivering antigen gene directly to the cytoplasm via membrane fusion. In this context, we investigated fusogenic liposome (FL) encapsulating DNA as a possible genetic immunization vehicle. RT-PCR analysis indicated that a FL could introduce and express encapsulating OVA gene efficiently and rapidly *in vitro*. Consistent with this observation, an *in vitro* assay showed that FL-mediated antigen-gene delivery can induce potent presentation of antigen via the MHC class I-dependent pathway. Accordingly, immunization with FL containing the OVA-gene induced potent OVA-specific Th1 and Th2 cytokine production. Additionally, OVA-specific CTL responses and antibody production were also observed in systemic compartments including the spleen, upon immunization with the OVA-gene encapsulating FL. These findings suggest that FL is an effective genetic immunization carrier system for the stimulation of antigen-specific immune responses against its encoding antigen.

© 2004 Elsevier Inc. All rights reserved.

**Keywords:** DNA vaccine; Tumor vaccine; Liposome; Drug delivery system

Genetic immunization using plasmid DNA (pDNA) encoding antigens from bacteria, viruses, and cancers has often led to protective cell-mediated (Th1) and humoral (Th2) immunity [1]. This system has some practical advantages over conventional vaccines, such as safety, stability, cost-effectiveness for manufacturing, and the ease of modifying and customizing the gene sequence to produce the desired type of recombinant pro-

tein for expression *in vivo*. Although naked DNA vaccines (i.e., pDNA in saline) are effective in small animal models [2–4], recent results from large-animal and non-human primate studies have been disappointing due to sub-optimal immune responses, despite the use of multi-milligram doses of naked pDNA [5]. Therefore, the development of adjuvants and excipients to increase immune responses to DNA vaccines has become an active area of research. The potential of genetic immunization to exert an effective antigen-specific immune response is directly related to both the level of

\* Corresponding author. Fax: +81 6 6879 8176.

E-mail address: [nakagawa@phs.osaka-u.ac.jp](mailto:nakagawa@phs.osaka-u.ac.jp) (S. Nakagawa).

expression of the encoded protein and the immunomodulatory activity generated by DNA vaccine formulations [6–8].

Two main causes for the ineffectiveness of genetic immunization have been proposed. First, the pDNAs are generally unstable in vivo. In other words, most injected pDNAs are degraded by extracellular deoxyribonucleases in situ [9–11]. In addition, a limited amount of undegraded pDNA is taken up via endocytosis by neighboring cells around the injection site, including antigen presenting cells (APCs). The endocytosed pDNAs are thus generally degraded in intracellular compartments such as endosomes or lysosomes. Consequently, extremely low levels of pDNAs that escaped from these host factors could express their encoded antigens, resulting in inefficient induction of antigen-specific immunity. Furthermore, in terms of eliciting antibody responses, DNA vaccines are poorly immunogenic relative to other vaccines such as peptide and protein vaccines [12,13]. Because of the extra- and intra-cellular degradation and low immunogenicity of naked pDNAs, genetic immunization exhibits poor performance when administered by routes other than intramuscular [5].

In this context, various approaches, such as combination with adjuvant or cytokines and particulating techniques, have been applied to enhance immune responses to the encoding antigen [9,11]. The carrier-mediated approach is particularly promising because of enhanced pDNA stability and immunogenicity, and also due to the combination with molecular adjuvants. To address these issues, several DNA-particulating approaches have been evaluated [14]. Recently, liposomes and micro- or nano-scale particles have been tested for genetic vaccination [14–20]. However, due to the low immunogenicity and degradation of pDNA by APCs, a novel carrier-mediated approach to improve the potency of DNA vaccines is required.

We previously developed a unique antigen delivery system based on liposomes fused with UV-inactivated Sendai virus, known as the fusogenic liposome, FL [21–25]. The FL efficiently delivered the encapsulated antigens into the cytoplasm of antigen presenting cells via membrane fusion for use in the MHC class I-presentation pathway [26,27]. Furthermore, FL possesses adjuvant activity derived from Sendai virus accessory proteins. It stimulates antigen presenting cells to up-regulate cell surface markers and enhances the expression of inflammatory cytokines by APCs. Additionally, we demonstrated that subcutaneous immunization with antigen-encapsulated FL induced high levels of antigen-specific immune responses at systemic immune compartments such as the spleen [26].

This information and our previous results allowed us to hypothesize that FL could be utilized as genetic vaccine carrier. Thus, the purpose of the present study was

to analyze the efficiency of immunization through dermal delivery of model antigen protein-encoding DNA (OVA) delivered by FL.

## Materials and methods

**Mice and cells.** Six- to eight-week-old male C57BL/6 (H-2<sup>b</sup>) mice were purchased from SLC (Hamamatsu, Japan). EL4 (Tohoku University, Sendai, Japan) is a C57BL/6 T lymphoma and EG7 is an ovalbumin (OVA)-transfected clone of EL4. IC21 cells are C57BL/6 macrophage clones, H-2Kb, CD8OVA1.3 (provided by Dr. Clifford V. Harding, Case Western Reserve University, Cleveland, OH) is a T-T hybrid cell, which is specific for OVA257-264-Kb.

EL4 and IC21 cells were grown in RPMI1640 medium supplemented with 10% FCS. The CTL-2 cells were maintained in RPMI1640 medium supplemented with 10% FCS and 1 U/ml human recombinant IL-2. The EG7 cells were maintained in RPMI1640 medium supplemented with 10% FCS and 400 µg/ml G418. CD8OVA1.3 was grown in DMEM supplemented with 10% FCS. All culture media were purchased from Invitrogen (Carlsbad, CA) and supplemented with non-essential amino acids, antibiotics and,  $5 \times 10^{-5}$  M 2-mercaptoethanol (2-ME).

**Plasmids.** The EcoRI fragment of pAc-neo-OVA [28] was cloned into the EcoRI site of pBluescriptII KS(-), resulting in pBluescriptII KS(-)/OVA. To construct an OVA gene expression vector, the BamHI/SalI fragment of pBluescript II KS(-)/OVA was ligated into a BamHI/SalI cut pCMV-script (Stratagene), resulting in pCMV-script/OVA (Fig. 1), which is driven by the cytomegalovirus promoter and contains the SV40 poly(A) signal.

**Preparation of fusogenic liposome.** pCMV-script/OVA containing unilamellar liposomes was prepared by a modified reverse-phase evaporation method using 46 µmol of lipids (egg phosphatidylcholine: L- $\alpha$ -dimyristyl phosphatidic acid: cholesterol = 5:1:4, molar ratio). After three cycles of freezing and thawing, the liposomes were sized by extrusion through a 0.8 µm polycarbonate membrane (Nucleopore; Coaster, Cambridge, USA) and pelleted by ultracentrifugation to remove unencapsulated plasmids. Then, FL encapsulated pCMV-script/OVA was prepared by fusing the liposomes with UV (2000 J/cm<sup>2</sup>)-inactivated Sendai virus as described previously [21,26]. The amount of plasmid DNA encapsulated within liposomes was determined by means of fluorometric assay using 3,5-diaminobenzoic acid.

**RT-PCR for OVA gene detection.** To examine the expression of OVA mRNA, IC-21 cells were incubated in serum free RPMI1640




Fig. 1. Structure of OVA expression plasmid pCMV-script/OVA. OVA cDNA was inserted into an expression plasmid (pCMV-script) containing a cytomegalovirus (CMV) promoter and a SV40 polyadenylation site.






Fig. 3. Plasma clearance and tissue distribution of native SOD and PVD-SODs 3 h after intravenous injection. Mice were intravenously injected with  $^{125}\text{I}$ -labeled native SOD and PVD-SODs. (A) After administration, blood was collected from the tail at indicated time, and the radioactivity was counted. (B) Each organ was removed 3 h after intravenous injection, and radioactivity was counted. Data was represented at % injected dose. The radioactivity was measured by a  $\gamma$ -counter. Data represent mean  $\pm$  S.D. ( $n=5$ ).

in the supernatant of homogenized kidneys was measured after acid precipitation to distinguish between bound polymer and free tyramine, it has been confirmed that the PVD did not release the free tyramine and it was not degraded in the kidneys (data not shown). The blood retention increased as the molecular weight increased (Fig. 1A). On the other hand, PVD with an average molecular weight of 6–8 kDa [PVD<sub>6k</sub> and PVD<sub>8k</sub>] showed highest renal accumulation and about 80% of the administered dose accumulate in the kidneys at 3 h after injection (Fig. 1B). Accumulation rates decreased to 60% for PVD<sub>14k</sub> and to 30% for PVD<sub>3k</sub>. Table 1 summarizes the clearance calculated based on the radioactivity at 3 h after intravenous injection of various PVDs in mice. The uptake clearance of PVD<sub>6k</sub> were the highest among various PVDs. Similar trends in results were noted at 1 and 24 h (Fig. 2A). PVD<sub>6k</sub> and PVD<sub>8k</sub> were rapidly eliminated from blood and specifically accumulated in the kidneys only 1 h after intravenous injection without distribution to other tissues. In addition, PVD<sub>6k</sub> and PVD<sub>8k</sub> showed high retention in the kidneys and about 60% of injected dose retained in the kidneys 24 h after intravenous administration. By the measurement of the urinary radioactivity excretion, it became clear that the PVD which accumulated in the kidney was gradually excreted urinary. Furthermore, measurement of urinary radioactivity excretion

revealed a significantly higher value for PVD<sub>3k</sub> with the lowest  $M_n$  (Fig. 2B).

### 3.2. Characteristics of PVD-SOD

The PVD-SOD was prepared via formation of amide bond between SOD lysine residues and carboxyl groups of PVD<sub>6k</sub>. The resultant PVD-SOD was separated into three fraction of different molecular size (high=H, middle=M, low=L) by gel filtration HPLC, and then, specific activities were measured. The separated PVD-SODs, with molecular size of 73, 120, and 220 kDa, were termed L-PVD-SOD, M-PVD-SOD, H-PVD-SOD, respectively. Table 2 shows characterization of PVD-SODs. Although specific activity decreased with increasing molecular size, even H-PVD-SOD with the largest molecular size still had 60% activity compared with native SOD.

Table 3  
Clearance of native SOD and PVD-SODs

	Clearance ( $\mu\text{l/h}$ )			
	Kidney	Liver	Lung	Spleen
H-PVD-SOD	18.0 $\pm$ 0.3	20.1 $\pm$ 0.5	1.7 $\pm$ 0.2	1.1 $\pm$ 0.1
M-PVD-SOD	31.9 $\pm$ 1.2	20.9 $\pm$ 0.2	2.0 $\pm$ 0.1	1.1 $\pm$ 0.1
L-PVD-SOD	182.6 $\pm$ 9.3	20.3 $\pm$ 2.3	3.6 $\pm$ 0.3	1.4 $\pm$ 0.1
Native SOD	34.1 $\pm$ 4.9	19.9 $\pm$ 3.0	5.3 $\pm$ 0.9	1.4 $\pm$ 0.2

Mice were used in groups of five. Each value is the mean  $\pm$  S.E.

**Table 4**  
Therapeutic effects of L-PVD-SOD to HgCl<sub>2</sub>-induced ARF

		Intact mice	ARF mice	Native SOD-treated ARF mice	L-PVD-SOD-treated ARF mice
Urinary levels	hemoglobin	– (<0.06 mg/dl)	++ (>0.75 mg/dl)	+	–
	ketone	– (<5 mg/dl)	++ (>20 mg/dl)	±	–
	glucose	– (<100 mg/dl)	++ (>2000 mg/dl)	+	±
	protein	– (<10 mg/dl)	++ (>1000 mg/dl)	+	±
	γ-GTP	– (<0.7 IU/L.OG)	++ (>550 IU/L)	++	+
	ALP	– (<14 IU/l)	++ (>400 IU/l)	+	±
	NAG	– (<12 IU/l)	++ (>17 IU/l)	++	–
	creatinine	– (<0.5 mg/dl)	++ (>1.5 mg/dl)	+	±

We used a mouse model of HgCl<sub>2</sub>-induced ARF to assess the therapeutic effects of L-PVD-SOD. ARF was induced by subcutaneous injection of 8 mg/kg HgCl<sub>2</sub>, and then, these mice were given 4 mg protein/kg native SOD or L-PVD-SOD intravenously. Therapeutic efficacy was qualitatively assessed 48 h after treatment with HgCl<sub>2</sub> by measuring the level of urinary alkaline phosphatase (ALP), urinary γ-glutamyl transferase (GTP), urinary *N*-acetyl-β-glucosaminidase (NAG), urinary glucose, urinary keton, urinary protein, urinary hemoglobin, and serum creatinine as ARF markers. Mice were used in groups of five. –: negative; ±: quasi-positive; +: weak positive; ++: strong positive.

### 3.3. Pharmacokinetics of PVD-SOD

We then evaluated pharmacokinetics of three kinds of PVD-SODs after intravenous administration. Native SOD was rapidly cleared from the blood circulation (Fig. 3A). Three hours after injection, little accumulation of native SOD into the kidneys was observed (Fig. 3B), and almost all native SOD was found to be eliminated in the urine (data not shown). On the other hand, the blood residency and renal distribution of PVD-SOD increased with decreasing their molecular size. For L-PVD-SOD with almost full activity compared with native SOD, its renal accumulation was about six times higher than that of native SOD, and L-PVD-SOD did not show with selective distribution to other major organs such as the liver or spleen. M-PVD-SOD and H-PVD-SOD showed higher distribution to liver than native SOD and L-PVD-SOD, probably due to their high blood concentration. Table 3 summarizes the clearance calculated based on the radioactivity at 3 h after intravenous injection of native SOD and three kinds of PVD-SODs in mice. This data also showed that L-PVD-SOD accumulated in kidney specifically.

### 3.4. Therapeutic effect of L-PVD-SOD on ARF

We also assessed the therapeutic effect of L-PVD-SOD on ARF (Table 4). ARF was induced by subcutaneous injection of HgCl<sub>2</sub> at a dose of 8 mg/

kg. Not only the levels of urinary ALP, γ-GTP, NAG, and serum creatinine, but also the urinary content of hemoglobin, keton, glucose, and protein rapidly increased 12 h later. Native SOD and L-PVD-SOD at a dose of 4 mg protein/kg were injected intravenously 12 h after injection of HgCl<sub>2</sub>. The therapeutic efficacy were assessed 48 h after administration of HgCl<sub>2</sub> (Table 4), because ARF markers reached the highest levels in untreated ARF mice. Native SOD showed weak therapeutic effects, but L-PVD-SOD effectively accelerated recovery from ARF.

## 4. Discussion

The *in vivo* pharmacokinetics of polymer-conjugated drugs, such as bioactive proteins, may be markedly influenced by the properties of the polymeric carriers, such as electric charge and hydrophilic/hydrophobic balances, attached to the surface of the drugs. Therefore, to achieve optimization of drug therapy by polymer conjugation typified by PEGylation, we must initially design polymeric carrier with useful functions such as targeting and controlled release capability, which can regulate closely their behavioral characteristics *in vivo*. We previously reported that polyvinylpyrrolidone (PVP) was more suitable polymeric carrier for enhancing the blood residency of drugs than PEG [8]. Using this PVP as a backbone polymer, we have evaluated

the *in vivo* pharmacokinetics of synthesized PVP derivatives with various electrical charge or hydrophilic/hydrophobic balance [20]. For instance, the co-polymer between vinylpyrrolidone (VP) and vinyl-lauric acid showed a marked increase in distribution in the spleen compared to that in the liver. Recently, we synthesized a novel polymeric drug carrier, PVD, which was a powerful candidate as a targeting carrier for a renal drug delivery system. In this study, we attempted to optimize the renal drug delivery system using PVD.

To assess the usefulness of PVD as a renal targeting carrier, the relationship between the  $M_n$  of PVD and its renal accumulation after intravenous injection was investigated (Fig. 1). Renal accumulation was highest for molecules with an  $M_n$  of 6–8 kDa, about 80% of the administered dose accumulate in the kidneys 3 h after intravenous injection. Increase or decrease of an  $M_n$  lowered the renal accumulation (Fig. 2). We have previously showed that the safety of PVD was similar to that of PEG and PVP, which are used clinically [17]. Thus, PVD with 6–8 kDa of  $M_n$  seems to be an extremely safe polymeric carrier with much higher renal targeting and retention capacity. It has previously been found that PVD was selectively accumulated in renal proximal tubular epithelial cells after intravenous injection [17]. Additionally, the uptake of PVD by renal tubular cells was inhibited by the energy inhibitor (NaN<sub>3</sub>) and was not affected by cytochalasin B. Thus, PVD may be taken in by an energy-dependent process except for endocytosis. Several specific molecules play a role in renal transport, and various organic anions transporter exist in the kidney, but these transporters generally carry low-molecular-weight drugs. It is unclear why PVD with a molecular weight of 6–8 kDa was selectively accumulated in the kidneys. To address this question, we are currently in the process of doing some experiments.

We further evaluated the usefulness of PVD as a renal targeting carrier by polymer conjugation to SOD, which has been expected to be a potential drug to renal disease. Many recent studies have reported an association between activated oxygen species such as superoxide radical (O<sub>2</sub><sup>-</sup>), hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), hydroxyl radical (·OH), and NO with various pathologic disease processes such as

cancer, inflammation, septicemia, and necrosis associated with ischemic reperfusion [21,22]. Several studies have investigated the use of activated oxygen metabolic enzymes and antioxidants as therapeutic agents in diseases where stress oxidation plays a prominent role [23,24]. SOD has shown promise as a therapeutic agent, eliminating O<sub>2</sub><sup>-</sup> in the early stages of formation of highly reactive oxygen species such as ·OH. Developments in genetic engineering have now enabled the production of large quantities of human Cu/Zn-SOD, which has attracted attention as a therapeutic agent [25]. With respect to kidney disease, activated oxygen is known to play an indispensable role in the mechanisms of ARF, complications associated with long-term maintenance dialysis, drug toxicity, and various inflammatory conditions [26]. We prepared PVD-SODs with various molecular size (Table 2) and then evaluated their tissue distribution 3 h after intravenous injection (Fig. 3). The renal accumulation of PVD-SODs decreased with increasing their molecular size. About sixfold more L-PVD-SOD with 73 kDa of  $M_n$ , which had almost full activity compared with native SOD, was distributed to the kidneys than native SOD. The renal accumulation of PVP-modified SOD with the same molecular size as L-PVD-SOD was similar to that of native SOD (data not shown). This confirms that renal accumulation of L-PVD-SOD is attributable to the properties of PVD. However, extensive PVD modification of SOD did not increase its renal accumulation. These results indicated that L-PVD-SOD was the optimal derivative which had a potential renal targeting capability. Thus, our results revealed that the renal accumulation probably involves not only an optimal introduction rate of electrical charges and/or hydrophobicity to PVD, but also molecular size of PVD-SOD.

Renal disease is one of the most serious problems in the world. Many researchers have attempted to deliver drug to the kidney. For example, Hashida et al. reported that cationized SOD and PEGylated SOD exhibited significant therapeutic effects on ischemic acute renal failure [27,28]. However, there is no report as to delivery of drug to the kidney specifically. On the other hand, PVD accumulated in the kidney about 80% of the administered dose. Therefore, we consider that PVD may be the most superior carrier for delivering drug to the kidney. Thus, the development

of new therapeutic approaches is expected for treating patients with renal disease. As shown in Table 4, L-PVD-SOD showed great potential as a renal antioxidant agent to ARF. These results suggest that L-PVD-SOD may be a candidate as a novel therapeutic agent with high renal targeting capability.

### Acknowledgements

This study was supported in part by a Grant-in-Aid for Scientific Research (No. 15680014) from the Ministry of Education, Science and Culture of Japan, in part by Health Sciences Research Grants for Research on Health Sciences focusing on Drug Innovation from the Japan Health Sciences Foundation (KH63124), and in part by Takeda Science Foundation.

### References

- [1] M. Blick, S.A. Sherwin, M. Rosenblum, J. Gutterman, Phase I study of recombinant tumor necrosis factor in cancer patients, *Cancer Res.* 47 (1987) 2986–2989.
- [2] S.A. Rosenberg, M.T. Lotze, L.M. Muul, A.E. Chang, F.P. Avis, S. Leitman, W.M. Linchan, C.N. Robertson, R.E. Lee, J.T. Rubin, et al., A progress report on the treatment of 157 patients with advanced cancer using lymphokine-activated killer cells and interleukin-2 or high-dose interleukin-2 alone, *N. Engl. J. Med.* 316 (1987) 889–897.
- [3] J.H. Donohue, S.A. Rosenberg, The fate of interleukin-2 after in vivo administration, *J. Immunol.* 130 (1983) 2203–2208.
- [4] A.P. Bollon, S.L. Berent, R.M. Torczynski, N.O. Hill, Y. Lemeshev, J.M. Hill, F.L. Jia, A. Joher, S. Pichyangkul, A. Khan, Human cytokines, tumor necrosis factor, and interferons: gene cloning, animal studies, and clinical trials, *J. Cell. Biochem.* 36 (1988) 353–367.
- [5] H. Tanaka, T. Tokiwa, Influence of renal and hepatic failure on the pharmacokinetics of recombinant human granulocyte colony-stimulating factor (KRN8601) in the rat, *Cancer Res.* 50 (1990) 6615–6619.
- [6] Y. Tsutsumi, M. Onda, S. Nagata, B. Lee, R.J. Kreitman, I. Pastan, Site-specific chemical modification with polyethylene glycol of recombinant immunotoxin anti-Tac(Fv)-PE38 (LMB-2) improves antitumor activity and reduces animal toxicity and immunogenicity, *Proc. Natl. Acad. Sci. U. S. A.* 97 (2000) 8548–8553.
- [7] Y. Kaneda, Y. Yamamoto, H. Kamada, S. Tsunoda, Y. Tsutsumi, T. Hirano, T. Mayumi, Antitumor activity of tumor necrosis factor- $\alpha$  conjugated with divinyl ether and maleic anhydride copolymer on solid tumors in mice, *Cancer Res.* 58 (1998) 290–295.
- [8] H. Kamada, Y. Tsutsumi, Y. Yamamoto, T. Kihira, Y. Kaneda, Y. Mu, H. Kodaira, S. Tsunoda, S. Nakagawa, T. Mayumi, Antitumor activity of tumor necrosis factor- $\alpha$  conjugated with polyvinylpyrrolidone on solid tumors in mice, *Cancer Res.* 60 (2000) 6416–6420.
- [9] Y. Yamamoto, Y. Tsutsumi, Y. Yoshioka, T. Nishibata, K. Kobayashi, T. Okamoto, Y. Mukai, T. Shimizu, S. Nakagawa, S. Nagata, T. Mayumi, Site-specific PEGylation of a lysine-deficient TNF- $\alpha$  with full bioactivity, *Nat. Biotechnol.* 21 (2003) 546–552.
- [10] Y. Tsutsumi, S. Tsunoda, H. Kamada, T. Kihira, Y. Kaneda, Y. Ohsugi, T. Mayumi, PEGylation of interleukin-6 effectively increases its thrombopoietic potency, *Thromb. Haemost.* 77 (1997) 168–173.
- [11] C.A. Jones, G.M. McQuillan, J.W. Kusek, M.S. Eberhardt, W.H. Herman, J. Coresh, M. Salive, C.P. Jones, L.Y. Agodoa, Serum creatinine levels in the US population: third National Health and Nutrition Examination Survey, *Am. J. Kidney Dis.* 32 (1998) 992–999.
- [12] F.N. Ziyadeh, S. Goldfarb, The diabetic renal tubulointerstitium, *Curr. Top. Pathol.* 88 (1995) 175–201.
- [13] A.A. Elfarra, et al., Targeting 6-thioguanine to the kidney with S-(guanin-6-yl)-L-cysteine, *J. Pharmacol. Exp. Ther.* 274 (1995) 1298–1304.
- [14] R.F. Haverdings, et al., Potentials and limitations of the low-molecular-weight protein lysozyme as a carrier for renal drug targeting, *Renal Failure* 23 (2001) 397–409.
- [15] B. Schechter, R. Amon, C. Colas, T. Burakova, M. Wilchek, Renal accumulation of streptavidin: potential use for targeted therapy to the kidney, *Kidney Int.* 47 (1995) 1327–1335.
- [16] R.J. Kok, F. Grijpstra, R.B. Walthuis, F. Moolenaar, D. de Zeeuw, D.K. Meijer, Specific delivery of captopril to the kidney with the prodrug captopril-lysozyme, *J. Pharmacol. Exp. Ther.* 288 (1999) 281–285.
- [17] H. Kamada, Y. Tsutsumi, K. Sato-Kamada, Y. Yamamoto, Y. Yoshioka, T. Okamoto, S. Nakagawa, S. Nagata, T. Mayumi, Synthesis of a poly(vinylpyrrolidone-co-dimethyl maleic anhydride) co-polymer and its application as renal targeting carrier, *Nat. Biotechnol.* 21 (2003) 399–404.
- [18] M. Nishikawa, Y. Takakura, M. Hashida, Pharmacokinetic evaluation of polymeric carriers, *Adv. Drug Deliv. Rev.* 21 (1996) 135–155.
- [19] M. Nishikawa, T. Nakano, T. Okabe, N. Hamaguchi, Y. Yamasaki, Y. Takakura, F. Yamashita, M. Hashida, Residualizing indium-111-radiolabel for plasmid DNA and its application to tissue distribution study, *Bioconjug. Chem.* 14 (2003) 955–961.
- [20] H. Kodaira, Y. Kaneda, Y. Yamamoto, T. Namba, Y. Tsutsumi, T. Hirano, T. Mayumi, Fundamental study on molecular design of bioconjugated drugs with water-soluble polymeric modifiers; influence of electric charge on pharmacokinetics of water-soluble polymers, *Drug Deliv. Syst.* 12 (1997) 431–437.
- [21] A. Hamvas, R. Palazzo, L. Kaiser, J. Cooper, T. Shuman, M. Velazquez, B. Freeman, D.P. Schuster, Inflammation and oxygen free radical formation during pulmonary ischemia-reperfusion injury, *J. Appl. Physiol.* 72 (1992) 621–628.