

	Potentially Significant Drug Interactions: Alterations in Dose or Regimen May Be Recommended Based on Drug Interaction Studies or Predicted Interaction.
Neuroleptic: pimozide	CONTRAINDICATED due to the potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Sedative/Hypnotics: midazolam, triazolam	CONTRAINDICATED due to potential for serious and/or life-threatening reactions such as prolonged or increased sedation or respiratory depression.

**Table 11: Established and Other Potentially Significant Drug Interactions:
Alterations in Dose or Regimen May Be Recommended
Based on Drug Interaction Studies or Predicted Interaction
(See CLINICAL PHARMACOLOGY for Magnitude of Interaction, Tables 4 and 5)**

Concomitant Drug Class: Drug Name	Effect on Concentration of Darunavir or Concomitant Drug	Clinical Comment
HIV-Antiviral Agents: Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs)		
Efavirenz	↓ darunavir ↑ efavirenz	Co-administration of darunavir/rtv and efavirenz decreased darunavir AUC by 13% and C _{min} by 31%. The AUC of efavirenz increased by 21% and C _{min} increased by 17%. The clinical significance has not been established. The combination of PREZISTA/rtv and efavirenz should be used with caution.
Nevirapine	↔ darunavir ↑ nevirapine	PREZISTA/rtv and nevirapine can be co-administered without any dose adjustments.
HIV-Antiviral Agents: Nucleoside Reverse Transcriptase Inhibitors (NRTIs)		
Didanosine		It is recommended that didanosine be administered on an empty stomach. Therefore, didanosine should be administered one hour before or two hours after PREZISTA/rtv (which are administered with food).
Tenofovir Disoproxil Fumarate	↔ darunavir ↑ tenofovir	PREZISTA/rtv and tenofovir disoproxil fumarate can be co-administered without any

		dose adjustments.
HIV-Antiviral Agents: HIV-Protease Inhibitors (PIs)		
Atazanavir (The reference regimen for atazanavir was atazanavir/ritonavir 300/100 mg q.d.)	↔ darunavir ↔ atazanavir	PREZISTA/rtv and atazanavir (300 mg q.d.) can be co-administered.
Indinavir (The reference regimen for indinavir was indinavir/ritonavir 800/100 mg b.i.d.)	↑ darunavir ↑ indinavir	The appropriate dose of indinavir in combination with PREZISTA/rtv has not been established.
Lopinavir/ritonavir	↓ darunavir ↑ lopinavir	Due to decrease in the exposure (AUC) of darunavir by 53%, appropriate doses of the combination have not been established. Hence, it is not recommended to co-administer lopinavir/ritonavir and PREZISTA, with or without an additional low-dose of ritonavir.
Saquinavir	↓ darunavir ↔ saquinavir	Due to a decrease in the exposure (AUC) of darunavir by 26%, appropriate doses of the combination have not been established. Hence, it is not recommended to co-administer saquinavir and PREZISTA, with or without low-dose ritonavir.
Other Agents		
Antiarrhythmics: bepridil, lidocaine (systemic), quinidine, amiodarone	↑ antiarrhythmics	Concentrations of bepridil, lidocaine, quinidine and amiodarone may be increased when co-administered with

		PREZISTA/rtv. Caution is warranted and therapeutic concentration monitoring, if available, is recommended for antiarrhythmics when co-administered with PREZISTA/rtv.
Anticoagulant: warfarin	↓ warfarin ↔ darunavir	Warfarin concentrations may be affected when co-administered with PREZISTA/rtv. It is recommended that the international normalized ratio (INR) be monitored when warfarin is combined with PREZISTA/rtv.
Antidepressant: trazodone	↑ trazodone	Concomitant use of trazodone and PREZISTA/rtv may increase plasma concentrations of trazodone. Adverse events of nausea, dizziness, hypotension and syncope have been observed following co-administration of trazodone and ritonavir. If trazodone is used with a CYP3A inhibitor such as PREZISTA/rtv, the combination should be used with caution and a lower dose of trazodone should be considered.
Anti-infective: clarithromycin	↑ clarithromycin	No dose adjustment of darunavir or clarithromycin is required for patients with normal renal function. For patients with renal impairment, the

		<p>following dose adjustments should be considered:</p> <ul style="list-style-type: none"> • For subjects with CLcr of 30-60 mL/min, the dose of clarithromycin should be reduced by 50%. • For subjects with CLcr of < 30 mL/min, the dose of clarithromycin should be reduced by 75%.
<p>Antifungals: ketoconazole, itraconazole, voriconazole</p>	<p>↑ ketoconazole ↑ darunavir ↑ itraconazole (not studied) ↓ voriconazole (not studied)</p>	<p>Ketoconazole and itraconazole are potent inhibitors as well as substrates of CYP3A. Concomitant systemic use of ketoconazole, itraconazole, and darunavir/ritonavir may increase plasma concentration of darunavir.</p> <p>Plasma concentrations of ketoconazole or itraconazole may be increased in the presence of darunavir/ritonavir. When co-administration is required, the daily dose of ketoconazole or itraconazole should not exceed 200 mg.</p> <p>Co-administration of voriconazole with darunavir/ritonavir has not been studied. Administration of</p>

		<p>voriconazole with ritonavir (100 mg twice daily) decreased the AUC of voriconazole by an average of 39%. Voriconazole should not be administered to patients receiving darunavir/ritonavir unless an assessment of the benefit/risk ratio justifies the use of voriconazole.</p>
<p>Antimycobacterial: rifabutin</p>	<p>↑ rifabutin ↓ darunavir</p>	<p>Rifabutin is an inducer and substrate of CYP450 enzymes. Concomitant use of rifabutin and darunavir in the presence of ritonavir is expected to increase rifabutin plasma concentrations and decrease darunavir plasma concentrations. When indicated, it is recommended to administer rifabutin at a dosage of 150 mg once every other day when co-administered with PREZISTA/rtv.</p>
<p>Calcium Channel Blockers: felodipine, nifedipine, nicardipine</p>	<p>↑ calcium channel blockers</p>	<p>Plasma concentrations of calcium channel blockers (e.g. felodipine, nifedipine, nicardipine) may increase when PREZISTA/rtv are co-administered. Caution is warranted and clinical monitoring of patients is recommended.</p>
<p>Corticosteroid: dexamethasone fluticasone propionate</p>	<p>↓ darunavir ↑ fluticasone propionate</p>	<p>Use with caution. Systemic dexamethasone induces CYP3A and can thereby decrease darunavir</p>

		<p>plasma concentrations. This may result in loss of therapeutic effect to PREZISTA. Concomitant use of inhaled fluticasone propionate and PREZISTA/rtv may increase plasma concentrations of fluticasone propionate. Alternatives should be considered, particularly for long term use.</p>
<p>HMG-CoA Reductase Inhibitors: atorvastatin, pravastatin</p>	<p>↑ atorvastatin ↑ pravastatin</p>	<p>When atorvastatin and PREZISTA/rtv is co-administered, it is recommended to start with the lowest possible dose of atorvastatin with careful monitoring. A gradual dose increase of atorvastatin may be considered based on the clinical response.</p> <p>When PREZISTA/rtv was administered with pravastatin, the mean increase in pravastatin AUC was 81%. However, pravastatin AUC increased by up to 5-fold in some subjects. The mechanism of the interaction is not known.</p>
<p>H2-Receptor Antagonists and Proton Pump Inhibitors: omeprazole, ranitidine</p>	<p>↔ darunavir</p>	<p>PREZISTA/rtv can be co-administered with H2-receptor antagonists and proton pump inhibitors without any dose adjustments.</p>
<p>Immunosuppressants: cyclosporine,</p>	<p>↑ immunosuppressants</p>	<p>Plasma concentrations of cyclosporine, tacrolimus or</p>

<p>tacrolimus, sirolimus</p>		<p>sirolimus may be increased when co-administered with PREZISTA/rtv. Therapeutic concentration monitoring of the immunosuppressive agent is recommended for immunosuppressant agents when co-administered with PREZISTA/rtv.</p>
<p>Narcotic Analgesic: methadone</p>	<p>↓ methadone</p>	<p>When methadone is co-administered with PREZISTA/rtv, patients should be monitored for opiate abstinence syndrome, as ritonavir is known to induce the metabolism of methadone, leading to a decrease in its plasma concentrations. An increase in methadone dosage may be considered based on the clinical response.</p>
<p>Oral Contraceptives/estrogen: ethinyl estradiol norethindrone</p>	<p>↓ ethinyl estradiol ↓ norethindrone</p>	<p>Plasma concentrations of ethinyl estradiol may be decreased due to induction of its metabolism by ritonavir. Alternative or additional contraceptive measures should be used when estrogen-based contraceptives are co-administered with PREZISTA/rtv.</p>
<p>PDE-5 inhibitors: sildenafil, vardenafil, tadalafil</p>	<p>↑ PDE-5 inhibitors</p>	<p>Concomitant use of PDE-5 inhibitors with PREZISTA/rtv should be done with caution. If concomitant use of PREZISTA/rtv with sildenafil, vardenafil, or</p>

		tadalafil is required, sildenafil at a single dose not exceeding 25 mg in 48 hours, vardenafil at a single dose not exceeding 2.5 mg dose in 72 hours, or tadalafil at a single dose not exceeding 10 mg dose in 72 hours, is recommended.
Selective Serotonin Reuptake Inhibitors (SSRIs): sertraline, paroxetine	↔ darunavir ↓ sertraline ↓ paroxetine	If sertraline or paroxetine is co-administered with PREZISTA/rtv, the recommended approach is a careful dose titration of the SSRI based on a clinical assessment of antidepressant response. In addition, patients on a stable dose of sertraline or paroxetine who start treatment with PREZISTA/rtv should be monitored for antidepressant response.

Other NRTIs:

Based on the different elimination pathways of the other NRTIs (zidovudine, zalcitabine, emtricitabine, stavudine, lamivudine and abacavir) that are primarily renally excreted, no drug interactions are expected for these drugs and PREZISTA/rtv.

Other protease inhibitors:

The co-administration of PREZISTA/rtv and PIs other than lopinavir/ritonavir, saquinavir, atazanavir, and indinavir has not been studied. Therefore, such co-administration is not recommended.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis and Mutagenesis:

Long-term carcinogenicity studies of darunavir in rodents have not been completed. Darunavir, however, was tested negative in the *in vitro* Ames reverse mutation assay and *in vitro* chromosomal aberration assay in human lymphocytes, both tested in the absence and presence of metabolic activation system. Darunavir does not induce chromosomal damage in the *in vivo* micronucleus test in mice.

Impairment of Fertility:

There were no effects on fertility and early embryonic development with darunavir in rats and darunavir has shown no teratogenic potential in mice (in the presence or absence of ritonavir), rats and rabbits.

Pregnancy

Pregnancy Category B: Reproduction studies conducted with darunavir have shown no embryotoxicity or teratogenicity in mice, rats and rabbits. Because of limited bioavailability of darunavir in animals and/or dosing limitations, the plasma exposures (AUC values) were approximately 50% in mice and rats and 5% in the rabbit of those obtained in humans at the recommended clinical dose boosted with ritonavir.

In the rat pre- and postnatal development study, a reduction in pup body weight gain was observed with darunavir alone or in combination with ritonavir during lactation. This was due to exposure of pups to drug substances via the milk. Sexual development, fertility or mating performance of offspring was not affected by maternal treatment with darunavir alone or in combination with ritonavir. The maximal plasma exposures achieved in rats were approximately 50% of those obtained in humans at the recommended clinical dose boosted with ritonavir.

There are, however, no adequate and well-controlled studies in pregnant women. PREZISTA should be used during pregnancy only if the potential benefit justifies the potential risk.

Antiretroviral Pregnancy Registry: *To monitor maternal-fetal outcomes of pregnant women exposed to PREZISTA, an Antiretroviral Pregnancy Registry has been established. Physicians are encouraged to register patients by calling 1-800-258-4263.*

Nursing Mothers

The Centers for Disease Control and Prevention recommend that HIV-infected mothers not breastfeed their infants to avoid risking postnatal transmission of HIV. Although it is not known whether darunavir is secreted in human milk, darunavir is secreted into the milk of lactating rats. Because of both the potential for HIV transmission and the potential for serious adverse reactions in nursing infants, **mothers should be instructed not to breastfeed if they are receiving PREZISTA.**

Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

Geriatric Use

Clinical studies of PREZISTA did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. In general, caution should be exercised in the administration and monitoring of PREZISTA in elderly patients reflecting the greater frequency of decreased hepatic function, and of concomitant disease or other drug therapy.

ADVERSE REACTIONS

The safety assessment is based on all safety data from the Studies TMC114-C213 and TMC114-C202 and the TMC114-C215/C208 analysis reported with the recommended dose PREZISTA/rtv 600/100 mg b.i.d. in the 458 subjects who initiated treatment with the

recommended dose (*de novo* subjects). In Studies TMC114-C213 and TMC114-C202, the mean exposure in weeks for subjects in the PREZISTA/rtv 600/100 mg b.i.d. arm and comparator PI arm was 63.5 and 31.5, respectively. The mean exposure in weeks for subjects in the TMC114-C215/C208 analysis was 23.9.

The most common treatment-emergent adverse events (> 10%) reported in the *de novo* subjects, regardless of causality or frequency, were diarrhea, nausea, headache, and nasopharyngitis.

For subjects in the PREZISTA/rtv 600/100 mg b.i.d. arm and the comparator PI arm in the pooled analysis for Studies TMC114-C213 and TMC114-C202, diarrhea was reported in 19.8% and 28.2%, nausea in 18.3% and 12.9%, headache in 15.3% and 20.2%, and nasopharyngitis in 13.7% and 10.5%, of subjects, respectively. In the randomized trials, rates of discontinuation of therapy due to adverse events were 9% in subjects receiving PREZISTA/rtv and in 5% of subjects in the comparator PI arm.

Due to the need for co-administration of PREZISTA with 100 mg of ritonavir, please refer to ritonavir prescribing information for ritonavir-associated adverse reactions.

Drug-related clinical adverse events of moderate or severe intensity (\geq Grade 2) occurring in \geq 2% of subjects treated with PREZISTA/rtv for 1 to 96 weeks are presented in Table 12.

Table 12: Percentage of Subjects with Selected Treatment Emergent, Drug-Related[^] Adverse Events of at least Moderate Intensity (Grades 2-4) in \geq 2% of Adult Subjects in Any PREZISTA/rtv Treatment Groups[†]			
System Organ Class, Preferred Term, %	Randomized Studies TMC114-C213 and TMC114-C202		Non-randomized TMC114-C215/C208 Analysis
	PREZISTA/rtv 600/100 mg b.i.d. +OBR N = 131	Comparator PI +OBR N = 124	PREZISTA/rtv 600/100 mg b.i.d. +OBR N = 327
Gastrointestinal Disorders			
Diarrhea	2.3%	3.2%	2.8%
Vomiting	1.5%	1.6%	2.4%
Abdominal Pain	2.3%	0.8%	1.2%
Constipation	2.3%	0.8%	0.6%
Nervous System Disorders			
Headache	3.8%	2.4%	0.9%

[^] Includes adverse events at least possibly, probably, or very likely related to the drug
N=total number of subjects per treatment group
[†] Excludes laboratory abnormalities that were reported as Adverse Events (see Table 13: Treatment Emergent Grade 2 to 4 Laboratory Abnormalities Reported in \geq 2% of Subjects)

Treatment-emergent adverse events occurring in less than 2% of *de novo* subjects (n=458) receiving PREZISTA/rtv, considered at least possibly related to treatment and of at least moderate intensity are listed below by body system:

Body as a Whole:

folliculitis, asthenia, pyrexia, fatigue, rigors, hyperthermia, peripheral edema

Cardiovascular System:

myocardial infarction, tachycardia, hypertension

Digestive System:

flatulence, abdominal distension, dry mouth, dyspepsia, abdominal pain, nausea, constipation

Metabolic and Nutritional Disorders:

anorexia, hypercholesterolemia, hyperlipidemia, diabetes mellitus, decreased appetite, obesity, fat redistribution, hyponatremia, polydipsia

Musculoskeletal System:

arthralgia, pain in extremity, myalgia, osteopenia, osteoporosis

Nervous System:

peripheral neuropathy, hypoesthesia, memory impairment, paresthesia, somnolence, transient ischemic attack, confusional state, disorientation, irritability, altered mood, nightmare, anxiety, headache

Respiratory System:

dyspnea, cough, hiccups

Skin and Appendages:

lipoatrophy, night sweats, allergic dermatitis, eczema, toxic skin eruption, alopecia, dermatitis medicamentosa, hyperhidrosis, skin inflammation, maculopapular rash, erythema multiforme, Stevens-Johnson Syndrome (reported in another ongoing clinical study)

Special Senses:

vertigo

Urogenital System:

acute renal failure, renal insufficiency, nephrolithiasis, polyuria, gynecomastia

Laboratory abnormalities:

The percentages of adult subjects treated with PREZISTA/rtv 600/100 mg b.i.d. with treatment-emergent Grade 2 to 4 laboratory abnormalities are presented in Table 13.

Table 13: Treatment Emergent Grade 2 to 4 Laboratory Abnormalities Reported in $\geq 2\%$ of Subjects				
		Randomized Studies TMC114-C213 and TMC114-C202		Non- randomized TMC114- C215/C208 Analysis
Laboratory Parameter Preferred Term, %	Limit	PREZISTA/ rtv 600/100 mg b.i.d. + OBR N = 131	Comparator PI + OBR N = 124	PREZISTA/ rtv 600/100 mg b.i.d. N = 327
Biochemistry				
Aspartate Aminotransferase	> 2.5 X ULN	10.0%	13.0%	5.3%
Alanine Aminotransferase	> 2.5 X ULN	6.9%	9.8%	5.6%
Gamma Glutamyl Transferase	> 2.5 X ULN	9.2%	8.9%	8.4%
Hyperbilirubinemia	> 1.5 X ULN	2.3%	15.4%	0.9%
Alkaline Phosphatase	> 2.5 X ULN	4.6%	0%	2.8%
Pancreatic Amylase	> 1.5 X ULN	16.9%	8.9%	10.8%
Pancreatic Lipase	> 1.5 X ULN	8.5%	4.1%	6.2%
Hyperglycemia	≥ 161 mg/dL	2.3%	8.1%	5.9%
Hypoglycemia	≤ 54 mg/dL	1.5%	1.6%	3.7%
Total Cholesterol	≥ 240 mg/dL	9.2%	3.3%	8.0%
Triglycerides	> 400 mg/dL	25.4%	26.0%	18.9%
Hypoalbuminemia	< 3 g/dL	3.1%	1.6%	4.3%
Hyperuricemia	≥ 9.9 mg/dL	6.9%	6.5%	2.2%
Bicarbonate	< 15 mmol/L	3.1%	4.1%	3.4%
Hypocalcemia	≤ 7.8 mg/dL	0%	0.8%	4.0%
Hyponatremia	≤ 129 meq/L	0.8%	0%	2.5%
Hypematremia	≥ 151 meq/L	2.3%	0%	0%
Hematology				
White Blood Cell Count decrease	< 3000 count/mm ³	15.4%	18.7%	13.0%
Total Absolute Neutrophil Count decrease	≤ 999 mm ³	6.9%	9.8%	11.5%
Lymphocytes decrease	< 1000 count/mm ³	4.6%	19.5%	10.9%
Partial Thromboplastin	> 1.66 X ULN	7.8%	4.1%	4.3%

Time increase				
Plasma Prothrombin Time increase	> 1.25 X ULN	3.9%	0.8%	0.6%
Platelet Count decrease	< 75,000/mm ³	3.1%	1.6%	2.8%

Patients co-infected with hepatitis B and/or hepatitis C virus:

Subjects co-infected with hepatitis B or C virus receiving PREZISTA/rtv, did not experience higher incidence of adverse events or clinical chemistry abnormalities than subjects receiving PREZISTA/rtv who were not co-infected. The pharmacokinetic exposure in co-infected subjects was comparable to that in subjects without co-infection. Standard clinical monitoring of patients with chronic hepatitis B and/or C is considered adequate.

OVERDOSAGE

Human experience of acute overdose with PREZISTA/rtv is limited. Single doses up to 3200 mg of the oral solution of darunavir alone and up to 1600 mg of the tablet formulation of darunavir in combination with ritonavir have been administered to healthy volunteers without untoward symptomatic effects.

There is no specific antidote for overdose with PREZISTA. Treatment of overdose with PREZISTA consists of general supportive measures including monitoring of vital signs and observation of the clinical status of the patient. If indicated, elimination of unabsorbed active substance is to be achieved by emesis or gastric lavage. Administration of activated charcoal may also be used to aid in removal of unabsorbed active substance. Since PREZISTA is highly protein bound, dialysis is unlikely to be beneficial in significant removal of the active substance.

DOSAGE AND ADMINISTRATION

Adults: The recommended oral dose of PREZISTA tablets is 600 mg (two 300 mg tablets) twice daily taken with ritonavir 100 mg twice daily and with food. The type of food does not affect exposure to darunavir.

Pediatric Patients: The safety and efficacy of PREZISTA in pediatric patients has not been established (see CLINICAL PHARMACOLOGY, *Special Populations, Pediatric Patients*).

Hepatic Impairment: There are no data regarding the use of PREZISTA/rtv when co-administered to patients with varying degrees of hepatic impairment; therefore, specific dosage recommendations cannot be made. PREZISTA/rtv should be used with caution in patients with hepatic impairment (see CLINICAL PHARMACOLOGY, *Pharmacokinetics in Adults, Special Populations, Hepatic Impairment* and PRECAUTIONS, *Patients with co-existing conditions, Hepatic Impairment*).

Renal Impairment: No dose adjustment is required in patients with moderate renal impairment. There are no pharmacokinetic data available in HIV-1 infected patients with severe renal impairment or end stage renal disease (see CLINICAL PHARMACOLOGY, *Pharmacokinetics*).

in Adults, Special Populations, Renal Impairment and PRECAUTIONS, Patients with co-existing conditions, Renal Impairment).

HOW SUPPLIED

PREZISTA (darunavir) tablets are supplied as orange, oval-shaped, film-coated tablets containing darunavir ethanolate equivalent to 300 mg of darunavir per tablet. Each tablet is debossed with "300" on one side and "TMC114" on the other side. PREZISTA tablets are packaged in bottles in the following configuration:

300 mg tablets—bottles of 120 (NDC 59676-560-01)

Storage:

Store PREZISTA tablets at 25°C (77°F); with excursions permitted to 15°-30°C (59°-86°F).

Manufactured for Tibotec, Inc. by:

JOLLC, Gurabo, Puerto Rico

Distributed by:

Tibotec Therapeutics, Division of Ortho Biotech Products, L.P., Raritan NJ 08869

Issued: June 2006

Patent Numbers: 5,843,946; 6,248,775; 6,335,460 and other US patents pending

10101700

PREZISTA™* (darunavir) Tablets

Patient Information about
PREZISTA (pre-ZIS-ta)
for HIV (Human Immunodeficiency Virus) Infection
Generic name: darunavir (da-ROO-nuh-veer)

ALERT: Find out about medicines that should NOT be taken with PREZISTA. Please also read the section “Who should not take PREZISTA?”.

Please read this information before you start taking PREZISTA. Also, read the leaflet each time you renew your prescription, just in case anything has changed. Remember, this leaflet does not take the place of careful discussions with your doctor. You and your doctor should discuss your treatment with PREZISTA the first time you take your medicine and at regular checkups. You should remain under a doctor’s care when using PREZISTA and should not change or stop treatment without first talking with a doctor.

WHAT IS PREZISTA?

PREZISTA is an oral tablet used for the treatment of HIV (Human Immunodeficiency Virus) infection in adults. HIV is the virus that causes AIDS (Acquired Immune Deficiency Syndrome). PREZISTA is a type of anti-HIV drug called a protease (PRO-tee-ase) inhibitor.

HOW DOES PREZISTA WORK?

PREZISTA blocks HIV protease, an enzyme which is needed for HIV to multiply. When used with other anti-HIV medicines, PREZISTA may reduce the amount of HIV in your blood (called “viral load”) and increase your CD4 (T) cell count. HIV infection destroys CD4 (T) cells, which are important to the immune system. The immune system helps fight infection. Reducing the amount of HIV and increasing the CD4 (T) cell count may improve your immune system and, thus, reduce the risk of death or infections that can happen when your immune system is weak (opportunistic infections).

PREZISTA is always taken with and at the same time as 100 mg of ritonavir (NORVIR®), in combination with other anti-HIV medicines. PREZISTA should also be taken with food.

DOES PREZISTA CURE HIV OR AIDS?

PREZISTA does **not** cure HIV infection or AIDS. At present, there is no cure for HIV infection. People taking PREZISTA may still develop infections or other conditions associated with HIV infection. Because of this, it is very important for you to remain under the care of a doctor. Although PREZISTA is not a cure for HIV or AIDS, PREZISTA can help reduce your risks of getting illnesses associated with HIV infection (AIDS and opportunistic infection) and eventually dying from these conditions.

* Trademark of Tibotec, Inc.
COPYRIGHT ©2006 Tibotec, Inc.
All rights reserved

DOES PREZISTA REDUCE THE RISK OF PASSING HIV TO OTHERS?

PREZISTA does **not** reduce the risk of passing HIV to others through sexual contact, sharing needles, or being exposed to your blood. For your health and the health of others, it is important to always practice safer sex by using a latex or polyurethane condom or other barrier method to lower the chance of sexual contact with any body fluids such as semen, vaginal secretions, or blood. Never re-use or share needles.

Ask your doctor if you have any questions on how to prevent passing HIV to other people.

WHAT SHOULD I TELL MY DOCTOR BEFORE I TAKE PREZISTA?

Tell your doctor about all of your medical conditions, including if you:

- are allergic to sulfa medicines.
- have diabetes. In general, anti-HIV medicines, such as PREZISTA, might increase sugar levels in the blood.
- have liver problems.
- have hemophilia. Anti-HIV medicines, such as PREZISTA, might increase the risk of bleeding.
- are pregnant or planning to become pregnant. The effects of PREZISTA on pregnant women or their unborn babies are not known. You and your doctor will need to decide if taking PREZISTA is right for you. If you take PREZISTA while you are pregnant, talk to your doctor about how you can be included in the Antiretroviral Pregnancy Registry.
- are breastfeeding. Do not breastfeed if you are taking PREZISTA. You should not breastfeed if you have HIV because of the chance of passing HIV to your baby. Talk with your doctor about the best way to feed your baby.

WHO SHOULD NOT TAKE PREZISTA?*

Together with your doctor, you need to decide whether taking PREZISTA is right for you.

Do not take PREZISTA if you:

- are allergic to darunavir or any of the other ingredients in PREZISTA
- are allergic to ritonavir (NORVIR®)
- take any of the following types of medicines because you could experience serious side effects:

<u>Type of Drug</u>	<u>Examples of Generic Names (Brand Names)</u>
Antihistamines (to treat allergy symptoms)	astemizole (Hismanal®) terfenadine (Seldane®)
Ergot Derivatives (to treat migraine and headaches)	dihydroergotamine (D.H.E. 45®, Migranal®) ergonovine ergotamine (Wigraine®, Ergostat®, Cafergot®, Ergomar®) methylergonovine
Gastrointestinal Motility Agent (to treat some digestive conditions)	cisapride (Propulsid®)
Neuroleptic (to treat psychiatric conditions)	pimozide (Orap®)
Sedative/hypnotics (to treat trouble with sleeping and/or anxiety)	midazolam (Versed®) triazolam (Halcion®)

CAN PREZISTA BE TAKEN WITH OTHER MEDICATIONS?*

Tell your doctor about all the medicines you take including prescription and nonprescription medicines, vitamins, and herbal supplements, including St. John's wort (*Hypericum perforatum*). PREZISTA and many other medicines can interact. Sometimes serious side effects will happen if PREZISTA is taken with certain other medicines (see "Who should not take PREZISTA?").

Tell your doctor if you are taking estrogen-based contraceptives. PREZISTA might reduce the effectiveness of estrogen-based contraceptives. You must take additional precautions for birth control such as a condom.

Tell your doctor if you take other anti-HIV medicines. PREZISTA can be combined with some other anti-HIV medicines while other combinations are not recommended.

* * The brands listed are the registered trademarks of their respective owners and are not trademarks of Tibotec, Inc.

Tell your doctor if you are taking any of the following medicines:

<u>Type of Drug</u>	<u>Examples of Generic Names (Brand Names)</u>
Antiarrhythmics (to treat abnormal heart rhythms)	bepiridil (Vascor [®]) lidocaine (Lidoderm [®]) quinidine amiodarone (Cordarone [®])
Anticoagulants (to prevent the clotting of red blood cells called platelets)	warfarin (Coumadin [®])
Anticonvulsants (to treat epilepsy and prevent seizures)	carbamazepine (Tegretol [®] , Carbatrol [®]) phenobarbital phenytoin (Dilantin [®] , Phenytek [®])
Antidepressants	trazodone (Desyrel [®])
Anti-infectives (to treat bacterial infections)	clarithromycin (Biaxin [®])
Antifungals (to treat fungal infections)	ketoconazole (Nizoral [®]) itraconazole (Sporanox [®]) voriconazole (Vfend [®])
Antimycobacterials (to treat bacterial infections)	rifabutin (Mycobutin [®]) rifampin (Rifadin [®] , Rifater [®] , Rifamate [®])
Calcium Channel Blockers (to treat heart disease)	felodipine (Plendil [®]) nifedipine (Adalat [®]) nicardipine (Cardene [®])
Corticosteroids (to treat inflammation or asthma)	dexamethasone (Decadron [®]) fluticasone propionate (Advair Diskus [®] , Cutivate [®] , Flonase [®] , Flovent Diskus [®])
HMG-CoA Reductase Inhibitors (to lower cholesterol levels)	atorvastatin (Lipitor [®]) lovastatin (Mevacor [®]) pravastatin (Pravachol [®]) simvastatin (Zocor [®])
Immunosuppressants (to prevent organ transplant rejection)	cyclosporine (Sandimmune [®] , Neoral [®]) tacrolimus (Prograf [®]) sirolimus (Rapamune [®])
Narcotic Analgesics	methadone
PDE-5 Inhibitors	sildenafil (Viagra [®])

<u>Type of Drug</u>	<u>Examples of Generic Names (Brand Names)</u>
(to treat erectile dysfunction)	vardenafil (Levitra [®]) tadalafil (Cialis [®])
Selective Serotonin Reuptake Inhibitors (SSRIs) (to treat depression, anxiety, or panic disorder)	paroxetine (Paxil [®]) sertraline (Zoloft [®])

Tell your doctor if you are taking any medicines that you obtained without a prescription.

This is **not** a complete list of medicines that you should tell your doctor that you are taking. Know and keep track of all the medicines you take and have a list of them with you. Show this list to all of your doctors and pharmacists any time you get a new medicine. Both your doctor and your pharmacist can tell you if you can take these other medicines with PREZISTA. Do not start any new medicines while you are taking PREZISTA without first talking with your doctor or pharmacist. You can ask your doctor or pharmacist for a list of medicines that can interact with PREZISTA.

HOW SHOULD I TAKE PREZISTA?

Take PREZISTA tablets every day exactly as prescribed by your doctor. You must take ritonavir (NORVIR[®]) at the same time as PREZISTA. The usual dose is 600 mg (two 300 mg tablets) of PREZISTA, together with 100 mg (one 100 mg capsule) of ritonavir (NORVIR[®]), twice daily *every day*. It may be easier to remember to take PREZISTA and ritonavir (NORVIR[®]) if you take them at the same time every day. If you have questions about when to take PREZISTA and ritonavir (NORVIR[®]), your doctor can help you decide which schedule works for you.

Take PREZISTA and ritonavir (NORVIR[®]) with food. The type of food is not important. Swallow the whole tablets with a drink such as water or milk. Do not chew the tablets.

Continue taking PREZISTA and ritonavir (NORVIR[®]) unless your doctor tells you to stop. Take the exact amount of PREZISTA and ritonavir (NORVIR[®]) that your doctor tells you to take, right from the very start. To help make sure you will benefit from PREZISTA and ritonavir (NORVIR[®]), you must not skip doses or interrupt therapy. If you don't take PREZISTA and ritonavir (NORVIR[®]) as prescribed, the beneficial effects of PREZISTA and ritonavir (NORVIR[®]) may be reduced or even lost.

If you miss a dose of PREZISTA or ritonavir (NORVIR[®]) by more than 6 hours, wait and then take the next dose of PREZISTA and ritonavir (NORVIR[®]) at the regularly scheduled time. If you miss a dose of PREZISTA or ritonavir (NORVIR[®]) by less than 6 hours, take your missed dose of PREZISTA and ritonavir (NORVIR[®]) immediately. Then take your next dose of PREZISTA and ritonavir (NORVIR[®]) at the regularly scheduled time.

You should always take PREZISTA and ritonavir (NORVIR[®]) together with food.