

of the physiological differences between monkeys and humans, leading to more efficient utilization of monkeys as an animal model.

References

- [1] Stevens, J.C., Shipley, L.A., Cashman, J.R., Vandenbranden, M. and Wrighton, S.A. (1993) Comparison of human and rhesus monkey *in vitro* phase I and phase II hepatic drug metabolism activities. *Drug Metab. Dispos.* 21, 753–760.
- [2] Sharer, J.E., Shipley, L.A., Vandenbranden, M.R., Binkley, S.N. and Wrighton, S.A. (1995) Comparisons of phase I and phase II *in vitro* hepatic enzyme activities of human, dog, rhesus monkey, and cynomolgus monkey. *Drug Metab. Dispos.* 23, 1231–1241.
- [3] Guengerich, F.P. (1997) Comparisons of catalytic selectivity of cytochrome P450 subfamily enzymes from different species. *Chem. Biol. Interact.* 106, 161–182.
- [4] Shimada, T., Mimura, M., Inoue, K., Nakamura, S., Oda, H., Ohmori, S. and Yamazaki, H. (1997) Cytochrome P450-dependent drug oxidation activities in liver microsomes of various animal species including rats, guinea pigs, dogs, monkeys, and humans. *Arch. Toxicol.* 71, 401–408.
- [5] Weaver, R.J., Dickins, M. and Burke, M.D. (1999) A comparison of basal and induced hepatic microsomal cytochrome P450 monooxygenase activities in the cynomolgus monkey (*Macaca fascicularis*) and man. *Xenobiotica* 29, 467–482.
- [6] Bogaards, J.J., Bertrand, M., Jackson, P., Oudshoorn, M.J., Weaver, R.J., van Bladeren, P.J. and Walther, B. (2000) Determining the best animal model for human cytochrome P450 activities: a comparison of mouse, rat, rabbit, dog, micropig, monkey and man. *Xenobiotica* 30, 1131–1152.
- [7] Narimatsu, S., Kobayashi, N., Masubuchi, Y., Horie, T., Kakegawa, T., Kobayashi, H., Hardwick, J.P., Gonzalez, F.J., Shimada, N., Ohmori, S., Kitada, M., Asaoka, K., Kataoka, H., Yamamoto, S. and Satoh, T. (2000) Species difference in enantioselectivity for the oxidation of propranolol by cytochrome P450 2D enzymes. *Chem. Biol. Interact.* 127, 73–90.
- [8] Sakate, R., Osada, N., Hida, M., Sugano, S., Hayasaka, I., Shimohira, N., Yanagi, S., Suto, Y., Hashimoto, K. and Hirai, M. (2003) Analysis of 5'-end sequences of chimpanzee cDNAs. *Genome Res.* 13, 1022–1026.
- [9] Magness, C.L., Fellin, P.C., Thomas, M.J., Korth, M.J., Agy, M.B., Proll, S.C., Fitzgibbon, M., Scherer, C.A., Miner, D.G., Katze, M.G. and Iadonato, S.P. (2005) Analysis of the *Macaca mulatta* transcriptome and the sequence divergence between *Macaca* and human. *Genome Biol.* 6, R60.
- [10] Li, Y. and Su, B. (2006) No accelerated evolution of 3'UTR region in human for brain-expressed genes. *Gene* 383C, 38–42.
- [11] Osada, N., Hida, M., Kusuda, J., Tanuma, R., Iseki, K., Hirata, M., Suto, Y., Hirai, M., Terao, K., Suzuki, Y., Sugano, S. and Hashimoto, K. (2001) Assignment of 118 novel cDNAs of cynomolgus monkey brain to human chromosomes. *Gene* 275, 31–37.
- [12] Osada, N., Hida, M., Kusuda, J., Tanuma, R., Hirata, M., Suto, Y., Hirai, M., Terao, K., Sugano, S. and Hashimoto, K. (2002) Cynomolgus monkey testicular cDNAs for discovery of novel human genes in the human genome sequence. *BMC Genomics* 3, 36.
- [13] Chen, W.H., Wang, X.X., Lin, W., He, X.W., Wu, Z.Q., Lin, Y., Hu, S.N. and Wang, X.N. (2006) Analysis of 10,000 ESTs from lymphocytes of the cynomolgus monkey to improve our understanding of its immune system. *BMC Genomics* 7, 82.
- [14] The Rhesus Macaque Genome Sequencing and Analysis Consortium (2007) Evolutionary and biomedical insights from the rhesus macaque genome. *Science* 316, 222–234.
- [15] Suzuki, Y. and Sugano, S. (2003) Construction of a full-length enriched and a 5'-end enriched cDNA library using the oligo-capping method. *Method Mol. Biol.* 221, 73–91.
- [16] Yamashita, R., Suzuki, Y., Wakaguri, H., Tsuritani, K., Nakai, K. and Sugano, S. (2006) DBTSS: database of human transcription start sites, progress report 2006. *Nucl. Acid Res.* 34, D86–89.
- [17] Xu, X.R., Huang, J., Xu, Z.G., Qian, B.Z., Zhu, Z.D., Yan, Q., Cai, T., Zhang, X., Xiao, H.S., Qu, J., Liu, F., Huang, Q.H., Cheng, M., Li, N.G., Du, J.J., Hu, W., Shen, K.T., Lu, G., Fu, G., Zhong, M., Xu, S.H., Gu, W.Y., Huang, W., Zhao, X.T., Hu, G.X., Gu, J.R., Chen, Z. and Han, Z.G. (2001) Insight into hepatocellular carcinogenesis at transcriptome level by comparing gene expression profiles of hepatocellular carcinoma with those of corresponding noncancerous liver. *Proc. Natl. Acad. Sci. USA* 98, 15089–15094.
- [18] Yu, Y., Zhang, C., Zhou, G., Wu, S., Qu, X., Wei, H., Xing, G., Dong, C., Zhai, Y., Wan, J., Ouyang, S., Li, L., Zhang, S., Zhou, K., Zhang, Y., Wu, C. and He, F. (2001) Gene expression profiling in human fetal liver and identification of tissue- and developmental-stage-specific genes through compiled expression profiles and efficient cloning of full-length cDNAs. *Genome Res.* 11, 1392–1403.
- [19] Otsuka, M., Arai, M., Mori, M., Kato, M., Kato, N., Yokosuka, O., Ochiai, T., Takiguchi, M., Omata, M. and Seki, N. (2003) Comparing gene expression profiles in human liver, gastric, and pancreatic tissues using full-length-enriched cDNA libraries. *Hepatology Res.* 27, 76–82.
- [20] Guengerich, F.P. (2005) Human cytochrome P450 enzymes in: (Ortiz de Montellano, P., Ed.), third ed, *Cytochrome P450: Structure, Mechanism, and Biochemistry*, pp. 377–530, Kluwer Academic/Plenum Publishers, New York.
- [21] Goldstein, J.A. (2001) Clinical relevance of genetic polymorphisms in the human CYP2C subfamily. *Brit. J. Clin. Pharmacol.* 52, 349–355.
- [22] Mankowski, D.C., Laddison, K.J., Christopherson, P.A., Ekins, S., Tweedie, D.J. and Lawton, M.P. (1999) Molecular cloning, expression, and characterization of CYP2D17 from cynomolgus monkey liver. *Arch. Biochem. Biophys.* 372, 189–196.
- [23] Hichiya, H., Kuramoto, S., Yamamoto, S., Shinoda, S., Hanioka, N., Narimatsu, S., Asaoka, K., Miyata, A., Iwata, S., Nomoto, M., Satoh, T., Kiryu, K., Ueda, N., Naito, S., Tucker, G.T. and Ellis, S.W. (2004) Cloning and functional expression of a novel marmoset cytochrome P450 2D enzyme, CYP2D30: comparison with the known marmoset CYP2D19. *Biochem. Pharmacol.* 68, 165–175.
- [24] Uno, Y., Fujino, H., Kito, G., Kamataki, T. and Nagata, R. (2006) CYP2C76, a novel CYP in cynomolgus monkey, is a major CYP2C in liver, metabolizing tolbutamide and testosterone. *Mol. Pharmacol.* 70, 477–486.
- [25] Uno, Y., Hosaka, S., Matsuno, K., Nakamura, C., Kito, G., Kamataki, T. and Nagata, R. (2007) Characterization of cynomolgus monkey cytochrome P450 (CYP) cDNAs: Is CYP2C76 the only monkey-specific CYP gene responsible for species differences in drug metabolism? *Arch. Biochem. Biophys.* 466, 98–105.
- [26] Mackenzie, P.I., Bock, K.W., Burchell, B., Guillemette, C., Ikushiro, S., Iyanagi, T., Miners, J.O., Owens, I.S. and Nebert, D.W. (2005) Nomenclature update for the mammalian UDP glycosyltransferase (UGT) gene superfamily. *Pharmacogenet. Genomics* 15, 677–685.
- [27] Hayes, J.D., Flanagan, J.U. and Jowsey, I.R. (2005) Glutathione transferases. *Annu. Rev. Pharmacol. Toxicol.* 45, 51–88.
- [28] Blanchard, R.L., Freimuth, R.R., Buck, J., Weinsilboum, R.M. and Coughtrie, M.W. (2004) A proposed nomenclature system for the cytosolic sulfotransferase (SULT) superfamily. *Pharmacogenetics* 14, 199–211.
- [29] Cashman, J.R. and Zhang, J. (2006) Human flavin-containing monooxygenases. *Annu. Rev. Pharmacol. Toxicol.* 46, 65–100.
- [30] Uno, Y., Kumano, T., Kito, G., Nagata, R., Kamataki, T. and Fujino, H. (2007) CYP2C76-mediated species difference in drug metabolism: A comparison of pitavastatin metabolism between monkeys and humans. *Xenobiotica* 37, 30–43.
- [31] Belanger, G., Barbier, O., Hum, D.W. and Belanger, A. (1999) Molecular cloning, expression and characterization of a monkey steroid UDP-glucuronosyltransferase, UGT2B19, that conjugates testosterone. *Eur. J. Biochem.* 260, 701–708.
- [32] Girard, C., Barbier, O., Turgeon, D. and Belanger, A. (2002) Isolation and characterization of the monkey UGT2B30 gene that encodes a uridine diphosphate-glucuronosyltransferase enzyme active on mineralocorticoid, glucocorticoid, androgen and oestrogen hormones. *Biochem. J.* 365, 213–222.
- [33] Male, D.A., Ormsby, R.J., Ranganathan, S., Giannakis, E. and Gordon, D.L. (2000) Complement factor H: sequence analysis of

- 221 kb of human genomic DNA containing the entire *fH*, *fHR-1* and *fHR-3* genes. *Mol. Immunol.* 37, 41–52.
- [34] Geraghty, D.E., Daza, R., Williams, L.M., Vu, Q. and Ishitani, A. (2002) Genetics of the immune response: identifying immune variation within the MHC and throughout the genome. *Immunol. Rev.* 190, 69–85.
- [35] Venables, J.P., Strain, L., Routledge, D., Bourm, D., Powell, H.M., Warwicker, P., Diaz-Torres, M.L., Sampson, A., Mead, P., Webb, M., Pirson, Y., Jackson, M.S., Hughes, A., Wood, K.M., Goodship, J.A. and Goodship, T.H. (2006) Atypical haemolytic uraemic syndrome associated with a hybrid complement gene. *PLoS Med.* 3, e431.
- [36] Oyama, M., Itagaki, C., Hata, H., Suzuki, Y., Izumi, T., Natsume, T., Isobe, T. and Sugano, S. (2004) Analysis of small human proteins reveals the translation of upstream open reading frames of mRNAs. *Genome Res.* 14, 2048–2052.
- [37] Frith, M.C., Forrest, A.R., Nourbakhsh, E., Pang, K.C., Kai, C., Kawai, J., Carninci, P., Hayashizaki, Y., Bailey, T.L. and Grimmond, S.M. (2006) The abundance of short proteins in the mammalian proteome. *PLoS Genet.* 2, e52.
- [38] Gustincich, S., Sandelin, A., Plessy, C., Katayama, S., Simone, R., Lazarevic, D., Hayashizaki, Y. and Carninci, P. (2006) The complexity of the mammalian transcriptome. *J. Physiol.* 575, 321–332.
- [39] Prasanth, K.V. and Spector, D.L. (2007) Eukaryotic regulatory RNAs: an answer to the 'genome complexity' conundrum. *Genes Dev.* 21, 11–42.
- [40] Demuth, J.P., Bie, T.D., Stajich, J.E., Cristianini, N. and Hahn, M.W. (2006) The evolution of mammalian gene families. *PLoS ONE* 1, e85.
- [41] Mattick, J.S. and Makunin, I.V. (2006) Non-coding RNA. *Hum. Mol. Genet.* 15, R17–R29.