

Figure 2. Generation of TCR γ -deficient mice and subsequent production of TCR $\alpha\gamma$ double mutant mice. (A) Schematic representation of WT and mutant (pC γ 4 Δ NL) genomic C γ 4 loci together with the 3 DNA fragments used to construct the mutant pC γ 4 Δ NL vector. The resulting targeting vector (pC γ 4 Δ NL) carrying a *loxP*-flanked *pgk-neo* gene cassette in place of exon 1 of C γ 4 gene used a neomycin resistance gene driven by the *pgk* promoter as positive selection marker is shown. Restriction enzyme sites, *Sph*I and *Kpn*I (solid bars), exon structures, V γ and J γ (open boxes), and *loxP* site (solid triangle) are indicated. (B) Schematic representation of the ES clone carrying WT TCR γ gene and mutant V γ 6 Δ L ES clone carrying the allele in which the V γ 6 region was replaced by a single *loxP* site and mutant pC γ 4 Δ NL targeting vector. Exon structures, V γ and J γ (open boxes) and *loxP* site (solid triangle) are indicated. (C) Schematic representation of generation of the mutant mice that carry the TCR γ -deficient (C γ Δ) allele by Cre-mediated recombination during embryonic development. Exon structures, V γ and J γ (open boxes), and *loxP* site (solid triangle) are indicated. (D) The mutant mice that carry the TCR γ -deficient (C γ Δ) allele were intercrossed to produce TCR $\gamma^{+/+}$ (WT), TCR $\gamma^{+/-}$ ($\gamma^{+/-}$), and TCR $\gamma^{-/-}$ ($\gamma^{-/-}$) mice, and the corresponding WT and mutant alleles were typed by PCR analysis of tail DNA with each set of primers indicated. (E) $\gamma\delta$ T cells are absent from the IEL compartment of $\gamma^{-/-}$ and $\alpha\gamma^{-/-}$ mice. $\gamma^{-/-}$ Mice were crossed with $\alpha^{-/-}$ mice to obtain WT ($\alpha^{+/+} \times \gamma^{+/+}$), $\gamma^{-/-}$ ($\alpha^{+/+} \times \gamma^{-/-}$), $\alpha^{-/-}$ ($\alpha^{-/-} \times \gamma^{+/+}$), and $\alpha\gamma^{-/-}$ ($\alpha^{-/-} \times \gamma^{-/-}$) littermate mice.


mation characterized by elongation of crypts was much milder in $\alpha\gamma^{-/-}$ mice as compared with $\alpha^{-/-}$ mice (Figure 3A). Although the body weight was comparable between $\alpha\gamma^{-/-}$ and $\alpha^{-/-}$ mice, it was evident that colonic weight was significantly decreased in $\alpha\gamma^{-/-}$ mice as compared with $\alpha^{-/-}$ mice (Figure 3B). The disease score characterized by the thickening of colonic mucosa with crypt elongation and inflammatory cell infiltration was also significantly lower in $\alpha\gamma^{-/-}$ mice than that rated in $\alpha^{-/-}$ mice (Figure 3C). Although approximately 20% of 20- to 60-week-old $\alpha^{-/-}$ mice displayed anorectal prolapse (ARP), it was not discerned in any of age-matched $\alpha\gamma^{-/-}$ mice (Figure 3D). Notably, no difference was observed in the age of onset of colitis and in the incidence of colitis (~80%) among 20- to 32-week-old $\alpha\gamma^{-/-}$ and $\alpha^{-/-}$ mice. In addition, in comparison with administration of PBS (as control), adoptive transfer of $\gamma\delta$ T cells that were purified from $\alpha^{-/-}$ mice did not increase the incidence of colitis in the recipient $\alpha\gamma^{-/-}$ mice. However,

the transfer of $\gamma\delta$ T cells exacerbated the severity of colitis in the recipient $\alpha\gamma^{-/-}$ mice. As shown in Figure 3E, more severe inflammatory cell infiltration was observed in the inflamed colon of the recipient $\alpha\gamma^{-/-}$ mice with $\gamma\delta$ T-cell transfer as compared with control $\alpha\gamma^{-/-}$ mice. Therefore, it is possible that $\gamma\delta$ T cells may be involved in the exacerbation, but not induction, of UC-like colitis.

Decrease in the Colonic Neutrophils and Monocytes in the Absence of $\gamma\delta$ T Cells

The above results indicate that the spontaneous colitis in $\alpha^{-/-}$ mice is ameliorated by the absence of $\gamma\delta$ T cells in $\alpha\gamma^{-/-}$ mice. With these findings in mind, flow cytometric analysis of colonic LP cells isolated from WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ littermate mice at approximately 28 weeks of age was performed, and the representative results of 5 independent experiments are presented in Figure 4A. In this experiment, WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ littermate mice yielded 5.1×10^5 , 6.1

Figure 3. Colonic mucosal inflammation is milder in $\alpha\gamma^{-/-}$ mice lacking $\gamma\delta$ T cells than in $\alpha^{-/-}$ mice possessing $\gamma\delta$ T cells. (A) Representative histologic pictures of colon from WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ littermate mice. These mice were maintained in the same cages under specific pathogen-free conditions and examined at 28 weeks of age. (B) Body weight, wet weight of large intestine (LI weight), and ratio (%) of LI weight to body weight in WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ littermate mice at 20 to 32 weeks of age are shown. Values in parentheses are numbers of mice examined. *** $P < .001$. (C) Disease scores (0–10) of colitis in 24- to 32-week-old WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ mice were assessed. Values in parentheses are numbers of mice examined. * $P < .05$. (D) Incidence of ARP in 24- to 60-week-old WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ mice. (E) The histologic findings (original magnification, $\times 20$) of colon of recipient $\alpha\gamma^{-/-}$ mouse groups that received transfer of $\gamma\delta$ T cells (left panel) and that received PBS (right panel).



$\times 10^5$, 32.3×10^5 , and 12.0×10^5 colonic LP cells, respectively. Based on the absolute numbers of infiltrated cells and the flow cytometry results shown in Figure 4A, it was evident that fewer $\text{Mac-1}^+\text{Ly-6G}^-$ cells and $\text{Mac-1}^+\text{Ly-6G}^+$ cells were present in the colonic LP cell population of $\alpha\gamma^{-/-}$ mice as compared with those of $\alpha^{-/-}$ mice. Monocytes express Mac-1 but not Ly-6G, whereas neutrophils express both Mac-1 and Ly-6G.²¹ Therefore, our results suggest that, in addition to monocyte infiltration (Figure 4A), there is a marked infiltration of neutrophils in the inflamed colonic LP of $\alpha^{-/-}$ mice. We also confirmed our previous finding¹⁵ that a remarkable increase in $\gamma\delta$ T cells was observed in the inflamed colonic LP of $\alpha^{-/-}$ mice (Figure 4A).

Immunohistochemical examination of inflamed colons from $\alpha^{-/-}$ and $\alpha\gamma^{-/-}$ mice at approximately 28 weeks of age was performed to further confirm flow cytometric results. Consistent with the flow cytometric observations, significantly smaller numbers of Mac-1^+ cells and Ly-6G^+ cells were observed in the colonic LP of $\alpha\gamma^{-/-}$ mice as compared with those in the colonic LP of $\alpha^{-/-}$ mice (Figure 4B).

To investigate whether $\gamma\delta$ T cells contribute to the generation of colonic environment for enhancing the

migration of neutrophils and monocytes into the inflamed colon, we examined chemotactic activity of colonic extracts from $\alpha^{-/-}$ and $\alpha\gamma^{-/-}$ mice to neutrophils and monocytes (Figure 5A). As a result, chemotactic activity to neutrophils of colonic extracts from $\alpha\gamma^{-/-}$ mice was significantly weaker than that from $\alpha^{-/-}$ mice, whereas the chemotactic activities to monocytes of both extracts were almost comparable (Figure 5B). The marked infiltration of neutrophils into the inflamed colonic LP of $\alpha^{-/-}$ mice is most likely mediated by some factors, such as MIP-2 (Figure 5A), that are enhanced in the presence of $\gamma\delta$ T cells.

Taking all of these results together, and in conjunction with our previous findings,²² colonic $\gamma\delta$ T cells of $\alpha^{-/-}$ mice exert aggravating effect on the UC-like colitis by increasing primarily the influx of neutrophils into the inflamed mucosa.

Attenuation of Colonic Proinflammatory Cascades by the Absence of $\gamma\delta$ T Cells

In view of the severe colitis, increased infiltration of $\text{Mac-1}^+\text{Ly-6G}^+$ and $\text{Mac-1}^+\text{Ly-6G}^-$ cells, and marked production of neutrophil chemotactic factor(s) in the inflamed colonic LP of $\alpha^{-/-}$ mice, quantitative real-time

BASIC-ALIMENTARY TRACT




Figure 4. Decrease in colonic Mac-1⁺Ly-6G⁻ and Mac-1⁻Ly-6G⁺ cells in the absence of $\gamma\delta$ T cells. WT (n = 3), $\gamma^{-/-}$ (n = 3), $\alpha^{-/-}$ (n = 4), and $\alpha\gamma^{-/-}$ (n = 4) mice from 28 weeks of age were examined. (A) Flow cytometric profiles of colonic LP cells. Absolute numbers of LP cells isolated from these WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ individuals were 5.1×10^5 , 6.1×10^5 , 32.3×10^5 and 12.0×10^5 , respectively. (B) Representative immunohistochemical verification of the prominent infiltrations of Mac-1⁺ and Ly-6G⁺ cells into the inflamed colonic LP of $\alpha^{-/-}$ mice. Five tissue sections prepared from ascending colon to rectum per mouse, namely, 20 sections obtained from inflamed large intestines of $\alpha^{-/-}$ mice and those obtained from inflamed large intestines of $\alpha\gamma^{-/-}$ mice, were examined in a blinded fashion by 5 independent investigators, and the statistical difference in absolute numbers of Mac-1⁺ and Ly-6G⁺ cells between large intestinal mucosa from $\alpha^{-/-}$ and $\alpha\gamma^{-/-}$ mice were determined by 2-sided Mann-Whitney U test. *P < .05.

RT-PCR analysis and measurement of the amounts of representative proinflammatory cytokines as well as chemokines were performed to dissect further the role of $\gamma\delta$ T cells in the UC-like colitis in $\alpha^{-/-}$ mice.

To this end, messenger RNA (mRNA) and proteins prepared from the large intestines of WT, $\gamma^{-/-}$, $\alpha^{-/-}$, and

$\alpha\gamma^{-/-}$ mice were examined. Inflamed colonic tissues from $\alpha^{-/-}$ and $\alpha\gamma^{-/-}$ mice contained at least 10 times higher levels of cytokine (Table 1)- and chemokine (Table 2)-specific mRNA than those of WT and $\gamma^{-/-}$ mice except for IL-7 and IL-10 mRNA. In contrast to the mRNA from colonic tissues of $\alpha^{-/-}$ mice, those of $\alpha\gamma^{-/-}$ mice contained significantly smaller amounts of cytokine (TNF- α , IL-1 β , IL-6, and TGF- β)- and chemokine (KC and MIP-2)-specific mRNA. With these observations in mind, we measured the amounts of representative cytokines as well as chemokines that had exhibited the differences in mRNA levels between the colonic tissues of $\alpha^{-/-}$ and $\alpha\gamma^{-/-}$ mice. First, in situ production of TNF- α , IL-1 β , and IL-6 but not TGF- β proteins was significantly down-regulated in the inflamed colonic mucosa of $\alpha\gamma^{-/-}$ mice as compared with that of $\alpha^{-/-}$ mice (Table 1). Second, KC and MIP-2 chemokines that are involved in the chemoattract of neutrophils and/or monocytes²³ were significantly decreased in large intestines of $\alpha\gamma^{-/-}$ mice compared with those in large intestines of $\alpha^{-/-}$ mice (Table 2).

To investigate the cell types responsible for the increases in these proinflammatory cytokines and chemokines, real-time RT-PCR analysis of mRNA present in the purified cell subsets from the inflamed colonic LP of $\alpha^{-/-}$ and $\alpha\gamma^{-/-}$ mice was performed (see Supplementary Figure 1 online at www.gastrojournal.org). The IL-1 β and MIP-2 mRNA were expressed preferentially by Gr-1⁺ cells, F4/80⁺ cells, and CD11c⁺ cells in the colon, whereas IL-6 mRNA was mainly expressed by Gr-1⁻F4/80⁻CD11c⁻ cell populations. Expression levels of TNF- α and KC-specific mRNA were comparable between all cell populations (Gr-1⁺ cells, F4/80⁺ cells, CD11c⁺ cells, and Gr-1⁻F4/80⁻CD11c⁻ cells) examined. Finally, the expression levels of these cytokine- and chemokine-specific mRNA in every cell subset were lower in cells from $\alpha\gamma^{-/-}$ mice than those in cells from $\alpha^{-/-}$ mice (see Supplementary Figure 1 online at www.gastrojournal.org).

Discussion

The $\alpha^{-/-}$ mice spontaneously develop colitis that shares many features with human UC.^{16,24} Commensal enteric flora is required for the development of this colitis as indicated by the absence of colitis in $\alpha^{-/-}$ mice that are maintained under germ-free conditions.^{22,25} The number of colonic $\gamma\delta$ T cells drastically decreases in the $\alpha^{-/-}$ mice under germ-free conditions.²² However, the study to identify the role of $\gamma\delta$ T cells in the UC-like chronic colitis in $\alpha^{-/-}$ mice has been hampered by the difficulty in generating TCR $\alpha\delta$ double mutant ($\alpha\delta^{-/-}$) mice because of the genomic organization of these TCR genes.²⁰ In the present study, we overcame this problem by newly generating $\gamma^{-/-}$ mice and subsequently crossing these mice with $\alpha^{-/-}$ mice to generate TCR α double mutant mice that lacked $\gamma\delta$ T cells. By using these $\alpha\gamma^{-/-}$ mice, we herein provide a novel insight into the role of $\gamma\delta$ T cells




Figure 5. Chemotactic activity of colonic extracts from WT, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ mice to neutrophils and monocytes. (A) The number of neutrophils and monocytes migrated in response to the increasing concentration of MIP-2 (open square) and MCP-1 (solid circle). The representative flow cytometric profiles are shown in the lower panels, and red and blue gates indicate the neutrophils and monocytes, respectively. Colored numbers represent means of the number of cells in each gate. (B) Chemotactic responses of neutrophils and monocytes to colonic extracts (0.7 mg/mL protein) from WT, $\alpha^{-/-}$, and $\alpha\gamma^{-/-}$ mice, and each circle represents an individual mouse. Horizontal bars show mean values, and dotted lines indicate the number of migrated cells in medium alone. The representative flow cytometric profiles of 3 individual animals indicated by the solid red and blue circles (upper panels) are shown in the lower panels, and red and blue gates indicate the neutrophils and monocytes, respectively.

that contributes to the exacerbation of UC-like colitis in $\alpha^{-/-}$ mice.

There is growing evidence supporting the fact that $\gamma\delta$ T cells are part of the innate immune system and play an active multifaceted immunoregulatory role in the coordinated innate and acquired immune responses that maintain the integrity of many organs containing epithelia.^{1,5,26,27} Nevertheless, the details of $\gamma\delta$ T-cell functions

are still not well understood as compared with those of $\alpha\beta$ T cells. $\gamma\delta$ T cells might play a defensive role against infections by various pathogenic microorganisms because exaggerated and severe infectious diseases occur in $\delta^{-/-}$ mice.²⁸⁻³³ However, the same $\delta^{-/-}$ mice have also been demonstrated to display an increased host resistance to infection.^{34,35} With regard to this, it is noteworthy that $V\gamma 1^+$ $\gamma\delta$ T cells are reported to eliminate the

Table 1. Real-Time RT-PCR Analysis and ELISA Assay of Cytokines in the Colonic Tissues

Mice (n)	Cytokine							
	TNF- α	IL-1 β	IL-6	TGF- β	IFN- γ	IL-7	IL-10	IL-12
RT-PCR (copies per 10 ³ HPRT)								
WT (5)	11.4 \pm 0.53	6.86 \pm 0.55	2.45 \pm 0.64	123 \pm 14.6	ND	11.7 \pm 1.03	2.63 \pm 0.28	0.82 \pm 0.21
$\gamma^{-/-}$ (5)	9.20 \pm 0.76	6.01 \pm 0.61	2.14 \pm 0.88	106 \pm 10.7	ND	10.2 \pm 0.78	2.09 \pm 0.31	0.33 \pm 0.09
$\alpha^{-/-}$ (7)	457 \pm 35.7**	260 \pm 18.6***	7.06 \pm 1.04*	538 \pm 48.6*	64.4 \pm 8.90	11.0 \pm 0.89	4.99 \pm 0.72	9.58 \pm 1.67
$\alpha\gamma^{-/-}$ (6)	205 \pm 51.8**	98.5 \pm 21.4***	2.93 \pm 0.73*	319 \pm 57.4*	45.0 \pm 15.6	9.29 \pm 1.11	6.42 \pm 2.11	6.63 \pm 1.01
ELISA (pg/mg protein)								
$\alpha^{-/-}$ (7)	100 \pm 12.2**	975 \pm 70.1**	8.24 \pm 1.39**	2.09 \pm 0.11	8.34 \pm 1.50			
$\alpha\gamma^{-/-}$ (6)	27.0 \pm 8.28**	647 \pm 24.6**	2.92 \pm 0.52**	2.57 \pm 0.31	5.92 \pm 2.56			

NOTE. All results are expressed as mean \pm SE.

* $P < .05$.

** $P < .01$.

*** $P < .001$.

ND, not detected

Table 2. Real-Time RT-PCR Analysis and ELISA Assay of Chemokines in the Colonic Tissues

Mice (n)	Chemokine					
	KC	MIP-2	GCP-2	MCP-1	MIP-1 α	MIP-1 β
RT-PCR (copies per 10 ³ HPRT)						
WT (5)	6.05 \pm 1.38	0.23 \pm 0.01	12.3 \pm 4.11	1.67 \pm 0.17	0.78 \pm 0.07	1.36 \pm 0.22
$\gamma^{-/-}$ (5)	11.8 \pm 1.19	0.22 \pm 0.03	4.03 \pm 1.49	1.77 \pm 0.26	0.80 \pm 0.04	1.42 \pm 0.11
$\alpha^{-/-}$ (7)	382 \pm 88.6*	101 \pm 12.8**	721 \pm 136	23.3 \pm 4.37	35.0 \pm 3.01	19.8 \pm 0.88
$\alpha\gamma^{-/-}$ (6)	65.2 \pm 17.4*	28.0 \pm 9.80**	307 \pm 168	16.0 \pm 4.11	23.9 \pm 7.34	14.2 \pm 3.97
ELISA (pg/mg protein)						
$\alpha^{-/-}$ (7)	144 \pm 22.4**	113 \pm 21.2*	737 \pm 139			
$\alpha\gamma^{-/-}$ (6)	32.4 \pm 13.4**	44.9 \pm 14.0*	414 \pm 219			

NOTE. All results are expressed as mean \pm SE.* $P < .05$.** $P < .01$.

macrophages infected with *Listeria monocytogenes*, whereas $\gamma\delta$ T cells using V γ elements other than V γ 1 gene appear to lack the ability to control macrophages but possess the ability to protect hosts from the infection-induced tissue injury.^{36,37} In contrast to the beneficial function of $\gamma\delta$ T cells by virtue of the fact that they can maintain the homeostasis of different types of organs,^{1-5,8,27} a deleterious effect of $\gamma\delta$ T cells on the regulation of neutrophil-mediated tissue damage after thermal (postburn) injury has been reported.³⁸ In various chronic and/or autoimmune inflammatory diseases, such as collagen-induced arthritis in mice³⁹ and murine insulin-dependent diabetes,⁴⁰ $\gamma\delta$ T cells have been shown to exert a protective effect. Conversely, $\gamma\delta$ T cells may directly contribute to autoimmune pathology of murine experimental allergic encephalomyelitis⁴¹ as well as lupus in MRL/lpr mice.⁴² Overall, both the beneficial and detrimental roles of $\gamma\delta$ T cells in inflammatory process are evident.⁴³

In chemically induced acute intestinal inflammation models (2,4,6-trinitrobenzene sulfonic acid- or dextran sulfate sodium-induced colitis), $\gamma\delta$ T cells have been reported to play a protective role.¹¹⁻¹⁴ Depletion of $\gamma\delta$ T cells by administration of anti-TCR $\gamma\delta$ mAb into TNF^{AARE} mice with a high frequency of spontaneous ileitis⁴⁴ did not lead to any histologic changes of ileitis.⁴⁵ However, transfer of bone marrow-derived $\gamma\delta$ T cells has been shown to induce CD-like colitis in the bone marrow transplanted CD3 ϵ tg colitis model.⁴⁶ Although the role of $\gamma\delta$ T cells in spontaneous chronic colitis remains to be explored to date, the results of the present study demonstrate the exacerbating effect of $\gamma\delta$ T cells on the UC-like chronic colitis in $\alpha^{-/-}$ mice (Figures 3 and 4 and Tables 1 and 2). Interestingly, approximately 20% of $\alpha^{-/-}$ mice during 20 to 60 weeks of age suffered from ARP, whereas none of age-matched $\alpha\gamma^{-/-}$ mice showed ARP (Figure 3D). Of note, there were no differences in the age of onset of colitis and in the incidence of colitis (~80%) among 20- to 32-week-old $\alpha^{-/-}$ and $\alpha\gamma^{-/-}$ mice, but much more severe colitis was observed in $\alpha^{-/-}$ mice as compared with $\alpha\gamma^{-/-}$ mice. Therefore, it is possible that ARP may reflect

increased severity of colitis and that $\gamma\delta$ T cells may participate in the development of ARP.

Absence of $\gamma\delta$ T cells in $\alpha\gamma^{-/-}$ mice leads to a significantly reduced production of TNF- α , IL-1 β , and IL-6 proteins in the colonic tissues. These findings are consistent with our previous results²⁴ showing the involvement of TNF- α , IL-1 β , and IL-6 in the perpetuation of inflammatory process in $\alpha^{-/-}$ mice. These inflammatory mediators have been shown to be important for host defense and wound repair.⁴⁷ Both KC and MIP-2 attract neutrophils to inflamed sites, and, in certain microbial infection, the collection of neutrophils leads to suppuration reflecting an active and vigorous host response against microbes. We also confirmed that colonic extracts from $\alpha\gamma^{-/-}$ mice exhibited the significantly weaker chemotactic activity to neutrophils than those from $\alpha^{-/-}$ mice. KC and MIP-2 mRNA expressions were lower in all purified cell subsets (Gr-1⁺, F4/80⁺, CD11c⁺, and Gr-1⁻F4/80⁻CD11c⁻ cells) from $\alpha\gamma^{-/-}$ mice than those from $\alpha^{-/-}$ mice. Therefore, in the presence of $\gamma\delta$ T cells, many types of immune cells may be triggered to produce more chemokines, followed by infiltration of neutrophils into the colonic mucosa in $\alpha^{-/-}$ mice. $\gamma\delta$ T-cell responsiveness that is manifested by recruitment and activation of inflammatory cells in which neutrophils predominate has also been reported.^{1,31} In this context, it is of importance to note that the activity and severity of UC patients with increase in $\gamma\delta$ T cells in the inflamed mucosa^{9,10} (Figure 1) can be judged by the activation state of neutrophils in circulation⁴⁸ as well as by regional accumulation of neutrophils in the colonic crypt walls (cryptitis) or in the lumen of crypts (crypt abscess).⁴⁹

The suppressive role of B cells⁵⁰ and the aggravating role of TCR β^{dim} T cells^{22,51} in the pathogenesis of colitis in $\alpha^{-/-}$ mice have been reported. Therefore, it is possible that $\gamma\delta$ T cells may contribute to the exacerbation of this colitis by dampening regulatory B-cell function or by cooperating the colitogenic TCR β^{dim} T cells. The possible complicated mechanism remains to be explored in the future. Levels of TNF- α and IL-1 β mRNA in F4/80⁺ cells

are higher in $\alpha^{-/-}$ mice compared with $\alpha\gamma^{-/-}$ mice (see Supplementary Figure 1 online at www.gastrojournal.org), suggesting that $\gamma\delta$ T cells may activate macrophages to secrete large amounts of proinflammatory cytokines.

In conclusion, although $\gamma\delta$ T cells at the inflamed colonic LP of $\alpha^{-/-}$ mice may protect intestinal epithelial injury, proinflammatory cytokines and neutrophil- and/or monocyte-chemoattractant chemokines induced by $\gamma\delta$ T cells may directly and/or indirectly contribute to increased severity of UC-like chronic colitis in $\alpha^{-/-}$ mice. Further understanding of the molecular mechanisms of $\gamma\delta$ T cell-mediated exacerbation of colitis in $\alpha^{-/-}$ mice will lead us to work out better therapeutic strategies for human UC.

Supplementary Data

Note: To access the supplementary material accompanying this article, visit the online version of *Gastroenterology* at www.gastrojournal.org, and at doi: 10.1053/j.gastro.2007.11.056.

References

- Hayday A, Tigelaar R. Immunoregulation in the tissues by $\gamma\delta$ T cells. *Nat Rev Immunol* 2003;13:233–242.
- Jameson J, Ugarte K, Chen N, et al. A role for skin $\gamma\delta$ T cells in wound repair. *Science* 2002;296:747–749.
- Boismenu R, Havran WL. Modulation of epithelial cell growth by intraepithelial $\gamma\delta$ T cells. *Science* 1994;266:1253–1255.
- Komano H, Fujiura Y, Kawaguchi M, et al. Homeostatic regulation of intestinal epithelia by intraepithelial $\gamma\delta$ T cells. *Proc Natl Acad Sci USA* 1995;92:6147–6151.
- Mak TW, Ferrick DA. The $\gamma\delta$ T-cell bridge: linking innate and acquired immunity. *Nat Med* 1998;4:764–765.
- Podolsky DK. Inflammatory bowel disease. *N Engl J Med* 2002;347:417–429.
- Targan SR, Karp LC. Defects in mucosal immunity leading to ulcerative colitis. *Immunol Rev* 2005;206:296–305.
- Shiohara T, Moriya N, Hayakawa J, et al. Resistance to cutaneous graft vs host disease is not induced in T-cell receptor δ gene-mutant mice. *J Exp Med* 1996;183:1483–1489.
- McVay LD, Li B, Biancaniello R, et al. Changes in human mucosal $\gamma\delta$ T cell repertoire and function associated with the disease process in inflammatory bowel disease. *Mol Med* 1997;3:183–203.
- Yeung MM-W, Melgar S, Baranov V, et al. Characterization of mucosal lymphoid aggregates in ulcerative colitis: immune cell phenotype and TCR- $\gamma\delta$ expression. *Gut* 2000;47:215–227.
- Hoffmann JC, Peters K, Henschke S, et al. Role of T lymphocytes in rat 2,4,6-trinitrobenzene sulphonic acid (TNBS) induced colitis: increased mortality after $\gamma\delta$ T cell depletion and no effect of $\alpha\beta$ T cell depletion. *Gut* 2001;48:489–495.
- Inagaki-Ohara K, Chinen T, Matsuzaki G, et al. Mucosal T cells bearing TCR- $\gamma\delta$ play a protective role in intestinal inflammation. *J Immunol* 2004;173:1390–1398.
- Chen Y, Chou K, Fuchs E, et al. Protection of the intestinal mucosa by intraepithelial $\gamma\delta$ T cells. *Proc Natl Acad Sci U S A* 2002;99:14338–14343.
- Tsuchiya T, Fukuda S, Hamada H, et al. Role of $\gamma\delta$ T cells in the inflammatory response of experimental colitis mice. *J Immunol* 2003;171:5507–5513.
- Mizoguchi A, Mizoguchi E, Chiba C, et al. Cytokine imbalance and autoantibody production in T cell receptor- α mutant mice with inflammatory bowel disease. *J Exp Med* 1996;183:847–856.
- Mombaerts P, Mizoguchi E, Grusby MJ, et al. Spontaneous development of inflammatory bowel disease in T cell receptor mutant mice. *Cell* 1993;75:274–282.
- Hokama A, Mizoguchi E, Sugimoto K, et al. Induced reactivity of intestinal CD4⁺ T cells with an epithelial cell lectin, galectin-4, contributes to exacerbation of intestinal inflammation. *Immunity* 2004;20:681–693.
- de Bruijn MF, van Vianen W, Ploemacher RE, et al. Bone marrow cellular composition in *Listeria monocytogenes* infected mice detected using ER-MP12 and ER-MP20 antibodies: a flow cytometric alternative to different counting. *J Immunol Methods* 1998;217:27–39.
- Itohara S, Mombaerts P, Lafaille J, et al. T cell receptor δ gene mutant mice: independent generation of $\alpha\beta$ T cells and programmed rearrangements of $\gamma\delta$ TCR genes. *Cell* 1993;72:337–348.
- Davis MM, Bjorkman PJ. T-cell antigen receptor genes and T-cell recognition. *Nature* 1988;334:395–402.
- Goren I, Kampfer H, Muller E, et al. Oncostatin M expression is functionally connected to neutrophils in the early inflammation phase of skin repair: implications for normal and diabetes-impaired wounds. *J Invest Dermatol* 2006;126:628–637.
- Kawaguchi-Miyashita M, Shimada S, Kurosu H, et al. An accessory role of TCR- $\gamma\delta$ ⁺ cells in the exacerbation of inflammatory bowel disease in TCR α mutant mice. *Eur J Immunol* 2001;31:980–988.
- Charo IF, Ransohoff RM. The many roles of chemokines and chemokine receptors in inflammation. *N Engl J Med* 2006;354:610–621.
- Mizoguchi A, Mizoguchi E, Bhan AK. Immune networks in animal models of inflammatory bowel disease. *Inflamm Bowel Dis* 2003;9:246–259.
- Dianda L, Hanby AM, Wright NA, et al. T cell receptor- $\alpha\beta$ -deficient mice fail to develop colitis in the absence of a microbial environment. *Am J Pathol* 1997;150:91–97.
- Tonegawa S, Berns A, Bonneville M, et al. Diversity, development, and probable functions of $\gamma\delta$ T cells. *Cold Spring Harbor Symp Quant Biol* 1989;54:31–44.
- Havran WL. A role for epithelial $\gamma\delta$ T cells in tissue repair. *Immunol Res* 2000;21:63–69.
- Mombaerts P, Arnoldi J, Russ F, et al. Different roles of $\alpha\beta$ and $\gamma\delta$ T cells in immunity against an intracellular bacterial pathogen. *Nature* 1993;365:53–56.
- Roberts SJ, Smith AL, Wet AB, et al. T-cell $\alpha\beta$ ⁺ and $\gamma\delta$ ⁺ deficient mice display abnormal but distinct phenotypes toward a natural, widespread infection of the intestinal epithelium. *Proc Natl Acad Sci U S A* 1996;93:11774–11779.
- D'Souza CD, Cooper AM, Frank AA, et al. An anti-inflammatory role for $\gamma\delta$ T lymphocytes in acquired immunity to *Mycobacterium tuberculosis*. *J Immunol* 1997;158:1217–1221.
- King DP, Hyde DM, Jackson KA, et al. Cutting edge: protective response to pulmonary injury requires $\gamma\delta$ lymphocytes. *J Immunol* 1999;162:5033–5036.
- Moore TA, Moore BB, Newstead NW, et al. $\gamma\delta$ -T cells are critical for survival and early proinflammatory cytokine gene expression during murine *Klebsiella pneumoniae*. *J Immunol* 2000;165:2643–2650.
- Selin LK, Santolucito PA, Pinto AK, et al. Innate immunity to viruses: control of vaccinia virus infection by $\gamma\delta$ T cells. *J Immunol* 2001;166:6784–6794.
- Emoto M, Nishimura H, Sakai T, et al. Mice deficient in $\gamma\delta$ T cells are resistant to lethal infection with *Salmonella choleraesuis*. *Infect Immun* 1995;63:3736–3738.

35. Uezu K, Kawakami K, Miyagi K, et al. Accumulation of $\gamma\delta$ T cells in the lungs and their regulatory roles in Th1 response and host defense against pulmonary infection with *Cryptococcus neoformans*. *J Immunol* 2004;172:7629–7634.
36. Andrew EM, Newton DJ, Dalton JE, et al. Delineation of the function of a major $\gamma\delta$ T cell subset during infection. *J Immunol* 2005;175:1741–1750.
37. Newton DJ, Andrew EM, Dalton JE, et al. Identification of novel $\gamma\delta$ T-cell subsets following bacterial infection in the absence of $V\gamma 1^+$ T cells: homeostatic control of $\gamma\delta$ T-cell responses to pathogen infection by $V\gamma 1^+$ T cells. *Infect Immun* 2006;74:1097–1105.
38. Toth B, Alexander M, Daniel T, et al. The role of $\gamma\delta$ T cells in the regulation of neutrophil-mediated tissue damage after thermal injury. *J Leukoc Biol* 2004;76:545–552.
39. Peterman GM, Spencer C, Sperling AI, et al. Role of $\gamma\delta$ T cells in murine collagen-induced arthritis. *J Immunol* 1993;151:6546–6558.
40. Harrison LC, Dempsey-Collier M, Kramer DR, et al. Aerosol insulin induces regulatory CD8 $\gamma\delta$ T cells that prevent murine insulin-dependent diabetes. *J Exp Med* 1996;184:2167–2174.
41. Rajan AJ, Gao Y-L, Raine CS, et al. A pathogenic role of $\gamma\delta$ T cells in relapsing-remitting experimental allergic encephalomyelitis in the SJL mouse. *J Immunol* 1996;157:941–949.
42. Peng SL, Madaio MP, Highes DP, et al. Murine lupus in the absence of $\alpha\beta$ T cells. *J Immunol* 1996;156:4041–4049.
43. Hayday A, Geng L. $\gamma\delta$ cells regulate autoimmunity. *Curr Opin Immunol* 1997;9:884–889.
44. Kontoyiannis D, Pasparakis M, Piazorzo TT, et al. Impaired on/off regulation of TNF biosynthesis in mice lacking TNF AU-rich elements: implications for joint and gut-associated immunopathologies. *Immunity* 1999;10:387–398.
45. Kuhl AA, Loddenkemper C, Westermann J, et al. Role of $\gamma\delta$ T cells in inflammatory bowel disease. *Pathobiology* 2002;70:150–155.
46. Simpson SJ, Hollander GA, Mizoguchi E, et al. Expression of pro-inflammatory cytokines by $TCR\alpha\beta^+$ and $TCR\gamma\delta^+$ T cells in an experimental model of colitis. *Eur J Immunol* 1997;27:17–25.
47. Nathan C. Points of control in inflammation. *Nature* 2002;420:846–852.
48. Suematsu M, Suzuki M, Kitahora T, et al. Increased respiratory burst of leukocytes in inflammatory bowel diseases: the analysis of free radical generation by using chemiluminescence probe. *J Clin Lab Immunol* 1987;24:125–128.
49. Simmonds NJ, Allen RE, Stevens TR, et al. Chemiluminescence assay of mucosal reactive oxygen metabolites in inflammatory bowel disease. *Gastroenterology* 1992;103:186–196.
50. Sugimoto K, Ogawa A, Shimomura Y, et al. Inducible IL-12-producing B cells regulate Th2-mediated intestinal inflammation. *Gastroenterology* 2007;133:124–136.
51. Takahashi I, Kiyono H, Hamada S. $CD4^+$ T-cell population mediates development of inflammatory bowel disease in T-cell receptor α chain-deficient mice. *Gastroenterology* 1997;112:1876–1886.

Received April 11, 2007. Accepted November 15, 2007.

Address requests for reprints to: Hiromichi Ishikawa, MD, PhD, Department of Microbiology and Immunology, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan. e-mail: h-ishika@sc.itc.keio.ac.jp; fax: (81) 3-5360-1508.

Supported in part by a Grant-in-Aid for Creative Scientific Research, the Japan Society for the Promotion of Science (13GS0015); by Special Coordination Funds for Promoting Science and Technology from the Japanese Ministry of Education, Culture, Sports, Science, and Technology; and by Research on Specific Diseases, Japanese Ministry of Health, Labor, and Welfare (to H.I.); by Keio University Special Grant-in-Aid for Innovative Collaborative Research Projects (to T.H.); by National Institutes of Health grants DK47677 (to A.K.B.) and DK064351 (to A.M.); by the Center for the Study of Inflammatory Bowel Disease, Massachusetts General Hospital; and by 21st Century Center-of-Excellence (COE) Program for Life Science from MEXT (to M.S.).

The authors thank Dr Daniel K. Podolsky for his many valuable discussions and comments, Dr Leo Lefrancois for his critical reading of the manuscript, and Dr Atsuhiko Ogawa for his excellent technical assistance.

M.N. and Y.K. contributed equally to this work.

Y.K.'s current location is Department of Immunology, Kinki University School of Medicine, Osaka, Japan.

H.Y.'s current location is Health Research Foundation, Kyoto, Japan.

Conflicts of interest: No conflicts of interest exist.

T.N. was a research fellow supported by 21st Century COE Program for Life Science from MEXT and is now supported by Global COE Program for Human Metabolomic Systems Biology from MEXT (to M.S.).

Hiromichi Ishikawa
Tomoaki Naito
Toshihiko Iwanaga
Hiromi Takahashi-Iwanaga
Makoto Suematsu
Toshifumi Hibi
Masanobu Nanno

Authors' address

Hiromichi Ishikawa¹, Tomoaki Naito^{1,2}, Toshihiko Iwanaga³,
Hiromi Takahashi-Iwanaga³, Makoto Suematsu²,
Toshifumi Hibi⁴, Masanobu Nanno⁵

¹Department of Microbiology and Immunology,
Keio University School of Medicine, Shinjuku-ku,
Tokyo, Japan.

²Department of Biochemistry and Integrative
Medical Biology, Keio University School of
Medicine, Shinjuku-ku, Tokyo, Japan.

³Laboratory of Cytology and Histology, Graduate
School of Medicine, Hokkaido University, Kita-
ku, Sapporo, Japan.

⁴Department of Internal Medicine, Keio University
School of Medicine, Shinjuku-ku, Tokyo, Japan.

⁵Yakult Central Institute for Microbiological
Research, Kunitachi, Tokyo, Japan.

Correspondence to:

Hiromichi Ishikawa
Department of Microbiology and Immunology
Keio University School of Medicine
35 Shinanomachi, Shinjuku-ku
Tokyo 160-8582, Japan
Tel.: 81 3 5363 3766
Fax: 81 3 5360 1508
E-mail: h-ishika@sc.itc.keio.ac.jp

Acknowledgements

We would like to thank all members of our laboratories who have contributed to portions of our work embodied in this review paper. The data provided in the preparation of this review were supported in part by a Grant-in-Aid for Creative Scientific Research, the Japan Society for the Promotion of Science (13GS0015), by Special Coordination Funds for Promoting Science and Technology from the Japanese Ministry of Education, Culture, Sports, Science and Technology, by Research on Specific Diseases, Japanese Ministry of Health, Labor and Welfare, and by Keio University Special Grant-in-Aid for Innovative Collaborative Research Projects (T.H.). T.N. is a research fellow supported by the 21st Century Center-of-Excellence Program for Life Science from MEXT (M.S.).

Immunological Reviews 2007

Vol. 215: 154–165

Printed in Singapore. All rights reserved

© 2007 The Authors
Journal compilation © 2007 Blackwell Munksgaard
Immunological Reviews
0105-2896

Curriculum vitae of intestinal intraepithelial T cells: their developmental and behavioral characteristics

Summary: The alimentary tract has an epithelial layer, consisting mainly of intestinal epithelial cells (IECs), that is exposed to the exterior world through the intestinal lumen. The IEC layer contains many intestinal intraepithelial T cells (IELs), and the total number of IELs constitutes the largest population in the peripheral T-cell pool. Virtually all $\gamma\delta$ -IELs and many $\alpha\beta$ -IELs in the mouse small intestine are known to express CD8 $\alpha\alpha$ homodimers. A wide range of evidence that supports extrathymic development of these CD8 $\alpha\alpha$ ⁺ IELs has been collected. In addition, while several studies identified cells with precursor T-cell phenotypes within the gut epithelium, how these precursors, which are dispersed along the length of the intestine, develop into $\gamma\delta$ -IELs and/or $\alpha\beta$ -IELs has not been clarified. The identification of lymphoid cell aggregations named 'cryptopatches' (CPs) in the intestinal crypt lamina propria of mice as sites rich in T-cell precursors in 1996 by our research group, however, provided evidence for a central site, whereby precursor IELs could give rise to T-cell receptor-bearing IELs. In this review, we discuss the development of IELs in the intestinal mucosa and examine the possibility that CPs serve as a production site of extrathymic IELs.

Keywords: CD8 $\alpha\alpha$ ⁺-IEL, $\gamma\delta$ -IEL, $\alpha\beta$ -IEL, extrathymic development of IEL, cryptopatches

Introduction

Surfaces in the body in contact with the outside world include the epidermis and the mucous epithelia. Directly below both the epidermis and the mucous epithelia is an extensive basement membrane (Bm) that serves as a thin wall separating them from the interior of the body, and no capillaries or lymphatic vessels are present in the epidermis and mucous epithelia. Therefore, any lymphomyeloid cells distributed in epidermis and mucous epithelia are extravasated from the postcapillary venules in the interior of Bm, and they must move into the epidermis and mucous epithelia by crossing Bm. T cells and B cells evolved as key players in the immune system of vertebrates, and an infinite number of antigen-specific receptors are produced by a mechanism called somatic gene rearrangement. It has been known for some time that lymphocytes are distributed in the epidermis and mucous epithelia, and in about the middle of the 1970s, it became clear that most intestinal intraepithelial lymphocytes settling in the small intestine of mice are T cells [intestinal

intraepithelial T cells (IELs)] (1). Furthermore, almost all T cells in the epidermis of laboratory mice are those expressing homogenous $\gamma\delta$ -type T-cell receptors (TCRs), also known as dendritic epidermal T cells (DETCs). The surprising finding concerning these $\gamma\delta$ -DETCs is that they are produced in the thymus at about day 15 of embryonic life and are thus derived from the first wave of fetal $\gamma\delta$ thymocytes (2). In this review article, we focus our discussion mainly on findings obtained in mice concerning development of IELs distributed among intestinal epithelial cells (IECs).

Surprising evidence, showed by studies using a monoclonal antibody to TCR, is that almost all mouse IELs are T cells (3–9). IELs are radically different from T cells residing in other sites of the body; most of them are ill-defined T cells with unusual but distinctive characteristics. These cells are located at the front line of defense, at the point which the interior of the body comes in contact with the greatest numbers of antigens from the exterior world.

The interior of the Bm consists of lamina propria (LP) that contains abundant immunoglobulin A (IgA)⁺ B cells, CD3⁺ T cells (Fig. 1), and various lymphomyeloid cells. In contrast (as discussed later), the exterior of the Bm contains an IEC layer with prominent colonization of CD8 α -expressing T cells (Fig. 1). The marked differences between the inside and the outside of the Bm are very important in connection with clarification of *in vivo* physiological functions and development of IELs on the front line of the intestinal mucosa. Research over the past 30 years has shown that IELs in mice and humans, especially those in the small intestine of mice, are a phenotypically and functionally distinctive subpopulation of peripheral T cells that is distinguished from so-called proper T cells, which are distributed in peripheral lymphoid tissues such as the spleen and lymph nodes (LNs) after development in the thymus (10). A vast majority of T cells found in peripheral lymphoid tissues of mice and humans are $\alpha\beta$ T cells, while only a few $\gamma\delta$ T cells are present. In contrast, IELs in mice and humans include large numbers of cells expressing $\alpha\beta$ TCRs and those expressing $\gamma\delta$ TCRs. From a study of IELs in athymic (*nu/nu*) mice, it is clear that many $\gamma\delta$ -IELs are present, although the population size is decreased. In addition, in spite of the sharp decrease in $\alpha\beta$ -IELs, meaningful numbers of these cells can be detected. Thus, a substantial proportion of $\gamma\delta$ -IELs seems to be generated and/or expanded in the absence of the thymus. In contrast, it is well known that both $\gamma\delta$ T cells and $\alpha\beta$ T cells are virtually undetectable in the spleen and LNs of *nu/nu* mice (11, 12). Functional aspects of IELs have been adequately explained in other reviews in this volume, and this review contains personal insights concerning the past, present, and future of extrathymic




Fig. 1. Immunohistochemical visualization of IgA-, CD8 α -, and CD3-expressing cells in jejunal villi. Note that IgA⁺ B cells are localized exclusively in the LP, whereas that CD8 α ⁺ T cells, namely intestinal IELs, are compartmentalized above the Bm in the IEC layer of the small intestine. In contrast, in addition to numerous IELs in the IEC layer, CD3⁺ T cells, mostly CD4⁺ T cells, are also found in the LP of the villi.

development of IELs and where the research is heading. Furthermore, we discuss how IELs settle down in the IEC layer through Bm and emphasize how they behave and survive in the IEC layer *in situ*.

IEL development in the intestinal epithelium: evolutionary perspective

The intestine was the first organ to appear when animals became multicellular; even though some multicellular animals lacked brains, there were none without intestines. To defend the intestines against pathogenic microorganisms and harmful substances from the exterior, macrophage-like lymphoid cells developed directly under the intestinal epithelium. The first organ to appear in our living body is the primordial gut, and many organs, including lungs, liver, pancreas, and thyroid gland, are derived from this apparatus. Some marine animals breathe through gills that develop from the upper digestive tract, and pulmonary respiration evolved with the change from marine to terrestrial life. It is well known that the thymus was derived from part of the gill. Therefore, all these organs have

a kindred relation, and latent production of lymphocytes appears possible. In agreement with this argument, fetal liver of mammals including mice is a primary lymphoid organ producing lymphomyeloid cells.

The IEC layer was proposed as a lymphocyte-producing organ as early as 1967 (13). Gut-associated lymphoid tissue (GALT), which contains about 60% of all peripheral lymphocytes, monitors and defends the intestinal mucosa in most vertebrates. T cells and antibodies, the key players in adaptive immunity, have not been found in the jawless fish *Agnatha*, the oldest phylogenetic vertebrate lacking a thymus, spleen, and LNs (14, 15). However, GALT, characterized by many lymphoid cells, is found in *Agnatha* such as lampreys and hagfish, and the intestinal mucosa of such animals appears to serve as lymphocyte production sites (14, 15). Furthermore, the bursa of Fabricius, a GALT of birds, and Peyer's patches (PPs) of ruminants are also well known as primary lymphoid tissues responsible for the development of B cells (16). If we consider the large amount of knowledge based on animal evolution, there is nothing remarkable about development of IELs in intestinal mucosa in situ in mice and humans.

Findings supporting extrathymic development of murine IELs

Type a and type b IELs

IELs of the murine gut have been identified as ill-defined T cells that lurk in the anatomical front of the intestine (1, 3-9), with a primarily cytotoxic T-cell phenotype (4, 17-19). A large fraction of murine IELs bearing CD8 $\alpha\alpha$ homodimers (CD8 $\alpha\alpha$ ⁺ IELs) have been proposed to originate locally through a differentiation process initiated in c-kit⁺IL-7R⁺ lineage marker (Lin)⁻ gut precursors. Hayday et al. (10) have proposed that the functional complexity and phenotype heterogeneity of IELs might be simplified if IELs are classified into just two cell types: 'a' and 'b'. Type a includes CD4⁺ and CD8 $\alpha\beta$ ⁺ $\alpha\beta$ -IELs that primarily recognize antigens presented by classical major histocompatibility complex (MHC) class I and class II molecules and are primed within the systemic circulation. Type b IELs include CD8 $\alpha\alpha$ ⁺ $\alpha\beta$ - and $\gamma\delta$ -IELs that respond to antigens not restricted by classical MHC molecules. Although CD8 $\alpha\alpha$ ⁺ $\alpha\beta$ - and $\gamma\delta$ -IELs are clearly different from one another, type b IELs share many 'unconventional' features that distinguish them from type a IELs.

Although dependence of the type a CD8 $\alpha\beta$ ⁺ and type b CD8 $\alpha\alpha$ ⁺ $\alpha\beta$ -IELs but not type b CD8 $\alpha\alpha$ ⁺ $\gamma\delta$ -IELs on MHC class I molecules was reported using β 2-microglobulin (β 2m)-deficient mice (20, 21), a recent analysis of gene expression

profiles between type b CD8 $\alpha\alpha$ ⁺ $\alpha\beta$ - and $\gamma\delta$ -IELs showed a high degree of similarity (22). These two classes of IELs are not only related functionally but also have a kindred relation.

The total number of $\alpha\beta$ -IELs decreased sharply in β 2m^{-/-} mice due to the disappearance of both CD8 $\alpha\beta$ ⁺ (type a) and CD8 $\alpha\alpha$ ⁺ (type b) subsets. In β 2m/TCR- δ double-mutant mice, which lack β 2m and $\gamma\delta$ -IELs, the CD8 $\alpha\alpha$ ⁺ subset expanded dramatically, while the CD8 $\alpha\beta$ ⁺ subset did not (Fig. 2). Thus, in the absence of $\gamma\delta$ -IELs, $\alpha\beta$ -IELs in β 2m-deficient mice outnumbered those in wildtype littermates due to considerable expansion of type b CD8 $\alpha\alpha$ ⁺ $\alpha\beta$ -IELs (Fig. 2). These results (23) indicate that generation of type b CD8 $\alpha\alpha$ ⁺ $\alpha\beta$ - and $\gamma\delta$ -IELs is essentially β 2m independent, while generation of type a CD8 $\alpha\beta$ ⁺ $\alpha\beta$ -IELs is highly dependent on β 2m-MHC class I




Fig. 2. Composition of $\alpha\beta$ -IEL subsets in wildtype (WT), β 2m-deficient, TCR- δ mutant (δ), and β 2m \times TCR- δ double-mutant (β 2m \times δ) mice. These four different mice were littermates of the F₂ generation of an intercross between β 2m^{-/-} and δ ^{-/-} mice. IELs isolated from these mutant mice were incubated first with anti-CD8 α monoclonal antibody (biotinylated) and then with streptavidin-allophycocyanin. After washing, the IELs were counterstained with two combinations of two phycoerythrin-conjugated monoclonal antibodies (anti-CD4 and anti-CD8 β) and two fluorescein-isothiocyanate-conjugated monoclonal antibodies (anti- $\alpha\beta$ TCR and anti- $\gamma\delta$ TCR, respectively). Absolute numbers of double-negative (CD4⁻CD8⁻), single positive (CD4⁺, CD8 $\alpha\alpha$ ⁺, or CD8 $\alpha\beta$ ⁺), and double positive (CD4⁺CD8⁺) subsets in the $\alpha\beta$ -IEL population were calculated on the basis of total number of $\alpha\beta$ -IELs. Note that CD8 $\alpha\alpha$ ⁺ and CD8 $\alpha\beta$ ⁺ $\alpha\beta$ -IEL subsets are absent from the small intestine of β 2m mutant mice, whereas the CD8 $\alpha\alpha$ ⁺ but not CD8 $\alpha\beta$ ⁺ $\alpha\beta$ -IEL subset expands markedly in the small intestine of double-mutant β 2m \times δ mice, namely β 2m mutant mice that lack $\gamma\delta$ -IELs.

molecules expressed by the controlling cells at the type a IEL precursor development site. These findings suggest the possibility that type b IELs, CD8 α^+ $\alpha\beta$ - and $\gamma\delta$ -IELs, develop in the same anatomical site(s). When no $\beta 2m$ -MHC class I molecules are present, development of CD8 α^+ $\alpha\beta$ -IELs is likely inhibited because development of CD8 α^+ $\gamma\delta$ -IELs surpasses that of CD8 α^+ $\alpha\beta$ -IELs competitively.

CD8 α exerts a specific and high affinity for interaction with the non-classical MHC class I molecule thymus leukemia (TL) antigen, which is expressed abundantly by murine thymic stromal cells and by IECs (24). It was also proposed that CD8 α^+ TCR- $\alpha\beta$ T cells originated from the thymus through agonist-dependent positive selection (25). The mechanism of development of CD8 α^+ T cells and *in vivo* physiological functions, including whether or not this scenario is correct, remain to be clarified.

Evidence obtained in a study of athymic *nu/nu* mice

The evidence that most clearly supports thymus-independent development of gut-oriented type b IELs is obtained from a study of T cells in the athymic (*nu/nu*) mouse. Almost no $\gamma\delta$ T cells or $\alpha\beta$ T cells are observed in the spleen and LNs of *nu/nu* mice. A considerable population of $\gamma\delta$ -IELs is present in IELs of *nu/nu* mice, and $\alpha\beta$ -IELs can also be detected (Fig. 3). Since these $\alpha\beta$ -IELs are CD8 α^+ type b IELs and no $\alpha\beta$ -IELs are found in TCR- $\beta^{-/-}$ mice (Fig. 3), it is evident that a few type b $\alpha\beta$ -IELs develop independently of the thymus.

Many reports have been published on the thymus-independent development of type b IELs. These findings include the presence of a few CD3 $^-$ lymphocytes in the IELs and the




Fig. 3. Composition of $\alpha\beta$ - and $\gamma\delta$ -IELs isolated from wildtype (WT), athymic (*nu/nu*) and TCR- $\beta^{-/-}$ mice. Flow cytometric analysis of IELs isolated from five individuals each of three different strains of mice was performed, and the representative profiles of IELs are presented. In this case, absolute numbers of IELs recovered were 5.4×10^6 from WT mice, 2.3×10^6 from *nu/nu* mice, and 5.3×10^6 from TCR- $\beta^{-/-}$ mice. The percentage of $\alpha\beta$ - and $\gamma\delta$ -IELs in the corresponding quadrants is shown. Note that $\alpha\beta$ -IELs are drastically reduced in the athymic condition. Nevertheless, a meaningful number of $\alpha\beta$ -IELs are still present in the small intestine of the *nu/nu* mouse compared with the total absence of $\alpha\beta$ -IELs from the TCR- $\beta^{-/-}$ mouse. In contrast, there are few, if any (<1%), $\alpha\beta$ T cells in the spleen and MLNs of this same *nu/nu* mouse (data not shown).

resulting possession of precursor T-cell-like properties, i.e. the fact that these tentative precursors observed in mice and humans retain messenger RNA (mRNA) for recombination-activating gene-1 (RAG-1) and RAG-2 and pre-T α molecules. For details, the reader is referred to previously published articles (8, 9, 26–30).

Here, we introduce our results (31) from screening lymphoid tissues of athymic (*nu/nu*) RAG-1^{GFP/+} animals, which have the green fluorescence protein (GFP) gene in the RAG-1 locus. Only CD19⁺ B cells (32) in the bone marrow (BM), spleen, mesenteric lymph nodes (MLNs), and PPs express RAG-1, while in IELs, a meaningful number of CD19⁻ cells express RAG-1, although the amount of RAG-1 molecules expressed is low (Fig. 4). Since CD19⁻RAG-1^{low} IELs are CD3⁻ and are not observed in IELs of RAG-1^{+/+} mice, this finding supports the distribution of small numbers of RAG-1^{low} precursor T cells in IELs. DETCs of wildtype mice that express homogenous $\gamma\delta$ TCRs (V γ 5J γ 4C γ 1 and V δ 1J δ 2C δ) are known to be derived from the first wave of fetal $\gamma\delta$ thymocytes, which are produced in the thymus at about antenatal day 15 (2). Therefore, even though V γ 5⁺ DETCs are not present naturally, it has been shown that V γ 1/V δ 6⁺ DETCs are present in the epidermis of *nu/nu* mice (33). These findings together with those reported by Matis et al. (34) and Yoshikai et al. (35) prove that the thymus-dependent type a subset (2) and the thymus-independent type b subset (33–35) such as $\gamma\delta$ T cells are present in peripheral anatomical sites other than the IEC layer.

A new member of GALT: cryptopatches

Discovery of cryptopatches in mouse small intestine as the precursor IEL-producing site

We have shown that multiple tiny lymphoid cell aggregations, filled with about 1000 closely packed c-kit⁺IL-7R⁺Thy1⁺ CD3⁻B220⁻ lymphocytes, colonize throughout the small intestinal mucosa of C57BL/6 mice (36). The location is in the crypt LP [cryptopatches (CPs)]. They are first detected in the third week of postnatal life in C57BL/6 mice. In terms of morphogenesis, cellular composition, and fine tissue structure, neither PPs nor isolated lymphoid follicles (ILFs) are identical to CPs (30, 37). CPs contain neither cells undergoing apoptosis nor cells bearing RAG-1 molecules but do contain dendritic stromal cells bearing CD11c/CD18 molecules. The presence of transcripts for germ line TCR genes and mRNA for proteins involved in TCR rearrangement (38) and the ability of c-kit⁺Lin⁻ CP cells to generate TCR⁺ IELs in T-cell-deficient mice (39) indicate that at least some CP cells are committed to the T-cell lineage and are competent for generation of




Fig. 4. A small number of RAG-1-expressing lymphocytes are present in the intestinal intraepithelial compartment of *nu/nu* mice. Lymphocytes from BM, spleen, MLNs, PPs, and IEC layer (IELs) were isolated from *nu/nu* mice carrying a GFP gene in place of the RAG-1 gene (*nu/nu* RAG-1^{GFP/+} mice) (31) and then were subjected to flow

cytometric analysis. Note that only in the IEL preparation, a meaningful number of GFP-dull-positive cells are present. Importantly, these cells are not B-lineage cells because they are CD19 negative.

thymus-independent type b IELs, especially CD8 α ⁺ γ δ -IELs (12).

To obtain direct evidence for generation of type b IELs from the precursors that settle in CPs, cytokine receptor γ chain mutant *nu/nu* mice that lack a thymus, PPs, CPs, and intestinal T cells (40) were reconstituted with wildtype Ly5.1⁺ BM cells. BM-derived TCR⁻ IELs first appeared within villous epithelia of small intestine overlaying regenerated CPs, and these TCR⁻ IELs subsequently emerged throughout the epithelia. Thereafter, TCR⁺ IELs increased sluggishly to a number comparable with that in athymic *nu/nu* mice and consisted of both $\alpha\beta$ - and $\gamma\delta$ -IELs (38). Taking all these results together, CPs are the first murine GALT to be identified that most likely serve as the site of development of lymphohemopoietic precursors for type b IEL descendents at commencement of weaning (36, 38, 40). Recent data showed a clonal relationship between CP T cells and $\gamma\delta$ -IELs (41).

Evidence that CPs are not the precursor IEL-producing site. By characterizing phenotypically distinct lineage-negative populations in the CPs and gut epithelium, Lambolez *et al.* (42) showed that only 3% of CP cells were clearly involved in T-cell differentiation and suggested that these CP structures may have an additional physiological role in the gut. In contrast, Guy-Grand *et al.* (43) clarified the following from an examination of RAG-2 expression using GFP transgenic mice evaluated by GFP (carrying a GFP reporter gene driven by the RAG-2 promoter). In *nu/nu* mice, T lymphopoiesis occurs mainly in MLNs, less in PPs, and not in CPs. Importantly, this extrathymic T lymphopoiesis is totally repressed in euthymic mice (43). Based on these findings, Guy-Grand *et al.* (43) concluded that in normal euthymic mice, all gut $\alpha\beta$ -IELs, including type b CD8 α ⁺ $\alpha\beta$ -IELs, are of thymic origin. It has also been shown that thymus transplantation into *nu/nu* mice results in the appearance of thymus graft-derived $\alpha\beta$ - and $\gamma\delta$ -IELs in *nu/nu* recipients (44). To evaluate this important issue, we generated *nu/nu* *aly/aly* (*alyphoplasia*) double-mutant mice

lacking thymus, all LNs, PPs, and ILFs but possessing CPs (12). Substantial colonization by $\gamma\delta$ -IELs comprising the major CD8 α ⁺ subset took place, and use of TCR- γ -chain variable gene segments by these $\gamma\delta$ -IELs was unaltered (12). These findings indicate that MLNs and PPs are not an absolute requirement for development of $\gamma\delta$ -IELs but instead support the notion that gut CPs generate progenitor $\gamma\delta$ -IELs, even under athymic conditions. However, absolute numbers of $\gamma\delta$ -IELs from *nu/nu* *aly/aly* mice are smaller than those from the corresponding control *nu/nu* mice (12). These features indicate that LNs and PPs in fact determine the number of $\gamma\delta$ -IELs under the *nu/nu* conditions. In any event, there would be a hierarchy of T-cell production in terms of anatomical sites, although the detailed mechanism is still not well understood. By a mechanism such as the tropic effect of thyrotropin-releasing hormone on IEL development (45, 46), however, almost all $\alpha\beta$ -IELs appear to originate in the thymus in euthymic mice. In *nu/nu* mice, however, IELs appear to be generated mainly by the MLNs and PPs, and in *nu/nu* *aly/aly* mice that lack a thymus, all LNs, PPs, ILFs, and $\alpha\beta$ - and $\gamma\delta$ -IELs appear to be generated from other anatomical sites, probably CPs. In this context, it has been reported that T-cell-committed precursors are distributed not only in the thymus but also in the BM, liver, and spleen (47–49). It is inferred based on these findings that extrathymic T-cell generation has not been proven to be repressed completely in normal euthymic mice. In other words, there is no solid evidence denying a possibility that extrathymic T cells are generated in euthymic conditions.

It has recently been shown that retinoic acid-related orphan receptors (ROR γ t) detected in fetal lymphoid tissue-inducer (Lti) cells are also expressed in cells within gut CPs and that, by fate mapping of ROR γ t⁺ cells, type b IELs, such as $\gamma\delta$ -IELs, are not the progeny of ROR γ t⁺ CP cells (50). However, it remains an open question whether a small fraction of lymphoid cells in CPs does not express ROR γ t or all CP cells express ROR γ t. To investigate this point in detail, we generated *nu/+* ROR γ t^{GFP/+}

mice and *nu/nu* ROR γ t^{GFP/+} mice and obtained the results shown in Fig. 5.

In agreement with the report of Eberl and Littman (50), many ROR γ t⁺ (GFP⁺) cells were present in thymocytes from *nu/+* ROR γ t^{GFP/+} mice, although GFP expression was weak (unpublished observation). Analysis of CP cells from *nu/nu* ROR γ t^{GFP/+} mice showed that almost all GFP⁺ cells were interleukin-7 receptor positive (IL-7R)⁺ and CD8 α ⁻. However, CD3⁻ cell subsets showing various phenotypes such as IL-7R⁺GFP⁻, c-kit⁺GFP⁺, c-kit⁺GFP⁻, c-kit⁻GFP⁺, Thy-1⁺GFP⁺, Thy-1⁺GFP⁻, Thy-1⁻GFP⁺, CD4⁺GFP⁺, CD4⁺GFP⁻, CD4⁻GFP⁺, CD8⁺GFP⁻, Lin⁺GFP⁺ (majority of them CD4⁺GFP⁺), Lin⁺GFP⁻, and Lin⁻GFP⁺ were all present but with variable population sizes (Fig. 5A). In addition to the colonization of $\gamma\delta$ -IELs and small numbers of $\alpha\beta$ -IELs, of particular note was the presence of CD3⁻ IELs showing c-kit⁺GFP⁺ and IL-7R⁺GFP⁺ phenotypes in IEC compartments of *nu/nu* ROR γ t^{GFP/+} mice (Fig. 5B). These new findings do not necessarily support the conclusion of Eberl and Littman (50) and show distribution of ROR γ t⁻ c-kit⁺, IL-7R⁺, Thy-1⁺, CD4⁺, CD8⁺, and/or Lin⁺ lymphocytes in murine gut CPs. In this context, we previously reported (36) that CPs are not detectable in IL-7R^{-/-} mice. However, although $\gamma\delta$ -IELs are absent owing to selective blockage of TCR- γ gene rearrangements (51), we noticed only a slight decrease in development of

type b $\alpha\beta$ -IEL subsets in IL-7R^{-/-} mice. With these observations in mind, we reinvestigated hundreds of cryosections prepared from small intestines of IL-7R^{-/-} mice by immunohistochemistry and verified that conspicuously emaciated CPs filled with c-kit⁺ cells and decreased by more than 16-fold in number were present in this mutant intestine (40). Similarly, although mice genetically deficient in lymphotoxin α (LT α) have been reported to lack CPs (52), we observed that histogenesis of CPs and intestinal development of $\alpha\beta$ - and $\gamma\delta$ -IELs remained almost intact in LT α ^{-/-} mice (37). In consideration of our research results, i.e. CPs are observed not only in IL-7R^{-/-} mice but also in LT α ^{-/-} mice, the conclusion that CP development is not found in ROR γ t^{GFP/GFP} mice lacking ROR γ t (50) seems to require careful reexamination. It is easy to ascertain the presence of CPs, but it is quite difficult to conclude that CPs are totally absent. At the same time, we should examine if type b IELs, such as $\gamma\delta$ -IELs, develop from the ROR γ t⁻ c-kit⁺GFP⁻ or IL-7R⁺GFP⁻ subset (Fig. 5A) distributed in the CPs of *nu/nu* ROR γ t^{GFP/+} mice.

What are these CPs?

Based on the results obtained using ROR γ t^{GFP/+} mice, Eberl and Littman (50) proposed that the principal function of murine c-kit⁺Lin⁻ ROR γ t⁺ CP cells is to induce formation of lymphoid




Fig. 5. Flow cytometric analysis of CP cells and IELs isolated from *nu/nu* mice carrying the wildtype ROR γ t genes (ROR γ t^{+/+} mice) and carrying a GFP gene in place of ROR γ t gene (ROR γ t^{GFP/+} mice). CP cells were isolated according to the method described previously (37). Although CP cells (A) and IELs (B) isolated from *nu/nu* ROR γ t^{+/+} mice do not contain GFP⁺ cells, those isolated from *nu/nu* ROR γ t^{GFP/+} mice contain a substantial population of GFP⁺ ROR γ t-expressing

cells. Furthermore, these ROR γ t-expressing cells from CP and IEL compartments of *nu/nu* ROR γ t^{GFP/+} mice appear to be composed of two discrete ROR γ t^{high} and ROR γ t^{low} cell subsets. 'Lin' in A represents lineage markers CD3, B220, Mac-1, Gr-1, TER119, CD11c, CD4, and CD8 α . Lymphocytes isolated from spleen, MLNs, and PPs of *nu/nu* ROR γ t^{GFP/+} mice lack GFP⁺ ROR γ t-expressing cells (data not shown).

follicles, namely ILFs, in the LP in a manner similar to induction of LNs and PPs by ROR γ t⁺ Lti cells. It is also possible that CP cells are precursor cells of gut-oriented lymphomyeloid cells other than IELs (42). The CPs of mice have been discounted as the anatomical site where precursor IELs congregate, as no CP-like lymphoid clusters have been reported in the intestinal mucosa of mammals other than mice. However, since evidence indicates development of gut-oriented T cells, mainly IELs, in the intestinal mucosa of humans (53–59) and rats (60), to determine whether or not CPs and CP-like lymphoid tissues are present in enteric mucosa of animals is an important goal for future experiments. We (61) showed that lymphocyte clusters, just like the structure named lymphocyte-filled villi (LFV) (62), populated predominantly with c-kit⁺IL-7R⁺ cells and less with $\alpha\beta$ TCR cells, were found distributed throughout the length of the small intestine of rats. Nevertheless, we were unable to verify whether these rat LFV, containing undifferentiated lymphocytes, represented clusters of extrathymic precursor T cells. With regard to this same issue, several groups have actively sought evidence of CPs in the human gastrointestinal tract, illuminating distinctive T-cell facets of human fetal gut lymphocytes (58, 59). In agreement with these findings, we have identified multiple tiny c-kit⁺ lymphoid cell clusters in human fetal intestine at the second trimester of fetal life (unpublished observation). Overall, we take it for granted that the differentiation of type b IELs is not exactly the same among different vertebrates. For instance, if epidermal immune regulation by $\gamma\delta$ -DETCs in mice is of considerable physiological importance, how do other mammals including humans cope without $\gamma\delta$ -DETCs? In these animals, immunoregulatory function should be provided by some lymphoid cells other than $\gamma\delta$ -DETCs.

Based on these findings, precursors of type b IELs that develop extrathymically in humans and rats appear to be produced in CP-like lymphoid tissues of mice during fetal life. Then, the precursors are dispersed throughout the LP or IEC layer, or self-renewal is localized in the IEC layer after expression of the $\alpha\beta$ TCR or $\gamma\delta$ TCR and completion of development, as in the case of B-1 B cells. Mouse DETCs (2) expressing homogenous V γ 5/V δ 1⁺ $\gamma\delta$ TCRs are produced only in fetal thymus in a very limited period at about day 15 of fetal life. No such V γ 5/V δ 1⁺ $\gamma\delta$ -DETCs are found in the epidermis of athymic nu/nu mice (33), while homogenous V γ 5/V δ 1⁺ $\gamma\delta$ -DETCs of euthymic mice are present throughout life (1.5–2 years). If the supply of V γ 5/V δ 1⁺ $\gamma\delta$ -DETCs occurs only in the first wave of fetal $\gamma\delta$ thymocytes development, the life of V γ 5/V δ 1⁺ $\gamma\delta$ -DETCs must be as long as the life of the mouse. Is this really the case? We do not think that it is. It is possible that

precursor V γ 5/V δ 1⁺ $\gamma\delta$ -DETCs generated by fetal thymus at about day 15 of fetal life lurk somewhere in the body (such as in the epidermis or dermis), develop at a fixed pace, and produce new V γ 5/V δ 1⁺ $\gamma\delta$ -DETCs. It is also possible that $\gamma\delta$ -DETCs expressing homogenous V γ 5/V δ 1-TCRs continuously undergo self-renewal in the epidermis. Indeed, $\gamma\delta$ -DETCs are shown to recognize and respond to antigens expressed on damaged, stressed, or transformed keratinocytes by means of their TCRs and produce keratinocyte growth factors (63), and it has been indicated that continuous stimulation of V γ 5/V δ 1-TCR⁺ $\gamma\delta$ -DETCs by the relevant ligand(s) is critical for maintenance of $\gamma\delta$ -DETCs throughout the life of animals (64). In this context, it is conceivable that extrathymically generated mature human and rat IELs, which originate from cells settling in CP-like lymphoid tissues of mice during fetal life, undergo continuous self-renewal over the entire postnatal life of these animals by stimulation with relevant gut-associated antigens. In conclusion, although much remains to be learned about the mysterious development of extrathymic T cells in general and also the enigmatic features of CPs before we evaluate them as anatomical sites in which murine type b precursor IELs develop, many aspects of extrathymic T-cell immunobiology are now coming together step-by-step.

Where do these IELs come from?

Many types of cells are present outside the Bm of the small intestine (Fig. 6). Cells other than IELs present in the IEC layer are known to develop from stem cells located in the crypt. As described in the Introduction, IELs must migrate into the IEC layer from LP across the Bm. Immunohistochemical examination has shown that there are at least 2, or many more, IELs for every 10 IECs in mouse small intestine (65, 66). Thus, absolute numbers




Fig. 6. Schematic illustration of cells that settle in the IEC layer of the small intestine. Every cell that settles in the IEC compartment, except IELs on the right of this figure (magnified), is the progeny of the stem cells shown on the left of this figure. Endocrine cells a and b contain different gut hormones.

of murine IELs are estimated to account for about half of the peripheral T-cell pool (67). However, it is not known if this huge IEL population enters the IEC layer across the Bm at sites such as crypts or villi, as shown in Fig. 6, or if IELs that have entered the IEC layer later cross the Bm and return to the LP. A large fraction of IELs is composed of $\gamma\delta$ -IELs, and the fact that $\gamma\delta$ T cells are almost absent from the LP compartment of villi in normal mice indicates that once they have entered the IEC layers, IELs and/or precursor IELs might return to LP in rare cases. By electron microscopy, we (39) showed that numerous lymphocytes cross the Bm that comes into contact with the CPs. Furthermore, by analysis of BM-chimeric mice, we (38) verified that donor BM-cell-derived IELs first appeared within the IEC layer of villi around regenerated CPs filled with BM-cell-derived c-kit⁺ cells. These experimental results, in conjunction with a wide range of evidence showing the presence of small numbers of CD3⁻ precursor IEL-like lymphocytes in the IEC compartment (8, 9, 26–30, 68), suggest a scenario in which precursor IELs that developed in CPs enter the IEC layer from the Bm that overlays CPs, and then these cells very sluggishly develop into mature type b IELs. In contrast, no definite findings have been obtained on whether type a $\alpha\beta$ -IELs, such as CD4⁺ or CD8 $\alpha\beta$ ⁺ T cells derived from the thymus, cross the Bm by some route and enter the IEC layer.

Parabiotic C57BL/6 Ly5.1 and Ly5.2 mice sharing circulation have been used by several groups (69, 70) to examine whether c-kit⁺ stem cells that settle in local organs give rise to lymphoid cells in these organs in situ. As anticipated, the partner cells mixed rapidly in the spleen, all LNs, and PPs. In contrast, there were no or very few mixtures of partner cells in the thymus, CPs, and IELs (70). Poussier *et al.* (69) reported that IELs in the murine intestine did not mix together in parabiotic Ly5.1 and Ly5.2 mice. Overall, these findings appear to once again support the notion that IELs arise from their own pre-existing local precursor cells and that neighboring CPs continuously supply these local precursor cells at unknown rates.

What are these IELs?

There are still many riddles concerning the behavior and biological function of IELs that settle in the anatomical front of the IEC compartment in situ. Some of these unresolved issues are discussed below. We also discuss the development of research yet to be undertaken and future perspectives.

Behavior

Epithelial stem cells (Fig. 6) proliferate at the base of the crypts. Newly formed cells move upward and differentiate into various

types of cells, mostly IECs (Fig. 6), in a process called migration-associated differentiation. The entire process of migration toward the top of the villi takes only several days, and the cells die there, most likely by apoptosis (71).

Electron microscopic examination of immersion-fixed tissue sections shows that IEL interdigitate tightly with IEC at the basolateral faces of IEC (Fig. 7A). Most IELs are terminally differentiated cells or those in the G0/G1 phase of the cell cycle.




Fig. 7. Electron microscopic analysis of small intestinal epithelium.

(A) Transmission electron micrograph showing IELs in intestinal epithelium obtained from an immersion-fixed sample. At least five IELs (L, edged with yellow line) are encased within the epithelium. No free space is visible among epithelial cells in the immersion-fixed samples; IELs appear to be tightly packed in the epithelium. (B) Scanning electron micrograph of IELs in a perfusion-fixed material. Two lymphocytes (yellow) equipped with spine-like processes are detected among epithelial cells. Note the broad free space at the basal region of the epithelium. A perfusion-fixed sample retains more precise and physiological three-dimensional structure of the tissues than that obtained by an immersion-fixed sample because the tissues shrink totally during and after the immersion-fixation. [This image is a modified reproduction of our figure that appeared in *Cell & Tissue Research*, used with kind permission of Springer Science and Business Media (83)].

It has been shown by bromodeoxyuridine labeling experiments that the average half-life of murine IELs is 3 weeks or much longer (72). If we assume that IELs are also migrating to the top of the villi, IECs must continuously get ahead of IELs, suggesting that individual IELs are capable of coming in contact with and/or surveying a significant number of IECs. Are IELs really migrating to the top of the villi? If so, at what speed are they migrating? At present, we do not know whether IELs can actually move upward or whether they maintain a rather stationary position.

Fig. 7B shows a vertically fractured face of perfusion-fixed small intestine by scanning electron microscopy. As is clear from this image, the surface in contact with intestinal lumen (apical surface of IEC) forms a tight junction between the IECs and is covered with microvilli with no gaps. However, there are relatively wide gaps between cells just on the Bm, and IELs are not always fixed by tight interdigitation with the lateral faces of the IECs. They appear to move to and fro relatively freely. IELs are also often in contact with IECs through process-like structures (Fig. 7B). In this image, it looks like the IELs are completely covered with trees, as if it were a mighty jungle viewed from the air, but there are considerable gaps near the ground under the trees where lions (\approx IELs) sleep and animals such as leopards (\approx IELs) lurk and move around (Fig. 8). In any case, clarification of the life and behavior of these mature IELs and the small number of precursor IELs in the IEC compartment are important topics for future study.




Fig. 8. A pictorial representation of the mighty jungle of the IEC layer. Based on the three-dimensional scanning electron microscopic picture of the IEC layer presented in Fig. 7B, we illustrated a mighty jungle in which a lion (\approx IEL, with potentially cytotoxic function?) is resting on the ground (Bm) and a leopard (\approx IEL, like an indiscriminate predator?) is moving around a tree trunk (lateral face of IEC). It is evident that there is considerable room between the tightly packed treetop and the surface of the ground where these animals (\approx IELs) might be freely moving to and fro.

Oligoclonality

Whereas IELs are potentially able to use multiple $V\beta$ TCR genes, both human (73–75) and murine (76) $\alpha\beta$ -IELs are known to be derived from a limited number of oligoclonal T-cell clones. This oligoclonality of IELs points to the presence of a restricted set of foreign and/or self-antigens in the gut that may be ligands involved in the stimulation and expansion of these gut T cells. The large numbers of T cells distributed in the spleen and LNs appear to be inexperienced or virgin T cells that have not received antigen stimulation in normal specific pathogen-free mice. In contrast, T cells in the gut of specific pathogen-free mice are activated cells, continuously receiving antigen stimulation. As a result of selection, only competent clones expand and become dominant. Under these circumstances, even though the types of antigens are very diverse and are not limited to a restricted set of foreign and/or self-antigens, the TCR of $\alpha\beta$ -IELs eventually results in oligoclonality. One mysterious and unexpected finding is that genetically identical individuals, even from the same litter and housed in the same cage, show distinct and apparently non-overlapping oligoclonal repertoires of both type a and type b $\alpha\beta$ -IELs (76). Is the immune response to a diverse range of external antigens in the intestinal lumen really involved in the establishment of the oligoclonality of $\alpha\beta$ -IELs? There is no firm experimental evidence to answer this question. In this respect, the finding that $\alpha\beta$ -IELs in germ-free mice are also oligoclonal (77) is very intriguing and shows that $\alpha\beta$ -IELs are oligoclonal, even when huge numbers of intestinal flora-derived antigens and microbe-associated immunoactive substances are not present.

Cytotoxicity

IELs are mostly terminally differentiated and activated T cells that possess a granular cytoplasmic structure containing perforin and granzyme (78), capable of killing Fc-receptor-bearing target cells after bridging them with anti-CD3, anti- $\alpha\beta$ TCR, or anti- $\gamma\delta$ TCR monoclonal antibodies (78–80). However, at most 10^7 IELs per mouse from the small intestine can be isolated for *in vitro* analysis, despite the fact that $5\text{--}10 \times 10^7$ IELs have been found to settle in the IEC compartment by immunohistochemistry (67) (Fig. 1). Therefore, we might have studied *in vitro* only a portion of the IELs, which can be isolated easily from intestinal mucosa as a subpopulation maintaining their typical granular cytoplasmic structure and cytotoxic activity. In this context, of great importance in the future is to examine *in vitro* characteristics of the large number of IELs that cannot be recovered using current techniques and/or are lost in the process of purification in a test tube. Such a 'missing' population should be examined,

if the cells maintain their characteristics of terminal differentiation. $\gamma\delta$ -IELs from germ-free mice are known to show cytotoxicity (81) in the same way as $\alpha\beta$ -IELs in germ-free mice show oligoclonality. Electron and light microscopy shows that murine IELs from *scid/scid* (severe combined immunodeficient) mice that are unable to generate $\alpha\beta$ - and $\gamma\delta$ -IELs have granulated IELs similar to those in normal mice (82), are $CD3^-CD8\alpha^+$ IELs are present in *scid/scid* mice and *nu/nu scid/scid* mice (40). In addition, findings showing that $CD3^-CD8\alpha^+$ IELs are not present in cytokine receptor γ chain mutant *nu/nu* mice lacking CPs (38, 40) suggest that a special but as yet unknown microenvironment in CP and IEC compartments could dictate the ability of precursor IELs entering the IEC layer to possess a granular cytoplasmic structure and to express $CD8\alpha$ molecules with no relation to presence or absence of a thymus, TCR expression, or intestinal flora.

It is well known that antigen-specific $\alpha\beta$ -IELs have a protective role against infection by pathogenic microorganisms, but ligands of cytotoxicity expressed by $\alpha\beta$ -IELs and $\gamma\delta$ -IELs from specific pathogen-free mice and $\gamma\delta$ -IELs from germ-free mice are still not clear. Do these cytotoxic IELs present in the harsh microenvironments at the front line of intestinal mucosa correctly identify target cells by TCR? Even without very strict specificity, are IECs that have been damaged by stress, viral

infections, bacterial infections, or transformed IECs rapidly detected and are these dubious IECs eliminated? Are these cytotoxic IELs indiscriminate predators lurking savagely in the jungle of the IEC layer (Fig. 8)? What T cells are these IELs anyway? In any event, elucidation of these conditions is very important to clarify the development and physiological function of IELs.

Conclusion

IELs are known as peripheral T cells with marked specificity and a very large population size, but their development and physiological function remain a mystery. Findings obtained through research over the past 30 years are very important. We view these diverse experimental observations as many tips of an iceberg that must firmly interconnect with its hidden part. Understanding the immunobiology of IELs must begin in the context of intestinal flora, which outnumber the total number of cells in our body and which have evolved with us in a commensal or symbiotic state. Elucidation of the characteristics of immune responses in the intestines, as the frontline defense against pathogens from the outer world, through clarification of the development and physiological function of IELs is essential for manipulation of intestinal immunity for our benefit.

References

- Guy-Grand D, Griscelli C, Vassalli P. The gut-associated lymphoid system: nature and properties of the large dividing cells. *Eur J Immunol* 1974;4:435-443.
- Tonegawa S, et al. Diversity, development, ligands, and probable function of $\gamma\delta$ T cells. *Cold Spring Harb Symp Quant Biol* 1989;54:31-44.
- Cerf-Bensussan N, Guy-Grand D, Griscelli C. Intraepithelial lymphocytes of human gut: isolation, characterization and study of natural killer activity. *Gut* 1985;26:81-88.
- Klein JR. Ontogeny of the $Thy-1^-$, $Lyt-2^+$ murine intestinal intraepithelial lymphocyte. Characterization of unique population of thymus-independent cytotoxic effector cells in the intestinal mucosa. *J Exp Med* 1986;164:309-314.
- Dillon SB, MacDonald TT. Functional characterization of Con A-responsive mouse small intestinal intraepithelial lymphocytes. *Immunology* 1986;59:389-396.
- Goodman T, Lefrancois L. Expression of the $\gamma\delta$ T-cell receptor on intestinal $CD8^+$ intraepithelial lymphocytes. *Nature* 1988;333:855-858.
- Klein JR. Thymus-independent development of gut T cells. *Chem Immunol* 1998;71:88-102.
- Lefrancois L, Puddington L. Basic aspects of intraepithelial lymphocyte immunobiology. In: Ogra PL, Mestecky J, Lamm ME, Strober W, Bienenstock L, McGhee JR, eds. *Mucosal Immunology*. San Diego: Academic Press, 1999:413-428.
- Kronenberg M, Cheroutre H. Development, function, and specificity of intestinal intraepithelial lymphocytes. In: Mestecky J, Lamm ME, Strober W, Bienenstock J, McGhee JR, Mayer L, eds. *Mucosal Immunology*. San Diego: Academic Press, 2005:565-581.
- Hayday A, Theodoridis E, Ramsburg E, Shires J. Intraepithelial lymphocytes: exploring the third way in immunology. *Nat Immunol* 2001;2:997-1003.
- Bandeira A, et al. Extrathymic origin of intestinal intraepithelial lymphocytes bearing T-cell antigen receptor $\gamma\delta$. *Proc Natl Acad Sci USA* 1991;88:43-47.
- Nonaka S, et al. Intestinal $\gamma\delta$ T cells develop in mice lacking thymus, all lymph nodes, Peyer's patches, and isolated lymphoid follicles. *J Immunol* 2005;174:1906-1912.
- Fichtelius KE. The gut epithelium - a first level lymphoid organ? *Exp Cell Res* 1968;49:87-104.
- Matsunaga T. Did the first adaptive immunity evolve in the gut of ancient jawed fish? *Cytogenet Cell Genet* 1998;80:138-141.
- Du Pasquier L, Flajnik M. Origin and evolution of the vertebrate immune system. In: Paul W, ed. *Fundamental Immunology*. Philadelphia: Lippincott-Raven Publisher, 1999:605-650.
- Griebel PJ, Hein WR. Expanding the role of Peyer's patches in B-cell ontogeny. *Immunol Today* 1996;17:30-39.

17. Davies M, Parrott D. The early appearance of specific cytotoxic T cells in murine gut mucosa. *Clin Exp Immunol* 1980;**42**:273–279.
18. Ernst PB, Clark DA, Rosenthal KL, Befus AD, Bienenstock J. Detection and characterization of cytotoxic T lymphocyte precursors in the murine intestinal intraepithelial leukocyte populations. *J Immunol* 1986;**136**:2121–2126.
19. Viney J, Kilshaw P, MacDonald T. Cytotoxic α/β^+ and γ/δ^+ T cells in murine intestinal epithelium. *Eur J Immunol* 1990;**20**:1623–1626.
20. Correa I, Bix M, Liao N, Zijlstra M, Jaenisch R, Raulat D. Most $\gamma\delta$ T cells develop normally in β_2 -microglobulin-deficient mice. *Proc Natl Acad Sci USA* 1992;**89**:653–657.
21. Neuhaus O, Emoto M, Blum C, Yamamoto S, Kaufmann S. Control of thymus-independent intestinal intraepithelial lymphocytes by β_2 -microglobulin. *Eur J Immunol* 1995;**25**:2332–2339.
22. Pennington DJ, et al. The inter-relatedness and interdependence of mouse T cell receptor $\gamma\delta^+$ and $\alpha\beta^+$ cells. *Nat Immunol* 2003;**4**:991–998.
23. Fujiura Y, et al. Development of $CD8\alpha^+$ intestinal intraepithelial T cells in β_2 -microglobulin- and/or TAP1-deficient mice. *J Immunol* 1996;**156**:2710–2715.
24. Leishman AL, et al. T cell responses modulated through interaction between $CD8\alpha$ and the nonclassical MHC molecules, TL. *Science* 2001;**294**:1936–1939.
25. Leishman AJ, et al. Precursor of functional MHC class I- or class II-restricted $CD8\alpha^+$ T cells are positively selected in the thymus by agonist self-peptides. *Immunity* 2002;**16**:355–364.
26. Lefrancois L, Puddington L. Extrathymic intestinal T-cell development: virtual reality? *Immunol Today* 1995;**16**:16–21.
27. Rocha B, Guy-Grand D, Vassalli P. Extrathymic T cell differentiation. *Curr Opin Immunol* 1995;**7**:235–242.
28. Poussier P, Julius M. Speculation on the lineage relationship among $CD4^+8^+$ gut-derived T cells and their role(s). *Semin Immunol* 1999;**11**:293–303.
29. Guy-Grand D, Vassalli P. Gut intraepithelial lymphocyte development. *Curr Opin Immunol* 2002;**14**:255–259.
30. Ishikawa H, Kanamori Y, Hamada H, Kiyono H. Development and function of organized gut-associated lymphoid tissues. In: Mestecky J, Lamm ME, Strober W, Bienenstock J, McGhee JR, Mayer L, eds. *Mucosal Immunology*. San Diego: Academic Press, 2005:285–405.
31. Kuwata N, Igarashi H, Ohmura T, Aizawa S, Sakaguchi N. Absence of expression of RAG1 in peritoneal B-1 cells detected by knocking into RAG1 locus with green fluorescent protein gene. *J Immunol* 1999;**163**:6355–6359.
32. Fujimoto M, et al. Complementary role for CD19 and Bruton's tyrosine kinase in B lymphocyte signal transduction. *J Immunol* 2002;**168**:5465–5476.
33. Ota Y, et al. Extrathymic origin of $V\gamma 1/V\delta 6$ T cells in the skin. *Eur J Immunol* 1992;**22**:595–598.
34. Matis LA, Cron R, Bluestone JA. Major histocompatibility complex-linked specificity of $\gamma\delta$ receptor-bearing T lymphocytes. *Nature* 1987;**330**:262–264.
35. Yoshikai Y, et al. Functional T cell receptor δ chain gene messages in athymic nude mice. *Eur J Immunol* 1988;**18**:1039–1043.
36. Kanamori Y, et al. Identification of novel lymphoid tissues in murine intestinal mucosa where clusters of c-kit⁺IL-7R⁺Thy1⁺ lympho-hemopoietic progenitors develop. *J Exp Med* 1996;**184**:1449–1459.
37. Hamada H, et al. Identification of multiple isolated lymphoid follicles on the antimesenteric wall of the mouse small intestine. *J Immunol* 2002;**168**:57–64.
38. Suzuki K, et al. Gut cryptopatches: direct evidence of extrathymic anatomical sites for intestinal T lymphopoiesis. *Immunity* 2000;**13**:691–702.
39. Saito H, et al. Generation of intestinal T cells from progenitors residing in gut cryptopatches. *Science* 1998;**280**:275–278.
40. Oida T, et al. Role of gut cryptopatches in early extrathymic maturation of intestinal intraepithelial T cells. *J Immunol* 2000;**164**:3616–3626.
41. Podd BS, et al. T cells in cryptopatch aggregates share TCR γ variable region junctional sequences with $\gamma\delta$ T cells in the small intestinal epithelium of mice. *J Immunol* 2006;**176**:6532–6542.
42. Lambolez F, et al. Characterization of T cell differentiation in the murine gut. *J Exp Med* 2002;**195**:437–449.
43. Guy-Grand D, et al. Extrathymic T cell lymphopoiesis: ontogeny and contribution to gut intraepithelial lymphocytes in athymic and euthymic mice. *J Exp Med* 2003;**197**:333–341.
44. Lin T, Matsuzaki G, Kenai H, Nomoto K. Progenies of fetal thymocytes are the major source of $CD4^+CD8\alpha^+$ intestinal intraepithelial lymphocytes early in ontogeny. *Eur J Immunol* 1994;**24**:1785–1791.
45. Wang J, Klein JR. Thymus-neuroendocrine interactions in extrathymic T cell development. *Science* 1994;**265**:1860–1862.
46. Wang J, Whetsell M, Klein JR. Local hormone networks and intestinal T cell homeostasis. *Science* 1997;**275**:1937–1939.
47. Dejbakhsh-Jones S, Jerabek L, Weissman IL, Strober S. Extrathymic maturation of $\alpha\beta$ T cells from hemopoietic stem cells. *J Exp Med* 1999;**155**:3338–3344.
48. Simizu T, et al. The majority of lymphocytes in the bone marrow, thymus and extrathymic T cells in the liver are generated in situ from their own preexisting precursors. *Microbiol Immunol* 1999;**43**:595–608.
49. Arcangeli M-L, et al. Extrathymic hemopoietic progenitors committed to T cell differentiation in the adult mouse. *J Immunol* 2005;**174**:1980–1988.
50. Eberl G, Littman DR. Thymic origin of intestinal $\alpha\beta$ T cells revealed by fate mapping of ROR γ ^t cells. *Science* 2004;**305**:248–251.
51. Maki K, Sunaga S, Ikuta K. The V-J recombination of T cell receptor- γ genes is blocked in interleukin-7 receptor-deficient mice. *J Exp Med* 1996;**184**:2423–2427.
52. Taylor RT, Lugerling A, Newell AN, Williams IR. Intestinal cryptopatch formation in mice requires lymphotoxin α and the lymphotoxin β receptor. *J Immunol* 2004;**173**:7183–7189.
53. Deusch K, Luling F, Reich K, Classen M, Wagner H, Pfeffer K. A major fraction of human intraepithelial lymphocytes simultaneously expresses the γ/δ T cell receptor, the CD8 accessory molecule and preferentially uses the $V\delta 1$ gene segment. *Eur J Immunol* 1991;**21**:1053–1059.
54. Lundqvist C, Baranov V, Hammarstrom S, Athlin L, Hammarstrom ML. Intra-epithelial lymphocytes. Evidence for regional specialization and extrathymic T cell maturation in the human gut epithelium. *Int Immunol* 1995;**7**:1473–1487.
55. Jarry A, Cerf-Bensussan N, Brousse N, Selz F, Guy-Grand D. Subsets of $CD3^+$ (T cell receptor α/β or γ/δ) and $CD3^-$ lymphocytes isolated from normal human gut epithelium display phenotypical features different from their counterparts in peripheral blood. *Eur J Immunol* 1990;**20**:1097–1103.
56. Lathe M, Terry L, MacDonald TT. High frequency of $CD8\alpha$ homodimer-bearing T cells in human fetal intestine. *Eur J Immunol* 1994;**24**:1703–1705.
57. Lynch S, Kelleher D, McManus R, O'Farrelly C. RAG1 and RAG2 expression in human intestinal epithelium: evidence of extrathymic T cell differentiation. *Eur J Immunol* 1995;**25**:1143–1147.
58. Koningsberger JC, et al. TCR expression in human fetal intestine and identification of an early T cell receptor β -chain transcript. *J Immunol* 1997;**159**:1775–1782.
59. Howie D, et al. Extrathymic T cell differentiation in the human intestine early in life. *J Immunol* 1998;**161**:5862–5872.
60. Ramanathan S, Marandi L, Poussier P. Evidence for the extrathymic origin of intestinal TCR $\gamma\delta^+$ T cells in normal rats and for an impairment of this differentiation pathway in BB rats. *J Immunol* 2002;**168**:2182–2187.

61. Hitotsumatsu O, et al. Identification and characterization of novel gut-associated lymphoid tissues in rat small intestine. *J Gastroenterol* 2005;**40**:956–963.
62. Mayrhofer G, Moghaddami M, Murphy C. Lymphocyte-filled villi (LFV): non-classical organized lymphoid tissues in the mucosa of the small intestine. *Mucosal Immunol Update* 1999;**7**:9–13.
63. Jemeson J, et al. A role for skin $\gamma\delta$ T cells in wound repair. *Science* 2002;**296**:747–749.
64. Jemeson JM, Cauvi G, Deborah A, Witherden DA, Havran WL. A keratinocyte-responsive $\gamma\delta$ TCR is necessary for dendritic epidermal T cell activation by damaged keratinocytes and maintenance in the epidermis. *J Immunol* 2004;**172**:3573–3579.
65. Ferguson A. Intraepithelial lymphocytes of the small intestine. *Gut* 1977;**18**:921–937.
66. Mysorekar IU, Lorenz RG, Gordon JI. A gnotobiotic transgenic mouse model for studying interactions between small intestinal enterocytes and intraepithelial lymphocytes. *J Biol Chem* 2002;**277**:37811–37819.
67. Rocha B, Vassalli P, Guy-Grand D. The V β repertoire of mouse gut homodimeric α CD8⁺ intraepithelial T cell receptor α/β ⁺ lymphocytes reveals a major extrathymic pathway of T cell differentiation. *J Exp Med* 1991;**173**:483–486.
68. Page ST, et al. Intestinal intraepithelial lymphocytes include precursors committed to the T cell receptor $\alpha\beta$ lineage. *Proc Natl Acad Sci USA* 1998;**95**:9459–9464.
69. Poussier P, Edouard P, Lee C, Binnie M, Julius M. Thymus-independent development and negative selection of T cells expressing T cell receptor α/β in the intestinal epithelium: evidence for distinct circulation patterns of gut- and thymus-derived T lymphocytes. *J Exp Med* 1992;**176**:187–199.
70. Suzuki S, et al. Low level of mixing of partner cells seen in extrathymic T cells in the liver and intestine of parabiotic mice: its biological implication. *Eur J Immunol* 1998;**28**:3719–3729.
71. Stappenbeck TS, Wong MH, Saam JR, Mysorekar IU, Gordon JI. Notes from some crypt watchers: regulation of renewal in the mouse intestinal epithelium. *Curr Opin Cell Biol* 1998;**10**:702–709.
72. Penney L, Kilshaw PJ, MacDonald TT. Regional variation in the proliferative rate and lifespan of $\alpha\beta$ TCR⁺ and $\gamma\delta$ TCR⁺ intraepithelial lymphocytes in the murine small intestine. *Immunology* 1995;**86**:212–218.
73. Balk SP, et al. Oligoclonal expansion and CD1 recognition by human intestinal intraepithelial lymphocytes. *Science* 1991;**253**:1411–1415.
74. Van Kerckhove, et al. Oligoclonality of human intestinal intraepithelial T cells. *J Exp Med* 1992;**175**:57–63.
75. Blumberg RS, Yockey CE, Grass GG, Ebert EC, Balk SP. Human intestinal intraepithelial lymphocytes are derived from a limited number of T cell clones that utilize multiple V β T cell receptor genes. *J Immunol* 1993;**150**:5144–5153.
76. Regnault A, Cumano A, Vassalli P, Guy-Grand D, Kourilsky P. Oligoclonal repertoire of the CD8 $\alpha\alpha$ and the CD8 $\alpha\beta$ TCR- α/β murine intestinal intraepithelial T lymphocytes: evidence for the random emergence of T cells. *J Exp Med* 1994;**180**:1345–1358.
77. Regnault A, et al. The expansion and selection of T cell receptor $\alpha\beta$ intestinal intraepithelial clones. *Eur J Immunol* 1996;**26**:914–921.
78. Guy-Grand D, Malassis-Seris M, Briottet C, Vassalli P. Cytotoxic differentiation of mouse gut thymodependent and independent intraepithelial T lymphocytes is induced locally. Correlation between functional assays, presence of perforin and granzyme transcripts, and cytoplasmic granules. *J Exp Med* 1991;**173**:1549–1552.
79. Goodman T, Lefrancois L. Intraepithelial lymphocytes. Anatomical site, not T cell receptor form, dictates phenotype and function. *J Exp Med* 1989;**170**:1569–1581.
80. Ishikawa H, Li Y, Abeliovich A, Yamamoto S, Kaufmann HE, Tonegawa S. Cytotoxic and interferon γ -producing activities of $\gamma\delta$ T cells in the mouse intestinal epithelium are strain dependent. *Proc Natl Acad Sci USA* 1993;**90**:8204–8208.
81. Kawaguchi M, et al. Cytolytic activity of intestinal intraepithelial lymphocytes in germ-free mice is strain dependent and determined by T cells expressing $\gamma\delta$ T-cell antigen receptors. *Proc Natl Acad Sci USA* 1993;**90**:8591–8594.
82. Croitoru K, et al. Presence of intestinal lymphocytes in mice with severe combined immunodeficiency disease. *Eur J Immunol* 1990;**20**:645–651.
83. Takahashi-Iwanaga H, Iwanaga T, Sakamoto Y, Fujita T. Ultrastructural and time-lapse observation of intraepithelial lymphocytes in the small intestine of the guinea pig: their possible role in the removal of effete enterocytes. *Cell Tissue Res* 1995;**280**:491–497.

Lamina Propria c-kit⁺ Immune Precursors Reside in Human Adult Intestine and Differentiate Into Natural Killer Cells

HIROSHI CHINEN,^{*,‡} KATSUYOSHI MATSUOKA,^{*} TOSHIRO SATO,^{*} NOBUHIKO KAMADA,^{*} SUSUMU OKAMOTO,^{*} TADAKAZU HISAMATSU,^{*} TAKU KOBAYASHI,^{*} HIROTOSHI HASEGAWA,[§] AKIRA SUGITA,[¶] FUKUNORI KINJO,[†] JIRO FUJITA,[‡] and TOSHIFUMI HIBI^{*}

^{*}Division of Gastroenterology and Hepatology, Department of Internal Medicine, School of Medicine, Keio University, Tokyo, Japan; [‡]Department of Medicine and Therapeutics, Control and Prevention of Infectious Diseases, Graduate School of Medicine, University of the Ryukyus, Okinawa, Japan; [§]Department of Surgery, School of Medicine, Keio University, Tokyo, Japan; and [¶]Department of Surgery, Yokohama Municipal Citizen's Hospital, Yokohama, Japan

Background & Aims: Recent studies have revealed that murine intestinal mucosa contains several kinds of lineage markers (lin)⁻ c-kit⁺ immune precursor cells. However, immune precursors in the human adult intestine have not been studied extensively. **Methods:** Lamina propria mononuclear cells and intraepithelial lymphocytes from surgically resected human adult intestine were examined for the surface antigen expression and cytokine profile by immunohistochemistry and flow cytometry. The transcriptional profile of these cells was analyzed by reverse-transcription polymerase chain reaction. The phenotypic and functional characterization of the *in vitro* differentiating cells from the precursors was examined by flow cytometry. **Results:** We identified lin⁻ c-kit⁺ cells scattered throughout lamina propria of the human adult intestine. These intestinal immune precursors expressed CD34, CD38, CD33, interleukin-2R α , and interleukin-7R α , and they had much more abundant expression of Id2, PU.1, SpiB1, and lymphotoxin than thymocytes. The lin⁻ c-kit⁺ immune precursors mainly differentiated into CD56⁺ c-kit^{dim} cells during *in vitro* culture. These *in vitro* differentiating cells corresponded to intestinal natural killer (NK) cells, which had distinct characteristics from their peripheral counterparts, such as CD83 and integrin α_E expression, less cytotoxic activity, and higher interferon- γ production. Furthermore, both c-kit^{dim} cells and NK cells were increased in lamina propria of Crohn's disease, although there was no change for peripheral blood NK cells. **Conclusions:** The human intestine may have the unique NK cell differentiation system, which may contribute to maintenance of immune homeostasis in the intestine.

The cellular components of the immune system, such as T cells, B cells, monocytes, granulocytes, macrophages, dendritic cells, and natural killer (NK) cells, are derived from common hematopoietic stem cells (HSCs) in the bone marrow. As a first step, HSCs differentiate into 2 distinct subsets: common myeloid progenitors and common lymphoid progenitors. Although common my-

eloid progenitors ultimately differentiate into myeloid cells such as monocytes, granulocytes, macrophages, and dendritic cells,¹ common lymphoid progenitors differentiate into B-cell precursors and common T- and NK-cell precursors (T/NKPs).² T/NKPs subsequently differentiate into NKPs and T-cell precursors.³⁻⁵ These steps are assumed to proceed mainly in the bone marrow, which is regarded as the most important site for primary immune cell differentiation.

A unique immune system has developed in the intestine. The intestinal immune system includes Peyer's patches, isolated lymphoid follicles, mesenteric lymph nodes (MLN), lamina propria mononuclear cells (LPMCs), and intraepithelial lymphocytes (IELs). This intestinal immune system maintains immunologic homeostasis against gut luminal antigens. In addition to these components, the intestine has become recognized as a site for differentiation of immune cells. Recent studies have revealed that murine intestinal mucosa contains immune precursor cells, which are lymphoid tissue inducer cells (LTi)^{6,7} in the fetus and cryptopatch (CP) cells⁸ in the adult. Both LTi and CP cells express c-kit, IL-7 receptor α subunit (IL-7R α), IL-2R α , CD44, and CD4^{dim}. These surface phenotypes are similar to those of common lymphoid progenitors in bone marrow, and LTi and CP cells have been reported to develop *in situ* into Peyer's patches^{6,7,9} and extrathymic T cells of IELs,^{10,11} respectively. In addition, a recent study suggested that CP cells can function as adult LTi by developing into isolated lymphoid folli-

Abbreviations used in this paper: CP, cryptopatch; HSC, common hematopoietic stem cell; IENK, intraepithelial natural killer cell; IFN, interferon; IL, interleukin; IL-7R α , IL-7 receptor α subunit; Lin, lineage markers; LPMCs, lamina propria mononuclear cells; LPNKs, lamina propria natural killer cells; LTi, lymphoid tissue inducer cells; MLN, mesenteric lymph nodes; NK, natural killer; PBL, peripheral blood lymphocytes; PBNKs, peripheral blood natural killer cells; PCR, polymerase chain reaction; pT α , pre-T cell receptor chain α ; RAG, recombination activating gene; SEM, standard error of the mean; T/NKPs, common T and natural killer cell precursors; TNF, tumor necrosis factor.

© 2007 by the AGA Institute
0016-5085/07/\$32.00
doi:10.1053/j.gastro.2007.05.017