

研究成果の刊行に関する一覧表

1. Haruhiko Tokuda, Shinji Takai, Rie Matsushima-Nishiwaki, Shigeru Akamatsu, Yoshiteru Hanai, Takayuki Hosoi, Atsushi Harada, Toshiki Ohta, Osamu Kozawa. (-)-Epigallocatechin gallate enhances prostaglandin F_{2α}-induced VEGF synthesis via upregulating SAPK/JNK activation in osteoblasts. *J Cell Biochem* 100: 1146-1153, 2007.
2. Haruhiko Tokuda, Shinji Takai, Rie Matsushima-Nishiwaki, Shigeru Akamatsu, Yoshiteru Hanai, Takayuki Hosoi, Atsushi Harada, Toshiki Ohta, Osamu Kozawa. (-)-Epigallocatechin gallate suppresses endothelin-1-induced interleukin-6 synthesis in osteoblasts: Inhibition of p44/p42 MAP kinase activation. *FEBS Letters* 581: 1311-1316, 2007.
3. Haruhiko Tokuda, Yoshiteru Hanai, Rie Matsushima-Nishiwaki, Junichi Yamauchi, Tomoaki Doi, Atsushi Harada, Shinji Takai, Osamu Kozawa. Rho-kinase regulates endothelin-1-stimulated IL-6 synthesis via p38 MAP kinase in osteoblasts. *Biochemical and Biophysical Research Communications*. 362: 799-804, 2007.
4. Kazumasa Hayasaka, Takashi Nihashi, Toshihiro Matsuura, Tetsuya Yagi, Kazumitsu Nakashima, Yasuji Kawabata, Kengo Ito, Takashi Katoh, Keita Sakata, Atsushi Harada. Metastasis of the gastrointestinal tract: FDG-PET imaging. *Ann Nucl Med* 21: 361-365, 2007.
5. Tokuda H, Takai S, Hanai Y, Harada A, Matsushima-Nishiwaki R, Akamatsu S, Ohta T, Kozawa O. Platelet-derived growth factor-BB amplifies PGF_{2α}-stimulated VEGF synthesis in osteoblasts: function of phosphatidylinositol 3-kinase. *Prostaglandins Leukot Essent Fatty Acids*. 77:187-193, 2007.
6. Takai S, Tokuda H, Hanai Y, Harada A, Yasuda E, Matsushima-Nishiwaki R, Kato H, Ogura S, Ohta T, Kozawa O. Negative regulation by p70 S6 kinase of FGF-2-stimulated VEGF release through stress-activated protein kinase/c-Jun N-terminal kinase in osteoblasts. *J Bone Miner Res*. 22:337-346, 2007.

7. 原田敦. 運動器不安定症と今後の展開. 整形・災害外科 27-35, 2007.
8. 原田敦, 松井康素, 奥泉宏康, 竹村真里枝, 若尾典充, 長屋政博, 水野雅士. 転倒・骨折予防の立場からみた骨強度の評価. Osteoporosis Jpn 15: 152-154, 2007.
9. 竹村真里枝, 松井康素, 原田敦, 安藤富士子, 下方浩史. 地域在住中高年者の骨代謝マーカーによる骨量減少/骨粗鬆症予測. Osteoporosis Jpn 15: 28-32, 2007.
10. 加藤智香子, 猪田邦雄, 原田敦, 長屋政博, 徳田治彦. 施設入所高齢者の転倒恐怖とQOL、ADL、身体活動量との関連. Osteoporosis Japan 317-319, 2007.
11. Takai S, Tokuda H, Hanai Y, Kozawa O. Limitation by p70 S6 kinase of PDGF-BB-induced IL-6 synthesis in osteoblast-like MC3T3-E1 cells. Metabolism 56: 476-483, 2007.
12. Takai S, Tokuda H, Hanai Y, Kozawa O. Activation of phosphatidylinositol 3-kinase/Akt limits FGF-2-induced VEGF release in osteoblasts. Mol Cell Endocrinol. 267:46-54, 2007.
13. Yamauchi J, Takai S, Matsushima, Nishiwaki R, Hanai Y, Doi T, Kato H, Ogura S, Kato K, Tokuda H, Kozawa O. (-)-Epigallocatechin gallate inhibits prostaglandin D2-stimulated HSP27 induction via suppression of the p44/p42 MAP kinase pathway in osteoblasts. Prostaglandins Leukot Essent Fatty Acids. 77: 173-179, 2007.
14. 中澤信, 長屋政博. 疾患特有の評価法 パーキンソン病・総合リハビリテーション 35(1): 55-60, 2007.
15. 長屋政博. 転倒予防教室の意義と今後の展望. Calcitonin@Osteoporosis No.7: 1-3, 2007.
16. 奥泉宏康. 転倒による骨折のバイオメカニクス. 身体教育医学研究 8: 75-77, 2007.
17. 奥泉宏康. ヒッププロテクターの現状と課題. GERONTOLOGY NEW HORIZON 19(3): 185-189, 2007.
18. 奥泉宏康, 武藤芳照. 人はなぜ転ぶのか? 落ちるのか? Nursing Today 22(12): 28-33,

2007.

19. Chikako Kato, Kunio Ida, Morio Kawamura, Masahiro Nagaya, Haruhiko Tokuda, Akiko Tamakoshi, Atsushi Harada. Relation of falls efficacy scale (FES) to quality of life among nursing home female residents with comparatively intact cognitive function in Japan. Nagoya J. Med. Sci. 70. 19-27, 2008 (in press)
20. 内藤善規、加藤智香子. 地域在住高齢者の骨密度に関連する要因の検討. 名古屋大学医学部保健学科卒業論文集. 1-6, 2008.
21. 川村皓生、加藤智香子. 地域在住高齢者と介護施設高齢者の健康関連 QOL に関連する因子の検討. 名古屋大学医学部保健学科卒業論文集. 1-9, 2008.

研究成果の刊行物・別刷

(–)-Epigallocatechin Gallate Enhances Prostaglandin F_{2α}-Induced VEGF Synthesis Via Upregulating SAPK/JNK Activation in Osteoblasts

Haruhiko Tokuda,^{1,2*} Shinji Takai,² Rie Matsushima-Nishiwaki,² Shigeru Akamatsu,³ Yoshiteru Hanai,^{1,2} Takayuki Hosoi,⁴ Atsushi Harada,⁵ Toshiki Ohta,⁶ and Osamu Kozawa²

¹Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

²Department of Pharmacology, Gifu University Graduate School of Medicine, Gifu 501-1194, Japan

³Critical Care Center, Matsunami General Hospital, Kasamatsu 501-6062, Japan

⁴Department of Advanced Medicine, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

⁵Department of Restorative Medicine, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

⁶Department of Internal Medicine, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

Abstract Catechin, one of the major flavonoids presented in plants such as tea, reportedly suppresses bone resorption. We previously reported that prostaglandin F_{2α} (PGF_{2α}) stimulates the synthesis of vascular endothelial growth factor (VEGF) via p44/p42 mitogen-activated protein (MAP) kinase in osteoblast-like MC3T3-E1 cells. To clarify the mechanism of catechin effect on osteoblasts, we investigated the effect of (–)-epigallocatechin gallate (EGCG), one of the major green tea flavonoids, on the VEGF synthesis by PGF_{2α} in MC3T3-E1 cells. The PGF_{2α}-induced VEGF synthesis was significantly enhanced by EGCG. The amplifying effect of EGCG was dose dependent between 10 and 100 μM. EGCG did not affect the PGF_{2α}-induced phosphorylation of p44/p42 MAP kinase. SB203580, a specific inhibitor of p38 MAP kinase, and SP600125, a specific inhibitor of stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK), reduced the PGF_{2α}-induced VEGF synthesis. EGCG markedly enhanced the phosphorylation of SAPK/JNK induced by PGF_{2α} without affecting the PGF_{2α}-induced phosphorylation of p38 MAP kinase. SP600125 markedly reduced the amplification by EGCG of the SAPK/JNK phosphorylation. In addition, the PGF_{2α}-induced phosphorylation of c-Jun was amplified by EGCG. These results strongly suggest that EGCG upregulate PGF_{2α}-stimulated VEGF synthesis resulting from amplifying activation of SAPK/JNK in osteoblasts. *J. Cell. Biochem.* 100: 1146–1153, 2007. © 2006 Wiley-Liss, Inc.

Key words: catechin; PGF_{2α}; VEGF; MAP kinase; osteoblast

Grant sponsor: The Ministry of Education, Science, Sports and Culture of Japan, Grant-in-Aid for Scientific Research; Grant numbers: 16590873, 16591482; Grant sponsor: The Ministry of Health, Labour and Welfare of Japan, The Research Grants for Longevity Sciences, Research on Proteomics and Research on Fracture and Dementia; Grant numbers: 15A-1, 15C-2, 17A-3.

*Correspondence to: Haruhiko Tokuda, Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu, Aichi 474-8511, Japan. E-mail: tokuda@ncgg.go.jp

Received 19 July 2006; Accepted 20 July 2006

DOI 10.1002/jcb.21104

Published online 9 October 2006 in Wiley InterScience (www.interscience.wiley.com).

© 2006 Wiley-Liss, Inc.

It is generally recognized that compounds in foods such as vegetables and fruits have beneficial properties to human being. Among them, flavonoids reportedly show antioxidative, antibacterial, and antitumor effects [Jankun et al., 1997; Harbourne and Williams, 2000]. Catechins are one of the major flavonoids, which are present in various species of plants such as tea [Harbourne and Williams, 2000]. In bone metabolism, it has been reported that catechin suppresses bone resorption [Delaisse et al., 1986]. Bone metabolism is regulated by two functional cells, osteoblasts, and osteoclasts, responsible for bone formation and bone resorption, respectively [Nijweide et al., 1986]. The

formation of bone structures and bone remodeling results from the coupling process, bone resorption by activated osteoblasts with subsequent deposition of new matrix by osteoblasts. As for osteoblasts, it has been shown that catechin stimulates alkaline phosphatase activity, a mature osteoblast phenotype and reduces bone-resorptive cytokine production in osteoblast-like MC3T3-E1 cells [Choi and Hwang, 2003]. However, the exact role of catechin in osteoblasts has not yet been clarified.

Vascular endothelial growth factor (VEGF) has been characterized as a heparin-binding angiogenic growth factor displaying high specificity for vascular endothelial cells [Ferrara and Davis-Smyth, 1997]. It is well recognized that VEGF, which is produced and secreted from a variety of cell types, increases capillary permeability and stimulates proliferation of endothelial cells [Ferrara and Davis-Smyth, 1997]. As for bone metabolism, it has been shown that inactivation of VEGF causes complete suppression of blood vessel invasion concomitant with impaired trabecular bone formation and expansion of hypertrophic chondrocyte zone in mouse tibial epiphyseal growth plate [Gerber et al., 1999]. Accumulating evidence indicates that osteoblasts among bone cells produce and secrete VEGF in response to various humoral factors [Goad et al., 1996; Wang et al., 1996; Ferrara and Davis-Smyth, 1997; Schalaepi et al., 1997]. During bone remodeling, capillary endothelial cells provide the microvasculature, and osteoblasts and osteoprogenitor cells, which locally proliferate and differentiate into osteoblasts, migrate into the resorption lacuna. It is currently recognized that the activities of osteoblasts, osteoclasts and capillary endothelial cells are closely coordinated via humoral factors as well as by direct cell-to-cell contact, and these cells cooperatively regulate bone metabolism [Erlebacher et al., 1995]. Thus, there is no doubt that VEGF secreted from osteoblasts plays an important role in the regulation of bone metabolism. However, the mechanism underlying VEGF synthesis in osteoblasts has not yet been fully clarified.

It is well known that prostaglandins (PGs) act as autocrine/paracrine modulators in osteoblasts and play crucial roles in the regulation of bone metabolism [Nijweide et al., 1986; Pilbeam et al., 1996]. Among them, $\text{PGF}_{2\alpha}$ is known as a potent bone-resorptive agent and stimulates the proliferation of osteoblasts and

inhibits their differentiation [Pilbeam et al., 1996]. In our previous studies [Miwa et al., 1990; Kozawa et al., 1994], we have reported that $\text{PGF}_{2\alpha}$ stimulates both phosphoinositide-hydrolyzing phospholipase C (PI-PLC) and phosphatidylcholine-hydrolyzing phospholipase D (PC-PLD), recognized to be two major pathways of physiological protein kinase C (PKC) activation [Nishizuka, 1992; Exton, 1999], in osteoblast-like MC3T3-E1 cells. In addition, we have recently shown that $\text{PGF}_{2\alpha}$ stimulates the VEGF synthesis through PKC-dependent activation of p44/p42 mitogen-activated protein (MAP) kinase in these cells [Tokuda et al., 2003]. In the present study, we investigated the effect of (-)-epigallocatechin gallate (EGCG), one of the major green tea flavonoids, on the $\text{PGF}_{2\alpha}$ -induced VEGF synthesis in osteoblast-like MC3T3-E1 cells. We here show that EGCG upregulates $\text{PGF}_{2\alpha}$ -stimulated VEGF synthesis via enhancing SAPK/JNK activation among the MAP kinase superfamily in these cells.

MATERIALS AND METHODS

Materials

$\text{PGF}_{2\alpha}$ was purchased from Sigma Chemical Co. (St. Louis, MO). Mouse VEGF enzyme immunoassay kit was purchased from R&D Systems, Inc. (Minneapolis, MN). (-)-Epigallocatechin gallate (EGCG), SB203580, and SP600125 were obtained from Calbiochem-Novabiochem Co. (La Jolla, CA). Phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies, p38 MAP kinase antibodies, phospho-specific SAPK/JNK antibodies, SAPK/JNK antibodies, and c-Jun antibodies were purchased from Cell Signaling Technology (Beverly, MA). ECL Western blotting detection system was purchased from Amersham Japan (Tokyo, Japan). Other materials and chemicals were obtained from commercial sources. $\text{PGF}_{2\alpha}$ was dissolved in ethanol. SB203580 and SP600125 were dissolved in dimethyl sulfoxide. The maximum concentration of ethanol or dimethyl sulfoxide was 0.1%, which did not affect the assay for VEGF or the analysis of Western blot.

Cell Culture

Cloned osteoblast-like MC3T3-E1 cells derived from newborn mouse calvaria [Sudo

et al., 1983] were maintained as previously described [Kozawa et al., 1992]. Briefly, the cells were cultured in α -minimum essential medium (α -MEM) containing 10% fetal calf serum (FCS) at 37°C in a humidified atmosphere of 5% CO₂/95% air. The cells were seeded into 35-mm diameter dishes (5×10^4 /dish) or 90-mm diameter dishes (5×10^5 /dish) in α -MEM containing 10% FCS. After 5 days, the medium was exchanged for α -MEM containing 0.3% FCS. The cells were used for experiments after 48 h.

VEGF Assay

The cultured cells were stimulated by PGF_{2 α} in 1 ml of α -MEM containing 0.3% FCS for the indicated periods. When indicated, the cells were pretreated with EGCG, SB203580, or SP600125 for 60 min. The reaction was terminated by collecting the medium, and VEGF in the medium was measured by a VEGF enzyme immunoassay kit.

Western Blot Analysis

The cultured cells were stimulated by PGF_{2 α} in α -MEM containing 0.3% FCS for the indicated periods. The cells were washed twice with phosphate-buffered saline and then lysed, homogenized, and sonicated in a lysis buffer containing 62.5 mM Tris/HCl, pH 6.8, 2% sodium dodecyl sulfate (SDS), 50 mM dithiothreitol, and 10% glycerol. The cytosolic fraction was collected as a supernatant after centrifugation at 125,000g for 10 min at 4°C. SDS-polyacrylamide gel electrophoresis (PAGE) was performed by Laemmli [1970] in 10% polyacrylamide gel. Western blotting analysis was performed as described previously [Kato et al., 1996] by using phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies, p38 MAP kinase antibodies, phospho-specific SAPK/JNK antibodies, SAPK/JNK antibodies, or c-Jun antibodies, with peroxidase-labeled antibodies raised in goat against rabbit IgG being used as second antibodies. Peroxidase activity on the nitrocellulose sheet was visualized on X-ray film by means of the ECL Western blotting detection system. The pretreatment of EGCG or SP600125 was performed for 60 min before the addition of PGF_{2 α} .

Determination

The absorbance of enzyme immunoassay samples was measured at 450 nm with EL 340

Bio Kinetic Reader (Bio-Tek Instruments, Inc., Winooski, VT). The densitometric analysis was performed using Molecular Analyst/Macintosh (Bio-Rad Laboratories, Hercules, CA).

Statistical Analysis

The data were analyzed by ANOVA followed by the Bonferroni method for multiple comparisons between pairs, and a $P < 0.05$ was considered significant. All data are presented as the mean \pm SEM of triplicate determinations. Each experiment was repeated three times with similar results.

RESULTS

Effect of EGCG on the PGF_{2 α} -Stimulated VEGF Synthesis in MC3T3-E1 Cells

We have previously shown that PGF_{2 α} stimulates VEGF synthesis in osteoblast-like MC3T3-E1 cells [Tokuda et al., 2003]. We first examined the effect of EGCG on the PGF_{2 α} -stimulated VEGF synthesis. EGCG, which by itself had little effect on the VEGF levels, significantly enhanced the PGF_{2 α} -stimulated synthesis of VEGF (Table I). The amplifying effect of EGCG was dose dependent between 10 and 100 μ M (Table I).

Effect of EGCG on the Phosphorylation of p44/p42 MAP Kinase Induced by PGF_{2 α} in MC3T3-E1 Cells

We have previously reported that the PGF_{2 α} -stimulated VEGF synthesis is regulated by PGF_{2 α} -activated p44/p42 MAP kinase in MC3T3-E1 cells [Tokuda et al., 2003]. In order to investigate whether EGCG effect on the PGF_{2 α} -stimulated VEGF synthesis is mediated

TABLE I. Effect of EGCG on the PGF_{2 α} -Stimulated VEGF Synthesis in MC3T3-E1 Cells

EGCG (μ M)	PGF _{2α}	VEGF (pg/ml)
—	—	19 \pm 10
—	+	580 \pm 51
10	—	14 \pm 10
10	+	630 \pm 85
30	—	15 \pm 10
30	+	2680 \pm 167*
100	—	20 \pm 10
100	+	4100 \pm 249*

The cultured cells were pretreated with various doses of EGCG for 60 min, and then stimulated by 10 μ M PGF_{2 α} or vehicle for 24 h. Each value represents the mean \pm SEM of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

* $P < 0.05$, compared to the value of PGF_{2 α} alone.




Fig. 1. Effect of EGCG on the phosphorylation of p44/p42 MAP kinase induced by PGF_{2α} in MC3T3-E1 cells. The cultured cells were pretreated with the indicated doses of EGCG or vehicle for 60 min, and then stimulated by 10 μM PGF_{2α} or vehicle for 30 min. The extracts of cells were subjected to SDS-PAGE with subsequent Western blotting analysis with antibodies against phospho-specific p44/p42 MAP kinase or p44/p42 MAP kinase. The histogram shows quantitative representations of the levels of PGF_{2α}-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean ± SEM of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

through p44/p42 MAP kinase activation in MC3T3-E1 cells, we next examined the effect of EGCG on the PGF_{2α}-induced phosphorylation of p44/p42 MAP kinase. However, EGCG hardly affected the PGF_{2α}-induced phosphorylation of p44/p42 MAP kinase (Fig. 1).

Effect PGF_{2α} of on the Phosphorylation of p38 MAP Kinase and SAPK/JNK in MC3T3-E1 Cells

Among the MAP kinase superfamily, p38 MAP kinase and SAPK/JNK in addition to p44/p42 MAP kinase play a crucial role as intracellular components to transduce the various signals of agonists [Widmann et al., 1999]. Therefore, in order to investigate whether activates p38 MAP kinase or SAPK/JNK in MC3T3-E1 cells, we next examined the effect of PGF_{2α} on the phosphorylation of p38 MAP kinase and SAPK/JNK. PGF_{2α} induced both the phosphorylation of p38 MAP kinase (lane 2 in Fig. 2) and SAPK/JNK (lane 2 in Fig. 3).




Fig. 2. Effect of EGCG on the phosphorylation of p38 MAP kinase induced by PGF_{2α} in MC3T3-E1 cells. The cultured cells were pretreated with the indicated doses of EGCG or vehicle for 60 min, and then stimulated by 10 μM PGF_{2α} or vehicle for 10 min. The extracts of cells were subjected to SDS-PAGE with subsequent Western blotting analysis with antibodies against phospho-specific p38 MAP kinase or p38 MAP kinase. The histogram shows quantitative representations of the levels of PGF_{2α}-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean ± SEM of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

Effects of SB203580 or SP600125 on the PGF_{2α}-Stimulated VEGF Synthesis in MC3T3-E1 Cells

In order to clarify the involvement of the MAP kinases in the PGF_{2α}-stimulated VEGF synthesis in MC3T3-E1 cells, we examined the effect of SB203580, a specific inhibitor of p38 MAP kinase [Cuenda et al., 1995] or SP600125, a specific SAPK/JNK inhibitor [Bennett et al., 2001], on the VEGF synthesis. SB203580 or SP600125, which by itself had little effect on the VEGF levels, significantly suppressed the PGF_{2α}-stimulated synthesis of VEGF (Table II).

Effect of EGCG on the Phosphorylation of p38 MAP Kinase or SAPK/JNK Induced by PGF_{2α} in MC3T3-E1 Cells

In order to clarify whether p38 MAP kinase is involved in EGCG effect on the PGF_{2α}-




Fig. 3. Effect of EGCG on the phosphorylation of SAPK/JNK induced by PGF $_{2\alpha}$ in MC3T3-E1 cells. The cultured cells were pretreated with the indicated doses of EGCG or vehicle for 60 min, and then stimulated by 10 μM PGF $_{2\alpha}$ or vehicle for 20 min. The extracts of cells were subjected to SDS-PAGE with subsequent Western blotting analysis with antibodies against phospho-specific SAPK/JNK or SAPK/JNK. The histogram shows quantitative representations of the levels of PGF $_{2\alpha}$ -induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean \pm SEM of triplicate determinations. Similar results were obtained with two additional and different cell preparations. * $P < 0.05$, compared to the value of PGF $_{2\alpha}$ alone.

stimulated VEGF synthesis in MC3T3-E1 cells, we examined the effect of EGCG on the PGF $_{2\alpha}$ -induced phosphorylation of p38 MAP kinase. However, EGCG had little effect on the PGF $_{2\alpha}$ -

TABLE II. Effects of SB203580 or SP600125 on the PGF $_{2\alpha}$ -Stimulated VEGF Synthesis in MC3T3-E1 Cells

Inhibitor (μM)	PGF $_{2\alpha}$	VEGF (pg/ml)
Vehicle	-	20 \pm 10
Vehicle	+	559 \pm 51
SB203580 (3)	-	13 \pm 10
SB203580 (3)	+	221 \pm 35*
SB203580 (10)	-	15 \pm 10
SB203580 (10)	+	46 \pm 12*
SP600125 (1)	-	15 \pm 10
SP600125 (1)	+	399 \pm 37*
SP600125 (30)	-	13 \pm 10
SP600125 (30)	+	162 \pm 29*

The cultured cells were pretreated with SB203580, SP600125, or vehicle for 60 min, and then stimulated by 10 μM PGF $_{2\alpha}$ or vehicle for 24 h. Each value represents the mean \pm SEM of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

* $P < 0.05$, compared to the value of PGF $_{2\alpha}$ alone.

induced phosphorylation of p38 MAP kinase (Fig. 2). On the contrary, EGCG markedly enhanced the PGF $_{2\alpha}$ -induced phosphorylation of SAPK/JNK (Fig. 3). According to the densitometric analysis, EGCG (10 μM) caused about 100% enhancement of the PGF $_{2\alpha}$ effect on the SAPK/JNK phosphorylation.

Effect of SP600125 on the Enhancement by EGCG of PGF $_{2\alpha}$ -Induced Phosphorylation of SAPK/JNK in MC3T3-E1 Cells

SP600125 [Bennett et al., 2001], which by itself did not affect the basal levels of phosphorylation of SAPK/JNK, significantly reduced the enhancement by EGCG of PGF $_{2\alpha}$ -induced SAPK/JNK phosphorylation (Fig. 4). The enhanced phosphorylation levels by EGCG of PGF $_{2\alpha}$ -induced SAPK/JNK were suppressed by




Fig. 4. Effect of SP600125 on the enhancement by EGCG of the PGF $_{2\alpha}$ -induced phosphorylation of SAPK/JNK in MC3T3-E1 cells. The cultured cells were pretreated with 30 μM SP600125 or vehicle for 60 min, and then incubated by 10 μM EGCG for 60 min. The cells were stimulated by 10 μM PGF $_{2\alpha}$ or vehicle for 20 min. The extracts of cells were subjected to SDS-PAGE with subsequent Western blotting analysis with antibodies against phospho-specific SAPK/JNK or SAPK/JNK. The histogram shows quantitative representations of the levels of PGF $_{2\alpha}$ -induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean \pm SEM of triplicate determinations. Similar results were obtained with two additional and different cell preparations. * $P < 0.05$, compared to the value of PGF $_{2\alpha}$ alone. ** $P < 0.05$, compared to the value of PGF $_{2\alpha}$ alone. *** $P < 0.05$, compared to the value of PGF $_{2\alpha}$ with EGCG pretreatment.

SP600125 similar to the levels by PGF_{2α} with SP600125 treatment.

Effect of EGCG on the Phosphorylation of c-Jun Induced by PGF_{2α} in MC3T3-E1 Cells

It is well known that c-Jun acts as a downstream effector of SAPK/JNK [Widmann et al., 1999; Weston and Davis, 2002]. Thus, in order to investigate whether EGCG effect on the PGF_{2α}-stimulated VEGF synthesis is mediated through c-Jun activation in MC3T3-E1 cells, we next examined the effect of EGCG on the PGF_{2α}-induced phosphorylation of c-Jun. We found that PGF_{2α} time-dependently phosphorylated c-Jun and that SP600125 suppressed the PGF_{2α}-induced phosphorylation of c-Jun in these cells (data not shown). EGCG markedly amplified the PGF_{2α}-induced phosphorylation of c-Jun (Fig. 5). According to the densitometric

analysis, EGCG (10 μM) caused about 80% enhancement of the PGF_{2α} effect on the c-Jun phosphorylation.

DISCUSSION

In the present study, we demonstrated that EGCG, which alone did not affect the levels of VEGF, significantly enhanced the PGF_{2α}-stimulated VEGF synthesis in osteoblast-like MC3T3-E1 cells. We next investigated the mechanism of EGCG behind the amplifying effect on the VEGF synthesis. It is well recognized that the MAP kinase superfamily mediates intracellular signaling of extracellular agonists and plays a crucial role in cellular functions including proliferation, differentiation, and cell death in a variety of cells [Widmann et al., 1999]. Three major MAP kinase, p44/p42 MAP kinase, p38 MAP kinase, and SAPK/JNK are known as central elements used by mammalian cells to transduce the diverse messages [Widmann et al., 1999]. In our previous study [Tokuda et al., 2003], we have shown that PGF_{2α}-activated p44/p42 MAP kinase acts as a positive regulator in PGF_{2α}-induced VEGF synthesis in osteoblast-like MC3T3-E1 cells. In the present study, we showed that EGCG failed to affect the PGF_{2α}-induced phosphorylation of p44/p42 MAP kinase. Thus, it seems unlikely that EGCG amplifies the PGF_{2α}-induced VEGF synthesis through upregulating the activation of p44/p42 MAP kinase in osteoblast-like MC3T3-E1 cells.

We further investigated the involvement of p38 MAP kinase and SAPK/JNK in the amplifying effect of EGCG. We found that PGF_{2α} induced both the phosphorylation of p38 MAP kinase and SAPK/JNK. It has been shown that MAP kinases are activated by phosphorylation of threonine and tyrosine residues by dual specificity MAP kinase kinases [Widmann et al., 1999]. Therefore, our findings strongly suggest that PGF_{2α} activates p38 MAP kinase and SAPK/JNK in addition to p44/p42 MAP kinase in osteoblast-like MC3T3-E1 cells. In addition, we showed that the PGF_{2α}-stimulated VEGF synthesis was suppressed by a specific p38 MAP kinase inhibitor, SB203580 [Cuenda et al., 1995], or a specific SAPK/JNK inhibitor, SP600125 [Bennett et al., 2001]. Based on our findings, it is probable that PGF_{2α} stimulates the synthesis of VEGF via the three MAP kinases in osteoblast-like MC3T3-E1 cells.




Fig. 5. Effect of EGCG on the phosphorylation of c-Jun induced by PGF_{2α} in MC3T3-E1 cells. The cultured cells were pretreated with the indicated doses of EGCG or vehicle for 60 min, and then stimulated by 10 μM PGF_{2α} or vehicle for 20 min. The extracts of cells were subjected to SDS-PAGE with subsequent Western blotting analysis with antibodies against c-Jun or SAPK/JNK. The histogram shows quantitative representations of the levels of PGF_{2α}-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean ± SEM of triplicate determinations. Similar results were obtained with two additional and different cell preparations. **P* < 0.05, compared to the control value. ***P* < 0.05, compared to the value of PGF_{2α} alone.

However, EGCG had little effect on the $\text{PGF}_{2\alpha}$ -induced p38 MAP kinase phosphorylation. Therefore, it seems unlikely that the EGCG-induced enhancement of $\text{PGF}_{2\alpha}$ -stimulated VEGF synthesis is due to the upregulation of p38 MAP kinase activation. On the contrary, we showed that the $\text{PGF}_{2\alpha}$ -induced SAPK/JNK phosphorylation was markedly amplified by EGCG. Furthermore, SP600125 [Bennett et al., 2001] markedly reduced the enhancement by EGCG almost to the levels of $\text{PGF}_{2\alpha}$ alone with SP600125 in the phosphorylation of SAPK/JNK. These results suggest that EGCG upregulates the $\text{PGF}_{2\alpha}$ -stimulated activation of SAPK/JNK. Additionally, we demonstrated that EGCG strengthened the $\text{PGF}_{2\alpha}$ -induced phosphorylation of c-Jun, well known as a downstream effector of SAPK/JNK [Widmann et al., 1999; Weston and Davis, 2002], as well as SAPK/JNK. Taking our findings into account as a whole, it is most likely that EGCG upregulates $\text{PGF}_{2\alpha}$ -stimulated VEGF synthesis through enhancing the activation of SAPK/JNK in osteoblast-like MC3T3-E1 cells. Further investigations are necessary to clarify the precise mechanism of catechin behind the amplification of VEGF synthesis in osteoblasts.

It is generally recognized that the expansion of capillary network providing microvasculature is an essential process of bone remodeling [Erlebacher et al., 1995]. Since VEGF is a specific mitogen of vascular endothelium [Ferrara and Davis-Smyth, 1997], it is speculated that VEGF synthesized by osteoblasts functions as a pivotal intercellular mediator between osteoblasts and vascular endothelial cells. Moreover, it has been reported that VEGF is involved in trabecular bone formation and expansion of the hypertrophic chondrocyte zone in epiphyseal growth plate of mouse [Gerber et al., 1999], supporting the importance of VEGF in bone metabolism. On the other hand, it has been shown that catechin exerts an inhibitory effect on bone resorption [Delaisse et al., 1986]. Additionally, in osteoblasts, production of bone-resorptive cytokines such as tumor necrosis factor- α and interleukin-6 has been reported to be suppressed by catechin [Choi and Hwang, 2003]. Based on our results as a whole, it is probable that catechin-enhanced VEGF synthesized from osteoblasts acts a crucial role in the process of bone remodeling via regulating the capillary endothelial cells proliferation. Further investigations are

required to elucidate the role of catechin in bone metabolism.

In conclusion, our present results strongly suggest that catechin upregulates $\text{PGF}_{2\alpha}$ -stimulated VEGF synthesis via enhancing activation of SAPK/JNK among the MAP kinase superfamily in osteoblasts.

ACKNOWLEDGMENTS

We are very grateful to Yoko Kawamura and Seiko Sakakibara for their skillful technical assistance.

REFERENCES

- Bennett BL, Sasaki DT, Murray BW, O'Leary EC, Sakata ST, Xu W, Leisten JC, Motiwala A, Pierce S, Satoh Y, Bhagwat SS, Manning AM, Anderson DW. 2001. SP600125, an anthranyrazolone inhibitor of Jun N-terminal Kinase. *Proc Natl Acad Sci USA* 98:13681-13686.
- Choi E-M, Hwang J-K. 2003. Effects of (+)-catechin on the function of osteoblastic cells. *Biol Pharm Bull* 26:523-526.
- Cuenda A, Rouse J, Doza YN, Meier R, Cohen P, Gallagher TF, Young PR, Lee JC. 1995. SB203580 is a specific inhibitor of a MAP kinase homologue which is stimulated by cellular stresses and interleukin-1. *FEBS Lett* 364:229-233.
- Delaisse JM, Eeckhout Y, Vaes G. 1986. Inhibition of bone resorption in culture by (+)-catechin. *Biochem Pharmacol* 35:3091-3094.
- Erlebacher A, Filvaroff EH, Girelman SE, Derynck R. 1995. Toward a molecular understanding of skeletal development. *Cell* 80:371-378.
- Exton JH. 1999. Regulation of phospholipase D. *Biochim Biophys Acta* 439:121-133.
- Ferrara N, Davis-Smyth T. 1997. The biology of vascular endothelial growth factor. *Endocr Rev* 18:4-25.
- Gerber HP, Vu TH, Ryan AM, Kowalski J, Werb Z, Ferrara N. 1999. VEGF couples hypertrophic cartilage remodeling, ossification and angiogenesis during endochondral bone formation. *Nat Med* 5:623-628.
- Goad DL, Rubin J, Wang H, Jr., Tashjian AH, Patterson C. 1996. Enhanced expression of vascular endothelial growth factor in human SaOS-2 osteoblast-like cells and murine osteoblasts induced by insulin-like growth factor I. *Endocrinology* 137:2262-2268.
- Harbourne JB, Williams CA. 2000. Advances in flavonoid research since 1992. *Phytochemistry* 55:481-504.
- Jankun J, Selman SH, Swiercz R, Skrzypczak-Jankun E. 1997. Why drinking green tea could prevent cancer. *Nature* 387:561.
- Kato K, Ito H, Hasegawa K, Inaguma Y, Kozawa O, Asano T. 1996. Modulation of the stress-induced synthesis of hsp27 and α B-crystallin by cyclic AMP in C6 glioma cells. *J Neurochem* 66:946-950.
- Kozawa O, Tokuda H, Miwa M, Kotoyori J, Oiso Y. 1992. Cross-talk regulation between cyclic AMP production and phosphoinositide hydrolysis induced by prostaglandin E_2 in osteoblast-like cells. *Exp Cell Res* 198:130-134.

- Kozawa O, Suzuki A, Kotoyori J, Tokuda H, Watanabe Y, Ito Y, Oiso Y. 1994. Prostaglandin $\text{F}_{2\alpha}$ activates phospholipase D independently from activation of protein kinase C in osteoblast-like cells. *J Cell Biochem* 55:373-379.
- Laemmli UK. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685.
- Miwa M, Tokuda H, Tsushita K, Kotoyori J, Takahashi Y, Ozaki N, Kozawa O, Oiso Y. 1990. Involvement of pertussis toxin-sensitive GTP-binding protein in prostaglandin $\text{F}_{2\alpha}$ -induced phosphoinositide hydrolysis in osteoblast-like cells. *Biochem Biophys Res Commun* 171:1229-1235.
- Nijweide PJ, Burger EH, Feyen JHM. 1986. Cells of bone: Proliferation, differentiation and hormonal regulation. *Physiol Rev* 66:855-886.
- Nishizuka Y. 1992. Intracellular signaling by hydrolysis of phospholipids and activation of protein kinase C. *Science* 258:607-614.
- Pilbeam CC, Harrison JR, Raisz LG. 1996. Prostaglandins and bone metabolism. In: Bilezikian JP, Raisz LG, Rodan GA, editors. *Principles of bone biology*. San Diego: Academic Press. pp 715-728.
- Schlaeppi JM, Gutzwiller S, Finlenzeller G, Fournier B. 1997. 1,25-dihydroxyvitamin D_3 induces the expression of vascular endothelial growth factor in osteoblastic cells. *Endocr Res* 23:213-229.
- Sudo H, Kodama H, Amagai Y, Yamamoto S, Kasai S. 1983. *In vivo* differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria. *J Cell Biol* 96:191-198.
- Tokuda H, Harada A, Hirade K, Matsuno H, Ito H, Kato K, Oiso Y, Kozawa O. 2003. Incadronate amplifies prostaglandin $\text{F}_{2\alpha}$ -induced vascular endothelial growth factor synthesis in osteoblasts. Enhancement of MAPK activity. *J Biol Chem* 278:18930-18937.
- Wang DS, Yamazaki K, Nohtomi K, Shizume K, Ohsumi K, Shibuya M, Demura H, Sato K. 1996. Increase of vascular endothelial growth factor mRNA expression by 1,25-dihydroxyvitamin D_3 in human osteoblast-like cells. *J Bone Miner Res* 11:472-479.
- Weston CR, Davis RJ. 2002. The JNK signal transduction pathway. *Curr Opin Genet Dev* 12:14-21.
- Widmann C, Gibson S, Jarpe MB, Johnson GL. 1999. Mitogen-activated protein kinase: Conservation of a three-kinase module from yeast to human. *Physiol Rev* 79:143-180.

(–)-Epigallocatechin gallate suppresses endothelin-1-induced interleukin-6 synthesis in osteoblasts: Inhibition of p44/p42 MAP kinase activation

Haruhiko Tokuda^{a,b,*}, Shinji Takai^b, Yoshiteru Hanai^{a,b}, Rie Matsushima-Nishiwaki^b, Takayuki Hosoi^c, Atsushi Harada^d, Toshiki Ohta^e, Osamu Kozawa^b

^a Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

^b Department of Pharmacology, Gifu University Graduate School of Medicine, Gifu 501-1194, Japan

^c Department of Advanced Medicine, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

^d Department of Restorative Medicine, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

^e Department of Internal Medicine, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan

Received 17 February 2007; accepted 22 February 2007

Available online 1 March 2007

Edited by Beat Imhof

Abstract We previously showed that endothelin-1 (ET-1) stimulates the synthesis of interleukin-6 (IL-6), a potent bone resorptive agent, in osteoblast-like MC3T3-E1 cells, and that protein kinase C (PKC)-dependent p44/p42 mitogen-activated protein (MAP) kinase plays a part in the IL-6 synthesis. In the present study, we investigated the effect of (–)-epigallocatechin gallate (EGCG), one of the major flavonoids containing in green tea, on ET-1-induced IL-6 synthesis in osteoblasts and the underlying mechanism. EGCG significantly reduced the synthesis of IL-6 stimulated by ET-1 in MC3T3-E1 cells as well primary cultured mouse osteoblasts. SB203580, a specific inhibitor of p38 MAP kinase, but not SP600125, a specific SAPK/JNK inhibitor, suppressed ET-1-stimulated IL-6 synthesis. ET-1-induced phosphorylation of p38 MAP kinase was not affected by EGCG. On the other hand, EGCG suppressed the phosphorylation of p44/p42 MAP kinase induced by ET-1. Both the IL-6 synthesis and the phosphorylation of p44/p42 MAP kinase stimulated by 12-*O*-tetradecanoylphorbol 13-acetate (TPA), a direct activator of PKC, were markedly suppressed by EGCG. The phosphorylation of MEK1/2 and Raf-1 induced by ET-1 or TPA were also inhibited by EGCG. These results strongly suggest that EGCG inhibits ET-1-stimulated synthesis of IL-6 via suppression of p44/p42 MAP kinase pathway in osteoblasts, and the inhibitory effect is exerted at a point between PKC and Raf-1 in the ET-1 signaling cascade.

© 2007 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

Keywords: Catechin; Endothelin-1; Interleukin-6; MAP kinase; Osteoblast

1. Introduction

Interleukin-6 (IL-6) is a pleiotropic cytokine that has important physiological effects on a wide range of functions such as

promoting B cell differentiation, T cell activation and inducing acute phase proteins [1–4]. The bone metabolism is regulated by two functional cells, osteoblasts and osteoclasts, responsible for bone formation and bone resorption, respectively [5]. The formation of bone structures and bone remodeling results from the coupling process, bone resorption by activated osteoclasts with subsequent deposition of new matrix by osteoblasts. It is well recognized that IL-6 is one of the most potent osteoclastogenic factors in the bone metabolism [3,4]. Bone resorption is mediated by the increased local production of inflammatory cytokines such as tumor necrosis factor- α and IL-1. In osteoblasts [6–8], bone resorptive agents such as tumor necrosis factor- α and IL-1 have been reported to stimulate the synthesis of IL-6. As for the bone metabolism, IL-6 has been shown to stimulate bone resorption and induce osteoclast formation [3,4,6,9]. Therefore, accumulating evidence indicates that IL-6 secreted from osteoblasts plays a key role as a downstream effector of bone resorptive agents. In previous studies [10,11], we have shown that endothelin-1 (ET-1) induces the activation of p44/p42 mitogen-activated protein (MAP) kinase, a member of the MAP kinase superfamily [12], via protein kinase C (PKC), resulting in the IL-6 synthesis in osteoblast-like MC3T3-E1 cells. However, the mechanism of ET-1 behind the IL-6 synthesis in osteoblasts has not yet been precisely clarified.

It is well known that compounds in foods such as vegetables and fruits have beneficial properties on human beings. Among them, it has been reported that flavonoids possess antioxidative, antibacterial and antitumor effects [13,14]. Catechins are one of the major flavonoids, which are present in various species of plants such as green tea [14]. In bone metabolism, catechin has been shown to suppress bone resorption [15]. As for osteoblasts, it has been shown that catechin stimulates alkaline phosphatase activity, a mature osteoblast phenotype [5], and reduces apoptosis in osteoblast-like MC3T3-E1 cells [16]. However, the precise mechanism of catechin in osteoblasts is not fully known.

In the present study, we investigated whether (–)-epigallocatechin gallate (EGCG), one of the major flavonoids containing in green tea, affects the ET-1-stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells and the mechanism behind it. We herein show that EGCG reduces the ET-1-stimulated

*Corresponding author. Address: Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu 474-8511, Japan.
Fax: +81 562 46 8396.
E-mail address: tokuda@ncgg.go.jp (H. Tokuda).

IL-6 synthesis via attenuation of p44/p42 MAP kinase pathway in these cells.

2. Materials and methods

2.1. Materials

ET-1 and 12-*O*-tetradecanoylphorbol 13-acetate (TPA) were purchased from Sigma Chemical Co. (St. Louis, MO). IL-6 ELISA kit was purchased from R&D Systems, Inc. (Minneapolis, MN). EGCG, SB203580 and SP600125 were obtained from Calbiochem-Novabiochem Co. (La Jolla, CA). Phospho-specific p38 MAP kinase antibodies, p38 MAP kinase antibodies, phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific MEK1/2 antibodies, MEK1/2 antibodies, phospho-specific Raf-1 antibodies and β -actin antibodies were purchased from Cell Signaling Technology (Beverly, MA). ECL Western blotting detection system was purchased from Amersham Japan (Tokyo, Japan). Other materials and chemicals were obtained from commercial sources. SB203580 and SP600125 were dissolved in dimethyl sulfoxide. The maximum concentration of dimethyl sulfoxide was 0.1%, which did not affect either the assay for IL-6 activity or the Western blotting analysis.

2.2. Cell culture

The cloned osteoblast-like MC3T3-E1 cells, which have been derived from newborn mouse calvaria [17], were maintained as previously described [18]. Briefly, the cells were cultured in α -minimum essential medium (α -MEM) containing 10% fetal calf serum (FCS) at 37 °C in a humidified atmosphere of 5% CO₂/95% air. The cells were seeded into 35-mm diameter dishes (5×10^4 /dish) or 90-mm diameter dishes (5×10^5 /dish) in α -MEM containing 10% FCS. After 5 days, the medium was exchanged for α -MEM containing 0.3% FCS. The cells were used for experiments after 48 h.

Primary cultured mouse osteoblasts were obtained from neonatal balb/c mouse by sequential collagenase digestions as previously described [19]. In brief, the calvarias of the neonatal mice were aseptically dissected from neonatal mice. The calvarias which were divided into small pieces were sequentially digested with 5 ml of PBS containing 0.1% collagenase (Sigma, MO) for 5 min at 37 °C. The cells isolated in fractions 2–6 were seeded into 90-mm diameter dishes in α -MEM containing 10% FCS at 37 °C in a humidified atmosphere of 5% CO₂/95% air. The medium was exchanged at every 3 days. After reaching confluence, the cells were subcultured. After trypsinization using trypsin-EDTA (0.05%/0.53 mM), the cells were rinsed three times in 10 ml of PBS. The cells (1×10^5) were seeded into 35-mm diameter dishes in 2 ml of α -MEM containing 10% FCS. After reaching confluency, the medium was exchanged for α -MEM containing 0.3% FCS. The cells were used for experiments after 48 h. We confirmed that the cultured cells possessed the high alkaline phosphatase activity, a mature osteoblasts phenotype [20].

2.3. Assay for IL-6

The cultured cells were stimulated by ET-1 in 1 ml of α -MEM containing 0.3% FCS, and then incubated for the indicated periods. The conditioned medium was collected, and IL-6 in the medium was then measured by an IL-6 ELISA kit. When indicated, the cells were pretreated with EGCG, SB203580 or SP600125 for 60 min.

2.4. Analysis of Western blotting

The cultured cells were stimulated by ET-1 in α -MEM containing 0.3% FCS for the indicated periods. The cells were washed twice with phosphate-buffered saline and then lysed, homogenized and sonicated in a lysis buffer containing 62.5 mM Tris/HCl, pH 6.8, 2% sodium dodecyl sulfate (SDS), 50 mM dithiothreitol and 10% glycerol. The cytosolic fraction was collected as a supernatant after centrifugation at 125000 \times g for 10 min at 4 °C. SDS-polyacrylamide gel electrophoresis (PAGE) was performed by Laemmli [21] in 10% polyacrylamide gel. A Western blotting analysis was performed as described previously [22] by using phospho-specific p38 MAP kinase antibodies, p38 MAP kinase antibodies, phospho-specific p44/p42 MAP kinase or p44/p42 MAP kinase with peroxidase-labeled antibodies raised in goat against rabbit IgG being used as second antibodies. The peroxidase activity on

PVDG membrane was visualized on X-ray film by means of the ECL Western blotting detection system. When indicated, the cells were pretreated with various doses of EGCG for 60 min.

2.5. Determinations

The absorbance of enzyme immunoassay samples was measured at 450 nm with EL 340 Bio Kinetic Reader (Bio-Tek Instruments, Inc., Winooski, VT). A densitometric analysis was performed using the Molecular Analyst/Macintosh software program (Bio-Rad Laboratories, Hercules, CA).

2.6. Statistical analysis

The data were analyzed by ANOVA followed by the Bonferroni method for multiple comparisons between pairs, and a $P < 0.05$ was considered significant. All data are presented as the means \pm S.E.M. of triplicate determinations. Each experiment was repeated three times with similar results.

3. Results

3.1. Effects of EGCG on the ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells and primary cultured mouse osteoblasts

We have previously demonstrated that ET-1 stimulates IL-6 synthesis in osteoblast-like MC3T3-E1 cells [10]. We first examined the effect of EGCG on the ET-1-stimulated IL-6 synthesis. EGCG, which alone had little effect on the IL-6 levels, reduced the ET-1-stimulated synthesis of IL-6 in a dose-dependent manner in the range between 1 and 100 μ M (Table 1). In addition, we also examined the effect of EGCG in primary cultured mouse osteoblasts, and found that EGCG (30 μ M) significantly reduced the IL-6 synthesis induced by ET-1 (Table 2). EGCG (30 μ M) caused about a 55% reduction in the ET-1-effect.

3.2. Effects of SB203580 or SP600125 on the ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells

In our previous study [11], we reported that p44/p42 MAP kinase plays a part in the ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells. Among the MAP kinase superfamily, p38

Table 1
Effects of EGCG, SB203580 or SP600125 on the ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells

EGCG or inhibitor	ET-1	IL-6 (pg/ml)
Vehicle	–	16 \pm 5
Vehicle	+	468 \pm 38*
EGCG (1 μ M)	–	15 \pm 10
EGCG (1 μ M)	+	419 \pm 35
EGCG (10 μ M)	–	17 \pm 8
EGCG (10 μ M)	+	414 \pm 30
EGCG (50 μ M)	–	15 \pm 10
EGCG (50 μ M)	+	347 \pm 28**
EGCG (100 μ M)	–	23 \pm 8
EGCG (100 μ M)	+	69 \pm 12**
SB203580	–	10 \pm 7
SB203580	+	85 \pm 12**
SP600125	–	15 \pm 10
SP600125	+	456 \pm 29

The cultured cells were pretreated with various doses of EGCG, 10 μ M SB203580, 10 μ M SP600125 or vehicle for 60 min, and then stimulated by 0.1 μ M ET-1 or vehicle for 24 h. Each value represents the means \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

* $P < 0.05$, compared to the control value.

** $P < 0.05$, compared to the value of ET-1 alone.

Table 2

Effects of EGCG on the ET-1-stimulated IL-6 synthesis in primary cultured mouse osteoblasts

EGCG	ET-1	IL-6 (pg/ml)
–	–	248 ± 39
–	+	354 ± 26*
+	–	216 ± 8
+	+	263 ± 26**

The cultured cells were pretreated with 30 μ M EGCG or vehicle for 60 min, and then stimulated by 0.1 μ M ET-1 or vehicle for 24 h. Each value represents the means \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

* $P < 0.05$, compared to the control value.

** $P < 0.05$, compared to the value of ET-1 alone.

MAP kinase and stress-activated protein kinase (SAPK)/c-Jun N-terminal kinase (JNK) in addition to p44/p42 MAP kinase, which together play a crucial role as intracellular components to transduce the various signals of agonists [12]. We have previously shown that ET-1 also activates p38 MAP kinase and SAPK/JNK [23,24]. In order to clarify the involvement of the MAP kinases in the ET-1-stimulated IL-6 synthesis in these cells, we examined the effect of SB203580, a specific inhibitor of p38 MAP kinase [25], or SP600125, a specific SAPK/JNK inhibitor [26], on the IL-6 synthesis. SB203580, which by itself had little effect on the IL-6 levels, significantly suppressed the ET-1-stimulated synthesis of IL-6 (Table 1). On the other hand, SP600125 hardly affected the IL-6 synthesis induced by ET-1 (Table 1).

3.3. Effect of EGCG on the phosphorylation of p38 MAP kinase by ET-1 in MC3T3-E1 cells

In order to clarify whether p38 MAP kinase is involved in EGCG-effect on the ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells, we examined the effect of EGCG on the ET-1-induced phosphorylation of p38 MAP kinase. However, EGCG had little effect on the ET-1-induced phosphorylation of p38 MAP kinase (Fig. 1).

3.4. Effect of EGCG on the phosphorylation of p44/p42 MAP kinase induced by ET-1 in MC3T3-E1 cells

In order to investigate whether EGCG-effect on the ET-1-stimulated IL-6 synthesis is mediated through p44/p42 MAP kinase activation in MC3T3-E1 cells, we next examined the effect of EGCG on the ET-1-induced phosphorylation of p44/p42 MAP kinase. EGCG markedly attenuated the ET-1-induced phosphorylation of p44/p42 MAP kinase (Fig. 2). According to the densitometric analysis, EGCG (100 μ M) caused about 80% reduction in the ET-1-effect.

3.5. Effects of EGCG on the TPA-stimulated IL-6 synthesis in MC3T3-E1 cells

We previously reported that PKC functions at a point upstream from p44/p42 MAP kinase in the ET-1-stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells [11]. We next examined the effect of EGCG on the IL-6 synthesis stimulated by TPA, a direct activator of PKC [27]. In our previous study [28], we found that TPA alone stimulates IL-6 synthesis in MC3T3-E1 cells. EGCG, which by itself had little effect on the IL-6 levels, significantly reduced the TPA-stimulated syn-




Fig. 1. Effect of EGCG on the phosphorylation of p38 MAP kinase induced by ET-1 in MC3T3-E1 cells. The cultured cells were pretreated with the indicated doses of EGCG or vehicle for 60 min, and then stimulated by 3 nM ET-1 or vehicle for 20 min. The extracts of cells were subjected to SDS-PAGE with a subsequent Western blotting analysis with antibodies against phospho-specific p38 MAP kinase or p38 MAP kinase. The histogram shows quantitative representations of the levels of ET-1-induced phosphorylation obtained from a laser densitometric analysis of three independent experiments. Each value represents the means \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

thesis of IL-6 (Table 3). EGCG (100 μ M) caused about an 80% reduction in the TPA-effect.

3.6. Effect of EGCG on the phosphorylation of p44/p42 MAP kinase induced by TPA in MC3T3-E1 cells

In order to investigate whether EGCG-effect on the TPA-stimulated IL-6 synthesis is mediated through inhibition of p44/p42 MAP kinase activation in MC3T3-E1 cells, we next examined the effect of EGCG on the TPA-induced phosphorylation of p44/p42 MAP kinase. We previously found that TPA induces the phosphorylation of p44/p42 MAP kinase in these cells [29]. EGCG markedly suppressed the TPA-induced phosphorylation of p44/p42 MAP kinase (Fig. 3).

3.7. Effects of EGCG on the phosphorylation of MEK1/2 and Raf-1 induced by ET-1 or TPA in MC3T3-E1 cells

It is well known that p44/p42 MAP kinase is activated by an upstream kinase, MEK1/2, which is activated by Raf-1 [12]. To clarify the exact mechanism of EGCG behind the IL-6 synthesis, we further examined the effects of EGCG on the phosphorylation of MEK1/2 and Raf-1 induced by ET-1 or TPA in MC3T3-E1 cells. The levels of ET-1 or TPA-induced phosphorylation of MEK1/2 were reduced by EGCG (data not shown). EGCG suppressed the levels of both ET-1- and TPA-induced the phosphorylation of Raf-1 (Fig. 4).




Fig. 2. Effect of EGCG on the phosphorylation of p44/p42 MAP kinase induced by ET-1 in MC3T3-E1 cells. The cultured cells were pretreated with the indicated doses of EGCG or vehicle for 60 min, and then stimulated by 3 nM ET-1 or vehicle for 5 min. The extracts of the cells were subjected to SDS-PAGE with a subsequent Western blotting analysis with antibodies against phospho-specific p44/p42 MAP kinase or p44/p42 MAP kinase. The histogram shows quantitative representations of the levels of ET-1-induced phosphorylation obtained from a laser densitometric analysis of three independent experiments. Each value represents the means \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. * $P < 0.05$, compared to the value of ET-1 alone.

Table 3
Effect of EGCG on the TPA-stimulated IL-6 synthesis in MC3T3-E1 cells

EGCG	TPA	IL-6 (pg/ml)
-	-	19 \pm 7
-	+	415 \pm 31*
+	-	29 \pm 10
+	+	98 \pm 17**

The cultured cells were pretreated with 100 μ M EGCG or vehicle for 60 min, and then stimulated by 0.1 μ M TPA or vehicle for 24 h. Each value represents the means \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations.

* $P < 0.05$, compared to the control value.

** $P < 0.05$, compared to the value of TPA alone.

4. Discussion

In the present study, we showed that EGCG, which by itself had little effect on the levels of IL-6, significantly reduced the ET-1-stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells. We found that EGCG reduced the IL-6 synthesis induced by ET-1 also in primary cultured mouse osteoblasts. It is likely that the inhibitory effect of EGCG on the IL-6 synthesis induced by ET-1 is not specific in a clonal osteoblast-like




Fig. 3. Effect of EGCG on the phosphorylation of p44/p42 MAP kinase induced by TPA in MC3T3-E1 cells. The cultured cells were pretreated with 100 μ M EGCG or vehicle for 60 min, and then stimulated by 0.1 μ M TPA or vehicle for 60 min. The extracts of cells were subjected to SDS-PAGE with a subsequent Western blotting analysis with antibodies against phospho-specific p44/p42 MAP kinase or p44/p42 MAP kinase. The histogram shows quantitative representations of the levels of TPA-induced phosphorylation obtained from a laser densitometric analysis of three independent experiments. Each value represents the means \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. * $P < 0.05$, compared to the value of TPA alone.

MC3T3-E1 cells but it is common in osteoblasts. We next investigated the mechanism of EGCG underlying the suppressive effect on the IL-6 synthesis. It is generally known that the MAP kinase superfamily plays a pivotal role in cellular functions including proliferation, differentiation, and cell death in a variety of cells [12]. Three major MAP kinase, p44/p42 MAP kinase, p38 MAP kinase, and SAPK/JNK are known as central elements used by mammalian cells to transduce the diverse messages [12]. We have previously reported that ET-1-activated p44/p42 MAP kinase acts as a positive regulator in ET-1-induced IL-6 synthesis in osteoblast-like MC3T3-E1 cells [11]. In addition, we demonstrated that p38 MAP kinase and SAPK/JNK are activated by ET-1 in these cells [23,24]. In the present study, we showed that SB203580 [25], markedly suppressed the ET-1-induced IL-6 synthesis while SP600125 [26], failed to affect the IL-6 synthesis. We previously showed that ET-1-induced both synthesis of vascular endothelial growth factor and phosphorylation of SAPK/JNK are significantly suppressed by 10 μ M SP600125 in MC3T3-E1 cells [24]. It thus seems unlikely that SAPK/JNK is involved in the ET-1-induced IL-6 synthesis in osteoblast-like MC3T3-E1 cells. Based on our results, it is probable that ET-1 stimulates the synthesis of IL-6 via p38 MAP kinase in addition to p44/p42 MAP kinase in osteoblast-like MC3T3-E1 cells.

We further investigated the involvement of p38 MAP kinase and p44/p42 MAP kinase in the inhibitory effect of EGCG on the IL-6 synthesis. EGCG attenuated the ET-1-induced phosphorylation of p44/p42 MAP kinase without affecting the p38 MAP kinase phosphorylation. Therefore, our findings strongly




Fig. 4. Effects of EGCG on the phosphorylation of Raf-1 induced by ET-1 or TPA in MC3T3-E1 cells. The cultured cells were pretreated with 100 μ M EGCG or vehicle for 60 min, and then stimulated by 0.1 μ M ET-1 (A), 0.1 μ M TPA (B) or vehicle for 60 min. The extracts of cells were subjected to SDS-PAGE with a subsequent Western blotting analysis with antibodies against phospho-specific Raf-1 or β -actin. The histogram shows quantitative representations of the levels of ET-1- (A) or TPA-induced phosphorylation (B) obtained from a laser densitometric analysis of three independent experiments. Each value represents the means \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. * P < 0.05, compared to the value of ET-1 (A) or TPA (B) alone.

suggest that EGCG inhibits the ET-1-induced IL-6 synthesis via the down-regulating p44/p42 MAP kinase pathway in osteoblast-like MC3T3-E1 cells. In our previous studies [10,11], we demonstrated that p44/p42 MAP kinase acts at a point downstream of PKC in ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells. To clarify whether the effect of catechin

is exerted at a point downstream of PKC or not, we additionally examined the effect of EGCG on the IL-6 synthesis by TPA [27]. We have found that TPA alone stimulates IL-6 synthesis in MC3T3-E1 cells [28]. We herein presented that EGCG reduced the TPA-stimulated IL-6 synthesis as well as the TPA-induced phosphorylation of p44/p42 MAP kinase in these cells, suggesting that the effect of EGCG is exerted at a point upstream of p44/p42 MAP kinase. Furthermore, we found that EGCG actually suppressed the levels of both ET-1- and TPA-induced phosphorylation of MEK1/2 and Raf-1. Taking our findings into account as a whole, it is most likely that EGCG down-regulates ET-1-stimulated IL-6 synthesis in osteoblasts, and that the effect is exerted at a point between PKC and Raf-1. Further investigations are necessary to clarify the precise mechanism of catechin underlying the down-regulation of IL-6 synthesis in osteoblasts.

IL-6 is well recognized to be a potent bone resorptive agent and induces osteoclast formation [4,6]. On the other hand, catechin reportedly has an inhibitory effect on bone resorption [15]. In addition, it has recently been shown that catechin increases cell viability and alkaline phosphatase activity, a marker of mature osteoblast phenotype [5], in osteoblast-like MC3T3-E1 cells and apoptosis of these cells is suppressed by catechin [16]. Taking our results into account, it is probable that catechin-induced suppression of p44/p42 MAP kinase activation plays an effect against bone resorption via down-regulating IL-6 synthesis in osteoblasts. Our present data therefore provide new insight into the pharmacological effects of catechin action on bone cells. Further investigations are required to elucidate the exact role of catechin in the bone metabolism.

In conclusion, these results strongly suggest that catechin inhibits the ET-1-stimulated synthesis of IL-6 via suppression of p44/p42 MAP kinase pathway in osteoblasts, and this effect is exerted at a point between PKC and Raf-1 in the ET-1 signaling cascade.

Acknowledgements: We are very grateful to Yoko Kawamura and Seiko Sakakibara for their skillful technical assistance. This investigation was supported in part by Grant-in-Aid for Scientific Research (16590873 and 16591482) for the Ministry of Education, Science, Sports and Culture of Japan, the Research Grants for Longevity Sciences (15A-1, 15C-2 and 17A-3), Research Grant on Proteomics and Research Grant on Fracture and Dementia from the Ministry of Health, Labour and Welfare of Japan.

References

- [1] Akira, S., Taga, T. and Kishimoto, T. (1993) Interleukin-6 in biology and medicine. *Adv. Immunol.* 54, 1–78.
- [2] Heymann, D. and Rousselle, A.V. (2000) gp130 Cytokine family and bone cells. *Cytokine* 12, 455–468.
- [3] Kwan Tat, S., Padrines, M., Theoleyre, S., Heymann, D. and Fortun, Y. (2004) IL-6, RANKL, ET-1- α /IL-1: interrelations in bone resorption pathophysiology. *Cytokine Growth Factor Rev.* 15, 49–60.
- [4] Blair, H.C., Robinson, L.J. and Zaidi, M. (2005) Osteoclast signalling pathways. *Biochem. Biophys. Res. Commun.* 328, 728–738.
- [5] Nijweide, P.J., Burger, E.H. and Feyen, J.H.M. (1986) Cells of bone: proliferation, differentiation, and humoral regulation. *Physiol. Rev.* 66, 855–886.
- [6] Ishimi, T., Miyaura, C., Jin, C.H., Akatsu, T., Abe, E., Nakamura, Y., Yamaguchi, Y., Yoshiki, S., Matsuda, T., Hirano, T., Kishimoto, T. and Suda, T. (1990) IL-6 is produced by

- osteoblasts and induces bone resorption. *J. Immunol.* 145, 3297–3303.
- [7] Helle, M., Brakenhoff, J.P.J., DeGroot, E.R. and Aarden, L.A. (1988) Interleukin-6 is involved in interleukin-1-induced activities. *Eur. J. Immunol.* 18, 957–959.
- [8] Littlewood, A.J., Russil, J., Harvey, G.R., Hughes, D.E., Russel, R.G.G. and Gowen, M. (1991) The modulation of the expression of IL-6 and its receptor in human osteoblasts *in vitro*. *Endocrinology* 129, 1513–1520.
- [9] Roodman, G.D. (1992) Interleukin-6: an osteotropic factor? *J. Bone Miner. Res.* 7, 475–478.
- [10] Matsuno, M., Kozawa, O., Suzuki, A., Tokuda, H., Kaida, T., Matsuno, H., Niwa, M. and Uematsu, T. (1998) Involvement of protein kinase C activation in endothelin-1-induced secretion of interleukin-6 in osteoblast-like cells. *Cell. Signal.* 10, 107–111.
- [11] Kawamura, H., Otsuka, T., Tokuda, H., Matsuno, H., Niwa, M., Matsui, N., Uematsu, T. and Kozawa, O. (1999) Involvement of p42/p44 MAP kinase in endothelin-1-induced interleukin-6 synthesis in osteoblast-like cells. *Bone* 24, 315–320.
- [12] Widmann, C., Gibson, S., Jarpe, M.B. and Johnson, G.L. (1999) Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol. Rev.* 79, 143–180.
- [13] Jankun, J., Selman, S.H., Swiercz, R. and Skrzypczak-Jankun, E. (1997) Why drinking green tea could prevent cancer. *Nature* 387, 561.
- [14] Harbourne, J.B. and Williams, C.A. (2000) Advances in flavonoid research since 1992. *Phytochemistry* 55, 481–504.
- [15] Delaisse, J.M., Eeckhout, Y. and Vaes, G. (1986) Inhibition of bone resorption in culture by (+)-catechin. *Biochem. Pharmacol.* 35, 3091–3094.
- [16] Choi, E.-M. and Hwang, J.-K. (2003) Effects of (+)-catechin on the function of osteoblastic cells. *Biol. Pharm. Bull.* 26, 523–526.
- [17] Sudo, H., Kodama, H., Amagai, Y., Yamamoto, S. and Kasai, S. (1983) *In vivo* differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria. *J. Cell Biol.* 96, 191–198.
- [18] Kozawa, O., Tokuda, H., Miwa, M., Kotoyori, J. and Oiso, Y. (1992) Cross-talk regulation between cyclic AMP production and phosphoinositide hydrolysis induced by prostaglandin E₂ in osteoblast-like cells. *Exp. Cell Res.* 198, 130–134.
- [19] Ishisaki, A., Tokuda, H., Yoshida, M., Hirade, K., Kunieda, K., Hatakeyama, D., Shibata, T. and Kozawa, O. (2004) Activation of p38 mitogen-activated protein kinase mediates thyroid hormone-stimulated osteocalcin synthesis in osteoblasts. *Mol. Cell. Endocrinol.* 214, 189–195.
- [20] Robinson, R.J., Doty, S.B. and Cooper, R.R. (1973) Electron micrography of mammalian bone in: *Biological Mineralization* (Zipkin, I., Ed.), pp. 257–296, Academic Press, NY.
- [21] Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.
- [22] Kato, K., Ito, H., Hasegawa, K., Inaguma, Y., Kozawa, O. and Asano, T. (1996) Modulation of the stress-induced synthesis of hsp27 and α B-crystallin by cyclic AMP in C6 glioma cells. *J. Neurochem.* 66, 946–950.
- [23] Kawamura, H., Otsuka, T., Matsuno, H., Niwa, N., Matsui, N., Kato, K., Uematsu, T. and Kozawa, O. (1999) Endothelin-1 stimulates heat shock protein 27 induction in osteoblasts: involvement of p38 MAP kinase. *Am. J. Physiol.* 277, E1046–E1054.
- [24] Wang, X., Tokuda, H., Hirade, K. and Kozawa, O. (2002) Stress-activated protein kinase/c-Jun N-terminal kinase (JNK) plays a part in endothelin-1-induced vascular endothelial growth factor synthesis in osteoblasts. *J. Cell. Biochem.* 87, 417–423.
- [25] Cuenda, A., Rouse, J., Doza, Y.N., Meier, R., Cohen, P., Gallagher, T.F., Young, P.R. and Lee, J.C. (1995) SB203580 is a specific inhibitor of a MAP kinase homologue which is stimulated by cellular stresses and interleukin-1. *FEBS Lett.* 364, 229–233.
- [26] Bennett, B.L., Sasaki, D.T., Murray, B.W., O'Leary, E.C., Sakata, S.T., Xu, W., Leisten, J.C., Motiwala, A., Pierce, S., Satoh, Y., Bhagwat, S.S., Manning, A.M. and Anderson, D.W. (2001) SP600125, an anthrapyrazolone inhibitor of Jun N-terminal Kinase. *Proc. Natl. Acad. Sci. USA* 98, 13681–13686.
- [27] Nishizuka, Y. (1986) Studies and perspectives of protein kinase C. *Science* 233, 305–312.
- [28] Kozawa, O., Suzuki, A., Tokuda, H. and Uematsu, T. (1997) Prostaglandin F_{2 α} stimulates interleukin-6 via activation of PKC in osteoblast-like cells. *Am. J. Physiol.* 272, E208–E211.
- [29] Tokuda, H., Harada, A., Hirade, K., Matsuno, H., Ito, H., Kato, K., Oiso, Y. and Kozawa, O. (2003) Incadronate amplifies prostaglandin F_{2 α} -induced vascular endothelial growth factor in osteoblasts: enhancement of MAP kinase activity. *J. Biol. Chem.* 278, 18930–18937.

Rho-kinase regulates endothelin-1-stimulated IL-6 synthesis via p38 MAP kinase in osteoblasts

Haruhiko Tokuda ^{a,b}, Yoshiteru Hanai ^{a,b}, Rie Matsushima-Nishiwaki ^b, Junichi Yamauchi ^b, Tomoaki Doi ^b, Atsushi Harada ^c, Shinji Takai ^b, Osamu Kozawa ^{b,*}

^a Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu, Aichi 474-8511, Japan

^b Department of Pharmacology, Gifu University Graduate School of Medicine, Gifu 501-1194, Japan

^c Department of Functional Restoration, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu, Aichi 474-8511, Japan

Received 2 August 2007

Available online 13 August 2007

Abstract

We have previously reported that endothelin-1 (ET-1) stimulates interleukin-6 (IL-6), a potent bone resorptive agent, through p44/p42 mitogen-activated protein (MAP) kinase and p38 MAP kinase in osteoblast-like MC3T3-E1 cells. In the present study, we investigated the involvement of Rho-kinase in the ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells. ET-1 time-dependently induced the phosphorylation of myosin phosphatase targeting subunit (MYPT-1), a Rho-kinase substrate. Y27632, a specific inhibitor of Rho-kinase, significantly suppressed the IL-6 synthesis induced by ET-1 as well as the MYPT-1 phosphorylation. Fasudil, another inhibitor of Rho-kinase, reduced the ET-1-stimulated IL-6 synthesis. Y27632 as well as fasudil attenuated the ET-1-induced phosphorylation of p38 MAP kinase but not p44/p42 MAP kinase. These results strongly suggest that Rho-kinase regulates ET-1-stimulated IL-6 synthesis through p38 MAP kinase activation in osteoblasts.

© 2007 Elsevier Inc. All rights reserved.

Keywords: ET-1; Rho-kinase; IL-6; Osteoblast

It is well recognized that interleukin-6 (IL-6) is a multifunctional cytokine that has crucial effects on a wide range of functions such as promoting B cell differentiation, T cell activation, and inducing acute phase proteins [1–3]. The bone metabolism is regulated mainly by two functional cells, osteoblasts and osteoclasts, responsible for bone formation and bone resorption, respectively [4]. As for bone metabolism, IL-6 has been shown to stimulate bone resorption and promote osteoclast formation [2,3,5,6]. It has been reported that potent bone resorptive agents such as tumor necrosis factor- α and IL-1 stimulate IL-6 synthesis in osteoblasts [5,7,8]. Currently, evidence is accumulating that IL-6 secreted from osteoblasts plays a pivotal role as a downstream effector of bone resorptive agents.

Endothelin-1 (ET-1) is a potent vasoconstrictor produced by endothelial cells [9]. In bone metabolism, it is well recognized that ET-1 plays an important role in the regulation of bone metabolism and acts as a bone resorptive agent [10]. Accumulating evidence suggests that tumor-produced ET-1 mediates bone metastases by stimulating osteoblast proliferation and new bone formation [11]. As for intracellular signaling of ET-1 in osteoblasts, the activities of ET-1 are mediated via ET_A receptors and ET_B receptors [11]. We have shown that ET-1 activates phospholipase C and phospholipase D through ET_A receptors in osteoblast-like MC3T3-E1 cells [12]. Regarding about IL-6 synthesis, we have reported that ET-1 stimulates IL-6 synthesis via p44/p42 MAP kinase and p38 MAP kinase but not stress-activated protein kinase (SAPK)/c-Jun N terminal kinase (JNK) among the MAP kinase superfamily in these cells [13,14]. However, the exact mechanism

* Corresponding author. Fax: +81 58 230 6215.

E-mail address: okozawa@gifu-u.ac.jp (O. Kozawa).

underlying ET-1-stimulated IL-6 synthesis in osteoblasts is not fully known.

Recent studies suggest that Rho and the down-stream effector, Rho-associated kinase (Rho-kinase) play important roles in a variety of cellular functions such as cell motility and smooth muscle contraction [15–17]. As for osteoblasts, it has been demonstrated that Rho and p38 MAP kinase are involved in the ET-1-induced expression of prostaglandin endoperoxide G/H synthase mRNA in osteoblasts [18]. In addition, Rho/Rho-kinase pathway reportedly stimulates osteoblast proliferation whereas it inhibits osteoblast differentiation [19]. However, the exact role of Rho-kinase in osteoblasts has not yet been fully clarified.

In the present study, we further investigated the exact mechanism behind ET-1-stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells. We here show that Rho-kinase regulates ET-1-stimulated IL-6 synthesis through p38 MAP kinase activation in these cells.

Materials and methods

Materials. ET-1 and mouse IL-6 enzyme immunoassay (ELISA) kit were purchased from R&D Systems, Inc. (Minneapolis, MN). Y27632 was obtained from Calbiochem–Novabiochem Co. (LaJolla, CA). Hydroxyfasudil (fasudil) was from Sigma (St. Louis, MO). Phospho-specific MYPT-1 antibodies were purchased from Upstate (Lake Placid, NY). MYPT-1 antibodies were obtained from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies, and p38 MAP kinase antibodies were purchased from Cell Signaling, Inc. (Beverly, MA). ECL Western blotting detection system was purchased from Amersham Biosciences (Piscataway, NJ). Other materials and chemicals were obtained from commercial sources. Y27632 was dissolved in dimethyl sulfoxide (DMSO). The maximum concentration of DMSO was 0.1%, which did not affect the assay for IL-6 or Western blot analysis.

Cell culture. Cloned osteoblast-like MC3T3-E1 cells derived from newborn mouse calvaria [20] were maintained as previously described [21]. Briefly, the cells were cultured in α -minimum essential medium (α -MEM) containing 10% fetal calf serum (FCS) at 37 °C in a humidified atmosphere of 5% CO₂/95% air. The cells were seeded into 35- or 90-mm diameter dishes in α -MEM containing 10% FCS. After 5 days, the medium was exchanged for α -MEM containing 0.3% FCS. The cells were used for experiments after 48 h.

IL-6 assay. The cultured cells were stimulated by various doses of ET-1 in 1 ml of α -MEM containing 0.3% FCS for the indicated periods. When indicated, the cells were pretreated with various doses of Y27632 or fasudil for 60 min. The conditioned medium was collected at the end of the incubation, and the IL-6 concentration was measured by ELISA kit.

Western blot analysis. The cultured cells were pretreated with Y27632 or fasudil for 60 min, and then stimulated by ET-1 in α -MEM containing 0.3% FCS for the indicated periods. The cells were washed twice with phosphate-buffered saline and then lysed, homogenized, and sonicated in a lysis buffer containing 62.5 mM Tris/HCl, pH 6.8, 2% sodium dodecyl sulfate (SDS), 50 mM dithiothreitol and 10% glycerol. The cytosolic fraction was collected as a supernatant after centrifugation at 125,000g for 10 min at 4 °C. SDS-polyacrylamide gel electrophoresis (PAGE) was performed by Laemmli [22] in 10% polyacrylamide gel. Western blotting analysis was performed as described previously [23] by using phospho-specific MYPT-1 antibodies, MYPT-1 antibodies, phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies or p38 MAP kinase antibodies, with peroxidase-labeled antibodies raised in goat against rabbit IgG being used as second antibodies. Peroxidase activity on the PVDF sheet was visualized on X-ray film by means of the ECL Western blotting detection system.

Determination. The absorbance of enzyme immunoassay samples was measured at 450 nm with EL 340 Bio Kinetic Reader (Bio-Tek Instruments, Inc., Winooski, VT). The densitometric analysis was performed using Molecular Analyst/Macintosh (Bio-Rad Laboratories, Hercules, CA).

Statistical analysis. The data were analyzed by ANOVA followed by the Bonferroni method for multiple comparisons between pairs, and a $p < 0.05$ was considered significant. All data are presented as the means \pm SEM of triplicate determinations. Each experiment was repeated three times with similar results.

Results

Effects of ET-1 on the phosphorylation of MYPT-1 in MC3T3-E1 cells

It is well known that myosin phosphatase targeting subunit (MYPT-1), which is a component of myosin phosphatase, is a down-stream substrate of Rho-kinase [15,24]. To investigate whether ET-1 activates Rho-kinase in osteoblast-like MC3T3-E1 cells, we examined the effect of ET-1 on the phosphorylation of MYPT-1. ET-1 significantly induced the phosphorylation of MYPT-1 in a time-dependent manner (Fig. 1A). The effect of ET-1 on the MYPT-1 phosphorylation reached its peak within 2 min and decreased thereafter (Fig. 1A).

We found that Y27632, a specific inhibitor of Rho-kinase [16], markedly suppressed the ET-1-induced phosphorylation levels of MYPT-1 (Fig. 1B). Additionally, the phosphorylation levels of MYPT-1 induced by ET-1 were attenuated by fasudil, another inhibitor of Rho-kinase [16] (data not shown).

Effects of Y27632 or fasudil on the ET-1-stimulated IL-6 synthesis in MC3T3-E1 cells

We previously showed that ET-1 stimulates IL-6 synthesis in osteoblast-like MC3T3-E1 cells [13]. In order to investigate whether or not Rho-kinase is involved in the ET-1-induced synthesis of IL-6 in MC3T3-E1 cells, we next examined the effect of Y27632 on the synthesis of IL-6 induced by ET-1. Y27632, which alone failed to affect the IL-6 levels, significantly suppressed the ET-1-induced synthesis of IL-6 (Fig. 2). The inhibitory effect of Y27632 was dose-dependent in the range between 0.1 and 10 μ M. Y27632 at 10 μ M caused approximately 90% inhibition in the ET-1-effect.

Fasudil as well as Y27632 reduced the ET-1-stimulated IL-6 synthesis in these cells (Table 1). The effect of fasudil on the IL-6 synthesis was dose-dependent in the range between 0.3 and 3 μ M. Fasudil (3 μ M) caused approximately 70% inhibition in the ET-1-effect.

Effect of Y27632 on the ET-1-induced phosphorylation of p44/p42 MAP kinase in MC3T3-E1 cells

It is generally recognized that three MAP kinases, p44/p42 MAP kinase, p38 MAP kinase and SAPK/JNK are known as central elements used by mammalian cells to