

The goal of antihypertensive therapy is to reduce the incidence of hypertension-related events. Many large-scale clinical trials have already demonstrated the benefits of antihypertensive treatment with drug therapy (3, 4). Based on these results, guidelines for the clinical management of hypertension such as "Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2004)" and the recommendations of the Seventh Report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC-7) have been established and used in the daily management of hypertension (5, 6). According to such guidelines, angiotensin converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) are first-line agents for the treatment of hypertension, especially in hypertensive patients with diabetes.

Losartan potassium (losartan), a subtype 1 (AT1) selective angiotensin II (AII) receptor antagonist, has been widely prescribed worldwide. Several reports have suggested that losartan not only lowers the blood pressure (BP) values, but also has target organ protective effects. The Losartan Intervention for Endpoint Reduction (LIFE) study was a double-blind, prospective, parallel group trial that was designed to compare the effects of losartan with those of the β -blocker atenolol on cardiovascular morbidity and mortality in approximately 8,300 hypertensive patients with left ventricular hypertrophy. It demonstrated that losartan had a more favorable effect on cardiovascular events than atenolol (7). The Reduction of Endpoints in Non-Insulin-Dependent Diabetes Mellitus with the Angiotensin II Antagonist Losartan (RENAAL) study was a multinational, double-blind, randomized, and placebo-controlled trial that enrolled 1,513 patients with type 2 diabetes and nephropathy. It demonstrated a renoprotective effect of losartan (8).

Although studies conducted in Western countries have reported various beneficial effects of losartan therapy (7-11), the actual therapeutic benefit for Japanese patients has been unclear. Because the genetic and environmental background may differ between Japanese and Western patients (12, 13), an investigation of the effects of losartan in Japanese hypertensive patients would be of value. Accordingly, the Japan Hypertension Evaluation with Angiotensin II Antagonist Losartan Therapy (J-HEALTH) study was initiated in 2000 as a large-scale observational study of losartan therapy.

Recently, the significance of home BP monitoring has been an important topic in the management of hypertension (14-16). Since large-scale analysis of home BP data has not been performed in Japan, direct evidence of the significance of home BP values for future cardiovascular events is still lacking (17). Accordingly, the J-HEALTH study was performed to investigate the long-term antihypertensive efficacy and safety of losartan, and the incidence of cardiovascular events and mortality in this population. The study also aimed to investigate whether home BP monitoring would be effective for use in routine antihypertensive treatment.

Table 1. Exclusion Criteria

• Pregnant or could become pregnant, or breast-feeding
• Severe hepatic or renal disease
• Diseases of poor prognosis; malignant neoplasm, performing hemodialysis, or virus infections such as HIV
• Taking the study drug prior to the registration
• Recent stroke or myocardial infarction within 1 month
• Other inappropriate conditions judged by each investigator

HIV, human immunodeficiency virus.

Methods

Objectives

This study was designed to enroll 30,000 patients with hypertension throughout Japan, and the patients were treated with losartan on an open-label basis at a daily dose at 25-50 mg with standard clinical management for a maximum of 5 years. The aims of this study were to investigate the efficacy and safety profile of losartan during actual clinical use in the 5-year post-marketing period, the incidence of cardiovascular events and mortality, the value of lifestyle modification as antihypertensive therapy, and the relationship between the clinic BP and the home BP values in Japanese hypertensive patients primarily treated with losartan.

Patients Recruitment

The eligible patients were men or women ≥ 20 years of age who were diagnosed as having hypertension by their physicians and had not taken any antihypertensive agents within the previous 1 month. Patients who had previously taken losartan were excluded. The other exclusion criteria are shown in Table 1. Each patient was informed of the purpose and methods of the study, as well as the effects and possible risks of losartan therapy, the right to withdraw from the study at any time, and the measures for privacy protection before they were enrolled. Patients provided their verbal informed consent and then underwent a complete medical history review, physical examination, and laboratory evaluation.

Drug Treatment and Study Procedure

The patients were initially treated with losartan at a dose of 25-50 mg once daily (usually in the morning), which is the approved dosage in Japan. The dose was increased up to a dose of 100 mg once daily, if necessary. Addition of other antihypertensive agents was allowed from 3 months after the start of losartan treatment, if required. No restrictions were placed on the treatment of complications.

The enrolled patients were registered in a central study registry that included the following information at baseline:

Table 2. Patients' Characteristics at Baseline

	Men	Women	Total
Number of patients (n (%))	13,737	17,311	31,048
Age (years old)	60.0±12.0	64.3±11.8	62.4±12.1
SBP (mmHg)	164.4±17.0	165.9±17.4	165.3±17.3
DBP (mmHg)	96.2±11.6	92.8±11.6	94.3±11.7
BMI (kg/m ²)	24.3±3.3	23.9±3.8	24.1±3.6
Alcohol drinkers (n (%))	9,147 (66.6)	2,674 (15.4)	11,821 (38.1)
Current smokers (n (%))	6,085 (44.3)	1,664 (9.6)	7,749 (25.0)
Complications			
Hyperlipidemia (n (%))	4,935 (35.9)	7,005 (40.5)	11,940 (38.5)
Diabetes mellitus (n (%))	2,170 (15.8)	1,883 (10.9)	4,053 (13.1)
Cardiovascular disease (n (%))	1,097 (8.0)	1,400 (8.1)	2,497 (8.0)
Cerebrovascular disease (n (%))	616 (4.5)	745 (4.3)	1,361 (4.4)
Hepatic disease (n (%))	1,901 (13.8)	1,069 (6.2)	2,970 (9.6)
Renal disease (n (%))	509 (3.7)	496 (2.9)	1,005 (3.2)
ECG abnormality (n (%))	2,119 (15.4)	2,234 (12.9)	4,353 (14.0)
Concomitant drugs			
Lipid-lowering drugs (n (%))	2,632 (19.2)	5,033 (29.1)	7,665 (24.7)
Antidiabetics (n (%))	1,452 (10.6)	1,336 (7.7)	2,788 (9.0)
UA lowering drugs (n (%))	1,371 (10.0)	239 (1.4)	1,610 (5.2)
Aspirin or antiplatelets (n (%))	1,102 (8.0)	1,188 (6.9)	2,290 (7.4)

Mean±SD. SBP, systolic blood pressure; DBP, diastolic blood pressure; BMI, body mass index; UA, uric acid.

demographic data, physical data (height and body weight), history of hypertension, and use of antihypertensive drugs; BP values and pulse rate; complications and medical history (renal disease, hepatic disease, cerebrovascular disease, coronary heart disease, endocrine/metabolic disease, and other diseases); laboratory test results (complete blood count, biochemistry tests, and urinalysis); lifestyle modification if performed (physical exercise, restriction of alcohol consumption or salt intake, ceasing smoking, weight loss, etc.); and electrocardiograph findings.

The following patient information was recorded in the worksheets and collected every year after the start of losartan-based antihypertensive treatment: adverse events, clinic BP values, pulse rate, heart rate, weight, daily dose of losartan, concomitant drugs, laboratory tests, and ECG (if performed).

The clinic BP was measured by the routine method at each institution. At each time of measurement, one clinic BP value was reported at the discretion of the physician. The clinic BP data measured at a maximum of 3 different visits prior to starting losartan therapy was used for calculation of the mean baseline clinic BP. After starting losartan therapy, the clinic BP value was measured every 3 months. The clinic BP data thus obtained were used for analysis of the clinic BP values during treatment.

The home BP was measured during the study by patients who voluntarily agreed to monitor their BP themselves. Home BP was measured with an electronic automated sphygmomanometer based on the cuff-oscillometric principal (HEM-740A; Omron Healthcare Co., Ltd., Kyoto, Japan).

Patients who had already been using another device and insisted on continuing its use were permitted to do so. Patients were asked to measure the home BP at rest in the sitting position once every morning just after waking and urinating, and before medication. Home BP was measured once at one opportunity of measurement. If the patient measured home BP twice or more at one opportunity, the first measured value was reported. Home BP values obtained prior to the start of losartan therapy were used to calculate the mean baseline home BP. As a rule, morning home BP values measured each month, usually on the day of attending hospital, were used for analysis of the mean home BP value during treatment.

Standard laboratory tests (including ECG recording) were performed with the routine methods used at each institution, so standardization of measuring methods and reference values was not carried out. A maximum of 2 results of standard laboratory tests measured prior to losartan therapy were used to calculate the baseline values. After the start of losartan therapy, standard laboratory tests were performed every 6 months.

To assess the complications and the medical history, physicians judged the existence of diseases indicated in the registration form prior to the start of the study at their discretion.

In addition, the patients who were receiving drug treatment for hyperlipidemia or diabetes mellitus and met the definition of either disease indicated in the relevant guidelines were defined as having hyperlipidemia or diabetes.

All adverse events were recorded by the investigators and were classified as definitely related, possibly related, or defi-

Table 3. Distribution of Age and WHO Hypertension Grade, and Mean Blood Pressure at Baseline

	Men (N=12,698)		Women (N=16,250)		Total (N=28,948)	
	n (%)	SBP/DBP (mmHg)	n (%)	SBP/DBP (mmHg)	n (%)	SBP/DBP (mmHg)
Age (years old)						
20-39	556 (4.4)	160.1±16.0/102.2±11.3	261 (1.6)	163.5±16.7/101.9±10.6	817 (2.8)	161.2±16.3/102.1±11.1
40-59	5,518 (43.5)	163.5±16.7/100.1±10.4	5,494 (33.8)	167.1±18.2/97.9±10.5	11,012 (38.0)	165.3±17.6/99.0±10.5
60-79	6,029 (47.5)	165.5±17.0/92.9±11.1	8,963 (55.2)	165.3±16.8/90.6±10.7	14,992 (51.8)	165.4±16.9/91.5±10.9
≥80	595 (4.7)	165.4±18.9/87.0±11.5	1,532 (9.4)	166.0±18.4/85.7±12.1	2,127 (7.4)	165.9±18.5/86.0±12.0
BP classification						
Optimal to High-normal	349 (2.7)	130.2±8.4/77.0±8.2	499 (3.1)	130.3±8.3/75.7±8.6	848 (2.9)	130.3±8.3/76.2±8.5
Grade 1	3,282 (25.9)	149.9±6.7/89.1±7.5	4,349 (26.8)	150.6±6.1/86.7±8.1	7,631 (26.4)	150.3±6.4/87.7±7.9
Grade 2	6,027 (47.5)	164.2±8.6/95.9±8.5	7,619 (46.9)	165.9±7.2/92.8±9.1	13,646 (47.1)	165.2±7.9/94.1±8.9
Grade 3	3,040 (23.9)	184.3±15.7/106.6±12.0	3,783 (23.3)	188.3±14.5/102.0±12.6	6,823 (23.6)	186.5±15.2/104.1±12.5

Mean±SD. BP, blood pressure; SBP, systolic blood pressure; DBP, diastolic blood pressure.

nately unrelated to losartan, or as unknown. All losartan-related adverse events were pooled and classified as adverse drug reactions (ADRs).

Endpoint Evaluation

The primary endpoint of the study was a composite of cardiovascular events, including fatal or non-fatal stroke (new occurrence or recurrence of cerebral hemorrhage, cerebral infarction, or subarachnoid hemorrhage diagnosed on the basis of typical clinical symptoms persisting for more than 24 h and/or computerized tomography/magnetic resonance imaging findings), transient ischemic attack defined as a focal neurological deficit presumed to be vascular in origin persisting for less than 24 h, fatal or non-fatal myocardial infarction (new occurrence or recurrence) diagnosed on the basis of typical clinical symptoms, ECG changes and elevation of cardiac enzymes, or sudden cardiac death. In addition, the independent event classification committee reviewed adjudicated endpoint events on the basis of all available information documented in the case report form by the physicians.

Statistical Considerations

Determination of the Sample Size

When performing life-table analysis combined with the log-rank test, a 30% difference in the incidence of the primary endpoint (stroke, transient ischemic attack, acute myocardial infarction, or sudden cardiac death) was assumed between a subgroup of patients that represented 60% of the total population with higher BP and the remaining patients with lower BP. The incidence of stroke, myocardial infarction and sudden cardiac death in the Japanese population is 6.8/1,000 patient-years in men and 4.8/1,000 patient-years in women according to the Hisayama study (18), and the mean follow-up period for the J-HEALTH was 2.7 years. Thus, a total of 28,000 patients were required to detect the assumed between-group difference with a 90% power at $\alpha=0.05$ (2-sided). Therefore,

the target sample size was set at 30,000 patients.

Statistical Analysis

For the present interim analysis, variables were compared using the *t*-test, the χ^2 test, or analysis of variance (ANOVA). Results were expressed as the mean±SD, and differences were considered statistically significant at $p<0.05$.

Statistical analysis of the overall results was based on survival analysis. Subgroups were classified by the BP values at baseline or during treatment. Differences between subgroups were assessed by the log-rank test or the χ^2 test. Relationships between the endpoints and the BP values, as well as prognostic factors, were assessed by using the Cox proportional hazards model with adjustment for gender, age, alcohol drinker, current smoker, coexisting of cardiovascular disease, cerebrovascular disease, diabetes mellitus, and hyperlipidemia. For analysis of safety data, the number of ADRs, drug-related ADRs and other ADRs were calculated. For efficacy analysis, the antihypertensive effect of losartan with respect to both clinic BP and home BP values was assessed, and subgroup analyses were performed as described for the safety analysis. Comparison of safety and efficacy among the subgroups was performed with the χ^2 test, *t*-test, or ANOVA. Results were expressed as the mean±SD and a 2-sided $p<0.05$ was considered statistically significant. Statistical analysis was conducted with the SAS package (version 8.02; SAS Institute Inc., Cary, USA).

Organization

The organization and the members of the committees of the J-HEALTH study are given in the Appendix. These committees were responsible for performing the study or analyzing the data. The Monitoring Committee determined the validity of continuing the study based on the safety and effectiveness of losartan therapy from an ethical point of view. The Event Assessment Committee reviewed the events of cerebrovascular disease and coronary heart disease reported during the

Table 4. Lipid Profiles at Baseline

	Men (N=13,737)	Women (N=17,311)	Total (N=31,048)
Hyperlipidemia (n)	4,935	7,005	11,940
TC (mg/dL)	216.4±37.2	230.2±35.3	224.4±36.7
HDL-C (mg/dL)	51.5±15.5	59.2±16.3	56.0±16.4
TG (mg/dL)	214.2±142.9	159.9±96.3	182.8±121.3
Without hyperlipidemia (n)	8,802	10,306	19,108
TC (mg/dL)	189.3±29.0	199.5±29.6	194.7±29.8
HDL-C (mg/dL)	56.5±14.5	61.1±15.0	58.9±14.9
TG (mg/dL)	126.8±89.1	110.0±58.4	117.8±74.7

Mean±SD. TC, total cholesterol; HDL-C, high density lipoprotein-cholesterol; TG, triglycerides.

study. The Safety Assessment Committee assessed the causal relationship between the ADRs that are reported and the drugs that are administered during the study. The Medical Expert Advisory and Publication Committee was responsible for reviewing the results and writing the paper.

Results

Baseline Patients' Characteristics

Between June 2000 and December 2001, patients were screened in all 47 prefectures throughout Japan. The number of patients enrolled in this study per prefecture ranged from 165 in Okinawa to 2,667 in Tokyo. The distribution of patient enrollment was similar to the recent Japanese population statistics (19), and there were no major regional differences of BP values among the prefectures (data not shown).

A total of 31,515 patients were screened at 3,755 institutions by 4,149 investigators. Among them, 31,048 patients were enrolled in this study and 467 patients were excluded according to the exclusion criteria shown in Table 1 or withdrew their consent before actual enrollment. The baseline characteristics of the 31,048 enrolled patients (13,737 men [44.2%] and 17,311 women [55.8%]) are summarized in Table 2. The mean age of the patients was 62.4±12.1 years and the mean clinic systolic/diastolic BP (SBP/DBP) values were 165.3±17.3/94.3±11.7 mmHg.

Concomitant medications and complications are also listed in Table 2. All complications and ECG abnormality were diagnosed by the study investigators. The prevalences of hyperlipidemia, diabetes mellitus, cardiovascular disease, cerebrovascular disease, and ECG abnormality were 38.5%, 13.1%, 8.0%, 4.4%, and 14.0% respectively. Subjects taking anti-diabetic agents or lipid-lowering drugs were defined as having diabetes or hyperlipidemia, respectively.

Table 3 shows distributions of age groups and grade in the World Health Organization (WHO), and the mean BP values at baseline. Young patients (20–39 years) accounted for 2.8%, middle-aged patients (40–59 years) accounted for 38.0%, and elderly patients (60–79 years) made up 51.8% of the total patients. It is worth noting that there were 2,127

(7.4%) very elderly patients (≥80 years). The age distribution was generally similar between men and women. Then we analyzed the BP values of each age group. As shown in Table 3, the SBP values increased with age, but the difference was relatively small. On the other hand, the DBP values decreased markedly with age. Grade 2 hypertensive patients (based on the WHO classification) were most frequent in our cohort (n=13,646, 47.1%), while the numbers of grade 1 and 3 patients were almost equal (n=7,631, 26.4% vs. n=6,823, 23.6%, respectively).

The mean total cholesterol (TC) level of all patients was 209.6 mg/dL, while the mean TC levels of hyperlipidemic patients (n=11,940) and non-hyperlipidemic patients (n=19,108) were 224.4 mg/dL and 194.7 mg/dL, respectively. Details of the lipid profile are given in Table 4.

A total of 11,135 patients agreed to measure their BP values at home. Although data were limited at the time of enrollment (n=9,182), the scatter plot (Fig. 1) demonstrates the relationship between clinic BP and home BP for both the SBP and DBP values. The Pearson's correlation coefficients were 0.62 and 0.69, respectively. "White-coat hypertension" (WCHT) was defined by the following criteria: clinic SBP ≥140 or clinic DBP ≥90 mmHg and home SBP <135 and home DBP <85 mmHg. Based on these criteria, 4.2% of our patient cohort had so-called WCHT.

Discussion

The JSH 2004 have been published and updated periodically based on mainly Western epidemiological and clinical results (5). Although many large-scale investigational studies have been conducted worldwide to explore the management of hypertension (3, 4), it is difficult to determine which studies are best applicable to each individual case in clinical practice. Therefore, it is very important to clarify the characteristics, clinical effects, and safety profiles of various drugs in clinical practice. Many studies with Japanese hypertensives have been conducted, but these have usually employed small cohorts in rural areas. Practical information from large-scale investigational studies in clinical practice is limited in Japan (20). The J-HEALTH is a large-scale (30,000 patients)




Fig. 1. Scattered diagrams of clinic and home blood pressure at baseline ($n=9,182$). Each patient's clinic and home blood pressure was plotted for SBP (A) and DBP (B) with regression equations. SBP, systolic blood pressure; DBP, diastolic blood pressure.

nationwide multicenter observational study, which may provide us valuable epidemiological information on Japanese patients with hypertension.

The distribution of the hypertensive patients enrolled in the J-HEALTH study was similar to that of the recent Japanese population statistics (19). Thus, the patients in this study can be regarded as being representative of the overall Japanese population. Geographical differences in the prevalence of hypertension have previously been noted in Japan (21), with a higher incidence in the primarily rural northern part of Japan and a lower rate in the Western region (5). One of the reasons for the higher BP values in rural northern Japan is the high salt intake of the local population. However, no major regional differences of the mean clinic BP values were observed among the prefectures in the J-HEALTH study. This may reflect recent lifestyle changes and/or the widespread acceptance of antihypertensive therapy in Japan.

It is well known that vascular mortality increases with age, but the contribution of BP values to vascular mortality differs between age groups. The Prospective Studies Collaboration has published a meta-analysis of individual data for one million adults in 61 prospective observational studies of BP values and mortality (22). Although the proportional difference in the risk of vascular death is associated with an absolute difference in BP values, the proportional difference in vascular mortality is only about half as large at 80–89 years compared with that at 40–49 years. Therefore, the age of the subjects is an essential factor when analyzing study endpoints. The mean age of the J-HEALTH cohort is 62.4 ± 12.1 years. On the other hand, the mean age was 67 years in the LIFE study (7). This age difference between randomized trials and actual clinical practice should be taken into consideration when applying the results of randomized trials.

The risks and benefits of treatment with antihypertensive agents are uncertain in patients older than 80 years (23). Gueyffier *et al.* suggested that antihypertensive treatment

could prevent stroke, major cardiovascular events, and heart failure, but not cardiovascular death based on a meta-analysis of data from 1,670 patients aged 80 years or older (24). The Hisayama study suggested that the Sixth Report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC-6) recommendations were not applicable to elderly Japanese persons over 80 years of age (25). Many very elderly hypertensive patients (≥ 80 years) were enrolled (595 men and 1,532 women) in the J-HEALTH study. Such a large number of elderly Japanese has not been studied before, so the results of the J-HEALTH study should be informative for this age group.

Of course, the BP is the most important characteristic of the patients in this study. The mean clinic SBP/DBP values of the J-HEALTH cohort were 165.3/94.3 mmHg. In contrast, the mean SBP/DBP values were 174.4/97.8 mmHg in the LIFE study. Based on their age and mean BP values, the J-HEALTH cohort is younger and has milder hypertension compared with the subjects of the LIFE study (7). Therefore, the J-HEALTH study may be able to assess the beneficial effects of ARB-based treatment for relatively low-risk hypertensive patients, who are the most common type encountered in clinical practice in Japan.

Not only the mean BP value itself, but also the grade of hypertension, is an important factor to be taken into consideration when evaluating a large-scale study. Grade 2 hypertension is the most frequent type in the J-HEALTH study population (47.1%). In their sub-analysis of the Hisayama study, Arima *et al.* excluded treated hypertensive patients and followed up 588 cardiovascular disease-free residents who were at least 60 years of age for about 30 years (from 1961 to 1993) to evaluate their cardiovascular risk. Among these patients, BP grade 1, 2, and 3 accounted for 27.2%, 18.6%, and 14.1%, respectively (they included normal BP and high normal BP subjects). Since the Hisayama study was an observational investigation of the general population, the hyperten-

sion stage distribution of the Hisayama population is different from that of the J-HEALTH cohort (25).

Complications represent another important background factor. The prevalences of ECG abnormality, cardiovascular disease, and cerebrovascular disease are 14.0%, 8.0%, and 4.4%, respectively, in the J-HEALTH cohort. These rates are lower than those in the LIFE study, again indicating that the J-HEALTH enrolled healthier subjects than the LIFE (7). The J-HEALTH cohort includes a high percentage of hyperlipidemic patients (38.5%). However, as shown in Table 4, the mean TC level of these hyperlipidemic patients is not extremely high, possibly because 24.7% of all the patients were taking lipid-lowering drugs (Table 3).

It has been emphasized that home BP values measured by ambulatory blood pressure (ABP) monitoring or self-measurement can be an important tool for the optimal management of hypertension with respect to cardiovascular risk (26). In the *Pressioni Arteriose Monitorate e Loro Associazioni* (PAMELA) study of 2,051 subjects who were representative of the general population, the clinic BP, home BP, and 24-h ABP values were measured. This study demonstrated that the risk of death increased more with a given increase in home BP or ABP than clinic BP values (27). Den Hond *et al.* investigated the diagnostic values of self-measured BP vs. ABP, and concluded that the specificity and sensitivity of ABP values for detecting WCHT were better than those of home BP values (28). While ABP values have better prognostic accuracy, the American Society of Hypertension Ad Hoc Panel recommends the use of home BP values for screening. Self-measurement of the BP is easy to repeat and is useful for patients to assess their own control (29). Hozawa *et al.* investigated the BP measured by home, ambulatory, and conventional methods in 1,174 Japanese subjects (150 with untreated hypertension, 399 with treated hypertension, and 625 normotensives). They also concluded that it was useful to measure the non-clinic BP values (30). We therefore determined the distribution and relation between clinic BP and home BP values at enrollment (Fig. 1).

WCHT is diagnosed by comparing the clinic BP and non-clinic BP values, and whether it causes target organ damage and cardiovascular events is one of the most important issues in the treatment of hypertension (31). The Ohasama study examined the prognostic significance of WCHT based on ABP monitoring, and concluded that the predictive power of the ABP values for subsequent mortality was stronger than that of the clinic BP values (28). The prevalence of WCHT is 4.2% in the J-HEALTH cohort, but the reported prevalence of WCHT in other studies varies widely because of differences in the definition of WCHT, method of BP measurement (self-measurement vs. ABP monitoring), and characteristics of the study population (untreated hypertension vs. treated hypertension). Masked hypertension (MHT) is also a topic of interest for antihypertensive therapy. The Self-Measurement of Blood Pressure at Home in the Elderly: Assessment and Follow-up (SHEAF) study demonstrated that about 9% of treated

elderly patients had MHT and that the cardiovascular risk associated with MHT is significantly high (32).

A total of 9,182 patients measured their home BP values at the baseline in the J-HEALTH cohort. Therefore, we will use their data to identify WCHT and MHT and investigate the effect of home BP data on cardiovascular events. The home BP data obtained by self-measurement will display the time course effect of antihypertensive management and provide prognostic information for the hypertensive population.

The J-HEALTH study began in June 2000, and follow-up was completed in December 2005. The J-HEALTH study will clarify the long-term antihypertensive efficacy and safety of losartan-based therapy, and assess its preventive effect on hypertension-related diseases. It may provide new insights into therapeutic strategies for Japanese hypertensive patients.

Appendix

J-HEALTH Committees

Monitoring Committee: Takenori Yamaguchi (Chair), Tanenao Eto, Toshiharu Furukawa, Katsumi Yoshida.

Event Assessment Committee: Hiroaki Naritomi (Chair), Yoichiro Hashimoto, Uichi Ikeda, Mitsuaki Isobe, Toshio Kushiro, Ken Nagata, Kazuyuki Shimada, Takenori Yamawaki.

Safety Assessment Committee: Kendo Kiyosawa (Chair), Hiroshi Hirose, Sadayoshi Ito, Akinori Kasahara, Hiroshi Kawabe, Genjiro Kimura, Hirofumi Makino, Mitsuhiro Moriyama, Ikuo Saito, Hironichi Suzuki, Eiji Tanaka.

Medical Expert Advisory and Publication Committee: Hiroaki Naritomi (Chair), Toshiro Fujita, Sadayoshi Ito, Toshio Ogihara, Kazuyuki Shimada, Kazuaki Shimamoto, Heizo Tanaka, Nobuo Yoshiike.

The Administrative Office: The Post-Marketing Surveillance Department of Banyu Pharmaceutical Co., Ltd. (Tokyo, Japan).

References

1. World Health Organization, International Society of Hypertension Writing Group: 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens* 2003; 21: 1983–1992.
2. Fujiwara T, Nishimura T, Ohkubo T, Imai Y: Rationale and design of HOMED-BP Study: Hypertension Objective treatment based on Measurement by Electrical Devices of Blood Pressure Study. *Blood Press Monit* 2002; 7: 77–82.
3. The ALLHAT Officers and Coordination for the ALLHAT Collaborative Research Group: Major outcomes in high-risk hypertensive patients randomized to angiotensin-converting enzyme inhibitor or calcium channel blocker vs diuretic: the Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT). *JAMA* 2002; 288: 2981–2997.
4. Neal B, MacMahon S, Chapman N: Effects of ACE inhibitors, calcium antagonists, and other blood pressure-lowering drugs: results of prospectively designed overviews of randomised trials. *Lancet* 2000; 356: 1955–1964.

5. Japanese Society of Hypertension: Japanese Society of Hypertension Guidelines for the Management of Hypertension (JSH 2004). *Hypertens Res* 2006; 29 (Suppl): S1–S105.
6. Chobanian AV, Bakris GL, Black HR, *et al*: Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension* 2003; 42: 1206–1252.
7. Dahlöf B, Devereux RB, Kjeldsen SE, *et al*: Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002; 359: 995–1003.
8. Brenner BM, Cooper ME, Zeeuw D, *et al*: Effects of losartan on renal and cardiovascular outcomes in patients with type 2 diabetes and nephropathy. *N Engl J Med* 2001; 345: 861–869.
9. Lindholm LH, Ibsen H, Dahlöf B, *et al*: Cardiovascular morbidity and mortality in patients with diabetes in the Losartan Intervention for Endpoint reduction in Hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002; 359: 1004–1010.
10. Dickstein K, Kjekshus J: Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomised trial. *Lancet* 2002; 360: 752–760.
11. Pitt B, Poole-Wilson PA, Segal R, *et al*: Effects of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomised trial—the Losartan Heart Failure Survival Study ELITE II. *Lancet* 2000; 355: 1582–1587.
12. Ueshima H: Changes in dietary habit, cardiovascular risk factors and mortality in Japan. *Acta Cardiol* 1990; 45: 311–327.
13. Khor GL: Cardiovascular epidemiology in the Asia-Pacific region. *Asia Pac J Clin Nutr* 2001; 10: 76–80.
14. Tsuji I, Imai Y, Nagai K, *et al*: Proposal of reference values for home blood pressure measurement: prognostic criteria based on a prospective observation of the general population in Ohasama, Japan. *Am J Hypertens* 1997; 10: 409–418.
15. Tachibana R, Tabara Y, Kondo I, *et al*: Home blood pressure is a better predictor of carotid atherosclerosis than office blood pressure in community-dwelling subjects. *Hypertens Res* 2004; 27: 633–639.
16. Ohkubo T, Obara T, Funahashi J, *et al*: Control of blood pressure as measured at home and office, and comparison with physicians' assessment of control among treated hypertensive patients in Japan: first report of the Japan Home versus Office Blood Pressure Measurement Evaluation (J-HOME) study. *Hypertens Res* 2004; 27: 755–763.
17. Thijs L, Staessen JA, Celis H, *et al*: Reference values for self-recorded blood pressure: a meta-analysis of summary data. *Arch Intern Med* 1998; 158: 481–488.
18. Kubo M, Kiyohara Y, Kato I, *et al*: Trends in the incidence, mortality, and survival rate of cardiovascular disease in a Japanese community: the Hisayama study. *Stroke* 2003; 34: 2349–2354.
19. Current Population Estimates as of October 1, 2004. Tokyo. Ministry of Internal Affairs and Communications, Statistics Bureau, 2004.
20. Iida M, Ueda K, Okayama A, *et al*: Impact of elevated blood pressure on mortality from all causes, cardiovascular diseases, heart disease and stroke among Japanese: 14 year follow-up of randomly selected population from Japanese—Nippon data 80. *J Hum Hypertens* 2003; 17: 851–857.
21. Konishi M, Iso H, Iida M, *et al*: Trends for coronary heart disease and its risk factors in Japan: epidemiologic and pathologic studies. *Jpn Circ J* 1990; 54: 428–435.
22. Lewington S, Clarke R, Oizilbash N, Peto R, Collins R: Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies. *Lancet* 2002; 360: 1903–1913.
23. Bulpitt CJ, Beckett NS, Cooke J, *et al*: Results of the pilot study for the Hypertension in the Very Elderly Trial. *J Hypertens* 2003; 21: 2409–2417.
24. Gueyffier F, Bulpitt C, Boissel JP, *et al*: Antihypertensive drugs in very old people: meta-analysis of randomized controlled trials. *Lancet* 1999; 353: 793–796.
25. Arima H, Tanizaki Y, Kiyohara Y, *et al*: Validity of the JNC VI recommendations for the management of hypertension in a general population of Japanese elderly: the Hisayama study. *Arch Intern Med* 2003; 163: 361–366.
26. Celis H, Den Hond E, Staessen JA: Self-measurement of blood pressure at home in the management of hypertension. *Clin Med Res* 2005; 3: 19–26.
27. Mancia G, Facchetti R, Bombelli M, Grassi G, Sega R: Long-term risk of mortality associated with selective and combined elevation in office, home, and ambulatory blood pressure. *Hypertension* 2006; 47: 846–853.
28. Den Hond E, Celis H, Vandenhoven G, O'Brien E, Staessen JA: Determinants of white-coat syndrome assessed by ambulatory blood pressure of self-measured home blood pressure. *Blood Press Monit* 2003; 8: 37–40.
29. Den Hond E, Celis H, Fagard R, *et al*: Self-measured versus ambulatory blood pressure in the diagnosis of hypertension. *J Hypertens* 2003; 21: 717–722.
30. Hozawa A, Ohkubo T, Kikuya M, *et al*: Blood pressure control assessed by home, ambulatory and conventional blood pressure measurements in the Japanese general population: the Ohasama Study. *Hypertens Res* 2002; 25: 57–63.
31. Celis H, Fagard RH: White-coat hypertension: a clinical review. *Eur J Intern Med* 2004; 15: 348–357.
32. Bobrie G, Chatellier G, Genes N, *et al*: Cardiovascular prognosis of “masked hypertension” detected by blood pressure self-measurement in elderly treated hypertensive patients. *JAMA* 2004; 291: 1342–1349.

Extremely Early Computed Tomography Signs in Hyperacute Ischemic Stroke as a Predictor of Parenchymal Hematoma

Shuhei Okazaki Hiroshi Moriwaki Kazuo Minematsu Hiroaki Naritomi

Department of Cerebrovascular Medicine, National Cardiovascular Center, Osaka, Japan

© S. Karger AG, Basel
PROOF Copy
for personal
use only

ANY DISTRIBUTION OF THIS
ARTICLE WITHOUT WRITTEN
CONSENT FROM S. KARGER
AG, BASEL IS A VIOLATION
OF THE COPYRIGHT.

Key Words

Acute stroke diagnosis · Computed tomography · Hemorrhagic transformation · Intracerebral hemorrhage · Ischemic stroke

Abstract

Background: In acute ischemic stroke, thrombolysis with intravenous recombinant tissue plasminogen activator is more effective when given within 90 min of onset compared to that given later than 90 min. However, the significance of early CT signs (ECTs) during such early periods has not yet been fully clarified. We investigated the usefulness of ECTs within 90 min for predicting parenchymal hematoma (PH) in patients without thrombolysis. **Methods:** We evaluated 212 consecutive patients with initial ischemic stroke in the anterior cerebral circulation who underwent the first CT within 6 h of onset. The patients were divided into 3 groups according to the interval from onset to CT: within 90 min (group A, n = 90), 91–180 min (group B, n = 76) and 181–360 min (group C, n = 46). Patients who had received thrombolytic therapy were excluded. ECTs were evaluated according to the Alberta Stroke Program Early CT Score (ASPECTS). The relationships between ECTs and the subsequent development of PH were compared among the groups. **Results:** In patients with ASPECTS values between 0 and 7, PH was developed more frequently in group A (35%) than in groups B (14%) or C (15%) (group A vs. B: p = 0.036, group A vs. C: p = 0.094). In group A, atrial fibrillation, elevated pretreatment

blood pressure and ASPECTS ≤ 7 were independent predictors of PH. **Conclusions:** The manifestation of ECTs as represented by ASPECTS ≤ 7 within 90 min after stroke appears to indicate a high risk of PH.

Copyright © 2008 S. Karger AG, Basel

Introduction

In hyperacute ischemic stroke, thrombolysis with intravenous recombinant tissue plasminogen activator (rt-PA) was found to improve the 90-day functional outcome when given within 3 h of symptom onset in the National Institute of Neurological Disorders and Stroke (NINDS) trial [1]. On the other hand, the frequency of intracranial hemorrhage (ICH) was shown to increase significantly in stroke patients treated with thrombolysis [1–3]. Thus, the pretreatment predictors of ICH are currently one of the most important issues in clinical practice. At present, computed tomography (CT) is the most widely available and convenient technique for evaluating stroke, and the prediction of ICH using CT would be exceedingly valuable. The European Cooperative Acute Stroke Study (ECASS) investigators suggested that baseline early CT signs (ECTs), according to the so-called one third rule, could be predictive of ICH [2]. However, there was no reference to ECTs in the NINDS trial [1]. The subanalysis of NINDS data could not confirm the relationship between ECTs and ICH either, although they found such a ten-

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2008 S. Karger AG, Basel
1015-9770/08/0000-0000\$24.50/0

Accessible online at:
www.karger.com/ced

Shuhei Okazaki, MD
Department of Cerebrovascular Medicine, National Cardiovascular Center
5-7-1 Fujishirodai, Suita
Osaka 565-8565 (Japan)
Tel. +81 6 6833 5012, Fax +81 6 6872 7486, E-Mail s-okazaki@umin.ac.jp

dency that symptomatic hemorrhages develop more commonly in patients whose ECTs extend over more than one third of the middle cerebral artery territory [4]. In addition, a recent meta-analysis of 6 major intravenous thrombolytic therapy trials suggests that the benefit of rt-PA is greater if started within 90 min compared to that later than 90 min [5]. However, the significance of ECTs in such very early periods has not yet been fully clarified.

Recently, a new scoring system was developed to better locate and semiquantify the early ischemic changes on CT: the Alberta Stroke Program Early CT Score (ASPECTS) [3]. This semiquantitative scoring method improved the interobserver agreement ratio [6] and facilitated the detection of earlier and more subtle changes [7]. We hypothesized that the time from onset to the appearance of ECTs would correlate with the severity of cerebral ischemia and the frequency of subsequent parenchymal hematoma (PH). We investigated the usefulness of ECT evaluation using ASPECTS within 90 min for predicting the risk of spontaneous PH in the following acute or subacute phase in patients without thrombolysis.

Patients and Methods

Patients

Among 1,679 patients with ischemic stroke admitted to the stroke care unit of the National Cardiovascular Center, Japan, between January 1998 and June 2005, we retrospectively evaluated 655 consecutive patients who underwent the first CT within 6 h after onset. We excluded patients with a history of symptomatic stroke or intracranial surgery. We also excluded patients with unclear onset including stroke at awakening and those with infarction limited to the posterior circulation alone. Thirty-five patients who had received thrombolysis (intravenous or intra-arterial) were also excluded. In Japan, the use of intravenous rt-PA for acute ischemic stroke was approved in October 2005, by which time the present study had been completed. Therefore, during the study period, thrombolytic therapy was performed only in patients who were enrolled in other clinical trials. Consequently, 212 consecutive patients with initial ischemic stroke in the anterior cerebral circulation were enrolled in this study. The 212 patients were divided into 3 groups according to the interval from onset to the first CT: within 90 min (group A, $n = 90$), 91–180 min (group B, $n = 76$) and 181–360 min (group C, $n = 46$). Anticoagulants and/or antiplatelet agents were administered to the patients in case the attending physicians considered these therapies were sufficiently safe and valuable.

CT Studies

All patients underwent noncontrast CT scans using an X-Vigor (Toshiba, Tokyo) on arrival. The scanning parameters were nonhelical, 120 kV, 170 mA, 10-mm collimation, matrix size of 512×512 and 2- or 3-second scan time. Routine photography was performed at window level and a width of 30 and 70–80

Hounsfield units (HU), respectively. Follow-up CTs were performed at 24 h and 7–14 days after stroke, as well as at the discretion of treating neurologists and reviewed in this study when performed within 14 days. All baseline CTs and follow-up CTs were retrospectively evaluated by 2 expert neurologists (S.O. and H.M.) with knowledge of the side of the affected hemisphere but blinded to treatment assignment, stroke severity or other clinical details. ECTs were scored according to ASPECTS [3, 6]. ASPECTS was assessed by scoring 10 regions (caudate nucleus, internal capsule, lentiform nucleus, insula and 6 cortical regions) systematically on the CT scan. A score of 1 was assigned for normal and 0 for a region showing ECTs. A normal CT scan has ASPECTS values of 10 points. Score 0 indicates diffuse ischemia throughout the territory of the middle cerebral artery. The ASPECTS study group suggested that baseline ASPECTS values in 2 categories (≤ 7 and > 7) discriminated poor and good functional outcome, or high and low risk of ICH [3]. Therefore, we compared the incidence of ICH between a group with ASPECTS ≤ 7 and one with ASPECTS > 7 in each period.

ECTs were defined as X-ray hypoattenuation, loss of the gray-white boundary or effacement of cortical sulci [3, 6]. ICH was classified as hemorrhagic infarction and PH; hemorrhagic infarction was defined as petechial infarction without space-occupying effect, and PH was defined as hemorrhage with mass effect. PH was subdivided into PH1, as blood clots in $\leq 30\%$ of the infarcted area with some slight space-occupying effect, and PH2, as blood clots in $> 30\%$ of the infarcted area with a substantial space-occupying effect, corresponding to ECASS II criteria [2]. In cases of disagreement in the findings, these 2 observers reviewed the CTs together and discussed the findings until a consensus was established.

Clinical Evaluation

Risk factors for ischemic stroke (i.e. age, sex, hypertension, diabetes mellitus, hyperlipidemia, atrial fibrillation and smoking) were evaluated in all patients. According to the exclusion criteria of NINDS [1], we defined 'elevated pretreatment blood pressure' as systolic blood pressure > 185 mm Hg or diastolic blood pressure > 110 mm Hg. Because the second analysis of NINDS showed that patients > 75 years of age or with National Institutes of Health Stroke Scale (NIHSS) scores > 20 were associated with poor outcome and high risk of ICH [9], we decided on the cutoff points of age > 75 years and NIHSS score > 20 on univariate analysis. The trial of ORG10172 in Acute Stroke Treatment criteria [10] were used to classify stroke subtypes.

Data Analysis

We determined the predictive value of ASPECTS referring to the findings of follow-up CT obtained at 24 h as a standard. ECTs were judged 'true positive' if a definite infarct was correctly located on the follow-up CT. If it was absent on the second CT, the reading was 'false positive'. The readings were judged 'true negative' if both baseline and follow-up CT did not reveal any infarcted changes. The readings were defined as 'false negative' if ECTs were not detected on the baseline CT and a definite infarct was visible on the follow-up CT. On the basis of these definitions, we assessed the predictive value in each of the 10 regions in each patient, then summed the results all together and calculated the sensitivity and specificity.

Table 1. Baseline characteristics of the 3 groups classified by time from stroke onset to initial CT

Variables	Group A 0–90 min)	Group B 91–180 min)	Group C 181–360 min)	p value
Patients	90	76	46	
Age, years	71 ± 11	70 ± 11	70 ± 12	0.783
Male	61 (68)	56 (74)	31 (67)	0.655
Hypertension	64 (71)	54 (71)	31 (67)	0.889
Diabetes mellitus	21 (23)	22 (29)	15 (33)	0.480
Hyperlipidemia	27 (30)	26 (34)	17 (37)	0.690
Atrial fibrillation	45 (50)	37 (49)	22 (48)	0.968
Smoking	25 (28)	23 (30)	12 (26)	0.588
Pretreatment, mm Hg				
Systolic blood pressure	162 ± 26	160 ± 26	171 ± 25	0.072
Diastolic blood pressure	87 ± 15	87 ± 13	90 ± 14	0.377
Elevated pretreatment blood pressure	20 (22)	12 (16)	9 (20)	0.578
NIHSS scores, median	12	9	6	0.097
1–5	20 (22)	24 (32)	21 (46)	
6–10	22 (24)	18 (24)	10 (22)	
11–15	14 (16)	7 (9)	5 (11)	
16–20	20 (22)	19 (25)	6 (13)	
>20	14 (16)	8 (11)	4 (9)	
ASPECTS, median	9	8	8	0.111
≤7	34 (37)	35 (46)	20 (44)	

Categorical values are expressed as numbers with percentages in parentheses, continuous variables as means ± SD. The χ^2 test was conducted for categorical variables, ANOVA for continuous variables and the Kruskal-Wallis test for scoring variables.

Statistical Analysis

Continuous data are expressed as means ± SD. For analysis of baseline characteristics, categorical variables were compared by χ^2 test, continuous variables by 1-way ANOVA and scoring variables by Kruskal-Wallis test. For univariate analysis of the predictors of PH, categorical variables were compared by χ^2 test, continuous variables by unpaired t test and scoring variables by Mann-Whitney U test. Multivariable analysis was undertaken using a logistic regression model. The results were considered significant at $p < 0.05$. Dr. SPSS II for Windows 11.0.1J was used for all calculations.

Results

Two hundred and twelve patients (148 men, 64 women) with ages ranging from 41 to 101 years (70 ± 11) were enrolled into this study. The stroke subtype was cardio-genic embolism in 118 patients, large-artery atherosclerosis in 29, small-vessel disease in 16 and other or unknown etiology in 49. As shown in table 1, the baseline characteristics were not significantly different between groups A, B and C.

ECTs were positive in 142 (67%) of the 212 patients. The predictive value for any ECTs with ASPECTS in

each group was as follows: sensitivity 60% and specificity 95% in group A, sensitivity 79% and specificity 91% in group B, and sensitivity 75% and specificity 90% in group C. PH was observed in 25 (12%) of the 212 patients. PH1 occurred in 14 patients and PH2 in 11. There were no differences in baseline CT findings and clinical characteristics between PH1 and PH2. Antithrombotic agents were used in 188 patients (89%) within 2 weeks from onset; 143 patients (68%) were treated with intravenous heparin and 79 patients (37%) with antiplatelet agents. Among the 188 patients with antithrombotic therapy, only 15 (8%) developed PH. On the other hand, 10 (42%) of the 24 patients without antithrombotic therapy developed PH. Figure 1 shows the proportion of PH by ASPECTS category in each group. When the ASPECTS values were between 0 and 7, PH was developed in 12 (35%) of the 33 patients in group A, showing a significantly higher frequency as compared with group B (14%) or group C (15%) (group A vs. group B: $p = 0.036$, group A vs. group C: $p = 0.094$).

Table 2 details the univariate analysis of predictors of PH within 90 min. Atrial fibrillation, elevated pretreatment blood pressure, NIHSS score and ASPECTS ≤ 7

Table 2. Univariate analysis of PH predictors in cases of CT examination within 90 min

Variables	PH (16 patients)	no PH (74 patients)	p value	Odds ratio
Age	73 ± 11	70 ± 11	0.33	
>75 years	7 (44)	24 (32)	0.40	1.62 [0.54, 4.87]
Male	12 (75)	49 (66)	0.57	1.53 [0.45, 5.24]
Hypertension	14 (88)	50 (68)	0.14	3.36 [0.71, 15.98]
Diabetes mellitus	6 (38)	15 (20)	0.19	2.36 [0.74, 7.53]
Hyperlipidemia	5 (31)	22 (30)	1.00	1.07 [0.33, 2.65]
Smoking	8 (50)	39 (53)	1.00	0.90 [0.30, 2.65]
Atrial fibrillation	13 (81)	32 (43)	0.011	5.69 [1.49, 21.66]
Elevated pretreatment blood pressure	7 (44)	13 (18)	0.042	3.65 [1.15, 11.58]
NIHSS scores, median	16.5	10	0.006	
>20	4 (25)	10 (14)	0.26	2.13 [0.57, 7.93]
ASPECTS, median	5	9	<0.001	
≤7	12 (75)	21 (28)	0.001	7.36 [2.14, 25.33]

Categorical values are expressed as numbers with percentages in parentheses, continuous variables as means ± SD. Figures in square brackets are 95% confidence limits. Elevated pretreatment blood pressure is defined as systolic blood pressure >185 mm Hg or diastolic blood pressure >110 mm Hg. Antiplatelet therapy includes aspirin, cilostazol, ticlopidine, argatroban and sodium ozagrel. Anticoagulant therapy includes heparin and warfarin.




Fig. 1. Incidence of PH after ischemic stroke by categories of time from stroke onset to CT, comparing ASPECTS ≤7 with ASPECTS >7. Values are numbers of patients with PH divided by the total patients in each group (n/n).

were associated with PH. Multivariate logistic regression analysis for prediction of PH is presented in table 3. ASPECTS ≤7, elevated pretreatment blood pressure and atrial fibrillation were independent predictors of PH within 90 min of onset.

Table 3. Predictors of PH in cases of CT examination within 90 min: multivariable logistic regression analysis

	p value	OR	95% CI
NIHSS (per 1 score increase)	0.308	1.05	0.95–1.17
ASPECTS ≤7	0.013	6.14	1.47–25.66
Elevated pretreatment blood pressure	0.022	5.45	1.28–23.25
Atrial fibrillation	0.045	5.02	1.03–24.37

Elevated pretreatment blood pressure is defined as systolic blood pressure >185 mm Hg or diastolic blood pressure >110 mm Hg. OR = Odds ratio; CI = confidence interval.

Discussion

We conducted the present study to evaluate the clinical significance of ECTs in the hyperacute phase of ischemic stroke and found that the manifestation of ECTs within 90 min after stroke indicates a high risk of subsequent PH. We evaluated PH as a type of ICH which may be more closely related with symptomatic deterioration than hemorrhagic infarction [11]. In our study, PH was observed in 12% of the patients. The frequency of PH was higher than that in ECASS II (3.1%) [2]. Okada et al. [12] reported that PH was found in 10% of 160 Japanese patients with acute

cerebral embolism. Because our study enrolled more patients with cardiogenic embolism than the ECASS II trial, the frequency of PH may become higher. Another explanation for the dissociation of PH frequency between our study and ECASS II is the racial difference in blood coagulation-fibrinolysis factors, such as fibrinogen and factor XII, between Japanese and Caucasians [13].

Several factors related to ECTs were reported to be important. CT density decreases linearly over time, describing the course of water uptake after ischemia [14]. The severity of perfusion impairment also correlates with the degree of CT density reduction [15]. In a recent study using single-photon emission CT, the time from onset to CT and residual cerebral blood flow were independent factors that contributed to the presence of ECTs [16]. Another study using positron emission tomography also shows that ECTs reflect the coupling of the severity of ischemia [17]. These findings suggest that the earlier the manifestation of ECTs, the severer the depth of brain ischemia. ECTs in the extremely early stage likely indicate severe ischemic brain damage associated with poor collaterals, leading to damage to the integrity of the blood-brain barrier [18]. Disruption of the blood-brain barrier contributes to the risk of ICH [19]. Our study showed that the sensitivity of ECTs was relatively low during the initial 90 min, however, the specificity was sufficiently high in each period. These findings were remarkably similar to the results of the study which assessed the significance of X-ray hypoattenuation at CT using the ECASS II population [20]. Careful detection of these subtle findings in the very early periods of ischemic stroke may facilitate the prediction of ICH.

The dichotomized score of ASPECTS 0–7 versus 8–10 was previously validated and shown to have prognostic value among patients treated with intravenous rt-PA for acute ischemic stroke within 3 h of onset [3, 6]. Recently, Dzialowski et al. [21] examined 800 patients within 6 h of stroke onset in the subanalysis of ECASS II and reported that ASPECTS ≤ 7 was a substantially increased risk of thrombolysis-related PH. These results are compatible

with our study showing that in patients without thrombolysis, ASPECTS ≤ 7 was an independent predictor of PH, a sign of poor prognosis.

Several previous studies of intravenous rt-PA therapy suggested that ECTs were not associated with increased risk of ICH [4, 22]. These studies did not describe the interval between onset and CT, and the relationship between the ECT manifestation and the subsequent hemorrhagic transformation was not clarified. However, Demchuk et al. [22] concluded on the basis of such results that there was no evidence of treatment effect modification from the baseline ASPECTS value by thrombolysis. Data from animal models and human clinical trials have demonstrated that earlier stroke treatment is associated with better outcome [5, 23, 24]. There is a possibility that thrombolysis in the very early stage of ischemic stroke may diminish the disruption of the blood-brain barrier and reduce the risk of ICH.

The present study has several limitations. First, it was designed prior to the government approval of intravenous rt-PA in Japan. Therefore, we could not compare the neuroradiological and functional outcome between patients with thrombolysis and those without in spite of the fact that such a comparison may allow us to definitively define a subgroup of patients receiving benefit from thrombolysis. Second, we defined the ECTs according to ASPECTS criteria which include parenchymal hypoattenuation and brain swelling. There is increasing evidence that brain swelling without parenchymal hypoattenuation may be reversible [25]. On the other hand, brain swelling theoretically can represent early stages in a process that manifests finally as PH [26]. The actual relationship between brain swelling and PH remains unclear.

In conclusion, the manifestation of ECTs as represented by ASPECTS ≤ 7 within 90 min after stroke appears to indicate a high risk of parenchymal hemorrhage even when thrombolysis was not performed. Whether or not thrombolysis is indicated for such cases should be clarified in the future by a large-scale study.

References

- 1 National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group: Tissue plasminogen activator for acute ischemic stroke. *N Engl J Med* 1995;333:1581–1587.
- 2 Larrue V, von Kummer R, Muller A, Bluhmki E: Risk factors for severe hemorrhagic transformation in ischemic stroke patients treated with recombinant tissue plasminogen activator: a secondary analysis of the European-Australasian Acute Stroke Study (ECASS II). *Stroke* 2001;32:438–441.
- 3 Barber PA, Demchuk AM, Zhang J, Buchan AM; ASPECTS Study Group: Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. *Lancet* 2000;355:1670–1674.

- 4 Patel SC, Levine SR, Tilley BC, Grotta JC, Lu M, Frankel M, Haley EC Jr, Brott TG, Broderick JP, Horowitz S, Lyden PD, Lewandowski CA, Marler JR, Welch KM; National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group: Lack of clinical significance of early ischemic changes on computed tomography in acute stroke. *JAMA* 2001;286:2830-2838.
- 5 Hacke W, Donnan G, Fieschi C, Kaste M, von Kummer R, Broderick JP, Brott T, Frankel M, Grotta JC, Haley EC Jr, Kwiatkowski T, Levine SR, Lewandowski C, Lu M, Lyden P, Marler JR, Patel S, Tilley BC, Albers G, Bluhmki E, Wilhelm M, Hamilton S; ATLANTIS, ECASS, and NINDS rt-PA Study Group Investigators: Association of outcome with early stroke treatment: pooled analysis of ATLANTIS, ECASS, and NINDS rt-PA stroke trials. *Lancet* 2004;363:768-774.
- 6 Pexman JH, Barber PA, Hill MD, Sevick RJ, Demchuk AM, Hudon ME, Hu WY, Buchan AM: Use of the Alberta Stroke Program Early CT Score (ASPECTS) for assessing CT scans in patients with acute stroke. *AJNR Am J Neuroradiol* 2001;22:1534-1542.
- 7 Mak HK, Yau KK, Khong PL, Ching AS, Cheng PW, Au-Yeung PK, Pang PK, Wong KC, Chan BP: Hypodensity of >1/3 middle cerebral artery territory versus Alberta Stroke Programme Early CT Score (ASPECTS): comparison of two methods of quantitative evaluation of early CT changes in hyperacute ischemic stroke in the community setting. *Stroke* 2003;34:1194-1196.
- 8 Todo K, Moriwaki H, Saito K, Tanaka M, Oe H, Naritomi H: Early CT findings in unknown onset and wake-up strokes. *Cerebrovasc Dis* 2006;21:367-371.
- 9 NINDS t-PA Stroke Study Group: Generalized efficacy of t-PA for acute stroke: subgroup analysis of the NINDS t-PA Stroke Trial. *Stroke* 1997;28:2119-2125.
- 10 Adams HP Jr, Bendixen BH, Kapelle LJ, Biller J, Love BB, Gordon DL, Marsh III EE; TOAST Investigators: Classification of subtype of acute ischemic stroke: definitions for use in a multicenter clinical trial. *Stroke* 1993;24:35-41.
- 11 Trouillas P, von Kummer R: Classification and pathogenesis of cerebral hemorrhages after thrombolysis in ischemic stroke. *Stroke* 2006;37:556-561.
- 12 Okada Y, Yamaguchi T, Minematsu K, Miyashita T, Sawada T, Sadoshima S, Fujishima M, Omae T: Hemorrhagic transformation in cerebral embolism. *Stroke* 1989;20:598-603.
- 13 Ueshima S, Matsuo O: The differences in thrombolytic effects of administered recombinant t-PA between Japanese and Caucasians. *Thromb Haemost* 2002;87:544-546.
- 14 Kucinski T, Vaterlein O, Glauche V, Fiehler J, Klotz E, Eckert B, Koch C, Rother J, Zeumer H: Correlation of apparent diffusion coefficient and computed tomography density in acute ischemic stroke. *Stroke* 2002;33:1786-1791.
- 15 Kucinski T, Majumder A, Knab R, Naumann D, Fiehler J, Vaterlein O, Eckert B, Rother J, Zeumer H: Cerebral perfusion impairment correlates with the decrease of CT density in acute ischaemic stroke. *Neuroradiology* 2004;46:716-722.
- 16 Hirano T, Yonehara T, Inatomi Y, Hashimoto Y, Uchino M: Presence of early ischemic changes on computed tomography depends on severity and the duration of hypoperfusion: a single photon emission-computed tomographic study. *Stroke* 2005;36:2601-2608.
- 17 Sobesky J, von Kummer R, Frackowiak M, Zaro Weber O, Lehnhardt FG, Dohmen C, Neveling M, Moller-Hartmann W, Jacobs AH, Heiss WD: Early ischemic edema on cerebral computed tomography: its relation to diffusion changes and hypoperfusion within 6 h after human ischemic stroke - a comparison of CT, MRI and PET. *Cerebrovasc Dis* 2006;21:336-339.
- 18 Del Zoppo GJ, von Kummer R, Hamann GF: Ischaemic damage of brain microvessels: inherent risks for thrombolytic treatment in stroke. *J Neurol Neurosurg Psychiatry* 1998; 65:1-9.
- 19 Latour LL, Kang DW, Ezzeddine MA, Chalela JA, Warach S: Early blood-brain barrier disruption in human focal brain ischemia. *Ann Neurol* 2004;56:468-477.
- 20 Von Kummer R, Bourquain H, Bastianello S, Bozzao L, Manelfe C, Meier D, Hacke W: Early prediction of irreversible brain damage after ischemic stroke at CT. *Radiology* 2001; 219:95-100.
- 21 Dzialowski I, Hill MD, Coutts SB, Demchuk AM, Kent DM, Wunderlich O, von Kummer R: Extent of early ischemic changes on computed tomography (CT) before thrombolysis: prognostic value of the Alberta Stroke Program Early CT Score in ECASS II. *Stroke* 2006;37:973-978.
- 22 Demchuk AM, Hill MD, Barber PA, Silver B, Patel SC, Levine SR; NINDS rtPA Stroke Study Group, NIH: Importance of early ischemic computed tomography changes using ASPECTS in NINDS rtPA stroke study. *Stroke* 2005;36:2110-2115.
- 23 Zivin JA, Lyden PD, DeGirolami U, Kochhar A, Mazzarella V, Hemenway CC, Johnston P: Tissue plasminogen activator: reduction of neurologic damage after experimental embolic stroke. *Arch Neurol* 1988;45:387-391.
- 24 Marler JR, Tilley BC, Lu M, Brott TG, Lyden PC, Grotta JC, Broderick JP, Levine SR, Frankel MP, Horowitz SH, Haley EC Jr, Lewandowski CA, Kwiatkowski TP; NINDS rt-PA Stroke Study Group: Early stroke treatment associated with better outcome: the NINDS rt-PA stroke study. *Neurology* 2000; 55:1649-1655.
- 25 Na DG, Kim EY, Ryoo JW, Lee KH, Roh HG, Kim SS, Song IC, Chang KH: CT sign of brain swelling without concomitant parenchymal hypoattenuation: comparison with diffusion- and perfusion-weighted MR imaging. *Radiology* 2005;235:992-998.
- 26 Simard JM, Kent TA, Chen M, Tarasov KV, Gerzanich V: Brain oedema in focal ischaemia: molecular pathophysiology and theoretical implications. *Lancet Neurol* 2007;6: 258-268.

Brief Communication

Increase in circulating CD34-positive cells in patients with angiographic evidence of moyamoya-like vessels

Tomoyuki Yoshihara¹, Akihiko Taguchi¹, Tomohiro Matsuyama², Yoko Shimizu¹, Akie Kikuchi-Taura³, Toshihiro Soma³, David M Stern⁴, Hiroo Yoshikawa⁵, Yukiko Kasahara¹, Hiroshi Moriwaki¹, Kazuyuki Nagatsuka¹ and Hiroaki Naritomi¹

¹Department of Cerebrovascular Disease, National Cardiovascular Center, Osaka, Japan; ²Institute for Advanced Medical Sciences, Hyogo College of Medicine, Hyogo, Japan; ³Department of Hematology, Osaka Minami National Medical Center, Osaka, Japan; ⁴Dean's Office, College of Medicine, University of Cincinnati, Cincinnati, Ohio, USA; ⁵Department of Internal Medicine, Hyogo College of Medicine, Hyogo, Japan

Increasing evidence points to a role for circulating endothelial progenitor cells, including populations of CD34-positive (CD34⁺) cells, in maintenance of cerebral blood flow. In this study, we investigated the link between the level of circulating CD34⁺ cells and neovascularization at ischemic brain. Compared with control subjects, a remarkable increase of circulating CD34⁺ cells was observed in patients with angiographic moyamoya vessels, although no significant change was observed in patients with major cerebral artery occlusion (or severe stenosis) but without moyamoya vessels. Our results suggest that the increased level of CD34⁺ cells associated with ischemic stress is correlated with neovascularization at human ischemic brain.

Journal of Cerebral Blood Flow & Metabolism advance online publication, 30 January 2008; doi:10.1038/jcbfm.2008.1

Keywords: antigens; CD34; moyamoya vessel; neovascularization

Introduction

Increasing evidence points to a role for bone marrow-derived immature cells, such as endothelial progenitor cells, in maintenance of vascular homeostasis and repair. CD34-positive (CD34⁺) cells comprise a population enriched for endothelial progenitor cells whose contribution to neovascularization includes both direct participation in forming the neovessel and regulatory roles as sources of growth/angiogenesis factors (Majka *et al*, 2001). Previously, we have shown accelerated neovascularization after administration of CD34⁺ cells in an experimental model of stroke (Taguchi *et al*, 2004b) and induced by autologous bone marrow mononuclear cells (rich cell fraction of CD34⁺ cells)

transplanted locally into patients with limb ischemia (Taguchi *et al*, 2003). In addition, we have observed a positive correlation between the level of circulating CD34⁺ cells and regional blood flow (Taguchi *et al*, 2004a), and cognitive function (Taguchi *et al*, 2007) in patients with chronic cerebral ischemia.

In this study, we have evaluated the level of circulating CD34⁺ cells in patients with unusually accelerated neovascularization induced by progressive occlusion (or severe stenosis) of the supraclavicular portion of the internal carotid artery, the proximal region of the anterior, and/or middle cerebral artery characterized angiographically by the presence of moyamoya-like vessels (Natori *et al*, 1997) that supply ischemic brain as collaterals. We have investigated the hypothesis that circulating bone marrow-derived immature cells might be associated with neovascularization at ischemic sites in the human brain.

Correspondence: Dr A Taguchi, Department of Cerebrovascular Disease, National Cardiovascular Center, 5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan.
E-mail: taguchi@ri.ncvc.go.jp

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Health, Labour, and Welfare (H19-Choujyu-029).

Received 29 October 2007; revised 19 December 2007; accepted 26 December 2007

Patients and methods

The institutional review board of the National Cardiovascular Center approved this study. All subjects provided

informed consent. A total of 50 individuals, including 24 patients with occlusion or severe stenosis (>90%) at the C1 portion of the internal carotid artery or the M1 portion of the middle cerebral artery, and 26 age-matched healthy volunteers with cardiovascular risk factors, but without history of vascular disease, were enrolled. The diagnosis of cerebral artery occlusion or stenosis was made angiographically and four patients were found to have classical angiographic evidence of moyamoya-like vessels, including one with right C1 occlusion, one with right M1 occlusion, and two with bilateral C1 severe stenosis. All patients with cerebral artery occlusion or stenosis had a history of cerebral infarction. Individuals excluded from the study included patients who experienced a vascular event within 30 days of measurements, premenopausal women, and those with evidence of infection and/or malignant disease. The number of circulating CD34⁺ cells was quantified as described (Taguchi *et al*, 2007). In brief, blood samples (200 µl) were incubated with phycoerythrin-labeled anti-CD34 antibody, fluorescein isothiocyanate-labeled anti-CD45 antibody, 7-aminoactinomycin-D (7-AAD), and internal control (all of these reagents are in the Stem-Kit; BeckmanCoulter, Marseille, France). After incubation, samples were centrifuged, and supernatant was removed to obtain concentrated cell suspensions. 7-Aminoactinomycin-D-positive dead cells and CD45-negative cells were excluded, and the number of cells forming clusters characteristic of CD34⁺ cells (i.e., low side scatter and low-to-intermediate CD45 staining) was counted. The absolute number of CD34⁺ cells was calculated using the internal control. Mean cell number of duplicate measurements was used for quantitative analysis. Statistical comparisons among groups were determined using analysis of variance or χ^2 test. Individual comparisons were performed using a two-tailed unpaired Students' *t*-test or Mann-Whitney's *U*-test. Mean \pm s.e. is shown.

Results

Enrolled individuals were divided into three groups: control subjects, patients with cerebral occlusion or severe stenosis, but without the presence of vessels with angiographic characteristics of moyamoya disease, and patients with angiographic evidence of moyamoya-like vessels. Baseline characteristics of the groups are shown in Table 1. The modified Rankin scale evaluation of patients with and without moyamoya-like vessels was 0.5 ± 0.5 and 1.3 ± 0.2 , respectively ($P=0.15$). Comparing these groups, there was a significant difference in the ratio of gender and treatment with aspirin between groups. However, no significant difference was observed in the number of circulating CD34⁺ cells in control group between genders (male, $n=13$, CD34⁺ cells = $0.93 \pm 0.10/\mu\text{L}$; female, $n=13$, CD34⁺ cells = $0.85 \pm 0.11/\mu\text{L}$; $P=0.59$) and treatment with aspirin (aspirin (+), $n=6$, CD34⁺ cells = $0.76 \pm 0.12/\mu\text{L}$; aspirin (-), $n=20$, CD34⁺ cells = $0.93 \pm 0.09/\mu\text{L}$; $P=0.26$), indicating mild and nonsignificant effects of gender and treatment with aspirin on the level of circulating CD34⁺ cells. In univariate analysis of control subjects, each cerebrovascular risk factor and treatment with statins showed no significant difference in the number of circulating CD34⁺ cells (data not shown).

A representative angiogram showing characteristics of moyamoya-like vessels is shown in Figures 1A and 1B. Angiographic moyamoya-like vessels were observed around the M1 portion of an occluded middle cerebral artery. Compared with a normal subject (Figure 1C) and patients without angiographic evidence of moyamoya-like vessels (Figure 1D), a remarkable increase in levels of

Table 1 Baseline characteristics

	Total	Control	Major artery occlusion/stenosis		P-value for trend
			Moyamoya (-)	Moyamoya (+)	
N	50	26	20	4	
Age, years	60.8 \pm 1.1	60.5 \pm 1.9	61.5 \pm 1.0	59.3 \pm 5.9	0.85
Male, n (%)	33 (66)	13 (50)	18 (90)	2 (50)	0.01
<i>Risk factor, n (%)</i>					
Hypertension	35 (70)	16 (62)	15 (75)	4 (100)	0.24
Hyperlipidemia	26 (52)	14 (54)	10 (50)	2 (50)	0.96
Diabetes mellitus	11 (22)	7 (27)	4 (20)	0 (0)	0.46
Smoking	15 (30)	7 (27)	8 (40)	0 (0)	0.25
<i>Treatment, n (%)</i>					
Ca channel blockers	20 (40)	10 (38)	8 (40)	2 (50)	0.91
β -Blockers	5 (10)	3 (11)	1 (5)	1 (25)	0.44
ACE inhibitor	7 (14)	4 (15)	2 (10)	1 (25)	0.70
ARB	12 (24)	5 (19)	5 (25)	2 (50)	0.40
Diuretics	4 (8)	2 (7)	1 (5)	1 (25)	0.40
Statin therapy	14 (28)	9 (34)	4 (20)	1 (25)	0.54
Aspirin	19 (38)	6 (23)	10 (50)	3 (75)	0.05
Ticlopidine	12 (24)	3 (11)	8 (40)	1 (25)	0.08

Abbreviations: ACE, angiotensin-converting enzyme; ARB, angiotensin 2 receptor blocker.




Figure 1 Increased levels of circulating CD34⁺ cells in patients with angiographic evidence of moyamoya-like vessels. (A, B) Representative angiogram from a patient with moyamoya-like vessels. Unusually accelerated neovascularization (based on angiographic features of moyamoya-like vessels, arrowheads) was observed around an occlusive M1 lesion (arrow). Anterior-posterior view (A) and lateral view (B) of the right internal carotid artery showed angiographically. (C–E) After exclusion of 7-aminoactinomycin-D (7-AAD)-positive dead cells and CD45-negative cells (nonleukocytes), CD34⁺ cells cluster at low side scatter. Representative fluorescence-activated cell sorting analyses from a control subject (C), a patient without moyamoya-like vessels (D), and a patient with moyamoya-like vessels (E) are shown. (F) A more than two-fold increase in circulating CD34⁺ cells was observed in patients with moyamoya-like vessels, compared with control subjects and patients without moyamoya-like vessels (**P* < 0.001). SS Lin: side-scatter linear scale.

peripheral CD34⁺ cells was observed in patients with moyamoya-like vessels (Figure 1E) based on fluorescence-activated cell sorting. To confirm this impression, levels of circulating CD34⁺ cells were quantified (control, CD34⁺ cells = 0.89 ± 0.07/μL; moyamoya (-), CD34⁺ cells = 0.98 ± 0.13/μL; moyamoya (+), CD34⁺ cells = 2.28 ± 0.53/μL) and found to be significantly increased in patients with moyamoya-like vessels more than two-fold higher than in controls (Figure 1F, *P* < 0.001).

Discussion

In this study, we have found that a feature of unusually accelerated neovascularization, evidence of moyamoya-like vessels in the immediate locale of an occluded major cerebral artery, can be correlated with a robust increase in the level of circulating

CD34⁺ cells. The latter was determined using a newly developed method that enables quantification of few CD34⁺ cells in peripheral blood in a highly reproducible manner.

After acute cerebral ischemia, mobilization of CD34⁺ cells from bone marrow has been shown in stroke patients (Taguchi *et al*, 2004a). Furthermore, transplantation of CD34⁺ cells (Taguchi *et al*, 2004b) and bone marrow cells (Borlongan *et al*, 2004a,b) has been shown to restore cerebral blood flow in experimental models of stroke. In chronic ischemia, transplantation of CD34⁺ cells has also been shown to accelerate neovascularization, including formation of collateral vessels, in patients with chronic ischemic heart disease (Boyle *et al*, 2006) and limb ischemia (Kudo *et al*, 2003). In addition, there is a report regarding the correlation between inadequate coronary collateral development and reduced numbers of circulating endothelial progenitor cells in

patients with myocardial ischemia (Lambiase *et al*, 2004). In this study, we show, for the first time, a correlation between neovascularization of the cerebral arterial circulation and increased levels of circulating CD34⁺ cells. Our results support the hypothesis that circulating CD34⁺ cells potentially contribute to neovascularization at sites of ischemic brain injury.

Acknowledgements

We thank K Obata and Y Okinaka for technical assistance.

Conflict of interest

The authors state no conflict of interest.

References

Borlongan CV, Lind JG, Dillon-Carter O, Yu G, Hadman M, Cheng C, Carroll J, Hess DC (2004a) Bone marrow grafts restore cerebral blood flow and blood brain barrier in stroke rats. *Brain Res* 1010:108–16

Borlongan CV, Lind JG, Dillon-Carter O, Yu G, Hadman M, Cheng C, Carroll J, Hess DC (2004b) Intracerebral xenografts of mouse bone marrow cells in adult rats facilitate restoration of cerebral blood flow and blood-brain barrier. *Brain Res* 1009:26–33

Boyle AJ, Whitbourn R, Schlicht S, Krum H, Kocher A, Nandurkar H, Bergmann S, Daniell M, O'Day J, Skerrett D, Haylock D, Gilbert RE, Itescu S (2006) Intra-coronary high-dose CD34⁺ stem cells in patients with chronic ischemic heart disease: a 12-month follow-up. *Int J Cardiol* 109:21–7

Kudo FA, Nishibe T, Nishibe M, Yasuda K (2003) Autologous transplantation of peripheral blood endo-

thelial progenitor cells (CD34⁺) for therapeutic angiogenesis in patients with critical limb ischemia. *Int Angiol* 22:344–8

Lambiase PD, Edwards RJ, Anthopoulos P, Rahman S, Meng YG, Bucknall CA, Redwood SR, Pearson JD, Marber MS (2004) Circulating humoral factors and endothelial progenitor cells in patients with differing coronary collateral support. *Circulation* 109:2986–92

Majka M, Janowska-Wieczorek A, Ratajczak J, Ehrenman K, Pietrkowski Z, Kowalska MA, Gewirtz AM, Emerson SG, Ratajczak MZ (2001) Numerous growth factors, cytokines, and chemokines are secreted by human CD34⁺ cells, myeloblasts, erythroblasts, and megakaryoblasts and regulate normal hematopoiesis in an autocrine/paracrine manner. *Blood* 97:3075–85

Natori Y, Ikezaki K, Matsushima T, Fukui M (1997) 'Angiographic moyamoya' its definition, classification, and therapy. *Clin Neurol Neurosurg* 99(Suppl 2): S168–72

Taguchi A, Matsuyama T, Moriwaki H, Hayashi T, Hayashida K, Nagatsuka K, Todo K, Mori K, Stern DM, Soma T, Naritomi H (2004a) Circulating CD34-positive cells provide an index of cerebrovascular function. *Circulation* 109:2972–5

Taguchi A, Matsuyama T, Nakagomi T, Shimizu Y, Fukunaga R, Tatsumi Y, Yoshikawa H, Kikuchi-Taura A, Soma T, Moriwaki H, Nagatsuka K, Stern DM, Naritomi H (2007) Circulating CD34-positive cells provide a marker of vascular risk associated with cognitive impairment. *J Cereb Blood Flow Metab*; e-pub ahead of print 8 August 2007

Taguchi A, Ohtani M, Soma T, Watanabe M, Kinoshita N (2003) Therapeutic angiogenesis by autologous bone-marrow transplantation in a general hospital setting. *Eur J Vasc Endovasc Surg* 25:276–8

Taguchi A, Soma T, Tanaka H, Kanda T, Nishimura H, Yoshikawa H, Tsukamoto Y, Iso H, Fujimori Y, Stern DM, Naritomi H, Matsuyama T (2004b) Administration of CD34⁺ cells after stroke enhances neurogenesis via angiogenesis in a mouse model. *J Clin Invest* 114:330–8

Brief Communication

Increase in circulating CD34-positive cells in patients with angiographic evidence of moyamoya-like vessels

Tomoyuki Yoshihara¹, Akihiko Taguchi¹, Tomohiro Matsuyama², Yoko Shimizu¹, Akie Kikuchi-Taura³, Toshihiro Soma³, David M Stern⁴, Hiroo Yoshikawa⁵, Yukiko Kasahara¹, Hiroshi Moriwaki¹, Kazuyuki Nagatsuka¹ and Hiroaki Naritomi¹

¹Department of Cerebrovascular Disease, National Cardiovascular Center, Osaka, Japan; ²Institute for Advanced Medical Sciences, Hyogo College of Medicine, Hyogo, Japan; ³Department of Hematology, Osaka Minami National Medical Center, Osaka, Japan; ⁴Dean's Office, College of Medicine, University of Cincinnati, Cincinnati, Ohio, USA; ⁵Department of Internal Medicine, Hyogo College of Medicine, Hyogo, Japan

Increasing evidence points to a role for circulating endothelial progenitor cells, including populations of CD34-positive (CD34⁺) cells, in maintenance of cerebral blood flow. In this study, we investigated the link between the level of circulating CD34⁺ cells and neovascularization at ischemic brain. Compared with control subjects, a remarkable increase of circulating CD34⁺ cells was observed in patients with angiographic moyamoya vessels, although no significant change was observed in patients with major cerebral artery occlusion (or severe stenosis) but without moyamoya vessels. Our results suggest that the increased level of CD34⁺ cells associated with ischemic stress is correlated with neovascularization at human ischemic brain.

Journal of Cerebral Blood Flow & Metabolism advance online publication, 30 January 2008; doi:10.1038/jcbfm.2008.1

Keywords: antigens; CD34; moyamoya vessel; neovascularization

Introduction

Increasing evidence points to a role for bone marrow-derived immature cells, such as endothelial progenitor cells, in maintenance of vascular homeostasis and repair. CD34-positive (CD34⁺) cells comprise a population enriched for endothelial progenitor cells whose contribution to neovascularization includes both direct participation in forming the neovessel and regulatory roles as sources of growth/angiogenesis factors (Majka *et al*, 2001). Previously, we have shown accelerated neovascularization after administration of CD34⁺ cells in an experimental model of stroke (Taguchi *et al*, 2004b) and induced by autologous bone marrow mononuclear cells (rich cell fraction of CD34⁺ cells)

transplanted locally into patients with limb ischemia (Taguchi *et al*, 2003). In addition, we have observed a positive correlation between the level of circulating CD34⁺ cells and regional blood flow (Taguchi *et al*, 2004a), and cognitive function (Taguchi *et al*, 2007) in patients with chronic cerebral ischemia.

In this study, we have evaluated the level of circulating CD34⁺ cells in patients with unusually accelerated neovascularization induced by progressive occlusion (or severe stenosis) of the supraclinoid portion of the internal carotid artery, the proximal region of the anterior, and/or middle cerebral artery characterized angiographically by the presence of moyamoya-like vessels (Natori *et al*, 1997) that supply ischemic brain as collaterals. We have investigated the hypothesis that circulating bone marrow-derived immature cells might be associated with neovascularization at ischemic sites in the human brain.

Correspondence: Dr A Taguchi, Department of Cerebrovascular Disease, National Cardiovascular Center, 5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan.

E-mail: taguchi@ri.ncvc.go.jp

This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Health, Labour, and Welfare (H19-Choujyu-029).

Received 29 October 2007; revised 19 December 2007; accepted 26 December 2007

Patients and methods

The institutional review board of the National Cardiovascular Center approved this study. All subjects provided

informed consent. A total of 50 individuals, including 24 patients with occlusion or severe stenosis (>90%) at the C1 portion of the internal carotid artery or the M1 portion of the middle cerebral artery, and 26 age-matched healthy volunteers with cardiovascular risk factors, but without history of vascular disease, were enrolled. The diagnosis of cerebral artery occlusion or stenosis was made angiographically and four patients were found to have classical angiographic evidence of moyamoya-like vessels, including one with right C1 occlusion, one with right M1 occlusion, and two with bilateral C1 severe stenosis. All patients with cerebral artery occlusion or stenosis had a history of cerebral infarction. Individuals excluded from the study included patients who experienced a vascular event within 30 days of measurements, premenopausal women, and those with evidence of infection and/or malignant disease. The number of circulating CD34⁺ cells was quantified as described (Taguchi *et al*, 2007). In brief, blood samples (200 µl) were incubated with phycoerythrin-labeled anti-CD34 antibody, fluorescein isothiocyanate-labeled anti-CD45 antibody, 7-aminoactinomycin-D (7-AAD), and internal control (all of these reagents are in the Stem-Kit; BeckmanCoulter, Marseille, France). After incubation, samples were centrifuged, and supernatant was removed to obtain concentrated cell suspensions. 7-Aminoactinomycin-D-positive dead cells and CD45-negative cells were excluded, and the number of cells forming clusters characteristic of CD34⁺ cells (i.e., low side scatter and low-to-intermediate CD45 staining) was counted. The absolute number of CD34⁺ cells was calculated using the internal control. Mean cell number of duplicate measurements was used for quantitative analysis. Statistical comparisons among groups were determined using analysis of variance or χ^2 test. Individual comparisons were performed using a two-tailed unpaired Students' *t*-test or Mann-Whitney's *U*-test. Mean \pm s.e. is shown.

Results

Enrolled individuals were divided into three groups: control subjects, patients with cerebral occlusion or severe stenosis, but without the presence of vessels with angiographic characteristics of moyamoya disease, and patients with angiographic evidence of moyamoya-like vessels. Baseline characteristics of the groups are shown in Table 1. The modified Rankin scale evaluation of patients with and without moyamoya-like vessels was 0.5 ± 0.5 and 1.3 ± 0.2 , respectively ($P=0.15$). Comparing these groups, there was a significant difference in the ratio of gender and treatment with aspirin between groups. However, no significant difference was observed in the number of circulating CD34⁺ cells in control group between genders (male, $n=13$, CD34⁺ cells = $0.93 \pm 0.10/\mu\text{L}$; female, $n=13$, CD34⁺ cells = $0.85 \pm 0.11/\mu\text{L}$; $P=0.59$) and treatment with aspirin (aspirin (+), $n=6$, CD34⁺ cells = $0.76 \pm 0.12/\mu\text{L}$; aspirin (-), $n=20$, CD34⁺ cells = $0.93 \pm 0.09/\mu\text{L}$; $P=0.26$), indicating mild and nonsignificant effects of gender and treatment with aspirin on the level of circulating CD34⁺ cells. In univariate analysis of control subjects, each cerebrovascular risk factor and treatment with statins showed no significant difference in the number of circulating CD34⁺ cells (data not shown).

A representative angiogram showing characteristics of moyamoya-like vessels is shown in Figures 1A and 1B. Angiographic moyamoya-like vessels were observed around the M1 portion of an occluded middle cerebral artery. Compared with a normal subject (Figure 1C) and patients without angiographic evidence of moyamoya-like vessels (Figure 1D), a remarkable increase in levels of

Table 1 Baseline characteristics

	Total	Control	Major artery occlusion/stenosis		P-value for trend
			Moyamoya (-)	Moyamoya (+)	
<i>N</i>	50	26	20	4	
Age, years	60.8 \pm 1.1	60.5 \pm 1.9	61.5 \pm 1.0	59.3 \pm 5.9	0.85
Male, <i>n</i> (%)	33 (66)	13 (50)	18 (90)	2 (50)	0.01
<i>Risk factor, n (%)</i>					
Hypertension	35 (70)	16 (62)	15 (75)	4 (100)	0.24
Hyperlipidemia	26 (52)	14 (54)	10 (50)	2 (50)	0.96
Diabetes mellitus	11 (22)	7 (27)	4 (20)	0 (0)	0.46
Smoking	15 (30)	7 (27)	8 (40)	0 (0)	0.25
<i>Treatment, n (%)</i>					
Ca channel blockers	20 (40)	10 (38)	8 (40)	2 (50)	0.91
β -Blockers	5 (10)	3 (11)	1 (5)	1 (25)	0.44
ACE inhibitor	7 (14)	4 (15)	2 (10)	1 (25)	0.70
ARB	12 (24)	5 (19)	5 (25)	2 (50)	0.40
Diuretics	4 (8)	2 (7)	1 (5)	1 (25)	0.40
Statin therapy	14 (28)	9 (34)	4 (20)	1 (25)	0.54
Aspirin	19 (38)	6 (23)	10 (50)	3 (75)	0.05
Ticlopidine	12 (24)	3 (11)	8 (40)	1 (25)	0.08

Abbreviations: ACE, angiotensin-converting enzyme; ARB, angiotensin 2 receptor blocker.




Figure 1 Increased levels of circulating CD34⁺ cells in patients with angiographic evidence of moyamoya-like vessels. (A, B) Representative angiogram from a patient with moyamoya-like vessels. Unusually accelerated neovascularization (based on angiographic features of moyamoya-like vessels, arrowheads) was observed around an occlusive M1 lesion (arrow). Anterior-posterior view (A) and lateral view (B) of the right internal carotid artery showed angiographically. (C–E) After exclusion of 7-aminoactinomycin-D (7-AAD)-positive dead cells and CD45-negative cells (nonleukocytes), CD34⁺ cells cluster at low side scatter. Representative fluorescence-activated cell sorting analyses from a control subject (C), a patient without moyamoya-like vessels (D), and a patient with moyamoya-like vessels (E) are shown. (F) A more than two-fold increase in circulating CD34⁺ cells was observed in patients with moyamoya-like vessels, compared with control subjects and patients without moyamoya-like vessels (**P* < 0.001). SS Lin: side-scatter linear scale.

peripheral CD34⁺ cells was observed in patients with moyamoya-like vessels (Figure 1E) based on fluorescence-activated cell sorting. To confirm this impression, levels of circulating CD34⁺ cells were quantified (control, CD34⁺ cells = $0.89 \pm 0.07/\mu\text{L}$; moyamoya (-), CD34⁺ cells = $0.98 \pm 0.13/\mu\text{L}$; moyamoya (+), CD34⁺ cells = $2.28 \pm 0.53/\mu\text{L}$) and found to be significantly increased in patients with moyamoya-like vessels more than two-fold higher than in controls (Figure 1F, *P* < 0.001).

Discussion

In this study, we have found that a feature of unusually accelerated neovascularization, evidence of moyamoya-like vessels in the immediate locale of an occluded major cerebral artery, can be correlated with a robust increase in the level of circulating

CD34⁺ cells. The latter was determined using a newly developed method that enables quantification of few CD34⁺ cells in peripheral blood in a highly reproducible manner.

After acute cerebral ischemia, mobilization of CD34⁺ cells from bone marrow has been shown in stroke patients (Taguchi *et al*, 2004a). Furthermore, transplantation of CD34⁺ cells (Taguchi *et al*, 2004b) and bone marrow cells (Borlongan *et al*, 2004a, b) has been shown to restore cerebral blood flow in experimental models of stroke. In chronic ischemia, transplantation of CD34⁺ cells has also been shown to accelerate neovascularization, including formation of collateral vessels, in patients with chronic ischemic heart disease (Boyle *et al*, 2006) and limb ischemia (Kudo *et al*, 2003). In addition, there is a report regarding the correlation between inadequate coronary collateral development and reduced numbers of circulating endothelial progenitor cells in