

- patterns of recurrence after resection of hepatocellular carcinoma. *J Am Coll Surg* 2003; 197: 753–8.
- 7 Belghiti J, Panis Y, Farges O, Benhamou JP, Fekete F. Intrahepatic recurrence after resection of hepatocellular carcinoma complicating cirrhosis. *Ann Surg* 1991; 214: 114–7.
 - 8 Kitamoto M, Nakanishi T, Kira S *et al*. The assessment of proliferating cell nuclear antigen immunohistochemical staining in small hepatocellular carcinoma and its relationship to histologic characteristics and prognosis. *Cancer* 1993; 72: 1859–65.
 - 9 Osada T, Sakamoto M, Iino Y *et al*. E-cadherin is involved in the intrahepatic metastasis of hepatocellular carcinoma. *Hepatology* 1996; 24: 1460–7.
 - 10 Okada S, Shimada K, Yamamoto J *et al*. Predictive factors for postoperative recurrence of hepatocellular carcinoma. *Gastroenterology* 1994; 106: 1618–24.
 - 11 Tsuda H, Zhang WD, Shimosato Y *et al*. Allele loss on chromosome 16 associated with progression of human hepatocellular carcinoma. *Proc Natl Acad Sci USA* 1990; 87: 6791–4.
 - 12 Nakanishi K, Sakamoto M, Yamasaki S, Todo S, Hirohashi S. Akt phosphorylation is a risk factor for early disease recurrence and poor prognosis in hepatocellular carcinoma. *Cancer* 2005; 103: 307–12.
 - 13 Peng SY, Chen WJ, Lai PL, Jeng YM, Sheu JC, Hsu HC. High alpha-fetoprotein level correlates with high stage, early recurrence and poor prognosis of hepatocellular carcinoma: significance of hepatitis virus infection, age, p53 and beta-catenin mutations. *Int J Cancer* 2004; 112: 44–50.
 - 14 Ye QH, Qin LX, Forgues M *et al*. Predicting hepatitis B virus-positive metastatic hepatocellular carcinomas using gene expression profiling and supervised machine learning. *Nat Med* 2003; 9: 416–23.
 - 15 Iizuka N, Oka M, Yamada-Okabe H *et al*. Oligonucleotide microarray for prediction of early intrahepatic recurrence of hepatocellular carcinoma after curative resection. *Lancet* 2003; 361: 923–9.
 - 16 Yuan RH, Jeng YM, Chen HL *et al*. Stathmin overexpression cooperates with p53 mutation and osteopontin overexpression, and is associated with tumor progression, early recurrence, and poor prognosis in hepatocellular carcinoma. *J Pathol* 2006; 209: 549–58.
 - 17 Budhu A, Forgues M, YeQH *et al*. Prediction of venous metastases, recurrence, and prognosis in hepatocellular carcinoma based on a unique immune response signature of the liver microenvironment. *Cancer Cell* 2006; 10: 99–111.
 - 18 Griffin TJ, Gygi SP, Ideker T *et al*. Complementary profiling of gene expression at the transcriptome and proteome levels in *Saccharomyces cerevisiae*. *Mol Cell Proteomics* 2002; 1: 323–33.
 - 19 Chen G, Gharib TG, Huang CC *et al*. Discordant protein and mRNA expression in lung adenocarcinomas. *Mol Cell Proteomics* 2002; 1: 304–13.
 - 20 Gygi SP, Rochon Y, Franza BR, Aebersold R. Correlation between protein and mRNA abundance in yeast. *Mol Cell Biol* 1999; 19: 1720–30.
 - 21 Ding SJ, Li Y, Tan YX *et al*. From proteomic analysis to clinical significance: overexpression of cytokeratin 19 correlates with hepatocellular carcinoma metastasis. *Mol Cell Proteomics* 2004; 3: 73–81.
 - 22 Yokoo H, Kondo T, Fujii K, Yamada T, Todo S, Hirohashi S. Proteomic signature corresponding to alpha fetoprotein expression in liver cancer cells. *Hepatology* 2004; 40: 609–17.
 - 23 Fujii K, Kondo T, Yokoo H, Yamada T, Iwatsuki K, Hirohashi S. Proteomic study of human hepatocellular carcinoma using two-dimensional difference gel electrophoresis with saturation cysteine dye. *Proteomics* 2005; 5: 1411–22.
 - 24 Fujii K, Kondo T, Yokoo H *et al*. Database of two-dimensional polyacrylamide gel electrophoresis of proteins labeled with CyDye DIGE Fluor saturation dye. *Proteomics* 2006; 6: 1640–53.
 - 25 Chignard N, Shang S, Wang H *et al*. Cleavage of endoplasmic reticulum proteins in hepatocellular carcinoma: detection of generated fragments in patient sera. *Gastroenterology* 2006; 130: 2010–22.
 - 26 Kuramitsu Y, Harada T, Takashima M *et al*. Increased expression and phosphorylation of liver glutamine synthetase in well-differentiated hepatocellular carcinoma tissues from patients infected with hepatitis C virus. *Electrophoresis* 2006; 27: 2998.
 - 27 Shen H, Cheng G, Fan H *et al*. Expressed proteome analysis of human hepatocellular carcinoma in nude mice (LCI-D20) with high metastasis potential. *Proteomics* 2006; 6: 528–37.
 - 28 Song HY, Liu YK, Feng JT *et al*. Proteomic analysis on metastasis-associated proteins of human hepatocellular carcinoma tissues. *J Cancer Res Clin Oncol* 2006; 132: 92–8.
 - 29 Ramaswamy S, Tamayo P, Rifkin R *et al*. Multiclass cancer diagnosis using tumor gene expression signatures. *Proc Natl Acad Sci USA* 2001; 98: 15 149–54.
 - 30 Toyosaka A, Okamoto E, Mitsunobu M, Oriyama T, Nakao N, Miura K. Intrahepatic metastases in hepatocellular carcinoma: evidence for spread via the portal vein as an efferent vessel. *Am J Gastroenterol* 1996; 91: 1610–15.
 - 31 Regimbeau JM, Abdalla EK, Vauthey JN *et al*. Risk factors for early death due to recurrence after liver resection for hepatocellular carcinoma: results of a multicenter study. *J Surg Oncol* 2004; 85: 36–41.
 - 32 Snell K, Fell DA. Metabolic control analysis of mammalian serine metabolism. *Adv Enzyme Regul* 1990; 30: 13–32.
 - 33 Snell K, Weber G. Enzymic imbalance in serine metabolism in rat hepatomas. *Biochem J* 1986; 233: 617–20.
 - 34 Snell K. Enzymes of serine metabolism in normal, developing and neoplastic rat tissues. *Adv Enzyme Regul* 1984; 22: 325–400.
 - 35 Farazi PA, DePinho RA. Hepatocellular carcinoma pathogenesis: from genes to environment. *Nat Rev Cancer* 2006; 6: 674–87.
 - 36 Misek DE, Chang CL, Kuick R *et al*. Transforming properties of a Q18->E mutation of the microtubule regulator Op18. *Cancer Cell* 2002; 2: 217–28.
 - 37 Chen G, Gharib TG, Wang H *et al*. Protein profiles associated with survival in lung adenocarcinoma. *Proc Natl Acad Sci USA* 2003; 100: 13 537–42.
 - 38 Blanc JF, Lalame C, Plomion C *et al*. Proteomic analysis of differentially expressed proteins in hepatocellular carcinoma developed in patients with chronic viral hepatitis C. *Proteomics* 2005; 5: 3778–89.
 - 39 Amabi B, Currie JC, Bouzeghrane M *et al*. Contribution of the 37-kDa laminin receptor precursor in the anti-metastatic P5P94-derived peptide PCK3145 cell surface binding. *Biochem Biophys Res Commun* 2006; 346: 358–66.
 - 40 Choi JH, Kim TN, Kim S *et al*. Overexpression of mitochondrial thioredoxin reductase and peroxiredoxin III in hepatocellular carcinomas. *Anticancer Res* 2002; 22: 3331–5.
 - 41 Zhu XD, Li CL, Lang ZW, Gao GF, Tien P. Significant correlation between expression level of HSP gp96 and progression of hepatitis B virus induced diseases. *World J Gastroenterol* 2004; 10: 1141–5.
 - 42 Masaki T, Tokuda M, Ohnishi M *et al*. Enhanced expression of the protein kinase substrate annexin in human hepatocellular carcinoma. *Hepatology* 1996; 24: 72–81.
 - 43 Huang C, Wu M. Superoxide dismutase activity in tissues from 19 cases of hepatocellular carcinoma. *Zhonghua Yi Xue Za Zhi* 1990; 70: 138–9, 12.
 - 44 Stenner-Liewen F, Luo G, Sahin U *et al*. Definition of tumor-associated antigens in hepatocellular carcinoma. *Cancer Epidemiol Biomarkers Prev* 2000; 9: 285–90.
 - 45 Lim SO, Park SJ, Kim W *et al*. Proteome analysis of hepatocellular carcinoma. *Biochem Biophys Res Commun* 2002; 291: 1031–7.
 - 46 Li C, Tan YX, Zhou H *et al*. Proteomic analysis of hepatitis B virus-associated hepatocellular carcinoma: Identification of potential tumor markers. *Proteomics* 2005; 5: 1125–39.
 - 47 Pang R, Yuen J, Yuen MF *et al*. PIN1 overexpression and beta-catenin gene mutations are distinct oncogenic events in human hepatocellular carcinoma. *Oncogene* 2004; 23: 4182–6.
 - 48 Yokota S, Yamamoto Y, Shinizu K *et al*. Increased expression of cytosolic chaperonin CCT in human hepatocellular and colonic carcinoma. *Cell Stress Chaperones* 2001; 6: 345–50.
 - 49 Zheng S, Ruan Y, Wu Z, Tang J. The relationship between; 67KD: laminin receptor expression and metastasis of hepatocellular carcinoma. *J Tongji Med University* 1997; 17: 200–2, 24.
 - 50 Gorg A, Weiss W, Dunn MJ. Current two-dimensional electrophoresis technology for proteomics. *Proteomics* 2004; 4: 3665–85.
 - 51 Kondo T, Hirohashi S. Application of highly sensitive fluorescent dyes (CyDye DIGE Fluor saturation dyes) to laser microdissection and two-dimensional difference gel electrophoresis (2D-DIGE) for cancer proteomics. *Nat Protoc* 2007; 1: 2940–56.