

- importance of detection of recurring cytogenetic abnormalities and multilineage dysplasia on survival. *Am J Clin Pathol.* 2003;119:672–80.
12. Bloomfield CD, Shuma C, Regal L, et al. Long-term survival of patients with acute myeloid leukemia: a third follow-up of the Fourth International Workshop on Chromosomes in Leukemia. *Cancer.* 1997;80:2191–8.
 13. Schoch C, Schnittger S, Klaus M, Kern W, Hiddemann W, Haferlach T. AML with 11q23/MLL abnormalities as defined by the WHO classification: incidence, partner chromosomes, FAB subtype, age distribution, and prognostic impact in an unselected series of 1897 cytogenetically analyzed AML cases. *Blood.* 2003;102:2395–402.
 14. Dohner K, Tobis K, Ulrich R, et al. Prognostic significance of partial tandem duplications of the MLL gene in adult patients 16 to 60 years old with acute myeloid leukemia and normal cytogenetics: a study of the Acute Myeloid Leukemia Study Group Ulm. *J Clin Oncol.* 2002;20:3254–61.
 15. Byrd JC, Mrozek K, Dodge RK, et al. Pretreatment cytogenetic abnormalities are predictive of induction success, cumulative incidence of relapse, and overall survival in adult patients with de novo acute myeloid leukemia: results from Cancer and Leukemia Group B (CALGB 8461). *Blood.* 2002;100:4325–36.
 16. Mitterbauer-Hohendanner G, Mannhalter C. The biological and clinical significance of MLL abnormalities in haematological malignancies. *Eur J Clin Invest.* 2004;34:12–24.
 17. Miyawaki S, Sakamaki H, Ohtake S, et al. A randomized, postremission comparison of four courses of standard-dose consolidation therapy without maintenance therapy versus three courses of standard-dose consolidation with maintenance therapy in adults with acute myeloid leukemia: the Japan Adult Leukemia Study Group AML97 study. *Cancer.* 2005;104:2726–34.
 18. Miyazaki Y, Kuriyama K, Miyawaki S, et al. Cytogenetic heterogeneity of acute myeloid leukemia (AML) with trilineage dysplasia: Japan Adult Leukemia Study Group-AML 92 study. *Br J Haematol.* 2003;120:56–62.
 19. Matsuo T, Kuriyama K, Miyazaki Y, et al. The percentage of myeloperoxidase-positive blast cells is a strong independent prognostic factor in acute myeloid leukemia, even in the patients with normal karyotype. *Leukemia.* 2003;17:1538–43.
 20. Arber DA, Stein AS, Carter NH, Ilie D, Forman SJ, Slovak ML. Prognostic impact of acute myeloid leukemia classification. *Am J Clin Pathol.* 2003;119:672–80.
 21. Yanada M, Suzuki M, Kawashima K, et al. Long-term outcomes for unselected patients with acute myeloid leukemia categorized according to the World Health Organization classification: a single-center experience. *Eur J Haematol.* 2005;74:418–23.
 22. Bao L, Wang X, Ryder J, et al. Prospective study of 174 de novo acute myelogenous leukemias according to the WHO classification: subtypes, cytogenetic features and FLT3 mutations. *Eur J Haematol.* 2006;77:35–45.
 23. Mayer-Monard S, Parlier J, Passweg J, et al. Combination of broad molecular screening and cytogenetic analysis for genetic risk assignment and diagnosis in patients with acute leukemia. *Leukemia.* 2006;20:247–53.
 24. Olesen LH, Clausen N, Dimitrijevic A, Kerndrup G, Kjeldsen E, Hokland P. Prospective application of a multiplex reverse transcription-polymerase chain reaction assay for the detection of balanced translocations in leukaemia: a single-laboratory study of 390 paediatric and adult patients. *Br J Haematol.* 2004;127:59–66.
 25. Kuriyama K, Tomonaga M, Matsuo T, et al. Poor response to intensive chemotherapy in de novo acute myeloid leukaemia with trilineage myelodysplasia. *Br J Haematol.* 1994;86:767–73.
 26. Haferlach T, Schoch C, Loeffler H, et al. Morphologic dysplasia in de novo acute myeloid leukemia (AML) is related to unfavorable cytogenetics but has no independent prognostic relevance under the conditions of intensive induction therapy: results of a multi-parameter analysis from the German AML Cooperative Group studies. *J Clin Oncol.* 2003;21:256–65.
 27. Martineau M, Berger R, Lillington DM, Moorman AV, Secker-Walker LM. The t(6;11)(q27;q23) translocation in acute leukemia: a laboratory and clinical study of 30 cases. *Leukemia.* 1998;12:788–91.
 28. Dreyling MH, Schrader K, Fonatsch C, et al. MLL and CALM are fused to AF10 in morphologically distinct subsets of acute leukemia with translocation t(10;11): both rearrangements are associated with a poor prognosis. *Blood.* 1998;91:4662–7.
 29. Scholl C, Schlenk RF, Eiwen K, Dohner H, Frohlin S, Dohner K. The prognostic value of MLL-AF9 detection in patients with t(9;11)(p22;q23)-positive acute myeloid leukemia. *Haematologica.* 2005;90:1626–34.
 30. Mrozek K, Heinonen K, Lawrence D, et al. Adult patients with de novo acute myeloid leukemia and t(9;11)(p22;q23) have a superior outcome to patients with other translocations involving band 11q23: a Cancer and Leukemia Group B study. *Blood.* 1997;90:4532–8.
 31. Pasqualucci L, Liso A, Martelli PM, et al. Mutated nucleophosmin detects clonal multilineage involvement in acute myeloid leukemia: impact on WHO classification. *Blood.* 2006;108:4146–55.
 32. Choudhary C, Schwable J, Brandts C, et al. AML-associated Flt3 kinase domain mutations show signal translocation differences compared with Flt3 ITD mutations. *Blood.* 2005;106:265–73.
 33. Bienz M, Ludwig M, Mueller BÜ, et al. Risk assessment in patients with acute myeloid leukemia and a normal karyotype. *Clin Cancer Res.* 2005;11:1416–24.
 34. Suzuki T, Kiyoi H, Ozeki K, et al. Clinical characteristics and prognostic implications of NPM1 mutations in acute myeloid leukemia. *Blood.* 2005;106:2854–61.
 35. Ozeki K, Kiyoi H, Hirose Y, et al. Biologic and clinical significance of the FLT3 transcript level in acute myeloid leukemia. *Blood.* 2004;103:1901–8.