

MATERIALS AND METHODS

Patients

Sixty-one patients with IPMN who underwent surgery at the Department of Gastroenterological Surgery, Tohoku University Hospital, Sendai, Japan, from 1988 to 2006 were retrospectively analyzed. All patients agreed by written consent to the use of their anonymous records and tissues for scientific studies including this study. This study was approved by the Ethics Committee of Tohoku University School of Medicine. Surgical specimens from the patients were fixed in 10% buffered formalin solution and embedded in paraffin for histological analysis.

Immunohistochemistry

Immunohistochemical staining of MUC1-CORE, MUC2, and MUC5AC was performed as described previously.⁴

Statistical Analysis

Statistical analysis was performed on Statview 5.0 software (SAS Institute Inc, Cary, NC). Any probability values <0.05 were considered to be statistically significant.

RESULTS

Clinicopathological Features

Sixty-one patients with IPMNs operated upon consecutively at Tohoku University Hospital between 1988 and 2006 (18 years) were retrospectively analyzed. The patients included 45 men and 16 women with a mean age of 65.3 years (range, 48–79 years). Six patients underwent total pancreatectomy, 37 patients pancreatoduodenectomy, 17 patients distal pancreatectomy, and 1 patient segmental pancreatectomy. The macro-

scopic types of IPMNs were determined by clinical imaging studies and investigation of resected tumors, according to criteria described previously,⁵ as follows: main duct type, 24 tumors, with 14 uniformly dilated type and 10 focally dilated type; branch duct type, 37 tumors, with 31 cystic subbranch type and 6 dilated subbranch type. Histopathologic diagnoses of the resected tumors were made one of the authors (T.F.) according to criteria defined by the Japan Pancreas Society.¹ A diagnosis of intraductal papillary-mucinous adenoma (IPMA) was made for 26 tumors, intraductal papillary-mucinous carcinoma (IPMC) for 29 tumors, and invasive adenocarcinoma being derived from IPMN for 6 tumors, with 3 tubular adenocarcinomas and the remaining 3 mucinous colloid carcinomas. The macroscopic types were found to be significantly associated with histological diagnosis: 20 (83.3%) of 24 tumors with the main duct type and 21 (43.2%) of 37 tumors with the branch duct type were diagnosed as malignant ($P = 0.0019$, χ^2 test).

Types of IPMN

The resected tumors were classified into the 4 distinct histological types of IPMN (ie, gastric, intestinal, pancreatobiliary, and oncocytic types) by their morphology and immunohistochemical reactivity to MUC1-CORE, MUC2, and MUC5AC, according to criteria described previously.⁴ Of the 61 tumors, 27 (44.3%) were classified into the gastric type, 29 (47.5%) into the intestinal type, 4 (6.6%) into the pancreatobiliary type, and 1 (1.6%) into the oncocytic type (Fig. 1).

Association Between the Types of IPMN and Clinicopathological Features

The types of IPMN were significantly associated with the histological diagnoses ($P < 0.0001$, χ^2 test) (Table 1): 22




FIGURE 1. Representative histological images of the types of IPMNs: (A) gastric, (B) intestinal, (C) pancreatobiliary, and (D) oncocytic.

TABLE 1. Types of IPMNs and Clinicopathological Features

Type	n	Sex (Male/Female)	Mean Age (yrs)	Mean Survival Period* (mo)	Diagnosis				Macroscopic Type	
					IPMA	IPMC	INV-C	INV-T	Branch	Main
Gastric	27	23:4	66.0	ND†	22	4	0	1	22	5
Intestinal	29	18:11	64.1	52.1	4	21	3	1	11	18
Pancreatobiliary	4	3:1	70.0	ND†	0	3	0	1	3	1
Oncocytic	1	1:0	55.0	ND†	0	1	0	0	1	0
Total	61	45:16	65.3	59.8	26	29	3	3	37	24

*Mean survival period by Kaplan-Meier analysis.

†Not determined because all subjects were censored.

INV-C indicates invasive colloid mucinous adenocarcinoma; INV-T, invasive tubular adenocarcinoma; ND, not determined.

of 27 cases of the gastric type were diagnosed as benign (ie, IPMA), whereas 25 of 29 cases of the intestinal type were diagnosed as malignant, with 21 IPMC, 3 invasive mucinous colloid adenocarcinomas being derived from IPMC, and 1 invasive tubular adenocarcinoma being derived from IPMC. Five gastric-type IPMNs were diagnosed as malignant (4 IPMC and 1 invasive tubular adenocarcinoma being derived from IPMC). The gastric-type IPMCs showed relatively low papillae with severe/high-grade atypia. They diffusely expressed MUC5AC but hardly MUC1 and MUC2 (Fig. 2). The intestinal type IPMAs corresponded to the goblet type, a variation of the intestinal type⁴ (Fig. 2).

Four pancreatobiliary-type IPMNs were diagnosed as malignant, including one with invasive tubular adenocarcinoma. One oncocytic type IPMN was diagnosed as IPMC.

Compared with the macroscopic types of IPMN, 22 of the 27 gastric-type IPMNs corresponded to the branch duct type, whereas 18 of the 29 intestinal-type IPMNs to the main duct type, which indicated that the types were closely associated with the macroscopic type ($P = 0.0068$, χ^2 test) (Table 1). Furthermore, the types were associated with the diagnosis in each macroscopic type: 8 of 11 branch duct-type IPMNs of the intestinal type and 4 of 22 of the gastric-type

IPMNs were malignant, whereas 1 of 18 main duct-type IPMNs of the intestinal type and 3 of 5 of the gastric-type IPMNs were benign ($P = 0.0020$ for the former and $P = 0.0045$ for the latter, χ^2 test). The oncocytic-type IPMN belonged to the branch duct type. Of 4 pancreatobiliary-type IPMNs, 3 belonged to the branch duct type and 1 to the main duct type.

Invasion was not associated with any specific type of IPMN.

Survival of IPMN

The patients were followed up for their prognosis. The mean follow-up period was 38.1 months (range, 1–176 months). Survival analyses were carried out according to the Kaplan-Meier method. The types of IPMN were associated with survival (Fig. 3A). Patients with intestinal-type IPMNs had a poorer survival than those with gastric-type IPMNs. The 5-year relative survival rates were 62.3% for patients with the intestinal types, whereas 100% for those with the gastric types. The mean survival period for patients with the intestinal-type IPMNs was 52.1 months. A P value could not be calculated because all patients with the gastric-type IPMN were censored. Nine of 29 patients with intestinal-type IPMNs died of the




FIGURE 2. Examples of gastric-type IPMC (A–D) and intestinal-type IPMA (E–H). Hematoxylin-eosin staining (A, E) and immunohistochemical staining of MUC1-CORE (B, F), MUC2 (C, G), and MUC5AC (D, H).




FIGURE 3. Kaplan-Meier survival analysis of patients with gastric- (dashed line) and intestinal-type (solid line) IPMNs: (A) overall survival, (B) survival of patients without invasive carcinoma.

disease during follow-up. All these patients had been diagnosed as having IPMC, including 3 with invasive mucinous colloid carcinomas and 1 with invasive tubular adenocarcinoma. Without the patients with invasive carcinomas, patients with intestinal-type IPMNs still had a poorer survival than those with gastric-type IPMNs (Fig. 3B). Of 4 patients with the pancreatobiliary-type IPMNs, 1 patient died of complications related to the operation and the other 3 patients were alive for 15, 8, and 3 months, respectively, until each observation period was censored. The patient with the oncocytic-type IPMN was alive for 18 months until the observation period was censored.

Diagnosis and the macroscopic types were also associated with survival. According to the diagnosis, the 5-year relative survival rates were 100% for patients with IPMA, 51.9% for patients with IPMC, and 0% for those with invasive adenocarcinoma. A *P* value could not be calculated because all patients with IPMA were censored. The mean survival period was 55.5 months for patients with IPMC, 20.3 months for patients with invasive mucinous colloid carcinoma, and 5.0 months for patients with invasive tubular adenocarcinoma. The patients with the main duct type revealed significantly poorer survival than those with the branch duct type ($P = 0.0376$ by log-rank test). The mean survival periods were 52.2 months for the main duct type and 64.2 months for the branch duct type. Even in each of the macroscopic type, the histological type seemed to be associated with survival: 3 of 11 patients with the branch duct-type IPMN of the intestinal type died, whereas all 5 patients with the main duct-type IPMN of the gastric type survived.

Sex, age, and operative procedures were not associated with survival.

DISCUSSION

The present retrospective study indicated that the histological types of IPMN are associated with histological diagnoses, macroscopic types, and survival of patients with IPMN. Characteristically, the gastric-type IPMN was likely to be diagnosed as benign, to reveal a fair prognosis, and to be confined to a branch duct. On the other hand, the intestinal-type IPMN was likely to be diagnosed as malignant, to reveal a poor prognosis, and to occupy the main duct. Because the pancreatobiliary and oncocytic types were rare in our series,

only 4 cases and 1 case, respectively, we could not estimate any of their clinicopathological characteristics.

Prognoses of the gastric-type IPMNs were fairly good; no patient died of the disease, which indicates that the gastric-type IPMN is less aggressive clinically. Consistently, 22 (81.5%) of the 27 gastric type cases in our series were diagnosed as benign. The remaining 5 cases were diagnosed as malignant, with 4 IPMC and 1 IPMC with invasive tubular adenocarcinoma. These patients were followed up for less than 18 months until being censored in this study; therefore, the long-term prognoses of them have not yet been determined.

Patients with intestinal-type IPMNs had poor prognoses. Consistently, the tumors were associated with malignant diagnosis (ie, IPMC or IPMC with invasion). The intestinal IPMCs accompanied invasive adenocarcinoma more frequently than IPMNs of other types, which certainly worsened the prognosis. Some of the intestinal-type IPMCs without invasion also led to death of the patients, which indicates that the intestinal-type IPMC even without invasion is clinically aggressive and may require careful follow-up or an additional intervention rather than operation only. Adsay et al⁶ indicated that the intestinal-type IPMN was associated with mucinous colloid carcinoma that revealed a better prognosis than IPMNs with invasive ductal adenocarcinoma. Although 3 of 4 invasive intestinal-type IPMCs in our series accompanied mucinous colloid carcinoma in their invasive component, we could not statistically evaluate the trend because of the small number of invasive cases.

The gastric-type IPMNs were associated with the branch duct-type IPMN, whereas the intestinal-type IPMNs were associated with the main duct type. Diagnoses were also associated with the macroscopic types. These results indicate that the histological type, macroscopic type, and diagnosis are strongly associated with each other. Patients with the branch duct-type IPMN are known to have a better prognosis than those with the main duct-type IPMNs, which was consistent with the current analysis.⁷ Even more notably, the branch duct-type IPMN of the intestinal type was more malignant than that of the gastric type. The main duct-type IPMN of the gastric type was more benign than that of the intestinal type. Our results indicate that we can estimate that a branch duct-type IPMN is likely to be a gastric-type IPMA, whereas a main duct-type IPMN is likely to be an intestinal-type IPMC.

In conclusion, evaluation of the histological types of IPMN may help to predict the clinical course of patients with IPMN and to design better clinical management of these patients.

ACKNOWLEDGMENT

The authors thank Ms Naomi K. Ono for technical assistance and Ms Barbara Levene for editing this manuscript.

REFERENCES

1. Kawarada Y, Japan Pancreas Society. *Classification of Pancreatic Carcinoma*, 2nd ed. Tokyo: Kanehara & Co, Ltd; 2003.
2. Hruban RH, Takaori K, Klimstra DS, et al. An illustrated consensus on the classification of pancreatic intraepithelial neoplasia and intraductal papillary mucinous neoplasms. *Am J Surg Pathol*. 2004;28:977-987.
3. Ohhashi K, Murakami Y, Maruyama M, et al. Four cases of mucous secreting pancreatic cancer. *Prog Digest Endosc*. 1982;20:348-351.
4. Furukawa T, Kloppel G, Volkan Adsay N, et al. Classification of types of intraductal papillary-mucinous neoplasm of the pancreas: a consensus study. *Virchows Arch*. 2005;447:794-799.
5. Furukawa T, Takahashi T, Kobari M, et al. The mucus-hypersecreting tumor of the pancreas. Development and extension visualized by three-dimensional computerized mapping. *Cancer*. 1992;70:1505-1513.
6. Adsay NV, Merati K, Andea A, et al. The dichotomy in the preinvasive neoplasia to invasive carcinoma sequence in the pancreas: differential expression of MUC1 and MUC2 supports the existence of two separate pathways of carcinogenesis. *Mod Pathol*. 2002;15:1087-1095.
7. Kobari M, Egawa S, Shibuya K, et al. Intraductal papillary mucinous tumors of the pancreas comprise 2 clinical subtypes: differences in clinical characteristics and surgical management. *Arch Surg*. 1999;134:1131-1136.

Locally Advanced Intrahepatic Cholangiocarcinoma Successfully Resected after Transcatheter Arterial Chemoembolization with Degradable Starch Microspheres: Report of a Case

Yongyou Wu^{1,3}, Akio Saiura¹, Junji Yamamoto¹, Rintaro Koga¹
Shingo Asahara², Akira Kamei², Koichi Takano², Takaaki Ikari², Makoto Seki¹
Toshiharu Yamaguchi¹, Tetsuichiro Muto¹

¹Department of Gastrointestinal Surgery, Cancer Institute Hospital, and ²Department of Internal Medicine Cancer Institute Hospital, Tokyo, Japan; ³Department of General Surgery 2nd Hospital of Soochow University, Suzhou, China

Corresponding Author: Akio Saiura, MD, Department of Gastrointestinal Surgery
Cancer Institute Ariake Hospital, 3-10-6 Ariake, Koto-ku, Tokyo 135-8500, Japan
Tel: +81 3 3520 0111, Fax: +81 3 3570 0343, E-mail: saiura-tky@umin.ac.jp

SUMMARY

We describe a case of initially unresectable locally advanced intrahepatic cholangiocarcinoma that showed remarkable regression after transcatheter arterial chemoembolization with degradable starch microspheres, allowing for subsequent successful curative resection. A 75-year-old female was referred to our hospital with a large hepatic mass. Computerized tomography examination showed a huge mass in the right liver extended partially to the left liver. Intrahepatic cholangiocarcinoma was strongly suspected, but surgical resection was abandoned due to the local spread in the liver. Three courses of transcatheter arterial chemoembolization with degradable starch microspheres were performed. The anti-

cancer agents, mitomycin C and epirubicin, combined with degradable starch microspheres were injected from the catheter for chemoembolization. After three courses of transcatheter arterial chemoembolization, the tumor size decreased from 10cm to 5.5cm in diameter. Then right trisegmentectomy together with extra-hepatic bile duct excision was performed. At 25 months after the first therapy and 21 months after operation, the patient remains healthy without recurrence. Transcatheter arterial chemotherapy with degradable starch microspheres may be a treatment of choice with locally advanced intrahepatic cholangiocarcinoma.

KEY WORDS:

Degradable starch Microspheres; Intrahepatic cholangiocarcinoma; Liver resection

ABBREVIATIONS:

Intrahepatic Cholangiocarcinoma (ICC); Degradable Starch Microspheres (DSM); Transcatheter Arterial Chemoembolization (TACE)

INTRODUCTION

Intrahepatic cholangiocarcinoma (ICC) accounts for 3-5% of all primary liver malignancies (1,2). The incidence is increasing and its pathogenesis remains unclear (3-5). Currently, complete surgical resection with histologically negative margins is the only cure for this formidable entity and yields a five-year survival of 13-42% (2,6). By contrast, in patients not electing surgical therapy, the prognosis is extremely poor, with a median survival of less than 1 year (7,8). Unfortunately, the overall resectability rate of ICC is unsatisfactory, ranging from 19-70% (2,9). Noncurative operation offers no survival benefit, with a median survival of only approximately 3 months (10). Thus, increasing the rate of curative resection is of vital importance for patients with ICC.

In this paper, we report a case of initially unresectable locally advanced ICC that showed remarkable




FIGURE 1 A 75-year-old female with cholangiocarcinoma. Intravenous contrast-enhanced CT scan of the upper abdomen demonstrating a huge solid mass, 10.0x9.0cm in size with peripheral enhanced areas in the right lobe of the liver. Involvement of the umbilical portion and slight dilation of the bile duct in Segment 2 is visible (arrow).



FIGURE 2 Intravenous contrast-enhanced CT scan showing lipiodol accumulation in the tumor and decreased tumor size 1 week after the third chemoembolization.



FIGURE 3 Raw surface of the liver after right trisegmentectomy of the liver together with the extra-hepatic bile duct excision. The surgical margin was free from cancer.

regression after transcatheter arterial chemoembolization with degradable starch microspheres (DSM-TACE), allowing for subsequent successful curative resection.

CASE REPORT

A 75-year-old female was referred to our hospital with right-sided abdominal pain, liver dysfunction, and a large hepatic mass. Computerized tomography (CT) examination showed a solid mass 10cm in diameter with peripheral enhancement in the right liver. The tumor extended partially to the left liver, adjacent to the umbilical portion of the liver. CT scan also disclosed slight dilation of the bile duct in the left lateral segment of the liver (**Figure 1**). Gastrointestinal studies revealed no other abnormal findings. Serum alkaline phosphatase (ALP), lactate dehydrogenase (LDH), and CA19-9 were elevated at 1286 U/L, 2774 U/L, and 430.7 U/mL, respectively. ICC was strongly suspected, but surgical resection was deferred secondary to local spread in the liver.

Three courses of transcatheter arterial chemoembolization with degradable starch microspheres (DSM-TACE) were performed. On December 17, 2003, the first DSM-TACE treatment was undertaken. Arteriography of the hepatic artery was performed to char-

acterize the vascular supply of the lesion. The anti-cancer agents, mitomycin C (6mg) and epirubicin (60mg), combined with DSM (600mg), were injected from the catheter for chemoembolization. The right hepatic artery and left hepatic artery were embolized at a ratio of 3:1. On January 26, 2004, DSM-TACE was repeated using the same protocol with different doses of the chemo-embolic agents (DSM, 300mg; epirubicin, 50mg; mitomycin C, 6mg). The third course was performed on February 17, 2004, again using DSM (450mg), mitomycin C (6mg) and epirubicin (55mg) for chemoembolization. In this last treatment, the artery supplying the left medial segment of the liver was embolized. No major complications or side effects were noted during or after each DSM-TACE session.

After three courses of DSM-TACE, the serum level of CA19-9 decreased from 430.7 U/mL to 25.8 U/mL, and the tumor size decreased from 10cm to 5.5cm in diameter (**Figure 2**). Surgical resection was then elected. Right trisegmentectomy together with extra-hepatic bile duct excision was performed on April 16, 2004 (**Figure 3**). Postoperative pathology confirmed a margin negative resection and a moderately differentiated intrahepatic cholangiocarcinoma without extra-hepatic lymphatic metastasis (**Figure 4**).

Postoperative recovery was excellent without complications. At 21 months after the first DSM-TACE therapy and 25 months after operation, the patient remains healthy and tumor free with normal serum tumor marker levels.

DISCUSSION

Treatments for unresectable ICC include chemotherapy, radiotherapy, photodynamic therapy, radio-frequency thermal ablation, and liver transplantation (11). Although some investigators have reported an improved prognosis following such treatment, the benefits tend to be limited, with poor life expectancy, and outcomes have not been verified via prospective and randomized clinical studies. Thus, improving the resectability rate of these tumors is of great significance for patient outcomes.



FIGURE 4 Macroscopic findings of the resected specimen. A 7.5x5.0-cm intrahepatic cholangiocarcinoma was located in the hepatic hilum with an adequate surgical margin.

In the present case, a margin-negative resection was initially considered impossible due to the local advancement to the umbilical portion of the liver. Degradable starch microspheres (DSM) are 40-45 microns in diameter and can be degraded by serum amylase within 30 to 60 minutes. When a mixture of DSM and chemotherapeutic agents are applied via transcatheter arterial infusion, transient occlusion of the arterial blood flow can enhance regional retention of the anti-tumor agents and reduce systematic toxicity. Furthermore, transient vascular occlusion induced by DSM can lead to the redistribution of blood flow to hypovascular areas and thereby potentiate the chemotherapeutic effects on ICC, which is typically a hypovascular tumor (12). Moreover, repeated administrations are possible, and DSM-TACE is well tolerated even in cases with portal vein trunk obstruction.

The use of DSM-TACE for treatment of unresectable hepatocellular carcinoma (HCC) and metastatic liver diseases is increasing secondary to improved patient outcomes. For example, the response rate is 52.9-72.7% for patients with HCC, 42.8% for patients with metastatic gastric cancer, and 37.8-70% for patients with colon cancer, with complete responses

observed in some cases (13-16). However, there are comparatively few published reports regarding the efficacy of DSM-TACE in patients with unresectable ICC. In 1989, Hirai *et al.* described a series of 11 cases of HCC and one case of ICC treated by DSM-TACE. While 72.7% of the HCC cases experienced a decrease in tumor size in response to treatment, there was no notable change in the case with ICC (14). By contrast, in the present report, tumor size decreased by 70% after DSM-TACE, thereby enabling curative resection.

In conclusion, unresectable ICC is associated with extremely poor prognosis, however, DSM-TACE may be of utility in such cases in establishing local control, improving quality of life, downstaging the disease, and even facilitating curative resection in some cases. DSM-TACE is considered to be a treatment of choice in patients with locally advanced ICC.

ACKNOWLEDGEMENTS

This study was supported by the Grants-in-Aid for basic research from the Ministry of Health and Welfare of Japan.

REFERENCES

- Ikai I, Arii S, Ichida T, Okita K, Omata M, Kojiro M, Takayasu K, et al: Report of the 16th follow-up survey of primary liver cancer. *Hepatol Res* 2005; 32:163-172.
- Morimoto Y, Tanaka Y, Ito T, Nakahara M, Nakaba H, Nishida T, Fujikawa M, et al: Long-term survival and prognostic factors in the surgical treatment for intrahepatic cholangiocarcinoma. *J Hepatobiliary Pancreat Surg* 2003; 10:432-440.
- Khan SA, Taylor-Robinson SD, Toledano MB, Beck A, Elliott P, Thomas HC: Changing international trends in mortality rates for liver, biliary and pancreatic tumours. *J Hepatol* 2002; 37:806-813.
- Patel T: Increasing incidence and mortality of primary intrahepatic cholangiocarcinoma in the United States. *Hepatology* 2001; 33:1353-1357.
- Patel T: Worldwide trends in mortality from biliary tract malignancies. *BMC Cancer* 2002; 2:10.
- Nakeeb A, Pitt HA, Sohn TA, Coleman J, Abrams RA, Piantadosi S, Hruban RH, et al: Cholangiocarcinoma. A spectrum of intrahepatic, perihilar, and distal tumors. *Ann Surg* 1996; 224:463-473; discussion 473-465.
- Anderson CD, Pinson CW, Berlin J, Chari RS: Diagnosis and treatment of cholangiocarcinoma. *Oncologist* 2004; 9:43-57.
- Carriaga MT, Henson DE: Liver, gallbladder, extrahepatic bile ducts, and pancreas. *Cancer* 1995; 75:171-190.
- Suzuki S, Sakaguchi T, Yokoi Y, Okamoto K, Kurachi K, Tsuchiya Y, Okumura T, et al: Clinicopathological prognostic factors and impact of surgical treatment of mass-forming intrahepatic cholangiocarcinoma. *World J Surg* 2002; 26:687-693.
- Chu KM, Fan ST: Intrahepatic cholangiocarcinoma in Hong Kong. *J Hepatobiliary Pancreat Surg* 1999; 6:149-153.
- Shirabe K, Shimada M, Harimoto N, Sugimachi K, Yamashita Y, Tsujita E, Aishima S: Intrahepatic cholangiocarcinoma: its mode of spreading and therapeutic modalities. *Surgery* 2002; 131:S159-164.
- Chang D, Jenkins SA, Grime SJ, Nott DM, Cooke T: Increasing hepatic arterial flow to hypovascular hepatic tumours using degradable starch microspheres. *Br J Cancer* 1996; 73:961-965.
- Nakajima T, Nozu S, Saitou AY, Okayasu K, Majima T, Yagi A: [Clinical effectiveness of intra-arterial chemotherapy combined with degradable starch microspheres (DSM) for liver metastases and prediction of effectiveness by diagnostic imaging]. *Gan To Kagaku Ryoho* 2001; 28:1489-1492.
- Hirai K, Yamashita K, Aoki Y, Sakai T, Abe M, Tanikawa K, Noguchi H, et al: [An assessment of therapeutic effects by transcatheter arterial embolization (TAE) with degradable starch microspheres (DSM) for hepatic malignant tumors]. *Nippon Gan Chiryō Gakkai Shi* 1989; 24:772-780.
- Pohlen U, Mansmann U, Berger G, Germer CT, Galikowski U, Boese-Landgraf J, Buhr HJ: Multicenter pilot study of 5-fluorouracil, folinic acid, interferon alpha-2b and degradable starch microspheres via hepatic arterial infusion in patients with nonresectable colorectal liver metastases. *Anticancer Res* 2004; 24:3275-3282.
- Furuse J, Ishii H, Satake M, Onaya H, Nose H, Mikami S, Sakai H, et al: Pilot study of transcatheter arterial chemoembolization with degradable starch microspheres in patients with hepatocellular carcinoma. *Am J Clin Oncol* 2003; 26:159-164.

CT and MRI Findings with Contrast Enhancement of Small Pancreatic Adenocarcinoma in the Late Phase

Susumu Hijioka, Takaaki Ikari, Akira Kamei, Koichi Takano, Shingo Asahara
Naoya Fujita, Miyuki Shimizu, Junji Yamamoto, Rikiya Fujita, Keiko Sasaki

Department of Gastroenterology, The Cancer Institute Ariake Hospital, Tokyo, Japan

Corresponding Author: Susumu Hijioka, MD, Department of Internal Medicine
Kamiamakusa General Hospital, 1419-19 Takado Ryugatake-machi Kamiamakusa-si
Kumamoto 866-029,3 Japan

Tel: +81 969 62 1122, Fax: +81 969 62 1546, E-mail: rizasusu@isis.ocn.ne.jp

SUMMARY

A 60-year-old female was found to have high serum amylase concentrations at a medical check-up. Dynamic computed tomography and magnetic resonance imaging demonstrated a mass in the body of the pancreas, which was enhanced in the late phase of the scans by administration of a contrast medium. Endoscopic retrograde pancreatography showed a stenosis of the main pancreatic duct at the body, and brushing cytology from the region revealed adenocarcinoma. Distal pancreatectomy was performed. The tumor was a well-differentiated adenocar-

cinoma, measuring 15x10mm. Fibrous tissues were sparsely distributed in the tumor, and there was an increase of dilated veins, in particular at the margin. Late-phase enhancement of the tumor with computed tomography or magnetic resonance imaging was considered to be correlated with this abundant vascular structure in the tumor. Marked tumor enhancement in the late phase might be a characteristic finding suggesting an early-stage pancreatic adenocarcinoma, which should be carefully checked.

KEY WORDS:

Pancreatic carcinoma;
Computed tomography;
Pancreatic neoplasm; Late phase

ABBREVIATIONS:

Computed Tomography (CT);
Magnetic Resonance Imaging (MRI);
Endoscopic Retrograde Pancreatography (ERP);
Ultrasonography (US);
Multidetector Computed Tomography (MDCT)

INTRODUCTION

Despite the advent of sophisticated imaging modalities, a majority of pancreatic ductal adenocarcinomas are diagnosed at stages IVA or IVB. The five-year survival rate of pancreatic ductal adenocarcinomas is still less than 5% (1,2), and the median survival time for unresectable cases ranges from 4 to 8 months. For a better prognosis, a tumor should be detected at an early stage and excised curatively.

In pancreatic adenocarcinoma, which is usually hypovascular, computed tomography (CT) shows a low-attenuation mass in the early pancreatic phase after infusion of the contrast medium. Magnetic resonance imaging (MRI) demonstrates similar findings. However, few detailed reports have been published concerning CT or MRI findings in small pancreatic adenocarcinomas. Our case demonstrated a marked enhancement of a pancreatic tumor in the late phase. Late-phase enhancement of small pancreatic carcinoma has not been studied previously in terms of radiological and pathological correlation. We present here a case of small pancreatic adenocarcinoma for which we studied the mechanism of marked late-phase enhancement. Description of the tumor extent is made based on the sixth edition of the TNM Classification (UICC) (3).

CASE REPORT

A 60-year-old female was found to have high amy-

lase blood concentrations at a medical check-up in April 2004 and visited a clinic. Abdominal ultrasonography (US) and CT revealed dilatation of the main pancreatic duct at the pancreatic body and tail, and she was referred to our hospital. Biochemistry showed no abnormality, except high levels of serum and urinary amylase. Tumor markers were all within the normal range, except elastase I.

US at 3.5MHz frequency demonstrated a tumor measuring 9x8mm at the pancreatic body with dilatation of the pancreatic duct in the tail (Figure 1a). US at 7.5MHz revealed that the tumor had an ill-defined




FIGURE 1 US at 3.5MHz frequency (a) demonstrated a low-echoic mass measuring 9x8mm at the pancreatic body (arrow) with dilatation of the pancreatic duct in the tail. US at 7.5MHz (b) revealed that the tumor had an ill-defined margin and high echoic spots (arrowhead), probably reflecting a duct structure within the mass.



FIGURE 2 Multidetector CT shows a hypoenhancing mass at the pancreatic phase 40 sec after contrast administration (a). However, it showed homogeneous contrast enhancement at the portal phase, 80 sec after contrast administration (b), and the enhancement lasted until the late phase, 180 sec after contrast administration (c).



FIGURE 3 Dynamic MRI did not show the tumor 20 sec after infusion of the contrast medium (a), but the whole tumor was revealed by contrast enhancement 80 sec after the infusion (b), and the enhancement lasted until 180 sec (c).

margin and an isogenic internal echo with high echoic spots (**Figure 1b**). Multidetector CT (MDCT) with four detectors was performed using 100mL of a contrast medium at a concentration of 370mg/dL. The contrast medium was intravenously infused at a rate of 3mL/sec. Scans were started at 40, 80 and 180 sec after initiation of contrast administration. Images were taken at 1.25-mm collimation and reconstructed with a width of 0.5mm. The tumor was visualized as a hypo-enhancing mass (162HU) at the pancreatic phase 40 sec after contrast administration (**Figure 2a**), but it showed homogeneous contrast enhancement (204HU) at the portal phase, 80 sec after contrast administration, and the enhancement lasted (164HU) until the late phase, 180 sec after contrast administration (**Figure 2c**).

In MRI, both T1- and T2-weighted images demonstrated dilatation of the pancreatic duct, but did not show a tumor. Dynamic MRI was performed using gadrinium DTPA; the tumor was not visualized 20 sec after infusion of the contrast medium (**Figure 3a**), but the whole tumor was revealed by contrast enhancement 80 sec after the infusion (**Figure 3b**), and the enhancement lasted until 180 sec (**Figure 3c**).

Endoscopic retrograde pancreatography (ERP) showed a stricture of the main pancreatic duct at the pancreatic body, and brushing cytology from the region revealed atypical cells characterized by enlargement, size heterogeneity, increased chromatin content of the nuclei, enlargement of nucleoli, and mucus in the cytoplasm, leading to the diagnosis of adenocarcinoma. Although the mass was enhanced in the late phase of both CT and MRI, the tumor was diagnosed as a resectable pancreatic adenocarcinoma and distal pancreatectomy was performed.

Cut surfaces of resected specimens revealed that the tumor was slightly whitish and ill demarcated (**Figure 4a**). The tumor was 15x10mm in size. The low-magnified view is shown in **Figure 4b**. There was exfoliation of the epithelium in the main pancreatic duct, possibly due to the postoperative catheterization of the duct by a thin catheter, which was performed to obtain a pancreatogram for comparison with ERP. A histological examination showed a well-differentiated adenocarcinoma infiltrating the pancreatic parenchyma with tubular and glandular formations (**Figure 4c**). Fibrous tissues were sparsely distributed in the stroma. In addition, there was an increase in markedly dilated veins in the tumor, in particular at the margin of the tumor (**Figure 4d**).

The final pathological diagnosis was as follows: Invasive ductal adenocarcinoma of the pancreas; well differentiated tubular adenocarcinoma, T1, intermediate type, INF β , ly1, v0, N0 Stage I.

DISCUSSION

Despite the advent of sophisticated imaging methods, the five-year survival rate for patients with ductal adenocarcinomas has been less than 5% for decades (1,2), and even in resected cases the rate has been approximately 10~20% (4-8). The prognosis of pancreatic adenocarcinomas is poorer among malignant tumors because they are likely to extend beyond pancreatic ductal walls into parenchyma and invade vascular and lymphatic structures or perineural spaces. In pancreas, there is no barrier, such as muscle layer around pancreatic ductal walls, which probably could facilitate cancer invasion at an earlier stage (9-13). However, patients with respectable small carcinoma less than 2cm in size have the favorable survival of 30-40% at five years (8,14), and the diagnosis of as small

carcinoma as possible should be made for better prognosis.

CT and MRI are imaging modalities useful for the diagnosis of pancreatic cancer. So far, CT and MRI have been effective methods to stage pancreatic adenocarcinoma, but not for detection of small pancreatic tumors because these methods provide poorer resolution than does US. However, recent development of multi-detector CT remarkably has improved spatial resolution, and it has become possible to diagnose small pancreatic tumors.

Since ductal adenocarcinoma of the pancreas shows a poorly demarcated cut surface, hard consistency with fibrous stroma and poor vascular structures histologically, it is visualized by dynamic CT in the pancreatic phase as a hypoenhancing mass relative to the normal pancreas (15), 30-40 sec after administration of contrast medium, when the normal parenchyma is best enhanced. In the late phase, the contrast medium begins to wash out of the normal tissue and infiltrate into the fibrous stroma of the tumor, thus providing mild enhancement.

Lu, Vedantham, Boland, Furukawa *et al.* reported that visualization of pancreatic adenocarcinoma was significantly higher in the pancreatic phase, when the tumor-pancreas contrast difference was maximal, than in the portal phase (15-18). In general practice, detection of a low-attenuation area in the pancreatic phase is thought to be an important sign in the diagnosis of pancreatic adenocarcinoma.

However, in the present case, dynamic CT in the pancreatic phase not only demonstrated a slightly lower attenuation area than the adjacent pancreatic parenchyma, but showed remarkable tumor staining in the late phase. MRI showed similar late-phase enhancement of the tumor. As mentioned previously, tumor staining in the late phase is thought to derive from the abundant fibrous stroma of the pancreatic adenocarcinoma, but pathological findings in the present case revealed mild fibrosis with abundant fibroblasts and a limited number of hyalinized collagen fibers. Although venous invasion was negative, the number of dilated veins was increased, particularly at the tumor margin. Although the mechanism underlying the fibrosis of pancreatic adenocarcinomas has not been understood fully, it can be hypothesized that




FIGURE 4 Cut surfaces of resected specimens revealed a slightly whitish and ill demarcated tumor. The tumor was 15x10mm in size (a). The low-magnified view (b) revealed the main pancreatic duct (arrow) and branch ducts (arrowheads). A histological examination showed a well-differentiated adenocarcinoma infiltrating the pancreatic parenchyma with tubular and glandular formations (c). There was an increase in the number of dilated veins (arrows) in the tumor, particularly at the margin of the tumor (d).

after an early stage of fibrosis, which possibly is associated with an increased number of dilated veins, stenosis occurs in these veins in association with hyalinized fibrotic changes and tumor infiltration.

In the present case, CT and MRI demonstrated tumor enhancement in the late phase, which seemed to reflect an increased number of dilated veins, as is often recognized in hemangioma, and the remarkable tumor staining in the late phase might be a characteristic CT or MRI finding suggestive of relatively early-stage pancreatic adenocarcinomas. Indeed, Yamaguchi *et al.* (19) reported that tumor staining was observed in the late phase by CT in 70% of pancreatic cancer 20mm or smaller in diameter, compared with 20% of pancreatic tumors 21mm or larger. Therefore, it is important to pay attention to tumor staining in the late phase, and not to overlook a low-attenuation area in the pancreatic phase, for diagnosis of small pancreatic cancer.

REFERENCES

- 1 Faivre J, Foran D, Esteve J, Obradovic M, Sant M, EURO CARE working Group: Survival of patients with primary liver cancer, pancreatic cancer and biliary tract cancer in Europe. *Eur J Cancer* 1998; 34:2184-2190.
- 2 Greenlee RT, Murray T, Bolden S, Wingo PA: cancer statistics 2000. *CA Cancer J Clin* 2000; 50:7-33.
- 3 Sobin LH, Wittekind Ch: TNM classification of Malignant Tumours Sixth Ed. International Union Against Cancer (UICC). New York: Wiley-Liss 2002; pp. 93-96.
- 4 Sperti C, Pasquali C, Piccoli A, Pedrazzoli S: Survival after resection for ductal adenocarcinoma of the pancreas. *Br J Surg* 1996; 83:625-631.
- 5 Yeo CJ, Sohn TA, Cameron JL, Hruban RH, Lillemoe KD, Pitt HA.: periaampullary adenocarcinoma: analysis of 5-year survivors. *Ann Surg* 1998; 227:821-831.
- 6 Sohn TA, Yeo CJ, Cameron JL, Koniaris L, Kausbal S, Abrams RA, et al: Resected adenocarcinoma of the pancreas- 616 patients: results, outcomes, and prognostic indicators. *J Gastrointest Surg* 2000; 4:567-579.
- 7 Sean PC, Robert G, Maha G, Paul Greig, Lloyd Smith, Robert Mackenzie, et al: Prognosis factors in resected pancreatic adenocarcinoma: Analysis of actual 5-year survivors. *J Am Coll Surg* 2004; 198:722-731.
- 8 Desiree P, Ilaria R, Gabriella N: Improved survival in small pancreatic cancer. *Dig Surg* 2001; 18:41-46.
- 9 Matsukuma S, Suda K: Small pancreatic tubular adenocarcinomas: Clinicopathological analysis with immunohistochemical and histochemical evaluation. *Pathol Int* 1996; 46:581-588.
- 10 Matsuo S, Sato T: Surgical treatment for carcinoma of the

- pancreas. *Am. J. Surg* 1986; 152:499-504.
- 11 **Tsuchiya R, Tomioka T, Izawa K, Noda T, Yamamoto K, Tsukada T, et al:** Collective review of small carcinoma of the pancreas. *Ann Surg* 1986; 203:77-81.
 - 12 **Manabe T, Miyashita T, Ohshiro G, Nonaka A, Suzuki T, Endo K, et al:** Small carcinoma of the pancreas: Clinical pathologic evaluation of 17 patients. *Cancer* 1988; 62:135-141.
 - 13 **Satake K, Nishiwaki H, Yokomatsu H, Kawazoe Y, Kim K, Haku A, et al:** Surgical curability and prognosis for standard versus extended resection for T1 carcinoma of the pancreas. *Surg. Gynecol. Obstet* 1992; 175:259-265.
 - 14 **Egawa S, Takeda K, Fukuyama N, Abe N, Motoi F, Sunamura M, et al:** Staging and outcome of patients with pancreatic cancer through the national pancreatic cancer registry. *Kantansui* 2004; 48:115-125. (In Japanese with English abstract)
 - 15 **Lu DS, Vedantham S, Krasny RM, Kadell B, Berger WL, Reber HA:** Two-phase helical CT for pancreatic tumors: pancreatic versus hepatic phase encasement of tumor, pancreas, and vascular structures. *Radiology* 1996; 199:697-701.
 - 16 **Vedantham S, Lu DS, Reber HA, Kadell B:** Small peri-pancreatic veins; improved assessment in pancreatic cancer patients using thin-section pancreatic phase helical CT. *AJR* 1998; 170:377-383.
 - 17 **Boland GW, Malley ME, Saez M, Castillo CF, Warshaw AL, Mueller PR:** Pancreatic-phase versus portal vein-phase helical CT of the pancreas: optimal temporal window for evaluation of pancreatic adenocarcinoma. *AJR* 1999; 172:605-608.
 - 18 **Furukawa H, Takayasu K, Mukai K, Kanai Y, Inoue K, Kosuge T, et al:** Late contrast-enhanced CT for small pancreatic carcinoma: Delayed enhanced area on CT with histopathological correlation. *Hepatogastroenterology* 1996; 43:1230-1237.
 - 19 **Yamaguchi T, Matuura N, Ishihara T, Saisyo H:** Tumor forming pancreatitis - Differential diagnosis by dynamic CT scan. *Tantosui* 2002; 23:653-657. (In Japanese with English abstract)

Effects of tumor selective replication-competent herpes viruses in combination with gemcitabine on pancreatic cancer

Izuru Watanabe · Hideki Kasuya · Naohiro Nomura · Toshio Shikano · Takashi Shirota · Naohito Kanazumi · Shin Takeda · Shuji Nomoto · Hiroyuki Sugimoto · Akimasa Nakao

Received: 29 April 2007 / Accepted: 25 July 2007 / Published online: 29 August 2007
© Springer-Verlag 2007

Abstract

Purpose Pancreatic cancer still has a poor prognosis, even if aggressive therapy is pursued. Currently, new modalities of oncolytic virus therapy are being tested against this cancer. The combination of one of two representative mutant herpes simplex viruses (R3616: γ_1 34.5 inactivated, hrR3: UL39 inactivated) with a standard anti-pancreatic cancer chemotherapy drug (gemcitabine), was investigated in this study.

Experimental design The intracellular concentration of ribonucleotide reductase was estimated by Western blotting. The effect of gemcitabine on viral replication and the total cytotoxic effect of the combination therapy were investigated on pancreatic cancer cell lines. We compared the results of two oncolytic viruses, R3616 and hrR3. A mouse model of pancreatic cancer with peritoneal dissemination was used to evaluate the in vivo effect of the combination therapy.

Results Although the replication of both viruses was inhibited by gemcitabine, the combination caused more tumor cell cytotoxicity than did virus alone in vitro. The results with R3616 were more striking. Although the difference was not statistically significant, R3616 with gemcitabine had a greater effect than did R3616 alone, while hrR3 with gemcitabine had a weaker effect than did hrR3 alone in vivo experiments.

Conclusion The combination of oncolytic virus with gemcitabine is a promising new strategy against advanced

pancreatic cancer. Each virus has different functional characteristics, and can affect the results of the combination of viruses and chemotherapy drugs. The results indicate that there is a complicated interaction among viruses, cells, and chemotherapy drugs and that the best combination of oncolytic virus and chemotherapeutic agents should be studied more extensively before embarking on a clinical trial.

Keywords Herpes oncolytic virus · Pancreatic cancer · Gemcitabine · Combination therapy

Introduction

Pancreatic cancer is a disease with an extremely poor prognosis. Surgical therapy for pancreatic cancer is still insufficient to cure most patients [1]. Recently gemcitabine has shown a modest survival advantage over 5-fluorouracil (5-FU) in patients with this cancer [2]. Gemcitabine has become one of the standard chemotherapy drugs against pancreatic cancer but more effective therapies must be devised in order to significantly improve survival. Oncolytic virus therapy has been highly trusted as a new type of therapy for advanced incurable pancreatic cancer, and may provide some clinical benefit to those patients in the near future. Currently, clinical trials using oncolytic viruses have been started against many types of cancer in world-wide [3], such as brain cancer [4, 5], prostate cancer [6, 7], pancreatic cancer [8], breast cancer [9], and head and neck cancer [10, 11]. This study investigated the possibility of combination therapy using gemcitabine and two herpes mutant oncolytic viruses (R3616 and hrR3) against pancreatic cancer.

Gemcitabine (difluorodeoxycytidine; dFdC) is intracellularly phosphorylated to difluorodeoxycytidine diphosphate (dFdCDP) and difluorodeoxycytidine triphosphate

I. Watanabe · H. Kasuya (✉) · N. Nomura · T. Shikano · T. Shirota · N. Kanazumi · S. Takeda · S. Nomoto · H. Sugimoto · A. Nakao
Department of Surgery II,
Nagoya University School of Medicine,
65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan
e-mail: hidekikasuya@aol.com

(dFdCTP). dFdCTP competes with deoxycytidine triphosphate (dCTP) for incorporation into DNA, and DNA synthesis is inhibited [2, 12, 13]. In addition, dFdCDP acts as an inhibitor of ribonucleotide reductase (RR) in cells, which in turn causes a major decrease in the dCTP pool. Therefore, gemcitabine reduces the activity of RR in cancer cell lines [14] (Fig. 1a). However, some cells have been known to acquire chemoresistance to gemcitabine due to over expression of RR [15–21].

R3616 and hrR3 are genetically engineered herpes simplex viruses [3]. R3616 lacks the γ_1 34.5 gene that produces the ICP34.5 protein. Replication of R3616 is severely restricted in normal cells, because the expression of ICP34.5 in normal cells prevent a protein shutoff mechanism that is associated with eIF2 α dephosphorylation through the protein kinase receptor (PKR). Most cancer cells lose this normal protein shutoff mechanism so that viral replication can proceed, which induces the virally infected cells to undergo apoptosis to protect the integrity of the cell's DNA and block viral replication [3, 22–24]. hrR3 lacks the UL39 gene that produces the ICP6 proteins (viral RR), a key enzyme in the biosynthesis of DNA in all prokaryotic and eukaryotic cells. The viral replication of hrR3 is severely restricted in cells that have high levels of holding proteins involved in nucleic acid synthesis such as cancer cells [25, 26] (Fig. 1b).

We investigated the effect of tumor-selective, replication-competent herpes viruses (R3616 and hrR3) against pancreatic cancer under the same conditions in which gemcitabine effects cancer cells. Our major concern was how gemcitabine may interrupt viral replication, and whether the combination of an oncolytic virus with gemcitabine can significantly improve anti-pancreatic cancer therapy.

Materials and methods

Viruses and cells

R3616 was kindly provided by Bernard Roizman Sc. D (University of Chicago, Chicago, IL, USA) and hrR3 was kindly provided by Sandra K. Weller Ph.D. (University of Connecticut, Storrs, CT, USA). SW1990, derived from a human pancreatic carcinoma, was kindly provided by Dr. T. Sawada (First Department of Surgery, Osaka City University, Osaka, Japan). CAPAN 1, also derived from a human pancreatic carcinoma, was obtained from the Japanese Cancer Research Resources Bank, Tokyo, Japan. PACA2, another cell line derived from a human pancreatic carcinoma, was obtained from the American Type Culture Collection, Manassas, VA, USA. Cells were grown in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal calf serum and 1% penicillin/streptomycin at 37°C (Sigma, Tokyo, Japan).




Fig. 1 a Gemcitabine structure and pathway. Gemcitabine HCl is a nucleoside analog that exhibits anti-tumor activity. Gemcitabine HCl is 2'-deoxy-2', 2'-difluorocytidine monohydrochloride (β -isomer). The empirical formula for gemcitabine HCl is C₉H₁₁F₂N₃O₄ × HCl. It has a molecular weight of 299.66. Gemcitabine is metabolized intracellularly by nucleoside kinases to the active diphosphate (dFdCDP) and triphosphate (dFdCTP) nucleosides. b Schematic illustration of hrR3 and R3616. hrR3 is a mutated herpes simplex virus (HSV) that has the LacZ gene inserted into the site of UL39 (ICP6), causing inactivation of ribonucleotide reductase activity that is associated with UL39. Ribonucleotide reductase is a key enzyme for viral DNA synthesis. R3616 is a mutated HSV that has a deletion of both γ_1 34.5 genes. The γ_1 34.5 gene produces ICP 34.5 that dephosphorylates eIF2 α -phosphate to permit continued viral protein synthesis. Those mutated HSVs replicate and destroy only the cancer cells

Western blot assay

A total of 10⁶ cells were harvested and rinsed twice with phosphate-buffered saline, pH 7.4. Cell extracts were prepared with lysis buffer (20 mM Tris, pH 7.5, 0.1% Triton-X, 0.5% deoxycholate, 1 mM phenylmethylsulfonyl fluoride, 10 μ g/ml aprotinin, and 10 μ g/ml leupeptin) and clarified by

centrifugation at 12,000g, for 15 min, at 4°C. Cell lysates containing equal amounts of protein as determined by a BCA assay kit were electrophoresed on a NuPAGE, Novex 4–12% Bis-Tris Gel (Invitrogen, Carlsbad, CA, USA), and the resolved proteins were transferred to PVDF membranes (Invitrogen). The membranes were blocked with 5% nonfat milk overnight at room temperature, and incubated with 0.2 µg/ml human anti-RRM1 antibody (CHEMICON International, Temecula, CA, USA) for 1 h. RRM1 was detected using an enhanced chemiluminescence (ECL) system following the manufacturer's instructions (Amersham Life Science, Uppsala, Sweden). β -actin also was detected on the same membrane to serve as a control for the amount of protein loaded.

Cytotoxic assay

Gemcitabine and viral-induced cytotoxicity assays were performed using the MTT assay as previously described [27, 28]. Briefly, 10^6 cells were plated in a 10-cm plate and 10 µg/ml of gemcitabine was added. After 24 h, a replication-competent virus (R3616 or hrR3) was added at multiplicity of infection (MOI) values ranging from 0.01 to 10 and incubated for an additional 48 h. The number of surviving cells was quantified by a colorimetric MTT assay. The results, expressed as mean \pm SD of four samples, were compared with the results from the cytotoxicity assays of gemcitabine alone and the virus alone. Statistical significance was determined by the two-sided Student's *t*-test using SPSS (SPSS, Chicago, IL, USA).

Viral replication assay

Viral replication assays were performed as described [28, 29]. Briefly, 10^6 cells were plated in a 10-cm plate and 10 µg/ml of gemcitabine was added. After 24 h, replication competent viruses (R3616 or hrR3) were added at MOI of 2. Forty-eight hours after infection, the supernatant and cells were harvested, exposed to three freeze-thaw cycles to release the virions, and titered. The results were compared with the assays of viral replication without gemcitabine.

Animal studies

Mice (6-week-old females BALB/c nu/nu) were obtained from the Charles River Japan, Yokohama, Japan. Animal studies were performed in accordance with the guidelines issued by the Nagoya University Animal Center. The mice, used in a peritoneal-disseminated carcinoma model, were injected with 10^6 PACA2 cells into the intraperitoneal cavity. The condition of the animals was checked once or twice a day for the duration of the study. The mice were divided randomly into six groups (A–F). Group A ($n = 10$), group D ($n = 10$), and group E ($n = 10$) were injected with 1 mg of

gemcitabine into the intraperitoneal cavity on day 14 after the injection of the PACA2 cells. The mice in groups A and B ($n = 10$) each were injected with 10^6 particles of R3616 on day 15 after the injection of the PACA2 cells. Group C ($n = 10$) and group D ($n = 10$) were injected with 10^6 particles of hrR3 on day 15 after the injection of the PACA2 cells. Group F ($n = 10$) was the control group, which was injected with only PACA2 cells into the intraperitoneal cavity.

Statistical differences between groups were determined by the log-rank test with the use of JMP 5.0 software (SAS Inc., Cary, NC, USA). $P < 0.05$ was considered statistically significant.

Results

Expression of RRM1 by Western blotting

As previously reported by many researchers on their papers, overexpression of ribonucleotide reductase subunit 1 (RRM1) is associated with chemoresistance to gemcitabine [15–17]. We examined the intensity of RRM1 protein expression in Capan1, PACA2, and SW1990 cells (Fig. 2). The intensity of RRM1 expression in the PACA2 cells was greater than in the other cell lines. The results from many previous related papers regarding chemoresistance to gemcitabine, indicated that PACA2 cells might have the highest potential of chemo resistance to gemcitabine among the three cell lines.

Comparison of cytotoxic assays between hrR3 and R3616, with or without gemcitabine

We compared the cytotoxicity of R3616 (γ_1 34.5 deficiency) and hrR3 (ICP6: RR gene deficiency) viruses' combination with gemcitabine by the MTT assay (Fig. 3). With both R3616 and hrR3, the cytotoxicity was increased by their





Fig. 2 Expression of ribonucleotide reductase M1 (RRM1) by Western blotting. PACA2 cells expressed the most ribonucleotide reductase M1 (RRM1) by Western blot assays among three pancreatic cancer cell lines tested. β -actin served as a control for the amount of protein loaded in each lane

Fig. 3 Comparison of cytotoxic assays between hrR3 and R3616, with or without gemcitabine. For both R3616 (γ 134.5 gene inactivated) and hrR3 (ICP6: ribonucleotide reductase gene inactivated), the cytotoxicity was increased when combined with gemcitabine. However, there was a trend toward greater cytotoxicity with R3616 than with hrR3 against all cell lines



combination with gemcitabine, but the more significant increase in cytotoxicity was observed with R3616 than hrR3. On the other hand, PACA2 cells, which expressed the most RRM1 by a Western blot assay, had the lowest increase in cytotoxicity with the combination of hrR3 and gemcitabine.

Comparison of cytotoxic assays between gemcitabine alone and gemcitabine with low titer virus

We also compared the cytotoxicity between gemcitabine alone and gemcitabine with low titer virus by the MTT assay (Fig. 4). Of all cell lines, the combination of gemcitabine and an MOI 0.01 of R3616 showed more cytotoxic tendency than did gemcitabine alone ($P = 0.04$ on PACA2 cell line), while the combination of gemcitabine with an MOI 0.01 of hrR3 tend to be less cytotoxic than gemcitabine alone. PACA2 cells.

Comparison of viral replication between hrR3 and R3616, with or without gemcitabine


We compared the viral replication between R3616 and hrR3 in the presence of gemcitabine by the plaque-forming

assay (Fig. 5). The replication of both viruses was inhibited by gemcitabine. The titer of hrR3 declined more than did R3616 in combination with gemcitabine. The replication of hrR3 was inhibited by gemcitabine in all cell lines. PACA2 cells expressed the most RRM1 by Western blot assay, and hrR3 replicated more vigorously with gemcitabine in the PACA2 cells than in the other two cell lines, while R3616 was also inhibited by gemcitabine in all cell lines but with somewhat weaker inhibition comparing to hrR3.

Animal studies

Long-term survival (LTS: 100 days) was achieved in 60% of mice treated with an intraperitoneal injection of R3616 followed by gemcitabine (group A). Mice treated with an intraperitoneal injection of R3616 had only a 50% LTS (group B). Mice treated with hrR3 had a 30% LTS (group C). Mice treated with hrR3 followed by gemcitabine had a 20% LTS (group D). Mice treated with gemcitabine alone had only a 10% LTS (group E). All mice in the control group died within 60 days (group F) (Fig. 6). Statistical differences in the survival rates were determined by log-rank analyses (group A versus group F, $P = 0.0011$; group A versus group D, $P = 0.0078$; group E versus group F,

Fig. 4 Comparison of cytotoxic assays between gemcitabine alone and gemcitabine with a very low titer of virus. For all cell lines, the combination of gemcitabine and a very low titer (MOI 0.01) of R3616 tended to exhibit greater cytotoxicity than did gemcitabine alone ($P = 0.04$ on PACA2 cell line), on the other hand, the combination of gemcitabine and a very low titer (MOI 0.01) of hrR3 tended to be less cytotoxic than gemcitabine alone



$P = 0.006$; group B versus group D, $P = 0.0174$; group B versus group F, $P = 0.0049$). There were no other statistically significant differences between the other groups except for shown above. Although it was not significantly different, R3616 with gemcitabine tended to have a stronger effect than did R3616 alone, while hrR3 with gemcitabine tended to be weaker than hrR3 alone.


Discussion

In this study, we compared the efficacy of hrR3 or R3616 plus gemcitabine against pancreatic cancer. An in vitro cytotoxic assay indicated that R3616 plus gemcitabine caused a significant increase in the cell-killing effect in all three pancreatic cancer cell lines than did hrR3. We postulate that this result was due to the functional differences caused by each deleted viral gene. The viral replication of hrR3 might be more interrupted by the effect of gemcitabine than that of R3616, and this reduction might have been responsible for the slight decrease in the cell-killing effect

of hrR3. Cellular RR is important for viral replication especially for hrR3 that has no RR [3, 25–27, 29, 30]. Gemcitabine is well known to reduce the activity of cellular RR in cancer cell lines [14]. Therefore, it is a possible that the effect of gemcitabine was greater in combination with hrR3 than with R3616 reducing the replication and cytotoxicity of the viruses.

Interestingly, infection with hrR3 at a very low concentration (MOI 0.01) in the presence of gemcitabine caused less cytotoxic than did gemcitabine alone. This may be the result of the virus protecting the cancer cells from the apoptosis caused by gemcitabine. The virus itself has some anti-apoptotic effects on cells in order to protect the host cells from bursting too early and until the virus particles have matured. Although gemcitabine reduced the replication of hrR3, some viral anti-apoptosis genes might still have worked in the infected cells without the burst-cell effect that is caused by an abundance of mature viruses. The apoptosis mechanism might malfunction as a result of this low virus concentration, causing an anti-apoptotic effect against gemcitabine. This effect might apply not only to HSV, a critical

Fig. 5 Comparison of viral replication between hrR3 and R3616, with or without gemcitabine. The viral replication of both R3616 and hrR3 were inhibited by the presence of gemcitabine. This phenomenon was more prominent with hrR3 and gemcitabine than with R3616



consideration when using a viral vector or an oncolytic virus with chemotherapy drugs, because most viruses have such an anti-apoptosis gene. Examples include US3 and US5 in HSV [31, 32], and E1b 19 kDa in adenovirus [33]. Furthermore, several distinct viruses have been shown to develop mechanisms to block premature apoptosis of infected cells [34–36]. This phenomenon should be considered when using any viral vector for gene therapy or oncolytic virus therapy with chemotherapy drugs. In our opinion, the anti-apoptosis genes in a virus should be studied more intensively if future development of oncolytic virus therapy is to proceed.

PACA2 cells had the highest density of RR by Western assays and also the lowest cytotoxic effect from single agent gemcitabine among the three pancreatic cancer cell lines tested as 60% cell survival in Fig. 4, which indicates that PACA2 cells have some type of resistance against the cytotoxicity of gemcitabine comparing to other two cell lines. For the combination of R3616 with gemcitabine, increased efficacy was observed against all the pancreatic cancer cell lines even if the cells had some resistance to the chemotherapy alone. On the other hand, the combination of

hrR3 with gemcitabine was of weak cytotoxicity toward PACA2 cells, which expressed the most RRM1 by Western blot assay, and the effect was less pronounced than when R3616 was used. These results suggest that, the combination of R3616 and gemcitabine might be suitable for the cancer cell type that is expected to offer resistance to gemcitabine.

In the in vivo experiments, the combination of R3616 with gemcitabine yielded a 60% LTS rate (100 days) in the mice. This was higher than in mice treated with an intraperitoneal injection of R3616 alone that resulted in a 50% LTS rate, while mice treated with only hrR3 had a 30% LTS rate; however, there was no statistically significant difference in the LTS rate between R3616 and R3616 with gemcitabine. Thus, combination therapy with R3616 and gemcitabine had the same or slightly higher efficacy than the virus alone. However, mice treated with hrR3 followed by gemcitabine showed a lower LTS rate (20%) than those treated with hrR3 alone. And moreover, there was a statistically significant difference between group A (R3616 + GEM) and group D (hrR3 + GEM) ($P = 0.0078$). From the results of our in vivo and in vitro, we determined that the




Fig. 6 a Cumulative survival curves of an in vivo mouse model. The PACA2 cells were injected into the peritoneal cavity. Each group was treated as shown below. Group A: R3616 + gemcitabine (GEM), group B: R3616 only, group C: hrR3 only, group D: hrR3 + GEM, group E: GEM only, and group F: no treatment. Differences in the survival rates were assessed by log-rank analysis (group A versus group F, $P = 0.0011$; group A versus group D, $P = 0.0078$; group E versus group F, $P = 0.006$; group B versus group D, $P = 0.0174$; and group B versus group F, $P = 0.0049$). There were no other statistically significant differences between the other groups except for shown above. There was a statistically significant difference between group A (R3616 + GEM) and group D (hrR3 + GEM) ($P = 0.0078$). b Long-term survival (LTS: over 100 days). LTS was achieved in 60% of mice treated with an intraperitoneal injection of R3616 followed by gemcitabine (group A). Mice treated with an intraperitoneal injection of R3616 only had a 50% LTS survival rate (group B). Mice treated with hrR3 only had a 30% LTS rate (group C). Mice treated with hrR3 followed by gemcitabine had a 20% LTS rate (group D). Mice treated with gemcitabine alone had only a 10% LTS rate (group E). The untreated group (group F) had a 0% LTS survival rate

combination of gemcitabine with R3616 (γ 134.5 inactivated) might be more effective than the combination with hrR3 (RR inactivated).

Potentially, chemotherapy drugs connote to inhibit oncolytic virus replication to some degree, but this effect may be influenced by the differences in the characteristics of each virus caused by gene mutation. UL 39 (ICP6)-deleted HSVs, such as G207 [3, 37], and Myb34.5 [38, 39] also have some kind of potential likely to be inhibited by

gemcitabine, as is hrR3, because of the genetic characteristics of RR. In other words, UL39-intact HSVs, such as HF10 [40, 41], RH105 [42], and DF γ 34.5 [43] are likely to interact differently from a UL 39-deleted HSV (e.g., hrR3), in combination therapy with gemcitabine. Additional studies must be needed for further confirmation of the efficacy depending upon the functional characteristics among the chemotherapy drugs, viruses, and the cancer cells.

In the future, oncolytic virus therapy in combination with chemotherapy drugs may become more popular for use in clinical trials. Therefore, the characteristics of each virus must be considered carefully to determine if they are suitable for use with the chemotherapy drugs chosen.

References

- Warshaw AL, Fernandez-del Castillo C (1992) Pancreatic carcinoma. *N Engl J Med* 326:455–464
- Burris HA III, Moore MJ, Andersen J, Green MR, Rothenberg ML, Modiano MR, et al (1997) Improvements in survival and clinical benefit with gemcitabine as first-line therapy for patients with advanced pancreas cancer: a randomized trial. *J Clin Oncol* 15:2403–2413
- Kasuya H, Takeda S, Nomoto S, Nakao A (2005) The potential of oncolytic virus therapy for pancreatic cancer. *Cancer Gene Ther* 12(9):725–736
- Germano IM, Fable J, Gultekin SH, Silvers A (2003) Adenovirus/herpes simplex-thymidine kinase/ganciclovir complex: preliminary results of a phase I trial in patients with recurrent malignant gliomas. *J Neuro oncol* 65:279–289
- Immonen A, Vapalahti M, Tyynela K, Hurskainen H, Sandmair A, Vanninen R, et al (2004) AdvHSV-tk gene therapy with intravenous ganciclovir improves survival in human malignant glioma: a randomized, controlled study. *Mol Ther* 10:967–972
- Teh BS, Ayala G, Aguilar L, Mai WY, Timme TL, Vlachaki MT, et al (2004) Phase I-II trial evaluating combined intensity-modulated radiotherapy and in situ gene therapy with or without hormonal therapy in treatment of prostate cancer- interim report on PSA response and biopsy data. *Int J Radiat Oncol Biol Phys* 58:1520–1529
- Kubo H, Gardner TA, Wada Y, Koeneman KS, Gotoh A, Yang L, et al (2003) Phase I dose escalation clinical trial of adenovirus vector carrying osteocalcin promoter-driven herpes simplex virus thymidine kinase in localized and metastatic hormone-refractory prostate cancer. *Hum Gene Ther* 14:227–241
- Mulvihill S, Warren R, Venook A, Adler A, Randlev B, Heise C, et al (2001) Safety and feasibility of injection with an E1B-55 kDa gene-deleted, replication-selective adenovirus (ONYX-015) into primary carcinoma of the pancreas: a phase I trial. *Gene Ther* 8:308–315
- Nakao A, Kimata H, Imai T, Kikumori T, Teshigahara O, Nagasaka T, et al (2004) Intratumoral injection of herpes simplex virus HF10 in recurrent breast cancer. *Ann Oncol* 15:987–991
- Lamont JP, Nemunaitis J, Kuhn JA, Landers SA, McCarty TM (2000) A prospective phase II trial of ONYX-015 adenovirus and chemotherapy in recurrent squamous cell carcinoma of the head and neck (the Baylor experience). *Ann Surg Oncol* 7:588–592
- Neunmaitis J, Cunningham C, Buchanan A, Edelman G, Maples P, Netto G, et al (2001) Intravenous infusion of a replication-selective adenovirus (ONYX-015) in cancer patients: safety, feasibility and biological activity. *Gene Ther* 8:746–759

12. Heinemann V, Hertel LW, Grindey GB, Plunkett W (1988) Comparison of the cellular pharmacokinetics and toxicity of 2',2'-difluorodeoxycytidine and 1- β -D-arabinofuranosylcytosine. *Cancer Res* 48:4024–4031
13. Huang P, Chubb S, Hertel LW, Plunkett W (1991) Action of 2',2'-difluorodeoxycytidine on DNA synthesis. *Cancer Res* 51:6110–6117
14. Heinemann V, Xu Y, Chubb S, Sen A, Hertel LW, Grindey GB, et al (1990) Inhibition of ribonucleotide reduction in CCRF-CEM cells by 2', 2'-difluorodeoxycytidine. *Mol Pharmacol* 38:567–572
15. Rosell R, Felip E, Taron M, Majo J, Mendez P, Sanchez-Ronco M, et al (2004) Gene expression as a predictive marker of outcome in stage (B-IIIa-IIIb non-small cell lung cancer after induction gemcitabine-based chemotherapy followed by resectional surgery. *Clin Cancer Res* 10:4215s–4219s
16. Davidson JD, Ma L, Flagella M, Geeganage S, Gelbert LM, Slapak CA (2004) An increase in the expression of ribonucleotide reductase large subunit 1 is associated with gemcitabine resistance in non-small cell lung cancer cell lines. *Cancer Res* 64:3761–3766
17. Rosell R, Danenberg KD, Alberola V, Bepler G, Sanchez JJ, Camps C, et al (2004) Ribonucleotide reductase messenger RNA expression and survival in gemcitabine/cisplatin-treated advanced non-small cell lung cancer patients. *Clin Cancer Res* 10:1318–1325
18. Goan YG, Zhou B, Hu E, Mi S, Yen Y (1999) Overexpression of ribonucleotide reductase as a mechanism of resistance to 2,2-difluorodeoxycytidine in the human KB cancer cell line. *Cancer Res* 59:4204–4207
19. Jung CP, Motwani MV, Schwartz GK (2001) Flavopiridol increases sensitization to gemcitabine in human gastrointestinal cancer cell lines and correlates with down-regulation of ribonucleotide reductase M2 subunit. *Clin Cancer Res* 7:2527–2536
20. Duxbury MS, Ito H, Zinner MJ, Ashley SW, Whang EE (2004) RNA interference targeting the M2 subunit of ribonucleotide reductase enhances pancreatic adenocarcinoma chemosensitivity to gemcitabine. *Oncogene* 23:1539–1548
21. Duxbury MS, Ito H, Benoit E, Zinner MJ, Ashley SW, Whang EE (2004) Retrovirally mediated RNA interference targeting the M2 subunit of ribonucleotide reductase: a novel therapeutic strategy in pancreatic cancer. *Surgery* 136:261–269
22. Andreansky SS, He B, Gillespie GY, Soroceanu L, Markert J, Chou J, et al (1996) The application of genetically engineered herpes simplex viruses to the treatment of experimental brain tumors. *Proc Natl Acad Sci USA* 93:11313–11318
23. Pyles RB, Warnick RE, Chalk CL, Szanti BE, Parysek LM (1997) A novel multiply-mutated HSV-1 strain for the treatment of human brain tumors. *Hum Gene Ther* 8:533–544
24. Coukos G, Makrigiannakis A, Kang EH, Rubin SC, Albelda SM, Molnar-Kimber KL (2004) Oncolytic herpes simplex virus-1 lacking ICP34.5 induces p53-independent death and is efficacious against chemotherapy-resistant ovarian cancer. *Clin Cancer Res* 6:3342–3353
25. Kasuya H, Nishiyama Y, Nomoto S, Hosono J, Takeda S, Nakao A (1999) Intraperitoneal delivery of hrR3 and ganciclovir prolongs survival in mice with disseminated pancreatic cancer. *J Surg Oncol* 72:136–141
26. Kasuya H, Mizuno M, Yoshida J, Nishiyama Y, Nomoto S, Nakao A (2000) Combined effects of adeno-associated virus vector and a herpes simplex virus mutant as neoplastic therapy. *J Surg Oncol* 74:214–218
27. Carroll NM, Chiocca EA, Takahashi K, Tanabe KK (1996) Enhancement of gene therapy specificity for diffuse colon carcinoma liver metastases with recombinant herpes simplex virus. *Ann Surg* 224:323–330
28. Kasuya H, Pawlik TM, Mullen JT, Donahue JM, Nakamura H, Chandrasekhar S, et al (2004) Selectivity of an oncolytic herpes simplex virus for cells expressing the DF3/MUC1 antigen. *Cancer Res* 64:2561–2567
29. Yoon SS, Nakamura H, Carroll NM, Bode BP, Chiocca EA, Tanabe KK (2000) An oncolytic herpes simplex virus type 1 selectively destroys diffuse liver metastases from colon carcinoma. *FASEB J* 14:301–311
30. Goldstein DJ, Weller S (1988) Factor(s) present in herpes simplex virus type 1-infected cells can compensate for the loss of the large subunit of the viral ribonucleotide reductase: characterization of an ICP6 deletion mutant. *Virology* 166:41–51
31. Ogg PD, McDonnell PJ, Ryckman BJ, Knudson CM, Roller RJ (2004) The HSV-1 Us3 protein kinase is sufficient to block apoptosis induced by overexpression of a variety of Bcl-2 family members. *Virology* 319:212–224
32. Benetti L, Roizman B (2004) Herpes simplex virus protein kinase Us3 activates and functionally overlaps protein kinase A to block apoptosis. *Proc Natl Acad Sci USA* 101:9411–9416
33. Huang DC, Cory S, Strasser A (1997) Bcl-2, Bcl-XL and adenovirus protein E1B19kD are functionally equivalent in their ability to inhibit cell death. *Oncogene* 14:405–414
34. Teodoro JG, Branton PE (1997) Regulation of p53-dependent apoptosis, transcriptional repression, and cell transformation by phosphorylation of the 55-kilodalton E1B protein of human adenovirus type 5. *J Virol* 71:3620–3627
35. Teodoro JG, Branton PE (1997) Regulation of apoptosis by viral gene products. *J Virol* 71:1739–1746
36. Tschopp J, Thome M, Hofmann K, Meink E (1998) The fight of viruses against apoptosis. *Curr Opin Genet Dev* 8:82–87
37. Cinatl J Jr, Michaelis M, Driever PH, Cinatl J, Hrabeta J, Suhan T, et al (2004) Mutated herpes simplex virus g207 is a potent inhibitor of angiogenesis. *Neoplasia* 6:725–735
38. Nakamura H, Kasuya H, Mullen JT, Yoon SS, Pawlik TM, Chandrasekhar S, et al (2002) Regulation of herpes simplex virus g₁ 34.5 expression and oncolysis of diffuse liver metastases by Myb34.5. *J Clin Invest* 109:871–882
39. Chung RY, Saeki Y, Chiocca EA (1999) β -myb promoter retargeting of herpes simplex virus gamma 34.5 gene-mediated virulence toward tumor and cycling cells. *J Virol* 73:7556–7564
40. Teshigahara O, Goshima F, Takao K, Kohno S, Kimata H, Nakao A, et al (2004) Oncolytic viral therapy for breast cancer with herpes simplex virus type 1 mutant HF10. *J Surg Oncol* 85:42–47
41. Kimata H, Takakuwa H, Goshima F, Teshigahara O, Nakao A, Kurata T, et al (2003) Effective treatment of disseminated peritoneal colon cancer with new replication-competent herpes simplex viruses. *Hepatogastroenterology* 50:961–966
42. Nilaver G, Muldoon LL, Kroll RA, Pagel MA, Breakefield XO, Davidson BL, et al (1995) Delivery of herpesvirus and adenovirus to nude rat intracerebral tumors after osmotic blood-brain barrier disruption. *Proc Natl Acad Sci USA* 92:9829–9833
43. Kasuya H, Pawlik TM, Mullen JT, Donahue JM, Nakamura H, Chandrasekhar S, et al (2004) Selectivity of an oncolytic herpes simplex virus for cells expressing the DF3/MUC1 antigen. *Cancer Res* 64:2561–2567

Analysis of Clinicopathological Features and Predictors of Malignancy in Intraductal Papillary Mucinous Neoplasms of the Pancreas

Tsutomu Fujii MD¹, Tadao Ishikawa MD¹, Naohito Kanazumi MD¹, Hiroyuki Sugimoto MD¹, Shuji Nomoto MD¹, Soichiro Inoue MD¹, Tetsuro Nagasaka MD², Shin Takeda MD¹, Akimasa Nakao MD, FACS¹

Department of ¹Surgery II and ²Laboratory Medicine, Nagoya University, Graduate School of Medicine Nagoya, Japan

Corresponding Author: Tsutomu Fujii, MD, Department of Surgery II, Nagoya University, Graduate School of Medicine, 65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan
Tel: +81 52 744 2245, Fax: +81 52 744 2255, E-mail: fjt@med.nagoya-u.ac.jp

KEY WORDS:

IPMN; IPMC;
predictor; PHRSD

ABBREVIATIONS:

Intraductal Papillary Mucinous Neoplasm (IPMN); Mucinous Cystic Neoplasm (MCN); World Health Organization (WHO); Intraductal Papillary Mucinous Carcinoma (IPMC); Computed Tomography (CT); Magnetic Resonance Cholangiopancreatography (MRCP); Endoscopic Retrograde Cholangiopancreatography (ERCP); Endoscopic Ultrasonography (EUS); Main Pancreatic Duct (MPD); Carcinoma *In Situ* (CIS); Carbohydrate Antigen 19-9 (CA19-9); Carcinoembryonic Antigen (CEA); Pancreaticoduodenectomy (PD); Pylorus-preserving Pancreaticoduodenectomy (PpPD); Pancreatic Head Resection with Segmental Duodenectomy (PHRSD); Segmental Resection of the Pancreatic Body (SR);

ABSTRACT

Background/Aims: Intraductal papillary mucinous neoplasms (IPMNs) are increasingly recognized, but it is very difficult to evaluate accurately the malignancy of these neoplasms by modern imaging. We reviewed our experience in order to elucidate predictors of tumor malignancy, invasiveness, and outcome.

Methodology: The clinicopathological features and surgical outcomes of 57 patients with IPMNs who underwent surgery in Nagoya University Hospital were analyzed.

Results: The histological diagnosis was adenoma in 40, borderline in 1, carcinoma *in situ* (CIS) in 7, and invasive carcinoma in 9 patients. Patients with invasive carcinomas had significantly shorter survival

rates than patients with benign IPMNs or CIS ($p < 0.0001$). Multivariate analyses revealed that the main duct or the combined type was significantly predictive of malignancy, and both main duct or combined type and diabetes mellitus were associated significantly with invasive carcinoma.

Conclusions: IPMNs generally grow slowly, but have a malignant potential that warrants radical surgical treatment when the tumor component invades the parenchyma. Our results suggest that the above factors should be considered in surgical management. The main duct type of IPMN or IPMN with mural nodules is potentially malignant or invasive. Therefore, radical operative management is indicated in these IPMNs.

INTRODUCTION

Intraductal papillary mucinous neoplasms (IPMNs) were described in the Japanese literature in 1982 (1) and in the English literature in 1986 (2). This entity is increasingly recognized and is characterized by an adenomatous proliferation of pancreatic duct epithelium that may involve the main pancreatic duct or ductal branches alone or in combination (3,4). IPMN is comprised of a spectrum of diseases from benign to malignant (5,6). This neoplasm is rare, constituting only 1% of pancreatic cancers, according to previous reports (7,8). Before the concept of IPMNs was established, confusion with mucinous cystic neoplasms (MCNs) or other cystic tumors of the pancreas existed (9). In 1996, the World Health Organization (WHO) established criteria to classify IPMNs and to distinguish them from MCNs (10). IPMNs exhibit different stages from adenoma with mild atypia to invasive carcinoma (10), but lack the ovarian stroma characteristics of MCNs. Most IPMNs, including non-invasive intraductal papillary mucinous carcinomas (IPMCs), have a more favorable prognosis than pan-

creatic ductal carcinoma, however, some cases of invasive IPMC have a poor prognosis (11-15). Therefore, appropriate management requires the differentiation of malignant tumors, but it is very difficult to evaluate accurately the extent of malignancy by modern imaging.

In this study, we retrospectively reviewed our experience and analyzed the clinicopathological features of IPMNs to elucidate predictive factors associated with malignancy, invasiveness, and tumor outcome.

METHODOLOGY

Fifty-seven patients who underwent surgery for IPMNs between March 1991 and July 2004 in Nagoya University Hospital were reviewed. We retrospectively examined the clinicopathological features, surgical outcomes, and factors predictive of malignancy. The study was approved by the Ethics Committee of the hospital. Informed consent was obtained from all patients for the subsequent use of resected tissues. All patients underwent radiographic and endoscopic examination such as computed tomography (CT), mag-