

Table II. Relationship between CD10 mRNA and metastasis in colorectal cancer patients.

	No. (n=167)	CD10 mRNA (T/N±S.D.)	P
Liver metastasis			
Negative	125	18.65±61.20	0.873
Positive	42	27.56±108.96	
All metastases			
Negative	94	20.13±69.11	0.886
Positive	73	21.87±84.12	

there was no significant difference between the groups. There were also no significant survival differences according to clinical stage (data not shown).

Discussion

We have recently demonstrated that CD10 protein expression in colorectal cancer cells was significantly associated with liver metastasis and that CD10 protein expression was an independent predictor of liver metastasis (10). Yao *et al.* have also demonstrated a significant association between CD10 protein expression and liver metastasis from colorectal cancer (8), and other reports have indicated a relationship between CD10 protein expression and the development and progression of colorectal cancer (7, 9). These results prompted us to examine the association between CD10 mRNA expression level and liver metastasis using real-time PCR. Although CD10 mRNA in tumor tissues was higher than that in matched normal tissues in more than 80% of colorectal cancers and was associated with tumor progression, there was no significant relationship between the level of CD10 mRNA expression and metastasis, including liver metastasis. This suggested that CD10 might play an important role in tumorigenesis and tumor progression and that measurement of CD10 mRNA in colorectal cancer tissues is not useful as a predictor of liver metastasis.

In this study, the level of CD10 mRNA was higher in pN1 and pN2 tumors than in pN0 tumors, and was also higher in stage III and IV tumors than in stage I and II tumors. These facts suggested that the CD10 mRNA level was associated with tumor progression. Many previous studies have demonstrated that overexpression of CD10 protein is associated with tumor progression (1-5, 7-11, 15, 17, 19-22) and with tumor proliferation and microvascular density (11), thus indicating that CD10 plays an important role in tumor progression. Although the actual function of CD10 in tumors is not known, it is a cell surface metalloendopeptidase with structural similarity to matrix




Figure 2. Survival curves for patients with $T/N \geq 5$ or $T/N < 5$. There was no significant difference between the groups.

metalloprotease, and is capable of degrading a number of bioactive peptide and cytokines. Therefore, CD10 at the tumor cell surface and in the areas adjacent to tumor glands is considered to activate or inactivate tumor-related substances and to create a microenvironment that facilitates tumor cell invasion and metastasis. On the other hand, several studies have demonstrated an association between reduced expression of CD10 and tumor progression in lymphoma (23), and renal cell (14), lung (18), prostate (16), ovarian (12) and cervical cancer (13). Tumorigenesis and proliferation are inhibited (12,13), and apoptosis is induced, by CD10 (23). When CD10 expression is reduced in a tumor, loss of these functions of CD10 was considered to promote tumor progression. Because these tumors arise from tissues that normally express CD10, loss of CD10 in an advanced cancer may be explained in terms of dedifferentiation during tumorigenesis. On the other hand, CD10 expression in tumors that arise from normal tissues without CD10 expression probably represent a phenomenon acquired during tumorigenesis and is considered to be associated with tumor progression. Colorectal cancer acquires CD10 expression during tumorigenesis because it is not expressed in normal colorectal mucosa or stroma.

CD10 protein is reportedly expressed even in tumor stroma (5, 7, 11, 17, 20-22). In colorectal cancer, CD10 protein is expressed both in tumor and stromal cells (6, 7). A similar pattern has been reported in gastric cancer (1, 5), melanoma (19, 20) and pancreatic endocrine tumor (11). CD10 expression in tumor and/or stromal cells was associated with tumor progression, except in pancreatic endocrine tumor. In the present study, because CD10 mRNA was extracted from both tumor and stroma, it was quantified in both tissues as a whole, and there was no clear relationship between CD10 mRNA expression and

liver metastasis. CD10 protein expression in tumor cells has been shown to be significantly associated with liver metastasis (8, 10). Therefore, CD10 mRNA in tumor cells might be associated with liver metastasis. Further investigation will be necessary to clarify the relationship between CD10 mRNA in tumor cells and colorectal cancer liver metastasis.

CD10 mRNA expression was higher in colon cancer than in rectal cancer. Because CD10 is expressed in the brush border of the small intestine (6), colon cancer is considered to have a higher tendency to differentiate to the small intestine than rectal cancer. In gastric cancer, phenotypic differences have been associated with prognosis and the pattern of recurrence (1, 3). These facts suggest that there are biological and oncological differences between colon and rectal cancer.

The level of CD10 mRNA was higher in well- or moderately-differentiated adenocarcinoma than in poorly differentiated or mucinous adenocarcinoma. However, CD10 mRNA expression in poorly differentiated or mucinous adenocarcinoma is still higher than in normal mucosa. On the other hand, CD10 protein is reportedly undetectable in poorly-differentiated adenocarcinoma (6, 26). This suggests that the expression of CD10 mRNA is not directly associated with protein production and that post-transcriptional regulation plays an important role in protein expression in cancer cells. This is one of the reasons why liver metastasis was associated with CD10 protein expression and not with CD10 mRNA.

In conclusion, CD10 mRNA shows significantly higher expression in tumor tissue than in matched normal tissue. Although CD10 mRNA is associated with depth of invasion, lymph node status and TNM stage, it is not associated with liver metastasis, any type of metastasis, or prognosis. Therefore it seems that CD10 mRNA extracted from tumor tissues might not be useful as a predictor of liver metastasis or a prognostic marker.

Acknowledgements

This work was supported by a Grant-in-Aid for Cancer Research from the Ministry of Health, Labor and Welfare of Japan.

References

- 1 Tajima Y, Shimoda T, Nakanishi Y, Yokoyama N, Tanaka T, Shimizu K, Saito T, Kawamura M, Kusano M and Kumagai K: Gastric and intestinal phenotypic marker expression in gastric carcinomas and its prognostic significance: immunohistochemical analysis of 136 lesions. *Oncology* 61: 212-220, 2001.
- 2 Naritomi K, Futami K, Arima S and Iwashita A: Malignant potential regarding mucin phenotypes and endocrine cell differentiation in gastric adenocarcinoma. *Anticancer Res* 23: 4411-4422, 2003.
- 3 Tajima Y, Yamazaki K, Nishino N, Morohara K, Yamazaki T, Kaetsu T, Suzuki S, Kawamura M, Kumagai K and Kusano M: Gastric and intestinal phenotypic marker expression in gastric carcinomas and recurrence pattern after surgery-immunohistochemical analysis of 213 lesions. *Br J Cancer* 91: 1342-1348, 2004.
- 4 Carl-McGrath S, Lendeckel U, Ebert M, Wolter AB, Roessner A and Rocken C: The ectopeptidases CD10, CD13, CD26, and CD143 are upregulated in gastric cancer. *Int J Oncol* 25: 1223-1232, 2004.
- 5 Huang WB, Zhou XJ, Chen JY, Zhang LH, Meng K, Ma HH and Lu ZF: CD10-positive stromal cells in gastric carcinoma: correlation with invasion and metastasis. *Jpn J Clin Oncol* 35: 245-250, 2005.
- 6 Yao T, Tsutsumi S, Akaiwa Y, Takata M, Nishiyama K, Kabashima A and Tsuneyoshi M: Phenotypic expression of colorectal adenocarcinomas with reference to tumor development and biological behavior. *Jpn J Cancer Res* 92: 755-761, 2001.
- 7 Ogawa H, Iwaya K, Izumi M, Kuroda M, Serizawa H, Koyanagi Y and Mukai K: Expression of CD10 by stromal cells during colorectal tumor development. *Hum Pathol* 33: 806-811, 2002.
- 8 Yao T, Takata M, Tsutsumi S, Nishiyama K, Taguchi K, Nagai E and Tsuneyoshi M: Phenotypic expression of gastrointestinal differentiation markers in colorectal adenocarcinomas with liver metastasis. *Pathology* 34: 556-560, 2002.
- 9 Iwase T, Kushima R, Mukai K, Mitsufuji S, Okanoue T and Hattori T: Overexpression of CD10 and reduced MUC2 expression correlate with the development and progression of colorectal neoplasms. *Pathol Res Pract* 201: 83-91, 2005.
- 10 Fujimoto Y, Nakanishi Y, Sekine S, Yoshimura K, Akasu T, Moriya Y and Shimoda T: CD10 expression in colorectal carcinoma correlates with liver metastasis. *Dis Colon Rectum* 48: 1883-1889, 2005.
- 11 Deschamps L, Handra-Luca A, O'Toole D, Sauvanet A, Ruzniewski P, Belghiti J, Bedossa P and Couvelard A: CD10 expression in pancreatic endocrine tumors: correlation with prognostic factors and survival. *Hum Pathol* 37: 802-808, 2006.
- 12 Kajiyama H, Shibata K, Terauchi M, Morita T, Ino K, Mizutani S and Kikkawa F: Neutral endopeptidase 24.11/CD10 suppresses progressive potential in ovarian carcinoma *in vitro* and *in vivo*. *Clin Cancer Res* 11: 1798-1808, 2005.
- 13 Terauchi M, Kajiyama H, Shibata K, Ino K, Mizutani S and Kikkawa F: Anti-progressive effect of neutral endopeptidase 24.11 (NEP/CD10) on cervical carcinoma *in vitro* and *in vivo*. *Oncology* 69: 52-62, 2005.
- 14 Gohring B, Holzhausen HJ, Meye A, Heynemann H, Rebmann U, Langner J and Riemann D: Endopeptidase 24.11/CD10 is down-regulated in renal cell cancer. *Int J Mol Med* 2: 409-414, 1998.
- 15 Langner C, Ratschek M, Rehak P, Schips L and Zigeuner R: CD10 is a diagnostic and prognostic marker in renal malignancies. *Histopathology* 45: 460-467, 2004.
- 16 Osman I, Yee H, Taneja SS, Levinson B, Zeleniuch-Jacquotte A, Chang C, Nobert C and Nanus DM: Neutral endopeptidase protein expression and prognosis in localized prostate cancer. *Clin Cancer Res* 10: 4096-4100, 2004.
- 17 Iwaya K, Ogawa H, Izumi M, Kuroda M and Mukai K: Stromal expression of CD10 in invasive breast carcinoma: a new predictor of clinical outcome. *Virchows Arch* 440: 589-593, 2002.

- 18 Tokuhara T, Adachi M, Hashida H, Ishida H, Taki T, Higashiyama M, Kodama K, Tachibana S, Sasaki S and Miyake M: Neutral endopeptidase/CD10 and aminopeptidase N/CD13 gene expression as a prognostic factor in non-small cell lung cancer. *Jpn J Thorac Cardiovasc Surg* 49: 489-496, 2001.
- 19 Kanitakis J, Narvaez D and Claudy A: Differential expression of the CD10 antigen (neutral endopeptidase) in primary *versus* metastatic malignant melanomas of the skin. *Melanoma Res* 12: 241-244, 2002.
- 20 Bilalovic N, Sandstad B, Golouh R, Nesland JM, Selak I and Torlakovic EE: CD10 protein expression in tumor and stromal cells of malignant melanoma is associated with tumor progression. *Mod Pathol* 17: 1251-1258, 2004.
- 21 Braham H, Trimeche M, Ziadi S, Mestiri S, Mokni M, Amara K, Hachana M, Sriha B and Korbi S: CD10 expression by fusiform stromal cells in nasopharyngeal carcinoma correlates with tumor progression. *Virchows Arch* 449: 220-224, 2006.
- 22 Piattelli A, Fioroni M, Iezzi G, Perrotti V, Stellini E, Piattelli M and Rubini C: CD10 expression in stromal cells of oral cavity squamous cell carcinoma: a clinic and pathologic correlation. *Oral Dis* 12: 301-304, 2006.
- 23 Bai M, Agnantis NJ, Skyrlas A, Tsanou E, Kamina S, Galani V and Kanavaros P: Increased expression of the bcl6 and CD10 proteins is associated with increased apoptosis and proliferation in diffuse large B-cell lymphomas. *Mod Pathol* 16: 471-480, 2003.
- 24 Chomczynski P and Sacchi N: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156-159, 1987.
- 25 Tsuji N, Kamagata C, Furuya M, Kobayashi D, Yagihashi A, Morita T, Horita S and Watanabe N: Selection of an internal control gene for quantitation of mRNA in colonic tissues. *Anticancer Res* 22: 4173-4178, 2002.
- 26 Sato Y, Itoh F, Hinoda Y, Ohe Y, Nakagawa N, Ueda R, Yachi A and Imai K: Expression of CD10/neutral endopeptidase in normal and malignant tissues of the human stomach and colon. *J Gastroenterol* 31: 12-17, 1996.

Received May 7, 2007

Revised June 20, 2007

Accepted July 3, 2007

A Case of Colon Cancer Detected by Carbon-11 Choline Positron Emission Tomography/Computed Tomography: An Initial Report

Takashi Terauchi¹, Ukihide Tateishi², Tetsuo Maeda², Daisuke Kanou¹, Hiromitsu Daisaki¹, Yoshihiro Moriya³, Noriyuki Moriyama⁴ and Tadao Kakizoe⁴

¹Division of Cancer Screening, Research Center for Cancer Prevention and Screening, National Cancer Center, ²Division of Radiology, ³Division of Colorectal Surgery, National Cancer Center Hospital and ⁴Center for Cancer Prevention and Screening, National Cancer Center, Tokyo, Japan

Received January 4, 2007; accepted May 25, 2007

[C-11] choline positron emission tomography ([C-11] choline PET) has been expected to be one of the new PET modalities similar to [F-18] fluorodeoxyglucose positron emission tomography (FDG-PET), which has spread worldwide as a gold standard of PET oncologic imaging. However, there has been no report on [C-11] choline PET used for detection of colorectal cancer, which is one of major targets of oncologic FDG-PET. We initiated the research to investigate the detectability of [C-11] choline PET for various tumors including colorectal cancer. This is the first report of a patient who underwent surgical resection for advanced colon cancer depicted by [C-11] choline positron emission tomography/computed tomography.

Key words: radiology-PET – diagnostic radiology – GI-colorectum-basic

INTRODUCTION

[C-11] choline positron emission tomography ([C-11] choline PET) has been expected as a new PET modality and reported to be useful for the detection of various tumors, such as brain tumor, lung cancer, esophageal cancer, prostate cancer, gynecological cancers, and bone and soft tissue sarcomas (1–6). We started the research in our institution on September 1, 2005 to clarify not only the detectability of [C-11] choline PET for various tumors but the mechanism of choline accumulation to cancer cells, which is approved by the Institutional Review Board. Here we report a successful detection of an advanced colon cancer by [C-11] choline PET, which seems to be the first case, and discuss the possibility of application of [C-11] choline PET to colorectal cancer.

CASE REPORT

A 50-year-old woman presented with melena and abdominal discomfort. The colonoscopy showed the elevated lesion in the sigmoid colon (Fig. 1). From endoscopic findings this tumor was diagnosed as type 1 advanced colon cancer with invasion into the subserosa. Pathologic diagnosis by

specimen of biopsy was well-differentiated adenocarcinoma. The computed tomography scan of the thorax, abdomen and pelvis revealed the thickening in the wall of the sigmoid colon and enlarged uterus suspected of leiomyoma. No specific enlarged lymph nodes and no definite metastases including the liver were detected. A whole body [C-11] choline positron emission tomography/computed tomography (PET/CT) was performed with the written informed consent to participate in this research approved by the Institutional Review Board. Emission scans from the skull to the mid thigh were obtained starting 14 min after intravenous injection of 444MBq of [C-11] choline, which was synthesized with a commercial module essentially using the method described by Hara and Yuasa (7). [C-11] choline PET images showed abnormal prominent uptake in the middle of the abdomen (Fig. 2a and c). The maximal standardized uptake value (SUV) of this uptake was 6.97. This uptake was suspected to correspond to the sigmoid cancer. However, there was another strong accumulation close to this uptake, suspected to be physiological accumulation to the small intestine (Fig. 2c). It was not so easy to differentiate between these uptakes only by PET images. Fused PET/CT images could show clearly that one prominent accumulation corresponded to the thickening in the wall of sigmoid colon in CT images (Fig. 2d). Low abnormal uptake was observed in the pelvic space, which corresponded to myoma uteri. There was no other abnormal accumulation in

For reprints and all correspondence: Takashi Terauchi, Division of Cancer Screening, Research Center for Cancer Prevention and Screening, National Cancer Center, Tsukiji, Chuo-ku, 104-0045, Tokyo, Japan. E-mail: tterauch@ncc.go.jp




Figure 1. Endoscopic finding. Elevated lesion can be seen. Morphologic diagnosis is advanced colon cancer, type 1. Biopsy revealed well-differentiated adenocarcinoma.

the whole body (Fig. 2a). The patient underwent surgical resection of sigmoid colon and simple hysterectomy. Pathology revealed well-differentiated adenocarcinoma invading the subserosa with two metastatic lymph nodes in N1 group (2/30), which was stage IIIa according to TNM classification (8). These two metastatic lymph nodes were less than 10 mm in diameter and diagnosed as normal lymph nodes on CT images. Lymph node metastases were not detected by [C-11] choline PET. Pathology of the uterus revealed multiple leiomyomas. The patient received adjuvant chemotherapy with 5-FU and leucovorin, and was discharged 13 days after surgery.

DISCUSSION

[C-11] choline has been considered as a new PET radiopharmaceutical for tumor detection since Hara et al. reported the usefulness of [C-11] choline PET for detection of brain tumor in 1997 (1). Choline is one of the components of phosphatidylcholine, an essential element of phospholipids in the cell membrane. Malignant tumors usually exhibit a high proliferation of cells and thus are associated with increased metabolism of cell membrane components. This biochemical background will lead to an increased uptake of choline to the cancer cells (9,10). Moreover, it is assumed that, whether tumor cells are in hypoxia or in normoxia, the rate of [C-11] choline uptake in tumors is an indicator of the tumor proliferation rate, whereas in [F-18] fluoro-deoxyglucose (FDG), tumor hypoxia is closely associated with tumor uptake (11). In this view point, [C-11] choline PET might detect malignancies at an earlier stage than FDG-PET, which has spread worldwide as a gold standard of PET oncologic imaging, although further investigation is

still needed. Compared with FDG-PET, [C-11] choline PET has the advantage of providing a clear image at an earlier period (5). In FDG-PET, patients have to wait for 60 min or longer after tracer injection for tumor activity to reach the peak count. With [C-11] choline, however, blood clearance is rapid and tumor activity reaches a maximum at 3–5 min after injection. The initial intense uptake remains at a nearly constant level afterwards, thus enabling the high activity ratio to remain for more than 30 min, compared with the background. Another advantage of [C-11] choline PET is the lower exposure dose, which is estimated at approximately 2.5 mSv/370 MBq in contrast with 7 mSv/370 MBq for FDG-PET (12).

[C-11] choline PET has been reported to be useful for the detection of various malignant tumors such as lung, esophageal and gynecological cancers, and bone and soft sarcomas, as well as FDG-PET (1–6). Furthermore, [C-11] choline PET is reported to be superior to FDG-PET in the detection of brain tumor and prostate cancer (1,4). Ramirez de Molina et al. reported that choline kinase, which catalyzes the phosphorylation of choline, is upregulated in lung, prostate and colorectal cancers (13). Therefore, [C-11] choline is speculated to also detect colorectal cancer, which is one of the major targets of FDG-PET. However, there has been no report on [C-11] choline PET for detection of colorectal cancer. This is because it is generally accepted that [C-11] choline often accumulates in the small intestine and/or colon mucosa, in which cell turnover is very rapid. As a consequence, the various degrees of physiological uptake obscure accumulation to the tumor, which is similarly observed in FDG-PET (5). Hara reported that rectal cancer was visualized with [C-11] choline PET (14). In our case, abnormal [C-11] choline deposit to the sigmoid colon could be detected, although there was physiological accumulation to the small intestine near the cancerous lesion. There was no other significant accumulation to the colon. In our experience, physiological colon uptake of [C-11] choline tends to be lower than accumulation in the small intestine, whereas physiological colon uptake of FDG is often so much higher than small intestine uptake that the cancerous lesion cannot be depicted. That might be an advantage of [C-11] choline PET in colorectal cancer. It might be due to the considerably rapid turnover of epithelial cells in the small intestine, but the precise reason is unknown, prompting further investigation. It is sometimes confusing whether abdominal uptake is corresponding to the small intestine or the colon on [C-11] choline PET images as well as on FDG-PET images. In such cases, fused PET/CT images give great assistance in diagnosing correctly the location of abnormal uptake. Besides the detection of the primary site, staging is another important role of oncologic PET as an initial examination. There are some reports on [C-11] choline PET as a staging procedure for prostate cancer and bone and soft tissue sarcomas (15,16). In our case, lymph node metastases could not be detected by [C-11] choline PET. This was considered to be due to the




Figure 2. (a) MIP image of [C-11] choline positron emission tomography. Abnormal uptake of [C-11] choline is observed in the middle of the abdomen (arrow). There is no significant physiological uptake in the colon. (b–d) Coronal computed tomography (b), coronal positron emission tomography (c), coronal co-registered positron emission tomography/computed tomography hybrid image (d). Thickening in the wall of the sigmoid colon is observed in the computed tomography (b, arrow). Abnormal uptake of [C-11] choline is observed in the sigmoid colon corresponding to the wall thickening (c, d, arrow). Differentiation between physiological uptake in the small intestine (c, d, arrow head) and tumor uptake can be made by fused PET/CT images. PET, positron emission tomography; CT, computed tomography.

size of the lymph nodes (less than 10 mm in diameter). Hepatic metastasis, which is frequent in colon cancer, might be hardly depicted by [C-11] choline PET because [C-11] choline is observed physiologically in the liver. However,

[C-11] choline PET might be useful for detecting other metastases of colorectal cancer such as brain metastasis, pulmonary metastasis and bone metastasis with low background uptake.

In summary, here we described a case of advanced colon cancer. A whole body [C-11] choline PET/CT permitted detection of the primary site. However, further studies must be performed on staging, diagnosis for recurrence and evaluation for effect of treatment to confirm the usefulness of [C-11] choline PET for colorectal cancer.

Acknowledgments

This work was supported in part by a Grant from Foundation for Promotion of Cancer Research and the Grant-in-Aid for Cancer Research (18-11) from the Ministry of Health, Labour and Welfare, Japan.

Conflict of interest statement

None declared.

References

- Hara T, Kosaka N, Shinoura N, Kondo T. PET imaging of brain tumor with [methyl-11C]choline. *J Nucl Med* 1997;38:842-7.
- Hara T, Inagaki K, Kosaka N, Morita T. Sensitive detection of mediastinal lymph node metastasis of lung cancer with 11C-choline PET. *J Nucl Med* 2000;41:1507-13.
- Kobori O, Kirihara Y, Kosaka N, Hara T. Positron emission tomography of esophageal carcinoma using 11C-choline and 18F-fluorodeoxyglucose. *Cancer* 1999;86:1638-48.
- Hara T, Kosaka N, Kishi H. PET imaging of prostate cancer using carbon-11-choline. *J Nucl Med* 1998;39:990-5.
- Torizuka T, Kanno T, Futatsubashi M, Okada H, Yoshikawa E, Nakamura F, Takekuma M, Maeda M, Ouchi Y. Imaging of gynecologic tumors: comparison of 11C-choline PET with 18F-FDG PET. *J Nucl Med* 2003;44:1051-6.
- Zang H, Tian M, Oriuchi N, Higuchi T, Watanabe H, Aoki J, Tanada S, Endo K. ¹¹C-choline PET for the detection of bone and soft tissue tumors in comparison with FDG PET. *Nucl Med Commun* 2003;24:273-9.
- Hara T, Yuasa M. Automated synthesis of [C-11] choline, a positron-emitting tracer for tumor imaging. *Appl Radiat Isot* 1999;50:531-3.
- Japanese Society for Cancer of the Colon and Rectum. Japanese Classification of Colorectal Carcinoma, 1st English edn. Tokyo: Kanehara, 1997.
- Zeisel SH. Dietary choline: biochemistry, physiology and pharmacology. *A Rev Nutr* 1981;1:95-121.
- Podo F. Tumor phospholipid metabolism. *NMR Biomed* 1999;12:413-39.
- Khan N, Oriuchi N, Ninomiya H, Higuchi T, Kamada H, Endo K. Positron emission tomographic imaging with ¹¹C-choline in differential diagnosis of head and neck tumors: comparison with ¹⁸F-FDG PET. *Ann Nucl Med* 2004;18:409-17.
- Kotzzerke J, Prang J, Neumaier B, Volkmer B, Guhlmann A, Keinschmidt K, Hautmann R, Reske SN. Experience with carbon-11 choline positron emission tomography in prostate carcinoma. *Eur J Nucl Med* 2000;27:1415-19.
- Ramirez de Molina A, Rodriguez-Gonzalez A, Gutierrez R, Martinez-Pineiro L, Sanchez J, Bonilla F, Rosell R, Lacal J. Overexpression of choline kinase is a frequent feature in human tumor-derived cell lines and in lung, prostate, and colorectal human cancers. *Biochem Biophys Res Commun* 2002;296:580-3.
- Hara T, Kosaka N, Kondo T, Kishi H, Kobori O. Imaging of brain tumor, lung cancer, esophagus cancer, colon cancer, prostate cancer, and bladder cancer with [C-11]choline [abstract]. *J Nucl Med* 1997;38(Suppl):250.
- Maeda T, Tateishi U, Komiyama M, Fujimoto H, Watanabe S, Terauchi T, Moriyama N, Arai Y, Sugimura K, Kakizoe D. Distant metastasis of prostate cancer: early detection of recurrent tumor with dual-phase carbon-11 choline positron emission tomography/computed tomography in two cases. *Jpn J Clin Oncol* 2006;36(9):598-601.
- Tateishi U, Yamaguchi U, Maeda T, Seki K, Terauchi T, Kawai A, Arai Y, Moriyama N, Kakizoe T. Staging performance of carbon-11 choline positron emission tomography/computed tomography in patients with bone and soft tissue sarcoma: comparison with conventional imaging. *Cancer Sci* 2006;97(10):1125-8.



and Other Interventional Techniques

Wound infection after elective laparoscopic surgery for colorectal carcinoma

Seiichiro Yamamoto, Shin Fujita, Takayuki Akasu, Seiji Ishiguro, Yutaka Kobayashi, Yoshihiro Moriya

Division of Colorectal Surgery, National Cancer Center Hospital, 5-1-1, Tsukiji, Chuo-ku, Tokyo, 104-0045, Japan

Received: 26 December 2006/Accepted: 22 January 2007/Online publication: 19 May 2007

Abstract

Background: The aim of this study was to evaluate various clinical parameters that would influence the occurrence of wound infection (WI) in elective laparoscopic surgery (LS) for colorectal carcinoma.

Methods: The study included 290 patients who underwent LS between June 2001 and December 2005. WI was diagnosed within 30 days of the operation, and both superficial and deep incision surgical site infection were evaluated together.

Results: Eighteen (6.2%) were diagnosed with WI. Of the infected patients, nine (50%) had WI at the extraction site, six (33%) at the port site of the drainage tube, and three (17%) at the supraumbilical incision. Following bivariate analysis, the variables of stoma creation, intraoperative hypotension, and length of operation were selected for multivariate analysis as their *P* values were <0.2, the predominant cutoff, and stoma creation and intraoperative hypotension were independently predictive of developing WI. Regarding the duration of postoperative hospital stay, there was no significant difference between patients with or without WI.

Conclusions: Stoma creation and intraoperative hypotension were independent risk factors for WI. The results obtained in this study should be considered in an effort to prevent WI in LS for colorectal carcinoma, although these risk factors need further evaluation.

Key words: Laparoscopic surgery — Colorectal carcinoma — Wound infection

In elective open colorectal surgery for colorectal malignancy, the most frequent complication is wound infection (WI) [14, 22]. It goes without saying that if WI

occurs, hospital costs (increased length of stay, antibiotics, dressing supplies, and nursing charges) and outpatient care (outpatient follow-up, self-dressing supplies, and antibiotics) will increase, thus increasing the total medical cost. In addition, it will delay the patient's return to work, causing a social loss. For these reasons, various risk factors possibly related to the occurrence of WI have been investigated mainly in cases of elective open colorectal surgery for colorectal malignancy [16–18].

In recently reported randomized controlled trials (RCTs) that compared the oncological outcomes of conventional open surgery (OS) versus laparoscopic surgery (LS) in colorectal carcinoma cases, LS was proven to be comparable to OS, and hence the number of patients receiving LS is expected to increase in the coming years [4, 12, 13]. In RCTs comparing OS versus LS for colorectal carcinoma in terms of short-term outcome that have been reported to date, the incidence of WI was 4–14% for LS versus 5–17% for OS; one third of those reports showed a significantly lower incidence of WI for LS compared to OS, and the rest showed no difference [2, 7, 8, 12, 13, 23]. As LS involves the pneumoperitoneum with CO₂ and delivery of the intestinal tract through a small incision, risk factors influencing WI occurrence in LS may be different from those in OS; however, only a few studies to date have investigated the risk factors related to the occurrence of WI in cases of LS for colorectal carcinoma.

At our institution, after having accumulated enough experience in LS for early colon carcinoma, we have gradually expanded the indications for LS. In June 2001, a single laparoscopic surgical team instructed by one surgeon (SY) began to perform LS, and postoperative management procedures were standardized, and began to expand the use of LS to include T3 colon/upper rectal carcinoma and T1-T2 middle/lower rectal carcinomas. As a consequence, the complication rate and mean length of hospitalization have been reduced [26, 27].

The purpose of this study was to evaluate various clinical parameters that would influence WI occurrence

Correspondence to: Seiichiro Yamamoto

in LS by accumulating occurrences of WI in elective LS cases via prospective entries at a single institution and evaluating its risk factors.

Patients and methods

Between June 2001 and December 2005, we performed 290 consecutive LS for selected patients with colorectal carcinoma. As the oncological safety of LS in colorectal carcinoma patients remains to be established, candidates for radical LS were patients who were preoperatively diagnosed with T1 or T2. Additionally, we included patients who were preoperatively diagnosed with T3 but wished to undergo LS, as well as those with colon or upper rectal carcinoma for which palliative resection was considered necessary. Contraindications for LS at our institution included the following groups: tumors larger than 6 cm, a prior history of extensive adhesions, severe obesity (body mass index $> 32 \text{ kg/m}^2$), intestinal obstruction, and patients who did not consent to LS. We defined conversion to OS as any incision greater than 7 cm, excluding cases in which the incision was enlarged due to a large specimen that could not be removed through a 7 cm incision.

Patients underwent mechanical bowel preparation with 2 liters of polyethylene glycol electrolyte solution one or two days before operation, and this decision differed according to the study period. Preoperative antimicrobial administration was given intravenously within 30 minutes before skin incision in all cases. Indications for postoperative and intraoperative repeated dosing differed according to the study period. Similarly, antimicrobial agents differed according to the study period; however, the antimicrobial agent given intraoperatively and/or postoperatively was the same antimicrobial prophylaxis given preoperatively in all patients.

Our LS techniques have previously been thoroughly described [26, 27]. During the externalized portion of the surgery, a plastic wound protector was applied to the abdominal extraction incision routinely. Wound irrigation was not performed in this series. All wounds were classified as clean-contaminated (bowel was opened without spilling the contents).

WI was diagnosed within 30 days of the operation according to the criteria of the Centers for Disease Control and Prevention (CDC) [9]. Briefly, superficial surgical site infections only involve the skin and subcutaneous tissue, whereas deep surgical site infections occur when the incisional wound involves the muscle and fascial layers but not the organ space. WI was characterized by wound erythema, cellulitis, localized pain, swelling, tenderness, or purulent or culture-positive wound discharge. All the patients visited outpatient clinic about three weeks after the day of discharge, and their wound conditions were checked. Superficial and deep incision surgical site infections were evaluated together under the umbrella term of WI [17].

Patients' theatre records were reviewed retrospectively. Other data were prospectively recorded in the divisional database. The parameters analyzed included: gender, age, body mass index, prior abdominal surgery, smoking status, ASA classification [15], preoperative hemoglobin, total protein, and hemoglobin A1c level. Pathological staging was performed according to Dukes' stage. Regarding the perioperative/operative characteristics, operative time, antimicrobial prophylaxis administration, minimum body temperature and intraoperative hypotension during surgery, intraoperative blood loss, presence or absence of a drainage tube for anastomoses, duration of drainage, stoma creation, type of anastomosis, date of preoperative mechanical bowel preparation, and postoperative hospital stay were analyzed. Stoma creation included both patients with endocolostomy who underwent laparoscopic abdominoperineal resection, and patients with covering ileostomy.

Perioperative steroid and insulin use for comorbid conditions was not evaluated because only three patients received perioperative steroid and only 10 patients required insulin, and WI did not occur in these patients. Similarly, the transfusion of cellular or plasma products was not evaluated, because only two patients without WI received transfusion in this series.

Statistical analysis was performed using Student's *t*-test, Fisher's exact test, the chi-square test, and the Mann-Whitney *U* test, as appropriate. Multivariate analysis was performed by logistic regression methods using independent variables with a *P* value < 0.2 by bivariate statistics. A *P* value of less than 0.05 was considered significant.

Results

During the study period, 290 patients underwent elective LS, and 18 (6.2%) were diagnosed with WI. Of the infected patients, nine (50%) had WI at the extraction site, six (33%) at the port site of the drainage tube, and three (17%) at the supraumbilical incision.

In bivariate analysis, patients were divided into those with or without WI and compared. Patient demographics are summarized in Table 1. There were no significant differences in baseline characteristics among groups.

Perioperative/operative results are shown in Table 2. All procedures were completed laparoscopically in this series, and we consider our exclusion criteria for LS (extensive adhesions and severe obesity) contributed to this lower conversion rate. Only stoma creation was significantly associated with the development of WI. However, there was a trend toward developing WI if the patient developed intraoperative hypotension ($P = 0.071$) or the operation exceeded 240 minutes ($P = 0.123$). Regarding the duration of postoperative hospital stay, there was no significant difference between the two groups ($P = 0.131$).

Following bivariate analysis, the variables of stoma creation, intraoperative hypotension, and length of operation were selected for multivariate analysis as their *P* values were < 0.2 , the predominant cutoff for inclusion. Table 3 summarizes the results of multivariate analysis, and stoma creation and intraoperative hypotension were independently predictive of developing WI.

Discussion

This is the first investigation demonstrating stoma creation and intraoperative hypotension as independent risk factors for WI in nearly 300 cases of colorectal carcinoma undergoing LS. In this study, clinical parameters that have conventionally been reported as risk factors for WI were evaluated for cases of elective LS for colorectal carcinoma. As a result, we found that stoma creation and intraoperative hypotension were independent risk factors. The results obtained in this study should be considered in an effort to prevent WI in LS for colorectal carcinoma, although these risk factors need further evaluation.

The incidence of WI for LS in this study was 7.6%, which is the same as in previous reports. In this series, the indications for LS were limited as mentioned, but at our hospital, the incidence of WI for OS is more than 10%, which is higher than that for LS. Regarding the incidence of WI in LS and OS, based on the National Nosocomial Infections Surveillance System's surgical patient surveillance component protocol, CDC reported that the incidence of WI after colon surgery was significantly lower in the LS group than in the OS group [6]. In RCTs comparing OS with LS for colorectal carcinoma in terms of short-term outcome that have been reported to date, the incidence of WI was 4–14% for LS versus 5–17% for OS; one third of those reports showed a significantly lower incidence of WI for LS compared to

Table 1. Correlation between wound infection and clinicopathologic features^a

	Wound infection		P value
	Positive (n = 18)	Negative (n = 272)	
Sex ratio (male:female)	13:5	153:119	.224
Mean age	57.9 (35-73)	61.0 (30-88)	.234
Mean body mass index (kg/m ²)	23.0 (17.3-27.9)	23.0 (15.1-32.4)	1.000
Prior abdominal surgery (percentage)	3 (16.7)	62 (22.8)	.772
Smoking status			.754
Nonsmoker	14	222	
Smoker	4	50	
ASA classification			.455
ASA I	13	165	
ASA II	5	82	
ASA III		25	
Mean preoperative hemoglobin (g/dL)	13.6 (12.2-15.4)	13.4 (8.0-16.4)	.578
Mean preoperative total protein (g/dL)	7.1 (6.5-8.1)	7.0 (5.5-8.7)	.313
Mean preoperative hemoglobin A1c (g/dL)	5.2 (4.6-5.9)	5.2 (3.1-9.5)	.917
Dukes' stage (n)			.446
A	14	185	
B	0	25	
C	3	49	
D	1	13	
Location (n)			.229
Colon:rectum	12:6	217:55	
Laparoscopic colorectal procedures (n)			
Ileocecal resection	1	28	
Right hemicolectomy	2	52	
Transverse colectomy		1	
Left hemicolectomy		2	
Descending colectomy	2	12	
Sigmoid colectomy	4	91	
Partial resection	5	30	
Anterior resection with DST	4	49	
Anterior resection with ISR-CAA	1	4	
Abdominoperineal resection		2	
Hartmann		1	
Transverse-coloplasty pouch	2	3	
Covering ileostomy	3	12	

ASA, American Society of Anesthesiologists; DST, double-stapling technique; ISR-CAA, intersphincteric rectal resection and handsewn coloanal anastomosis

OS, and the rest showed no difference [2, 7, 8, 12, 13, 23]. In addition, meta-analysis of these trials demonstrated a significantly lower incidence for LS [1]. From the results of these studies, the incidence of WI in LS is so far considered to be lower than that of OS.

At the same time, there are some reports that, when WI occurs, the treatment costs are significantly higher for OS than LS [2]. In this study, even with WI, patients undergoing LS did not require prolonged hospitalization because of smaller wounds and were able to dress their own wounds after being discharged. Even when WI occurs in LS, the maximum range of the infected area can only be 6-7 cm, which is a great advantage of LS.

To date, many studies have reported the immunological superiority of LS compared to OS [19, 25]. In our LS series, after completing manipulation of the pneumoperitoneum with CO₂, one port site is used to make a skin incision as the extraction site, through which necessary manipulations including tumor removal and anastomosis are done, and immediately thereafter, we close the wound to the fascia. Therefore, the extraction site was usually open for a total of 20-30 minutes; the

procedure rarely requires more than 60 minutes. Consequently, the net length of time during which a small laparotomy wound is made is shorter in LS than in OS, and this may possibly have a favorable influence on wound healing after LS. Regarding the plastic wound protectors used to prevent port site recurrence, Kercher et al. [10] reported that a wound protector did not significantly diminish the rate of WI at the extraction site; however, it is currently unthinkable to perform LS for malignant diseases without a plastic wound protector, and the significance of plastic wound protectors in terms of WI may need to be investigated in benign diseases.

With regard to the risk factors for WI in cases of colorectal carcinoma, various parameters have been evaluated in the past [16, 17, 18]. In this study, those many parameters were evaluated, and interestingly, we found that stoma creation and intraoperative hypotension were the only independent risk factors. Although there are some reports suggesting that stoma creation or intraoperative hypotension are risk factors for WI in elective open colorectal resections [11, 17, 20], the other many parameters that have conventionally been re-

Table 2. Correlation between wound infection and perioperative/operative characteristics^a

	Wound infection		P value
	Positive (n = 18)	Negative (n = 272)	
Length of operation			.123
≤ 240 minutes	9	186	
> 240 minutes	9	86	
Antimicrobial prophylaxis administration			
Single dosing:multiple dosing (intra- and post-operative)	11:7	176:96	.802
Intraoperative repeated dosing (LOO > 180 minutes)	2/14 (14.3%)	22/207 (10.6%)	.653
First generation: second generation	6:12	92:180	1.000
Mean minimum body temperature (°C)	35.8 (34.9–36.7)	35.9 (34.5–37.4)	.854
Intraoperative hypotension (SBP < 80mmHg)	7	54	.071
Mean operative blood loss (mL)	64 (6–250)	61 (3–217)	.839
Drainage of anastomoses	17	268	.276
Mean duration of drainage (days)	4.5 (0–8)	4.6 (0–8)	.638
Stoma creation	4	14	.019
Anastomosis			.470
Double-stapling technique	8	123	
FETE + Handsewn anastomosis	10	142	
Handsewn per anum coloanal anastomosis	1	4	
Colon preparation			.380
One day before surgery	16	210	
Two days before surgery	2	62	
Median hospital stay (days)	8 (7–23)	8 (7–21)	.131

LOO, length of operation; SBP, systolic blood pressure; FETE, functional end-to-end anastomosis

^a Values in parentheses are ranges unless indicated otherwise

Table 3. Multivariable analysis of factors affecting wound infection

Independent Predictors	Odds ratio	Confidence intervals	P value
Stoma creation (yes/no)	5.275	1.196–23.263	.028
Intraoperative hypotension (SBP < 80mmHg/SBP ≥ 80mmHg)	3.171	1.099–9.147	.033

SBP, systolic blood pressure

garded as risk factors for WI were negative. Unfortunately, we were unable to evaluate matters such as the necessity of mechanical bowel preparation and the significance of preoperative oral prophylaxis, because all the patients underwent mechanical bowel preparation without oral antibiotics. In recent years, the significance of mechanical bowel preparation has been investigated in RCTs, and there are increasing reports that mechanical bowel preparation is not necessary; however many of those trials investigated cases of elective OS [3, 16]. As to whether the same result would be obtained in LS cases, further investigation is needed.

In this study, surgery was performed without oral antimicrobial prophylaxis administration in all patients. We do not administer oral antimicrobial prophylaxis in either OS or LS, for fear that clostridium difficile colitis or gastrointestinal symptoms including nausea, vomiting, abdominal pain, and diarrhea might be caused by oral antibiotics [5, 24]. In such cases, intravenous antimicrobial prophylaxis administration is required, but factors such as the type of antibiotic administered and its dosing frequency may influence the occurrence of WI. As the type of antibiotic used and its dosing frequency vary according to the study period, the influence of these

factors was evaluated in this study. We found no difference in the incidence of WI between single dosing⁶ and multiple dosing as well as between the presence and absence of intraoperative repeated dosing in cases with an operative time of four hours or more. With regard to the type of antibiotic administered, no difference was seen in the incidence of WI between first and second generation. Whether these items influence the occurrence of WI in LS cases is a question that requires further investigation.

In addition, the significance of drainage tubes should also be investigated. In our study, WI at the port site through which a drainage tube was inserted accounted for one-third of all WIs. Some reports say that drainage tubes are not necessary in elective colorectal surgery [21], but we leave drainage tubes in place for 4–5 days after operation to obtain information on postoperative bleeding and postoperative peritonitis associated with anastomotic leakage. However, because of the low frequency of these complications in LS, we now remove drainage tubes as early as the second or third postoperative day. Further investigation is also required with regard to the necessity of drainage tubes in LS.

The results of this study suggest the importance of maintaining normotension intraoperatively to reduce the incidence of WI, and we must recognize that this variable needs to be further investigated [17]. Moreover, if a stoma needs to be created, the port site should be positioned away from the stoma site as far as possible, and in patients with a created stoma, the wound may need to be completely covered with plastic film or other material. On the other hand, if WI occurs, hospital costs will be reduced by opening the wound early and managing it with self-dressing gauze changes.

In conclusion, as a result of our investigation into cases of LS, we found that stoma creation and intra-operative hypotension were independent risk factors for WI. Various factors are related to the occurrence of WI. It is necessary to further investigate risk factors for WI in LS cases so that efforts can be made to effectively prevent WI by measures such as increasing the dosing frequency of antimicrobial prophylaxis only in patients at higher risk for WI.

References

1. Abraham NS, Young JM, Solomon MJ (2004) Meta-analysis of short-term outcomes after laparoscopic resection for colorectal cancer. *Br J Surg* 91: 1111–1124
2. Braga M, Vignali A, Zuliani W, Frasson M, Di Serio C, Di Carlo V (2005) Laparoscopic versus open colorectal surgery: cost-benefit analysis in a single-center randomized trial. *Ann Surg* 242: 890–896
3. Bucher P, Gervaz P, Soravia C, Mermillod B, Erne M, Morel P (2005) Randomized clinical trial of mechanical bowel preparation versus no preparation before elective left-sided colorectal surgery. *Br J Surg* 92: 409–414
4. Clinical Outcomes of Surgical Therapy Study Group (2004) A comparison of laparoscopically assisted and open colectomy for colon cancer. *N Engl J Med* 350: 2050–2059
5. Espin-Basany E, Sanchez-Garcia JL, Lopez-Cano M, Lozoya-Trujillo R, Medarde-Ferrer M, Armadans-Gil L, Alemany-Vilches L, Armengol-Carrasco M (2005) Prospective, randomized study on antibiotic prophylaxis in colorectal surgery. Is it really necessary to use oral antibiotics? *Int J Colorectal Dis* 20: 542–546
6. Gaynes RP, Culver DH, Horan TC, Edwards JR, Richards C, Tolson JS (2001) National Nosocomial Infections Surveillance System. Surgical site infection (SSI) rates in the United States, 1992–1998: the National Nosocomial Infections Surveillance System basic SSI risk index. *Clin Infect Dis* 33: S69–77
7. Guillou PJ, Quirke P, Thorpe H, Walker J, Jayne DG, Smith AM, Heath RM, Brown JM, MRC CLASICC trial group (2005) Short-term endpoints of conventional versus laparoscopic-assisted surgery in patients with colorectal cancer (MRC CLASICC trial): multicentre, randomised controlled trial. *Lancet* 365: 1718–1726
8. Hasegawa H, Kabeshima Y, Watanabe M, Yamamoto S, Kitajima M (2003) Randomized controlled trial of laparoscopic versus open colectomy for advanced colorectal cancer. *Surg Endosc* 17: 636–640
9. Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG (1992) CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Infect Control Hosp Epidemiol* 13: 606–608
10. Kercher KW, Nguyen TH, Harold KL, Poplin ME, Matthews BD, Sing RF, Heniford BT (2004) Plastic wound protectors do not affect wound infection rates following laparoscopic-assisted colectomy. *Surg Endosc* 18: 148–151
11. Konishi T, Watanabe T, Kishimoto J, Nagawa H (2006) Elective colon and rectal surgery differ in risk factors for wound infection: results of prospective surveillance. *Ann Surg* 244: 758–763
12. Lacy AM, Garcia-Valdecasas JC, Delgado S, Castells A, Taura P, Piqué JM, Visa J (2002) Laparoscopy-assisted colectomy versus open colectomy for treatment of non-metastatic colon cancer: a randomised trial. *Lancet* 359: 2224–2229
13. Leung KL, Kwok SPY, Lam SCW, Lee JFY, Yiu RYC, Ng SSM, Lai PBS, Lau WY (2004) Laparoscopic resection of rectosigmoid carcinoma: prospective randomised trial. *Lancet* 363: 1187–1192
14. Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR (1999) The Hospital Infection Control Practices Advisory Committee. Guideline for prevention of surgical site infection, 1999. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 20: 250–278
15. Owens WD, Felts JA, Spitznagel EL Jr (1978) ASA physical status classifications: a study of consistency of ratings. *Anesthesiology* 49: 239–243
16. Slim K, Vicaut E, Panis Y, Chipponi J (2004) Meta-analysis of randomized clinical trials of colorectal surgery with or without mechanical bowel preparation. *Br J Surg* 91: 1125–1130
17. Smith RL, Bohl JK, McElearney ST, Friel CM, Barclay MM, Sawyer RG, Foley EF (2004) Wound infection after elective colorectal resection. *Ann Surg* 239: 599–607
18. Sørensen LT, Hemmingsen U, Kallehave F, Wille-Jørgensen P, Kjaergaard J, Møller LN, Jørgensen T (2005) Risk factors for tissue and wound complications in gastrointestinal surgery. *Ann Surg* 241: 654–658
19. Tang CL, Eu KW, Tai BC, Soh JG, MacHin D, Seow-Choen F (2001) Randomized clinical trial of the effect of open versus laparoscopically assisted colectomy on systemic immunity in patients with colorectal cancer. *Br J Surg* 88: 801–807
20. Tang R, Chen H, Wang Y, Changchien C, Chen JS, Hsu KC, Chiang JM, Wang JY (2001) Risk factors for surgical site infection after elective resection of the colon and rectum: a single-center prospective study of 2,809 consecutive patients. *Ann Surg* 234: 181–189
21. Urbach DR, Kennedy ED, Cohen MM (1999) Colon and rectal anastomoses do not require routine drainage: a systematic review and meta-analysis. *Ann Surg* 229: 174–180
22. Weiss CA 3rd, Statz CL, Dahms RA, Remucal MJ, Dunn DL, Beilman GJ (1999) Six years of surgical wound infection surveillance at a tertiary care center: review of the microbiologic and epidemiological aspects of 20,007 wounds. *Arch Surg* 134: 1041–1048
23. Winslow ER, Fleshman JW, Birnbaum EH, Brunt LM (2002) Wound complications of laparoscopic vs open colectomy. *Surg Endosc* 16: 1420–1425
24. Wren SM, Ahmed N, Jamal A, Safadi BY (2005) Preoperative oral antibiotics in colorectal surgery increase the rate of *Clostridium difficile* colitis. *Arch Surg* 140: 752–756
25. Wu FP, Sietses C, von Blomberg BM, van Leeuwen PA, Meijer S, Cuesta MA (2003) Systemic and peritoneal inflammatory response after laparoscopic or conventional colon resection in cancer patients: a prospective, randomized trial. *Dis Colon Rectum* 46: 147–155
26. Yamamoto S, Fujita S, Akasu T, Moriya Y (2004) A comparison of the complication rates between laparoscopic colectomy and laparoscopic low anterior resection. *Surg Endosc* 18: 1447–1451
27. Yamamoto S, Fujita S, Akasu T, Moriya Y (2005) Safety of laparoscopic intracorporeal rectal transection with double-stapling technique anastomosis. *Surg Laparosc Endosc Percutan Tech* 15: 70–74



and Other Interventional Techniques

Wound infection after elective laparoscopic surgery for colorectal carcinoma

Seiichiro Yamamoto, Shin Fujita, Takayuki Akasu, Seiji Ishiguro, Yutaka Kobayashi, Yoshihiro Moriya

Division of Colorectal Surgery, National Cancer Center Hospital, 5-1-1, Tsukiji, Chuo-ku, Tokyo, 104-0045, Japan

Received: 26 December 2006/Accepted: 22 January 2007/Online publication: 19 May 2007

Abstract

Background: The aim of this study was to evaluate various clinical parameters that would influence the occurrence of wound infection (WI) in elective laparoscopic surgery (LS) for colorectal carcinoma.

Methods: The study included 290 patients who underwent LS between June 2001 and December 2005. WI was diagnosed within 30 days of the operation, and both superficial and deep incision surgical site infection were evaluated together.

Results: Eighteen (6.2%) were diagnosed with WI. Of the infected patients, nine (50%) had WI at the extraction site, six (33%) at the port site of the drainage tube, and three (17%) at the supraumbilical incision. Following bivariate analysis, the variables of stoma creation, intraoperative hypotension, and length of operation were selected for multivariate analysis as their *P* values were <0.2, the predominant cutoff, and stoma creation and intraoperative hypotension were independently predictive of developing WI. Regarding the duration of postoperative hospital stay, there was no significant difference between patients with or without WI.

Conclusions: Stoma creation and intraoperative hypotension were independent risk factors for WI. The results obtained in this study should be considered in an effort to prevent WI in LS for colorectal carcinoma, although these risk factors need further evaluation.

Key words: Laparoscopic surgery — Colorectal carcinoma — Wound infection

In elective open colorectal surgery for colorectal malignancy, the most frequent complication is wound infection (WI) [14, 22]. It goes without saying that if WI

occurs, hospital costs (increased length of stay, antibiotics, dressing supplies, and nursing charges) and outpatient care (outpatient follow-up, self-dressing supplies, and antibiotics) will increase, thus increasing the total medical cost. In addition, it will delay the patient's return to work, causing a social loss. For these reasons, various risk factors possibly related to the occurrence of WI have been investigated mainly in cases of elective open colorectal surgery for colorectal malignancy [16–18].

In recently reported randomized controlled trials (RCTs) that compared the oncological outcomes of conventional open surgery (OS) versus laparoscopic surgery (LS) in colorectal carcinoma cases, LS was proven to be comparable to OS, and hence the number of patients receiving LS is expected to increase in the coming years [4, 12, 13]. In RCTs comparing OS versus LS for colorectal carcinoma in terms of short-term outcome that have been reported to date, the incidence of WI was 4–14% for LS versus 5–17% for OS; one third of those reports showed a significantly lower incidence of WI for LS compared to OS, and the rest showed no difference [2, 7, 8, 12, 13, 23]. As LS involves the pneumoperitoneum with CO₂ and delivery of the intestinal tract through a small incision, risk factors influencing WI occurrence in LS may be different from those in OS; however, only a few studies to date have investigated the risk factors related to the occurrence of WI in cases of LS for colorectal carcinoma.

At our institution, after having accumulated enough experience in LS for early colon carcinoma, we have gradually expanded the indications for LS. In June 2001, a single laparoscopic surgical team instructed by one surgeon (SY) began to perform LS, and postoperative management procedures were standardized, and began to expand the use of LS to include T3 colon/upper rectal carcinoma and T1–T2 middle/lower rectal carcinomas. As a consequence, the complication rate and mean length of hospitalization have been reduced [26, 27].

The purpose of this study was to evaluate various clinical parameters that would influence WI occurrence

Correspondence to: Seiichiro Yamamoto

in LS by accumulating occurrences of WI in elective LS cases via prospective entries at a single institution and evaluating its risk factors.

Patients and methods

Between June 2001 and December 2005, we performed 290 consecutive LS for selected patients with colorectal carcinoma. As the oncological safety of LS in colorectal carcinoma patients remains to be established, candidates for radical LS were patients who were preoperatively diagnosed with T1 or T2. Additionally, we included patients who were preoperatively diagnosed with T3 but wished to undergo LS, as well as those with colon or upper rectal carcinoma for which palliative resection was considered necessary. Contraindications for LS at our institution included the following groups: tumors larger than 6 cm, a prior history of extensive adhesions, severe obesity (body mass index $> 32 \text{ kg/m}^2$), intestinal obstruction, and patients who did not consent to LS. We defined conversion to OS as any incision greater than 7 cm, excluding cases in which the incision was enlarged due to a large specimen that could not be removed through a 7 cm incision.

Patients underwent mechanical bowel preparation with 2 liters of polyethylene glycol electrolyte solution one or two days before operation, and this decision differed according to the study period. Preoperative antimicrobial administration was given intravenously within 30 minutes before skin incision in all cases. Indications for postoperative and intraoperative repeated dosing differed according to the study period. Similarly, antimicrobial agents differed according to the study period; however, the antimicrobial agent given intraoperatively and/or postoperatively was the same antimicrobial prophylaxis given preoperatively in all patients.

Our LS techniques have previously been thoroughly described [26, 27]. During the externalized portion of the surgery, a plastic wound protector was applied to the abdominal extraction incision routinely. Wound irrigation was not performed in this series. All wounds were classified as clean-contaminated (bowel was opened without spilling the contents).

WI was diagnosed within 30 days of the operation according to the criteria of the Centers for Disease Control and Prevention (CDC) [9]. Briefly, superficial surgical site infections only involve the skin and subcutaneous tissue, whereas deep surgical site infections occur when the incisional wound involves the muscle and fascial layers but not the organ space. WI was characterized by wound erythema, cellulitis, localized pain, swelling, tenderness, or purulent or culture-positive wound discharge. All the patients visited outpatient clinic about three weeks after the day of discharge, and their wound conditions were checked. Superficial and deep incision surgical site infections were evaluated together under the umbrella term of WI [17].

Patients' theatre records were reviewed retrospectively. Other data were prospectively recorded in the divisional database. The parameters analyzed included: gender, age, body mass index, prior abdominal surgery, smoking status, ASA classification [15], preoperative hemoglobin, total protein, and hemoglobin A1c level. Pathological staging was performed according to Dukes' stage. Regarding the perioperative/operative characteristics, operative time, antimicrobial prophylaxis administration, minimum body temperature and intraoperative hypotension during surgery, intraoperative blood loss, presence or absence of a drainage tube for anastomoses, duration of drainage, stoma creation, type of anastomosis, date of preoperative mechanical bowel preparation, and postoperative hospital stay were analyzed. Stoma creation included both patients with endocolostomy who underwent laparoscopic abdominoperineal resection, and patients with covering ileostomy.

Perioperative steroid and insulin use for comorbid conditions was not evaluated because only three patients received perioperative steroid and only 10 patients required insulin, and WI did not occur in these patients. Similarly, the transfusion of cellular or plasma products was not evaluated, because only two patients without WI received transfusion in this series.

Statistical analysis was performed using Student's *t*-test, Fisher's exact test, the chi-square test, and the Mann-Whitney *U* test, as appropriate. Multivariate analysis was performed by logistic regression methods using independent variables with a *P* value < 0.2 by bivariate statistics. A *P* value of less than 0.05 was considered significant.

Results

During the study period, 290 patients underwent elective LS, and 18 (6.2%) were diagnosed with WI. Of the infected patients, nine (50%) had WI at the extraction site, six (33%) at the port site of the drainage tube, and three (17%) at the supraumbilical incision.

In bivariate analysis, patients were divided into those with or without WI and compared. Patient demographics are summarized in Table 1. There were no significant differences in baseline characteristics among groups.

Perioperative/operative results are shown in Table 2. All procedures were completed laparoscopically in this series, and we consider our exclusion criteria for LS (extensive adhesions and severe obesity) contributed to this lower conversion rate. Only stoma creation was significantly associated with the development of WI. However, there was a trend toward developing WI if the patient developed intraoperative hypotension ($P = 0.071$) or the operation exceeded 240 minutes ($P = 0.123$). Regarding the duration of postoperative hospital stay, there was no significant difference between the two groups ($P = 0.131$).

Following bivariate analysis, the variables of stoma creation, intraoperative hypotension, and length of operation were selected for multivariate analysis as their *P* values were < 0.2 , the predominant cutoff for inclusion. Table 3 summarizes the results of multivariate analysis, and stoma creation and intraoperative hypotension were independently predictive of developing WI.

Discussion

This is the first investigation demonstrating stoma creation and intraoperative hypotension as independent risk factors for WI in nearly 300 cases of colorectal carcinoma undergoing LS. In this study, clinical parameters that have conventionally been reported as risk factors for WI were evaluated for cases of elective LS for colorectal carcinoma. As a result, we found that stoma creation and intraoperative hypotension were independent risk factors. The results obtained in this study should be considered in an effort to prevent WI in LS for colorectal carcinoma, although these risk factors need further evaluation.

The incidence of WI for LS in this study was 7.6%, which is the same as in previous reports. In this series, the indications for LS were limited as mentioned, but at our hospital, the incidence of WI for OS is more than 10%, which is higher than that for LS. Regarding the incidence of WI in LS and OS, based on the National Nosocomial Infections Surveillance System's surgical patient surveillance component protocol, CDC reported that the incidence of WI after colon surgery was significantly lower in the LS group than in the OS group [6]. In RCTs comparing OS with LS for colorectal carcinoma in terms of short-term outcome that have been reported to date, the incidence of WI was 4–14% for LS versus 5–17% for OS; one third of those reports showed a significantly lower incidence of WI for LS compared to

Table 1. Correlation between wound infection and clinicopathologic features^a

	Wound infection		P value
	Positive (n = 18)	Negative (n = 272)	
Sex ratio (male:female)	13:5	153:119	.224
Mean age	57.9 (35-73)	61.0 (30-88)	.234
Mean body mass index (kg/m ²)	23.0 (17.3-27.9)	23.0 (15.1-32.4)	1.000
Prior abdominal surgery (percentage)	3 (16.7)	62 (22.8)	.772
Smoking status			.754
Nonsmoker	14	222	
Smoker	4	50	
ASA classification			.455
ASA I	13	165	
ASA II	5	82	
ASA III		25	
Mean preoperative hemoglobin (g/dL)	13.6 (12.2-15.4)	13.4 (8.0-16.4)	.578
Mean preoperative total protein (g/dL)	7.1 (6.5-8.1)	7.0 (5.5-8.7)	.313
Mean preoperative hemoglobin A1c (g/dL)	5.2 (4.6-5.9)	5.2 (3.1-9.5)	.917
Dukes' stage (n)			.446
A	14	185	
B	0	25	
C	3	49	
D	1	13	
Location (n)			.229
Colon:rectum	12:6	217:55	
Laparoscopic colorectal procedures (n)			
Ileocecal resection	1	28	
Right hemicolectomy	2	52	
Transverse colectomy		1	
Left hemicolectomy		2	
Descending colectomy	2	12	
Sigmoid colectomy	4	91	
Partial resection	5	30	
Anterior resection with DST	4	49	
Anterior resection with ISR-CAA	1	4	
Abdominoperineal resection		2	
Hartmann		1	
Transverse-coloplasty pouch	2	3	
Covering ileostomy	3	12	

ASA, American Society of Anesthesiologists; DST, double-stapling technique; ISR-CAA, intersphincteric rectal resection and handsewn coloanal anastomosis

OS, and the rest showed no difference [2, 7, 8, 12, 13, 23]. In addition, meta-analysis of these trials demonstrated a significantly lower incidence for LS [1]. From the results of these studies, the incidence of WI in LS is so far considered to be lower than that of OS.

At the same time, there are some reports that, when WI occurs, the treatment costs are significantly higher for OS than LS [2]. In this study, even with WI, patients undergoing LS did not require prolonged hospitalization because of smaller wounds and were able to dress their own wounds after being discharged. Even when WI occurs in LS, the maximum range of the infected area can only be 6-7 cm, which is a great advantage of LS.

To date, many studies have reported the immunological superiority of LS compared to OS [19, 25]. In our LS series, after completing manipulation of the pneumoperitoneum with CO₂, one port site is used to make a skin incision as the extraction site, through which necessary manipulations including tumor removal and anastomosis are done, and immediately thereafter, we close the wound to the fascia. Therefore, the extraction site was usually open for a total of 20-30 minutes; the

procedure rarely requires more than 60 minutes. Consequently, the net length of time during which a small laparotomy wound is made is shorter in LS than in OS, and this may possibly have a favorable influence on wound healing after LS. Regarding the plastic wound protectors used to prevent port site recurrence, Kercher et al. [10] reported that a wound protector did not significantly diminish the rate of WI at the extraction site; however, it is currently unthinkable to perform LS for malignant diseases without a plastic wound protector, and the significance of plastic wound protectors in terms of WI may need to be investigated in benign diseases.

With regard to the risk factors for WI in cases of colorectal carcinoma, various parameters have been evaluated in the past [16, 17, 18]. In this study, those many parameters were evaluated, and interestingly, we found that stoma creation and intraoperative hypotension were the only independent risk factors. Although there are some reports suggesting that stoma creation or intraoperative hypotension are risk factors for WI in elective open colorectal resections [11, 17, 20], the other many parameters that have conventionally been re-

Table 2. Correlation between wound infection and perioperative/operative characteristics^a

	Wound infection		P value
	Positive (n = 18)	Negative (n = 272)	
Length of operation			.123
≤ 240 minutes	9	186	
> 240 minutes	9	86	
Antimicrobial prophylaxis administration			
Single dosing: multiple dosing (intra- and post-operative)	11:7	176:96	.802
Intraoperative repeated dosing (LOO > 180 minutes)	2/14 (14.3%)	22/207 (10.6%)	.653
First generation: second generation	6:12	92:180	1.000
Mean minimum body temperature (°C)	35.8 (34.9–36.7)	35.9 (34.5–37.4)	.854
Intraoperative hypotension (SBP < 80mmHg)	7	54	.071
Mean operative blood loss (mL)	64 (6–250)	61 (3–217)	.839
Drainage of anastomoses	17	268	.276
Mean duration of drainage (days)	4.5 (0–8)	4.6 (0–8)	.638
Stoma creation	4	14	.019
Anastomosis			.470
Double-stapling technique	8	123	
FETE + Handsewn anastomosis	10	142	
Handsewn per anum coloanal anastomosis	1	4	
Colon preparation			.380
One day before surgery	16	210	
Two days before surgery	2	62	
Median hospital stay (days)	8 (7–23)	8 (7–21)	.131

LOO, length of operation; SBP, systolic blood pressure; FETE, functional end-to-end anastomosis

^a Values in parentheses are ranges unless indicated otherwise

Table 3. Multivariable analysis of factors affecting wound infection

Independent Predictors	Odds ratio	Confidence intervals	P value
Stoma creation (yes/no)	5.275	1.196–23.263	.028
Intraoperative hypotension (SBP < 80mmHg/SBP ≥ 80mmHg)	3.171	1.099–9.147	.033

SBP, systolic blood pressure

garded as risk factors for WI were negative. Unfortunately, we were unable to evaluate matters such as the necessity of mechanical bowel preparation and the significance of preoperative oral prophylaxis, because all the patients underwent mechanical bowel preparation without oral antibiotics. In recent years, the significance of mechanical bowel preparation has been investigated in RCTs, and there are increasing reports that mechanical bowel preparation is not necessary; however many of those trials investigated cases of elective OS [3, 16]. As to whether the same result would be obtained in LS cases, further investigation is needed.

In this study, surgery was performed without oral antimicrobial prophylaxis administration in all patients. We do not administer oral antimicrobial prophylaxis in either OS or LS, for fear that clostridium difficile colitis or gastrointestinal symptoms including nausea, vomiting, abdominal pain, and diarrhea might be caused by oral antibiotics [5, 24]. In such cases, intravenous antimicrobial prophylaxis administration is required, but factors such as the type of antibiotic administered and its dosing frequency may influence the occurrence of WI. As the type of antibiotic used and its dosing frequency vary according to the study period, the influence of these

factors was evaluated in this study. We found no difference in the incidence of WI between single dosing and multiple dosing as well as between the presence and absence of intraoperative repeated dosing in cases with an operative time of four hours or more. With regard to the type of antibiotic administered, no difference was seen in the incidence of WI between first and second generation. Whether these items influence the occurrence of WI in LS cases is a question that requires further investigation.

In addition, the significance of drainage tubes should also be investigated. In our study, WI at the port site through which a drainage tube was inserted accounted for one-third of all WIs. Some reports say that drainage tubes are not necessary in elective colorectal surgery [21], but we leave drainage tubes in place for 4–5 days after operation to obtain information on postoperative bleeding and postoperative peritonitis associated with anastomotic leakage. However, because of the low frequency of these complications in LS, we now remove drainage tubes as early as the second or third postoperative day. Further investigation is also required with regard to the necessity of drainage tubes in LS.

The results of this study suggest the importance of maintaining normotension intraoperatively to reduce the incidence of WI, and we must recognize that this variable needs to be further investigated [17]. Moreover, if a stoma needs to be created, the port site should be positioned away from the stoma site as far as possible, and in patients with a created stoma, the wound may need to be completely covered with plastic film or other material. On the other hand, if WI occurs, hospital costs will be reduced by opening the wound early and managing it with self-dressing gauze changes.

In conclusion, as a result of our investigation into cases of LS, we found that stoma creation and intra-operative hypotension were independent risk factors for WI. Various factors are related to the occurrence of WI in LS cases so that efforts can be made to effectively prevent WI by measures such as increasing the dosing frequency of antimicrobial prophylaxis only in patients at higher risk for WI.

References

- Abraham NS, Young JM, Solomon MJ (2004) Meta-analysis of short-term outcomes after laparoscopic resection for colorectal cancer. *Br J Surg* 91: 1111-1124
- Braga M, Vignali A, Zuliani W, Frasson M, Di Serio C, Di Carlo V (2005) Laparoscopic versus open colorectal surgery: cost-benefit analysis in a single-center randomized trial. *Ann Surg* 242: 890-896
- Bucher P, Gervaz P, Soravia C, Mermillod B, Erne M, Morel P (2005) Randomized clinical trial of mechanical bowel preparation versus no preparation before elective left-sided colorectal surgery. *Br J Surg* 92: 409-414
- Clinical Outcomes of Surgical Therapy Study Group (2004) A comparison of laparoscopically assisted and open colectomy for colon cancer. *N Engl J Med* 350: 2050-2059
- Espin-Basany E, Sanchez-Garcia JL, Lopez-Cano M, Lozoya-Trujillo R, Medarde-Ferrer M, Armadans-Gil L, Alemany-Vilches L, Armengol-Carrasco M (2005) Prospective, randomized study on antibiotic prophylaxis in colorectal surgery. Is it really necessary to use oral antibiotics? *Int J Colorectal Dis* 20: 542-546
- Gaynes RP, Culver DH, Horan TC, Edwards JR, Richards C, Tolson JS (2001) National Nosocomial Infections Surveillance System. Surgical site infection (SSI) rates in the United States, 1992-1998: the National Nosocomial Infections Surveillance System basic SSI risk index. *Clin Infect Dis* 33: S69-77
- Guillou PJ, Quirke P, Thorpe H, Walker J, Jayne DG, Smith AM, Heath RM, Brown JM, MRC CLASICC trial group (2005) Short-term endpoints of conventional versus laparoscopic-assisted surgery in patients with colorectal cancer (MRC CLASICC trial): multicentre, randomised controlled trial. *Lancet* 365: 1718-1726
- Hasegawa H, Kabeshima Y, Watanabe M, Yamamoto S, Kitajima M (2003) Randomized controlled trial of laparoscopic versus open colectomy for advanced colorectal cancer. *Surg Endosc* 17: 636-640
- Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG (1992) CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Infect Control Hosp Epidemiol* 13: 606-608
- Kercher KW, Nguyen TH, Harold KL, Poplin ME, Matthews BD, Sing RF, Heniford BT (2004) Plastic wound protectors do not affect wound infection rates following laparoscopic-assisted colectomy. *Surg Endosc* 18: 148-151
- Konishi T, Watanabe T, Kishimoto J, Nagawa H (2006) Elective colon and rectal surgery differ in risk factors for wound infection: results of prospective surveillance. *Ann Surg* 244: 758-763
- Lacy AM, Garcia-Valdecasas JC, Delgado S, Castells A, Taura P, Piqué JM, Visa J (2002) Laparoscopy-assisted colectomy versus open colectomy for treatment of non-metastatic colon cancer: a randomised trial. *Lancet* 359: 2224-2229
- Leung KL, Kwok SPY, Lam SCW, Lee JFY, Yiu RYC, Ng SSM, Lai PBS, Lau WY (2004) Laparoscopic resection of rectosigmoid carcinoma: prospective randomised trial. *Lancet* 363: 1187-1192
- Mangram AJ, Horan TC, Pearson ML, silver LC, Jarvis WR (1999) The Hospital Infection Control Practices Advisory Committee. Guideline for prevention of surgical site infection, 1999. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 20: 250-278
- Owens WD, Felts JA, Spitznagel EL Jr (1978) ASA physical status classifications: a study of consistency of ratings. *Anesthesiology* 49: 239-243
- Slim K, Vicaut E, Panis Y, Chipponi J (2004) Meta-analysis of randomized clinical trials of colorectal surgery with or without mechanical bowel preparation. *Br J Surg* 91: 1125-1130
- Smith RL, Bohl JK, McElearney ST, Friel CM, Barclay MM, Sawyer RG, Foley EF (2004) Wound infection after elective colorectal resection. *Ann Surg* 239: 599-607
- Sørensen LT, Hemmingsen U, Kallehave F, Wille-Jørgensen P, Kjaergaard J, Møller LN, Jørgensen T (2005) Risk factors for tissue and wound complications in gastrointestinal surgery. *Ann Surg* 241: 654-658
- Tang CL, Eu KW, Tai BC, Soh JG, MacHin D, Seow-Choen F (2001) Randomized clinical trial of the effect of open versus laparoscopically assisted colectomy on systemic immunity in patients with colorectal cancer. *Br J Surg* 88: 801-807
- Tang R, Chen H, Wang Y, Changchien C, Chen JS, Hsu KC, Chiang JM, Wang JY (2001) Risk factors for surgical site infection after elective resection of the colon and rectum: a single-center prospective study of 2,809 consecutive patients. *Ann Surg* 234: 181-189
- Urbach DR, Kennedy ED, Cohen MM (1999) Colon and rectal anastomoses do not require routine drainage: a systematic review and meta-analysis. *Ann Surg* 229: 174-180
- Weiss CA 3rd, Statz CL, Dahms RA, Remucal MJ, Dunn DL, Beilman GJ (1999) Six years of surgical wound infection surveillance at a tertiary care center: review of the microbiologic and epidemiological aspects of 20,007 wounds. *Arch Surg* 134: 1041-1048
- Winslow ER, Fleshman JW, Birnbaum EH, Brunt LM (2002) Wound complications of laparoscopic vs open colectomy. *Surg Endosc* 16: 1420-1425
- Wren SM, Ahmed N, Jamal A, Safadi BY (2005) Preoperative oral antibiotics in colorectal surgery increase the rate of *Clostridium difficile* colitis. *Arch Surg* 140: 752-756
- Wu FP, Sietes C, von Blomberg BM, van Leeuwen PA, Meijer S, Cuesta MA (2003) Systemic and peritoneal inflammatory response after laparoscopic or conventional colon resection in cancer patients: a prospective, randomized trial. *Dis Colon Rectum* 46: 147-155
- Yamamoto S, Fujita S, Akasu T, Moriya Y (2004) A comparison of the complication rates between laparoscopic colectomy and laparoscopic low anterior resection. *Surg Endosc* 18: 1447-1451
- Yamamoto S, Fujita S, Akasu T, Moriya Y (2005) Safety of laparoscopic intracorporeal rectal transection with double-stapling technique anastomosis. *Surg Laparosc Endosc Percutan Tech* 15: 70-74

Cancer Invasion to Auerbach's Plexus is an Important Prognostic Factor in Patients with pT3-pT4 Colorectal Cancer

Shin Fujita, M.D.,¹ Yukihiro Nakanisi, M.D.,² Hirokazu Taniguchi, M.D.,³
Seiichiro Yamamoto, M.D.,¹ Takayuki Akasu, M.D.,¹ Yoshihiro Moriya, M.D.,¹
Tadakazu Shimoda, M.D.³

¹ Department of Surgery, National Cancer Center Hospital, Tokyo, Japan

² Department of Pathology, National Cancer Center Hospital, National Cancer Center Research Institute, Tokyo, Japan

³ Clinical Laboratory Division, National Cancer Center Hospital, Tokyo, Japan

PURPOSE: By defining perineural invasion of colorectal cancer as invasion to Auerbach's plexus, we examined the usefulness of this pathologic finding as a prognostic factor. **METHODS:** A total of 509 consecutive patients who underwent curative surgery for pT3 or pT4 colorectal cancer between May 1997 and December 2001 were reviewed. All the surviving patients were followed for more than five years. All the pathologic findings, including perineural invasion, were described prospectively in the pathology report forms. **RESULTS:** Perineural invasion was detected in 132 of 509 patients (26 percent) and was significantly associated with lymph node status, lymphatic invasion, and venous invasion. Incidences of local and systemic recurrence were significantly higher in patients with perineural invasion than in those without perineural invasion. The disease-free survival of the perineural invasion-positive group was significantly poorer than that of the perineural invasion-negative group for Stages II and III colon cancer, irrespective of the use of adjuvant chemotherapy. This improved disease-free survival also was seen in patients with Stage II rectal cancer not treated with adjuvant chemotherapy. There was a nonsignificant difference in disease-free survival for Stage II rectal cancer and Stage III rectal cancer treated with chemotherapy, that of

the perineural invasion-positive group being poorer. Multivariate analysis showed that lymph node status, perineural invasion, depth of invasion, and cancer site were significant prognostic factors. **CONCLUSIONS:** Perineural invasion defined as cancer invasion to Auerbach's plexus is an important prognostic factor for colorectal cancer. [Key words: Colorectal cancer; Perineural invasion; Auerbach's plexus; Prognostic factor]

Several reports have shown that perineural invasion (PNI) is an important prognostic factor in colorectal cancer¹⁻⁵ and rectal cancer.⁶⁻¹⁷ Therefore, the colorectal working group of the American Joint Committee on Cancer (AJCC) prognostic factors consensus conference has classified PNI as category IIA, which means that PNI has been extensively studied biologically and/or clinically and is considered to have sufficient predictive value for outcome to be noted in pathology reports.¹⁸ However, because many reports on PNI have been based on retrospective studies, and PNI has not been clearly defined, there is still no definitive conclusion about the degree to which PNI is a prognostic factor, especially in colon cancer. Therefore, in pathology reports compiled at the National Cancer Center Hospital from May 1997, we defined PNI as cancer invasion to Auerbach's plexus, because this feature is a prominent and easily detectable type of PNI, and

Supported by a Grant-in-Aid for Cancer Research from the Ministry of Health, Labor and Welfare of Japan.

Correspondence to: Shin Fujita, M.D., Department of Surgery, National Cancer Center Hospital, 1-1 Tsukiji 5-chome, Chuo-ku, Tokyo, 104-0045, Japan, e-mail: sfujita@ncc.go.jp

Dis Colon Rectum 2007; 50: 1860-1866

DOI: 10.1007/s10350-007-9072-8

© The American Society of Colon and Rectal Surgeons

Published online: 27 September 2007

PNI was reported prospectively. Although we had already reported that PNI was an important prognostic factor,¹⁹ follow-up time in the previous study was short and the number of patients examined was small. In the present study, all surviving patients were followed for more than five years and the number of examined patients was larger than in our previous study. Moreover, only pT3 or pT4 tumors were examined in the present study, because PNI was not found in pT1 tumors and was rare in pT2 tumors.

PATIENTS AND METHODS

Consecutive patients who underwent curative surgery for pT3 or pT4 colorectal cancer at the National Cancer Center Hospital between May 1997 and Dec 2001 were reviewed. Synchronous or metachronous multiple cancers were excluded from the analysis. One patient who died four days after surgery because of anastomotic leakage and sepsis also was excluded. A total of 509 patients were examined. The patients were followed up at three-month intervals for two years and at six-month intervals thereafter. Tumor markers were examined at every patient visit. CT scans of the liver and lung or abdominal ultrasonography with chest x-ray were performed at least every six months. Colonoscopy was performed twice within five years after surgery. All the surviving patients were followed for more than five years. Fifty-one of 266 patients with Stage III tumors received postoperative adjuvant chemotherapy as part of a clinical trial. Adjuvant radiotherapy was not used for rectal cancer during the study period.

Pathologic Examination

All the specimens were reviewed by two pathologists (TS and YN). Perineural invasion was defined as the presence of cancer cells inside the perineurium in Auerbach's plexus adjacent to the tumor front, and the results and other pathologic findings were described prospectively in the pathology report forms.

Statistical Analysis

Statistical analysis was performed by using the chi-squared test. Survival rates were calculated by the Kaplan-Meier method and survival curves were compared by using the log-rank test. Cox proportional hazards model was used for multivariate

analysis. Data differences between groups were considered statistically significant at $P < 0.05$.

RESULTS

PNI and Clinicopathologic Characteristics

A representative case of PNI is shown in Figure 1. Cancer cells invaded the perineurium in Auerbach's plexus. PNI was detected in 132 of 509 patients (26 percent). PNI and clinicopathologic characteristics of the patients are shown in Table 1. PNI was significantly associated with lymph node status, lymphatic invasion, and venous invasion ($P < 0.01$).

PNI in Relation to Recurrence and Survival

In colon cancer, the incidence of liver metastasis in the PNI-positive group was significantly higher than that in the PNI-negative group ($P < 0.01$; Table 2). In rectal cancer, the incidences of liver and lung metastasis and local recurrence in the PNI-positive group were significantly higher than in the PNI-negative group ($P \leq 0.01$). The five-year, disease-free survival rate in the PNI-positive group was 53 percent and that in the PNI-negative group was 80 percent (Fig. 2). Outcome was significantly poorer in the PNI-positive group than in the PNI-negative group ($P < 0.01$). Disease-free survival rates were examined according to tumor site (colon and rectum) and Stage (Stages II and III). Disease-free survival in the PNI-positive group was significantly poorer than that in the PNI-negative group for Stage II and III colon cancer ($P = 0.02, 0.03$, respectively) and Stage III rectal cancer ($P < 0.01$; Table 3, Fig. 3). Although disease-free survival in the PNI-positive group also was poorer than that of the PNI-negative group for Stage II rectal cancer, the difference was not statistically significant ($P = 0.21$). Because 51 of 266 patients with Stage III tumors received adjuvant chemotherapy, which is known to affect survival, the effect of adjuvant chemotherapy on disease-free survival was analyzed (Table 3). Patient survival in the PNI-positive group was poorer than that in the PNI-negative group, irrespective of whether adjuvant chemotherapy was given. Multivariate analysis of PNI, lymph node status, depth of invasion, tumor differentiation, lymphatic invasion, venous invasion, tumor site, preoperative CEA, gender, age, and adjuvant chemotherapy showed that lymph node status, PNI, depth of invasion, and tumor site were significant prognostic factors ($P < 0.01$; Table 4).




Figure 1. Representative PNI. a. Arrows shows cancer cells inside the perineurium in Auerbach's plexus. This is a case of massive PNI. b. Eighty percent of cases of PNI involve only slight invasion to Auerbach's plexus. In this case, one or two plexuses adjacent to the tumor front were invaded by cancer cells (arrow). Arrowhead shows Auerbach's plexus without cancer invasion. PNI = perineural invasion.

DISCUSSION

PNI has been reported to be a prognostic factor in colorectal cancer,¹⁻⁵ colon cancer,²⁰⁻²² and rectal cancer.⁶⁻¹⁷ However, there is still no definitive conclusion about the degree to which PNI is a prognostic factor, especially in colon cancer, because many of the previous studies of PNI were retrospective, and PNI was not clearly defined. Although many of the reports did not define PNI, PNI was considered

to be perineural cancer invasion within and outside the bowel wall in some of them,^{1,6,9,12} and only extramural PNI was examined in other studies.^{7,10,14} We defined PNI as cancer invasion to Auerbach's plexus, and on this basis prospectively examined more than 500 patients. Our findings clearly demonstrated that PNI was a significant prognostic factor in pT3 or pT4 colorectal cancer. Therefore, this study provides strong evidence that cancer invasion to Auerbach's plexus is a prognostic factor for colorectal cancer.