

Suzuki, R., Yoshikawa, T., Ihira, M., Enomoto, Y., Inagaki, S., Matsumoto, K., Kato, K., Kudo, K., Kojima, S., Asano, Y., 2006. Development of the loop-mediated isothermal amplification method for rapid detection of cytomegalovirus DNA. *J. Virol. Meth.* 132, 216–221.

Tedeschi, R., Enbom, M., Bidoli, E., Linde, A., De Paoli, P., Dillner, J., 2001. Viral load of human herpesvirus 8 in peripheral blood of human immunod-

efficiency virus-infected patients with Kaposi's sarcoma. *J. Clin. Microbiol.* 39, 4269–4273.

Yoshikawa, T., Ihira, M., Akimoto, S., Usui, C., Miyake, F., Suga, S., Enomoto, Y., Suzuki, R., Nishiyama, Y., Asano, Y., 2004. Rapid diagnosis of human herpesvirus 7 infection by a novel DNA amplification method, Loop-mediated isothermal amplification. *J. Clin. Microbiol.* 42, 1348–1352.

研究成果の刊行に関する一覧表

平成 19 年度 慶應義塾大学医学部 熱帯医学・寄生虫学教室 竹内 勤

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Asao Makioka, Masahiro Kumagai, Seiki Kobayashi, Tsutomu Takeuchi	Differences in protein profiles of the isolates of Entamoeba histolytica and E. dispar by surface-enhanced laser desorption ionization time-of- flight mass spectrometry (SELDI-TOF MS) ProteinChip assays	Parasitol Res	102	103-110	2007
H. TACHIBANA, X.-J. CHENG, S. KOBAYASHI, Y. OKADA, J. ITOH and T. TAKEUCHI	Primary structure, expression and localization of two intermediate subunit lectins of Entamoeba dispar that contain multiple CXXC motifs	Parasitology	134	1989-1999	2007
Sadatomo Tasaka, MD, FCCP; Naoki Hasegawa, MD; Seiki Kobayashi, MD; Wakako Yamada, MD; Tomoyasu Nishimura, MD; Tsutomu Takeuchi, MD; and Akitoshi Ishizaka, MD	Serum Indicators for the Diagnosis of Pneumocystis Pneumonia	CHEST	131	1173-1180	2007
Jun Suzuki, Seiki Kobayashi, Ph.D., Rie Murata, Yoshitoki Yanagawa, D.V.M., and Tsutomu Takeuchi, M.D., Ph.D.	PROFILES OF A PATHOGENIC ENTAMOEBIA HISTOLYTICA-LIKE VARIANT WITH VARIATIONS IN THE NUCLEOTIDE SEQUENCE OF THE SMALL SUBUNIT RIBOSOMAL RNA ISOLATED FROM A PRIMATE (DE BRAZZA'S GUENON)	Journal of Zoo and Wildlife Medicine	38	471-474	2007
Taro Fukao, Yoko Fukuda, Kotaro Kiga, Jafar Sharif, Kimihiro Hino, Yutaka Enomoto, Aya Kawamura, Kaito Nakamura, Tsutomu Takeuchi, and Masanobu Tanabe	An Evolutionarily Conserved Mechanism for MicroRNA-223 Expression Revealed by MicroRNA Gene Profiling	Cell	129	617-631	2007

# Differences in protein profiles of the isolates of *Entamoeba histolytica* and *E. dispar* by surface-enhanced laser desorption ionization time-of-flight mass spectrometry (SELDI-TOF MS) ProteinChip assays

Asao Makioka · Masahiro Kumagai ·  
Seiki Kobayashi · Tsutomu Takeuchi

Received: 4 August 2007 / Accepted: 17 August 2007 / Published online: 11 September 2007  
© Springer-Verlag 2007

**Abstract** Surface-enhanced laser desorption ionization time of flight mass spectrometry (SELDI-TOF MS) ProteinChip assays with weak cationic exchange chips were used for protein profiling of different isolates of *Entamoeba histolytica* and *E. dispar*. When SELDI-TOF MS spectra of cell lysates from *E. histolytica* strain HM-1:IMSS were compared with those from four other laboratory strains (200:NIH, HK-9, DKB, and SAW755CR) grown under the same culture conditions, different peak patterns of SELDI-TOF MS were observed among these strains, independent of their zymodeme types. Similarly, five Japanese isolates of *E. histolytica* grown under the same culture conditions revealed different peak patterns among themselves. The SELDI-TOF MS spectra of cell lysates from two isolates of *E. dispar* strain AS16IR and CYNO 09:TPC showed the presence of peaks specific for *E. dispar* isolates and the absence of peaks common to *E. histolytica* isolates. This is not only the first use of SELDI-TOF MS ProteinChip technology for protein profiling of different strains of *Entamoeba* but also the use for parasitic protozoa. The SELDI-TOF MS spectra

show a realistic view of proteins with a biological status of *E. histolytica* and *E. dispar* isolates, contributing to show their phenotypic differences of proteins and provide a unique means of distinguishing them.

## Introduction

Surface-enhanced laser desorption ionization time-of-flight mass spectrometry (SELDI-TOF MS) ProteinChip assay is a relatively recent technology for exploring proteomes combining chromatography and mass spectrometry (Hutchens and Yip 1993; Issaq et al. 2002). The ProteinChip arrays contain sample spots of 1 or 2 mm diameter, with each chip having a different surface chemistry. These may be chemical (e.g., ionic, hydrophobic, hydrophilic) or biochemical (antibody, receptor, deoxyribonucleic acid [DNA], etc.) and are designed to capture proteins of interest and then analyzed directly on the SELDI-TOF mass spectrometer. The ProteinChip technology has been used for protein profiling and biomarker discovery for diseases such as cancer, neurological disorders, and pathogenic organisms including human African trypanosomiasis (Issaq et al. 2002; Papadopoulos et al. 2004).

*Entamoeba histolytica*, a protozoan parasite, is responsible for an estimated 40–50 million cases of amebic colitis and liver abscess (WHO/PAHO/UNESCO 1997). There are a large number of isolates of *E. histolytica* that differ in their phenotypes. Although genetic diversity of isolates of *E. histolytica* has been extensively studied, there are few reports on differences in their protein profiles as the phenotype. In this study, we used the SELDI-TOF MS ProteinChip

A. Makioka (✉) · M. Kumagai  
Department of Tropical Medicine,  
Jikei University School of Medicine,  
3-25-8 Nishi-shinbashi, Minato-ku,  
Tokyo 105-8461, Japan  
e-mail: makioka@jikei.ac.jp

S. Kobayashi · T. Takeuchi  
Department of Tropical Medicine and Parasitology,  
Keio University School of Medicine,  
35 Shinanomachi, Shinjuku-ku,  
Tokyo 160-8582, Japan

assays to examine protein profiles of different isolates of *E. histolytica*. We also compared protein profiles of *E. histolytica* with those of nonpathogenic *E. dispar*.

## Materials and methods

Five laboratory strains and five Japanese isolates of *E. histolytica* and two isolates of *E. dispar* were used in this study (Table 1). The Japanese isolates of *E. histolytica* were obtained from clinical specimens collected from amebiasis patients in Japan. Axenic in vitro cultures were established and maintained in Diamond's BI-S-33 medium as previously described (Diamond et al. 1978). The two *E. dispar* isolates included one human isolate and one nonhuman primate isolate. The *E. dispar* trophozoites were cultivated axenically in newly designed medium (Kobayashi et al. 2005) with a modification of replacement of gluconic acid with maltose (Kobayashi et al. manuscript in preparation). All cases with intestinal amebiasis or liver abscess were diagnosed by microscopic demonstration of trophozoites or cysts in stool or of trophozoites in liver aspirates, respectively. Zymodeme analysis (Sargeant 1988) and polymerase chain reaction analysis (Tachibana et al. 1991; Cheng et al. 1993) of the amoeba isolates were performed to characterize them.

Trophozoites of *E. histolytica* and *E. dispar* in log-phase culture were harvested by centrifugation at 400×g for 5 min and washed by centrifugation three times in phosphate-buffered saline and stored in -80°C until use. The pellets

of amoeba were dissolved in 2% CHAPS (Sigma-Aldrich, St Louis, MO), 5 mM Tris-HCl buffer, pH 8.0 containing proteinase cocktail (Sigma-Aldrich) with 140 μM E-64 by vortexing. After centrifugation at 15,000×g for 15 min, the resulting supernatants were used as whole-cell lysates. Protein concentration was determined by Bio-Rad DC protein assay kit (Bio-Rad Laboratories, Hercules, CA) according to the manufacturer's instructions.

One microgram of cell extract diluted into 4 μl of starting buffer was spotted onto each ProteinChip array (originally Ciphergen Biosystems, Fremont, CA; now Bio-Rad Laboratories) with CM10 (weak cation exchanger: Carboxylate). The chips were then incubated at room temperature for 20 min on a shaker. Nonbound proteins and other contaminants were washed from the CM10 ProteinChip arrays with a buffer of 0.1 M ammonium acetate, pH 4.0, three times. Finally, all chips were washed with MilliQ water to remove interfering salts and detergents. After drying, 0.5 μl of saturated energy-absorbing molecule solution (sinapinic acid in acetonitrile [v/v] and trifluoroacetic acid [v/v]) was added twice, and the chips were allowed to air dry. Mass spectrometry analysis was performed by time-of-flight mass spectrometry in a PBS-II mass reader (Ciphergen Biosystems). Spectra were collected using an average 80 nitrogen laser shots. Spectrum analysis was performed using the ProteinChip software version 2.1b (Ciphergen Biosystems). The optimal detection size range was set between 2,000 and 20,000 Da because the system is most effective at profiling low-molecular-weight proteins (i.e., <20 kDa; Issaq et al. 2002). The intensity of each

**Table 1** *Entamoeba histolytica* and *E. dispar* isolates used in this study

Isolate	Isolation		Clinical diagnosis	Serology <sup>a</sup> (S)/PCR <sup>b</sup> (P)	Zymodeme <sup>c</sup>
	Location	Year			
<i>E. histolytica</i>					
HM-1:JMSS	Mexico	1967	Dysentery	NA <sup>d</sup>	II
200:NIH	USA	1949	Dysentery	NA	II
HK-9	Korea	1951	Dysentery	NA	II
DKB	UK	1924	Dysentery	NA	II
SAW755CR	UK	1979	Hematophagous trophozoites in feces	NA	XIV
KU43	Japan	2002	Colitis	S +	II
KU46	Japan	2004	Colitis	S +	XXI
KU2	Japan	1988	Colitis	S +	XIX
KU38	Japan	2002	Asymptomatic	S +	II
KU14	Japan	1999	Asymptomatic	S +	XIX
<i>E. dispar</i>					
AS161R	Iran	1997	Abdominal pain	P +	I
CYNO 09:TPC	Philippines	1994	NA	P +	I

<sup>a</sup> Serology was done by enzyme-linked immunosorbent assay, gel diffusion test, and/or indirect fluorescent-antibody test.

<sup>b</sup> PCR analysis using two sets of oligonucleotide primers each (p11 plus p12 and p13 plus p14, respectively) for amplification of the DNAs of *E. histolytica* and *E. dispar* was performed.

<sup>c</sup> Zymodemes type I and III are classified as *E. dispar*.

<sup>d</sup> NA Not available

of the peaks to be quantified was determined according to externally calibrated standards (Ciphergen Biosystems). According to the manufacturer, the mass accuracy of the spectrometer is 0.1%. The raw intensity data were normalized using the total ion current of  $m/z$  between 2,000 and 20,000 for all study sample profiles.

The mean  $\pm$  SE of the intensity of each peak in SELDI-TOF MS spectra from three independent cultures was calculated for all the isolates. The percentages of the number of peaks whose intensities were comparable to between the two isolates were shown as percentage similarity of the peak pattern. From these values, a dendrogram was generated by the unweighted pair-group method with arithmetic mean (UPGMA).

## Results

### Reproducibility of SELDI-TOF MS spectra tested with *E. histolytica* HM-1:IMSS

SELDI-TOF MS spectra of the whole cell lysates of *E. histolytica* HM-1:IMSS as a standard strain from three independent cultures are shown in Fig. 1. The peak patterns were almost reproducible. The two groups of peaks in which 4.305 and 8.274 kDa as major peaks were detected in a mass range of 3–15 kDa.

### Comparison of SELDI-TOF MS spectra between *E. histolytica* HM-1:IMSS and other laboratory strains

SELDI-TOF MS spectra of *E. histolytica* strain HM-1:IMSS were compared with those from four other laboratory strains, 200:NIH, HK-9, DKB, and SAW755CR, grown under the same culture conditions. As shown in Fig. 2, peaks of HM-1:IMSS in the mass range of 3–6 kDa were significantly higher than those of the other four strains. The peaks of 6–8 kDa were low in all the strains. In the mass range of 8–10 kDa, a peak of 8.274 kDa of HM-1:IMSS was significantly higher

than those of the other four strains, whereas peaks of 8.568, 8.791, and 9.267 kDa of HM-1:IMSS were lower than those of the other four strains. In addition, the peak of 9.267 kDa revealed different intensities among the other four strains. There were low peaks in the range of 10–15 kDa in all the strains.


### Comparison of SELDI-TOF MS spectra between *E. histolytica* HM-1:IMSS and Japanese isolates

SELDI-TOF MS spectra of *E. histolytica* strain HM-1:IMSS were compared with those from five Japanese isolates, KU43, KU46, KU2, KU38, and KU14. As shown in Fig. 3, only KU2 showed lower peaks than HM-1:IMSS in the range of 3–6 kDa. The peak of 5.423 kDa, which was little seen in the laboratory strains except HM-1:IMSS, was detected in the Japanese isolates except KU2. The intensity of the major peak of 4.305 kDa was comparable to or higher than that of HM-1:IMSS in the Japanese isolates except KU2. The Japanese isolates also showed low peaks in the range of 6–8 kDa, although several peaks of KU2 were higher than those of the other four Japanese isolates. In the range of 8–10 kDa, the peak patterns of KU43, KU38, and KU14 were relatively similar to that of HM-1:IMSS, whereas a high peak of 9.267 kDa was observed only in KU2. The peak of 8.568 kDa in KU46 and KU2 was significantly higher than those of HM-1:IMSS and other Japanese isolates. No higher peaks were detected in the range of 10–15 kDa of all the Japanese isolates like the laboratory strains of *E. histolytica*.


### Comparison of SELDI-TOF MS spectra between *E. histolytica* HM-1:IMSS and *E. dispar* isolates

SELDI-TOF MS spectra of *E. histolytica* HM-1:IMSS grown in the medium for *E. dispar* were similar to those grown in BI-S-33 medium (Fig. 1; data not shown). When SELDI-TOF MS spectra of HM-1:IMSS were compared with those from *E. dispar* isolates, AS161R and CYNO 09:TPC, the


**Fig. 1** SELDI-TOF MS spectra of *E. histolytica* strain HM-1:IMSS from three independent cultures. The molecular masses are shown above the peaks




**Fig. 2** Representative SELDI-TOF MS spectra of five laboratory strains of *E. histolytica*. The molecular masses are shown above the peaks



**Fig. 3** Representative SELDI-TOF MS spectra of five Japanese isolates of *E. histolytica*. The molecular masses are shown above the major peaks



**Fig. 4** Representative SELDI-TOF MS spectra of *E. dispar* isolates. The molecular masses are shown above the major peaks



peaks of 4.305 and 5.423 kDa were significantly lower than those of HM-1:IMSS, which was similar to the *E. histolytica* laboratory strains except HM-1:IMSS. The peak of 5.887 kDa was detected only in CYNO 09:TPC. The peaks of AS16IR and CYNO 09:TPC were mostly low in the range of 6–8 kDa, and the peak of 7.571 kDa was observed only in CYNO 09:TPC. In the range of 8–10 kDa, the major peak of 8.274 kDa common to *E. histolytica* isolates was not seen in either isolate of *E. dispar*, but the peak of 8.246 kDa was observed. Furthermore, like the *E. histolytica* laboratory strains except HM-1:IMSS, the peak of 8.568 kDa was significantly higher than that of HM-1:IMSS. The peak of 9.267 kDa, which was not seen in HM-1:IMSS, was observed in both isolates. In the range of 10–15 kDa, two peaks of 11.531 and

11.734 kDa not detected in *E. histolytica* were detected in both isolates of *E. dispar* (Fig. 4).


Similarity of peak patterns by SELDI-TOF MS among *E. histolytica* and *E. dispar* isolates and a dendrogram by the UPGMA method

The HM-1:IMSS showed 38.4 (200:NIH) to 69.2% (KU38 and KU14) similarity against other *E. histolytica* isolates, and KU38 and KU14 showed the highest similarity of 96.9% (Fig. 5). *E. dispar* AS16IR and CYNO 09:TPC showed 81.5% similarity to each other and were most associated with *E. histolytica* DKB, showing 72.3% similarity. The dendrogram representing relations among *E. histolytica* and

**Fig. 5** Similarity of protein profiles among *E. histolytica* and *E. dispar* isolates. Percentage similarities were determined as described in Materials and methods

	HM-1:IMSS	200:NIH	HK-9	DKB	SAW755CR	KU43	KU46	KU2	KU38	KU14	AS16IR	CYNO09:TPC
HM-1:IMSS												
200:NIH	38.4											
HK-9	50.8	81.5										
DKB	53.8	76.9	81.5									
SAW755CR	44.6	70.8	83.1	87.7								
KU43	56.9	50.8	64.6	63.1	58.5							
KU46	49.2	56.9	70.8	70.8	63.1	67.7						
KU2	41.5	83.1	75.4	78.5	70.8	44.6	61.5					
KU38	69.2	55.4	72.3	70.8	67.7	84.6	83.1	60.0				
KU14	69.2	58.5	73.8	80.0	72.3	83.1	83.1	63.1	96.9			
AS16IR	38.5	61.5	61.5	72.3	61.5	47.7	44.6	55.4	58.5	56.9		
CYNO09:TPC	29.2	53.8	58.5	72.3	63.1	44.6	46.2	43.1	47.7	55.4	81.5	

**Fig. 6** Dendrogram representing relations among *E. histolytica* and *E. dispar* isolates by the UPGMA method



*E. dispar* isolates indicates that HM-1:IMSS is in an independent clade separated from other *E. histolytica* isolates, and two isolates of *E. dispar* were in one clade (Fig. 6). In two other clades, one includes KU46, KU43, KU38, and KU14, and the other includes HK-9, DKB, SAW755CR, 200:NIH, and KU2. DKB and SAW755CR, and 200:NIH and KU2, as well as KU38 and KU14 were included in the same clades, respectively (Fig. 6).

## Discussion

ProteinChip technology is a new technology used to study proteomic profiles in biological samples, such as serum, cerebrospinal fluid, and cell or tissue extracts (Merchant and Weinberger 2000; Wulfskuhle et al. 2001; Carrette et al. 2003; Luo et al. 2003; Carlson et al. 2004) and is used in biomarker discovery and protein profiling (Ball et al. 2002; Choe et al. 2002; Yasui et al. 2003; Hayman and Przyborski 2004). With regard to parasitic protozoa, this technology was first used to detect antigens in sera of patients with African trypanosomiasis as a novel diagnostic method (Papadopoulos et al. 2004).

The heterogeneity of several DNA loci, including protein-coding sequences, has been extensively characterized among *E. histolytica* and *E. dispar* isolates (Clark and Diamond 1993; Ghosh et al. 2000; Zaki and Clark 2001; Zaki et al. 2002; Haghghi et al. 2002; Haghghi et al. 2003), and the overall genomic diversity among them has also been reported (Shah et al. 2005). However, there have been few reports on protein profiling among isolates of the two

*Entamoeba* species. In this study, we used SELDI-TOF MS ProteinChip technology to identify protein patterns in different isolates of *E. histolytica* and showed phenotypic heterogeneity of proteins among them even under identical culture conditions. The results clearly indicate differences in SELDI-TOF MS spectra between the HM-1:IMSS and the other four laboratory strains and also between the HM-1:IMSS and the Japanese isolates, independent of their zymodemes. The HM-1:IMSS, which is most widely used as a standard strain, was in a separate clade in the dendrogram from the other *E. histolytica* isolates. Although the reason for this remains unclear, it should be taken into consideration when *E. histolytica* HM-1:IMSS as a standard strain is compared with other *Entamoeba* species. The percentage similarity in peak patterns among the isolates did not distinguish *E. dispar* from *E. histolytica*. It is clear from comparison of protein profiles between *E. histolytica* and *E. dispar* that there are peaks specific for each species of amoeba. These could be useful markers for distinguishing the two species, although the number of *E. dispar* isolates used in this study is limited.

Two-dimensional polyacrylamide gel electrophoresis (2D-PAGE) has been the most commonly used method for proteomic analysis (Görg et al. 2004), but the need for protein staining and the subsequent sample handling limits the sensitivity of the overall approach (Issaq et al. 2002). The 2D-PAGE is also laborious, difficult to automate, has poor resolution for extreme masses and extreme acidic or basic proteins, and requires large amount of starting material. The 2D-PAGE has recently been used for proteome of *E. histolytica* HM-1:IMSS trophozoites (Leitsch et al. 2005).



If 2D-PAGE is used for analysis of a number of different isolates of *E. histolytica* and *E. dispar*, it would be very laborious, time consuming, and difficult to analyze. In this respect, SELDI-TOF MS ProteinChip technique has an advantage over 2D-PAGE.

There is a wide variety of clinical manifestations observed among individuals infected with *E. histolytica* and/or *E. dispar*. What determines these differences remains unclear. In this respect, studies on phenotypic differences in proteins other than genetic heterogeneity are important and would contribute to resolving the question because the proteome, compared with the genome, provides a more realistic view of a biological status and is, therefore, expected to be more useful than gene analysis for evaluating disease presence and progression and response to treatment (Engwegen et al. 2006). Thus, proteomics can bridge the gap between the genome sequence and cellular behavior.

Finally, SELDI-TOF MS ProteinChip technique was successfully used to analyze different isolates of *E. histolytica* and *E. dispar*. Using the different array surfaces, a complete picture of each strain of *Entamoeba* may be drawn, and thus a better set of protein fingerprint profiles for each strain would be provided. Like the successful application of this technology to bacterial proteomes (Barzaghi et al. 2004), the results show the usefulness of ProteinChip technology for studying the proteomics of parasitic protozoa as well, alone or in combination with other technologies.

**Acknowledgments** We thank N Watanabe for his valuable discussion with us, T Obata for SELDI-TOF MS ProteinChip analysis, and T Yamashita for her technical assistance. This work was supported by a Health Science Research Grant for Research on Emerging and Re-emerging Infectious Diseases from the Ministry of Health, Labor and Welfare of Japan.

## References

- Ball G, Mian S, Holding F, Allibone RO, Lowe J, Ali S, Li G, McCardle S, Ellis IO, Creaser C, Rees RC (2002) An integrated approach utilizing artificial neural networks and SELDI mass spectrometry for the classification of human tumours and rapid identification of potential biomarkers. *Bioinformatics* 18:395–404
- Barzaghi D, Isbister JD, Lauer KP, Born TL (2004) Use of surface-enhanced laser desorption/ionization-time of flight to explore bacterial proteomes. *Proteomics* 4:2624–2628
- Carlson KA, Ciborowski P, Schellpeper CN, Biskup TM, Shen RF, Luo X, Destache CJ, Gendelman HE (2004) Proteomic fingerprinting of HIV-1-infected human monocyte-derived macrophages: a preliminary report. *J Neuroimmunol* 147:35–42
- Carrette O, Demalte J, Scherl A, Yalkinoglu O, Corthals G, Burkhard P, Hochstrasser DF, Sanchez JC (2003) A panel of cerebrospinal fluid potential biomarkers for the diagnosis of Alzheimer's disease. *Proteomics* 3:1486–1494
- Cheng X-J, Tachibana H, Kobayashi S, Kaneda Y, Huang M-Y (1993) Pathogenicity of *Entamoeba histolytica* isolates from Shanghai, China. *Parasitol Res* 79:608–610
- Choe LH, Dutt MJ, Relkin N, Lee KH (2002) Studies of potential cerebrospinal fluid molecular markers for Alzheimer's disease. *Electrophoresis* 23:2247–2251
- Clark CG, Diamond LS (1993) *Entamoeba histolytica*: a method for isolate identification. *Exp Parasitol* 77:450–455
- Diamond LS, Hartow DR, Cunnick CC (1978) A new medium for the axenic cultivation of *Entamoeba histolytica* and other *Entamoeba*. *Trans R Soc Trop Med Hyg* 72:431–432
- Engwegen JYMN, Gast MC, Schellens JHM, Beijnen JH (2006) Clinical proteomics: searching for better tumour markers with SELDI-TOF mass spectrometry. *Trends Pharmacol Sci* 27:251–259
- Ghosh S, Frisardi M, Ramirez-Avila L, Descoteaux S, Sturm-Ramirez K, Newton-Sanchez OA, Santos-Preciado JJ, Ganguly C, Lohia A, Reed S, Samuelson J (2000) Molecular epidemiology of *Entamoeba* spp.: evidence of a bottleneck (demographic sweep) and trans-continental spread of diploid parasites. *J Clin Microbiol* 38:3815–3821
- Görg A, Weiss W, Dunn MJ (2004) Current two-dimensional electrophoresis technology for proteomics. *Proteomics* 4:3665–3685
- Haghighi A, Kobayashi S, Takeuchi T, Masuda G, Nozaki T (2002) Remarkable genetic polymorphism among *Entamoeba histolytica* isolates from a limited geographic area. *J Clin Microbiol* 40:4081–4090
- Haghighi A, Kobayashi S, Takeuchi T, Thammapalerd N, Nozaki T (2003) Geographic diversity among genotypes of *Entamoeba histolytica* field isolates from a limited geographic area. *J Clin Microbiol* 41:3748–3756
- Hayman MW, Przyborski SA (2004) Proteomic identification of biomarkers expressed by human pluripotent stem cells. *Biochem Biophys Res Commun* 316:918–923
- Hutchens TW, Yip TT (1993) New desorption strategies for the mass spectrometric analysis of macromolecules. *Rapid Commun Mass Spectrom* 7:576–580
- Issaq HJ, Veenstra TD, Cornard TP, Felschow D (2002) The SELDI-TOF MS approach to proteomics: protein profiling and biomarker identification. *Biochem Biophys Res Commun* 292:587–592
- Kobayashi S, Imai E, Haghighi A, Khalifa SA, Tachibana H, Takeuchi T (2005) Axenic cultivation of *Entamoeba dispar* in newly designed yeast extract-iron-gluconic acid-dihydroxyacetone-serum medium. *J Parasitol* 91:1–4
- Leitsch D, Radauer C, Paschinger K, Wilson IB, Breiteneder H, Scheiner O, Duchene M (2005) *Entamoeba histolytica*: analysis of the trophozoite proteome by two-dimensional polyacrylamide gel electrophoresis. *Exp Parasitol* 110:191–195
- Luo X, Carlson KA, Wojna V, Mayo R, Biskup TM, Stoner J, Anderson J, Gendelman HE, Melendez LM (2003) Macrophage proteomic fingerprinting predicts HIV-1-associated cognitive impairment. *Neurology* 60:1931–1937
- Merchant M, Weinberger SR (2000) Recent advancements in surface enhanced laser desorption/ionization-time of flight-mass spectrometry. *Electrophoresis* 21:1164–1177
- Papadopoulos MC, Abel PM, Stich A, Taralli E, Bell BA, Planche T, Loosemore A, Saadoun S, Wilkins P, Krishna S (2004) A novel and accurate diagnostic test for human African trypanosomiasis. *Lancet* 363:1358–1363
- Sargeant PG (1988) Zymodemes of *Entamoeba histolytica*. In: Radvin JJ (ed) *Amebiasis: human infection by Entamoeba histolytica*. Wiley, New York, pp 370–387
- Shah PH, MacFarlane RC, Bhattacharya D, Matese JC, Demeter J, Stroup SE, Singh U (2005) Comparative genomic hybridizations of *Entamoeba* strains reveal unique genetic fingerprints that correlate with virulence. *Eukaryotic Cell* 4:504–515
- Tachibana H, Kobayashi S, Takekoshi M, Ihara S (1991) Distinguishing pathogenic isolates of *Entamoeba histolytica* by polymerase chain reaction. *J Infect Dis* 164:825–826

- WHO/PAHO/UNESCO (1997) A consultation with experts on amebiasis. *Epidemiol Bull* 18:13–14
- Wulfkuhle JD, McLean KC, Paweletz CP, Sgroi DC, Trock BJ, Steeg PS, Petricoin EF III (2001) New approaches to proteomic analysis of breast cancer. *Proteomics* 1:1205–1215
- Yasui Y, Pepe M, Thompson ML, Adam BL, Wright GL Jr, Qu Y, Potter JD, Winget M, Thornquist M, Feng Z (2003) A data-analytic strategy for protein biomarker discovery: profiling of high-dimensional proteomic data for cancer detection. *Biostatistics* 4:449–463
- Zaki M, Clark CG (2001) Isolation and characterization of polymorphic DNA from *Entamoeba histolytica*. *J Clin Microbiol* 39:897–905
- Zaki M, Meelu P, Sun W, Clark CG (2002) Simultaneous differentiation and typing of *Entamoeba histolytica* and *Entamoeba dispar*. *J Clin Microbiol* 40:1271–1276

# Primary structure, expression and localization of two intermediate subunit lectins of *Entamoeba dispar* that contain multiple CXXC motifs

H. TACHIBANA<sup>1\*</sup>, X.-J. CHENG<sup>1</sup>, S. KOBAYASHI<sup>2</sup>, Y. OKADA<sup>3</sup>, J. ITOH<sup>3</sup>  
and T. TAKEUCHI<sup>2</sup>

<sup>1</sup>Department of Infectious Diseases and <sup>3</sup>Teaching and Research Support Center, Tokai University School of Medicine, Isehara, Kanagawa 259-1193, Japan

<sup>2</sup>Department of Tropical Medicine and Parasitology, Keio University School of Medicine, Shinjuku-ku, Tokyo 160-8582, Japan

(Received 29 May 2007; revised 25 June 2007; accepted 4 July 2007; first published online 6 September 2007)

## SUMMARY

We have recently identified 2 surface proteins in *Entamoeba histolytica* as intermediate subunits of galactose- and *N*-acetyl-D-galactosamine-inhibitable lectin (EhIgl1 and EhIgl2); these proteins both contain multiple CXXC motifs. Here, we report the molecular characterization of the corresponding proteins in *Entamoeba dispar*, which is neither pathogenic nor invasive. Two *Igl* genes encoding 1110 and 1106 amino acids (EdIgl1 and EdIgl2) were cloned from 2 strains of *E. dispar*. The amino acid sequence identities were 79% between EdIgl1 and EdIgl2, 75–76% between EdIgl1 and EhIgl1, and 73–74% between EdIgl2 and EhIgl2. However, all the CXXC motifs were conserved in the EdIgl proteins, suggesting that the fold conferred by this motif is important for function. Comparison of the expression level of the *Igl* genes by real-time RT-PCR showed 3–5 times higher expression of *EdIgl1* compared to *EdIgl2*. Most EdIgl1 and EdIgl2 proteins were co-localized on the surface and in the cytoplasm of trophozoites, based on confocal microscopy. However, a different localization of EdIgl1 and EdIgl2 in intracellular vacuoles and a different level of phenotypic expression of the two Igl were also observed. These results demonstrate that Igl are important proteins even in non-pathogenic amoeba and that Igl1 and Igl2 may possess different functions.

Key words: *Entamoeba dispar*, *Entamoeba histolytica*, intermediate subunit of Gal/GalNAc lectin (Igl), cysteine-rich protein.

## INTRODUCTION

It has been estimated that 480 million people worldwide are infected with *Entamoeba histolytica* or *Entamoeba dispar* (Walsh, 1986). *E. histolytica* is the causative agent of human amoebic colitis and liver abscess, which result in up to 110 000 deaths annually. *E. dispar* is morphologically indistinguishable from *E. histolytica*, but is non-pathogenic and non-virulent (Diamond and Clark, 1993). Adherence of *E. histolytica* trophozoites to host cells is an essential step in its pathogenicity, and it is well known that the 170 kDa heavy subunit of galactose- and *N*-acetyl-D-galactosamine (Gal/GalNAc)-inhibitable lectin (Hgl) is the key factor in adherence and subsequent pathogenesis of the amoeba (Petri *et al.* 2002). Hgl is a transmembrane protein that forms a heterodimer with a glycosylphosphatidylinositol (GPI)-anchored 35 kDa light subunit (Lgl) via

disulfide bonds (Petri *et al.* 1989). Recently, we have identified a GPI-anchored 150 kDa intermediate subunit (Igl) of lectin, which is non-covalently associated with Hgl (Cheng *et al.* 1998, 2001). There are 2 isoforms of Igl, which consist of 1101 and 1105 amino acids and are referred to as Igl1 and Igl2, respectively; both are cysteine-rich proteins containing multiple CXXC motifs. A mouse monoclonal antibody (mAb) to Igl significantly inhibits adherence and cytotoxicity of trophozoites to mammalian cells *in vitro* and also inhibits liver abscess formation in hamsters (Tachibana *et al.* 1997; Cheng *et al.* 1997, 1999). Antibodies to Igl have been detected not only in symptomatic patients with amoebiasis but also in asymptomatic cyst passers of *E. histolytica* (Tachibana *et al.* 2004). Immunization of hamsters with affinity-purified Igl can prevent amoebic liver abscess formation (Cheng and Tachibana, 2001), and Igl has also been detected in the *E. histolytica* fraction that interacts with the brush border of enterocytes (Seigneur *et al.* 2005). Therefore, Igl seems to be one of the key molecules in amoebic adherence to host cells and pathogenicity; however, the correlation of these effects with

\* Corresponding author: Department of Infectious Diseases, Tokai University School of Medicine, Isehara, Kanagawa 259-1193, Japan. Tel: +81 463 93 1121. Fax: +81 463 95 5450. E-mail: htachiba@is.icc.u-tokai.ac.jp

each Igl isoform and the differences between the isoforms are not known.

Comparison of *E. histolytica* and *E. dispar* is also of importance for understanding the pathogenicity of amoeba. One well-known difference between the two species is associated with the family of cysteine proteases that are pathogenic factors in *E. histolytica*; in *E. dispar*, the *EhCP1* gene is absent and *EhCP5* is degenerate (Bruchhaus *et al.* 1996; Willhoeft *et al.* 1999). Concerning lectins, it has been reported that Hgl is present in *E. dispar*, but that its expression level is lower than that in *E. histolytica* (Pillai *et al.* 1997, 2001). Therefore, it is of interest to determine if Igl1 and Igl2 are expressed in *E. dispar*. We report here the primary structure of the 2 Igl isoforms in 2 *E. dispar* strains isolated from human and cynomolgus monkey, respectively. We also compared the expression levels of Igl genes between *E. dispar* and *E. histolytica*, and examined the expression and localization of Igl1 and Igl2 in *E. dispar*.

#### MATERIALS AND METHODS

##### *Cultivation of parasites*

Trophozoites of the *E. dispar* SAW1734RcLAR strain were grown axenically or monoxenically with sterilized *Crithidia fasciculata* in YIGADHA-S medium supplemented with 15% adult bovine serum at 37 °C (Kobayashi *et al.* 2005; Khalifa *et al.* 2006). Trophozoites of the *E. dispar* CYNO9:TPC strain were axenically cultured in the YIGADHA-S medium. Trophozoites of *E. histolytica* HM-1:IMSS were axenically cultured in TYI-S-33 medium supplemented with 15% adult bovine serum at 37 °C (Diamond *et al.* 1978). Cultured trophozoites were harvested in the logarithmic phase of growth and used in subsequent experiments.

##### *Construction of a cDNA library and cloning of the Igl gene*

Poly(A) RNA of *E. dispar* SAW1734RcLAR trophozoites was isolated using a QuickPrep mRNA purification kit (Amersham Pharmacia). A cDNA library was constructed from 5 µg of poly(A) RNA using a cDNA synthesis kit (Amersham Pharmacia) and a λgt11 vector kit (Stratagene). The library was screened with a 657 bp probe using the Gene Images AlkPhos Direct labelling and detection system (Amersham Pharmacia). The probe was prepared from plasmid DNA containing the *E. histolytica* Igl1 gene by PCR amplification using primers EhIgl1-S877 (5'-CCC TCG AGT CAA ATG GTG AAT GTA AGC C-3') and EhIgl1-AS1088 (5'-CCC TCG AGT TAA ATG CCT TTA GCT CCA TT-3') (Tachibana *et al.* 2004). The positive clone containing the longest insert was subcloned into a pUC19 vector and sequenced. To extend the sequence of the 5' end, rapid amplification of the

cDNA end was performed with a 5'-Full RACE Core Set (Takara). For the cloning of the other Igl gene, the cDNA library was subjected to PCR using primers 5'-CAA TTT CAC TTG GTG AGT ACA AAG CTG-3' (forward) and 5'-GAA AAT TCC TTT ACT TCC ATT GCA GTT TCC-3' (reverse). These primers were prepared based on the sequence of the first cloned *E. dispar* Igl gene, with reference to the location of common sequences between the two *E. histolytica* Igl genes. The amplified genes were cloned using a TOPO TA Cloning Kit (Invitrogen) and sequenced. To extend the sequence of the 5' and 3' ends of the cloned DNA, a 5'-Full RACE Core Set and 3'-Full RACE Core Set were used (Takara). For the cloning of Igl genes from the CYNO9:TPC strain, genomic DNA isolated as previously described (Tachibana *et al.* 1991) was used as a template for PCR, using the forward primers 5'-ATG TTT ATT ATT CTT TTA TTC ATA TCA ATT TCA C-3' (Igl1) and 5'-ATG TTT ATT CTT CTT TTA TTT ATA TCA ATT TCA C-3' (Igl2), and the reverse primer 5'-TTA GAA CAT AAA TGA TAA CAT GAC TAT CAC CAT C-3'. Thirty-five cycles of PCR using *Pyrobest* DNA polymerase (Takara) were performed as follows: denaturation at 94 °C for 15 s (195 s in cycle 1), annealing at 58 °C for 30 s, and polymerization at 72 °C for 180 s (600 s in cycle 35). Amplified DNA was cloned using a Zero Blunt TOPO PCR Cloning Kit (Invitrogen) and sequenced. Nucleotide sequence data were analysed using Genetyx-Mac ver. 11.

##### *Southern blot analysis*

Genomic DNA was isolated from *E. dispar* SAW-1734RcLAR trophozoites as described previously (Tachibana *et al.* 1991). Three µg of genomic DNA was digested with restriction enzymes *DraI*, *TaqI* and *HindIII*. The fragments were separated on a 1% agarose gel, transferred to a Hybond N<sup>+</sup> membrane (Amersham Pharmacia) by capillary action, and fixed by alkaline denaturation. The membrane was hybridized at 55 °C in buffer containing a Gene Images AlkPhos Direct-labelled probe (Amersham Pharmacia) prepared by PCR amplification of cloned cDNA. The primers used for amplification were 5'-AGA TGG ATT CTA TTT TGA TGA-3' (forward) and 5'-CAT ATG TCT TGA ACA TGG-3' (reverse). The blots were detected using a CDP-star detection reagent (Amersham Pharmacia) and exposed to autoradiography films.

##### *Real-time RT-PCR analysis*

Total RNAs of *E. dispar* and *E. histolytica* trophozoites isolated using an RNeasy mini kit (Qiagen) were used for cDNA synthesis with a GeneAmp RNA PCR kit (Applied Biosystems). Reaction mixtures for quantitative real-time PCR analysis were

prepared using SYBR Premix Ex *Taq* (Takara), specific primers, Rox dye, and the cDNAs. The primers used were as follows: 5'-TGA CAA AGA CAA TAC TTG TAA AAA GTG-3' (forward) and 5'-ATT ACT AAC ACA TGC ACA TTT TTT GTC-3' (reverse) for *E. dispar Igl1* genes; 5'-TCG ATG AAA ATA ATG TAT GCC AGA AAT-3' (forward) and 5'-TCA TCA AGG CAA GCA CAT TGA CTG-3' (reverse) for *E. dispar Igl2* genes; 5'-GTT CAC AGG TTG GTG CTT GTA CG-3' (forward) and 5'-ACA GTA CAT GGC TTT TCT CCG GTA-3' (reverse) for *E. histolytica Igl1* genes; 5'-GAT TCA CAA ACA AAG GAG TGT GCC-3' (forward) and 5'-GTG CAT TTG AAC CAC TAG CAG CAA-3' (reverse) for *E. histolytica Igl2* genes; and 5'-CCA GCT ATG TAT GTT GGA ATT CAA G-3' (forward) and 5'-GAT CAA GTC TAA GAA TAG CAT GTG G-3' (reverse) for *actin* genes. Forty cycles of amplification with recording of fluorescence intensity in each cycle were performed using an ABI PRISM 7700 Sequence Detection System (Applied Biosystems). After initial denaturation at 95 °C for 10 sec, a shuttle PCR protocol consisting of denaturation at 95 °C for 5 sec and annealing-extension at 60 °C for 30 sec was applied. Relative quantitation with data from the ABI PRISM 7700 Sequence Detection System software version 1.7 was performed by the comparative  $C_T$  method, using the *actin* gene as an internal standard. The experiments were repeated 3 times, including the steps of culture of trophozoites and isolation of RNA.

#### Expression of recombinant Igl1 and Igl2

DNA fragments encoding full-length Igls, except for the N-terminus and C-terminus signal sequences, were obtained by PCR amplification of cloned *Igl* genes. Primers EdIgl1-S14-Xho (5'-CCC TCG AGG AGT ACA AAG CTG ATA AAC T-3') and EdIgl-AS-Xho (5'-CCC TCG AGT TAA ATT CCT TTA CTT CCA TT-3') were used for amplification of the *Igl1* gene of SAW1734RcLAR. For amplification of the *Igl2* gene of SAW1734RcLAR, primers EdIgl2-S14-Xho (5'-CCC TCG AGG ATT ACA AAG CTG ATA AAC TCA TC-3') and EdIgl-AS-Xho were used. PCR was performed as previously described (Tachibana *et al.* 2004). Each amplified DNA fragment was digested with *Xho*I, purified, and ligated with pET19b vector (Novagen). The plasmid was introduced into competent *Escherichia coli* JM109 cells and the direction and sequence of inserts were confirmed. *E. coli* BL21Star(DE3)pLysS cells (Invitrogen) were transformed with the cloned plasmids. Each clone was cultured in 400 ml of Luria-Bertani medium containing ampicillin until the culture reached an optical density of 0.6 at 600 nm. Isopropyl- $\beta$ -D-thiogalactopyranoside was added to the cultures

at a final concentration of 1 mM, and the cultures were incubated at 37 °C for 3 h. Preparation of inclusion bodies and refolding of the proteins were performed as previously described (Tachibana *et al.* 2004).

#### Production of specific mAbs

MAbs to Igl1 and Igl2 of *E. dispar* SAW1734RcLAR were prepared as follows. Six-week-old male BALB/c mice were inoculated intraperitoneally with 10  $\mu$ g of recombinant proteins in Freund's complete adjuvant and were inoculated again after 2 weeks. After an additional 3 weeks, the mice received only recombinant proteins. Four days later, spleen cells of immunized mice were isolated and fused with X63 Ag8.653 mouse myeloma cells using 50% polyethylene glycol 1500. Hybridomas secreting mAbs against *E. dispar* Igls were screened by immunofluorescent staining and ELISA, and were cloned by limiting dilution. Immunoglobulin isotypes of mAbs were determined by immunofluorescent staining using subtype-specific antibodies. Ascites was obtained by intraperitoneal inoculation of hybridomas into pristinely-primed mice, and immunoglobulin was purified using an Affi-Gel protein A MAPS II kit (Bio-Lab).

#### Dot blot analysis

Recombinant Igls and sonicated trophozoites of *E. dispar* SAW1734RcLAR were blotted on the nitrocellulose membrane using a Bio-Dot microfiltration apparatus (Bio-Rad). Filter strips were blocked with 3% bovine serum albumin (BSA) in phosphate-buffered saline (PBS) and reacted with mouse anti-*E. dispar* Igl mAbs for 30 min. After being washed with PBS containing 0.05% Tween-20 (PBS-Tween), the strips were incubated with horseradish peroxidase (HRP)-labelled goat anti-mouse IgG antibody (MP Biomedicals) for 30 min. The strips were then washed with PBS-Tween and developed with a Konica Immunostaining HRP-1000 kit.

#### SDS-PAGE and Western blot analysis

Recombinant Igl proteins or *E. dispar* trophozoites were treated with Laemmli's sample buffer (Laemmli, 1970) containing 2 mM phenylmethylsulfonyl fluoride, 2 mM *N*- $\alpha$ -*p*-tosyl-L-lysine chloromethylketone, 2 mM *p*-hydroxymercuriphenylsulfonic acid, and 4  $\mu$ M leupeptin for 5 min at 95 °C and then subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Western blot analysis was performed as previously described (Tachibana *et al.* 2004).

### Flow cytometry

Immunophenotypic surface staining of Igl1 and Igl2 using flow cytometry was performed on trophozoites of *E. dispar* SAW1734RclAR strain. Intact cells were incubated on ice with 3% BSA in PBS for 15 min, and then with a mixture of mAbs ED1-13 and ED2-1 for 15 min. After washing with ice-cold PBS, the cells were incubated with a mixture of Alexa Fluor 488-labelled goat anti-mouse IgG2b-specific antibody (Molecular Probes) and PE-labelled goat anti-mouse IgG1-specific antibody (Santa Cruz Biotechnology) for 15 min on ice. The cells were washed with ice-cold PBS and then fixed in 4% paraformaldehyde. Aliquots of approximately 5000 cells per sample were analysed using FACS Calibur (Becton Dickinson), with data analysis using CellQuest Software (BD Immunocytometry systems).

### Confocal microscopy

*E. dispar* SAW1734RclAR trophozoites were fixed with 4% paraformaldehyde in PBS and attached to silane-coated glass slides using Shandon Cytospin 2. After washing with PBS, the glass slides were incubated with 10% sucrose in PBS for 1 h and then stored at  $-80^{\circ}\text{C}$  until use. For double staining of Igl1 and Igl2, fixed trophozoites on the slides were treated with 0.1% Triton X-100 in PBS for 5 min, blocked with 3% BSA in PBS for 30 min and then incubated for 1 h at room temperature with a mixture of 2 mAbs, ED1-13 and ED2-1. After washing, the slides were incubated with a mixture of Alexa Fluor 488-labelled goat anti-mouse IgG2b-specific antibody and Alexa Fluor 594-labelled goat anti-mouse IgG1-specific antibody (Molecular Probes) for 1 h. The stained trophozoites were mounted using glycerol containing 1.25 mg/ml 1,4-diazabicyclo(2,2,2)octane and 10% PBS, and the samples were observed using a Zeiss LSM510 META confocal laser scanning microscope.

## RESULTS

### Cloning of genes encoding *E. dispar* Igl1 and Igl2

Two *Igl* genes cloned from the cDNA library from the *E. dispar* SAW1734RclAR strain encoded proteins of 1110 and 1106 amino acids, respectively, with calculated molecular masses of 120.9 kDa and 120.3 kDa and theoretical *pI* values of 5.5 and 4.87, respectively (DDBJ, EMBL, and GenBank Accession numbers AB287423 and AB287424). Two *Igl* genes were also cloned from the genomic DNA of the *E. dispar* CYNO9:TPC strain. These genes also encoded proteins of 1110 and 1106 amino acids, respectively, with calculated molecular masses of 121.0 kDa and 120.4 kDa and theoretical *pI* values of 5.41 and 4.74, respectively (DDBJ, EMBL,

and GenBank Accession numbers AB287425 and AB287426). Based on the similarity of the *pI* values to those of Igl of *E. histolytica* (5.52 for Igl1 and 5.17 for Igl2) (Cheng *et al.* 2001), the former protein was designated as Igl1 and the latter as Igl2 in both *E. dispar* strains. Multiple alignments of amino acid sequences among these Igl proteins and other *E. histolytica* Igl are shown in Fig. 1; the amino acid identity between *E. dispar* Igl1 and Igl2 was 79% in both strains. Differences in amino acids between the proteins in the two *E. dispar* strains were greater for Igl2 than for Igl1. In comparison with the *E. histolytica* Igl isoforms, the amino acid sequence identities were 75–76% for Igl1 and 73–74% for Igl2. Insertions of 6 amino acids in Igl1 and Igl2 are present in the two *E. dispar* strains around position 840. However, all cysteine residues found in *E. histolytica* Igl proteins were conserved in the *E. dispar* proteins. Both *E. dispar* Igl isoforms also contained hydrophobic amino- and carboxy-terminal signal sequences consistent with a GPI-anchored plasma membrane protein, and all Igl sequences contained a signature epidermal growth factor-like domain close to the C-terminus. The nucleotide sequence identities were 88% between the *E. dispar* Igl1 and Igl2 genes, 83% between the *E. dispar* and *E. histolytica* Igl1 genes, and 82% between the *E. dispar* and *E. histolytica* Igl2 genes.

A BLAST search of *E. dispar* SAW760 strain genomic sequences (<http://www.ncbi.nlm.nih.gov/BLAST/>) was performed, although the *E. dispar* genome project is incomplete. An identical sequence to that of the Igl1 gene of the SAW1734RclAR strain was found (Genbank Accession no. AANV01000026), and 2 sequences (AANV01000644 and AANV01001389) were identified that showed 99% identity with the Igl2 gene of SAW1734RclAR in partially overlapping regions.

### Southern blot analyses of Igl genes

Southern blot hybridization using a 420-bp PCR product as a probe was performed on *E. dispar* SAW1734RclAR genomic DNA digested with *Dra*I, *Taq*I and *Hind*III (Fig. 2). The results indicated the presence of 2 *Igl* genes in the *E. dispar* genomic DNA.

### Real-time RT-PCR analysis of Igl genes

Expression levels of Igl1 and Igl2 are compared in Fig. 3. In the SAW1734RclAR strain, expression of Igl1 was 3 times higher than that of Igl2 ( $P < 0.001$ ), and in the CYNO9:TPC strain, expression of Igl1 was 5 times higher than that of Igl2 ( $P < 0.01$ ). Higher expression of Igl1 compared to Igl2 was also observed in *E. histolytica* ( $P < 0.001$ ). Expression of Igl1 was lower in *E. dispar* than in *E. histolytica* (SAW1734RclAR *vs* HM-1:IMSS,  $P < 0.01$ ;




Fig. 1. Alignment of the deduced amino-acid sequences of the *Ig1* and *Ig2* genes from *Entamoeba dispar* SAW1734RclAR (Ed.SAW1734), *E. dispar* CYN09:T1PC (Ed.CYN09) and *E. histolytica* HM-1:IMSS (Eh.HM1). Identical and conserved amino-acid residues are highlighted in black and grey, respectively.

CYN09:T1PC vs HM-1:IMSS, not significant), whereas *Ig2* expression was similar in the two species.

*Dot blot and Western blot analyses of Igls*

Full length (except for the signal sequence) recombinant *E. dispar* *Igls* from the SAW1724RclAR strain were prepared in *Escherichia coli*. The

apparent molecular weight of the recombinant protein with the leader peptide was slightly larger for *Ig2* (190 kDa) than for *Ig1* (170 kDa) in SDS-PAGE under reducing conditions (data not shown). Recombinant *Ig1* and *Ig2* proteins were used for immunization of mice to prepare mAbs specific for each *Ig*. In dot blot analysis, mAb ED1-13 reacted specifically with recombinant *Ig1*, but not with recombinant *Ig2*; in contrast, mAb ED2-1 was




Fig. 2. Southern blot analysis of *Igl* genes in *Entamoeba dispar* SAW1734RclAR. Genomic DNA was digested with *Dra*I (D, lane 1), *Taq*I (T, lane 2) and *Hind*III (H, lane 3) and hybridized with the probe. The blot is representative of 2 independent experiments. Numbers to the right indicate the sizes of DNA markers (in kilobases).




Fig. 3. Real-time reverse transcription PCR analysis of *Igl* genes from *Entamoeba dispar* and *E. histolytica*. Expression levels of *Igl1* (open bars) and *Igl2* (filled bars) in trophozoites from *E. dispar* SAW1734RclAR, *E. dispar* CYN09:TPC and *E. histolytica* HM-1:IMSS are expressed as values relative to the expression level of *actin*. Vertical bars indicate the s.e. of the mean from 3 experiments.

reactive specifically with recombinant *Igl2* (Fig. 4A). In conditions under which the reactivity of these mAbs to *E. dispar* trophozoites was comparable, differences in reactivity to equal amounts of recombinant Igls were observed. The reactivity of mAb ED1-13 to  $10^4$  trophozoites was similar to the reactivity to  $1 \mu\text{g}$  of EdIgl1, whereas the reactivity of mAb ED2-1 to  $10^4$  trophozoites were comparable to the reactivity to  $0.1 \mu\text{g}$  of EdIgl2, suggesting that the amount of *Igl2* was approximately one-tenth that of *Igl1* in the trophozoites. In Western blot analysis using trophozoites from the SAW1734RclAR strain, mAbs ED1-13 and ED2-1 recognized a 100 kDa and





Fig. 4. (A) Reactivity of anti-*Igl* monoclonal antibodies to recombinant Igls and crude antigens of *Entamoeba dispar* in dot blot analysis. Various concentrations (0.01, 0.1 and  $1 \mu\text{g}$ ) of recombinant *Igl1* (rIgl1) and recombinant *Igl2* (rIgl2) and various numbers ( $10^2$ ,  $10^3$  and  $10^4$ ) of sonicated trophozoites from strain SAW1734RclAR were spotted on nitrocellulose membranes. One strip was stained with Coomassie brilliant blue (CBB). Other strips were treated with anti-*E. dispar* *Igl* monoclonal antibodies (ED1-13, ED2-1) and anti-*E. histolytica* *Igl* monoclonal antibody (EH3015). HRP-conjugated goat antibody to mouse IgG was used as a secondary antibody. The blot is representative of 2 independent experiments. (B) Western immunoblot analysis of native Igls of *E. dispar*. Lysates of SAW1734RclAR trophozoites were subjected to SDS-PAGE in a 7.5% gel under non-reducing conditions and transferred to polyvinylidene difluoride membranes. Protein bands of the size marker in lane 1 were stained with Coomassie brilliant blue. The strips were treated with monoclonal antibodies as follows: lane 2, ED1-13; lane 3, ED2-1; and lane 4, EH3015. HRP-conjugated goat antibody to mouse IgG was used as a secondary antibody. The blot is representative of 3 independent experiments. The numbers to the left indicate molecular masses (in kilodaltons).

a 120 kDa band, respectively, under non-reducing conditions (Fig. 4B). No bands were detected in the Western blot under reducing conditions, indicating that the mAbs recognized discontinuous epitopes on the *Igl* proteins.

*Phenotypic expression of Igls on the surface of trophozoites*

To compare the amounts of *Igl1* and *Igl2* expressed on the surface of trophozoites from the *E. dispar* SAW1724RclAR strain, flow cytometric analysis was performed using specific mAbs for each Igls (Fig. 5). The results demonstrated that almost all






Fig. 5. Flow cytometric analysis of phenotypic expression of Igl1 and Igl2 on the surface of *Entamoeba dispar* trophozoites. Intact trophozoites from the SAW1734Rc1AR strain were double stained with monoclonal antibodies ED1-13 (specific for Igl1, IgG2b) and ED2-1 (specific for Igl2, IgG1), followed by Alexa Fluor 488-labelled goat anti-mouse IgG2b-specific antibody and PE-labelled goat anti-mouse IgG1-specific antibody (A). The control was stained only with secondary antibodies (B). The results are representative of 3 independent experiments.

trophozoites expressed both Igl1 and Igl2 on the cell surface.

#### Localization of Igls on trophozoites

Localization of Igl1 and Igl2 on *E. dispar* trophozoites was examined by confocal laser scanning microscopy using specific mAbs. Both Igls were localized on the plasma membrane and in cytoplasm in all trophozoites (Figs 6 and 7). However, the amount of each Igl in the trophozoites was variable, especially in the cytoplasm; that is, Igl1- and Igl2-dominant cells were present (arrows and arrowheads in Fig. 6). In addition, Igl1- and Igl2-dominant vacuoles were also observed within a single trophozoite (arrow and arrowhead in Fig. 7).

#### DISCUSSION

Comparison of the 2 *Igl* genes cloned from *E. dispar* with those from *E. histolytica* indicated differences in the sequences of the corresponding proteins in *E. dispar* and *E. histolytica* and in the 2 strains of *E. dispar*. However, all the cysteine residues, including the CXXC motifs, were conserved between species and between strains, which suggests that the fold of the protein is maintained and is important for its function. Amino acid identities of Igl1 and Igl2 within species (79% in *E. dispar* and 81% in *E. histolytica*; Cheng *et al.* 2001) were higher than those for each Igl between the 2 species (75–76% for Igl1 and 73–74% for Igl2). Insertions of 6 amino acids around position 840 in *E. dispar* were present in both Igl1 and Igl2, as shown in Fig. 1. Nucleotide identities of the 2 *Igl* genes were also higher within

species, compared to the respective identities of the *Igl* genes between *E. dispar* and *E. histolytica*. These results suggest that duplication of the genes may have occurred after divergence of the species.

The properties of the Gal/GalNAc lectin of *E. histolytica* were demonstrated in a 150 kDa fraction purified by affinity chromatography using the *E. histolytica*-specific mAb EH3015 (Cheng *et al.* 1998). In addition, Igl1 and Igl2 of *E. histolytica* have been detected, in addition to Hgl and Lgl, in the protein fraction that binds specifically to GalNAc-BSA-coated magnetic beads (McCoy and Mann, 2005). However, when we performed a preliminary examination of the reactivity of recombinant Igls from *E. histolytica* and *E. dispar* with GalNAc<sub>27</sub>-BSA by dot blot analysis and surface plasmon resonance, we could not prove that the recombinant proteins had sugar-binding properties (data not shown). Therefore, the Igls may exist as part of the lectin complex, perhaps with non-covalent association to another protein containing a sugar-binding site, with this association occurring either directly or being mediated by a third protein. Recently, it has been reported that Igl of *E. histolytica* is found in the protein fraction that interacts with purified brush border from human enterocytes (Seigneur *et al.* 2005). It has also been demonstrated that the 140 kDa fibronectin-binding molecule of *E. histolytica* (Talamas-Rohana *et al.* 1992) is identical with Igl2 (Hernandez-Ramirez *et al.* 2007). These observations indicate that Igls are important proteins for amoebic adherence to host cells, and since Igls are also expressed in the non-pathogenic amoeba, it seems likely that these proteins are important for colonization of amoebae in the large intestine.




Fig. 6. Localization of Ig11 and Ig12 on trophozoites of *Entamoeba dispar* SAW1734RclAR observed by confocal laser scanning microscopy. Fixed trophozoites were stained with ED1-13 specific for Ig11 and a secondary Alexa Fluor 488-labelled anti-mouse IgG2b antibody (green) (A) or ED2-1 specific for Ig12 and a secondary Alexa Fluor 594-labelled anti-mouse IgG1 antibody (red) (B). Differential interference contrast microscopy is shown in (C). A merged image is shown in (D). Arrows and arrowheads indicate Ig11- and Ig12-dominant cells, respectively.

One of the interesting observations in this study is the difference in expression between Ig11 and Ig12. Since higher expression of Ig11 was observed at both protein and mRNA levels, the difference between the isoforms seems to be regulated mostly at the transcriptional level. The expression level of *Igl1* was also lower in *E. dispar* than in *E. histolytica*, whereas that of *Igl2* was comparable in the two species. It has been demonstrated that expression of *Hgl* (*Hgl2*) and *Lgl* (*Lgl1*) in *E. dispar* is lower than in *E. histolytica* (Pillai *et al.* 1997, 2001); therefore, Ig11 may be more closely associated with *Hgl* and *Lgl*. However, it is unknown whether the 2 Ig1 isoforms are

associated with different isoforms of *Hgl* or *Lgl*. DNA microarray analyses have shown that a large number of genes are expressed differently in *E. histolytica* and *E. dispar*, and that there is a difference in gene expression between strains of *E. histolytica* of high and low virulence (Shah *et al.* 2005; MacFarlane and Singh, 2006; Davis *et al.* 2007). Lower expression of *Hgl* genes in *E. dispar* compared to *E. histolytica* has also been confirmed (MacFarlane and Singh, 2006), but *Lgl3* expression was found to be higher in an *E. histolytica* strain of low virulence compared to a strain of high virulence (Davis *et al.* 2007). *Igl* genes are down-regulated by




Fig. 7. Stereo images of localization of Ig11 (A) and Ig12 (B) on *Entamoeba dispar* trophozoites observed by confocal laser scanning microscopy. Fixed trophozoites were stained as described in Fig. 6. Differential interference contrast microscopy is shown in (C). A merged image is shown in (D). Arrow and arrowhead indicate single localization of Ig11 and Ig12, respectively.

heat shock stress, as are most *Hgl* and *Lgl* genes (Weber *et al.* 2006). Collectively, these observations suggest that the lectin as a whole is important for adherence and subsequent pathogenesis.

We also compared phenotypic expression of the two Igl in each trophozoite using flow cytometry and confocal microscopy. Interestingly, some trophozoites expressed Igl1 or Igl2 dominantly, although most cells expressed the two Igl proteins to a comparable extent, suggesting that both Igl are important for the amoeba. Igl were also localized in intracellular vacuoles of *E. dispar*. Recently, it has been demonstrated that Igl is contained in phagosomes of *E. histolytica* and that the quantity of Igl varies during phagosome maturation (Okada *et al.* 2005, 2006). The different localization of Igl1 and Igl2 in vacuoles suggests that the functions of the two Igl may differ or that their expression may vary during maturation of phagosomes or depending on certain cellular conditions. However, there may also be differences in the involvement of microtubules and proteases in phagosome maturation and degradation in *E. histolytica* and *E. dispar* (Mitra *et al.* 2005). In *E. histolytica*, different subcellular localization of the two Igl isoforms has yet to be shown.

In conclusion, this is the first study of the differences between Igl1 and Igl2 of *E. dispar*. Igl seems to be a vaccine candidate for amebiasis and may also be a useful antigenic molecule for specific serodiagnosis of amoebiasis (Cheng and Tachibana, 2001; Tachibana *et al.* 2004). Therefore, further studies of Igl are required to clarify its role in the host-parasite relationship.

This work was supported by a Grant-in-Aid for Scientific Research from the Japanese Society for the Promotion of Science, and a grant for Research on Emerging and Re-emerging Infectious Diseases from the Ministry of Health, Labour and Welfare of Japan. X.-J. C. is a recipient of a Postdoctoral Fellowship for Foreign Researchers from the Japanese Society for the Promotion of Science.

## REFERENCES

- Bruchhaus, I., Jacobs, T., Leippe, M. and Tannich, E.** (1996). *Entamoeba histolytica* and *Entamoeba dispar*: differences in numbers and expression of cysteine proteinase genes. *Molecular Microbiology* **22**, 255–263.
- Cheng, X.-J., Hughes, M. A., Huston, C. D., Loftus, B., Gilchrist, C. A., Lockhart, L. A., Ghosh, S., Miller-Sims, V., Mann, B. J., Petri, W. A., Jr. and Tachibana, H.** (2001). Intermediate subunit of the Gal/GalNAc lectin of *Entamoeba histolytica* is a member of a gene family containing multiple CXXC sequence motifs. *Infection and Immunity* **69**, 5892–5898.
- Cheng, X.-J., Kaneda, Y. and Tachibana, H.** (1997). A monoclonal antibody against the 150-kDa surface antigen of *Entamoeba histolytica* inhibits adherence and cytotoxicity to mammalian cells. *Medical Science Research* **25**, 159–161.
- Cheng, X.-J. and Tachibana, H.** (2001). Protection of hamsters from amebic liver abscess formation by immunization with the 150- and 170-kDa surface antigens of *Entamoeba histolytica*. *Parasitology Research* **87**, 126–130.
- Cheng, X.-J., Tachibana, H. and Kaneda, Y.** (1999). Protection of hamsters from amebic liver abscess formation by a monoclonal antibody to a 150-kDa surface lectin of *Entamoeba histolytica*. *Parasitology Research* **85**, 78–80.
- Cheng, X.-J., Tsukamoto, H., Kaneda, Y. and Tachibana, H.** (1998). Identification of the 150-kDa surface antigen of *Entamoeba histolytica* as a galactose- and *N*-acetyl-D-galactosamine-inhibitable lectin. *Parasitology Research* **84**, 632–639.
- Davis, P. H., Schulze, J. and Stanley, S. L., Jr.** (2007). Transcriptomic comparison of two *Entamoeba histolytica* strains with defined virulence phenotypes identifies new virulence factor candidates and key differences in the expression patterns of cysteine proteases, lectin light chains, and calmodulin. *Molecular and Biochemical Parasitology* **151**, 118–128.
- Diamond, L. S. and Clark, C. G.** (1993). A redescription of *Entamoeba histolytica* Schaudinn, 1903 (Emended Walker, 1911) separating it from *Entamoeba dispar* Brumpt, 1925. *Journal of Eukaryotic Microbiology* **40**, 340–344.
- Diamond, L. S., Harlow, D. R. and Cunnick, C. C.** (1978). A new medium for the axenic cultivation of *Entamoeba histolytica* and other *Entamoeba*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **72**, 431–432.
- Hernández-Ramírez, V. I., Rios, A., Angel, A., Magos, M. A., Pérez-Castillo, L., Rosales-Encina, J. L., Castillo-Henkel, E. and Talamás-Rohana, P.** (2007). Subcellular distribution of the *Entamoeba histolytica* 140 kDa FN-binding molecule during host-parasite interaction. *Parasitology* **134**, 169–177.
- Khalifa, S. A., Imai, E., Kobayashi, S., Haghghi, A., Hayakawa, E. and Takeuchi, T.** (2006). Growth-promoting effect on iron-sulfur proteins on axenic cultures of *Entamoeba dispar*. *Parasite* **13**, 51–58.
- Kobayashi, S., Imai, E., Haghghi, A., Khalifa, S. A., Tachibana, H. and Takeuchi, T.** (2005). Axenic cultivation of *Entamoeba dispar* in newly designed yeast extract-iron-gluconic acid-dihydroxyacetone-serum medium. *Journal of Parasitology* **91**, 1–4.
- Laemmli, U. K.** (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature, London* **227**, 680–685.
- MacFarlane, R. C. and Singh, U.** (2006). Identification of differentially expressed genes in virulent and nonvirulent *Entamoeba* species: potential implications for amebic pathogenesis. *Infection and Immunity* **74**, 340–351.
- McCoy, J. J. and Mann, B. J.** (2005). Proteomic analysis of Gal/GalNAc lectin-associated proteins in *Entamoeba histolytica*. *Experimental Parasitology* **110**, 220–225.
- Mitra, B. N., Yasuda, T., Kobayashi, S., Saito-Nakano, Y. and Nozaki, T.** (2005). Differences in morphology of phagosomes and kinetics of acidification and degradation in phagosomes between the pathogenic *Entamoeba histolytica* and the non-pathogenic *Entamoeba dispar*. *Cell Motility and the Cytoskeleton* **62**, 84–99.