

FIG. 1. HCV core protein associates with E6AP in vivo. (A) 293T cells were transfected with pcDNA3-MEF-core or empty plasmid, incubated for 48 h, and then harvested. The expressed MEF-core and binding proteins were recovered using the MEF purification procedure. Proteins bound to the MEF-core immobilized on anti-FLAG beads were dissociated with FLAG peptides, resolved by 9% SDS-PAGE, and visualized by silver staining. Control experiments were performed using 293T cells transfected with vector alone. The positions of E6AP, DDX3, and PA28 γ are indicated by arrows. (B to E) The proteins detected in panel A were confirmed by immunoblotting with appropriate antibodies: E6AP (B), DDX3 (C), PA28 γ (D), and MEF-core (E).

ability of GST-E6AP (418–517) to bind to the HCV core protein was confirmatory (Fig. 2C, lane N) and led to the conclusion that the HCV core-binding domain of E6AP was aa 418 to aa 517.

The HCV core-binding domain for E6AP. By use of a panel of HCV core deletion mutants (Fig. 3A), GST-E6AP was found to coimmunoprecipitate with all of the FLAG-core proteins (Fig. 3A, lanes A to H) except FLAG-core (72–191) or FLAG-core (92–191) (Fig. 3A, lanes I and J). No association of control GST protein with any FLAG-core proteins was observed (data not shown). These data suggest that the aa-58-to-aa-71 segment of the HCV core binds to E6AP. The ability of GST-core (58–71) to associate with purified MEF-E6AP confirmed that the core (aa 58–71) was the site for E6AP binding on the HCV core protein (Fig. 3B).

E6AP decreases steady-state levels of HCV core protein in 293T cells and HepG2 cells. One of the features of HECT domain ubiquitin ligases is direct association with their substrates (50). Thus, we hypothesized that E6AP would function as an E3 ubiquitin ligase for the HCV core protein. We as-

TABLE 1. Identification of E6AP by tandem mass spectrometry^a

Peptide <i>m/z</i>	Sequence determined	Residues
720.9	VFSSAEALVQSFRR	156–168
922.4	AACSAAAMEEDSEASSSR	196–213
774.9	MMETFQQLITYK	339–350
1,053.1	ITVLYSLVQGQQLNPYLRR	507–524
809.4	EFVISYSYDIILNK	712–724

^a The protein was ubiquitin protein ligase E3A (E6AP) isoform 2 (GenBank accession no. NP_000453).




FIG. 2. Mapping of the HCV core-binding domain for E6AP. (A) Structure of E6AP. Shown is a schematic representation of the regions of E6AP isoform II that mediate E6 binding (aa 401 to 418), E6-dependent association with p53 (aa 290 to 791), and the HECT catalytic domain (aa 525 to 875). The catalytic cysteine residue is located at aa 843. (B) Schematic representation of GST-E6AP proteins. GST proteins A through N contain the E6AP amino acids indicated to the right. The shaded region of each represents the GST sequence. Closed boxes represent proteins that are bound specifically to HCV core protein, and open boxes represent those that are not bound. (C) Binding of HCV core protein to GST-E6AP proteins A through N. In vitro-translated core protein (aa 1 to 173) was assayed for association with GST (-) or the GST-E6AP fusion proteins A through N. Association of core protein was detected by immunoblotting with anti-core MAb.

essed the effects of E6AP on the HCV core protein in 293T cells. FLAG-core (1–191) together with HA-tagged wild-type E6AP, catalytically inactive mutant E6AP, E6AP C-A (19), or WWP1 (another HECT domain ubiquitin ligase) (22) was introduced into 293T cells, and the levels of the core protein were examined by immunoblotting. The steady-state levels of the core protein decreased with an increase in the amount of E6AP plasmids (Fig. 4A and B). However, neither E6AP C-A mutant nor WWP1 decreased the steady-state levels of the core protein, suggesting that E6AP enhances degradation of the core protein.

To verify the critical need for endogenous E6AP in the core degradation, expression of E6AP was knocked down by siRNA and the expression of the core protein and E6AP was assayed by immunoblotting. Transfection of the E6AP-specific siRNA




FIG. 3. Mapping of the E6AP binding domain for HCV core protein. (A) In vitro binding of E6AP to HCV core protein. 293T cells were transfected with each plasmid indicated in the upper panel. At 48 h posttransfection, cell lysates were mixed with purified GST-E6AP, immunoprecipitated with anti-FLAG beads, and then immunoblotted with anti-GST PAb (middle panel) or anti-FLAG MAb (bottom panel). The last lane (input) represents GST-E6AP used in this assay (middle panel). (B) Binding of GST-core (aa 58 to aa 71) to purified MEF-E6AP. GST served as a negative control for binding. Upper panel, Coomassie blue-stained SDS-PAGE of GST and GST-core (58-71). Lower panel, results of the GST pull-down assay. MEF-E6AP was detected by anti-myc MAb. CBB, Coomassie brilliant blue; IB, immunoblot.

duplex reduced the protein level of E6AP by 90% at 48 h posttransfection (Fig. 4C, middle panel). Immunoblotting revealed a 4.1-fold increase in the level of the core protein in the cells transfected with E6AP siRNA (Fig. 4C, top panel), suggesting that endogenous E6AP plays a role in the proteolysis of the HCV core protein.

Then we examined whether E6AP reduces the steady-state




FIG. 4. E6AP decreases steady-state levels of HCV core protein in 293T cells and in HepG2 cells. (A) 293T cells (1×10^6 cells/10-cm dish) were transfected with 1 μ g of pCAG FLAG-core (1-191) along with either pCAG-HA-E6AP, pCAG-HA-E6AP C-A, or pCAG-HA-WWP1 as indicated. At 48 h posttransfection, protein extracts were separated by SDS-PAGE and analyzed by immunoblotting with anti-HA PAb (top panel), anti-FLAG MAb (middle panel), and anti-GAPDH MAb (bottom panel). (B) Quantitation of data shown in panel A. Intensities of the gel bands were quantitated using the NIH Image 1.62 program. The level of GAPDH served as a loading control. Circles, E6AP; triangles, E6AP C-A; squares, WWP1. (C) Knockdown of endogenous E6AP by siRNA inhibits degradation of HCV core protein in 293T cells. 293T cells (3×10^5 cells/six-well plate) were transfected with 40 pmol of E6AP-specific duplex siRNA (or control siRNA) as described in Materials and Methods. The cells were transfected with 2 μ g of FLAG-core (1-191) expression plasmid and cultured for 24 h, harvested, and analyzed by immunoblotting. Shown is immunoblot detection of FLAG-tagged core protein (top panel), E6AP protein (middle panel), and GAPDH (bottom panel) in control siRNA-treated 293T cells or E6AP-siRNA-treated 293T cells. The relative levels of protein expression were quantitated by densitometry and indicated below in the respective lanes. GAPDH served as a loading control. (D) HepG2 cells (2×10^5 cells/six-well plate) were transfected with pCAG FLAG-core (1-152) along with either empty vector or pCMV E6AP as indicated. The cells were harvested at 44 h posttransfection. Where indicated, cells were treated with 25 μ M MG132 or with dimethyl sulfoxide control 14 h prior to collection. Equivalent amounts of the whole-cell lysates were separated by SDS-PAGE and analyzed by immunoblotting with anti-FLAG MAb (upper panel) or anti-GAPDH MAb (lower panel).




FIG. 5. Kinetic analysis of E6AP-dependent degradation of HCV core protein. (A) 293T cells (1×10^6 cells/10-cm dish) were transfected with 1 µg of pCAG-FLAG core (1–152) plus 4 µg of empty vector, pCMV-HA-E6AP, or pCMV-HA-E6AP C-A. The cells were treated with 50 µg/ml CHX at 44 h after transfection. Cell extracts were collected at 0, 3, 6, and 9 h after treatment with CHX, followed by immunoblotting. (B) Specific signals were quantitated by densitometry, and the percent remaining core at each time was compared with that at the starting point. The level of GAPDH served as a loading control. Open circles, E6AP; closed circles, empty plasmid; closed triangles, E6AP C-A; closed squares, E6AP with MG132 treatment. Data are representative of three independent experimental determinations.

levels of the core protein in hepatic cells as well as in 293T cells. Exogenous expression of E6AP resulted in reduction of the core protein in human hepatoblastoma HepG2 cells (Fig. 4D). Treatment of the cells with the proteasome inhibitor MG132 increased the core protein level, suggesting that the core protein was degraded through the ubiquitin-proteasome pathway. These results indicate that E6AP enhances proteasomal degradation of the HCV core protein in both hepatic cells and nonhepatic cells.

Kinetic analysis of E6AP-dependent degradation of HCV core protein. To determine whether the E6AP-induced reduction of the core protein is due to an increase in the rate of core degradation, we performed kinetic analysis using the protein synthesis inhibitor CHX. HCV core protein together with wild-type E6AP or inactive mutant E6AP C-A was expressed in 293T cells. At 44 h after transfection, cells were treated with either 50 µg/ml CHX alone or 50 µg/ml CHX plus 25 µM MG132 to inhibit proteasome function. Cells were collected at 0, 3, 6, and 9 h following treatment and analyzed by immunoblotting (Fig. 5A). Overexpression of E6AP resulted in rapid degradation of the core protein, whereas inactive mutant




FIG. 6. E6AP promotes degradation of full-length HCV core protein in Huh-7 cells. Huh-7 cells (2×10^5 cells/six-well plate) were transfected with 0.5 µg of pCAG-core (1–191) together with 2 µg of pCMV-HA-E6AP or pCMV-HA-E6AP C-A. At 48 h posttransfection, cells were harvested and analyzed by immunoblotting with anticore MAb (top panel), anti-E6AP PAb (middle panel), or anti-GAPDH MAb (bottom panel).

E6AP C-A increased the half-life of the core protein (Fig. 5B), suggesting that the inactive E6AP inhibited degradation of the core protein in a dominant-negative manner, which is in agreement with previous studies (19, 55). Treatment of the cells with MG132 inhibited the degradation of the core protein (Fig. 5B). Reverse transcription-PCR to determine mRNA levels of the HCV core gene and GAPDH gene found that neither wild-type E6AP nor inactive E6AP changed mRNA levels of the HCV core gene and GAPDH gene (data not shown). These results indicate that E6AP enhances proteasomal degradation of the core protein.

E6AP promotes degradation of the full-length core protein in Huh-7 cells. To determine whether the full-length HCV core protein expressed in hepatic cells is degraded through an E6AP-dependent pathway, human hepatoma Huh-7 cells were transfected with pCAG HCV core (1–191) along with either E6AP or E6AP C-A. To rule out the effects of N-terminal FLAG tag on the core degradation, HCV core protein was expressed as untagged protein. Expression of wild-type E6AP resulted in reduction of the core protein (Fig. 6). On the other hand, HCV core protein was not decreased after transfection of inactive E6AP, indicating that the full-length core protein expressed in Huh-7 cells is also degraded through an E6AP-dependent pathway.

E6AP mediates ubiquitylation of HCV core protein in vivo. To determine whether E6AP can induce ubiquitylation of HCV core protein in cells, we performed in vivo ubiquitylation assays. 293T cells were cotransfected with FLAG-core (1–191) and either E6AP or empty plasmid, together with a plasmid encoding HA-tagged ubiquitin to facilitate detection of ubiquitylated core protein. Cell lysates were immunoprecipitated with anti-FLAG MAb and immunoblotted with anti-HA PAb to detect ubiquitylated core protein (Fig. 7A). Only a little ubiquitin signal was observed on the core protein in the absence of cotransfected E6AP (Fig. 7A, lane 3). In contrast, coexpression of E6AP led to readily detectable ubiquitylated forms of the core protein as a ladder and a smear of higher-molecular-weight bands (Fig. 7A, compare lane 3 with lane 4). Immunoblot analysis with anticore PAb confirmed that FLAG-core proteins were immunoprecipitated (Fig. 7B, lanes 2 to 4, short exposure) and that higher-molecular-weight bands con-




FIG. 7. E6AP-dependent ubiquitylation of HCV core protein in vivo. 293T cells (1×10^6 cells/10-cm dish) were transfected with 1 μ g of pCAG FLAG-core (1–191) together with 2 μ g of plasmid encoding E6AP as indicated. Each transfection also included 2 μ g of plasmid encoding HA-ubiquitin. The cell lysates were immunoprecipitated with FLAG beads and analyzed by immunoblotting with anti-HA PAb (A) or anti-core PAb (B). A shorter exposure of the core blot shows immunoprecipitated FLAG-core protein (B, right panel). A longer exposure of the core blot shows the presence of a ubiquitin smear (B, left panel). Asterisks indicate cross-reacting immunoglobulin light chain or heavy chain. Arrows indicate FLAG-core. IB, immunoblot; IP, immunoprecipitation.

jugated with HA-ubiquitin were indeed ubiquitylated forms of the core protein (Fig. 7B, lanes 3 and 4, long exposure).

E6AP mediates ubiquitylation of HCV core protein in vitro. To rule out the possibility that E6AP contributes to core protein degradation by inducing degradation of inhibitors of core turnover, we determined whether E6AP functions directly as a ubiquitin ligase by testing the ability of purified MEF-E6AP to mediate in vitro ubiquitylation of the purified recombinant HCV core protein. HCV core protein was expressed as a fusion protein containing N-terminal GST tag and C-terminal His tag and purified as described in Materials and Methods. GST-C173HT (aa 1–173) and GST-C152HT (aa 1–152) (see Materials and Methods) were used to determine whether the mature core protein and the C-terminally truncated core protein are targeted for ubiquitylation in vitro. The validity of this assay was established by demonstrating that E6AP but not E6AP C-A induced ATP-dependent ubiquitylation of GST-core protein. When in vitro ubiquitylation reactions were carried out either in the absence of MEF-E6AP or in the presence of MEF-E6AP C-A, no ubiquitylation signal was detected (Fig. 8A, lanes 4 and 5). However, inclusion of purified MEF-E6AP in the reaction mixture resulted in marked ubiquitylation of GST-C173HT (Fig. 8A, lane 6), while no ubiquitylation was observed in the absence of ATP (Fig. 8A, lane 7). No signal was detected when GST-HT was used as a substrate (Fig. 8A, lane 8). The higher-molecular-weight species of GST-core proteins were reactive with both anti-ubiquitin MAb (Fig. 8B, right panel, lanes 2 and 4) and anti-GST MAb (Fig. 8B, left panel, lanes 2 and 4). Both GST-C152HT and GST-C173HT were polyubiquitylated by E6AP in vitro (Fig. 8B), indicating that both the C-terminally truncated core and the mature core are polyubiquitylated by E6AP in vitro. These results revealed

that E6AP directly mediated ubiquitylation of HCV core proteins in an ATP-dependent manner.

Exogenous expression of E6AP reduces intracellular HCV core protein levels and supernatant infectivity titers in HCV-infected Huh-7 cells. We used a recently developed system for the production of infectious HCV particles using the HCV JFH1 strain (28, 56, 61) to examine whether E6AP can promote degradation of HCV core protein expressed from infectious HCV. E6AP-dependent core degradation was assessed in Huh-7 cells inoculated with the culture supernatant containing HCV JFH1. Levels of HCV core protein were detectable at day 3 postinfection and increased with time. Immunofluorescence staining for the core protein indicated that the percentage of HCV core-positive cells in the Huh-7 cells was almost 100 at day 7 postinfection. Transfection efficiency was 50 to 60% as measured with GFP-expressing plasmid. At day 7 postinfection, exogenous expression of E6AP reduced the intracellular core protein level by about 60% compared to the empty plasmid-transfected control cells (Fig. 9A). Inactive E6AP had little effect on the core protein levels. Total protein levels in the cells (Fig. 9B) and intracellular HCV RNA levels (Fig. 9C) did not change after transfection of wild-type E6AP or inactive E6AP. The immunofluorescence study revealed that HCV core protein was variably detected and the intensity of core staining was reduced in the cells staining positive for wild-type E6AP compared with neighboring cells staining negative for E6AP (Fig. 9E). Using inactive E6AP revealed colocalization of the core protein and E6AP in the perinuclear region (Fig. 9F) of HCV-infected cells. These results suggest that E6AP enhanced degradation of HCV core protein expressed from infectious HCV. Then we titrated HCV infectivity in the culture supernatant at day 7 postinfection by limiting




FIG. 8. In vitro ubiquitylation of HCV core protein by recombinant E6AP. For in vitro ubiquitylation of HCV core protein, purified GST-C173HT and GST-C152HT were used as substrates. Purified GST-HT was used as a negative control. Assays were done in 40- μ l volumes containing each component as indicated. The reaction mixture is described in Materials and Methods. The reaction was carried out at 37°C for 120 min followed by purification with glutathione-Sepharose beads and analysis by immunoblotting with the indicated antibodies. Arrows indicate GST-C173HT, GST-C152HT, and GST-HT, respectively. Ubiquitylated species of GST-core proteins are marked by brackets. IB, immunoblot.

dilution assays. Exogenous expression of E6AP reduced the supernatant infectivity titer, whereas inactive E6AP had no effect on its infectivity titer (Fig. 9D), suggesting that the E6AP-dependent ubiquitin proteasome pathway affects the production of HCV particles through downregulation of the core protein.

E6AP silencing increases the levels of intracellular HCV core protein and supernatant infectivity titers in HCV-infected Huh-7 cells. Finally, to further validate the role of E6AP in HCV production, expression of endogenous E6AP was knocked down by siRNA and the HCV infectivity titers released from HCV JFH1-infected cells were examined. Knock-down of E6AP by siRNA led to an increase in intracellular core protein levels (Fig. 10A) and supernatant HCV infectivity titers (Fig. 10B). Taken together, our results suggest that E6AP mediates ubiquitylation and degradation of HCV core protein in HCV-infected cells, thereby affecting the production of HCV particles.

DISCUSSION

HCV core protein is a major component of viral nucleocapsid, plays a central role in viral assembly (25, 40), and contributes to viral pathogenesis and hepatocarcinogenesis (9). Therefore, it is important to clarify the molecular mechanisms that govern the cellular stability of this viral protein. We have previously reported that processing at the C-terminal hydrophobic domain of the core protein leads to efficient polyubiquitylation of the core protein (52). In this study, we identified E6AP as an HCV core-binding protein and showed that HCV core protein interacts with E6AP in vivo and in vitro, that E6AP enhances ubiquitylation and degradation of the mature core protein as well as the C-terminally truncated core protein, and that HCV core protein expressed from infectious HCV is

degraded via E6AP-dependent proteolysis. HCV core protein and E6AP were found to colocalize in the cytoplasm, especially in the perinuclear region. Moreover, exogenous expression of E6AP reduces intracellular core protein levels and supernatant HCV infectivity titers in HCV-infected Huh-7 cells. Knock-down of endogenous E6AP by siRNA increases intracellular core protein levels and supernatant infectivity titers in HCV-infected cells. These findings suggest that E6AP mediates ubiquitylation and degradation of HCV core protein, thereby affecting the production of HCV particles.

HCV core protein interacts with E6AP through the region of the core protein between aa 58 and aa 71. These 14 amino acids are highly conserved, with the first nine amino acids (PRGRRQPIP) present in the core protein of all the HCV genotypes (3). This result suggests that E6AP-dependent degradation of HCV core protein is common to all HCV genotypes and plays an important role in the HCV life cycle or viral pathogenesis. Our data indicated that HCV core proteins of genotypes 1b and 2a are subjected to proteolysis through an E6AP-mediated degradation pathway. We are currently examining whether E6AP promotes degradation of HCV core proteins of other genotypes.

Studies in addition to ours have reported that other HCV proteins, such as NS5B (8), the unglycosylated cytosolic form of E2 (39), NS2 (7), and F protein (58), are degraded through the ubiquitin-proteasome pathway. These studies suggest that the ubiquitin-proteasome pathway plays a role in the HCV life cycle or viral pathogenesis. To our knowledge, the present study is the first to demonstrate that the ubiquitin-proteasome pathway affects the HCV life cycle.

PA28 γ was found to interact with HCV core protein in hepatocytes and promote proteasomal degradation of HCV core protein (30). PA28 γ , however, has been shown to function




FIG. 9. Exogenous expression of E6AP reduces intracellular HCV core protein levels and supernatant infectivity titers in HCV-infected Huh-7 cells. Naïve Huh-7 cells were seeded as described in Materials and Methods; inoculated with 2.5 ml of the inoculum including infectious HCV JFH1 (6.5×10^3 TCID₅₀/ml); and transfected with 6 μg of empty plasmid, pCAG-HA-E6AP, or pCAG-HA-E6AP C-A. The culture supernatant and the cells were collected at days 3 and 7 postinfection. (A) Intracellular HCV core protein levels. (B) Levels of total protein. (C) Levels of intracellular HCV RNA in HCV-infected Huh-7 cells. Data represent the averages of three experiments with error bars. (D) Supernatant infectivity titers. At day 7 postinfection, culture supernatants were collected and assayed for TCID₅₀ determinations. The difference between empty vector and E6AP or between E6AP and E6AP C-A was significant (*, $P < 0.05$, Student's *t* test). (E and F) HCV JFH1-infected Huh-7 cells were transfected with either MEF-E6AP plasmid or MEF-E6AP C-A plasmid, grown on coverslips, fixed, and processed for double-label immunofluorescence for HCV core and MEF-E6AP (E) or MEF-E6AP C-A (F). Anticore MAb (2H9) and anti-FLAG PAb were used as primary antibodies. Nuclei were visualized by staining the cells with DAPI. All the samples were examined with a BZ-8000 microscope. Representative images of individual cells are shown with merge images. emp, empty vector.




FIG. 10. E6AP silencing leads to an increase in the level of intracellular HCV core protein and supernatant infectivity titer in HCV-infected Huh-7 cells. (A) HCV JFH1-infected cells were replated in a six-well plate at 3×10^5 cells/well and transfected with 40 pmol of E6AP siRNA or control siRNA. The culture medium was changed at 24 h after transfection. The cells were harvested at day 2 after transfection, and the intracellular core protein levels were quantitated using the HCV core antigen ELISA. Equivalent amounts of the whole-cell lysates were separated by SDS-PAGE and analyzed by immunoblotting with anti-E6AP MAb or anti-GAPDH MAb. (B) Culture supernatants were collected at day 2 after transfection and assayed for TCID₅₀ determinations. For both panels, the difference between E6AP siRNA and control siRNA was significant (*, $P < 0.05$, Student's *t* test).

in a ubiquitin-independent, ATP-independent, and 20S proteasome-dependent pathway (27). There have been reports that several cellular factors, such as p53 (2), p73 (2), and RPN4 (18), are degraded through two alternative pathways, the ubiquitin-dependent 26S proteasome-dependent pathway and the ubiquitin-independent 20S proteasome-dependent pathway. Here we provide evidence that E6AP mediates ubiquitylation of HCV core protein. Still unclear is whether the PA28 γ -dependent pathway requires polyubiquitylation of HCV core protein. HCV core protein is predominantly localized in the cytoplasm, especially at the endoplasmic reticulum membrane, on the surface of lipid droplets, and on mitochondria and mitochondrion-associated membranes (51). In HCV JFH1-infected cells, HCV core was found to localize in the cytoplasm and frequently to accumulate in the perinuclear region and the lipid droplets (44). Our results indicated that E6AP colocalized with HCV core protein especially in the perinuclear region. PA28 γ was found to colocalize with HCV core protein in the nucleus. Functional differences may exist between the E6AP-dependent pathway and the PA28 γ -dependent pathway in the stability control of HCV core protein. The functional role of the E6AP-dependent pathway and the PA28 γ -dependent pathway remains to be elucidated.

The HCV core-binding region of E6AP was mapped to the region between aa 418 and aa 517. The multicopy maintenance protein 7, Mcm7, interacts with E6AP through a short motif,

termed the L2G box (aa 412 to 414), that lies within the E6 binding site of E6AP (23). Our data indicated that the E6 binding region containing the L2G motif is not required for interaction between HCV core protein and E6AP (Fig. 2C, lane M).

We propose here that E6AP may affect the production of HCV particles through controlling the amounts of HCV core protein. This mechanism may contribute to persistent infection. The E6AP binding domain of the core protein resides in the RNA-binding domain and binding domains for many host factors (40). These factors may affect the binding between E6AP and HCV core protein, resulting in control of E6AP-dependent core degradation. Another possibility is that HCV core protein may affect the normal function of E6AP, thereby contributing to pathogenesis. It will be intriguing to investigate whether HCV core protein has any effect on E6AP-dependent degradation of host factors. The other intriguing possibility is that HCV core-E6AP complex may function as an E3 ligase-like E6-E6AP complex to target host factors for proteasomal degradation and contribute to viral pathogenesis.

In conclusion, we have demonstrated that E6AP interacts with HCV core protein in vitro and in vivo and mediates ubiquitin-dependent degradation of the core protein, leading to downregulation of HCV particles. We propose that the E6AP-mediated ubiquitin-proteasome pathway may play a role in affecting the production of HCV particles through controlling the amounts of viral nucleocapsid protein. Identification of the specific E3 ubiquitin ligase may contribute to gaining a better understanding of the biology of the HCV life cycle as well as molecular details of the ubiquitin-dependent degradation of HCV core protein.

ACKNOWLEDGMENTS

We thank D. Bohmann (EMBL) for providing pMT123, K. Miyazono (University of Tokyo) for pcDEF3-6Myc-WWP1, and K. Iwai (Osaka City University) for recombinant baculovirus carrying His 6-mouse E1. Huh-7.5.1 cells and Huh-7 cells were kindly provided by F. V. Chisari (Scripps Research Institute). We also thank P. Zhou (Weill Medical College of Cornell University), S. I. Wells (Cincinnati Children's Hospital Medical Center), and A. W. Hudson (Medical College of Wisconsin) for critical readings of the manuscript; M. Matsuda, S. Yoshizaki, M. Ikeda, and M. Sasaki for technical assistance; Y. Sugiyama and S. Senzui for plasmid construction; and T. Mizoguchi for secretarial work.

This work was supported in part by a grant for Research on Health Sciences focusing on Drug Innovation from the Japan Health Sciences Foundation; by grants-in-aid from the Ministry of Health, Labor and Welfare; by grants-in-aid from the Ministry of Education, Culture, Sports, Science and Technology; and by the program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO), Japan. T.I. was supported in part by a grant from Novartis Foundation (Japan) for the Promotion of Science and by the Tokyo Metropolitan University President's Fund, Special Emphasis Research Project of Japan.

REFERENCES

1. Aizaki, H., Y. Aoki, T. Harada, K. Ishii, T. Suzuki, S. Nagamori, G. Toda, Y. Matsuura, and T. Miyamura. 1998. Full-length complementary DNA of hepatitis C virus genome from an infectious blood sample. *Hepatology* 27: 621-627.
2. Asher, G., P. Tsvetkov, C. Kahana, and Y. Shaul. 2005. A mechanism of ubiquitin-independent proteasomal degradation of the tumor suppressors p53 and p73. *Genes Dev.* 19:316-321.
3. Bukh, J., R. H. Purcell, and R. H. Miller. 1994. Sequence analysis of the core gene of 14 hepatitis C virus genotypes. *Proc. Natl. Acad. Sci. USA* 91:8239-8243.

4. Chen, C., and H. Okayama. 1987. High-efficiency transformation of mammalian cells by plasmid DNA. *Mol. Cell. Biol.* 7:2745-2752.
5. Choo, Q. L., G. Kuo, A. J. Weiner, L. R. Overby, D. W. Bradley, and M. Houghton. 1989. Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 244:359-362.
6. Choo, Q. L., K. H. Richman, J. H. Han, K. Berger, C. Lee, C. Dong, C. Gallegos, D. Coit, R. Medina-Selby, P. J. Barr, et al. 1991. Genetic organization and diversity of the hepatitis C virus. *Proc. Natl. Acad. Sci. USA* 88:2451-2455.
7. Franck, N., J. Le Seyec, C. Guguen-Guillouze, and L. Erdtmann. 2005. Hepatitis C virus NS2 protein is phosphorylated by the protein kinase CK2 and targeted for degradation to the proteasome. *J. Virol.* 79:2700-2708.
8. Gao, L., H. Tu, S. T. Shi, K. J. Lee, M. Asanaka, S. B. Hwang, and M. M. Lai. 2003. Interaction with a ubiquitin-like protein enhances the ubiquitination and degradation of hepatitis C virus RNA-dependent RNA polymerase. *J. Virol.* 77:4149-4159.
9. Giannini, C., and C. Brechot. 2003. Hepatitis C virus biology. *Cell Death Differ.* 10(Suppl. 1):S27-S38.
10. Grakoui, A., D. W. McCourt, C. Wychowski, S. M. Feinstone, and C. M. Rice. 1993. Characterization of the hepatitis C virus-encoded serine proteinase: determination of proteinase-dependent polyprotein cleavage sites. *J. Virol.* 67:2832-2843.
11. Harris, K. F., I. Shoji, E. M. Cooper, S. Kumar, H. Oda, and P. M. Howley. 1999. Ubiquitin-mediated degradation of active Src tyrosine kinase. *Proc. Natl. Acad. Sci. USA* 96:13738-13743.
12. Hijikata, M., H. Mizushima, T. Akagi, S. Mori, N. Kakiuchi, N. Kato, T. Tanaka, K. Kimura, and K. Shimotohno. 1993. Two distinct proteinase activities required for the processing of a putative nonstructural precursor protein of hepatitis C virus. *J. Virol.* 67:4665-4675.
13. Huijbregtse, J. M., M. Scheffner, S. Beaudenon, and P. M. Howley. 1995. A family of proteins structurally and functionally related to the E6-AP ubiquitin-protein ligase. *Proc. Natl. Acad. Sci. USA* 92:2563-2567.
14. Huijbregtse, J. M., M. Scheffner, and P. M. Howley. 1993. Cloning and expression of the cDNA for E6-AP, a protein that mediates the interaction of the human papillomavirus E6 oncoprotein with p53. *Mol. Cell. Biol.* 13:775-784.
15. Hussy, P., H. Langen, J. Mous, and H. Jacobsen. 1996. Hepatitis C virus core protein: carboxy-terminal boundaries of two processed species suggest cleavage by a signal peptide peptidase. *Virology* 224:93-104.
16. Ichimura, T., H. Yamamura, K. Sasamoto, Y. Tominaga, M. Taoka, K. Kakiuchi, T. Shinkawa, N. Takahashi, S. Shimada, and T. Isobe. 2005. 14-3-3 proteins modulate the expression of epithelial Na⁺ channels by phosphorylation-dependent interaction with Nedd4-2 ubiquitin ligase. *J. Biol. Chem.* 280:13187-13194.
17. Iwai, K., K. Yamanaka, T. Kamura, N. Minato, R. C. Conaway, J. W. Conaway, R. D. Klausner, and A. Pause. 1999. Identification of the von Hippel-Lindau tumor-suppressor protein as part of an active E3 ubiquitin ligase complex. *Proc. Natl. Acad. Sci. USA* 96:12436-12441.
18. Ju, D., and Y. Xie. 2004. Proteasomal degradation of RPN4 via two distinct mechanisms, ubiquitin-dependent and -independent. *J. Biol. Chem.* 279:23851-23854.
19. Kao, W. H., S. L. Beaudenon, A. L. Talis, J. M. Huijbregtse, and P. M. Howley. 2000. Human papillomavirus type 16 E6 induces self-ubiquitination of the E6AP ubiquitin-protein ligase. *J. Virol.* 74:6408-6417.
20. Kato, T., M. Miyamoto, A. Furusaka, T. Date, K. Yasui, J. Kato, S. Matsushima, T. Komatsu, and T. Wakita. 2003. Processing of hepatitis C virus core protein is regulated by its C-terminal sequence. *J. Med. Virol.* 69:357-366.
21. Kishino, T., M. Lalaude, and J. Wagstaff. 1997. UBE3A/E6-AP mutations cause Angelman syndrome. *Nat. Genet.* 15:70-73.
22. Komuro, A., T. Imamura, M. Saitoh, Y. Yoshida, T. Yamori, K. Miyazono, and K. Miyazawa. 2004. Negative regulation of transforming growth factor-beta (TGF-beta) signaling by WW domain-containing protein 1 (WWP1). *Oncogene* 23:6914-6923.
23. Kuhne, C., and L. Banks. 1998. E3-ubiquitin ligase/E6-AP links multicopy maintenance protein 7 to the ubiquitination pathway by a novel motif, the L2G box. *J. Biol. Chem.* 273:34302-34309.
24. Kumar, S., A. L. Talis, and P. M. Howley. 1999. Identification of HHR23A as a substrate for E6-associated protein-mediated ubiquitination. *J. Biol. Chem.* 274:18785-18792.
25. Kunkel, M., M. Lorincz, R. Rijnbrand, S. M. Lemon, and S. J. Watowich. 2001. Self-assembly of nucleocapsid-like particles from recombinant hepatitis C virus core protein. *J. Virol.* 75:2119-2129.
26. Kuo, G., Q. L. Choo, H. J. Alter, G. L. Gitnick, A. G. Redeker, R. H. Purcell, T. Miyamura, J. L. Dienstag, M. J. Alter, C. E. Stevens, et al. 1989. An assay for circulating antibodies to a major etiologic virus of human non-A, non-B hepatitis. *Science* 244:362-364.
27. Li, X., D. M. Lonard, S. Y. Jung, A. Malovannaya, Q. Feng, J. Qin, S. Y. Tsai, M. J. Tsai, and B. W. O'Malley. 2006. The SRC-3/AIB1 coactivator is degraded in a ubiquitin- and ATP-independent manner by the REGγ proteasome. *Cell* 124:381-392.
28. Lindenbach, B. D., M. J. Evans, A. J. Syder, B. Wolk, T. L. Tellinghuisen, C. C. Liu, T. Maruyama, R. O. Hynes, D. R. Burton, J. A. McKeating, and C. M. Rice. 2005. Complete replication of hepatitis C virus in cell culture. *Science* 309:623-626.
29. McLauchlan, J., M. K. Lemberg, G. Hope, and B. Martoglio. 2002. Intramembrane proteolysis promotes trafficking of hepatitis C virus core protein to lipid droplets. *EMBO J.* 21:3980-3988.
30. Moriishi, K., T. Okabayashi, K. Nakai, K. Moriya, K. Koike, S. Murata, T. Chiba, K. Tanaka, R. Suzuki, T. Suzuki, T. Miyamura, and Y. Matsuura. 2003. Proteasome activator PA28γ-dependent nuclear retention and degradation of hepatitis C virus core protein. *J. Virol.* 77:10237-10249.
31. Moriya, K., H. Fujie, Y. Shintani, H. Yotsuyanagi, T. Tsutsumi, K. Ishibashi, Y. Matsuura, S. Kimura, T. Miyamura, and K. Koike. 1998. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat. Med.* 4:1065-1067.
32. Moriya, K., H. Yotsuyanagi, Y. Shintani, H. Fujie, K. Ishibashi, Y. Matsuura, T. Miyamura, and K. Koike. 1997. Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J. Gen. Virol.* 78:1527-1531.
33. Natsume, T., Y. Yamauchi, H. Nakayama, T. Shinkawa, M. Yanagida, N. Takahashi, and T. Isobe. 2002. A direct nanoflow liquid chromatography-tandem mass spectrometry system for interaction proteomics. *Anal. Chem.* 74:4725-4733.
34. Niwa, H., K. Yamamura, and J. Miyazaki. 1991. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* 108:193-199.
35. Oda, H., S. Kumar, and P. M. Howley. 1999. Regulation of the Src family tyrosine kinase Blk through E6AP-mediated ubiquitination. *Proc. Natl. Acad. Sci. USA* 96:9557-9562.
36. Ogino, T., H. Fukuda, S. Imajoh-Ohmi, M. Kohara, and A. Nomoto. 2004. Membrane binding properties and terminal residues of the mature hepatitis C virus capsid protein in insect cells. *J. Virol.* 78:11766-11777.
37. Okamoto, K., K. Moriishi, T. Miyamura, and Y. Matsuura. 2004. Intramembrane proteolysis and endoplasmic reticulum retention of hepatitis C virus core protein. *J. Virol.* 78:6370-6380.
38. Owsianka, A. M., and A. H. Patel. 1999. Hepatitis C virus core protein interacts with a human DEAD box protein DDX3. *Virology* 257:330-340.
39. Pavo, N., D. R. Taylor, and M. M. Lai. 2002. Detection of a novel unglycosylated form of hepatitis C virus E2 envelope protein that is located in the cytosol and interacts with PKR. *J. Virol.* 76:1265-1272.
40. Polyak, S. J., K. C. Klein, I. Shoji, T. Miyamura, and J. R. Lingappa. 2006. Assemble and interact: pleiotropic functions of the HCV core protein, p. 89-119. *In* S.-L. Tan (ed.), *Hepatitis C viruses: genomes and molecular biology*. Horizon Bioscience, Norwich, United Kingdom.
41. Poynard, T., M. F. Yuen, V. Ratziu, and C. L. Lai. 2003. Viral hepatitis C. *Lancet* 362:2095-2100.
42. Ravaggi, A., G. Natoli, D. Primi, A. Albertini, M. Levrero, and E. Cariani. 1994. Intracellular localization of full-length and truncated hepatitis C virus core protein expressed in mammalian cells. *J. Hepatol.* 20:833-836.
43. Ray, R. B., L. M. Lagging, K. Meyer, and R. Ray. 1996. Hepatitis C virus core protein cooperates with *ras* and transforms primary rat embryo fibroblasts to tumorigenic phenotype. *J. Virol.* 70:4438-4443.
44. Rouille, Y., F. Helle, D. Delgrange, P. Roingeard, C. Voisset, E. Blanchard, S. Belouard, J. McKeating, A. H. Patel, G. Maertens, T. Wakita, C. Wychowski, and J. Dubuisson. 2006. Subcellular localization of hepatitis C virus structural proteins in a cell culture system that efficiently replicates the virus. *J. Virol.* 80:2832-2841.
45. Saito, I., T. Miyamura, A. Ohbayashi, H. Harada, T. Katayama, S. Kikuchi, Y. Watanabe, S. Koi, M. Onji, Y. Ohta, et al. 1990. Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc. Natl. Acad. Sci. USA* 87:6547-6549.
46. Santolini, E., G. Migliaccio, and N. La Monica. 1994. Biosynthesis and biochemical properties of the hepatitis C virus core protein. *J. Virol.* 68:3631-3641.
47. Sato, S., M. Fukasawa, Y. Yamakawa, T. Natsume, T. Suzuki, I. Shoji, H. Aizaki, T. Miyamura, and M. Nishijima. 2006. Proteomic profiling of lipid droplet proteins in hepatoma cell lines expressing hepatitis C virus core protein. *J. Biochem. (Tokyo)* 139:921-930.
48. Scheffner, M., J. M. Huijbregtse, and P. M. Howley. 1994. Identification of a human ubiquitin-conjugating enzyme that mediates the E6-AP-dependent ubiquitination of p53. *Proc. Natl. Acad. Sci. USA* 91:8797-8801.
49. Scheffner, M., J. M. Huijbregtse, R. D. Vierstra, and P. M. Howley. 1993. The HPV-16 E6 and E6-AP complex functions as a ubiquitin-protein ligase in the ubiquitination of p53. *Cell* 75:495-505.
50. Scheffner, M., U. Nuber, and J. M. Huijbregtse. 1995. Protein ubiquitination involving an E1-E2-E3 enzyme ubiquitin thioester cascade. *Nature* 373:81-83.
51. Suzuki, R., S. Sakamoto, T. Tsutsumi, A. Rikimaru, K. Tanaka, T. Shimoike, K. Moriishi, T. Iwasaki, K. Mizumoto, Y. Matsuura, T. Miyamura, and T. Suzuki. 2005. Molecular determinants for subcellular localization of hepatitis C virus core protein. *J. Virol.* 79:1271-1281.
52. Suzuki, R., K. Tamura, J. Li, K. Ishii, Y. Matsuura, T. Miyamura, and T. Suzuki. 2001. Ubiquitin-mediated degradation of hepatitis C virus core pro-

- tein is regulated by processing at its carboxyl terminus. *Virology* 280:301-309.
53. Suzuki, T., K. Omata, T. Satoh, T. Miyasaka, C. Arai, M. Maeda, T. Matsuno, and T. Miyamura. 2005. Quantitative detection of hepatitis C virus (HCV) RNA in saliva and gingival crevicular fluid of HCV-infected patients. *J. Clin. Microbiol.* 43:4413-4417.
54. Takamizawa, A., C. Mori, I. Fuke, S. Manabe, S. Murakami, J. Fujita, E. Onishi, T. Andoh, I. Yoshida, and H. Okayama. 1991. Structure and organization of the hepatitis C virus genome isolated from human carriers. *J. Virol.* 65:1105-1113.
55. Talis, A. L., J. M. Huibregtse, and P. M. Howley. 1998. The role of E6AP in the regulation of p53 protein levels in human papillomavirus (HPV)-positive and HPV-negative cells. *J. Biol. Chem.* 273:6439-6445.
56. Wakita, T., T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H. G. Krausslich, M. Mizokami, R. Bartenschlager, and T. J. Liang. 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* 11:791-796.
57. Wertz, I. E., K. M. O'Rourke, Z. Zhang, D. Dornan, D. Arnott, R. J. Deshaies, and V. M. Dixit. 2004. Human de-etioloated-1 regulates c-Jun by assembling a CUL4A ubiquitin ligase. *Science* 303:1371-1374.
58. Xu, Z., J. Choi, W. Lu, and J. Ou. 2003. Hepatitis C virus F protein is a short-lived protein associated with the endoplasmic reticulum. *J. Virol.* 77:1578-1583.
59. Yamaguchi, R., S. Momosaki, G. Gao, C. C. Hsia, M. Kojiro, C. Scudamore, and E. Tabor. 2004. Truncated hepatitis C virus core protein encoded in hepatocellular carcinomas. *Int. J. Mol. Med.* 14:1097-1100.
60. Yasui, K., T. Wakita, K. Tsukiyama-Kohara, S. I. Funahashi, M. Ichikawa, T. Kajita, D. Moradpour, J. R. Wands, and M. Kohara. 1998. The native form and maturation process of hepatitis C virus core protein. *J. Virol.* 72:6048-6055.
61. Zhong, J., P. Gastaminza, G. Cheng, S. Kapadia, T. Kato, D. R. Burton, S. F. Wieland, S. L. Uprichard, T. Wakita, and F. V. Chisari. 2005. Robust hepatitis C virus infection in vitro. *Proc. Natl. Acad. Sci. USA* 102:9294-9299.

Available online at www.sciencedirect.com

Journal of Virological Methods 148 (2008) 174–181

www.elsevier.com/locate/jviromet

Dynamic behavior of hepatitis C virus quasispecies in a long-term culture of the three-dimensional radial-flow bioreactor system

Kyoko Murakami^a, Yasushi Inoue^{a,b}, Su-Su Hmwe^{a,c}, Kazuhiko Omata^{a,d}, Tomokatsu Hongo^e,
Koji Ishii^a, Sayaka Yoshizaki^a, Hideki Aizaki^a, Tomokazu Matsuura^f, Ikuo Shoji^a,
Tatsuo Miyamura^a, Tetsuro Suzuki^{a,*}

^a Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan

^b Pulmonary and Critical Care Unit, Mita Hospital, International University of Health and Welfare, Japan

^c Department of Infectious Diseases, Internal Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

^d Department of Oral and Maxillofacial Surgery, The Nippon Dental University School of Dentistry at Tokyo, Tokyo, Japan

^e ABLE Corporation, Shizuoka, Japan

^f Department of Laboratory medicine, The Jikei University School of Medicine, Tokyo, Japan

Received 25 July 2007; received in revised form 9 November 2007; accepted 21 November 2007

Abstract

Hepatitis C virus (HCV) exists in infected individuals as quasispecies, usually consisting of a dominant viral isolate and a variable mixture of related, yet genetically distinct, variants. A prior HCV infection system was developed using human hepatocellular carcinoma cells cultured in the three-dimensional radial-flow bioreactor (RFB), in which the cells retain morphological appearance and their differentiated hepatocyte functions for an extended period of time. This report studies the selection and alteration of the viral quasispecies in the RFB system inoculated with pooled serum derived from HCV carriers. Monitoring the viral RNA and core protein in the culture supernatants, together with nucleotide sequencing of hypervariable region 1 of the HCV genome, demonstrated that (1) the virus production intermittently fluctuated in the cultures, (2) the viral genetic diversity was markedly reduced 3 days post-infection (p.i.), and (3) dominant species changed on days 19–33 p.i., suggesting that the virus populations can be selected according to susceptibility to the viral infection and replication. A therapeutic effect of interferon- α also demonstrated the inhibition of HCV expression. Thus, this HCV infection model in the RFB system should be useful for investigating the dynamic behavior of HCV quasispecies in cultured cells and evaluating anti-HCV compounds.

© 2007 Elsevier B.V. All rights reserved.

Keywords: Hepatitis C virus; Three-dimensional culture; Radial-flow bioreactor; Dynamics; Quasispecies

1. Introduction

Hepatitis C virus (HCV) is a major cause of chronic liver diseases (Choo et al., 1989; Kuo et al., 1989; Saito et al., 1990) and has been estimated to infect more than 170 million people throughout the world (Poynard et al., 2003). Symptoms of persistent HCV infection extend from chronic hepatitis to cirrhosis and ultimately hepatocellular carcinoma (Choo et al., 1989; Kuo et al., 1989; Saito et al., 1990). HCV belongs to the genus *Hepacivirus*, included in the family of Flaviviridae, and possesses a viral genome of a single, positive-stranded RNA with

a nucleotide (nt) length of approximately 9.6 kb (Choo et al., 1991; Grakoui et al., 1993; Hijikata et al., 1991). It has been shown that HCV, like many other RNA viruses, circulates within infected individuals as a diverse population and closely related variants are referred to as quasispecies (Martell et al., 1992). This quasispecies model of mixed virus populations may imply a significant survival advantage because the simultaneous presence of multiple variant genomes and/or high rate of generation of new variants allow rapid selection of the mutants are better suited to new environmental conditions (Pawlotsky, 2006).

Studies on HCV replication and development of selective antiviral drugs have been hampered primarily by the lack of efficient cell culture systems. Establishment of selectable dicistronic HCV RNAs that are capable of autonomous replication to high levels in human hepatoma Huh-7 cells was a

* Corresponding author. Tel.: +81 3 5285 1111; fax: +81 3 5285 1161.

E-mail address: tesuzuki@nih.go.jp (T. Suzuki).

significant breakthrough in HCV research; however, virus production has not been observed in the conventional monolayer cultures (Blight et al., 2000; Lohmann et al., 1999). Recently, it has been described that infectious HCV particles are efficiently produced from a genotype 2a isolate JFH-1 in Huh-7 cells (Blight et al., 2000; Wakita et al., 2005; Zhong et al., 2005). This JFH-1 based HCV culture system is an invaluable achievement permitting a variety of studies on the complete HCV life cycle. However, HCV infection systems with human sera or plasmas containing intact virions are still limited because of low levels of propagation in the cultures. Reverse transcription (RT)-PCR was typically used to detect the viral RNA in cell extracts; however, synthesized viral proteins were not observed in these systems (Ikeda et al., 1998; Tagawa et al., 1995).

There are reports of differentiated human hepatoma FLC4 (functional liver cell 4) cells grown in a three-dimensional (3D) radial-flow bioreactor (RFB) that can be infected by HCV-positive serum and support viral replication (Aizaki et al., 2003). Furthermore, production and release of infectious HCV has been observed in the RFB system following transfection of FLC4 cells with *in vitro* transcribed HCV genomic RNA, as well as in a 3D system using Huh-7 cells harboring genome-length dicistronic RNAs (Murakami et al., 2006). The RFB system, in which the bioreactor column consists of a cylindrical matrix with porous bead microcarriers extended vertically, was aimed initially at developing artificial liver tissues and allows liver-derived cells to maintain morphological appearance as well as their physiological functions, such as the ability to synthesize albumin and drug-metabolizing activity mediated by cytochrome P450 (Iwahori et al., 2003). The radial-flow configuration permits full contact between culture medium and cells at a physiologic perfusion flow rate, and prevents excessive shear stresses and buildup of waste products, thus ensuring the long-term viability of 3D cell culture.

The aim of the present study was to characterize HCV dynamics in the RFB system during long-term cultures inoculated with pooled serum obtained from HCV carriers, and to examine the therapeutic effects of interferon-alpha (IFN- α) in this HCV infection model.

2. Materials and methods

2.1. Cell cultures

FLC4 cells (Aoki et al., 1998), which were derived from human hepatocellular carcinoma cells and negative for HCV RNA and HBV DNA, were maintained in serum-free ASF104 medium (Ajinomoto, Japan) supplemented with 4 g/L D-glucose on the collagen-coated dishes before inoculating into the RFB column. The RFB system (ABLE, Japan) was manipulated as described previously (Aizaki et al., 2003) with minor modifications. Briefly, RFB columns, which have bed volumes of 30 or 4 mL and are filled with porous glass microcarriers (diameter 0.6 mm, vacant capacity 50%, pore size <120 μ m) (Hongo et al., 2005), were seeded with FLC4 cells, which subsequently attached to the surface and inside of porous glass beads. ASF104 medium containing 2% fetal calf serum was added at a flow rate

of 50 mL/day, and the culture condition was automatically controlled by monitoring temperature, pH value and oxygen levels in the vessel throughout the duration of the study.

2.2. Infection of HCV-positive sera

HCV antibody-positive sera used in this study were blood donor samples supplied by The Japanese Red Cross Center, Tokyo, Japan. HCV RNA loads in the sera were as follows: serum A, 2.4×10^6 copies/mL; serum B, 8.6×10^6 copies/mL; serum C, 5.9×10^6 copies/mL; serum D, 2.5×10^6 copies/mL; serum E, 1.0×10^7 copies/mL; serum F, 1.4×10^7 copies/mL (Table 1). In the first experiment (Fig. 3), aliquots of each serum containing 2×10^6 copies of HCV RNA were mixed and pooled serum sample with 1.2×10^7 copies was prepared as an inoculum. The pooled serum (2.5 mL) was added to the 3D cultured-FLC4 cells in the 30-mL RFB column, and the culture medium was changed after 12 h of incubation. At various times during the culture period, culture medium (50 mL) was collected to determine HCV RNA and the core protein. Collected culture media were passed through a 0.20- μ m filter to remove the debris, and stored at -80°C . In the second experiment to evaluate a therapeutic effect of anti-HCV drug (Fig. 4), 4-mL RFB columns were used. IFN- α (Sumiferon 300; Sumitomo Pharmaceuticals, Japan) was added to one of two columns at a final concentration of 100 IU/mL after the infection. Culture medium was periodically collected for determination of HCV RNA, the core protein and transaminases, and was replaced with the same volume of fresh medium with or without IFN- α .

2.3. Quantitation of HCV RNA and core protein

HCV RNA was extracted from 140 μ L of each serum or culture medium using QIAamp Viral RNA Mini spin column (QIAGEN); RNA was eluted in 60 μ L of water and stored at -80°C . Real-time RT-PCR was performed using TaqMan EZ RT-PCR Core Reagents (PE Applied Biosystems), as described previously (Aizaki et al., 2003; Suzuki et al., 2005). The viral core antigen in the culture medium was quantified by immunoassay (Ortho HCV-Core ELISA Kit; Ortho-Clinical Diagnostics), according to the manufacturer's instruction (Murakami et al., 2006).

2.4. PCR amplification and nucleotide sequencing of HVR1 domain and its flanking region

Five microliters of RNA samples prepared as above were reverse transcribed using SuperScript II (Invitrogen) and a specific primer 5'-CATCCATGTGCAGCCGAACC-3' (corresponding to nucleotides [nt] 2006–1987 of HCV NIHJ1) (Aizaki et al., 1998). For the nested PCR, a genotype-independent set of primers specific for hypervariable region 1 (HVR1). The first round of PCR was performed with the outer sense primer 5'-GCATGGCTTGGGATATGATG-3' (nt 1291–1310) and with the reverse transcription primer described above as the outer antisense primer. After the initial 3.5-min denaturation step at 94°C , 35 PCR cycles, with each cycle

Table 1
HCV-positive sera used in this study

Serum	Clone	HCV HVR1 sequence	% in the serum	genotype
A	A1	KVLI VMLS FAGVDGSTRITIGGRTAHTTQGSASLFS SGPAQKIQLINTNGS	75	1
	A2	-----L-----N-H-V--AV-SS---FT---KL-----S---	12.5	
	A3	-----L-----N-YAS---AGLL-R-V--I-TA-----S---	12.5	
B	B1	KVVV ILLL AAGVDAGTNTIGGSAAQTTS GFTGLFRSGARQNIQLINTNGS	50	2
	B2	-----R	12.5	
	B3	-----S-----	12.5	
	B4	--L-V--F-----E-HVT--N-GR--A-LV--LTP--K-----	12.5	
	B5	--I-----	12.5	
C	C1	KVLI VMLL FAGVDGDTHVSGGTQGRAAYGLASLFALGPTQKIQLVNTNGS	83.3	1
	C2	-----A-----	16.7	
D	D1	KVLI VMLL FAGVDGVTHTSGAAAGHNARSLGSLFSLGSAQKQLINTNGS	40	1
	D2	-----A-Y--GT--Y-TRTFT-F--R-PS--I-----	20	
	D3	-----T--Y--T-T---P-----V-----	10	
	D4	-----V---T---P-----V-----	10	
	D5	-----V-----	10	
	D6	-----Y-T--FT---S---I--V-----	10	
E	E1	KVLI VMLL FAGVDGSTRVSGGQAGRVTKSLASFFS PGPQKIQLVNSNGS	40	1
	E2	-----HGFT-L--A-S-----	30	
	E3	-----QGFT-L--A-S-----	10	
	E4	-----S-FT-L-TV-----	10	
	E5	-----N-Y-----AH--T-L--A-S-----	10	
F	F1	KVLI VMLL FAGVDGETNVMGGGRAGHTTNTFTSLFSLVGPQKIQLVNSNGS	37	1
	F2	-----D-K-----S-L--N--S-----	27	
	F3	-----K---Q---S-L--N--S-----	18	
	F4	-----A-----A-TK-----D---	9	
	F5	-----G-----A--A--L---TR--S-----	9	

consisting of 1 min at 94 °C, 2 min at 45 °C, and 3 min at 72 °C, were carried out, followed by a 10-min extension step at 72 °C. The second round was performed with the inner sense primer 5'-GGTAAGCTTCCATGGTGGGGAAGTGGGC-3' (nt 1419–1447) and the inner antisense primer 5'-CTGGAATTCGCAGTCCTGTTGATGTGCCA-3' (nt 1627–1599). The amplified products were cloned into the pGEM-T vector (Promega), and at least 8 independent clones were sequenced with an automatic DNA sequencer (ABI PRISM 310, PE Applied Biosystems).

3. Results

3.1. The outline of the RFB system

The RFB system was initially aimed at developing artificial liver tissues and allows liver-derived cells to maintain morphological appearance as well as their physiological functions, such as the ability to synthesize albumin and drug-metabolizing activity mediated by cytochrome P450 (Iwahori et al., 2003). Fig. 1 shows the outline of the RFB system. The bioreactor column consists of a vertically extended cylindrical matrix with porous glass microcarriers, which were most suitable for FLC4 culture as described in Section 2. The conditioning vessel is connected to a circulation system including tanks either for supplying fresh medium or for recovering sample aliquots. Oxygen consump-

tion, temperature and pH of the culture medium are monitored continuously and conditioned in the vessel by computer and mass flow controller throughout the culture. Thus, the radial-flow configuration permits full contact between culture medium and cells at a physiologic perfusion flow rate, and prevents excessive shear stresses and a buildup of waste products, thus ensuring the long-term viability of 3D culture. For the long-term culture up to 110 days, temperature in the vessel gradually decreased from 37 to 30 °C as shown in Fig. 2A. The oxygen consumption, which indicates the cell growth condition, increased slowly from days 0 to 80 post-inoculation of the cells, and maintained a constant level afterwards. Under this condition, the production rate of albumin was found to be stable from days 15 to 105. The following experiments of HCV infection were done in such a stable phase of the cell condition after 3 weeks of pre-culture. Cell grown in the RFB column reached confluence at the end of culture (day 110) since the cells were observed outside the matrix bed (Fig. 2B).

3.2. Infection of HCV-positive sera to RFB cultured FLC4 cells

Previously, HCV RNA could be detected in FLC4 cells grown in the RFB up to 4 weeks of culture following inoculation with an HCV carrier plasmid (Aizaki et al., 2003). Establishment of a long-term stable culture system of human liver-derived cells




Fig. 1. Outline of the RFB system. RFB system consists of vessel, column and PC monitoring system. Culture condition was automatically controlled: oxygen concentration, temperature, pH, and oxygen level in the conditioning vessel are continuously monitored by PC and conditioned by mass flow controller.

retaining their differentiated hepatocyte function, as described above, enables evaluations of dynamic analysis of HCV replication and selection of viral variability and quasispecies. The potential of this culture system for screening HCV-positive sera was well suited for the viral infection.

Table 1 shows the serum samples (A–F) from six HCV carriers. The nucleotide complexity of HCV in serum samples was determined by sequencing the 1449–1598 nt region of the HCV genome, which includes HVR1 located at the N-terminal region of E2. Each serum was a mixture of a dominant HCV clone and related but distinct viral populations. The dominant species in

sera A, C, D, E, and F were found to be genotype 1, and that in serum B was genotype 2. Viral loads in A–F, respectively, were 2.4×10^6 , 8.6×10^6 , 5.9×10^6 , 2.5×10^6 , 1.0×10^7 and 1.4×10^7 copies/mL, which were determined by real-time RT-PCR, as previously described (Aizaki et al., 2003; Suzuki et al., 2005). HCV loads of 2×10^6 copies from each serum sample were mixed to prepare a pooled serum sample containing 1.2×10^7 copies of HCV RNA. After FLC4 cells were inoculated into the RFB and subjected to 2 weeks of pre-culture for the preparation of 3D culture, the cells were infected with the pooled serum. Cell number at infection was about 10^8 in the 30-




Fig. 2. Long-term culture of FLC4 cells in the RFB system. (A) Long-term culture of FLC4 cells in the RFB system. Temperature (closed circles) was gradually decreased from 37 to 30 °C. Oxygen consumption (open circles) was gradually increased from days 0 to 80 and reached the steady-state level. Albumin concentration (closed triangles) was constant from days 15 to 105. (B) The appearance of the RFB column at the beginning (day 0) and at the end (day 110) of culture.




Fig. 3. HCV propagation in FLC4 cells cultured in the RFB system following inoculation with pooled sera obtained from HCV carriers. The 3D-cultured FLC4 cells were incubated with a pooled serum sample for 12 h, followed by changing the culture medium to fresh one. Culture medium was periodically collected for 42 days after inoculation, and HCV RNA and the viral core protein were quantified, respectively, by real-time RT-PCR and ELISA. (A) HCV RNA level in culture supernatant. (B) HCV-core protein (closed circles) and oxygen consumption (open triangles) levels in culture supernatant. (C) Changes in the viral quasispecies distribution after the inoculation. Percentages in the inoculum or in the culture medium at each time point (day 3, 9, 19, or 33 p.i.) are indicated at the right side. *, termination codon.

mL RFB column, as estimated from the glucose consumption (Kawada et al., 1998). Culture medium in the RFB was replaced with fresh medium 12 h post-infection (p.i.) and periodically sampled for 42 days.

Fig. 3A and B shows the levels of HCV RNA and viral core protein in the culture medium, respectively. HCV RNA was not observed on the first 2 days following infection, but was detectable from day 3 p.i. Viral RNA levels fluctuated, with peaks on days 3, 9, 19–21 and 33–36 p.i. At days 19–21 p.i., the average amount of HCV RNA detected in the culture supernatant was approximately 3×10^6 copies/day. Intermittent peaks were observed in HCV core protein levels in the culture supernatant, and the peak pattern of the core protein was largely consistent with that of viral RNA. During the infection experiment, the level of oxygen consumption was constant at approximately 12 ppm, thus suggesting that the desired conditions (constant or very gradually increasing cell number) were maintained.

3.3. Quasispecies analysis in RFB culture

The above results suggest that, although the environment was consistent in the pooled serum infection, there were periods in which the viruses actively replicated and released from the cells and periods in which they poorly replicated. The pooled serum used for the infection exhibited HCV populations had at least 26 distinct quasispecies (Table 1). To investigate whether the quasispecies distribution was altered due to infection, and whether HCV populations are selected during long-term culture in the RFB, total RNA was extracted from the culture supernatant samples collected on days 3, 9, 19 and 33 p.i., and the nucleotide sequence of the region containing HVR1 was deter-

mined, as described above. As shown in Fig. 3C, it is of interest that only two HCV species were detected in the sample at day 3 p.i.; the dominant clone C1-1, comprising approximately 70% of the viral population, and clone B4, comprising 30%. Although clone C1-1 was not detected in the sequence of the inoculum shown in Table 1, it was most similar to clone C1, a dominant clone in plasma C, among the HCV population observed in the inoculum; thus, it is possible that clone C1-1 is one of the minor species in serum C. Clone B4 was found to be derived from serum B. An almost identical HCV population was observed in the sample at day 9 p.i. In this sample, the dominant clone C1-1 and clone B4-1, which differs from clone B4 by only one amino acid, were detected. In contrast, more significant variation in quasispecies structure of the HCV species was observed in the sample at day 19 p.i. than that at day 9 p.i. With B4 as the dominant clone, the serum B-derived HCV species, clones B4 and B4-2, which differs from clone B4 by one amino acid, comprised 58% of the total population. Four types of HCV sequences derived from serum C were detected. Two of these (clones C1-3 and C1-4) contained lethal mutations. It was also found that the HCV species detected in the sample at day 33 p.i. included only two clones (clones B4 and B4-3), derived from serum B. The dominant clone, B4, was found to comprise 89% of the total population.

3.4. Potential use of the RFB system for evaluation of anti-HCV compounds

An experiment was carried out to determine whether this HCV infection experiment system was useful for the evaluation of anti-HCV drugs (Fig. 4). For this purpose, a small,




Fig. 4. A therapeutic effect of IFN in HCV infection model in the RFB cultures. HCV-infected FLC4 cells were treated with or without 100 IU/mL IFN- α . (A) Culture media were periodically collected, and HCV RNA levels were determined. Closed circles: without IFN treatment, open circles: treatment with IFN. (B) Changes in the viral quasispecies distribution in the cells without IFN treatment. Percentages in the inoculum or in the culture medium on day 32 p.i. are indicated at the right side. *, termination codon.

4-mL RFB column was adopted and a pair of RFB cultures infected with the HCV-positive pooled plasma (Table 1) was prepared. IFN- α was added to one culture at a final concentration of 100 IU/mL at 12 h p.i. No cytotoxicity was observed in FLC4 cells under these conditions (data not shown). Culture media from two cultures (12.5 mL each) were sampled periodically for 35 days and replaced by the same volume of fresh medium in the presence or absence of IFN- α . HCV RNA in the collected media was quantified by real-time RT-PCR, as described above. As shown in Fig. 4A, in the no-treatment culture, fluctuations in the viral RNA levels with the peaks on days 7, 18, and 32 p.i. ($1.5\text{--}5 \times 10^4$ copies/mL) were observed. However, while HCV RNA at $0.5\text{--}0.8 \times 10^4$ copies/mL was detected in the IFN-treated culture at days 5–11 p.i., no HCV RNA was detected at days 12–30 p.i. Serum levels of hepatic transaminases such as ALT and AST are known to be markers of liver damage. In the HCV-infection model with FLC4 cells cultured in RFB, the AST levels in the culture medium, which ranged from 5 to 10 IU/L without HCV infection, increased to 20–50 IU/L according to the viral infection (data not shown). Such increased AST levels were found to fall by the IFN treatment to lower than 10 IU/L at day 28 p.i. As reported previously, the ALT levels in the culture medium were constantly low; its levels were less than 10 IU/mL, with or without HCV infection (Aizaki et al., 2003). The viral nucleotide sequence in the no-treatment culture medium at day 32 p.i. was determined. It was found that serum B-derived clone B4 was dominant, and serum C-derived clone C1 was present as a minor clone (Fig. 4B); thus, the results corresponded well with those demonstrated in Fig. 3. An increase in viral RNA in the IFN-treated culture after day 32 p.i. was observed; although the degree of increase was only slight (Fig. 4A). It will be interesting to test whether HCV species grown in the IFN-treated culture is a variant resistant to IFN- α .

4. Discussion

At present an important limitation of the *in vitro* HCV infection system is that the only established culture system is based on genotype 2a, JFH-1 isolate, and Huh-7-derived cell lines. The development of alternate infection systems in which other HCV strains and host cells are available has been needed for the study of HCV dynamics and virus–host interactions, and for testing antivirals. This paper demonstrates that a long-term culture of the 3D RFB system is a useful tool for investigating HCV dynamics. The present results revealed that the viral quasispecies distribution altered in the HCV infection system in the RFB system. The change probably occurs in the following two-stage process. The first change was observed on day 3 p.i.; thus, it is possible that the HCV species were selected according to infectivity in FLC4 cells. It has been reported that HCV particle populations in chronic hepatitis C patients consist of low-density virions and higher-density immune complex forms (Hijikata et al., 1993; Kanto et al., 1994). Inoculation of cultured cells with HCV has demonstrated that the immune complex forms were less infective than the antibody-unbound virions (Shimizu et al., 1994). Therefore, another hypothesis may be that a large number of HCV populations in sera A, D, E, and F are immune complex forms; thus, these sera are less susceptible to the cells than sera B and C. The second change was observed on days 19–33 p.i. While the serum C-derived clone was dominant in the early stages after infection, the serum B-derived HCV clone became dominant over time. In the absence of immunological selection pressure, viral nucleotide mutations at random positions are accumulated during viral replication, and the newly generated variant species are selected principally, if not solely, based on the intrinsic replicative advantages or disadvantages that these mutations confer. Thus, these results suggest that the use of pooled serum sample allowed for screening of infectious materials compatible for the RFB culture.

Evaluation methods for anti-HCV drugs using monolayer culture systems with various culture cells, such as the replicon system and the JFH-1 based virion production system, have been reported (Bartenschlager et al., 2003; Blight et al., 2000; Boriskin et al., 2006; Lanford et al., 2003; Lindenbach et al., 2005; Lohmann et al., 1999; Wakita et al., 2005; Zhong et al., 2005). These methods utilize viral markers, such as HCV RNA and antigens, as indicators of treatment efficacy. However, the utility of long-term cell culture systems for anti-HCV drug evaluation based on infection with human sera is still limited. The use of a chimpanzee model, the only non-human host for HCV infection, is restricted due to several reasons such as problematic availability and ethical consideration. Given intensive efforts to reduce and replace animal testing in the course of development of new therapies worldwide, the RFB-based HCV infection model is a potential alternative to animal models such chimpanzee for assessing anti-HCV compounds. According to the studies with regards to mathematical modeling of HCV kinetics (Dahari et al., 2005; Dixit et al., 2004; Layden et al., 2003; Layden-Almer et al., 2006; Perelson et al., 2005), IFN therapy against HCV infection generally generates a biphasic decline in viral load; there is a rapid decrease in the serum HCV RNA level over the

first 1 day of treatment, followed by the second phase, which is slower than the first-phase viral decline. To date, there were no such observable viral kinetics in the IFN treatment under such experimental settings. Further detailed kinetic analyses of the use of varying doses of IFN and of very early time points to evaluate the antiviral effect are in progress.

In summary, by investigating the dynamics of HCV populations in the RFB culture system, it was demonstrated that HCV was intermittently detected in the culture supernatants of long-term culture, and that changes in viral quasispecies appear to be related to this fluctuation in the virus level. It was also shown that an HCV-infection model using the RFB system is useful for evaluating potential antivirals. Further investigation on the infection and growth of various HCV-positive sera is currently being conducted in order to obtain an adaptive clone with higher replication efficiency in this culture system.

Acknowledgements

The authors thank T. Wakita and S. Nagamori for helpful discussion and suggestions. We also thank M. Matsuda, T. Shimoji and M. Yahata for technical assistance, and T. Mizoguchi for secretarial work. This work was supported in part by a grant for Research on Health Sciences focusing on Drug Innovation from the Japan Health Sciences Foundation; by grants-in-aid from the Ministry of Health, Labor and Welfare; and by the program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation, Japan.

References

Aizaki, H., Aoki, Y., Harada, T., Ishii, K., Suzuki, T., Nagamori, S., Toda, G., Matsuura, Y., Miyamura, T., 1998. Full-length complementary DNA of hepatitis C virus genome from an infectious blood sample. *Hepatology* 27, 621–627.

Aizaki, H., Nagamori, S., Matsuda, M., Kawakami, H., Hashimoto, O., Ishiko, H., Kawada, M., Matsuura, T., Hasumura, S., Matsuura, Y., Suzuki, T., Miyamura, T., 2003. Production and release of infectious hepatitis C virus from human liver cell cultures in the three-dimensional radial-flow bioreactor. *Virology* 314, 16–25.

Aoki, Y., Aizaki, H., Shimoike, T., Tani, H., Ishii, K., Saito, I., Matsuura, Y., Miyamura, T., 1998. A human liver cell line exhibits efficient translation of HCV RNAs produced by a recombinant adenovirus expressing T7 RNA polymerase. *Virology* 250, 140–150.

Bartenschlager, R., Kaul, A., Sparacio, S., 2003. Replication of the hepatitis C virus in cell culture. *Antivir. Res.* 60, 91–102.

Blight, K.J., Kolykhalov, A.A., Rice, C.M., 2000. Efficient initiation of HCV RNA replication in cell culture. *Science* 290, 1972–1974.

Boriskin, Y.S., Pecheur, E.I., Polyak, S.J., 2006. Arbidol: a broad-spectrum antiviral that inhibits acute and chronic HCV infection. *Virology* 343, 56.

Choo, Q.L., Kuo, G., Weiner, A.J., Overby, L.R., Bradley, D.W., Houghton, M., 1989. Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 244, 359–362.

Choo, Q.L., Richman, K.H., Han, J.H., Berger, K., Lee, C., Dong, C., Gallegos, C., Coit, D., Medina-Selby, R., Barr, P.J., et al., 1991. Genetic organization and diversity of the hepatitis C virus. *Proc. Natl. Acad. Sci. U.S.A.* 88, 2451–2455.

Dahari, H., Major, M., Zhang, X., Mihalik, K., Rice, C.M., Perelson, A.S., Feinstone, S.M., Neumann, A.U., 2005. Mathematical modeling of primary hepatitis C infection: noncytolytic clearance and early blockage of virion production. *Gastroenterology* 128, 1056–1066.

Dixit, N.M., Layden-Almer, J.E., Layden, T.J., Perelson, A.S., 2004. Modelling how ribavirin improves interferon response rates in hepatitis C virus infection. *Nature* 432, 922–924.

Grakoui, A., McCourt, D.W., Wychowski, C., Feinstone, S.M., Rice, C.M., 1993. Characterization of the hepatitis C virus-encoded serine proteinase: determination of proteinase-dependent polyprotein cleavage sites. *J. Virol.* 67, 2832–2843.

Hijikata, M., Kato, N., Ootsuyama, Y., Nakagawa, M., Shimotohno, K., 1991. Gene mapping of the putative structural region of the hepatitis C virus genome by in vitro processing analysis. *Proc. Natl. Acad. Sci. U.S.A.* 88, 5547–5551.

Hijikata, M., Shimizu, Y.K., Kato, H., Iwamoto, A., Shih, J.W., Alter, H.J., Purcell, R.H., Yoshikura, H., 1993. Equilibrium centrifugation studies of hepatitis C virus: evidence for circulating immune complexes. *J. Virol.* 67, 1953–1958.

Hongo, T., Kajikawa, M., Ishida, S., Ozawa, S., Ohno, Y., Sawada, J., Umezawa, A., Ishikawa, Y., Kobayashi, T., Honda, H., 2005. Three-dimensional high-density culture of HepG2 cells in a 5-ml radial-flow bioreactor for construction of artificial liver. *J. Biosci. Bioeng.* 99, 237–244.

Ikeda, M., Sugiyama, K., Mizutani, T., Tanaka, T., Tanaka, K., Sekihara, H., Shimotohno, K., Kato, N., 1998. Human hepatocyte clonal cell lines that support persistent replication of hepatitis C virus. *Virus Res.* 56, 157–167.

Iwahori, T., Matsuura, T., Maehashi, H., Sugo, K., Saito, M., Hosokawa, M., Chiba, K., Masaki, T., Aizaki, H., Ohkawa, K., Suzuki, T., 2003. CYP3A4 inducible model for in vitro analysis of human drug metabolism using a bioartificial liver. *Hepatology* 37, 665–673.

Kanto, T., Hayashi, N., Takehara, T., Hagiwara, H., Mita, E., Naito, M., Kasahara, A., Fusamoto, H., Kamada, T., 1994. Buoyant density of hepatitis C virus recovered from infected hosts: two different features in sucrose equilibrium density-gradient centrifugation related to degree of liver inflammation. *Hepatology* 19, 296–302.

Kawada, M., Nagamori, S., Aizaki, H., Fukaya, K., Niya, M., Matsuura, T., Sujino, H., Hasumura, S., Yashida, H., Mizutani, S., Ikenaga, H., 1998. Massive culture of human liver cancer cells in a newly developed radial flow bioreactor system: ultrafine structure of functionally enhanced hepatocarcinoma cell lines. *In Vitro Cell Dev. Biol. Anim.* 34, 109–115.

Kuo, G., Choo, Q.L., Alter, H.J., Gitnick, G.L., Redeker, A.G., Purcell, R.H., Miyamura, T., Dienstag, J.L., Alter, M.J., Stevens, C.E., et al., 1989. An assay for circulating antibodies to a major etiologic virus of human non-A, non-B hepatitis. *Science* 244, 362–364.

Lanford, R.E., Guerra, B., Lee, H., Averett, D.R., Pfeiffer, B., Chavez, D., Notvall, L., Bigger, C., 2003. Antiviral effect and virus-host interactions in response to alpha interferon, gamma interferon, poly(i)-poly(c), tumor necrosis factor alpha, and ribavirin in hepatitis C virus subgenomic replicons. *J. Virol.* 77, 1092–1104.

Layden, T.J., Layden, J.E., Ribeiro, R.M., Perelson, A.S., 2003. Mathematical modeling of viral kinetics: a tool to understand and optimize therapy. *Clin. Liver Dis.* 7, 163–178.

Layden-Almer, J.E., Cotler, S.J., Layden, T.J., 2006. Viral kinetics in the treatment of chronic hepatitis C. *J. Viral Hepat.* 13, 499–504.

Lindenbach, B.D., Evans, M.J., Syder, A.J., Wolk, B., Tellinghuisen, T.L., Liu, C.C., Maruyama, T., Hynes, R.O., Burton, D.R., McKeating, J.A., Rice, C.M., 2005. Complete replication of hepatitis C virus in cell culture. *Science* 309, 623–626.

Lohmann, V., Korner, F., Koch, J., Herian, U., Theilmann, L., Bartenschlager, R., 1999. Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science* 285, 110–113.

Martell, M., Esteban, J.I., Quer, J., Genesca, J., Weiner, A., Esteban, R., Guardia, J., Gomez, J., 1992. Hepatitis C virus (HCV) circulates as a population of different but closely related genomes: quasispecies nature of HCV genome distribution. *J. Virol.* 66, 3225–3229.

Murakami, K., Ishii, K., Ishihara, Y., Yoshizaki, S., Tanaka, K., Gotoh, Y., Aizaki, H., Kohara, M., Yoshioka, H., Mori, Y., Manabe, N., Shoji, I., Sata, T., Bartenschlager, R., Matsuura, Y., Miyamura, T., Suzuki, T., 2006. Production of infectious hepatitis C virus particles in three-dimensional cultures of the cell line carrying the genome-length dicistronic viral RNA of genotype 1b. *Virology* 351, 381–392.

- Pawlotsky, J.M., 2006. Hepatitis C virus population dynamics during infection. *Curr. Top. Microbiol. Immunol.* 299, 261–284.
- Perelson, A.S., Herrmann, E., Micol, F., Zeuzem, S., 2005. New kinetic models for the hepatitis C virus. *Hepatology* 42, 749–754.
- Poynard, T., Yuen, M.F., Ratziu, V., Lai, C.L., 2003. Viral hepatitis C. *Lancet* 362, 2095–2100.
- Saito, I., Miyamura, T., Ohbayashi, A., Harada, H., Katayama, T., Kikuchi, S., Watanabe, Y., Koi, S., Onji, M., Ohta, Y., et al., 1990. Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc. Natl. Acad. Sci. U.S.A.* 87, 6547–6549.
- Shimizu, Y.K., Hijikata, M., Iwamoto, A., Alter, H.J., Purcell, R.H., Yoshikura, H., 1994. Neutralizing antibodies against hepatitis C virus and the emergence of neutralization escape mutant viruses. *J. Virol.* 68, 1494–1500.
- Suzuki, T., Omata, K., Satoh, T., Miyasaka, T., Arai, C., Maeda, M., Matsuno, T., Miyamura, T., 2005. Quantitative detection of hepatitis C virus (HCV) RNA in saliva and gingival crevicular fluid of HCV-infected patients. *J. Clin. Microbiol.* 43, 4413–4417.
- Tagawa, M., Kato, N., Yokosuka, O., Ishikawa, T., Ohto, M., Omata, M., 1995. Infection of human hepatocyte cell lines with hepatitis C virus in vitro. *J. Gastroenterol. Hepatol.* 10, 523–527.
- Wakita, T., Pietschmann, T., Kato, T., Date, T., Miyamoto, M., Zhao, Z., Murthy, K., Habermann, A., Krausslich, H.G., Mizokami, M., Bartenschlager, R., Liang, T.J., 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* 11, 791–796.
- Zhong, J., Gastaminza, P., Cheng, G., Kapadia, S., Kato, T., Burton, D.R., Wieland, S.F., Uprichard, S.L., Wakita, T., Chisari, F.V., 2005. Robust hepatitis C virus infection in vitro. *Proc. Natl. Acad. Sci. U.S.A.* 102, 9294–9299.

The NS3 Helicase and NS5B-to-3'X Regions Are Important for Efficient Hepatitis C Virus Strain JFH-1 Replication in Huh7 Cells[∇]

Asako Murayama,¹ Tomoko Date,¹ Kenichi Morikawa,¹ Daisuke Akazawa,^{1,2} Michiko Miyamoto,³ Minako Kaga,¹ Koji Ishii,¹ Tetsuro Suzuki,¹ Takanobu Kato,^{4,5} Masashi Mizokami,⁴ and Takaji Wakita^{1*}

Department of Virology II, National Institute of Infectious Diseases, Tokyo, Japan¹; Pharmaceutical Research Lab, Toray Industries, Inc., Kanagawa, Japan²; Department of Microbiology, Tokyo Metropolitan Institute for Neuroscience, Tokyo, Japan³; Department of Clinical Molecular Informative Medicine, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan⁴; and Liver Disease Branch, NIDDK, National Institutes of Health, Bethesda, Maryland⁵

Received 23 September 2006/Accepted 8 May 2007

The JFH-1 strain of hepatitis C virus (HCV) is a genotype 2a strain that can replicate autonomously in Huh7 cells. The J6 strain is also a genotype 2a strain, but its full genomic RNA does not replicate in Huh7 cells. However, chimeric J6/JFH-1 RNA that has J6 structural-protein-coding regions and JFH-1 nonstructural-protein-coding regions can replicate autonomously and produce infectious HCV particles. In order to determine the mechanisms underlying JFH-1 RNA replication, we constructed various J6/JFH-1 chimeras and tested their RNA replication and virus particle production abilities in Huh7 cells. Via subgenomic-RNA-replication assays, we found that both the JFH-1 NS5B-to-3'X (NSBX) and the NS3 helicase (N3H) regions are important for the replication of the J6CF replicon. We applied these results to full-length genomic RNA replication and analyzed replication using Northern blotting. We found that a chimeric J6 clone with JFH-1 N3H and NSBX could replicate autonomously but that a chimeric J6 clone with only JFH-1 NSBX had no replication ability. Finally, we tested the virus production abilities of these clones and found that a chimeric J6 clone with JFH-1 N3H and NSBX could produce infectious HCV particles. In conclusion, the JFH-1 NS3 helicase and NS5B-to-3'X regions are important for efficient replication and virus particle formation of HCV genotype 2a strains.

Hepatitis C virus (HCV) is a major cause of chronic liver disease (7, 22). The lack of a robust cell culture system for producing virus particles has hampered the development of HCV research (2). Although the development of a subgenomic-replicon system enabled research into HCV RNA replication (32), infectious-virus-particle production remained impossible. Recently, an HCV cell culture system was developed using a JFH-1 genotype 2a strain of HCV cloned from a fulminant hepatitis patient (30, 48, 54), allowing investigation of the virus life cycle.

HCV is a positive-strand RNA virus that belongs to the *Hepacivirus* genus in the *Flaviviridae* family. The HCV genome comprises about 9,600 nucleotides that encode a single polyprotein of around 3,000 amino acids (8, 18, 44), which is processed by cellular and viral encoded proteases into at least 10 different structural and nonstructural proteins (11, 13, 14, 33).

The JFH-1 strain of HCV is a genotype 2a strain, and it is the first HCV strain that can produce HCV particles in Huh7 cells (48). Subgenomic replicons of JFH-1 replicate efficiently in Huh7 cells and do not require cell culture-adaptive mutations (19). The J6CF strain of HCV is also a genotype 2a strain and is known to be infectious in chimpanzees (49), but its

entire genomic RNA does not replicate in Huh7 cells, despite the ~90% nucleotide sequence homology between JFH-1 and J6CF. However, J6/JFH-1 chimeric RNA that has J6 structural-protein-coding regions and JFH-1 nonstructural-protein-coding regions can replicate autonomously and produce infectious HCV particles (30, 39). Why only the JFH-1 clone can replicate efficiently in Huh7 cells remains unclear.

In this study, to investigate the mechanisms underlying efficient JFH-1 replication, we focused on the differences in replication between JFH-1 and J6CF strains by using intragenotypic JFH-1 and J6CF chimeras and compared their respective abilities to replicate RNA and produce virus particles in Huh7 cells.

MATERIALS AND METHODS

Cell culture. Huh7 cells (36) were cultured at 37°C in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum under 5% CO₂ conditions.

Subgenomic-replicon constructs. pSGR-JCH1 and pSGR-JCH4 were constructed based on pSGR-JFH1 (19, 21). pSGR-J6CF was also constructed from pJ6CF (a kind gift from Jens Bukh) (49), using the same method used to construct pSGR-JFH1. Plasmids used in luciferase assays were constructed based on pSGR-JFH1/Luc (20). Chimeric replicons were constructed by substitution of the corresponding regions. For convenience, several restriction enzyme recognition sites (ClaI [2275], EcoT22I [3639], and BsrGI [6127]) were introduced into the pSGR-J6CF sequence via nucleotide substitutions. The substitutions of the corresponding regions were achieved as follows, with the 5' untranslated region (5'UTR) inserted between NotI and AgeI: NS3, PmeI-EcoT22I; NS3 protease, PmeI-ClaI; NS3 helicase, ClaI-EcoT22I; NS4, EcoT22I-MunI; NS5A, MunI-BsrGI; NS5B, BsrGI-StuI; and 3'UTR, StuI-XbaI (see Fig. 2A and 3A). pSGR-JCH1/Luc and pSGR-JCH4/Luc were also constructed using the same procedure as that for pSGR-JFH1/Luc (20, 21). The Con1 replicon (pSGR-Con1/Luc) was

* Corresponding author. Mailing address: Department of Virology II, National Institute of Infectious Diseases, Toyama 1-23-1, Shinjuku, Tokyo 162-8640, Japan. Phone: 81-3-5285-1111. Fax: 81-3-5285-1161. E-mail: wakita@nih.go.jp.

[∇] Published ahead of print on 23 May 2007.




FIG. 1. G418-resistant colony formation of JFH-1, J6CF, JCH-1, and JCH-4. Subgenomic RNAs were synthesized *in vitro*, using pSGR-JFH1, pSGR-J6CF, pSGR-JCH1, and pSGR-JCH4 as templates. Transcribed subgenomic RNAs were electroporated into Huh7 cells, and cells were cultured with G418 for 3 weeks before staining with crystal violet as described in Materials and Methods. JFH-1 subgenomic RNA (0.1 μ g) and 10 μ g of J6CF, JCH-1, and JCH-4 subgenomic RNAs were transfected into Huh7 cells. Experiments were performed in triplicate, and representative staining examples are shown.

constructed from pFK-I389/neo/NS3-3'/wt (a kind gift from Ralf Bartenschlager) (32), and the H77c replicon (pSGR-H77c/Luc) was constructed from pCV-H77c (a kind gift from Robert H. Purcell) (50). For convenience, ClaI (2275) and BsrGI (6127) recognition sites were introduced into the pSGR-Con1/Luc and pSGR-H77c/Luc sequences via nucleotide substitutions. Substitutions of the NS3 helicase region and NSBX regions were performed as described above.

Full-length genomic HCV constructs. Plasmids used in the analysis of genomic RNA replication were constructed based on pJFH1 (48) and pJ6CF (49). For convenience, several restriction enzyme recognition sites (ClaI [3929], EcoT22I [5293], and BsrGI [7781]) were introduced into the J6CF sequence via nucleotide substitutions. Substitutions of the NS3 helicase regions were performed by replacement of the ClaI-EcoT22I fragment, substitutions of the NSBX regions were performed by replacement of the BsrGI-XbaI fragment, substitutions of the NSB regions were performed by replacement of the BsrGI-StuI fragment, and a substitution of the 3'UTR was performed by replacement of the StuI-XbaI fragment (see Fig. 5A).

RNA synthesis and transfection. RNA synthesis and transfection were performed as described previously (48). In brief, plasmids were linearized with XbaI, treated with mung bean nuclease (New England Biolabs, Ipswich, MA), and purified. Linearized, purified DNAs were used as templates for *in vitro* RNA synthesis using a MEGascript T7 kit (Ambion, Austin, TX) in accordance with the manufacturer's instructions. Synthesized RNA was treated with DNase I (Ambion), followed by purification using ISOGEN-LS (Nippon Gene, Tokyo, Japan). The quality of synthesized RNA was examined by agarose gel electrophoresis. Ten micrograms of *in vitro*-synthesized RNA was used for each electroporation. Trypsinized Huh7 cells (3×10^6 cells) were washed with Opti-MEM I (Invitrogen, Carlsbad, CA) and resuspended in Cytomix buffer (47). RNA was mixed with 400 μ l of cell suspension, and the mixture was then transferred to an electroporation cuvette (Precision Universal Cuvettes, Thermo Hybaid, Middlesex, United Kingdom). The cells were then pulsed at 260 V and 950 μ F using a Gene Pulser II apparatus (Bio-Rad, Hercules, CA). Transfected cells were immediately transferred to 10-cm culture dishes or six-well plates, each containing culture medium, and incubated at 37°C under 5% CO₂. Luciferase mRNA was synthesized from luciferase T7 control DNA (Promega, Madison, WI) by using a mMESAGE mMACHINE T7 kit (Ambion). To monitor transfection efficiency, *in vitro*-synthesized luciferase RNA was cotransfected with HCV RNA and luciferase activity measured at 4 h after transfection.

G418-resistant colony formation assay. The G418-resistant colony formation assay was performed as described previously (19). In brief, 0.1 μ g or 10 μ g of transcribed RNAs was transfected into 3×10^6 Huh7 cells by electroporation. Transfected cells were immediately transferred to 10-cm culture dishes containing 10 ml of culture medium. G418 (1.0 mg/ml) (Nakalai Tesque, Kyoto, Japan) was added to the culture medium at 16 to 24 h after transfection. Culture medium supplemented with G418 was replaced every 3 days. Three weeks after transfection, cells were fixed with buffered formalin and stained with crystal violet.

Luciferase reporter assay. The luciferase activities of the JFH-1 subgenomic replicon and chimeras in Huh7 cells were measured as described previously (20). Briefly, 5 μ g of transcribed RNAs was transfected into 3×10^6 Huh7 cells by electroporation. Transfected cells were immediately resuspended in culture medium and seeded into six-well culture plates. Cells were harvested serially at 4, 24, and 48 h after transfection and lysed with 200 μ l of cell culture lysis reagent

(Promega). Debris was then removed by centrifugation. Luciferase activity was quantified using a Lumat LB9507 luminometer (EG & G Berthold, Bad Wildbad, Germany) and a luciferase assay system (Promega). Assays were performed three times independently, with each value corrected for transfection efficiency as determined by measuring luciferase activity 4 h after transfection. The data are expressed as relative luciferase units (RLU).

Quantification of HCV core protein. To estimate the concentration of HCV core protein in the culture medium, we performed an HCV core enzyme-linked immunosorbent assay (Ortho-Clinical Diagnostics, Tokyo, Japan) in accordance with the manufacturer's instructions.

Northern blot analysis. Northern blot analysis was performed as described previously (48). In brief, total cellular RNA from HCV RNA-transfected cells was extracted using ISOGEN (Nippon Gene) in accordance with the manufacturer's instructions. Isolated RNA (2 μ g) was separated on a 1% agarose gel containing formaldehyde, transferred to a Hybond N+ positively charged nylon membrane (GE Healthcare, Piscataway, NJ), and immobilized using a Stratalinker UV cross-linker (Stratagene, La Jolla, CA). Hybridization was performed with [α -³²P]dCTP-labeled DNA by using Rapid-Hyb buffer (GE Healthcare). The DNA probe was synthesized using the NSSB-to-3'X fragment of JFH1 excised from pJFH1 by BsrGI and XbaI and labeled using the Megaprime DNA labeling system (GE Healthcare).

Infection of cells with secreted HCV and determination of infectivity. Culture medium from RNA-transfected cells was collected at 72 h posttransfection. Huh7 cells were seeded at a density of 1×10^4 cells per well in poly-D-lysine-coated 96-well plates (CORNING, Corning, NY). On the following day, the collected culture media were serially diluted and used for inoculation of the seeded cells, and the plates were incubated for another 3 days at 37°C. The cells were fixed in methanol for 15 min at -20°C, and the infected foci were visualized by immunofluorescence as described below.

Cells were blocked for 1 h with BlockAce (Dainippon Sumitomo Pharma, Osaka, Japan) supplemented with 0.3% Triton X-100 and then washed with phosphate-buffered saline, followed by incubation with anti-core antibody at 50 μ g/ml in BlockAce. After incubation for 1 h at room temperature, the cells were washed and incubated with a 1:400 dilution of AlexaFluor 488-conjugated anti-mouse immunoglobulin G (Molecular Probes, Eugene, OR) in BlockAce. The cells were then washed and examined using fluorescence microscopy (Olympus, Tokyo, Japan). Infectivity was quantified by counting the infected foci and expressed as numbers of focus-forming units per milliliter (FFU/ml).

RESULTS

G418-resistant colony formation of JFH-1, J6CF, and other genotype 2a subgenomic replicons. First, to compare the replication efficiencies of the JFH-1 and J6CF strains, we performed a G418-resistant colony formation assay with JFH-1 and J6CF RNAs by using subgenomic replicons. The JFH-1 subgenomic replicon formed many colonies with transfection of only 0.1 μ g RNA, but the J6CF subgenomic replicon formed no colonies, even with transfection of 10 μ g RNA (Fig. 1). We also tested




FIG. 2. Luciferase activities of chimeric replicons with a JFH-1 backbone. (A) Structures of chimeric subgenomic replicons with a JFH-1 backbone. The restriction enzyme recognition sites used for the construction of plasmids are indicated. *P*, *Pme*I; *C*, *Cla*I; *E*, *Eco*T22I; *M*, *Mun*I; *B*, *Bsr*GI; *S*, *Stu*I; *X*, *Xba*I; wt, wild type. (B) Subgenomic RNAs were synthesized *in vitro* from wild-type or chimeric replicon constructs. Transcribed subgenomic RNAs (5 μ g) were electroporated into Huh7 cells, and cells were harvested serially at 4, 24, and 48 h after transfection. The harvested cells were lysed, and then luciferase activities in the cell lysates were measured. The assays were performed three times independently and the results expressed as luciferase activities (RLU). Each value was corrected for transfection efficiency as determined by measuring the luciferase activity 4 h after transfection. Data are presented as means and standard deviations for luciferase activity at 24 h (white bars) and 48 h (gray bars) after transfection.

other genotype 2a clones (the JCH-1 and JCH-4 strains), which were isolated from patients with chronic hepatitis C (21). Their subgenomic replicons did not form colonies either. Given that chimeric J6/JFH-1 RNA that has the J6 structural-protein-coding

regions and JFH-1 nonstructural-protein-coding regions reportedly replicates autonomously and produces infectious HCV particles (30, 39), we hypothesized that some of the JFH-1 nonstructural-protein-coding regions are important for JFH-1 replication.