

Synthetic small interfering RNA targeting heat shock protein 105 induces apoptosis of various cancer cells both *in vitro* and *in vivo*

Seiji Hosaka,¹ Tetsuya Nakatsura,^{1,4} Hirotake Tsukamoto,¹ Takumi Hatayama,³ Hideo Baba² and Yasuharu Nishimura^{1,5}

Departments of ¹Immunogenetics and ²Gastroenterological Surgery, Graduate School of Medical Sciences, Kumamoto University, 1-1-1 Honjo, Kumamoto 860-8556, and ³Department of Biochemistry, Kyoto Pharmaceutical University, 5 Nakauchi-cho, Misasagi, Yamashina-ku, Kyoto 607-8414, Japan

(Received December 17, 2005/Revised February 21, 2006/Accepted February 25, 2006/Online publication April 19, 2006)

We previously reported that heat shock protein 105 (HSP105), identified by serological analysis of a recombinant cDNA expression library (SEREX) using serum from a pancreatic cancer patient, was overexpressed in various human tumors and in the testis of adult men by immunohistochemical analysis. In the present study, to elucidate the biological function of the HSP105 protein in cancer cells, we first established NIH3T3 cells overexpressing murine HSP105 (NIH3T3-HSP105). The NIH3T3-HSP105 cells acquired resistance to apoptosis induced by heat shock or doxorubicin. The small interfering RNA (siRNA)-mediated suppression of HSP105 protein expression induced apoptosis in human cancer cells but not in fibroblasts. By a combination of siRNA introduction and doxorubicin or heat shock treatment, apoptosis was induced synergistically in a human colon cancer cell line, HCT116. *In vivo*, siRNA inoculation into the human gastric cancer cell line KATO-3 established in the flank of an NOD SCID mouse suppressed the tumor growth. This siRNA-induced apoptosis was mediated through caspases, but not the p53 tumor suppressor protein, even though the HSP105 protein was bound to wild-type p53 protein in HCT116 cells. These findings suggest that the constitutive overexpression of HSP105 in cancer cells is involved in malignant transformation by protecting tumor cells from apoptosis. HSP105 may thus be a novel target molecule for cancer therapy and a treatment regimen using synthetic siRNA to suppress the expression of HSP105 protein may provide a new strategy for cancer therapy. (*Cancer Sci* 2006; 97: 623–632)

Heat shock protein 105 (also called HSP110)⁽¹⁾ is a stress protein belonging to the HSP105/110 family that is expressed constitutively in most tissues at low levels, whereas HSP105 mRNA is expressed at high levels in mouse and rat brain. At the protein level, high expression levels have been reported only in the brain of mice.^(1–4) Like other heat shock proteins, HSP105 plays an important role as a chaperone under physiological conditions. HSP105 is induced by various stressors and plays an important role in protecting cells from the cytotoxic effects mediated by such stressors. HSP105 is composed of an ATP-binding domain, a β -sheet, a loop and α -helical domains similar to those observed in the HSP70 family of proteins, and it binds to non-native protein substrates, thereby preventing the aggregation of denatured protein through an interaction with the β -sheet domain of HSP105.^(2,5–7)

The rat neuronal cell line PC12 transfected stably with murine *HSP105* exhibited resistance to caspase-mediated apoptosis induced by stressors.⁽⁸⁾ In a spinal and bulbar muscular atrophy model, the transient expression of tAR containing an expanded polyglutamine tract caused aggregates of polyglutamine in COS-7 and SK-N-SH cells and then induced the cells to undergo apoptosis. In contrast, in cells cotransfected with tAR and *HSP105*, both the aggregation of polyglutamine and the degree of its cell toxicity decreased.⁽⁹⁾ In contrast, HSP105 has demonstrated apoptosis-enhancing activity during murine embryogenesis.^(10–13) These observations suggest that HSP105 is involved in the regulation of apoptosis.

We previously reported that HSP105 is overexpressed in various human tumors, including colon cancer cells but not colorectal adenomas, thus suggesting that the overexpression of HSP105 is a late event in the colorectal adenoma–carcinoma sequence.^(14,15) Recent studies have also demonstrated that the expression level of HSP105 is elevated in highly metastatic colon cancer cell lines, and is correlated with advanced clinical stages and positive lymph node involvement.⁽¹⁶⁾

RNA interference is used widely for manipulating biological systems, and has also been utilized successfully as a therapeutic material in experimental animals.^(17–19) Synthetic siRNA strongly inhibits the expression of target proteins when they are transfected with cationic liposomes, which are thought to be safer than viral vectors for human therapy. The local injection of synthetic siRNA against VEGF⁽²⁰⁾ or sphingosine 1-phosphate receptor-1⁽²¹⁾ into established tumors has been reported recently to suppress angiogenesis and tumor growth.

The role of HSP105 in cancer cells has yet to be elucidated. In the present study, we first transfected the *HSP105* gene

⁴T. Nakatsura is currently in the Immunotherapy Section, Investigative Treatment Division Research Center for Innovative Oncology, National Cancer Center Hospital East, Kashiwa City, Chiba, Japan.

⁵To whom correspondence should be addressed.

E-mail: mxnishim@gpo.kumamoto-u.ac.jp

Abbreviations: CML, chronic myelocytic leukemia; DMEM, Dulbecco's modified Eagle's medium; ER, endoplasmic reticulum; FCS, fetal calf serum; FITC, fluorescein-isothiocyanate; GFP, green fluorescent protein; HBSS, Hanks' balanced salt solution; HSP70, heat shock protein 70; HSP105, heat shock protein 105; IRES, internal ribosomal entry site; NIH3T3-HSP105, NIH3T3 cells overexpressing murine HSP105; PARP, poly ADP-ribose polymerase; PBS, phosphate-buffered saline; PI, propidium iodide; RT-PCR, reverse transcription-polymerase chain reaction; SEREX, serological analysis of a recombinant cDNA expression library; siRNA, small interfering RNA; tAR, truncated androgen receptor; VEGF, vascular endothelial growth factor.

into NIH3T3 cells to examine the function of those cells, while also treating those transfectants with several stressors. To investigate whether the suppression of HSP105 expression affects the growth of cancer cells, we then introduced synthetic siRNA specific to HSP105 into several human cancer cell lines both *in vitro* and *in vivo*. We herein report that HSP105 has an anti-apoptotic function and that an overexpression of HSP105 is essential for cancer cells to survive.

Materials and Methods

Cell lines and culture

The human pancreatic cancer cell line PK8 and the murine fibroblast cell line NIH3T3 were kindly provided by the Cell Resource Center for Biomedical Research Institute of Development, Aging and Cancer, Tohoku University (Sendai, Japan). The human hepatoma cell line SK-Hep1, human colon cancer cell line SW620, and human gastric cancer cell lines KATO-3 and MKN28 were kindly provided by Dr K. Itoh, Kurume University (Kurume, Japan). The human colon cancer cell line HCT116 was kindly provided by Dr B. Vogelstein, Johns Hopkins University (Baltimore, MD, USA). Primary normal fibroblast strains Turu and Mori were kindly provided by Dr M. Yamaizumi, Kumamoto University (Kumamoto, Japan).

NIH3T3, HCT116, SW620, Turu and Mori were all cultured *in vitro* in DMEM, and SK-Hep1, PK8, KATO-3 and MKN28 were cultured in RPMI medium supplemented with 10% FCS and 1% penicillin and streptomycin in a 5% CO₂ atmosphere at 37°C.

Mice

C57BL/6 NOD SCID mice were kindly provided by Dr S. Okada, Kumamoto University. The mice were maintained at the Center for Animal Resources and Development of Kumamoto University and they were handled in accordance with the animal care policy of Kumamoto University.

Plasmid construction and transfection

To obtain pCAGGS-IRES-neo-R, a DNA fragment containing an IRES, the neomycin-resistance gene *neo-R* was inserted into the mammalian expression vector pCAGGS. A cDNA fragment encoding the murine HSP105 protein was inserted into pCAGGS-IRES-neo-R. Cell transfection was carried out using Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's recommendations in six-well multiplates. At 48 h after transfection, the cells were replated and were then cultured in the presence of an appropriate concentration of G418 for 1 week.

RT-PCR analysis

Total RNA was isolated using either TRIZOL reagent (Gibco BRL, Rockville, MD, USA) or RNeasy spin column kits (Qiagen, Valencia, CA, USA). Total RNA from normal human colon tissue was purchased from Clontech (Palo Alto, CA, USA). RT-PCR analysis was carried out as described previously.⁽²²⁾ Briefly, 1 µg of total RNA was converted into cDNA in 20 µL of reaction buffer. Each PCR regime involved an initial denaturation step at 94°C for 5 min followed by 24–33 cycles for each type of cDNA. All samples were then processed at 94°C for 1 min, 58°C for 1 min and 72°C for 1 min

The primer sequences were as follows: *HSP105* forward 5'-ATGAAGTGATGGAATGGATG-3' and reverse 5'-TTTGGTT-

TCGGTTGTGTTAC; *NOXA* forward 5'-AGATGCCTGGGAA-GAAG-3' and reverse 5'-AGTCCCCTCATGCAAGT-3'; *PUMA* forward 5'-TGTAGAGGAGACAGGAATCCACGG-3' and reverse, 5'-AGGCACCTAATTGGGCTCCATCTC-3'; *Bax* forward 5'-AGCGGCGGTGATGGACGGGTC-3' and reverse 5'-TCCAAGGCAGCTGGGGCCTCA-3'; and *p53* forward 5'-CCATGGCCATCTACAAGCAG-3' and reverse 5'-AGGGT-GAAATATTCTCC-3'. PCR products were visualized by ethidium bromide staining after separation on a 2% agarose gel.

Detection of cell apoptosis

Cells in the early phase of apoptosis were detected by staining with annexin V, which binds to a phosphatidyl serine marker specific for the early phase of apoptosis, using an annexin V-FITC apoptosis detection kit (BioVision, Mountain View, CA, USA). The NIH3T3-mock cells and NIH3T3-HSP105 cells were incubated at 45°C for 90 min or with 200 ng/mL doxorubicin. The cancer cell lines were treated with either 100 nM or 200 nM siRNA, with 100 nM siRNA and 200 ng/mL doxorubicin or with 100 nM siRNA and heat shock (45°C, 30 min). The cells were harvested at the times indicated and were then stained with FITC-conjugated annexin V for flow cytometric analysis according to the manufacturer's recommendations. In the caspase inhibition assay, 100 µM Z-VAD-FMK (Sigma, St Louis, MO, USA) was added 1 h before siRNA transfection.

To detect the late phase of apoptosis, DNA fragmentation in the cells was evaluated by staining with PI and flow cytometric analysis as follows: NIH3T3-mock cells and NIH3T3-HSP105 cells treated with doxorubicin were collected by trypsinization, washed with PBS, fixed in cold 70% ethanol, and stored at -20°C until staining. After fixation the cells were washed in PBS and incubated with 100 µg/mL of RNaseA for 30 min at room temperature, before staining with 25 µg/mL of PI. Flow cytometry was carried out using a FACScan flow cytometer (BD Biosciences, San Jose, CA, USA) and the data were analyzed using CellQuest software (BD Biosciences).

Immunoprecipitation and western blot analysis

The cell samples were lysed in appropriate amounts of lysing buffer (200 mM NaCl, 20 mM Tris [pH 7.4], 1% Nonidet P-40, 1 mM sodium orthovanadate [WAKO, Osaka, Japan], 10% glycerol, plus one protease inhibitor tablet [Roche Applied Sciences, Penzberg, Germany]). Hsp105 and p53 were immunoprecipitated with rabbit polyclonal anti-HSP105 antibody and mouse monoclonal anti-p53 antibody (DO-1; Santa Cruz Biotechnology, Santa Cruz, CA, USA), respectively, together with protein A beads (Pierce, Rockford, IL, USA). The proteins were analyzed using 7% or 10% sodium dodecylsulfate-polyacrylamide gel electrophoresis and were then transferred onto a nitrocellulose membrane. HSP105 was blotted with the above-described antibody and PARP was blotted with rabbit polyclonal anti-PARP antibody (Santa Cruz Biotechnology). p53 was blotted with DO-1 or DO-1 and labeled with biotin using the Mini-biotin-XX Protein Labeling Kit (Molecular Probes, Eugene, OR, USA). Phospho-p53 (Ser46) and phospho-p53 (Ser15) were blotted with rabbit polyclonal antibody specific to phospho-p53 (Ser46) and mouse monoclonal antibody specific to phospho-p53 (Ser15) (New England Biolabs, Beverly, MA, USA), respectively, and then with horseradish peroxidase-conjugated rabbit antimouse IgG or donkey antirabbit IgG

(Amersham Biosciences, Piscataway, NJ, USA), respectively, as secondary antibodies. The bands were visualized by enhanced chemiluminescence (Amersham Biosciences).

siRNA and *in vivo* siRNA treatment

The siRNA duplexes were purchased from Dharmacon Research (HSP105-siRNA and luciferase; Lafayette, CO, USA), Qiagen (GFP) and Invitrogen (HSP105-siRNA-2). The siRNA sequences used were as follows: HSP105, UUGGCUGCAACUCCG-AUUGTT; HSP105-siRNA-2, UGUACAUUACCUUUUUU-CCAACUCC; luciferase, CGUACGCGGAUACUUGATT; and GFP, GCAAGCUGACCCUGAAGUUA. The transfection of siRNA oligonucleotides was carried out using oligofectamine (Invitrogen) according to the manufacturer's recommendations in a six-well plate.

KATO-3 cells (2×10^6) were suspended in 100 μ L of HBSS solution (Gibco, Langley, OK, USA), injected subcutaneously into the dorsal skin of NOD SCID mice and were then allowed to grow. After 10 days, siRNA solutions were injected locally into tumors every third day. siRNA solutions were prepared by incubating 1 nmol of siRNA and 20 μ L of oligofectamine in 5% glucose solution for 15 min at room temperature. The tumor volume was calculated using the equation

$$V = (L \times W^2) \times 0.5,$$

where V = volume, L = length and W = width.

Statistical analysis

The two-tailed Student's *t*-test was used to determine the statistical significance of differences in the percentage of the cell fraction evaluated by flow cytometric analysis, and in tumor size between the treatment groups. A value of $P < 0.05$ was considered to be significant. Statistical analyses were carried out using the StatView 5.0 software package (Abacus Concepts, Calabasas, CA, USA).

Results

NIH3T3-HSP105 cells acquired resistance to apoptosis induced by stressors

To elucidate the biological function of HSP105 in cancer cells, we first transfected the murine HSP105 expression vectors or an empty vector into murine embryonal fibroblast NIH3T3 cells, which are known to express smaller amounts of HSP105 than cancer cells. Subsequently, NIH3T3-mock cell lines and NIH3T3-HSP105 cell lines overexpressing HSP105 protein were established after selecting cells with G418 (Fig. 1A). There was no difference in either the morphology or the proliferative characteristics between these two cell lines cultured in DMEM supplemented with 1, 5 or 10% FCS (data not shown).

In a previous study, Hatayama *et al.* reported that rat neuronal cells overexpressing murine HSP105 were able to avoid undergoing apoptosis induced by several kinds of stressors.⁽⁸⁾ We therefore examined the anti-apoptotic effects of HSP105 overexpression in NIH3T3-HSP105 cells. We exposed NIH3T3-mock cells and NIH3T3-HSP105 cells to heat shock at 45°C for 90 min and then detected annexin V-positive early apoptotic cells by flow cytometric analysis. The proportion of annexin V-positive NIH3T3-HSP105 cells was smaller than that of NIH3T3-mock cells ($P < 0.01$) (Fig. 1B).

We further treated these two cell lines with 200 ng/mL doxorubicin, which is known to be an inducer of apoptosis. After incubating NIH3T3-mock cells with doxorubicin for 12 h, annexin V-positive cells were detected by flow cytometric analysis. The number of annexin V-positive cells increased gradually thereafter, and almost all cells were stained with annexin V by 24 h after the treatment. However, even after 36 h incubation of cells with doxorubicin, only a small fraction (<18%) of NIH3T3-HSP105 cells were stained with annexin V. The difference in the number of annexin V-positive cells at 48 h after treatment was statistically significant ($P < 0.001$) between NIH3T3-mock and NIH3T3-HSP105 (Fig. 1C).

We next examined the DNA fragmentation of those two cell lines, which is observed during the late phase of apoptosis by staining with PI. The percentage of NIH3T3-HSP105 cells exhibiting DNA fragmentation was less than that of NIH3T3-mock cells, and the difference was statistically significant at both 36 h ($P < 0.001$) and 48 h ($P < 0.0001$) after doxorubicin treatment (Fig. 1D). These data suggest that HSP105 has an anti-apoptotic effect against apoptosis induced by heat shock and doxorubicin.

HSP105 siRNA induced various human cancer cell lines to undergo apoptosis

We previously reported that HSP105 protein is overexpressed in various human tumors and in the testis of normal adult men, but not in colon adenoma, by immunohistochemical analysis.⁽¹⁵⁾ We thus examined the function of HSP105 protein in cancer cells by downregulating *HSP105* gene expression with siRNA. We used two human colon cancer cell lines, HCT116 and SW620, in which the expression of HSP105 mRNA was significantly elevated in comparison to that in normal human colon epithelium (Fig. 2A). At approximately 24 h after transfection, the adherent HCT116 cells treated with HSP105 siRNA started peeling off, and almost all of the cells had peeled off and were observed as dying cells at 48 h after transfection. In contrast, most of the HCT116 cells treated with luciferase siRNA proliferated normally (Fig. 2B). In a western blot analysis of HCT116 cells, the expression of HSP105 protein was markedly suppressed after treatment with two different HSP105-specific siRNA (HSP105-siRNA and HSP105-siRNA-2), and those cells were significantly stained with FITC-annexin V based on flow cytometric analysis (Fig. 2B,C).

Similarly, in SW620 cells after treatment with HSP105-siRNA, the expression of HSP105 protein was suppressed and a significant number of annexin V-positive cells were detected in proportion to the concentration of siRNA administered to the SW620 cells. The siRNA effect was more notable at 24 h than at 48 h after siRNA treatment in SW620 (Fig. 2C). This finding may be associated with the fact that the expression of HSP105 in SW620 was much higher than any of the other cancer cell lines tested (Fig. 3A). In addition, in other cancer cell lines, including the human hepatoma cell line SK-Hep1, human pancreatic cancer cell line PK8, and two human gastric cancer cell lines KATO-3 and MKN28, HSP105 protein expression was suppressed and all of these cell lines underwent apoptosis at 48 h after transfection of HSP105 siRNA (Fig. 2D). These data indicate that the suppression of HSP105 protein expression by siRNA can thus induce human cancer cells originating from various tissues to undergo apoptosis.




Fig. 1. Anti-apoptotic effect of HSP105 overexpression on NIH3T3 cells. (A) Western blot analysis of HSP105 expressed in NIH3T3 cells transfected with pCAGGS-IRES-neo-R or pCAGGS-IRES-neo-R-HSP105. β -Actin is shown as a control for the equal loading of protein. (B–D) Flow cytometric analyses of apoptotic cells. NIH3T3-mock cells and NIH3T3-HSP105 cells were treated at 45°C for 90 min (B) or with 200 ng/mL doxorubicin (C,D). To detect early apoptotic cells, the cells were harvested at the times indicated, stained with fluorescein-isothiocyanate-annexin V and analyzed by flow cytometry (C,D). These data are representative of at least three independent experiments. Percentages shown in the panel indicate percentage of annexin V-positive cells in heat-treated cells (B) and doxorubicin-treated cells (C). (D) Detection of DNA fragmentation by propidium iodide staining. The percentage of sub-G₁ fractions at the times indicated is shown, and the representative data of flow cytometric analysis at 48 h is shown in the panel. Data are mean \pm SD ($n = 3$). The asterisk indicates that the difference in the percentages of the sub-G₁ fractions is statistically significant between the two values indicated by lines ($P < 0.001$).

HSP105 siRNA did not induce human fibroblasts to undergo apoptosis

HSP105 is a stress-induced protein that is usually expressed ubiquitously at low levels, except in the brain and testis. To investigate the effect of HSP105 siRNA on normal cells, we applied HSP105 siRNA to human fibroblasts, Turu and Mori, generated from healthy donors. In western blot analysis, the level of HSP105 protein expressed in the Turu and Mori cells was lower than in the cancer cell lines (Fig. 3A). The expression of HSP105 protein was suppressed with HSP105 siRNA treatment, as shown in Fig. 3B,C, and the shape of fibroblast cells changed from fusiform to a round shape, but those cells were not induced to undergo apoptosis. Approximately 10 days after HSP105 siRNA transfection, both the expression of HSP105 protein and cell shape were restored (data not shown). These results indicate that the suppressive effect of HSP105 siRNA on the HSP105 expression is transient and reversible, and that the marked reduction of HSP105 protein does not have any harmful effect on normal fibroblasts under non-stressed conditions.

HSP105 siRNA treatment suppressed the growth of tumors overexpressing HSP105 *in vivo*

Small interfering RNA treatment using liposomes suppresses tumor growth *in vivo*.^(21,23) To elucidate the effects of HSP105 siRNA on growing tumors *in vivo*, we injected either HSP105 siRNA or irrelevant siRNA locally into 5–7 mm KATO-3 tumors transplanted in NOD SCID mice every 3 days. As shown in Fig. 4, HSP105 siRNA suppressed tumor growth significantly in comparison to the irrelevant siRNA ($P < 0.01$). On day 15 after the first injection of HSP105 siRNA into tumors, the volumes of the tumors remained almost the same as those on day 0. During this observation period, neither abnormal behaviors nor neurological abnormalities were observed in these HSP105 siRNA-treated mice, and the expression of HSP105 protein in the brain was not suppressed by immunohistochemical analysis (data not shown).

Synergistic effects of siRNA and doxorubicin or heat shock on the *in vitro* induction of apoptosis in tumor cells

For the treatment of cancer patients with advanced, unresectable or recurrent focus, chemotherapy, radiation and other therapies are applied singly or in combination. To investigate the feasibility of combined treatment of tumor cells with HSP105 siRNA and other treatments, we treated HCT116 cells with HSP105 siRNA and either heat stress or doxorubicin. At 12 h after siRNA transfection into HCT116 cells, we added 200 ng/mL doxorubicin or treated the cells with heat shock at 45°C for 30 min, and detected the presence of apoptotic cells by staining with annexin V after 36 or 24 h incubation, respectively. As shown in Fig. 5, both combined treatments synergistically induced HCT116 cells to undergo apoptosis in comparison with the single treatment ($P < 0.001$).

Apoptosis induced by HSP105 siRNA was dependent on the caspase cascade but not on the p53 pathway

To investigate whether caspases are involved in apoptosis induced with siRNA, we examined PARP cleavage by western blot analysis using HCT116 cells with wild-type p53. PARP, a nuclear enzyme involved in DNA repair, is a well-known substrate for caspase-3, and is cleaved from a 112-kDa protein

to an 85-kDa protein by the activation of caspase-3. As shown in Fig. 6A, cleavage of PARP was observed in those cells transfected with HSP105 siRNA. Moreover, this cleavage was completely inhibited by adding a pan-caspase inhibitor, Z-VAD-FMK. As a result, the apoptosis induced by HSP105 siRNA was also inhibited by Z-VAD-FMK.

We next examined whether the p53 tumor suppressor protein is involved in HSP105 siRNA-induced apoptosis. The p53 protein is located upstream of the caspase cascade and is associated with heat shock proteins such as HSP70 and HSP90.⁽²⁴⁾ As shown in Fig. 6C, the HSP105 and p53 proteins were coimmunoprecipitated with anti-HSP105 antibodies and the DO-1 in non-treated HCT116 cells. These results indicate that a proportion of HSP105 protein is bound to p53 protein in non-treated HCT116 cells. Furthermore, the expression of p53 protein decreased with HSP105 siRNA treatment at the post-transcriptional level (the mRNA expression of p53 was not suppressed), and p53 protein was not phosphorylated at serine 15 or serine 46 by this treatment (Fig. 6D,E). Furthermore, to confirm the suppression of p53 transcriptional activity, the mRNA expression of Bax, NOXA and PUMA, which are the transcriptional targets of p53-mediated apoptosis, were assessed by RT-PCR and found to be suppressed (Fig. 6E). These data suggest that HSP105 protein is thus associated with wild-type p53 protein in HCT116 cells under non-stress conditions, and HSP105 siRNA-induced apoptosis is dependent on caspases but independent of the p53-mediated apoptosis pathway.

Discussion

In the present study, we obtained the following results: (1) HSP105 protein protects tumor cells from apoptosis; (2) constitutive overexpression of HSP105 protein is essential for the survival of various kinds of cancer cells; and (3) apoptosis induced with HSP105 siRNA treatment is dependent on caspases but not p53.

Recent studies, including ours, have shown that HSP105 is overexpressed in various human tumors and that HSP105 is thus speculated to be involved in both tumorigenesis and protection of cells from apoptosis.^(6,14,15) Our data obtained using NIH3T3-HSP105 cells are consistent with the findings of a recent study on neuronal PC12 cells in which the overexpression of HSP105 did not affect the growth rate of PC12 cells, but the apoptosis induced by stressors was inhibited in those cells.⁽⁶⁾ These data indicate that HSP105 is involved in tumorigenesis through protection of cells against apoptosis.

Among the heat shock proteins, HSP70 is well characterized and HSP105 shares functional properties with HSP70. HSP70 also inhibits apoptosis induced by various stimuli.^(25–27) Furthermore, HSP70 is also overexpressed in human breast cancer.^(28–31) One difference in function between HSP70 and HSP105 is that HSP105 has an increased capacity to bind to denatured polypeptides in comparison to HSP70.⁽³²⁾ In addition, HSP105 suppresses the aggregation of denatured proteins under stress conditions in the presence of ADP, whereas HSP70 suppresses it in the presence of ATP.⁽⁷⁾ In breast cancer cell lines, inhibition of HSP70 expression by antisense cDNA causes those cells to undergo apoptosis, and this action is




Fig. 2. Small interfering RNA (siRNA)-mediated inhibition of HSP105 expression enhanced the apoptotic cell death of human cancer cell lines. (A) Reverse transcription-polymerase chain reaction analysis of HSP105 mRNA expression in normal human colon epithelium, and in human cancer cell lines HCT116 and SW620. (B) Light microscopic pictures of HCT116 cells introduced with or without siRNA and representative flow cytometric analysis data of apoptotic cells stained with annexin V at 48 h after transfection. (C,D) Western blot analysis of HSP105 protein expression and flow cytometric analysis of apoptotic cells. HCT116 cells and SW620 cells were treated with oligofectamine, control siRNA, HSP105-siRNA (100 nM or 200 nM) or HSP105-siRNA-2 (B,C). (C) For western blot analysis, the cells were lysed at 24 or 48 h after transfection and analyzed. β -Actin is shown as a quantitative control. For flow cytometric analysis, cells were harvested at 24 or 48 h after transfection and then stained with fluorescein-isothiocyanate-annexin V and analyzed by flow cytometry. (D) Western blot analysis of HSP105 protein expression in cancer cell lines including SK-Hep1, PK8, KATO-3 and MKN28, and flow cytometric analysis of apoptotic cells at 48 h after transfection with 100 nM green fluorescent protein siRNA (\square) or HSP105 siRNA (\blacksquare). Data are the mean of three independent experiments \pm SD. The asterisks indicate that the differences in the percentages of annexin V-positive cells are statistically significant between the two values indicated by lines (* P < 0.01; ** P < 0.001).




Fig. 3. No apoptosis-inducing effects of HSP105 small interfering RNA (siRNA) on human fibroblasts. (A) Western blot analysis of HSP105. The lysates of human fibroblasts Turu and Mori, and human cancer cell lines HCT116, SW620, SK-Hep1, PK8, KATO-3 and MKN28 were used and blotted with an anti-HSP105 antibody. β -Actin is shown as a quantitative control. (B) Light microscopic pictures of Turu at 72 h after transfection with 100 nM green fluorescent protein (GFP) siRNA or HSP105 siRNA. (C) Effects of siRNA on Turu and Mori. Western blot analysis of HSP105 and flow cytometric analysis of apoptotic cells detected by annexin V staining at 72 h after transfection of 100 nM GFP siRNA or HSP105 siRNA. These data are representative of at least three independent experiments. The percentages shown in the panel indicate percentage of annexin V-positive cells in HSP105 siRNA-treated cells.

independent of caspases.⁽³³⁾ In the present study, transfection of HSP105 siRNA caused HCT116 cells to undergo apoptosis in a caspase-dependent manner without suppressing the expression of HSP70 protein (data not shown). Our data suggest that HSP105 has a different character regarding caspase dependency in comparison to HSP70.

In the present study, HSP105 siRNA transfection induced various cancer cell lines to undergo apoptosis. These observations raise the question of how such apoptosis is induced. One possible explanation is that suppression of HSP105 activates the apoptotic pathway mediated by the p53 tumor suppressor protein. Molecular chaperones such as HSP70 and HSP90 are overexpressed in various tumor cells,⁽³⁴⁾ associating with wild-type or mutated p53 tumor suppressor proteins. Such heat shock proteins mediate stabilization, cytoplasmic sequestration and localization of p53 proteins.^(35,36) In our study, HSP105 protein was bound to wild-type p53 in HCT116 cells under non-stress conditions, as shown in Fig. 6B. However, when the expression of HSP105 protein was suppressed with HSP105 siRNA, the expression of p53 protein also decreased and the p53-mediated apoptotic pathway was not activated (Fig. 6C,D). These results suggest that HSP105 stabilizes the p53 protein

and protects it from degradation, but the apoptosis induced by HSP105 siRNA treatment is not mediated by the p53-dependent apoptotic pathway. We herein observed that every cancer cell line with wild-type p53 (HCT116), mutated p53 (SW620) or without p53 (KATO-3) was induced to undergo apoptosis. Further studies are needed to elucidate the biological significance of the interaction between HSP105 and p53 protein in cancer cells.

The second possible mechanism of cancer cell apoptosis induced with HSP105 siRNA treatment is the involvement of ER stress. Heat shock proteins have housekeeping functions, such as folding and degradation of various proteins. ER stress, induced by the accumulation of unfolded or misfolded proteins, induces the unfolded protein response, characterized by the induction of chaperones, the translation block and ER-associated degradation. However, if such degradation is not sufficient, then prolonged ER stress activates various apoptotic pathways, including caspase activation.⁽³⁷⁻³⁹⁾

Abnormal protein aggregation has been suspected to cause many neurodegenerative diseases, including Alzheimer's disease, Parkinson's disease and trinucleotide repeat disease. In the brain in Alzheimer's disease, HSP90 facilitates the clearance




Fig. 4. The inhibitory effect of HSP105 small interfering RNA (siRNA) on the growth of established tumors in mice. (A) KATO-3 cells (2×10^6) were implanted subcutaneously into the dorsal skin of NOD SCID mice to establish growing tumors, and siRNA was injected into the tumors every 3 days (indicated by arrows). The tumor volume was measured and plotted (Luciferase siRNA, \blacktriangle , HSP105-siRNA, \blacksquare). Data are mean \pm SD ($n = 4$). The asterisk indicates that the difference in the tumor volume on day 15 is statistically significant between the two values as indicated by lines ($P < 0.01$).

of amyloid-beta.⁽⁴⁰⁾ In our study, HSP105 siRNA treatment induced HCT116 cells to undergo apoptosis through caspase activation. Considering these findings, we speculate that HSP105 siRNA treatment may induce aggregation of unfolded protein while also causing insufficient protein degradation, consequently leading to ER stress-mediated apoptosis, especially in cancer cells carrying mutations and aberrant expression of oncoproteins. Recent reports demonstrating that HSP105 prevents the aggregation of thermal-denatured protein *in vitro*⁽⁶⁾ and that overexpression of HSP105 suppresses aggregation and cell toxicity in a spinal and bulbar muscular atrophy model⁽⁹⁾ support our speculation. Regarding caspase dependency, the cleavage of PARP has been reported to be suppressed in PC12 cells overexpressing HSP105 protein and those cells were also protected from apoptosis caused by several stressors.⁽⁸⁾ These observations are consistent with our results.

Cancer cells often have aberrantly expressed or mutated genes that lead to uncontrolled cell growth and the prevention of apoptosis, and the usage of siRNA against such targets is thus considered to be promising for cancer therapy. Several recent studies have demonstrated the effective silencing by siRNA of targets, such as β -catenin for colon cancer,⁽⁴¹⁾ mutated K-ras for pancreatic carcinoma⁽⁴²⁾ and BCR/ABL fusion protein for CML.⁽⁴³⁾ In those reports, the injection of siRNA either induced target cells to undergo apoptosis or caused inhibition of their proliferation. In the present study, HSP105 siRNA





Fig. 5. The synergistic effect of HSP105 small interfering RNA (siRNA) with doxorubicin or heat shock regarding the induction of apoptotic cell death in HCT116 cells. At 12 h after transfection with 100 nM siRNA, the cells were incubated with 200 ng/mL doxorubicin (A) or treated with heat shock at 45°C for 30 min (B). Subsequently, the cells were stained with fluorescein-isothiocyanate-annexin V and analyzed by flow cytometry. Data are the mean of three independent experiments \pm SD ($n = 3$). The asterisks indicate that the differences in the percentages of annexin V-positive cells are statistically significant between the two values as indicated by lines ($P < 0.001$).

induced various human cancer cell lines to undergo apoptosis both *in vitro* and *in vivo* without side effects. It is notable that the effect of HSP105 siRNA treatment was transient and not lethal in normal fibroblasts, whereas the effects of known chemical agents tend to be cytotoxic for normal cells. Indeed, human fibroblast cells treated with doxorubicin were induced to undergo apoptosis (data not shown). These data suggest that HSP105 siRNA treatment is useful for cancer therapy and it may thus be applied to various kinds of cancer patients with minimal side effects.

For patients with advanced or metastatic cancer, combination therapies using some cytotoxic agents and radiation are now often performed clinically. We expected synergistic effects of combination therapy using HSP105 siRNA and doxorubicin, which have different mechanisms of action. siRNA suppresses the expression of targeted proteins by RNA cleavage, whereas doxorubicin a DNA intercalating agent that induces apoptosis by damaging DNA. In the present study, treatment combining HSP105 siRNA with doxorubicin synergistically induced cancer cells to undergo apoptosis. We also suspected that heat shock is effective when it is combined with HSP105 siRNA because HSP105 is essential in order to protect cells from heat stress.⁽⁸⁾

Fig. 6. Caspase-dependent and p53-independent induction of apoptosis in HCT116 cells administered HSP105 small interfering RNA (siRNA). (A) Western blot analysis of poly ADP-ribose polymerase (PARP) expression and (B) flow cytometric analysis of apoptosis induced in HCT116 cells transfected with siRNA in the presence of dimethylsulfoxide (DMSO) or Z-VAD-FMK. HCT116 cells treated with luciferase siRNA + DMSO, HSP105 siRNA + DMSO or HSP105 siRNA + 100 μ M Z-VAD-FMK were cultured for 48 h and apoptotic cells were stained with annexin V. Cells were lysed and blotted with either anti-HSP105 or anti-PARP antibody. Data are the mean values of three independent experiments \pm SD. The asterisks indicate that the differences in the percentages of annexin V-positive cells were statistically significant between the two values as indicated by lines ($P < 0.001$). (C,D) Western blot analysis of HSP105 and p53. HCT116 cells were lysed and immunoprecipitated with an anti-HSP105 antibody or an anti-p53 antibody (DO-1), and the proteins were blotted with either anti-HSP105 antibody or a biotin-labeled DO-1. The immunoprecipitates with rabbit polyclonal IgG and mouse monoclonal IgG2a were used as negative controls for anti-HSP105 antibody and DO-1, respectively (C). HCT116 cells transfected with luciferase siRNA or HSP105 siRNA were lysed at 48 h after transfection and blotted with the anti-HSP105 antibody DO-1, anti-phospho-p53 (Ser46) and antiphospho-p53 (Ser15). Ultraviolet light-irradiated HCT116 cell lysates were used as a positive control for p53 phosphorylation. (D) β -Actin is shown as a quantitative control. (E) Reverse transcription-polymerase chain reaction (RT-PCR) analysis of HSP105, p53, Bax, NOXA and PUMA expression in HCT116 cells transfected with siRNA. HCT116 cells transfected with luciferase siRNA or HSP105 siRNA were harvested at 24 h after transfection and the cDNAs were used for PCR analysis. cDNA extracted from the untreated HCT116 cells was used as a negative control. β -Actin is shown as a quantitative control.



As shown in Fig. 5B, the combination of HSP105 siRNA with heat shock, which is clinically applied to cancer patients as hyperthermia, exhibited a synergistic apoptotic effect in cancer cells.

In conclusion, our findings suggest that HSP105 is involved in tumorigenesis by protecting cancer cells from apoptosis, and the constitutive overexpression of HSP105 protein was

found to be essential for various cancer cells to survive. We also suggest that the apoptosis-inducing effect of HSP105 siRNA is specific for cancer, therefore HSP105 siRNA may be useful as a novel therapeutic tool for patients with cancers originating from various tissues. By using effective drug delivery systems and combining this treatment with existing cytotoxic agents, an enhanced effect is thus expected.

Acknowledgments

This work was supported in part by Grants-in-Aid 12213111 (Y. Nishimura) and 14770142 (T. Nakatsura) from the Ministry of Education, Science, Technology, Sports and Culture,

Japan. We thank Drs S. Senju, M. Monji and Y. Yoshitake (Department of Immunogenetics, Graduate School of Medical Science, Kumamoto University) for their helpful advice, and Dr B. Vogelstein, Johns Hopkins University, for providing the cell line HCT116.

References

- 1 Lee-Yoon D, Easton D, Murawski M, Burd R, Subjeck JR. Identification of a major subfamily of large hsp70-like proteins through the cloning of the mammalian 110-kDa heat shock protein. *J Biol Chem* 1995; **270**: 15 725–33.
- 2 Yasuda K, Nakai A, Hatayama T, Nagata K. Cloning and expression of murine high molecular mass heat shock proteins, HSP105. *J Biol Chem* 1995; **270**: 29 718–23.
- 3 Wakatsuki T, Hatayama T. Characteristic expression of 105-kDa heat shock protein (HSP105) in various tissues of non-stressed and heat-stressed rats. *Biol Pharm Bull* 1998; **21**: 905–10.
- 4 Satoh J, Yukitake M, Kuroda Y. Constitutive and heat-inducible expression of HSP105 in neurons and glial cells in culture. *Neuroreport* 1998; **9**: 2977–83.
- 5 Ishihara K, Yasuda K, Hatayama T. Molecular cloning, expression and localization of human 105 kDa heat shock protein, hsp105. *Biochim Biophys Acta* 1999; **1444**: 138–42.
- 6 Yamagishi N, Nishihori H, Ishihara K, Ohtsuka K, Hatayama T. Modulation of the chaperone activities of Hsc70/Hsp40 by Hsp105 α and Hsp105 β . *Biochem Biophys Res Commun* 2000; **272**: 850–5.
- 7 Yamagishi N, Ishihara K, Saito Y, Hatayama T. Hsp105 but not Hsp70 family proteins suppress the aggregation of heat-denatured protein in the presence of ADP. *FEBS Lett* 2003; **555**: 390–6.
- 8 Hatayama T, Yamagishi N, Minobe E, Sakai K. Role of hsp105 in protection against stress-induced apoptosis in neuronal PC12 cells. *Biochem Biophys Res Commun* 2001; **288**: 528–34.
- 9 Ishihara K, Yamagishi N, Saito Y *et al.* Hsp105 α suppresses the aggregation of truncated androgen receptor with expanded CAG repeats and cell toxicity. *J Biol Chem* 2003; **278**: 25 143–50.
- 10 Hatayama T, Takigawa T, Takeuchi S, Shiota K. Characteristic expression of high molecular mass heat shock protein HSP105 during mouse embryo development. *Cell Struct Funct* 1997; **22**: 517–25.
- 11 Evrard L, Vanmuylder N, Dourov N, Glineur R, Louryan S. Cytochemical identification of HSP110 during early mouse facial development. *J Craniofac Genet Dev Biol* 1999; **19**: 24–32.
- 12 Yamagishi N, Ishihara K, Saito Y, Hatayama T. Hsp105 α enhances stress-induced apoptosis but not necrosis in mouse embryonal F9 cells. *J Biochem (Tokyo)* 2002; **132**: 271–8.
- 13 Yamagishi N, Saito Y, Ishihara K, Hatayama T. Enhancement of oxidative stress-induced apoptosis by Hsp105 α in mouse embryonal F9 cells. *Eur J Biochem* 2002; **269**: 4143–51.
- 14 Nakatsura T, Senju S, Yamada K, Jotsuka T, Ogawa M, Nishimura Y. Gene cloning of immunogenic antigens overexpressed in pancreatic cancer. *Biochem Biophys Res Commun* 2001; **281**: 936–44.
- 15 Kai M, Nakatsura T, Egami H, Senju S, Nishimura Y, Ogawa M. Heat shock protein 105 is overexpressed in a variety of human tumors. *Oncol Rep* 2003; **10**: 1777–82.
- 16 Hwang TS, Han HS, Choi HK *et al.* Differential, stage-dependent expression of Hsp70, Hsp110 and Bcl-2 in colorectal cancer. *J Gastroenterol Hepatol* 2003; **18**: 690–700.
- 17 Yang G, Thompson JA, Fang B, Liu J. Silencing of H-ras gene expression by retrovirus-mediated siRNA decreases transformation efficiency and tumor growth in a model of human ovarian cancer. *Oncogene* 2003; **22**: 5694–701.
- 18 Gaggar A, Shayakhmetov DM, Lieber A. CD46 is a cellular receptor for group B adenoviruses. *Nat Med* 2003; **9**: 1408–12.
- 19 Wilson JA, Jayasena S, Khvorova A *et al.* RNA interference blocks gene expression and RNA synthesis from hepatitis C replicons propagated in human liver cells. *Proc Natl Acad Sci USA* 2003; **100**: 2783–8.
- 20 Takei Y, Kadomatsu K, Yuzawa Y, Matsuo S, Muramatsu T. A small interfering RNA targeting vascular endothelial growth factor as cancer therapeutics. *Cancer Res* 2004; **64**: 3365–70.
- 21 Chae SS, Paik JH, Fumeaux H, Hla T. Requirement for sphingosine 1-phosphate receptor-1 in tumor angiogenesis demonstrated by *in vivo* RNA interference. *J Clin Invest* 2004; **114**: 1082–9.
- 22 Monji M, Senju S, Nakatsura T *et al.* Head and neck cancer antigens recognized by the humoral immune system. *Biochem Biophys Res Commun* 2002; **294**: 734–41.
- 23 Nogawa M, Yuasa T, Kimura S *et al.* Intravesical administration of small interfering RNA targeting PLK-1 successfully prevents the growth of bladder cancer. *J Clin Invest* 2005; **115**: 978–85.
- 24 King FW, Wawrzynow A, Höhfeld J, Zylicz M. Co-chaperones Bag-1, Hop and Hsp40 regulate Hsc70 and Hsp90 interactions with wild-type or mutant p53. *EMBO J* 2001; **20**: 6297–305.
- 25 Jäättelä M, Wissing D, Bauer PA, Li GC. Major heat shock protein hsp70 protects tumor cells from tumor necrosis factor cytotoxicity. *EMBO J* 1992; **11**: 3507–12.
- 26 Mosser DD, Caron AW, Bourget L, Denis-Larose C, Massie B. Role of the human heat shock protein hsp70 in protection against stress-induced apoptosis. *Mol Cell Biol* 1997; **17**: 5317–27.
- 27 Jäättelä M, Wissing D, Kokholm K, Kallunki T, Egeblad M. Hsp70 exerts its anti-apoptotic function downstream of caspase-3-like proteases. *EMBO J* 1998; **17**: 6124–34.
- 28 Ciocca DR, Clark GM, Tandon AK, Fuqua SA, Welch WJ, McGuire WL. Heat shock protein hsp70 in patients with axillary lymph node-negative breast cancer: prognostic implications. *J Natl Cancer Inst* 1993; **85**: 570–4.
- 29 Lazaris A, Chatzigianni EB, Panoussopoulos D, Tzimas GN, Davaris PS, Golenatis B. Proliferating cell nuclear antigen and heat shock protein 70 immunolocalization in invasive ductal breast cancer not otherwise specified. *Breast Cancer Res Treat* 1997; **43**: 43–51.
- 30 Vargas-Roig LM, Fanelli MA, López LA *et al.* Heat shock proteins and cell proliferation in human breast cancer biopsy samples. *Cancer Detect Prev* 1997; **21**: 441–51.
- 31 Vargas-Roig LM, Gago FE, Tello O, Aznar JC, Ciocca DR. Heat shock protein expression and drug resistance in breast cancer patients treated with induction chemotherapy. *Int J Cancer* 1998; **79**: 468–75.
- 32 Oh HJ, Chen X, Subjeck JR. Hsp110 protects heat-denatured proteins and confers cellular thermoresistance. *J Biol Chem* 1997; **272**: 31 636–40.
- 33 Nylandsted J, Rohde M, Brand K, Bastholm L, Elling F, Jäättelä M. Selective depletion of heat shock protein 70 (Hsp70) activates a tumor-specific death program that is independent of caspases and bypasses Bcl-2. *Proc Natl Acad Sci USA* 2000; **97**: 7871–6.
- 34 Helmbrecht K, Zeise E, Rensing L. Chaperones in cell cycle regulation and mitogenic signal transduction: a review. *Cell Prolif* 2000; **33**: 341–65.
- 35 Walerych D, Kudla G, Gutkowska M *et al.* Hsp90 chaperones wild-type p53 tumor suppressor protein. *J Biol Chem* 2004; **279**: 48 836–45.
- 36 Whitesell L, Sutphin PD, Pulcini EJ, Martinez JD, Cook PH. The physical association of multiple molecular chaperone proteins with mutant p53 is altered by geldanamycin, an hsp90-binding agent. *Mol Cell Biol* 1998; **18**: 1517–24.
- 37 Welihinda AA, Tirasophon W, Kaufman RJ. The cellular response to protein misfolding in the endoplasmic reticulum. *Gene Expr* 1999; **7**: 293–300.
- 38 Nakagawa T, Zhu H, Morishima N *et al.* Caspase-12 mediates endoplasmic reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 2000; **403**: 98–103.
- 39 Momoi T. Caspases involved in ER stress-mediated cell death. *J Chem Neuroanat* 2004; **28**: 101–5.
- 40 Kakimura J, Kitamura Y, Takata K *et al.* Microglial activation and amyloid-beta clearance induced by exogenous heat-shock proteins. *FASEB J* 2002; **16**: 601–3.
- 41 Verma UN, Surabhi RM, Schmalting A, Becerra C, Gaynor RB. Small interfering RNAs directed against β -catenin inhibit the *in vitro* and *in vivo* growth of colon cancer cells. *Clin Cancer Res* 2003; **9**: 1291–300.
- 42 Brummelkamp TR, Bernards R, Agami R. Stable suppression of tumorigenicity by virus-mediated RNA interference. *Cancer Cell* 2002; **2**: 243–7.
- 43 Wilda M, Fuchs U, Wössmann W, Borkhardt A. Killing of leukemic cells with a BCR/ABL fusion gene by RNA interference (RNAi). *Oncogene* 2002; **21**: 5716–24.

Immunization with heat shock protein 105-pulsed dendritic cells leads to tumor rejection in mice [☆]

Kazunori Yokomine ^{a,b}, Tetsuya Nakatsura ^{a,*,1}, Motozumi Minohara ^c, Jun-ichi Kira ^c, Tatsuko Kubo ^d, Yutaka Sasaki ^b, Yasuharu Nishimura ^{a,*}

^a Department of Immunogenetics, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

^b Department of Gastroenterology and Hepatology, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

^c Department of Neurology, Neurological Institute, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

^d Department of Molecular Pathology, Graduate School of Medical Sciences, Kumamoto University, Kumamoto, Japan

Received 18 February 2006

Available online 3 March 2006

Abstract

Recently, we reported that heat shock protein 105 (HSP105) DNA vaccination induced anti-tumor immunity. In this study, we set up a preclinical study to investigate the usefulness of dendritic cells (DCs) pulsed with mouse HSP105 as a whole protein for cancer immunotherapy *in vivo*. The recombinant HSP105 did not induce DC maturation, and the mice vaccinated with HSP105-pulsed BM-DCs were markedly prevented from the growth of subcutaneous tumors, accompanied with a massive infiltration of both CD4⁺ T cells and CD8⁺ T cells into the tumors. In depletion experiments, we proved that both CD4⁺ T cells and CD8⁺ T cells play a crucial role in anti-tumor immunity. Both CD4⁺ T cells and CD8⁺ T cells specific to HSP105 were induced by stimulation with HSP105-pulsed DCs. As a result, vaccination of mice with BM-DCs pulsed with HSP105 itself could elicit a stronger tumor rejection in comparison to DNA vaccination. © 2006 Elsevier Inc. All rights reserved.

Keywords: Heat shock protein 105; Cancer antigen; Dendritic cells; Th; CTL

Heat shock proteins (HSPs) are soluble intracellular proteins, which are ubiquitously expressed, and their expression can be induced at much higher levels as a result of heat shock or other forms of stress. HSPs have essential functions in the regulation of protein folding, conformation, assembly, and sorting. HSPs have been shown to be molecular chaperones that function to maintain the native

conformational states of proteins and prevent protein–protein aggregation [1]. HSPs can also induce the response of antigen-specific effector CD8⁺ T cells which can protect hosts from both infection and tumor challenge [2]. Srivastava and co-workers [3,4] led to a proposal that the tumor-derived HSP–peptide complex elicits a protective immunity that is specific to a particular cancer, while HSPs derived from normal tissues did not elicit any protective immunity to the cancers tested. Immunotherapeutic clinical trials targeted at autologous tumor-derived gp96–peptide complexes are still ongoing in metastatic melanoma and colorectal carcinoma patients [5].

Dendritic cells (DCs) are powerful antigen-presenting cells (APCs) that are considered to be potent immunotherapeutic agents to promote the host immune response against tumor antigen. DCs become efficient tumor vaccines when pulsed with synthetic or natural tumor-derived peptides, transduced with tumor-derived RNA or vectors

[☆] **Abbreviations:** BM-DC, bone marrow-derived DC; HSP105, heat shock protein 105; Th, helper T cell; CTL, cytotoxic T lymphocyte; MHC, major histocompatibility complex; C26 (C20), colon 26 clone 20.

* Corresponding authors. Fax: +81 96 373 5314 (Y. Nishimura); +81 4 7131 5490 (T. Nakatsura).

E-mail addresses: tnakatsu@east.ncc.go.jp (T. Nakatsura), mxnishim@gpo.kumamoto-u.ac.jp (Y. Nishimura).

¹ Present address: Immunotherapy Section, Investigative Treatment Division, Research Center for Innovative Oncology, National Cancer Center Hospital East, 6-5-1 Kashiwanoha, Kashiwa City, Chiba 277-8577, Japan.

encoding for tumor-associated proteins, or directly fused to or incubated with tumor cells [6]. For effective induction of cytotoxic T lymphocytes (CTLs) by vaccination, “Cross-presentation” mediated by DCs often plays an important role. Such cross-presentation includes the antigen presentation of exogenous antigens by major histocompatibility complex (MHC) class I molecules as well as by MHC class II molecules [7,8]. HSP-chaperoned peptides were cross-presented by the MHC class I molecules of the DCs several 100-fold more efficiently than unchaperoned peptides [9]. In addition, CD91, also called α_2 -macroglobulin receptor is expressed on DCs and has been shown to act as one of the receptors for HSP-chaperoned peptides to efficiently incorporate the HSP-peptide complexes [10].

We earlier reported that heat shock protein 105 (HSP105) was overexpressed in a variety of human cancers but it is not expressed in normal tissue except for the testes [11,12], thus suggesting that HSP105 itself may be a potential candidate as a target antigen for cancer immunotherapy. The amino acid sequences and expression patterns of HSP105 are very similar between humans and mice. HSP105 has been found to be immunogenic in mice and an effective anti-tumor immunity has been observed after *HSP105* DNA vaccination [13]. In the present study, we set up a preclinical study to investigate the usefulness of HSP105 as a target for cancer immunotherapy using DCs. It has been reported that HSPs can induce DC maturation and activation as determined by the upregulation of MHC class II and CD86 molecules, the secretion of IL-12 and TNF α [14,15], and migration into draining lymphoid organs [16]. On the contrary, some investigators reported that HSP-mediated maturation of DCs was caused by contaminating lipopolysaccharide (LPS) fraction because endotoxin-free HSP70 failed to induce DC maturation [17]. We herein show that the highly purified HSP105 did not induce DC maturation and that the immunization of HSP105-pulsed DC led to the tumor rejection of melanoma and colorectal cancer in mice. These findings suggested that HSP105 itself could be a valuable tumor-associated antigen applicable for DC-based immunotherapy of tumors overexpressing it.

Materials and methods

Cell lines and mice. A subline of BALB/c-derived colorectal cancer cell line Colon 26, C26 (C20), was provided by Dr. Kyoichi Shimomura (Astellas Pharmaceutical Co., Tsukuba, Japan). Other cancer cell lines were kindly provided by the Cell Resource Center for Biomedical Research Institute of Development, Aging, and Cancer, Tohoku University (Sendai, Japan). All these cell lines were cultured in RPMI 1640 medium supplemented with 10% heat-inactivated fetal calf serum at 37 °C in a humidified 5% CO₂ atmosphere. We used the B16-F10 melanoma cell line syngeneic to C57BL/6 mice and C26 (C20) for the tumor challenge. Female 6- to 8-week-old C57BL/6 mice (H-2^b) and BALB/c mice (H-2^d) were purchased from Charles River Japan (Yokohama, Japan). These mice were kept under specific pathogen-free conditions. These experiments were approved by the Animal Research Committee of Kumamoto University.

Production of recombinant proteins. We produced highly purified recombinant mouse HSP105 from the *Escherichia coli* strain BL21 cells transduced with the mouse *HSP105* gene expression vector, as described previously [18]. Purified proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and Coomassie brilliant blue (CBB)-stained bands were quantified by densitometry. Thereafter, by using affinity chromatography on a polymyxin B agarose gel (Sigma Chemical Co., St. Louis, MO), the endotoxin levels were decreased. We also produced highly purified recombinant myelin basic protein (MBP) as described previously [19]. Both recombinant HSP105 and MBP were estimated to be almost endotoxin free by using Limulus amoebocyte lysate assay kit (BioWhittaker, Walkersville, MD), and endotoxin contents in these two materials were below 10 endotoxin U/mg.

Immunizations and tumor challenge. Bone marrow-derived DCs (BM-DC) were prepared as described previously [20]. BM-DCs were pulsed with 2 μ g/ml HSP105 at 37 °C for 16 h, non-adherent and loosely adherent proliferating DCs were collected and used as HSP105-pulsed BM-DC. In tumor prevention experiments, mice were intraperitoneally inoculated with HSP105-pulsed BM-DC (5×10^5) suspended in 200 μ l PBS on days -14 and -7. In parallel, groups of mice were injected with BM-DC alone or PBS as controls. Tumor challenge was initiated by subcutaneous injection with B16-F10 cells (1×10^4) or C26 (C20) cells (3×10^4) suspended in 100 μ l HBSS (Gibco, Grand Island, NY) in shaved right flanks on day 0. Tumor occurrence was observed twice a week. The tumor size was evaluated by measuring two perpendicular diameters using calipers.

Flow cytometric analysis. Staining of cells and analysis on a flow cytometer (FACScan; BD Biosciences) were done as described previously [21]. Antibodies and reagents used for staining were as follows: FITC-conjugated anti-I-A^b (clone 28-16-8S; mouse IgG2a; Caltag, Burlingame, CA), R-PE-conjugated anti-mouse CD80 (clone RMMP-1; rat IgG2a; Caltag), R-PE-conjugated anti-mouse CD86 (clone RMMP-2; rat IgG2a; Caltag), FITC-conjugated anti-mouse CD4 (clone L3T4; rat IgG2a; BD PharMingen, San Diego, CA), FITC-conjugated anti-mouse CD8 (clone Ly-2; rat IgG2a; BD PharMingen), FITC-conjugated mouse IgG2a control (clone G155-178; BD PharMingen), and R-PE-conjugated rat IgG2a control (clone LO-DNP-16; Caltag).

Depletion of CD4⁺ T cells and CD8⁺ T cells in mice. Rat monoclonal antibodies (mAbs) GK1.5 specific to mouse CD4 and 2.43 specific to mouse CD8 were used to deplete CD4⁺ T cells and CD8⁺ T cells in vivo, respectively. The mice were injected with ascites (0.1 ml/mouse) from hybridoma-bearing nude mice intraperitoneally on days -18, -15, -11, -8, -4, and -1 and the tumor cells were inoculated on day 0. Normal rat IgG (Sigma, St. Louis, MO; 200 μ g/mouse) was used as a control. The depletion of T cell subsets was monitored by a flow cytometric analysis, which showed more than a 90% specific depletion in the number of splenocytes.

Immunohistochemical analysis. Immunohistochemical detection of HSP105 was done as previously described [11,12]. Rabbit polyclonal anti-human HSP105 (Santa Cruz, Santa Cruz, CA) was used as the primary antibody in this study. Immunohistochemical staining of CD4 and CD8 was done as previously described [22]. For the terminal deoxynucleotidyl transferase-mediated nick end labeling (TUNEL) method, we used ApopTag Fluorescein In Situ Apoptosis Detection Kits (Serologicals Corporation, Norcross, GA).

Induction of CD4⁺ T cells and CD8⁺ T cells specific to HSP105. The mice were inoculated intraperitoneally with HSP105-pulsed BM-DC on days -14 and -7. Spleen cells were harvested on day 0, and CD4⁺ T cells and CD8⁺ T cells were purified using the magnetic cell sorting system (MACS) with anti-mouse CD4 (L3T4) mAb and anti-mouse CD8 α (Ly-2) mAb, respectively. The purity of these T cell subsets exceeded 95% by a flow cytometric analysis. CD4⁺ T cells or CD8⁺ T cells (3×10^5 /well) were separately incubated in RPMI 1640 medium supplemented with 10% horse serum, IL-2 (100 U/ml), and 2-ME (50 μ M) together with the irradiated (4500 Gy) HSP105-pulsed BM-DC in 24-well culture plates. BM-DCs (3×10^4 /well) pulsed with 2 μ g/ml HSP105 for 16 h were irradiated (4500 Gy) and added to the culture wells for the restimulation once a week. After the third restimulation in vitro, both proliferation and cytotoxicity assays were performed as described previously [23]. For the

control of ^{51}Cr -release assay, CD8^+ T cells isolated from the mice immunized with BM-DCs alone were restimulated in vitro with BM-DCs alone once a week and used as effector cells.

ELISPOT assay. HSP105-specific IFN- γ production of T cells was quantified using the appropriate ELISPOT kit (BD Biosciences, San Diego, CA) according to the manufacturer's instructions. CD4^+ T cells or CD8^+ T cells were incubated with the BM-DC alone, BM-DCs pre-pulsed with HSP105, or BM-DCs pre-pulsed with myelin basic protein (MBP) as a control at 37 °C for 24 h. Each BM-DC was pre-pulsed with 2 $\mu\text{g}/\text{ml}$ protein at 37 °C for 16 h. The spots were automatically counted and subsequently analyzed using the Eliphoto system (MINERVA TECH, Tokyo, Japan).

Statistical analysis. The statistical significance of the differences in the findings between the experimental groups was determined by Student's *t* test. The overall survival rate was calculated using the Kaplan–Meier method, and statistical significance was evaluated using Wilcoxon's test. A value of $P < 0.05$ was considered to be statistically significant.

Results

HSP 105 does not induce maturation of DCs

To analyze the direct effect of HSP 105 used in this study on BM-DCs, BM-DCs were incubated with HSP105, LPS as a positive control, and left untreated for 16 h. As shown in Fig. 1, no significant difference was observed in the levels of surface expression of CD80, CD86, and I-A^b between untreated BM-DCs and HSP105-pulsed BM-DCs. Moreover, HSP105-pulsed BM-DCs microscopically did not show any morphological changes in comparison to the untreated BM-DC. On the contrary, LPS-pulsed BM-DCs exhibited markedly increased expression of these three molecules. Although it is reported that HSPs could induce

DC maturation and activation [14–16], the recombinant HSP105 used in this study including little LPS (below 10 endotoxin U/mg) did not show such activity. Thereafter, we evaluated the antigenicity of HSP105 to induce anti-tumor immunity.

The HSP105-pulsed BM-DC vaccine induced anti-tumor immunity against the lethal challenge of B16-F10 and C26 (C20)

We recently reported that mouse HSP105 was also over-expressed in the liver metastasis of C26 (C20) cells, and lung metastase of the B16-F10 cells, and that HSP105 DNA vaccination inhibited the growth of these tumors [13]. In this study, we investigated the effects of HSP105 vaccination based on DCs on the growth of B16-F10 and C26 (C20) tumor cells in vivo. The objective was to determine whether prophylactic vaccination induced significant immunity against tumor growth and a prolonged survival. The protocol of vaccination in this study is shown in Fig. 2A. The results shown in Fig. 2B demonstrate that immunization with HSP105-pulsed BM-DC markedly inhibited the growth of B16-F10 tumors in comparison to other groups ($P < 0.01$). As shown in Fig. 2C, five of eight (62.5%) mice immunized with HSP105-pulsed BM-DC remained tumor free and survived for 100 days after the tumor challenge. In contrast, the mice vaccinated with BM-DC alone (12.5%) or PBS (0%) showed little protection against the growth of B16-F10 tumor in comparison to the observations in mice treated with HSP105-pulsed




Fig. 1. Expression levels of cell surface I-A^b, CD80, and CD86 on BM-DCs, HSP105-pulsed BM-DCs, and LPS-pulsed BM-DCs were analyzed by flow cytometric analysis. BM-DCs were pulsed with 2 $\mu\text{g}/\text{ml}$ HSP105, 1 $\mu\text{g}/\text{ml}$ LPS or left untreated for 16 h. (A) The expression levels in HSP105-pulsed BM-DCs (filled histogram), LPS-pulsed BM-DCs (dotted line), and untreated BM-DCs (thick line), and the profiles of cells treated with isotype matched Ig as a negative control for staining (thin line). (B) The mean fluorescence intensity (MFI) of I-A^b, CD80, and CD86 staining in the cells. The results are representative of three independent experiments with similar results.




Fig. 2. Protection against tumor growth of B16-F10 and C26 (C20) cells by immunization with HSP105-pulsed BM-DC vaccine. (A) Protocol of the vaccination. The mice were immunized with PBS, BM-DC alone, and HSP105-pulsed BM-DC on 14 and 7 days before the tumor challenge. Seven days after the second immunization, the mice were challenged with B16-F10 cells s.c. (B,C), or C26 (C20) cells s.c. (D,E). (B,D) The tumor size was evaluated by measuring two perpendicular diameters. The result is presented as the mean area of tumor \pm SE, and we evaluated statistical significance of the differences between each group using the unpaired Student's *t* test. (C,E) The mice in each group were observed for their survival period. The statistical significance of the differences between each group was evaluated using Wilcoxon's test.

BM-DC ($P < 0.05$). Similar results were observed in a prophylactic immunotherapy model using C26 (C20). Four of five (80%) mice immunized with HSP105-pulsed BM-DC completely rejected the C26 (C20) (3×10^4) cells (Figs. 2D and E), whereas tumors grew rapidly and all five mice died within 70 days in control mice treated with PBS or BM-DC alone. These results suggest that the HSP105-pulsed BM-DC vaccine is a potent vaccine that can efficiently induce specific anti-tumor immunity.

Both CD4⁺ T cells and CD8⁺ T cells are required for anti-tumor immunity

To determine the role of CD4⁺ T cells and CD8⁺ T cells in the protection against B16-F10 and C26 (C20) tumor cells induced by HSP105 vaccination, we depleted mice of CD4⁺ T cells or CD8⁺ T cells by the treatment with anti-CD4 or anti-CD8 mAb *in vivo*, respectively. During the depletion procedure, the mice were immunized with HSP105-pulsed BM-DC vaccine and challenged with B16-F10 or C26 (C20) cells (Fig. 3A). In both B16-F10 and C26 (C20) models, mice depleted of CD4⁺ T cells, and CD8⁺ T cells developed aggressively growing tumors

after the challenge in comparison to the findings in control mice treated with rat IgG ($P < 0.05$) (Figs. 3B and D). The mice depleted of CD4⁺ T cells or CD8⁺ T cells all died by 52–65 days, whereas more than 50% of the control mice survived for 70 days ($P < 0.05$) (Figs. 3C and E). These results suggest that both CD4⁺ T cells and CD8⁺ T cells play crucial roles in the protective anti-tumor immunity induced by the HSP105-pulsed BM-DC vaccine.

Vaccination of HSP105-pulsed BM-DCs induced infiltrations of both CD4⁺ T cells and CD8⁺ T cells into tumor cells, but not into normal organs

Four of five (80%) mice immunized with the HSP105-pulsed BM-DCs completely rejected challenges of C26 (C20) cells (3×10^4) (Fig. 2E). To ascertain whether these rejections were induced by CD4⁺ T cells or CD8⁺ T cells, the subcutaneous inoculation of many C26 (C20) cells (1×10^6) into the right flank was done at 7 days after the second vaccination. After tumor formation, we removed the tumor and immunohistochemically stained it using anti-CD4 mAb, anti-CD8 mAb, and the TUNEL method. The infiltration of CD4⁺ T cells and CD8⁺ T cells into C26




Fig. 3. Both CD4⁺ and CD8⁺ T cells are involved in the antitumor immunity elicited by the HSP105-pulsed DC vaccine. (A) Protocol for the vaccination and the depletion of T cells. C57BL/6 mice and BALB/c mice were challenged s.c. with B16-F10 cells and C26 (C20) cells, respectively. (B,D) The tumor size was evaluated by measuring two perpendicular diameters. The result is presented as the mean area of tumor \pm SE, and we evaluated the statistical significance of the differences between each group using the unpaired Student's *t* test. (C,E) The mice in each group were observed for their survival period. The statistical significance of the differences between each group was evaluated using Wilcoxon's test.

(C20) tumors and some apoptotic C26 (C20) tumor cells were observed in the mice vaccinated with HSP105-pulsed BM-DCs, but never in the mice vaccinated with unpulsed BM-DCs (Fig. 4A). These results suggest that HSP105-pulsed BM-DCs have the potential to sensitize many HSP105-specific CD4⁺ T cells and CD8⁺ T cells to kill C26 (C20) tumor cells.

We evaluated the risk of autoimmunity by immunization against self-HSP105. Both BALB/c and C57BL/6 mice immunized with HSP105-pulsed BM-DC were apparently healthy without any abnormality such as dermatitis, arthritis, or neurological disorders. The tissues of the mice immunized with HSP105-pulsed BM-DC were histologically examined. The brain, liver, heart, kidneys, and spleen had normal structures and did not show any pathological changes suggestive of an immune response, such as the infiltration of CD4⁺ T cells and CD8⁺ T cells or tissue

destruction and repair. Although we used female mice for the experiments described above, we also immunized male mice with HSP105-pulsed BM-DC to ascertain whether immunization with HSP105-pulsed BM-DC induced autoimmunity in the testis in which HSP105 is strongly expressed. However, no sign of autoimmunity was observed in the testis (Fig. 4B).

Induction of HSP105-specific CD4⁺ T cells and CD8⁺ T cells by immunization with HSP105-pulsed BM-DC

CD4⁺ T cell lines specific to HSP105 were established from spleen cells derived from mice vaccinated with HSP105-pulsed BM-DC. CD4⁺ T cells were separated from spleen cells and the purity of these cells was more than 95% by flow cytometric analysis. These cells were restimulated with irradiated and HSP105-pulsed DCs once




Fig. 4. Vaccination of HSP105-pulsed BM-DCs induced infiltrations of both CD4⁺ T cells and CD8⁺ T cells into C26 (C20) tumor and induced the apoptosis of C26 (C20) tumor cells. (A) C26 tumors removed from the mice vaccinated with BM-DCs or HSP105-pulsed BM-DCs were analyzed using immunohistochemical staining with anti-CD4 mAb, anti-CD8 mAb, and the TUNEL method on 4 days after the inoculation of tumor cells (1×10^6). (B) Normal tissue specimens of mice vaccinated with HSP105-pulsed BM-DCs were examined histologically and immunohistochemically. Objective magnification was 200 \times . The spleen was used as a positive control for the staining of both CD4⁺ and CD8⁺ cells.

a week. After three restimulations, both an ELISPOT assay and a proliferation assay were performed. The ELISPOT assay showed that HSP105-sensitized CD4⁺ T cells produced IFN- γ in response to BM-DCs prepulsed with HSP105 but not an irrelevant MBP (Fig. 5A). As shown in Fig. 5B, HSP105-sensitized CD4⁺ T cells proliferated in the presence of BM-DCs prepulsed with HSP105 but not MBP. These observations clearly indicated that HSP105-specific CD4⁺ T cells were included in the T cell line.

We investigated whether HSP105-specific CD8⁺ T cells were also induced with HSP105-pulsed DC vaccination. CD8⁺ T cells were purified (>95%) from spleen cells of vaccinated mice and restimulated with irradiated and HSP105-pulsed DCs once a week. After three restimulations, the ELISPOT assay and 6 h ⁵¹Cr-release assay were performed to detect the HSP105-specific CTL responses (Figs. 5C and D). The CD8⁺ T cell line exhibited a HSP105-specific production of IFN- γ in an ELISPOT assay when cells were stimulated with BM-DCs prepulsed with HSP105 but not MBP ($P < 0.01$), however, the number of spots was smaller than that of CD4⁺ T cells (Fig. 5C). CD8⁺ T cells immunized with HSP 105-pulsed DC demonstrated a significant cytolytic activity against the B16-F10 cells pretreated with IFN- γ to induce the expression of MHC class I molecules on the cell surface, whereas CD8⁺ T cells from mice immunized with BM-DC alone revealed little cytolytic activity ($P < 0.005$) (Fig. 5D). The induction of HSP105-specific CD8⁺ T cells by the immunization in vivo with HSP105-

pulsed BM-DC and the stimulation of the CD8⁺ T cell line in vitro with the HSP105-pulsed BM-DC strongly suggested that these HSP105-specific CD8⁺ T cells were induced by the cross-presentation of HSP105 by BM-DCs.

Discussion

HSPs are classified into several families based on their apparent molecular weights, such as HSP105/110, HSP90, HSP70, HSP60, HSP40, and HSP27 [24]. HSP105 consists of HSP105 α and HSP105 β . HSP105 α is a constitutively expressed 105-kDa HSP that is induced by a variety of stresses, whereas HSP105 β is a 90-kDa truncated form of HSP105 α that is specifically induced by heat shock at 42 °C [24]. In this study, we used the mouse HSP105 α protein. The cDNA sequence of murine HSP105 is almost the same as that of the Chinese hamster HSP110 [25,26], so HSP105 belongs to a member of the HSP105/110 family. We recently reported by the immunohistochemical analysis that HSP105 is overexpressed in a variety of human tumors [12], the liver metastasis of the C26 (C20) cells in the BALB/c mice, and lung metastasis of the B16-F10 cells in the C57BL/6 mice [13]. We examined the expression of HSP105 in the mouse cancer cell lines using a Western blotting analysis and found that HSP105 was strongly expressed in all 7 mouse cell lines tested (data not shown).

Many studies have shown that certain HSPs purified from a tumor can function as an effective vaccine against the same




Fig. 5. Induction of HSP105-specific CD4⁺ T cells and CD8⁺ T cells by stimulation with HSP105-pulsed BM-DCs. (A) An ELISPOT assay for IFN- γ production by CD4⁺ T cell lines stimulated with HSP105 protein-pulsed BM-DCs. CD4⁺ T cells derived from the mice vaccinated with HSP105-pulsed BM-DC were stimulated in vitro with HSP105-pulsed BM-DC three times. For the ELISPOT assay, these CD4⁺ T cells were co-cultured with BM-DC prepulsed with HSP105, MBP, or unpulsed BM-DC for 24 h. (B) Cell proliferation of CD4⁺ T cell lines stimulated with HSP105-pulsed BM-DCs was determined by measuring [³H]thymidine incorporation. CD4⁺ T cells were co-cultured with BM-DC prepulsed with HSP105, MBP, or unpulsed BM-DC for 72 h. (C) An ELISPOT assay for IFN- γ production by CD8⁺ T cell lines stimulated with HSP105-pulsed BM-DCs. CD8⁺ T cells derived from mice vaccinated with HSP105-pulsed BM-DC were stimulated with HSP105-pulsed BM-DC three times in vitro. For the ELISPOT assay, these CD8⁺ T cells were co-cultured with BM-DC prepulsed with HSP105, MBP, or unpulsed BM-DC for 24 h. (D) CD8⁺ T cells stimulated with HSP105-pulsed BM-DC or BM-DC alone (control) were examined for their CTL activity against B16-F10 cells treated with IFN- γ (10^3 U/ml) using 6 h ⁵¹Cr-release assay. The results were analyzed using the mean values of a triplicate or a quadruplicate assay. The data shown in A–D are each representative of three independent experiments with similar results.

tumor by stimulating T cells with tumor-specific peptides bound to HSPs. Subjeck and co-workers [27,28] reported that tumor-derived HSP110-peptide complexes also stimulated tumor immunity as other HSP families did in mice. Despite studies establishing a chaperoning effect of HSPs, one impediment to the full-fledged acceptance of HSPs as peptide-transporting vehicles is the lack of mass spectrometric data directly identifying HSP-associated peptides [29]. Stress-inducible proteins can be recognized by natural killer cells and CTLs as whole antigens expressed on the surface of stressed cells in humans [30]. Proteins dramatically upregulated or modified under stressful conditions should lead to increased presentation as do peptides presented by HLA class I molecules. About 25 HSP-derived peptides bound by HLA class I molecules have been identified through mass spectrometry [30]. Cancer patients have been reported to possess CTLs specific to HSP60-derived peptide [31], while HLA-A*0201-restricted HSP70-derived CTL epitopes have been identified in both an HLA-A*0201 transgenic mouse model and in humans [32]. In this study, although we did

not identify HSP105-derived epitope peptides for CD4⁺ T cells and CD8⁺ T cells, we did prove that HSP105 itself could induce both CD4⁺ Th-cells and CD8⁺ CTLs specific to HSP105 as a cancer antigen. Contrary to our findings, however, Subjeck and co-workers [28] reported that HSP110 immunization did not elicit anti-tumor immunity. This discrepancy could be attributed to the difference in the methods of immunization.

It has been reported that HSPs can induce the maturation and activation of DCs as determined by upregulation of MHC class II and CD86 molecules, secretion of the IL-12 and TNF α [14,15]. However, HSP105-pulsed BM-DCs did not show any changes in comparison to the untreated BM-DC, thus suggesting that HSP105 did not induce DC maturation and activation. It is unlikely that HSP105 brought tumor-derived peptides into the culture system, because the HSP105 used in this study was the recombinant protein produced in *E. coli*. Furthermore, we recently identified HSP105-derived CTL epitopes restricted by HLA-A*0201 or -A*2402 using HLA

transgenic mouse model (unpublished data). These results also supported that HSP105 served not as a mediator for maturation of DCs, but as a cancer antigen eliciting tumor immunity.

The results of the T cell depletion study showed that the depletion of either CD4⁺ T cells or CD8⁺ T cells abrogated the anti-tumor immune response induced by the HSP 105-pulsed BM-DC vaccine, and that both CD4⁺ and CD8⁺ T cells play crucial roles in the protective anti-tumor immunity. CD8⁺ T cells are thought to serve as the dominant effector cell mediating tumor killing, in contrast, CD4⁺ T cells are thought to have an indirect role in providing help to CTL as well as a direct role in tumor rejection [33]. It is interesting that B16-F10 tumor cells that lack MHC class I were killed in *in vivo* study. We suppose that CD4⁺ T cells may have an important role in this case. Peptides derived from HSP105 bound by MHC class II on the surface of HSP105-pulsed BM-DCs activate CD4⁺ T cells. The activated CD4⁺ T cells can secrete IFN- γ upon stimulation with tumor local DCs presenting tumor-derived HSP105 peptides, which contribute not only to activation of CD8⁺ T cells but also to restoration of MHC class I expression in B16-F10 cells. The activated HSP105 specific CD8⁺ T cells can recognize the peptides derived from HSP105 in the context of MHC class I and kill the B16-F10 cells.

In the field of cancer immunotherapy, most enthusiasm has been directed toward the use of various cancer vaccines; peptide vaccines alone, peptide plus cytokines, vaccination either with recombinant virus or with naked DNA encoding tumor antigen, and peptide pulsed on DCs [34]. DCs represent the most potent antigen presenting cells and also play an important role in the induction of specific T cell response [35]. Peptides pulsed on DCs have been reported to be the most effective vaccine in comparison to DNA vaccine or peptide–adjuvant mixture [36]. In this study, 62.5% and 80.0% of the mice immunized with HSP105-pulsed BM-DC completely rejected B16-F10 cells and C26 (C20) cells, respectively. On the other hand, only 50.0% of the mice immunized with the HSP105-DNA vaccine rejected these tumor cells in our previous study [13]. Although a further comparative analysis of the vaccination properties of these two strategies is required, our results suggested that protein-pulsed DCs are a more powerful vaccine than the DNA vaccine.

In this study, we used BM-DCs pulsed with HSP105 but not with HSP105-derived peptide as a cancer vaccine. We think that protein-pulsed DCs thus have an advantage over peptide-pulsed DCs. DCs are the major cell type known for its capacity to cross-present antigens [37]. In this study, HSP105-sensitized CD8⁺ T cells responded to HSP105 *in vitro* by the stimulation of purified CD8⁺ T cells with HSP105-pulsed DCs. This result strongly suggested that the HSP105-specific CD8⁺ T cells were activated via the cross-presentation of HSP105 by BM-DCs. Although it became evident that gp96- and HSP70-chaperoned peptides can be presented to CTLs by DCs in the context of MHC class I molecules [38,39], we herein provide the first

evidence that HSP itself can be cross-presented to CTLs by DCs. HSP105-pulsed DC can present peptides derived from exogenously added HSP105 in the context of not only MHC class II molecules on the surface of DCs to activate CD4⁺ T cells, but also MHC class I molecules by cross-presentation to activate CD8⁺ T cells. We herein showed the induction of specific CD4⁺ T cells and CD8⁺ T cells *in vivo* by stimulation with HSP105-pulsed DCs. The application of the peptide-pulsed DC as potential vaccine is limited to patients with the appropriate HLA alleles. To circumvent this limitation, we have used HSP105-pulsed DC to induce a HSP105 specific T cell response. HSP105-pulsed DCs offer the advantage of potentially presenting multiple immunogenic T cell epitopes without the need of prior knowledge of the individual patient's HLA type.

The mechanism of action of HSP105-pulsed BM-DCs injected intraperitoneally is still unclear. We think that DCs injected in the abdominal cavity might immigrate into mesenteric lymphatic vessels. Some DCs stay in mesenteric lymph nodes, others circulate in the blood via the thoracic duct and finally reach the spleen and bone marrow. Recent experimental evidence suggested that peripheral DCs migrate through the lymphatic vessels to the blood [40]. Although the present study showed that intraperitoneal injection of DCs induced an effective anti-tumor immunity in mice, comparison of effectiveness to other routes of immunization with DCs, such as intravenous, subcutaneous, and intranodal, remains to be investigated.

In conclusion, our results indicate that HSP105 itself is a tumor rejection antigen which may possibly be useful for cancer immunotherapy, and that HSP105-pulsed BM-DC vaccinations can prime HSP105-specific T cells *in vivo*, to prevent the subcutaneous growth of B16-F10 and C26 cancer cells expressing HSP105, without inducing autoimmune destruction. Our findings suggest that HSP105-pulsed BM-DC vaccination is a novel strategy for the prevention of cancer in patients treated surgically, who are at high risk for a recurrence of the cancer. Because of the overexpression of HSP105 in a variety of human tumors [12], clinical trial of immunotherapy targeted against HSP105 may well be applicable to various cancers.

Acknowledgments

We thank Dr. Kyoichi Shimomura (Astellas Pharmaceutical Co.) for providing the cancer cell line. This work was supported in part by Grants-in-Aid (No. 12213111 for Y. Nishimura, and No. 14770142 for T. Nakatsura) from the Ministry of Education, Science, Technology, Sports and Culture, Japan, and The Sagawa Foundation for Promotion of Cancer Research and Meiji Institute of Health Science.

References

- [1] M.E. Feder, G.E. Hofmann, Heat-shock proteins, molecular chaperones, and the stress response: evolutionary and ecological physiology, *Annu. Rev. Physiol.* 61 (1999) 243–282.

- [2] P. Srivastava, Interaction of heat shock proteins with peptides and antigen presenting cells: chaperoning of the innate and adaptive immune responses, *Annu. Rev. Immunol.* 20 (2002) 395–425.
- [3] H. Udono, P.K. Srivastava, Heat shock protein 70-associated peptides elicit specific cancer immunity, *J. Exp. Med.* 178 (1993) 1391–1396.
- [4] Y. Tamura, P. Peng, K. Liu, M. Daou, P.K. Srivastava, Immunotherapy of tumors with autologous tumor-derived heat shock protein preparations, *Science* 278 (1997) 117–120.
- [5] G. Parmiani, A. Testori, M. Maio, C. Castelli, L. Rivoltini, L. Pilla, F. Belli, V. Mazzaferro, J. Coppa, R. Patuzzo, M.R. Sertoli, A. Hoos, P.K. Srivastava, M. Santinami, Heat shock proteins and their use as anticancer vaccines, *Clin. Cancer Res.* 10 (2004) 8142–8146.
- [6] J.M. Timmerman, R. Levy, Dendritic cell vaccines for cancer immunotherapy, *Annu. Rev. Med.* 50 (1999) 507–529.
- [7] A.J. Adler, D.W. Marsh, G.S. Yochum, J.L. Guzzo, A. Nigam, W.G. Nelson, D.M. Pardoll, CD4⁺ T cell tolerance to parenchymal self-antigens requires presentation by bone marrow-derived antigen-presenting cells, *J. Exp. Med.* 187 (1998) 1555–1564.
- [8] C. Kurts, W.R. Heath, F.R. Carbone, J. Allison, J.F. Miller, H. Kosaka, Constitutive class I-restricted exogenous presentation of self antigens in vivo, *J. Exp. Med.* 184 (1996) 923–930.
- [9] F. Castellino, P.E. Boucher, K. Eichelberg, M. Mayhew, J.E. Rothman, A.N. Houghton, R.N. Germain, Receptor-mediated uptake of antigen/heat shock protein complexes results in major histocompatibility complex class I antigen presentation via two distinct processing pathways, *J. Exp. Med.* 191 (2000) 1957–1964.
- [10] R.J. Binder, D.K. Han, P.K. Srivastava, CD91: a receptor for heat shock protein gp96, *Nat. Immunol.* 1 (2000) 151–155.
- [11] T. Nakatsura, S. Senju, K. Yamada, T. Jotsuka, M. Ogawa, Y. Nishimura, Gene cloning of immunogenic antigens overexpressed in pancreatic cancer, *Biochem. Biophys. Res. Commun.* 281 (2001) 936–944.
- [12] M. Kai, T. Nakatsura, H. Egami, S. Senju, Y. Nishimura, M. Ogawa, Heat shock protein 105 is overexpressed in a variety of human tumors, *Oncol. Rep.* 10 (2003) 1777–1782.
- [13] M. Miyazaki, T. Nakatsura, K. Yokomine, S. Senju, M. Monji, S. Hosaka, H. Komori, Y. Yoshitake, Y. Motomura, M. Minohara, T. Kubo, K. Ishihara, T. Hatayama, M. Ogawa, Y. Nishimura, DNA vaccination of Hsp105 leads to tumor rejection of colorectal cancer and melanoma in mice through activation of both CD4⁺ T cells and CD8⁺ T cells, *Cancer Sci.* 96 (2005) 695–705.
- [14] H. Singh-Jasuja, H.U. Scherer, N. Hilf, D. Arnold-Schild, H.G. Rammensee, R.E. Toes, H. Schild, The heat shock protein gp96 induces maturation of dendritic cells and down-regulation of its receptor, *Eur. J. Immunol.* 30 (2000) 2211–2215.
- [15] S.B. Flohé, J. Brüggemann, S. Lendemans, M. Nikulina, G. Meierhoff, S. Flohé, H. Kolb, Human heat shock protein 60 induces maturation of dendritic cells versus a Th1-promoting phenotype, *J. Immunol.* 170 (2003) 2340–2348.
- [16] R.J. Binder, K.M. Anderson, S. Basu, P.K. Srivastava, Heat shock protein gp96 induces maturation and migration of CD11c⁺ cells in vivo, *J. Immunol.* 165 (2000) 6029–6035.
- [17] H. Bausinger, D. Lipsker, U. Ziyilan, S. Manié, J.P. Briand, J.P. Cazenave, S. Muller, J.F. Haeuw, C. Ravanat, H. de la Salle, D. Hanau, Endotoxin-free heat-shock protein70 fails to induce APC activation, *Eur. J. Immunol.* 32 (2002) 3708–3713.
- [18] N. Yamagishi, H. Nishihori, K. Ishihara, K. Ohtsuka, T. Hatayama, Modulation of the chaperone activities of Hsc70/Hsp40 by Hsp105 α and Hsp105 β , *Biochem. Biophys. Res. Commun.* 272 (2000) 850–855.
- [19] M. Minohara, H. Ochi, S. Matsushita, A. Irie, Y. Nishimura, J. Kira, Differences between T-cell reactivities to major myelin protein-derived peptides in opticospinal and conventional forms of multiple sclerosis and healthy controls, *Tissue Antigens* 57 (2001) 447–456.
- [20] T. Nakatsura, H. Komori, T. Kubo, Y. Yoshitake, S. Senju, T. Katagiri, Y. Furukawa, M. Ogawa, Y. Nakamura, Y. Nishimura, Mouse homologue of a novel human oncofetal antigen, Glypican-3, evokes T-cell-mediated tumor rejection without autoimmune reaction in mice, *Clin. Cancer Res.* 10 (2004) 8630–8640.
- [21] S. Senju, S. Hirata, H. Matsuyoshi, M. Masuda, Y. Uemura, K. Araki, K. Yamamura, Y. Nishimura, Generation and genetic modification of dendritic cells derived from mouse embryonic stem cells, *Blood* 101 (2003) 3501–3508.
- [22] H. Matsuyoshi, S. Senju, S. Hirata, Y. Yoshitake, Y. Uemura, Y. Nishimura, Enhanced priming of antigen-specific CTLs in vivo by embryonic stem cell-derived dendritic cells expressing chemokine along with antigenic protein: application to antitumor vaccination, *J. Immunol.* 172 (2004) 776–786.
- [23] H. Fujita, S. Senju, H. Yokomizo, H. Saya, M. Ogawa, S. Matsushita, Y. Nishimura, Evidence that HLA class II-restricted human CD4⁺ T cells specific to p53 self peptides respond to p53 proteins of both wild and mutant forms, *Eur. J. Immunol.* 28 (1998) 305–316.
- [24] E.A. Craig, J.S. Weissman, A.L. Horwich, Heat shock proteins and molecular chaperones: mediators of protein conformation and turnover in the cell, *Cell* 78 (1994) 365–372.
- [25] K. Yasuda, A. Nakai, T. Hatayama, K. Nagata, Cloning and expression of murine high molecular mass heat shock proteins, HSP105, *J. Biol. Chem.* 270 (1995) 29718–29723.
- [26] D. Lee-Yoon, D. Easton, M. Murawski, R. Burd, J.R. Subjeck, Identification of a major subfamily of large hsp70-like proteins through the cloning of the mammalian 110-kDa heat shock protein, *J. Biol. Chem.* 270 (1995) 15725–15733.
- [27] X.Y. Wang, L. Kazim, E.A. Repasky, J.R. Subjeck, Characterization of heat shock protein 110 and glucose-regulated protein 170 as cancer vaccines and the effect of fever-range hyperthermia on vaccine activity, *J. Immunol.* 165 (2001) 490–497.
- [28] X.Y. Wang, X. Chen, M.H. Manjili, E. Repasky, R. Henderson, J.R. Subjeck, Targeted immunotherapy using reconstituted chaperone complexes of heat shock protein 110 and melanoma-associated antigen gp100, *Cancer Res.* 63 (2003) 2553–2560.
- [29] C.V. Nicchitta, Re-evaluating the role of heat-shock protein-peptide interactions in tumour immunity, *Nat. Rev. Immunol.* 3 (2003) 427–432.
- [30] H.D. Hickman-Miller, W.H. Hildebrand, The immune response under stress: the role of HSP-derived peptides, *Trends Immunol.* 25 (2004) 427–433.
- [31] K. Azuma, S. Shichijo, H. Takedatsu, N. Komatsu, H. Sawamizu, K. Itoh, Heat shock cognate protein 70 encodes antigenic epitopes recognized by HLA-B4601-restricted cytotoxic T lymphocytes from cancer patients, *Br. J. Cancer* 89 (2003) 1079–1085.
- [32] O. Faure, S. Graff-Dubois, L. Bretaudeau, L. Derré, D.A. Gross, P.M. Alves, S. Cornet, M.T. Duffour, S. Chouaib, I. Miconnet, M. Gregoire, F. Jotereau, F.A. Lemonnier, J.P. Abastado, K. Kosmatopoulos, Inducible HSP70 as target of anticancer immunotherapy: identification of HLA-A*0201-restricted epitopes, *Int. J. Cancer* 108 (2004) 863–870.
- [33] K. Hung, R. Hayashi, A. Lafond-Walker, C. Lowenstein, D. Pardoll, H. Levitsky, The central role of CD4⁺ T cells in the antitumor immune response, *J. Exp. Med.* 188 (1998) 2357–2368.
- [34] S.A. Rosenberg, J.C. Yang, N.P. Restifo, Cancer immunotherapy: moving beyond current vaccines, *Nat. Med.* 10 (2004) 909–915.
- [35] J. Banchereau, R.M. Steinman, Dendritic cells and the control of immunity, *Nature* 392 (1998) 245–252.
- [36] M. Bellone, D. Cantarella, P. Castiglioni, M.C. Crosti, A. Ronchetti, M. Moro, M.P. Garancini, G. Casorati, P. Dellabona, Relevance of the tumor antigen in the validation of three vaccination strategies for melanoma, *J. Immunol.* 165 (2000) 2651–2656.
- [37] M.L. Albert, B. Sauter, N. Bhardwaj, Dendritic cells acquire antigen from apoptotic cells and induce class I-restricted CTLs, *Nature* 392 (1998) 86–89.
- [38] E. Noessner, R. Gastpar, V. Milani, A. Brandl, P.J. Hutzler, M.C. Kuppner, M. Roos, E. Kremmer, A. Asea, S.K. Calderwood, R.D. Issels, Tumor-derived heat shock protein 70 peptide complexes are cross-presented by human dendritic cells, *J. Immunol.* 169 (2002) 5424–5432.

- [39] H. Singh-Jasuja, R.E. Toes, P. Spee, C. Münz, N. Hilf, S.P. Schoenberger, P. Ricciardi-Castagnoli, J. Neefjes, H.G. Rammensee, D. Arnold-Schild, H. Schild, Cross-presentation of glycoprotein 96-associated antigens on major histocompatibility complex class I molecules requires receptor-mediated endocytosis, *J. Exp. Med.* 191 (2000) 1965–1974.
- [40] L.L. Cavanagh, R. Bonasio, I.B. Mazo, C. Halin, G. Cheng, A.W. van der Velden, A. Cariappa, C. Chase, P. Russell, M.N. Starnbach, P.A. Koni, S. Pillai, W. Weninger, U.H. von Andrian, Activating of bone marrow-resident memory T cells by circulating, antigen-bearing dendritic cells, *Nat. Immunol.* 6 (2005) 1029–1037.