

against MC38 targets, whereas splenocytes from mice treated with PBS failed to exhibit detectable reactivity against this cell line. In contrast, splenocytes harvested from mice treated with Eph-DCs displayed far stronger anti-MC38 cytolytic reactivity than any control treatment group. CD8-positive, T cell-depleted splenocytes (harvested from mice treated with Eph-DCs) displayed significantly weaker anti-MC38 cytolytic reactivity than whole splenocytes; however, CD4-positive T cell-depleted splenocytes did not (Fig. 4B). Conversely, cytolytic activity was not observed against EphA2-negative cells (BL6) in any of the control/treatment arms, as shown in Figure 4A. We also harvested splenocytes 1 day after the second immunization (ie, Day 1 after tumor inoculation) to examine the potential early activation of NK cells by immunization with Eph-DCs. Figure 4C shows that no cytolytic activity was observed against the NK-sensitive YAC-1 cells in any treatment arm. These results suggest that principal antitumor effector cells in vaccinated mice are CD8-positive CTLs.

Requirement of Both CD4-positive T Cells and CD8-positive T Cells, but not NK Cells, for the Antitumor Effect of Immunization With Eph-DCs

To examine which lymphocyte subsets contributed to Eph-DC or unpulsed DC treatment, we performed depletion studies on a subset of CD4-positive T cells, CD8-positive T cells, and NK cells. Figure 5A shows that the therapeutic efficacy of Eph-DC therapy was reduced strongly in CD8-positive, T cell-depleted mice and was reduced partially in CD4-positive, T cell-depleted mice. In contrast, tumor growth still was suppressed in vaccinated mice that had been depleted of NK cells. In addition, the therapeutic efficacy of unpulsed DC therapy in CD4-positive or CD8-positive, T cell-depleted mice also was reduced, whereas that in NK cell-depleted mice was not reduced (Fig. 5B). These results suggest that both CD8-positive T cells and, to a lesser degree, CD4-positive T cells are required for the observed antitumor effects noted for Eph-DC vaccination. Both CD8-positive and CD4-positive T cells also were involved in unpulsed DC vaccination in our model.

Tumor Rechallenge

We then sought to determine whether prior Eph-DC treatment would have a durable effect on a subcutaneous rechallenge with MC38 tumor cells. C57BL/6 mice were immunized with subcutaneous injections of Eph-DCs and challenged with subcutaneous MC38 tumors. Forty-two days after the primary tumor inoculation, 2×10^5 MC38 cells were injected subcutaneously into the contralateral flank of these mice




FIGURE 5. Effective Eph peptide-pulsed dendritic cell (Eph-DC) or unpulsed DC vaccination required both CD4-positive and CD8-positive T cells, but not natural killer (NK) cells. The antibody-mediated, in vivo depletion of CD4-positive T cells (open triangles), CD8-positive T cells (open squares), and NK cells (open circles) was achieved as described in the text then, and the depleted mice were treated with either Eph-DC vaccines (A) or unpulsed DC vaccines (B) on Days -7 and 0, and they received subcutaneous murine colorectal adenocarcinoma (MC38) cells (2×10^5) on Day 0. Solid triangles indicate the phosphate-buffered saline-injected group. (A) The depletion of CD8-positive T cells markedly reduced the therapeutic efficacy of Eph-DC therapy (solid diamonds) ($P < .05$ on Days 14, 21, and 28 vs Eph-DCs), and the depletion of CD4-positive T cells partially reduced its therapeutic efficacy ($P < .05$ on Days 21 and 28 vs Eph-DCs); whereas the depletion of NK cells did not reduce its therapeutic efficacy ($N = 10$ mice per group). (B) The depletion of CD4-positive and CD8-positive T cells also reduced the therapeutic efficacy of unpulsed DC therapy (solid diamonds) ($P < .05$ on Days 21 and 28 vs unpulsed DCs), whereas the depletion of NK cells did not reduce its therapeutic efficacy ($N = 8$ mice per group). Each data point represents the mean tumor size \pm standard deviation.




FIGURE 6. Animals treated successfully with Eph peptide-pulsed dendritic cell (Eph-DC) vaccines rejected a subsequent rechallenge with murine colorectal adenocarcinoma (MC38) (Eph receptor A2 [EphA2]-positive) tumors. Animals treated with Eph-DC vaccines prior to MC38 subcutaneous injection were rechallenged subcutaneously 42 days after the initial tumor inoculation with MC38 cells in the flank contralateral to the initial challenge. In parallel with this, naive mice were injected subcutaneously with MC38 cells as a control. Rechallenged MC38 tumors in mice that received the Eph-DC regimen (diamonds) were inhibited significantly through the chosen endpoint of these experiments on Day 28 (after MC38 rechallenge) compared with tumor growth in naive mice (triangles) ($P < .05$ on Days 14, 21, 28; $N = 9$ mice per group).

on Day 0. Figure 6 shows that rechallenged tumors in mice that received the Eph-DC treatment regimen were inhibited significantly in their progression through the chosen endpoint of these experiments on Day 28 ($P < .05$ vs naive mice on Days 14, 21, and 28).

DISCUSSION

Modified, DC-based vaccines using synthetic peptides derived from known tumor antigens, such as CEA, have been reported for colon cancer; although, to date, objective clinical responses have been observed only in a minority of patients who received treatment with these modalities.¹⁰⁻¹² This may be explained in part by the application of DC-based vaccines to immunosuppressed patients with advanced colon cancer and/or to the modest immunogenicity of tumor antigens (ie, CEA) that have been applied to date in this setting. Recently, a novel tumor antigen, EphA2, has been identified that has specific characteristics and that frequently is overexpressed in advanced cancers, suggesting that this antigen may have great potential as a target for immunotherapy, especially in

patients with advanced-stage or metastatic cancer. In the current study, 60% of colon cancer tissue samples overexpressed EphA2, consistent with a recent report by Saito et al.²⁵ We demonstrated that Eph-DC vaccines effectively promoted antitumor effects in a colon cancer model, suggesting that EphA2-derived CTL epitopes have the potential to serve as relevant components of novel DC-based vaccines for colon cancer.

IFN- γ ELISPOT assays revealed that immunization with Eph-DCs in normal mice resulted in the induction of specific CD8-positive T cells. Based on these results, we examined the antitumor effectiveness of Eph-DC vaccines in a syngenic, EphA2-positive MC38 colon cancer model. The Eph-DC vaccines induced antitumor effects against EphA2-positive MC38 colon carcinoma, but not against EphA2-negative BL6 melanoma, suggesting that EphA2-specific antitumor immunity was generated by Eph-DC vaccines, consistent with the results from our earlier IFN- γ ELISPOT assays.

In vitro assays revealed that the main antitumor effector cells for killing MC38 colon cancer cells were CD8-positive T cells and, possibly, CTLs. This cytolytic activity was specific for MC38 cells, because splenocytes did not kill BL6 cells. These results suggested that Eph-DC vaccines could efficiently generate specific CTLs that recognize and kill relevant EphA2-positive (but not irrelevant EphA2-negative) tumor targets.

Our in vivo lymphocyte-depletion studies demonstrated that CD8-positive T cells contributed to the inhibition of tumor growth in Eph-DC immunization and that CD4-positive T cells contributed to a lesser extent. Moreover, our tumor experimental data demonstrated that immunization with Eph-DCs maintains the antitumor effect against MC38 tumor over an extended period of time, despite the use of a single EphA2-derived CD8-positive T cell epitope in the DC-based vaccine. Typically, effector CD8-positive T cells induced by minimal CTL epitope peptides do not persist for a long time, and the induction of durable-memory CD8-positive T cells requires the support of CD4-positive T cells.³¹⁻³³ Therefore, these results suggest that Eph-DC vaccines may activate CD8-positive T cells (that recognize EphA2-derived CTL epitopes) and CD4-positive T cells (that recognize tumor antigens related to MC38 colon cancer cells), which are taken up by specifically dedicated antigen-presenting cells, and that the activated CD4-positive T cells likely contribute to the generation and maintenance of memory in EphA2-specific, CD8-positive T cells.

Immunization with control, unpulsed DCs was inhibited both EphA2-positive MC38 tumor growth

compared with PBS. Generally, unpulsed DC vaccines are not expected to generate CTLs. However, our lymphocyte-depletion studies in the MC38 tumor model demonstrated that the therapeutic efficacy of unpulsed DC therapy in CD4-positive or CD8-positive T cell-depleted mice was reduced equally. These results suggest that unpulsed DCs can induce protective antitumor effects in mice through the presentation of "self" peptides in MHC complexes to specific autoreactive CTLs, which are capable of recognizing tumor cells that also present these peptides, consistent with the previous report by Dworacki et al.³⁴ Moreover, unexpectedly, Eph-DC or unpulsed DC vaccines had weak antitumor effects against EphA2-negative BL6 tumors compared with PBS treatment. BL6 cells do not express EphA2, and EphA2-specific CTLs do not have cytolytic activity against BL6 tumors. Dworacki et al. reported that immunization with unpulsed DCs inhibited a variety of syngeneic tumors through the activation of both CD4-positive T cells and CD8-positive T cells,³⁴ suggesting that Eph-DC or unpulsed DC vaccines may activate CD4-positive and CD8-positive T cells weakly and that these cells may play a role in weakly inhibiting BL6 tumor growth.

Recent research in DC biology has revealed that DCs also contribute to innate immune responses by activating NK cells through IL-12 secretion and direct cellular interaction.³⁵ However, our current data demonstrated that NK cells were not involved in generating antitumor effect of Eph-DC or unpulsed DC vaccination in our lymphocyte-depletion studies. We speculate that subcutaneous, local NK cells may not be activated efficiently by immunization of Eph-DCs or unpulsed DCs, because NK cells are not so abundant in the subcutaneous tissue. Instead, if we were to apply this DC vaccine in sites rich in NK cells (ie, the liver), then this strategy may prove to be more effective in treating NK cell-sensitive liver cancers, for instance. Currently, we are evaluating these possibilities and performing histopathologic evaluations to confirm that treated animals do not exhibit auto-immune pathology in organs (such as lung, kidney, etc) that constitutively express low levels of EphA2.

Despite the recent progress and early successes reported for DC-based cancer immunotherapies, there is significant room for improvement in these regimens, especially with respect to advanced colon cancer. In this study, we demonstrated that Eph-DC vaccines revealed antitumor effects against colon cancers. In addition to CEA-based vaccines, EphA2-derived peptide-pulsed DC vaccines may represent a promising therapeutic modality against advanced colon cancers. Recently, it has been reported that some clinical trials using peptide cocktail-pulsed DCs may be useful strategies

for treating patients with malignant tumors.^{36,37} Therefore, DCs pulsed with multiple peptides derived from various tumor-associated antigens, including both EphA2 and CEA, may improve the therapeutic effects against advanced colon cancers.

REFERENCES

1. Fields RC, Shimizu K, Mule JJ. Murine dendritic cells pulsed with whole tumor lysates mediate potent antitumor immune response in vitro and in vivo. *Proc Natl Acad Sci USA*. 1998;95:9482-9487.
2. Paglia P, Chiodoni C, Rodolfo M, Colombo MP. Murine dendritic cells loaded in vitro with soluble protein prime cytotoxic T lymphocytes against tumor antigen in vivo. *J Exp Med*. 1996;183:317-322.
3. Mayordome JL, Zorina T, Storkus WJ, et al. Bone marrow-derived dendritic cells pulsed with synthetic tumor peptide elicits protective and therapeutic antitumor immunity. *Nat Med*. 1995;12:1297-1302.
4. Boczkowski D, Nair SK, Snyder D, Gilboa E. Dendritic cells pulsed with RNA are potent antigen-presenting cells in vitro and in vivo. *J Exp Med*. 1996;184:465-472.
5. Condon C, Watkins SC, Celluzzi CM, Thompson K, Falo LD Jr. DNA-based immunization by in vitro transfection of dendritic cells. *Nat Med*. 1996;2:1122-1128.
6. Hsu FJ, Benike C, Fagnoni F, et al. Vaccination of patients with B-cell lymphoma using autologous antigen-pulsed dendritic cells. *Nat Med*. 1996;2:52-58.
7. Nestle FO, Alijagic S, Gilliet M, et al. Vaccination of melanoma patients with peptide- or tumor lysate-pulsed dendritic cells. *Nat Med*. 1998;4:328-322.
8. Salgaller ML, Tjoa BA, Lodge PA, et al. Dendritic cell-based immunotherapy of prostate cancer. *Crit Rev Immunol*. 1998;18:109-119.
9. Kugler A, Stuhler G, Walden P, et al. Regression of human metastatic renal cell carcinoma after vaccination with tumor cell-dendritic cell hybrids. *Nat Med*. 2000;6:332-336.
10. Simone M, Carlo RR, Mario L, Donato N. Colorectal cancer vaccines: principles, results, and perspectives. *Gastroenterology*. 2004;27:1821-1837.
11. Alters SE, Gader Jr, Philip R. Immunology of cancer. Generation of CEA specific CTL using CEA peptide pulsed dendritic cells. *Adv Exp Med Biol*. 1997;417:519-524.
12. Vermorken JB, Classen AM, van Tinteren H, et al. Active specific immunotherapy for stage 2 and 3 human colon cancer: randomized trial. *Lancet*. 1999;353:345-350.
13. Wilkinson DG. Multiple roles of Eph receptors and ephrins in neural development. *Nat Rev Neurosci*. 2001;2:155-164.
14. Mellitzer G, Xu Q, Wilkinson DG. Control of cell behavior by signaling through Eph receptors and ephrins. *Curr Opin Neurobiol*. 2000;10:400-408.
15. Knoll B, Zarbalis K, Wurst W, Drescher U. A role for the EphA family in the topographic targeting of vomeronasal axons. *Development*. 2000;128:895-906.
16. Oba S, Wang Y, Song J, et al. Genomic structure and loss of heterozygosity of EphB2 in colorectal cancer. *Cancer Lett*. 2001;164:97-104.
17. Dodelet V, Pasquale E. Eph receptors and Ephrin ligands: embryogenesis to tumorigenesis. *Oncogene*. 2000;19:5614-5619.
18. Bittner M, Meltzer P, Chen Y, et al. Molecular classification of cutaneous malignant melanoma by gene expression profiling. *Nature*. 2000;406:536-540.

19. Pawel D, Kathryn H, Susan J, et al. Antiangiogenic and antitumor efficacy of EphA2 receptor antagonist. *Cancer Res.* 2004;64:910-919.
20. Cheng N, Brantley DM, Chen J. The ephrins and Eph receptors in angiogenesis. *Cytokine Growth Factor Rev.* 2002;13:75-85.
21. Easty D, Gurthrie B, Maung K, et al. Protein B61 as a new growth factor: expression of B61 and up-regulation of its receptor epithelial cell kinase during melanoma progression. *Cancer Res.* 1995;55:2528-2532.
22. Zantek ND, Walker-Daniels J, Stewart J, Wang B, Brackenbury R, Kinch MS. MCF-10A-NeoST: a new cell system for studying cell-ECM and cell-cell interaction in breast cancer. *Clin Cancer Res.* 2001;7:3640-3648.
23. Kinch MS, Moore MB, Harpole DH Jr. Predictive value of the EphA2 receptor tyrosine kinase in lung cancer recurrence and survival. *Clin Cancer Res.* 2003;9:613-618.
24. Christopher JH, Tatsumi T, Kathleen SO, et al. Expression of EphA2 is prognostic of disease-free interval and overall survival in surgically treated patients with renal cell carcinoma. *Clin Cancer Res.* 2005;11:226-231.
25. Saito T, Masuda N, Miyazaki T, et al. Expression of EphA2 and E-cadherin in colorectal cancer with cancer metastasis. *Oncol Rep.* 2004;11:605-611.
26. Tatsumi T, Herrem CJ, Olson WC, et al. Disease stage variation in CD4+ and CD8+ T-cell reactivity to the receptor tyrosine kinase EphA2 in patients with renal cell carcinoma. *Cancer Res.* 2003;63:4481-4489.
27. Alves PM, Fature O, Graff-Dubois S, et al. EphA2 as target of anticancer immunotherapy: identification of HLA-A0201-restricted epitopes. *Cancer Res.* 2003;63:8476-8480.
28. Hatano M, Kuwashima N, Tatsumi T, et al. Vaccination with EphA2-derived T cell-epitopes promotes immunity against both EphA2-expressing and EphA2-negative tumors [electronic resource]. *J Transl Med.* 2004;2:40.
29. Tatsumi T, Gambotto A, Robbins PD, Storkus WJ. Interleukin 18-gene transfer expands the repertoire of anti-tumor Th1-type immunity elicited by dendritic cell-based vaccines in association with enhanced therapeutic efficacy. *Cancer Res.* 2002;62:5853-5858.
30. Mayordomo BJI, Loftus DJ, Sakamoto H, et al. Therapy of murine tumors with p53 wild-type and mutant sequence peptide-based vaccines. *J Exp Med.* 1996;183:1357-1365.
31. Janssen EM, Lemmens EE, Wolfe T, Christen U, von Herrath MG, Schoenberger SP. CD4+ T cells are required for secondary expansion and memory CD8+ T lymphocytes. *Nature.* 2003;421:852-856.
32. Sun JC, Bevan MJ. Detective CD8 T cell memory following acute infection without CD4 T cell help. *Science.* 2003;300:339-342.
33. Bennet RM, Carbone FR, Karemalls F, Miller FAP, Heath WR. Induction of CD8+ cytotoxic T lymphocyte response by cross-priming requires cognate CD4+ T cell help. *J Exp Med.* 1997;186:65-70.
34. Dworacki G, Cicinnati VR, Beckebaum S, Pizzoferrato E, Hoffmann TK, De Leo SB. Unpulsed dendritic cells induce broadly applicable anti-tumor immunity in mice. *Cancer Biol Ther.* 2005;4:50-56.
35. Fernandez NC, Lozier A, Flament C, et al. Dendritic cells directly trigger NK cell functions: cross-talk relevant in innate anti-tumor immune responses in vivo. *Nat Med.* 1999;5:405-411.
36. Fuessel S, Meye A, Schmitz M, et al. Vaccination of hormone-refractory prostate cancer patients with peptide cocktail-loaded dendritic cells: results of a phase I clinical trial. *Prostate.* 2006;66:811-821.
37. Akiyama Y, Tanosaki R, Inoue N, et al. Clinical response in Japanese metastatic melanoma patients treated with peptide cocktail-pulsed dendritic cells [electronic resource]. *J Transl Med.* 2005;3:4.

Signal Transducer and Activator of Transcription 3 Signaling Within Hepatocytes Attenuates Systemic Inflammatory Response and Lethality in Septic Mice

Ryotaro Sakamori,^{1*} Tetsuo Takehara,^{1*} Chihiro Ohnishi,¹ Tomohide Tatsumi,¹ Kazuyoshi Ohkawa,¹ Kiyoshi Takeda,² Shizuo Akira,³ and Norio Hayashi¹

Sepsis is an infection-induced syndrome with systemic inflammatory response leading to multi-organ failure and occasionally death. During this process, signal transducer and activator of transcription 3 (STAT3) is activated in the liver, but the significance of this molecule has not been established. We generated hepatocyte-specific STAT3-deficient mice (L-STAT3 KO) and examined the susceptibility of these mice to cecal ligation and puncture-induced peritonitis, a well-established septic model. L-STAT3 KO mice showed significantly higher mortality and produced lesser amounts of various acute phase proteins than control littermates. Although blood bacterial infection did not differ between L-STAT3 KO mice and control mice, the former showed deterioration of the systemic inflammatory response as evidenced by a significant increase in various cytokines such as tumor necrosis factor α , IFN- γ , IL-6, IL-10, monocyte chemoattractant protein 1, and macrophage inflammatory protein 1 β . A similar hyperinflammatory response was observed in another septic model caused by lipopolysaccharide (LPS) injection. *In vitro* analysis revealed that soluble substances derived from hepatocytes and dependent on STAT3 were critical for suppression of cytokine production from LPS-stimulated macrophage and splenocytes. **Conclusion:** STAT3 activation in hepatocytes can attenuate a systemic hyperinflammatory response and lethality in sepsis, in part by suppressing immune cell overactivation, implying a critical role of hepatocyte STAT3 signaling in maintaining host homeostasis. (HEPATOLOGY 2007;46: 1564-1573.)

Signal transducer and activator of transcription 3 (STAT3) mediates a signal from the IL-6 family of cytokines such as IL-6, oncostatin M, leukemia inhibitory factor, and ciliary neurotrophic factor, and acti-

vates transcription of various target genes.¹ Although a STAT3 is now known to be ubiquitously expressed in variety of cells and has pleiotropic functions, it was formerly termed *acute phase response factor* and was first identified in the liver as an inducible DNA binding protein binding to type 2 IL-6-responsive elements within the promoter of hepatic acute phase protein (APP) genes.^{2,3} Because deletion of STAT3 leads to embryonic lethality in mice, the significance of STAT3 in adult organs has been investigated using conditional knockout animals generated by the Cre/loxP recombination system.⁴ Research has shown that STAT3 signaling within hepatocytes controls a variety of physiological or pathological processes, including hepatocyte proliferation after partial hepatectomy,⁵ apoptosis resistance of hepatocytes during Fas-mediated liver injury,⁶ and regulation of hepatic gluconeogenic genes.⁷ Although STAT3 is activated in response to a rise of circulating cytokines, the significance of hepatic STAT3 has not been elucidated under systemic inflammatory conditions.

Sepsis is an infection-induced systemic syndrome, the incidence of which is estimated at 750,000 cases annually in North America with overall mortality being approxi-

Abbreviations: APP, acute phase protein; CLP, cecal ligation and puncture; LPS, lipopolysaccharide; L-STAT3 KO, hepatocyte-specific STAT3-deficient mice; STAT3, signal transducer and activator of transcription 3; TNF- α , tumor necrosis factor α ; TUNEL, terminal deoxynucleotidyl transferase-mediated dUTP nick end-labeling.

From the ¹Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, Osaka, Japan; the ²Department of Molecular Genetics, Medical Institute of Bioregulation, Faculty of Medical Sciences, Kyushu University, Fukuoka, Japan; and the ³Department of Host Defense, Research Institute for Microbial Diseases, Osaka University, Osaka, Japan.

Received March 13, 2007; accepted May 25, 2007

Supported by a grant-in-aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

*These authors contributed equally to this study.

Address reprint requests to: Dr. Norio Hayashi, Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, 2-2 Yamada-oka, Suita, Osaka 565-0871, Japan. E-mail: hayashin@gh.med.osaka-u.ac.jp; fax: (81)-6-6879-3629.

Copyright © 2007 by the American Association for the Study of Liver Diseases. Published online in Wiley InterScience (www.interscience.wiley.com).

DOI 10.1002/hep.21837

Potential conflict of interest: Nothing to report.

Supplementary material for this article can be found on the HEPATOLOGY website (<http://interscience.wiley.com/jpages/0270-9139/suppmat/index.html>).

mately 30%, but rising to 40% in the elderly.⁸ Sepsis develops when the initial, appropriate host response to an infection becomes amplified and then dysregulated.⁹ Among those harmful or damaging responses is the rise of a variety of circulating cytokines such as IL-6, tumor necrosis factor α (TNF- α), IL-10, and IFN- γ . These cytokines lead directly to the development of systemic inflammatory response syndrome. During this process, an increasing proportion of patients will develop adult respiratory distress syndrome, disseminated intravascular coagulation, and/or acute renal failure, leading to the multiple organ dysfunction syndrome.¹⁰ The liver is also one of the target organs of multiple organ dysfunction syndrome, although liver dysfunction may cause patient death less frequently than cardiovascular dysfunction.¹¹ Conversely, sepsis is a serious complication of severe liver diseases such as fulminant hepatitis¹² and decompensated cirrhosis.¹³ Thus, research on the relevance of signal transduction in liver cells in the septic condition would not only satisfy basic scientific interest but would also have clinical implications.

In the present study, we used hepatocyte-specific STAT3-deficient (L-STAT3 KO) mice and examined the significance of STAT3 signaling within hepatocytes in a well-established murine model of sepsis. We found that STAT3 deficiency in hepatocytes causes exacerbation of the hyperinflammatory response by attenuating hepatic production of soluble substances that can suppress immune cell activation and also increases mortality in septic mice. This study identified an anti-inflammatory function of hepatic STAT3 signaling and its protective role against systemic inflammation, providing genetic evidence for a close link between hepatocytes and the immune system.

Materials and Methods

Animals. Mice carrying a STAT3 gene with 2 *loxP* sequences flanking exon 22 and a STAT3 null allele (*STAT3* fl/−) have been described previously.¹⁴ To generate mice with hepatocyte-specific STAT3 deficiency, we crossed *STAT3* fl/− mice and Alb-Cre transgenic mice,¹⁵ which express the Cre recombinase gene under the regulation of the albumin gene promoter. We crossed Alb-Cre *STAT3* fl/fl mice and *STAT3* fl/− mice. The resulting Alb-Cre *STAT3* fl/− mice were used as L-STAT3 KO mice. Sex-matched *STAT3* fl/− mice obtained from the same litter were used as control mice. All mice were used at the age of 12–15 weeks. All animals were housed under specific pathogen-free conditions and were treated with humane care under approval from the Animal Care and Use Committee of Osaka University Medical School.

Cecal Ligation and Puncture and Lipopolysaccharide Injection. Cecal ligation and puncture (CLP) is a well-established murine model of septic shock. The mice underwent CLP surgery as described previously.¹⁶ In brief, the mice were anesthetized via intraperitoneal injection of sodium pentobarbital. Under sterile condition, the cecum was assessed via a 1-cm midline incision of the lower abdomen, ligated with a suture below the ileocecal valve, and punctured once with a 23-gauge needle. The cecum was replaced in the peritoneum, and the abdomen was closed with sutures. The mice were injected with 1 mL of lactate Ringer's solution subcutaneously for fluid resuscitation. As another septic model, lipopolysaccharide (LPS) (form *Escherichia coli* 055: B5; Sigma, St. Louis, MO) was injected intraperitoneally at a dose of 4 mg/kg body weight.

Preparation of Peritoneal Macrophage. To isolate peritoneal macrophages, we injected mice intraperitoneally with 2 mL of 4% thioglycollate. Peritoneal exudates cells were isolated from the peritoneal cavity 4 days after injection. The cells were incubated for 4 hours in 96-well plates and washed 3 times with phosphate-buffered saline. We used the adherent cells as peritoneal macrophages for further experiments.

Determination of the Bacterial Load. Mice were sacrificed 24 hours after CLP surgery. Samples of blood were obtained in sterile condition. Fifty microliters of the blood were then plated on heart-infusion plates. The heart-infusion plates were incubated at 37°C overnight, and the number of bacteria colonies was counted. Results were expressed as log₁₀ of CFU.

Blood Biochemistry. Blood samples were obtained 24 hours after CLP or LPS injection. Acute phase proteins, cytokines, and chemokines in plasma were determined via MultiAnalyte Profile testing (Rules Based Medicine, Austin, TX). Levels of serum ALT and creatinine were measured with a standard UV method using a Hitachi type 7170 automatic analyzer (Tokyo, Japan).

Measurement of Culture Supernatant. Levels of cytokines (TNF- α , IL-6, IL-10, and IFN- γ) in the culture supernatants were measured using commercially available ELISA kits in accordance with the manufacturer's instructions (BD Biosciences-Pharmingen, San Diego, CA). Haptoglobin was determined in cell-free supernatants by using a commercially available ELISA kit (Immunology Constants Laboratory, Newberg, OR).

Western Blot Analysis. The total cellular protein was extracted with the RIPA buffer containing 1% Nonidet P-40, 0.5% sodium deoxycholate, 0.1% SDS, 50 μ g/mL aprotinin, 100 μ g/mL phenylmethylsulfonyl fluoride, and 50 mM sodium fluoride in phosphate-buffered saline (pH 7.4). Twenty micrograms of protein were separated

via 7.5% SDS-PAGE and blotted onto a polyvinylidene difluoride membrane. After blocking with Tris-buffered saline 0.1% Tween 20 containing 5% skim milk or Blocking One-P (Nacalai Tesque, Kyoto, Japan) for 1 hour at room temperature, the membrane was incubated overnight at 4°C with antibodies to STAT3 or tyrosine⁷⁰⁵-phosphorylated STAT3 (Cell Signaling Technology, Danvers, MA), respectively. After washing with Tris-buffered saline 0.1% Tween 20, the membrane was incubated with anti-horseradish peroxidase-linked antibody for 1 hour at room temperature. The immune complex was detected by an enhanced chemiluminescent assay. In some experiments, tyrosine⁷⁰¹-phosphorylated STAT1 antibody (Cell Signaling Technology) was also used. This antibody recognizes the phosphorylated form of both STAT1 α and STAT1 β .

Histology and Terminal Deoxynucleotidyl Transferase-Mediated dUTP Nick End-Labeling. The formalin-fixed livers were paraffin-embedded, and liver sections were analyzed by hematoxylin-eosin staining. Terminal deoxynucleotidyl transferase-mediated deoxyuridine triphosphate nick-end labeling (TUNEL) was performed using an ApopTag kit according to the manufacturer's instructions (Serological Corporation, Norcross, GA).

Primary Culture of Hepatocytes. Livers were digested using a standard *in situ* 2-step collagenase perfusion procedure (Gibco BRL, Rockville, MD). Hepatocytes were isolated from nonparenchymal cells via subsequent centrifugation at 50g for 1 minute. In a selected experiment, nonparenchymal cells in the supernatants were pelleted at 1,500 rpm for 5 minutes and subjected to western blot analysis. Isolated hepatocytes with >90% viability were cultured in Williams' medium E containing 10% fetal bovine serum overnight. On the next day, the cells were stimulated with recombinant IL-6 (PeproTech, London, UK). The cells were harvested after 2 hours for the analysis of STAT3 activation. In another experiment, supernatants were harvested after 48 hours.

Cytokine Production by Macrophage and Splenocytes. The murine macrophage cell line RAW 264.7 was obtained from the American Type Culture Collection (Manassas, VA). RAW cells were plated at a density of 5×10^5 /well in a 96-well plate and were incubated at 37°C in culture supernatants of hepatocyte from L-STAT3 KO mice or control mice. As a control, RAW cells were also cultured in Williams' medium E. After 24 hours, LPS was added to achieve a final concentration of 100 ng/mL. After 24 hours of incubation at 37°C in an atmosphere of 5% CO₂, the supernatant was collected and stored at -80°C for measurement of TNF- α , IL-6, and IL-10. Splenocytes were isolated by way of a standard

procedure for wild-type mice¹⁷ and incubated with hepatocyte culture supernatant. Twenty-four hours after incubation, the cells were stimulated with LPS (1,000 ng/mL) for 24 hours. The resultant culture supernatant was subjected to IFN- γ ELISA.

Statistics. Kaplan-Meier curves were used to show survival over time. Data are expressed as interquartile range and median and compared using the Mann-Whitney *U* test. Statistical significance was set at $P < 0.05$.

Results

Mice with hepatocyte-specific STAT3 deficiency were produced by crossing floxed STAT3 mice and Alb-Cre transgenic mice carrying the Cre recombinase gene under the regulation of the albumin gene promoter. L-STAT3 KO mice were born and grew without any gross abnormality. Western blot analysis revealed that STAT3 expression was substantially decreased in the liver but not in other organs (Fig. 1A). Isolation of hepatocytes from nonparenchymal cells by liver perfusion followed by centrifugation confirmed that STAT3 deficiency is specific in hepatocytes (Fig. 1B). In addition, STAT3 expression did not differ in peritoneal macrophages between L-STAT3 KO mice and control littermates (Fig. 1C). Those cells isolated from L-STAT3 KO mice produced similar levels of TNF- α in response to LPS compared with those from control littermates (Fig. 1D).

L-STAT3 KO Mice Are More Vulnerable to Septic Shock. To examine the role of hepatic STAT3 during septic shock, we used a well-examined clinically relevant murine model of sepsis performed by CLP.¹⁶ CLP clearly activated liver STAT3, which was determined via phosphorylation of STAT3 in control mice (Fig. 2A), in agreement with a previous report.¹⁸ Liver STAT3 activation during sepsis is mostly due to the activation of STAT3 in hepatocytes, because liver STAT3 was only marginally activated in L-STAT3 KO mice. CLP activated liver STAT1 both in L-STAT3 KO mice and wild-type mice, suggesting that the absence of STAT3 does not affect the activation of other STATs. Given that STAT3 is a well-known mediator for APP,¹⁹ we measured APPs such as fibrinogen and haptoglobin in plasma after CLP (Fig. 2B). The levels of fibrinogen and haptoglobin clearly increased after CLP in wild-type mice. In contrast, induction of fibrinogen was completely diminished in L-STAT3 KO mice, whereas that of haptoglobin was partially inhibited. This is consistent with the previous notion that fibrinogen is a class 2 gene and haptoglobin is a class 1 gene; the class 2 gene is predominantly regulated by type 2 IL-6 responsive elements binding to STAT²⁰ and the class 1 gene by both type 1 IL-6 responsive elements




Fig. 1. Hepatocyte-specific STAT3 deficiency in mice. Floxed STAT3 mice were crossed with Alb-Cre transgenic mice. Floxed STAT3 mice having the Alb-Cre transgene were regarded as L-STAT3 KO mice (KO); those not having the Alb-Cre gene were used as a wild-type control (WT). (A) STAT3 expression in a variety of organs from L-STAT3 KO mice and wild-type mice via western blot analysis. Expression of GAPDH was served as a loading control. Representative blots are shown. (B) Expression of STAT3 of isolated hepatocytes and nonhepatocytes. Liver of L-STAT3 KO mice or wild-type mice was collagenase-perfused and separated into hepatocyte and nonhepatocyte fractions. STAT3 expression was determined via western blot analysis. Expression of GAPDH was served as a loading control. Representative blots are shown. (C) Expression of STAT3 in isolated macrophage. Peritoneal macrophage was isolated from L-STAT3 KO mice or wild-type mice and subjected to western blot analysis of STAT3 expression. Representative blots are shown. (D) LPS-stimulated TNF- α production of peritoneal macrophages. Peritoneal macrophages were isolated from L-STAT3 KO mice or wild-type mice ($n = 6$ for each group) and stimulated with LPS (100 ng/mL) for 24 hours. TNF- α production was determined via ELISA in culture supernatants.

binding to CCAAT enhancer-binding protein (C/EBP) and type 2 IL-6 responsive elements.²¹

To address the issue of whether hepatic STAT3 is involved in the outcome of CLP-induced lethality, we performed CLP blinded to the genetic background and checked the survival of the mice every 6 hours. L-STAT3 KO mice were significantly more vulnerable to CLP-induced lethality than wild-type littermates (Fig. 2C). To examine the possible difference in bacterial infection after CLP, we measured colony forming unit of blood bacteria 24 hours after CLP. Because there was no significant difference in bacterial amount between L-STAT3 KO mice and wild-type mice (Fig. 2D), we considered hepatic STAT3 to have had a beneficial effect on the outcome of septic shock without affecting bacterial infection.

Hepatic STAT3-Deficient Mice Show Exacerbated Liver Injury. To examine liver injury and renal dysfunction in CLP-induced sepsis, we measured ALT and creatinine levels. L-STAT3 KO mice showed increased levels of serum ALT and creatinine compared with wild-type littermates, although the difference in creatinine did not reach a significant level (Fig. 3A). TUNEL of the liver revealed that the number of apoptotic hepatocytes was significantly higher in L-STAT3 KO mice than in wild-type littermates (Fig. 3B,C). However, the liver injury itself presumably is not a direct cause of animal death, because histologic abnormality was modest. Furthermore,

LPS injection, which is another model of septic shock, induced more hepatocyte apoptosis than CLP but did not kill any mice tested (Fig. 3A-C and data not shown), supporting the idea that increased liver injury could not explain the increased lethality in L-STAT3 KO mice.

Exacerbated Systemic Inflammatory Response in L-STAT3 KO Mice. Hypercytokinemia underlying systemic inflammatory response syndrome may play an important role in the development of multiple organ dysfunction syndrome and lethality.⁹ We measured several circulating cytokines and chemokines in septic mice and found that TNF- α , IFN- γ , IL-6, IL-10, monocyte chemoattractant protein-1 (MCP-1) and macrophage inflammatory protein-1 β (MIP-1 β) had clearly increased 24 hours after CLP in L-STAT3 KO mice. Of importance is the finding that the plasma levels of these cytokines and chemokines were significantly higher in L-STAT3 KO mice than in wild-type mice, although they did not differ before CLP. This result indicates that the increased lethality found in L-STAT3 KO mice is associated with hypercytokinemia (Fig. 4A). Although plasma insulin levels significantly increased 24 hours after CLP, there was no significant difference between L-STAT3 KO mice and wild-type mice, suggesting that insulin levels do not affect the difference in animal lethality (Supplementary Fig. 1).

Given that bacterial infection did not differ between L-STAT3 KO mice and wild-type mice, we examined the




Fig. 2. STAT3 activation, APP production, and survival in CLP mice. (A) Western blot analysis of phosphorylated STAT3 and STAT1 in CLP mice. L-STAT3 KO mice (KO) and wild-type mice (WT) were treated with CLP and sacrificed at the indicated time points. Their liver tissues were subjected to analysis of Tyrosine 705 phosphorylation of STAT3 or Tyrosine 701 phosphorylation of STAT1 via western blot analysis. GAPDH expression served as a control. Representative blots are shown. *7 days. (B) Levels of circulating haptoglobin and fibrinogen before and 24 hours after CLP (n = 8 for each group). *P < 0.05. (C) Comparison of survival after CLP between L-STAT3 KO (n = 27) mice and wild-type littermates (n = 28). (D) Colony-forming units of blood bacteria after CLP. L-STAT3 KO or wild-type mice were sacrificed 24 hours after CLP. Blood samples were subjected to analysis of bacterial growth (n = 10 for knockout mice and n = 9 for wild-type mice).

response of cytokine production upon endotoxin stimulation. To this end, we injected the same amount of LPS to L-STAT3 KO mice and control mice and measured circulating cytokines. LPS injection into L-STAT3 KO mice upregulated those cytokines to a lesser extent than CLP. In agreement with the finding on the CLP model, the levels of TNF- α , IL-10, MCP-1, and MIP-1 β were significantly higher in L-STAT3 KO mice than in wild-type mice after LPS injection (Fig. 4B), indicating that L-STAT3 KO mice were highly sensitive to endotoxin and prone to show hypercytokinemia.

STAT3-Regulated Soluble Factors Produced by Hepatocytes Suppress Cytokine Production From Immune Cells. To examine the underlying mechanisms of the hyperimmune response in L-STAT3 KO mice, we hypothesized that STAT3-mediated soluble factors from hepatocytes repress cytokine production from immune cells. We isolated hepatocytes from L-STAT3 KO mice

and control mice and stimulated them with or without IL-6, collecting the conditional medium of hepatocytes. Wild-type hepatocytes displayed STAT3 activation in primary culture without stimulation, but the levels increased upon IL-6 exposure, whereas KO hepatocytes did not show any STAT3 activation (Fig. 5A). Consistent with this was the finding that the wild-type hepatocytes produced more haptoglobin than KO hepatocytes, even in the absence of IL-6 (Fig. 5B).

Next, we cultured RAW cells, a murine macrophage cell line, in the presence or absence of culture supernatant of hepatocytes. RAW cells produced TNF- α , IL-6, and IL-10 but not IFN- γ upon stimulation of LPS, and hepatocyte culture supernatant suppressed the production of these cytokines (Fig. 5C). Importantly, the suppression was significantly weaker in the presence of conditional medium of KO hepatocytes than in the presence of conditional medium of wild-type hepatocytes.




Fig. 3. Organ injury in septic mice. (A) Serum ALT and creatinine levels in L-STAT3 KO mice (KO) and control mice (WT) 24 hours after CLP. * $P < 0.05$. (B) Representative histology (left part of each panel) and TUNEL (right part of each panel) of liver sections 24 hours after CLP or LPS injection. (C) Comparison of TUNEL-positive hepatocytes for at least 9 mice in each group. * $P < 0.05$.

cytes. Furthermore, murine primary splenocytes produced IFN- γ upon LPS stimulation, and the production was also suppressed in the presence of conditional medium of hepatocytes. Again, IFN- γ production was significantly higher in splenocytes cultured with KO hepatocyte supernatant than in those with wild-type hepatocyte supernatant (Fig. 5D). These data indicate that soluble substances from hepatocytes

suppressed activation of immune cells, which was critically dependent on STAT3.

Discussion

The present study clearly demonstrated that the absence of STAT3 in hepatocytes leads to high levels of circulating cytokines and increased mortality of CLP-in-




Fig. 4. Circulating cytokines before and after CLP or LPS injection. (A) L-STAT3 KO mice (KO) or wild-type littermates (WT) were treated with CLP (n = 8 in each group). Before and 24 hours after CLP, blood samples were obtained from mice and subjected to analysis of each cytokine indicated. *P < 0.05. (B) Mice were injected with 4 mg/kg of LPS (n = 8 in each group). Before and 24 hours after LPS injection, blood samples were obtained and subjected to the analysis of each cytokine indicated. *P < 0.05.

duced septic mice without affecting bacterial infection. L-STAT3 KO mice produced high levels of cytokines when injected with LPS, confirming that the absence of STAT3 signaling within hepatocytes induces a hyperinflammatory response even if the extent of the input stimuli remains constant. This phenomenon is similar to a previous report of macrophage-specific disruption of STAT3 in which serum cytokines such as TNF-α, IL-6, and IL-10 increased upon LPS stimulation.²² In those

mice, immune cells could not respond to IL-10, which potentially inhibit TNF-α production via STAT3 signaling, and thus produced high levels of TNF-α. Further study revealed those mice to be vulnerable to CLP-induced sepsis.^{23,24} However, in our L-STAT3 KO mice, the levels of STAT3 in macrophage did not differ from control mice and produced the same amount of TNF-α in response to LPS (Fig. 1C-D). Thus, suppression of the inflammatory response in wild-type mice was critically




Fig. 5. Suppression of cytokine production from immune cells by hepatocyte culture supernatant. Hepatocytes were isolated from L-STAT3 KO (KO) or wild-type (WT) mice and cultured in the presence or absence of IL-6 for 2 hours (for western blot analysis) or 48 hours (for collection of culture supernatants). (A) STAT3 phosphorylation (pSTAT3) and STAT3 expression of hepatocytes via western blot analysis. GAPDH expression served as a control. Representative blots are shown. (B) Haptoglobin production from primary hepatocytes. Haptoglobin concentration was determined in the hepatocyte supernatants via ELISA. Comparison of haptoglobin production between knockout hepatocytes and wild-type hepatocytes ($n = 5$ mice/group) cultured in the absence of IL-6. $*P < 0.05$. (C,D) Suppression of cytokine production in RAW cells or splenocytes by the hepatocyte supernatants. Hepatocytes were cultured for 48 hours. RAW cells (C) or splenocytes freshly isolated from wild-type mice (D) were cultured in the presence (KO or WT) or absence (Medium) of hepatocyte supernatants for 24 hours and then stimulated with LPS for another 24 hours. TNF- α , IL-6, IL-10, and IFN- γ production was determined via ELISA. $*P < 0.05$.

dependent on hepatic STAT3 signaling. Indeed, *in vitro* analysis revealed that soluble factors from hepatocytes repress cytokine production from activated macrophage and splenocytes in a hepatic STAT3-dependent manner. Whereas research has established that STAT3 mediates a variety of effects on hepatocytes, including proliferation,⁵ apoptosis protection,⁶ and glucose metabolism,⁷ the present study reveals that hepatic STAT3 has an important extrahepatic effect. This effect is activated by a variety of cytokines produced from immune cells such as IL-6 but, in turn, suppresses immune cell activation via production of soluble factors, providing a negative feedback loop. Thus, the present study describes a role of hepatic STAT3 in maintaining host homeostasis by negatively regulating the immune system.

APPs are liver plasma proteins whose levels of expression are either positively or negatively regulated by cytokines during inflammation. It has been established that STAT3 regulates the expression of most, if not all, APPs

in the liver.¹⁹ Consistent with this, L-STAT3 KO mice displayed impaired production of APPs in response to CLP (Fig. 2B). Some APPs such as C-reactive protein,²⁵ serum amyloid P,²⁶ and $\alpha 2$ -macroglobulin²⁷ have been shown to bind bacteria and to positively or negatively affect their eradication. Several reports also suggest that APPs exert proinflammatory as well as anti-inflammatory effects.^{25,28} C-reactive protein binds to the phosphocholine of some foreign pathogens as well as phospholipid constituents of damaged cells and can activate the complement system, whereas the antioxidants haptoglobin and hemopexin protect against reactive oxygen species. Thus, each APP has a unique role in the complex mechanism controlling infection-induced inflammation. The L-STAT3 KO mice used in the present study offer a unique model for identifying the net effect of STAT3-regulated APPs during the septic condition. Our work has revealed that the most prominent effect of STAT3-regulated APPs is suppression of the hyperinflammatory re-

sponse and lethality without an effect on bacterial infection. The soluble factors from hepatocytes that suppress cytokine production from immune cells are still unknown. Although there may be several substances involved in this phenomenon, one candidate might be haptoglobin, which was recently demonstrated to suppress TNF- α , IL-12, and IL-10 from human peripheral blood mononuclear cells *in vitro*.²⁹ We also obtained a similar finding that RAW cells produced a lesser amount of TNF- α upon LPS stimulation in the presence of haptoglobin (Supplementary Fig. 2). Identification of these substances may have important therapeutic implications for controlling the hyperinflammatory condition. Further study is needed to clarify this point.

The liver is one of the target organs of sepsis-induced multiple organ dysfunction syndrome. Evidence for this comes from the fact that CLP mice or LPS mice showed liver injury as evidenced by increases in serum ALT and TUNEL-positive hepatocytes scattered in the liver lobule. Furthermore, L-STAT3 KO mice displayed more hepatocyte apoptosis in mice subjected to CLP or LPS injection. Previous research has indicated that the absence of hepatic STAT3 renders hepatocytes more vulnerable to Fas-mediated apoptosis.⁶ It is possible that STAT3-null hepatocytes are more vulnerable to apoptosis in the septic model. However, at the same time, L-STAT3 KO mice showed higher levels of proinflammatory cytokines such as TNF- α , which is a direct inducer of hepatocyte apoptosis. In our model, it is difficult to differentiate which contributed more to increased liver injury: the decrease in apoptosis resistance or the increase in proinflammatory cytokine. It can be said that the increase of proinflammatory cytokines is presumably one of the causes, but not a result, of liver injury. In addition, as discussed in the Results section, liver injury was relatively modest and probably not a direct cause of animal death.

In the present study, the lack of hepatic STAT3 caused increased mortality in CLP mice. Although we did not address the direct link between hypercytokinemia and animal death, accumulating evidence suggests that an increase in a variety of cytokines is involved in lethality in CLP mice. For example, it was shown that IL-6 plays an important role in the increased expression of the C5a receptor in the lung, liver, kidney, and heart during the development of sepsis in CLP mice and that interception of IL-6 leads to reduced expression of the C5a receptor and improved survival.³⁰ In addition, enforced expression of the IL-6 gene in wild-type mice led to high mortality (unpublished data). TNF- α and other cytokines increase expression of inducible nitric oxide synthase, and increased production of nitric oxide causes further vascular instability and may also contribute to the direct myocar-

dial depression that occurs in sepsis.³¹ Thus, dysregulation of cytokines may be harmful for host organs and is probably linked to animal death.

The present study revealed an important role of hepatocytes in repressing the hyperinflammatory response in pathologic conditions. This raises the possibility that hyperinflammation may be ill-controlled when liver function is severely impaired. Although sepsis itself is not a frequent cause of liver failure, it is a serious complication of acute or chronic liver failure. Systemic inflammatory response syndrome is an important determinant of prognosis in fulminant hepatitis.¹² Sepsis originating from spontaneous bacterial peritonitis or renal infection is one of the causes of patient death with decompensated cirrhosis.¹³ In patients with limited function of the liver, possible impairment of STAT3-regulated hepatocyte function may be involved in their poor prognosis when complicated with severe inflammation. Careful liver-supporting therapy or early liver transplantation should be considered not only for maintaining liver function but also from the aspect of controlling dysregulated hyperinflammatory responses.

In conclusion, hepatic STAT3 represses systemic hyperinflammatory response by stimulating hepatic production of soluble substances that can attenuate immune cell overactivation and also improves host survival during septic condition. This sheds light on hepatocytic STAT3 as a negative regulator for immune cell overactivation and its role in host defense during systemic severe inflammation.

References

1. Akira S. IL-6-regulated transcription factors. *Int J Biochem Cell Biol* 1997; 29:1401-1418.
2. Akira S, Nishio Y, Inoue M, Wang XJ, Wei S, Matsusaka T, et al. Molecular cloning of APRF, a novel IFN-stimulated gene factor 3 p91-related transcription factor involved in the gp130-mediated signaling pathway. *Cell* 1994;77:63-71.
3. Zhong Z, Wen Z, Darnell JE Jr. Stat3: a STAT family member activated by tyrosine phosphorylation in response to epidermal growth factor and interleukin-6. *Science* 1994;264:95-98.
4. Takeda K, Noguchi K, Shi W, Tanaka T, Matsumoto M, Yoshida N, et al. Targeted disruption of the mouse Stat3 gene leads to early embryonic lethality. *Proc Natl Acad Sci U S A* 1997;94:3801-3804.
5. Li W, Liang X, Kellendonk C, Poli V, Taub R. STAT3 contributes to the mitogenic response of hepatocytes during liver regeneration. *J Biol Chem* 2002;277:28411-28417.
6. Haga S, Terui K, Zhang HQ, Enosawa S, Ogawa W, Inoue H, et al. Stat3 protects against Fas-induced liver injury by redox-dependent and -independent mechanisms. *J Clin Invest* 2003;112:989-998.
7. Inoue H, Ogawa W, Ozaki M, Haga S, Matsumoto M, Furukawa K, et al. Role of STAT-3 in regulation of hepatic gluconeogenic gene and carbohydrate metabolism *in vivo*. *Nat Med* 2004;10:168-174.
8. Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med* 2001;29:1303-1310.
9. Cohen J. The immunopathogenesis of sepsis. *Nature* 2002;420:885-891.

10. Hotchkiss RS, Karl IE. The pathophysiology and treatment of sepsis. *N Engl J Med* 2003;348:138-150.
11. Wang P, Chaudry IH. Mechanism of hepatocellular dysfunction during hyperdynamic sepsis. *Am J Physiol* 1996;270:R927-R938.
12. Rolando N, Wade J, Davalos M, Wendon J, Philpott-Howard J, Williams R. The systemic inflammatory response syndrome in acute liver failure. *HEPATOLOGY* 2000;32:734-739.
13. Fasolato S, Angeli P, Dallagnese L, Maresio G, Zola E, Mazza E, et al. Renal failure and bacterial infections in patients with cirrhosis: epidemiology and clinical features. *HEPATOLOGY* 2007;45:223-229.
14. Takeda K, Kaisho T, Yoshida N, Takeda J, Kishimoto T, Akira S. Stat3 activation is responsible for IL-6-induced T cell proliferation through preventing apoptosis: generation and characterization of T cell-specific Stat3-deficient mice. *J Immunol* 1998;161:4652-4660.
15. Takehara T, Tatsumi T, Suzuki T, Rucker EB 3rd, Hennighausen L, Jinushi M, et al. Hepatocyte-specific disruption of Bcl-xL leads to continuous hepatocyte apoptosis and liver fibrotic responses. *Gastroenterology* 2004;127:1189-1197.
16. Fink MP, Heard SO. Laboratory models of sepsis and septic shock. *J Surg Res* 1990;49:186-196.
17. Takehara T, Uemura A, Tatsumi T, Suzuki T, Kimura R, Shiotani A, et al. Natural killer cell-mediated ablation of metastatic liver tumors by hydrodynamic injection of IFN α gene to mice. *Int J Cancer* 2007;120:1252-1260.
18. Andrejko KM, Chen J, Deutschman CS. Intrahepatic STAT-3 activation and acute phase gene expression predict outcome after CLP sepsis in the rat. *Am J Physiol* 1998;275:G1423-G1429.
19. Alonzi T, Maritano D, Gorgoni B, Rizzuto G, Libert C, Poli V. Essential role of STAT3 in the control of the acute-phase response as revealed by inducible gene inactivation in the liver. *Mol Cell Biol* 2001;21:1621-1632.
20. Zhang Z, Fuentes NL, Fuller GM. Characterization of the IL-6 responsive elements in the gamma fibrinogen gene promoter. *J Biol Chem* 1995;270:24287-24291.
21. Kim H, Baumann H. The carboxyl-terminal region of STAT3 controls gene induction by the mouse haptoglobin promoter. *J Biol Chem* 1997;272:14571-14579.
22. Takeda K, Clausen BE, Kaisho T, Tsujimura T, Terada N, Forster I, et al. Enhanced Th1 activity and development of chronic enterocolitis in mice devoid of Stat3 in macrophages and neutrophils. *Immunity* 1999;10:39-49.
23. Matsukawa A, Takeda K, Kudo S, Maeda T, Kagayama M, Akira S. Aberrant inflammation and lethality to septic peritonitis in mice lacking STAT3 in macrophages and neutrophils. *J Immunol* 2003;171:6198-6205.
24. Matsukawa A, Kudo S, Maeda T, Numata K, Watanabe H, Takeda K, et al. Stat3 in resident macrophages as a repressor protein of inflammatory response. *J Immunol* 2005;175:3354-3359.
25. Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* 1999;340:448-454.
26. Noursadeghi M, Bickerstaff MC, Gallimore JR, Herbert J, Cohen J, Pepys MB. Role of serum amyloid P component in bacterial infection: protection of the host or protection of the pathogen. *Proc Natl Acad Sci U S A* 2000;97:14584-14589.
27. Hocheppied T, Van Leuven F, Libert C. Mice lacking alpha2-macroglobulin show an increased host defense against Gram-negative bacterial sepsis, but are more susceptible to endotoxemic shock. *Eur Cytokine Netw* 2002;13:86-91.
28. Tilg H, Dinarello CA, Mier JW. IL-6 and APPs: anti-inflammatory and immunosuppressive mediators. *Immunol Today* 1997;18:428-432.
29. Arredouani MS, Kasran A, Vanoirbeek JA, Berger FG, Baumann H, Ceuppens JL. Haptoglobin dampens endotoxin-induced inflammatory effects both in vitro and in vivo. *Immunology* 2005;114:263-271.
30. Riedemann NC, Neff TA, Guo RF, Bernacki KD, Laudes IJ, Sarma JV, et al. Protective effects of IL-6 blockade in sepsis are linked to reduced C5a receptor expression. *J Immunol* 2003;170:503-507.
31. Landry DW, Oliver JA. The pathogenesis of vasodilatory shock. *N Engl J Med* 2001;345:588-595.

Original Article

Early decline of hemoglobin can predict progression of hemolytic anemia during pegylated interferon and ribavirin combination therapy in patients with chronic hepatitis C

Naoki Hiramatsu,¹ Nao Kurashige,¹ Tsugiko Oze,¹ Tetsuo Takehara,¹ Shinji Tamura,¹ Akinori Kasahara,¹ Masahide Oshita,² Kazuhiro Katayama,³ Harumasa Yoshihara,⁴ Yasuharu Imai,⁵ Michio Kato,⁶ Sumio Kawata,⁷ Hirohito Tsubouchi,⁸ Hiromitsu Kumada,⁹ Takeshi Okanoue,¹⁰ Shinichi Kakumu¹¹ and Norio Hayashi¹

¹Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, Osaka, ²Osaka Police Hospital, Osaka, ³Osaka Kouseinenkin Hospital, Osaka, ⁴Osaka Rousai Hospital, Sakai, ⁵Ikeda Municipal Hospital, Ikeda, ⁶National Hospital Organization Osaka National Hospital, Osaka, ⁷Department of Gastroenterology, Yamagata University School of Medicine, Yamagata, ⁸Department of Digestive and Lifestyle Related Disease, Kagoshima University Graduate School of Medical and Dental Sciences, Kagoshima, ⁹Toranomon Hospital, Tokyo, ¹⁰Molecular Gastroenterology and Hepatology, Graduate School of Medical Science, Kyoto Prefectural University of Medicine, Kyoto and ¹¹Department of Internal Medicine, Division of Gastroenterology, Aichi Medical University School of Medicine, Aichi, Japan

Aim: Ribavirin, used to treat chronic hepatitis C, can induce hemolytic anemia, forcing the discontinuance of treatment. To establish a predictive measure to help circumvent this, we evaluated the relationship of hemoglobin (Hb) decline with the discontinuance of treatment during the progression of ribavirin-induced anemia.

Methods: One hundred and sixteen patients (71% male) with genotype 1 chronic hepatitis C were treated with pegylated interferon (PegIFN) α -2b and ribavirin. The mean age was 50.6 years and 55% were IFN naïve. A decline of Hb concentration by 2 g/dL at two weeks from the start of the treatment ("2 by 2" standard) was adopted as the predictive factor for the progression of anemia.

Results: By applying the "2 by 2" standard, with $\Delta\text{Hb} \geq 2$ g/dL (34%, $n = 39$), treatment was discontinued in 12 cases (31%), three of which (8%) because of severe anemia. For

$\Delta\text{Hb} < 2$ g/dL (64%, $n = 76$), treatment was discontinued in 11 (14%) cases; none due to severe anemia. Ten percent (4/39) of patients showed the minimum $\text{Hb} \leq 8.5$ g/dL in the $\Delta\text{Hb} \geq 2$ g/dL group, with none in the $\Delta\text{Hb} < 2$ g/dL group ($P = 0.001$). Furthermore, the patients with minimum $\text{Hb} \leq 8.5$ g/dL were found only in the "2 by 2" standard-positive and low CL/F (<15) group (4/29, 14%).

Conclusion: Monitoring the Hb decline using the "2 by 2" standard can identify patients who are prone to developing severe anemia. Further prospective studies are needed using ribavirin reduction based on the "2 by 2" standard.

Key words: "2 by 2" standard, chronic hepatitis C, pegylated interferon and ribavirin combination therapy, progression of anemia

INTRODUCTION

THE AIM OF antiviral therapy for hepatitis C virus (HCV) is to obtain a sustained viral response (SVR) and to reduce the occurrence rate of hepatocellular

carcinoma or hepatic disease-related mortality.^{1,2} The current optimal therapy for patients with chronic hepatitis C is a combination of pegylated interferon (PegIFN) and ribavirin. This combination can significantly improve the SVR rate and is recommended as a standard regimen worldwide.^{3–8} However, the SVR rates for the combination therapy of ribavirin with PegIFN for naïve patients with HCV genotype 1 has been reported to be 42–52%,^{6,9,10} which means that eradication of HCV is not complete in approximately half of these patients. Recently, long-term treatment¹¹ and a higher dosage

Correspondence: Dr Naoki Hiramatsu, Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita City, Osaka 565-0871, Japan.
Email: hiramatsu@gh.med.osaka-u.ac.jp
Received 29 December 2006; revision 11 April 2007; accepted 25 May 2007.

of drugs^{12,13} have been used to try to raise the SVR rate for patients with HCV genotype 1. However, it remains to be established what constitutes satisfactory efficacy. In this study we focused on a treatment strategy to enable the prediction of severe side-effects in order to avoid the need to discontinue treatment and raise the SVR rate by PegIFN and ribavirin combination therapy. It is important that ribavirin, the key drug for eradicating HCV, is continued until the end of treatment in order to attain the maximum SVR rate. Hemolytic anemia induced by ribavirin is known as one of the most important adverse effects in the combination therapy of PegIFN and ribavirin.^{14–17} To decrease the discontinuance rate of ribavirin due to severe anemia, epoetin alfa has been used for patients with progressing anemia, which can maintain the dose level of ribavirin as well as the quality of life of the patients.^{18–20} However, from a cost-effectiveness standpoint, it would be difficult for this treatment strategy to become standard. Also, side-effects other than anemia arising from an overload of ribavirin mainly due to renal dysfunction cannot be avoided by the additional administration of epoetin alfa.

Hemolysis induced by ribavirin has been suggested to be related to a high plasma concentration of ribavirin.²¹ The apparent clearance of ribavirin (CL/F), which reflects its plasma concentration at four weeks after the start of combination therapy, has been used as a predictive factor for ribavirin-induced hemolytic anemia before the start of treatment.^{22–24} However, the progression of hemolytic anemia occurs due not only to hemolysis, but also impaired hematogenous function. On the other hand, hemoglobin (Hb) dynamics directly reflect the degree of progression of anemia. We have reported that the early decline of Hb correlates with the progression of anemia during IFN and ribavirin combination therapy.²⁵ It is necessary to verify that a similar early predictor for the progression of anemia can be adopted in PegIFN and ribavirin combination therapy, since PegIFN is known to induce less depression of bone marrow function than usual IFN.

In this study, we evaluated the utility of the early decline of Hb in comparison with the CL/F to predict the progression of anemia in the combination therapy of PegIFN and ribavirin.

METHODS

Patients

THIS STUDY WAS conducted at 12 institutions in Japan. A total of 116 patients with chronic hepatitis C were enrolled and treated with a combination of

Table 1 Patient characteristics

Age (years)	50.6 ± 10.1 (24–70)
Gender (male/female)	82/34 (male 70.7%)
Body weight (kg)	64.5 ± 11.1
Previous IFN therapy (naïve/relapser/no responder)	64/38/14
HCV-RNA level (KIU/L) (<500/500–850/850<)	18/27/71
ALT (IU/L)	110 ± 60 (33–76)
Crn (mg/dL)	0.9 ± 0.2
Liver histology	
Fibrosis (F1/F2/F3/unknown)	35/49/31/1
Activity (A1/A2/A3/A4)	15/33/56/12
WBC (/mm ³)	5317 ± 1207
Neutrocytes (/mm ³)	2778 ± 902
Platelets (×10 ⁴ /mm ³)	17.4 ± 4.0
RBC (×10 ⁴ /mm ³)	459 ± 41
Hemoglobin (g/dL)	14.5 ± 1.2

Data are given as the mean ± SD.

ALT, alanine transaminase; RBC, red blood cells; WBC, white blood cells.

PegIFN and ribavirin. All patients were anti-hepatitis C virus antibody positive, had HCV-RNA detectable in their serum by the polymerase chain reaction (PCR) method, and showed elevated serum alanine transaminase (ALT) (above the upper limit of the normal), serum Hb concentration ≥12 g/dL, neutrocytes ≥1500/mm³ and platelets ≥10⁵/mm³ within six months before the treatment. Exclusion criteria were the presence of hepatitis B surface antigen, antihuman immunodeficiency virus antibody and other forms of liver disease (alcoholic liver disease, hepatotoxic drugs, autoimmune hepatitis).

The baseline characteristics of the patients are shown in Table 1. The mean age was 50.6 ± 10.1 years, and 71% (82 patients) were male. All patients had HCV-RNA with genotype 1 and high viral loads (more than 10⁵ copies/mL serum by Amplicor-HCV monitor assay). The mean ALT level was 110 ± 60 IU/L. Sixty-four patients (55%) were IFN naïve and the others were undergoing retreatment.

Treatment schedule

All patients were treated with a combination of PegIFN α-2b (Pegintron; Schering-Plough, Kenilworth, NJ, USA) and ribavirin (Rebetol; Schering-Plough) for 48 weeks. PegIFN was administered at a mean of 1.5 µg/kg body weight subcutaneously once a week. Ribavirin was given orally twice a day for the total dose. Dosages of both medications were decided based on the

body weight of the patients: those with a body weight of 40-60 kilograms (kg) were given PegIFN 75 µg/body and ribavirin 600 mg/day, those with a body weight of 60-80 kg were given PegIFN 105 µg/body and ribavirin 800 mg/day, and those with a body weight of 80-100 kg were given PegIFN 135 µg/body and ribavirin 1000 mg/day. The PegIFN dose was reduced by 50% if the neutrocyte count was below 750/mm³ or the platelet (Plt) count was below 8 × 10⁴/mm³. The PegIFN was discontinued if the neutrocyte count was below 500/mm³ or the Plt count was below 5.0 × 10⁴/mm³. The ribavirin dose of 200 mg was reduced when the Hb concentration decreased to less than 10 g/dL and the ribavirin was discontinued when the Hb concentration decreased to less than 8.5 g/dL, in accordance with the drug information for ribavirin. No ferric medicine or erythropoietin to prevent anemia was administered.

Patients with persistently undetectable HCV-RNA six-months after the end of treatment were considered to have achieved SVR.

Blood tests

All patients were examined for serum HCV-RNA level, hematological and biochemical tests just before therapy, at the end of week 2 and every four weeks during the treatment. When the treatment was completed, the patients were assessed every four weeks up to 24 weeks after the end of treatment.

Total ribavirin clearance

Using the method of Kamar *et al.*, CL/F at the start of the treatment was calculated as follows: CL/F (L/h) = 32.3 × BW × (1 - 0.0094 × age) × (1 - 0.42 × sex)/Scr (BW, body weight; sex = 0 for male and 1 for female; Scr = serum creatinine).¹⁷

Definition of "severe anemia" leading to the discontinuance of ribavirin

In this study, the "discontinuance of ribavirin due to severe anemia" was defined as follows: discontinuance of ribavirin due to a decrease of Hb to less than 8.5 g/dL or clinical symptoms of anemia associated with a decrease of Hb of more than 3 g/dL from the start of the combination therapy.

Statistical analysis

Age, body weight, ribavirin dosage/body weight, white blood cell count, red blood cell count, Hb concentration, Plt, serum ALT levels and serum creatinine are expressed as mean ± SD. The SVR rate was evaluated using the intention-to-treat analysis (ITT analysis). The

differences in proportions were tested by the χ^2 -test and Mantel-Haenszel χ^2 -test. A value of $P < 0.05$ (two-tailed) was considered to indicate significance. All calculations were performed by SAS program 9.1 (SAS Institute, Cary, NC, USA).

RESULTS

Frequency and reasons for dose reduction or discontinuance of PegIFN and/or ribavirin

OF THE 116 patients, 92 completed 48 weeks of therapy, but 24 patients (21%) had to discontinue both PegIFN and ribavirin. Thirty-nine patients (34%) completed the entire treatment schedule without reduction or discontinuance of either drug. The ribavirin dose was decreased for 39 patients (34%) and the PegIFN dose was decreased for 33 patients (28%), including 19 patients for whom both drugs had to be reduced. The reasons for discontinuance of both drugs included anemia, thyroid dysfunction, skin eruption and neutropenia, with the major reasons being anemia (17%) and thyroid dysfunction (17%).

Efficacy of the combination therapy with dose reduction or discontinuance of PegIFN and/or ribavirin

The SVR rate was 57% (66/116) for all according to ITT analysis. According to the category of response to previous IFN therapy, the SVR rates were 43% (6/14) in




Figure 1 SVR rate due to PegIFN/ribavirin dose reduction or discontinuance. (□), All patients; (▨), patients without dose reduction; (▩), patients with dose reduction; (■), patients with drug discontinuance. Significant levels: * $P = 0.003$; ** $P = 0.001$.

Table 2 Rate of the ribavirin reduction or discontinuance due to adverse effects according to CL/F level

	No reduction	Dose reduction	Discontinuance	
			All cases	Cases due to severe anemia
20 ≤ CL/F (n = 12)	67% (8/12)	25% (3/12)	8% (1/12)	0
15 ≤ CL/F < 20 (n = 23)	57% (13/23)	30% (7/23)	13% (3/23)	0
10 ≤ CL/F < 15 (n = 39)	46% (18/39)	31% (12/39)	23% (9/39)	5% (2/39)
CL/F < 10 (n = 42)	33% (14/42)	40% (17/42)	26% (11/42)	5% (2/42)

$P = 0.031$ (Mantel-Haenszel χ^2 -test).

Table 3 Minimum hemoglobin levels during PegIFN/ribavirin combination therapy according to CL/F level

	10 g/dL < Hb	8.5 < Hb ≤ 10 g/dL	Hb ≤ 8.5 g/dL
20 ≤ CL/F (n = 12)	92% (11/12)	12% (1/12)	0
15 ≤ CL/F < 20 (n = 23)	83% (19/23)	17% (4/23)	0
10 ≤ CL/F < 15 (n = 39)	72% (28/39)	23% (9/39)	5% (2/39)
CL/F < 10 (n = 42)	50% (21/42)	43% (18/42)	7% (3/42)

$P = 0.009$ (Mantel-Haenszel χ^2 -test).

non-responders, 61% (23/38) in relapsers, and 58% (37/64) in naïve patients. The relationship between dose reduction or discontinuance of PegIFN and ribavirin and the SVR rate on ITT analysis is shown in Figure 1. Similar SVR rates were obtained in the groups without dose reduction of PegIFN and ribavirin (64%, 25/39) and with reduction of PegIFN and/or ribavirin (66%, 35/53); in detail, the SVR rate was 79% (11/14) in the group with reduction of only PegIFN, 55% (11/20) with reduction of only ribavirin, and 63% (12/19) with reduction of both PegIFN and ribavirin. In the group where both drugs were discontinued, the SVR rate was 25% (6/24), significantly lower than the group without reduction of both drugs ($P = 0.003$), and the group with reduction of PegIFN and/or ribavirin ($P = 0.001$).

CL/F and dose reduction or discontinuance of ribavirin

CL/F calculated for all patients showed a median of 12.6 L/h (range 4.5–27.9). At the start of the treatment, 36% (42/116) were under 10 L/h, 34% (39/116) were 10–15 L/h, 20% (23/116) were 15–20 L/h and 10% (12/116) were 20 L/h or more.

The rate of dose reduction or discontinuance of ribavirin is shown in Table 2 for different levels of CL/F. The rate of discontinuance of ribavirin in all cases was 8% (1/12) for the CL/F ≥ 20, 13% (3/23) for the 15 ≤ CL/F < 20, 23% (9/39) for the 10 ≤ CL/F < 15, and

26% (11/42) for the CL/F < 10 group. Ribavirin did not have to be discontinued due to severe anemia among patients with 15 ≤ CL/F, but did for the 18% (2/11) of those with CL/F < 10 and 22% (2/9) of those with 10 ≤ CL/F < 15. The rate of reduction and discontinuance of ribavirin correlated significantly with the CL/F level.

CL/F and minimum hemoglobin level during treatment

To examine the relationship between anemia and the cessation of ribavirin in further detail, we evaluated the minimum hemoglobin level during treatment. Table 3 presents the different levels in relation to CL/F. The patients with minimum Hb ≤ 8.5 g/dL, the criterion for discontinuance of ribavirin, accounted for 7% (3/42) of the group of CL/F < 10, and 5% (2/39) of the group of 10 ≤ CL/F < 15. No patients of the group of CL/F ≥ 15 showed minimum Hb ≤ 8.5 g/dL.

Early decline of Hb and progression of anemia during combination therapy

Following the initiation of combination therapy, the Hb concentration decreased rapidly until the end of four-weeks. At the end of two weeks, Hb had decreased by 1.1 ± 1.0 g/dL among the patients without dose reduction of ribavirin ($n = 53$), 1.6 ± 1.2 g/dL among those with dose reduction ($n = 39$), and 1.8 ± 1.0 g/dL among




Figure 2 Course of ΔHb in the initial phase. (---), No reduction; (-.-.), reduction; (—), discontinuance. *Significantly different between patients with discontinuance and patients with no reduction ($P = 0.04$). **Significantly different between patients with discontinuance and patients with no reduction ($P = 0.008$), and between patients with discontinuance and patients with reduction ($P = 0.003$).

those who had discontinued ribavirin ($n = 24$). It was significantly different between the patients with no reduction and those with discontinuance of therapy ($P = 0.04$). At the end of four weeks, Hb had decreased by 1.9 ± 1.2 g/dL among the patients without dose reduction of ribavirin, 2.8 ± 1.2 g/dL among those with dose reduction, and 2.9 ± 1.2 g/dL among those who had discontinued ribavirin. Hb decline at the end of four weeks was significantly greater in the patients who had discontinued treatment and those who had reduced it, than in those with no reduction ($P = 0.008$, $P = 0.003$, respectively) (Fig. 2).

In this study, we selected the Hb decrease at the end of two weeks as the predictive factor for anemia progression. This is because the judgment of Hb decrease at the end of four weeks is too late to prevent progression of anemia or to perform appropriate counter-measures, such as the administration of epoetin or reduction of ribavirin. Next, we tried to use two borderlines of ΔHb:

ΔHb 2.0 indicates a 2 g/dL Hb decrease at the end of two weeks and ΔHb 1.5 indicates a 1.5 g/dL Hb decrease. When ΔHb 2.0 was adopted, the rate of discontinuance of drugs was 31% (12/39) in the ΔHb ≥ 2.0 and 14% (11/76) in the ΔHb < 2.0 . When ΔHb 1.5 was adopted, it was 23% (14/60) in the ΔHb ≥ 1.5 and 16% (9/55) in the ΔHb < 1.5 . Comparison of the ΔHb 2.0 and ΔHb 1.5 standards showed the sensitivity to be 52% (12/23) and 61% (14/23), and the specificity to be 71% (65/92) and 50% (46/92), respectively. With respect to discontinuance due to anemia, both ΔHb 2.0 and ΔHb 1.5 gave 100% sensitivity (3/3), and the specificities were 68% (76/112) using ΔHb 2.0 and 49% (55/112) using ΔHb 1.5. We decided to adopt the standard of ΔHb 2 g/dL at the end of two weeks from the start of the pegylated IFN and ribavirin combination therapy as the predictive factor for anemia progression ("2 by 2" standard), which has been taken as a predictive factor for anemia in the IFN and ribavirin combination therapy.²⁵

Applying the "2 by 2" standard to PegIFN plus ribavirin combination therapy, the rate of reduction or discontinuance of the ribavirin dose was examined with respect to the Hb decrease level (Table 4). Only one patient was excluded from this study, because the treatment was discontinued on the 11th day. In the group of ΔHb (the decrease in Hb concentration at two weeks from the baseline) ≥ 2 g/dL ($n = 39$), the doses were reduced for 18 patients (46%) and discontinued for 12 (31%), three of whom (8%) had severe anemia. For the group of ΔHb < 2 g/dL (76 patients), the doses were reduced for 21 patients (28%) and discontinued for 11 (14%); none due to severe anemia.

Early decline of Hb and minimum hemoglobin level during treatment

As in the case of ΔHb, we evaluated the minimum hemoglobin level during treatment, as shown in Figure 3. The patients with minimum Hb ≤ 8.5 g/dL accounted for 10% (4/39) of the group of ΔHb ≥ 2 g/dL, and there was no patient with minimum Hb ≤ 8.5 g/dL

Table 4 Rate of the ribavirin reduction or discontinuance due to adverse effects according to Hb decrease levels

	No reduction	Dose reduction	Discontinuance	
			All cases	Cases due to severe anemia
ΔHb < 2 g/dL ($n = 76$)	58% (44/76)	28% (21/76)	14% (11/76)	0
ΔHb ≥ 2 g/dL ($n = 39$)	23% (9/39)	46% (18/39)	31% (12/39)	8% (3/39)

$P = 0.004$ (Mantel-Haenszel χ^2 -test).




Figure 3 Minimum hemoglobin levels during PegIFN/ribavirin combination therapy. (□), 10 g/dL < minimum Hb; (▨), 8.5 < minimum Hb ≤ 10 g/dL; (■), minimum Hb ≤ 8.5 g/dL. (a) According to the "2 by 2" standard (Hb 2 g/dL decrease at two weeks from the baseline). $P = 0.009$ (Mantel-Haenszel χ^2 -test). (b) according to CL/F levels. $P = 0.001$ (Mantel-Haenszel χ^2 -test).

in the $\Delta Hb < 2 \text{ g/dL}$ group (Fig. 3a). The patients with minimum Hb ≤ 8.5 g/dL accounted for 6% (5/81) of the group of $CL/F < 15$, and there was no patient with minimum Hb ≤ 8.5 g/dL in the $15 \leq CL/F$ group (Fig. 3b). The number of patients with minimum Hb ≤ 8.5 g/dL during PegIFN and ribavirin combination therapy according to "2 by 2" standard and CL/F levels is shown in Table 5. The patients with minimum Hb ≤ 8.5 g/dL were found only in the "2 by 2" standard-positive and low CL/F (<15) group (4/29, 14%).

DISCUSSION

PREDICTION OF THE progression of anemia is necessary to decide whether drugs can be continued, with minimization of the disadvantages induced by anemia. Recently, CL/F has been used as a marker of

Table 5 The number of patients with minimum hemoglobin ≤ 8.5 g/dL during PegIFN/ribavirin combination therapy according to "2 by 2" standard and CL/F levels

	$\Delta Hb < 2 \text{ g/dL}$ (n = 76)	$\Delta Hb \geq 2 \text{ g/dL}$ (n = 39)
$CL/F \geq 15$ (n = 35)	0/25	0/10
$CL/F < 15$ (n = 80)	0/51	4/29 (14%)

progressing anemia that necessitates discontinuance of treatment. For example, if the patients have a low CL/F level, they should start treatment with a low ribavirin dose. In this study, we attempted to use the CL/F level measurement for our patients. To predict which patients might have to discontinue the treatment, the target range had to be $CL/F < 15$ because 6% of patients (n = 5) in this range showed minimum Hb ≤ 8.5 g/dL, which is the level at which ribavirin should be discontinued. No patients of the $CL/F \geq 15$ group showed minimum Hb ≤ 8.5 g/dL. Our findings showed that 70% of the patients (81/116) with $CL/F < 15$ should be discriminated from the others (Table 3). In the same manner, using ΔHb as the marker, 34% of the target patients in the $\Delta Hb \geq 2 \text{ g/dL}$ group were identified because 10% in this range showed minimum Hb ≤ 8.5 g/dL. No patients in the $\Delta Hb < 2 \text{ g/dL}$ group showed minimum Hb ≤ 8.5 g/dL. Compared to CL/F, ΔHb is considered to be more sensitive and convenient for identifying the high risk patients for whom treatment would need to be discontinued. Furthermore, the application of "2 by 2" standard in the group with low level of $CL/F < 15$ can be the most sensitive method for this (Table 5), since no patients with progression of anemia were found in the "2 by 2" standard-negative group with $CL/F < 15$.

In Japan, ribavirin doses are set at 600 mg for <60 kg, 800 mg for 60-80 kg, and 1000 mg for ≥80 kg, which are lower doses than those used in Europe and the USA. In this study, the mean ribavirin level at the start of treatment was 743 mg per day, while the AASLD practice guideline for genotype 1 hepatitis C is a daily dose of 1000 mg for body weight ≤ 75 kg and 1200 mg if >75 kg²⁶. In Japan, the use of lower doses is why fewer patients treated with PegIFN and ribavirin combination therapy are forced to discontinue the treatment due to severe anemia. Since the "2 by 2" model and/or CL/F can identify the patients who are prone to develop severe anemia, the other patients could be candidates for ribavirin dose-up strategies to raise SVR rates.

A considerable number of patients with chronic hepatitis C are over 60 years old in Japan (mean age is