

| | | | |
|--|---|-------------------|----------------------|
| Akiyama M, Sakai K, Ogawa M, McMillan JR, Sawamura D, <u>Shimizu H</u> | Novel duplication mutation in the patatin domain of adipose triglyceride lipase (PNPLA2) in neutral lipid storage disease with severe myopathy | Muscle Nerve | 36(6):856-9,2007 |
| Aoyagi S, Akiyama M, Mashiko M, Shibaki A, <u>Shimizu H</u> | Extensive proliferative nodules in a case of giant congenital naevus | Clin Exp Dermatol | 33(2):125-7,2008 |
| Natsuga K, Akiyama M, Kato N, Sakai K, Sugiyama-Nakagiri Y, Nishimura M, Hata H, Abe M, Arita K, Tsuji-Abe Y, Onozuka T, Aoyagi S, Kodama K, Ujiie H, Tomita Y, <u>Shimizu H</u> | Novel ABCA12 Mutations Identified in Two Cases of Non-Bullous Congenital Ichthyosiform Erythroderma Associated with Multiple Skin Malignant Neoplasia | J Invest Dermatol | 127(11):2669-73,2007 |
| Olasz EB, Roh J, Yee CL, Arita K, Akiyama M, <u>Shimizu H</u> | Vogel JC, Yancey KB: Human Bullous Pemphigoid Antigen 2 Transgenic Skin Elicits Specific IgG in Wild-Type Mice | J Invest Dermatol | in press |
| Shinkuma S, Nishie W, Shibaki A, Sawamura D, Ito K, Tsuji-Abe Y, Natsuga K, Chan P, <u>Amagai M, Shimizu H</u> | Cutaneous type pemphigus vulgaris with skin manifestations similar to the classical mucocutaneous type: a case report and review of the literature | Clin Exp Dermatol | in press |
| Tsubota A, Akiyama M, Sakai K, Yanagi T, McMillan J, Higashi A, <u>Shimizu H</u> | Congenital ichthyosiform erythroderma mimicking ichthyosis bullosa of Simens | Br J Dermatol | in press |
| Yasukawa K, Sawamura D, Goto M, Nakamura H, <u>Shimizu H</u> | Histone deacetylase inhibitors preferentially augment transient transgene expression in human dermal fibroblasts | Br J Dermatol | 157(4):662-9,2007 |
| Abe M, Sawamura D, Goto M, Nakamura H, Nagasaki A, Nomura Y, Kawasaki H, Isogai R, <u>Shimizu H</u> | ITGB4 missense mutation in a transmembrane domain causes non-lethal variant of junctional epidermolysis bullosa with pyloric atresia | J Dermatol Sci | 47:165-7, 2007 |
| Akiyama M, Sakai K, Arita K, Nomura Y, Ito K, Kodama K, McMillan JR, Kobayashi K, Sawamura D, <u>Shimizu H</u> | A novel GJB2 mutation p.Asn54His in a patient with palmoplantar keratoderma, sensorineural hearing loss and knuckle pads | J Invest Dermatol | 127:1540-3, 2007 |
| Akiyama M, Sakai K, Sato T, McMillan JR, Goto M, Sawamura D, <u>Shimizu H</u> | Compound heterozygous ABCA12 mutations including a novel nonsense mutation underlie harlequin ichthyosis | Dermatology | 215:155-9, 2007 |
| Akiyama M, Titeux M, Sakai K, McMillan JR, Tonasso L, Calvas P, Jossic F, Hovnanian A, <u>Shimizu H</u> | DNA-based prenatal diagnosis of harlequin ichthyosis and characterization of ABCA12 mutation consequences | J Invest Dermatol | 127:568-73, 2007 |

| | | | |
|---|--|-------------------|------------------|
| Arita K, Jacyk WK, Wessagowit V, van Rensburg EJ, Chaplin T, Mein CA, Akiyama M, Shimizu H, Happle R, McGrath JA | The South African "bathing suit ichthyosis" is a form of lamellar ichthyosis caused by a homozygous missense mutation, p.R315L, in transglutaminase 1 | J Invest Dermatol | 127:490-3, 2007 |
| Chen KR, Sakamoto M, Ikemoto K, Abe R, Shimizu H | Granulomatous arteritis in cutaneous lesions of Churg-Strauss syndrome | J Cutan Pathol | 34:330-7, 2007 |
| Hoshina D, Akiyama M, Hamasaka K, Shimizu H | An infantile case of pityriasis lichenoides et varioliformis acuta | Br J Dermatol | 157:194-6, 2007 |
| Hoshina D, Akiyama M, Hata H, Aoyagi S, Sato-Matsumura KC, Shimizu H | Eccrine porocarcinoma and Bowen's disease arising in a seborrheic keratosis | Clin Exp Dermatol | 32:54-6, 2007 |
| Hoshina D, Sawamura D, Nomura T, Tanimura S, Abe M, Onozuka T, Kodama K, Akiyama M, Shimizu H | Epidermolysis bullosa acquisita associated with psoriasis vulgaris | Clin Exp Dermatol | 32:516-8, 2007 |
| Hoshina D, Shibaki A, Aoyagi S, Kimura K, Shimizu H | Giant dermatofibroma: a rare variant of dermatofibroma preferentially developing on the lower limbs | Clin Exp Dermatol | 32:132-134, 2007 |
| Ito H, Akiyama M, Nakagawa H, Uematsu R, Deguchi K, McMillan JR, Nishimura S, Shimizu H | N-Linked neutral oligosaccharides in the stratum corneum of normal and ichthyotic skin | Arch Dermatol Res | 298:403-7, 2007 |
| McMillan JR, Akiyama M, Rouan F, Mellerio JE, Lane EB, Leigh IM, Owaribe K, Wiche G, Fujii N, Uitto J, Eady RA, Shimizu H | Plectin defects in epidermolysis bullosa simplex with muscular dystrophy | Muscle Nerve | 35:24-35, 2007 |
| McMillan JR, Akiyama M, Tanaka M, Yamamoto S, Goto M, Abe R, Sawamura D, Shimomura M, Shimizu H | Small-diameter porous poly (epsilon-caprolactone) films enhance adhesion and growth of human cultured epidermal keratinocyte and dermal fibroblast cells | Tissue Eng | 13:789-98, 2007 |
| Moriuchi R, Shibaki A, Yasukawa K, Onozuka T, Sato T, Kaneda M, Iguchi A, Kobayashi R, Shimizu H | Neonatal vesiculopustular eruption of the face: a sign of trisomy 21-associated transient myeloproliferative disorder | Br J Dermatol | 156:1373-4, 2007 |
| Motoda N, Akiyama M, Aoyagi S, Sawamura D, Shimizu H | Low-grade myxofibrosarcoma invaded into the underlying skeletal muscle | J Dermatol | 34:561-4, 2007 |
| Natsuga K, Abe R, Ujiie H, Shibaki A, Sawamura D, Nishio M, Fujimoto K, Koike T, Shimizu H | Non-Hodgkin lymphoma preceded by recalcitrant eczema | Eur J Haematol | 79:369-370, 2007 |

| | | | |
|--|---|------------------------|--------------------|
| Natsuga K, Sawamura D, Homma E, Nomura T, Abe M, Muramatsu R, Mochizuki T, Koike T, <u>Shimizu H</u> | Amicrobial pustulosis associated with IgA nephropathy and Sjogren's syndrome | J Am Acad Dermatol | 57:523-6, 2007 |
| Nemoto I, Shimizu T, Fujita Y, Tateishi Y, Tsuji-Abe Y, <u>Shimizu H</u> | Tumour-like muscular sarcoidosis | Clin Exp Dermatol | 32:298-300, 2007 |
| Nishie W, Sawamura D, Goto M, Ito K, Shibaki A, McMillan JR, Sakai K, Nakamura H, Olasz E, Yancey KB, Akiyama M, <u>Shimizu H</u> | Humanization of autoantigen | Nat Med | 13:378-83, 2007 |
| Nomura T, Sandilands A, Akiyama M, Liao H, Evans AT, Sakai K, Ota M, Sugiura H, Yamamoto K, Sato H, Palmer CN, Smith FJ, McLean WH, <u>Shimizu H</u> | Unique mutations in the filaggrin gene in Japanese patients with ichthyosis vulgaris and atopic dermatitis | J Allergy Clin Immunol | 119:434-40, 2007 |
| Sawamura D, Goto M, Sakai K, Nakamura H, McMillan JR, Akiyama M, Shirado O, Oyama N, Satoh M, Kaneko F, Takahashi T, Konno H, <u>Shimizu H</u> | Possible involvement of exon 31 alternative splicing in phenotype and severity of epidermolysis bullosa caused by mutations in PLEC1 | J Invest Dermatol | 127:1537-40, 2007 |
| Tsubota A, Akiyama M, Sakai K, Goto M, Nomura Y, Ando S, Abe M, Sawamura D, <u>Shimizu H</u> | Keratin 1 gene mutation detected in epidermal nevus with epidermolytic hyperkeratosis | J Invest Dermatol | 127:1371-4, 2007 |
| Yamanaka Y, Akiyama M, Sugiyama-Nakagiri Y, Sakai K, Goto M, McMillan JR, Ota M, Sawamura D, <u>Shimizu H</u> | Expression of the keratinocyte lipid transporter ABCA12 in developing and reconstituted human epidermis | Am J Pathol | 171:43-52, 2007 |
| Yamane N, Sawamura D, Nishie W, Abe M, Kodama K, Adachi K, Nakamura H, Ishii N, <u>Hashimoto T</u> , <u>Shimizu H</u> | Anti-p200 pemphigoid in a 17-year-old girl successfully treated with systemic corticosteroid and dapsone | Br J Dermatol | 156:1075-8, 2007 |
| Yaosaka M, Abe R, Ujiie H, Abe Y, <u>Shimizu H</u> | Unilateral periorbital oedema due to sarcoid infiltration of the eyelid: an unusual presentation of sarcoidosis with facial nerve palsy and parotid gland enlargement | Br J Dermatol | 157:200-202, 2007 |
| Yasukawa K, Sawamura D, Goto M, Nakamura H, <u>Shimizu H</u> | Histone deacetylase inhibitors preferentially augment transient transgene expression in human dermal fibroblasts | Br J Dermatol | 157:662-9, 2007 |
| Morita S, Shirakata Y, Shiraishi A, Kadota Y, <u>Hashimoto K</u> , Higashiyama S, Ohashi Y | Human corneal epithelial cell proliferation by epiregulin and its cross-induction by other EGF family members | Molecular Vision | 13:2119-2128, 2007 |

| | | | |
|---|---|-------------------|---------------------|
| Dai X, Sayama K, Shirakata Y, Tokumaru S, Yang L, Tohyama M, Hirakawa S, Hanakawa Y, <u>Hashimoto K</u> | PPARgamma is an important transcription factor in 1alpha,25-dihydroxyvitamin D3-induced involucrin expression | J Dermatol Sci | 50:53-60,2008 |
| 永井弥生, 荻野幸子, 石川 治, 是枝哲, 宮地良樹, 藤本晃英, 竹原和彦, 宮脇さおり, 橋本公二, 六戸亜希子, 花田勝美 | アトピーセラ－ME－01の臨床効果および基礎的検討 | 臨床皮膚科 | 61 : 286-295, 2007 |
| 青木恵美, 永松将吾, 中岡啓喜, 大塚壽, 橋本公二 | 皮膚に原発して紡錘形細胞型有棘細胞癌の1例 | 形成外科 | 50 : 89-96, 2007 |
| 井上有紀子, 八幡陽子, 村上信司, 橋本公二 | 好酸球増多症候群に伴い多発性指尖潰瘍を生じた1例 | 皮膚科の臨床 | 49 : 179-183, 2007 |
| 白石 研, 村上信司, 橋本公二 | 減汗症を伴ったコリン性蕁麻疹の1例 | 西日本皮膚科 | 4 : 359-364, 2007 |
| 藤山幹子, 橋本公二 | 【薬剤アレルギーの病態・診断・治療】中毒性表皮壊死剥離症とは? | Q&Aでわかるアレルギー疾患 | 3 : 77-78, 2007 |
| 橋本公二 | 薬剤性過敏症候群 (DIHS) - その発見と展開 - | マルホ皮膚科セミナー | 186 : 4-6, 2007 |
| 橋本公二 | 【重症薬疹】重症薬疹とは | アレルギー・免疫 | 14 : 414-421, 2007 |
| 藤山幹子, 橋本公二 | 【特集/広がる薬疹の世界 最新の概念・病態・治療】DIHSの診断におけるHHV-6再活性化の意義 HS, DRESS, DIHSの疾患概念 | 医学のあゆみ | 220 : 885-888, 2007 |
| 橋本公二, 竹田 潔 | 皮膚と自然免疫 | 皮膚アレルギーフロンティア | 2 : 98-105, 2007 |
| Dai X, Sayama K, Shirakata Y, Hanakawa Y, Yamasaki K, Tokumaru S, Yang L, Wang X, Hirakawa S, Tohyama M, Yamauchi T, Takashi K, Kagechika H, <u>Hashimoto K</u> | STAT5a/PPAR gamma pathway regulates involucrin expression in keratinocyte differentiation | J Invest Dermatol | 127:1728-1735, 2007 |
| Nagai H, Tokumaru S, Sayama K, Shirakata Y, Hanakawa Y, Hirakawa S, Dai X, Tohyama M, Yang L, <u>Hashimoto K</u> | Suppressor of cytokine signaling 3 negative regulation of signal transducer and activator of transcription 3 in platelet-derived growth factor-induced fibroblast migration | J Dermatol | 34:523-530, 2007 |
| Nanba D, Inoue H, Shigemi Y, Shirakata Y, <u>Hashimoto K</u> , Higashiyama S | An intermediary role of proHB-EGF shedding in growth factor-induced c-Myc gene expression | J Cell Physiol | 214(2):465-73,2008 |
| Tohyama M, <u>Hashimoto K</u> , Yasukawa M, Kimura H, Horikawa T, Nakajima K, Urano Y, Matsumoto K, Iijima M, Shear NH | Association of human herpesvirus 6 reactivation with the flaring and severity of drug-induced hypersensitivity syndrome | Br J Dermatol | 157(5)934-40,2007 |

| | | | |
|---|--|---------------------------------------|---------------------|
| Tohyma M, Sayama K, Komatsuzawa H, Hanakawa Y, Shirakata Y, Dai X, Yang L, Tokumaru S, Nagai H, Hirakawa S, Sugai M, <u>Hashimoto K</u> | CXCL 16 is a novel mediator of the innate immunity of epidermal keratinocytes | Int Immunol | 19:1095-1102, 2007 |
| Shiraishi K, Yamasaki K, Nanba D, Inoue H, Hanakawa Y, Shirakata Y, <u>Hashimoto K</u> , Higashiyama S | Pre-B-cell leukemia transcription factor 1 is a major target of promyelocytic leukemia zinc-finger-mediated melanoma cell growth suppression | Oncogene | 26:339-48, 2007 |
| Shirakata Y, Kishimoto J, Tokumaru S, Yamasaki K, Hanakawa Y, Tohyama M, Sayama K, <u>Hashimoto K</u> | Epiregulin, a member of the EGF family, is over-expressed in psoriatic epidermis | J Dermatol Sci | 45:69-72, 2007 |
| Ehama R, Ishimatsu-Tsuji Y, Iriyama S, Ideta R, Soma T, Yano K, Kawasaki C, Suzuki S, Shirakata Y, <u>Hashimoto K</u> , Kishimoto J | Hair follicle regeneration using grafted rodent and human cells | J Invest Dermatol | 127:2106-15, 2007 |
| Saga, K., Tamai, K., Kawachi, M., Shimbo, T.; Fujita, H., Yamazaki, T., and <u>Kaneda, Y</u> | Functional modification of Sendai virus by siRNA | J. Biotechnology | 133:386-394,2008 |
| Otsuru, S., Tamai, K., Yamazaki, T., Yoshikawa, H., and <u>Kaneda, Y</u> | Circulating bone marrow-derived osteoblast progenitor cells are recruited to the bone-forming site by CXCR4/SDF1 pathway | Stem Cells | 26(1):223-34,2008 |
| Takami, Y., Nakagami, H., Morishita, R., Katsuya, T.,m Cui, T=X, Ichikawa, T., Saito, Y., Hayashi, H., Kikuchi, Y., Nishikawa, T., Baba, Y., Yasudda, O., Rakugi, H., Ogihara, T., and <u>Kaneda, Y</u> | Ubiquitin carboxy-terminal hydrolase L1, a novel deubiquitinating enzyme in the vasculature, attenuates NF-kB activation | Arterioscler. Thromb. Vasc. Biol | 27:2184-2190, 2007 |
| Shimizu, N., Azuma, N., Nishikawa, T., Hirata, S., Morishita, R., <u>Kaneda, Y</u> , and Sasajima, T | Effect on vein graft intimal hyperplasia of nuclear factor-kB decoy transfection using the second generation of HVJ vector | The Journal of Cardiovascular Surgery | 48, 463-470, 2007 |
| Son, G., Iimuro, Y., Seki, E., Hirano, T., <u>Kaneda, Y</u> , and Fujimoto, J | Selective inactivation of NF-kB in the liver using NF-kB decoy suppresses CCL4-induced liver injury and fibrosis | Am. J. Phys. Gast. Liver Phys | 293, G631-639, 2007 |
| Shimbo, T., Kawachi, M., Saga, K., Fujita, H., Yamazaki, T., <u>Tamai, K.</u> , and <u>Kaneda, Y</u> | Development of a transferrin receptor-targeting HVJ-E vector | Biochem. Biophys. Res. Comm | 364(3):423-8,2007 |
| Kawachi, M., <u>Tamai, K.</u> , Saga, K., Yamazaki, T., Fujita, H., Shimbo, T., Nimura, K., Nishifuji, K., <u>Amagai, M.</u> , Uitto, J., and <u>Kaneda, Y</u> | Development of tissue-targeting HVJ envelope vector for successful delivery of therapeutic gene to mouse skin | Human Gene Therapy | 18, 881-894, 2007 |

| | | | |
|--|--|---------------------------------|----------------------|
| Hayashi, H., Nakagami, H., Takami, Y., Sato, N., Saito, Y., Nishikawa, T., Mori, M., Koriyama, H., <u>Tamai, K.</u> , Morishita, R., and <u>Kaneda, Y</u> | Involvement of g-secretase in postnatal angiogenesis | Biochem. Biophys. Res. Comm | in press |
| Kawano, H., Komaba, S., Kanamori, T., and <u>Kaneda, Y</u> | A new therapy for highly effective tumor eradication using HVJ-E combined with chemotherapy | BMC Medicine | in press |
| Saito, Y., Nakagami, H., Kurooka, M., Takami, Y., Kikuchi, Y., Hayashi, H., Nishikawa, T., Tamai, K., Morishita, R., Azuma, N., Sasajima, T., and <u>Kaneda, Y</u> | Cold shock domain protein A represses angiogenesis and lymphangiogenesis via inhibition of serum response element | Oncogene | in press |
| Fujihara, A., Kurooka, M., Miki, T., and <u>Kaneda, Y</u> | Intratumoral injection of inactivated Sendai virus particles elicits strong antitumor activity by enhancing local CXCL10 expression and systemic NK cell activation | Cancer Immunology Immunotherapy | 57(1):73-84,2008 |
| Kawano, H., Komaba, S., Yamasaki, T., Maeda, M., Kimura, Y., Maeda, A., and <u>Kaneda, Y</u> | New potential therapy for orthotopic bladder carcinoma by combining HVJ envelope with doxorubicin | Cancer Chemother. Pharmacol | in press |
| Izumo, T., Otsuru, A., Tokunaga, Y., Namba, H., <u>Kaneda, Y.</u> , Nagata, I., and Yamashita, S | Epstein-Barr virus-based vector improves the tumor cell killing effect of pituitary tumor in HVJ-liposome-mediated transcriptional targeting suicide gene therapy | Int J Oncol | 31(2):379-87,2007 |
| Yasuoka, E., Oshima, K., <u>Tamai, K.</u> , Kubo, T., and <u>Kaneda, Y</u> | Needleless intranasal administration of HVJ-E containing allergen attenuates experimental allergic rhinitis | J. Mol. Medicine | 85 279-288, 2007 |
| Okada, M., Kita, Y., Nakajima, T., Kanamaru, N., Hashimoto, S., Nagasawa, T., <u>Kaneda, Y.</u> , et al | Evaluation of a novel vaccine (HVJ-liposome/HSP65 DNA+IL-12 DNA) against tuberculosis using the cynomolgus monkey model of TB | Vaccine | 25(16):2990-3,2007 |
| Hayashi, M., Nimura, K., Kashiwagi, K., Harada, T., Takaoka, K., Kato, H., <u>Tamai, K.</u> , and <u>Kaneda, Y</u> | Comparative Roles of Twist-1 and Id-1 in Transcriptional Regulation by BMP Signaling | J. Cell Sci | 120, 1350-1357, 2007 |
| Umegaki, N., <u>Tamai, K.</u> , Nakano, H., Moritsugu, R., Yamazaki, T., Hanada, H., Katayama, I., and <u>Kaneda, Y</u> | Differential regulation of karyopherin alpha2 expression by TGF-b1 and IFN-gamma in normal human keratinocytes: evident contribution of KPNA2 for nuclear translocation of IRF-1 | J. Invest. Dermatol | 127, 1456-1464, 2007 |
| Otsuru, S., <u>Tamai, K.</u> , Yamazaki, T., Yoshikawa, H., and <u>Kaneda, Y</u> | Bone marrow-derived osteoblast progenitor cells in circulating blood contribute to ectopic bone formation in mice | Biochem. Biophys. Res. Comm | 354, 453-458, 2007 |

| | | | |
|---|--|---------------------------------------|-------------------|
| Kurooka, M. and <u>Kaneda, Y</u> | Inactivated Sendai virus particles eradicate tumors by inducing immune responses through blocking regulatory T cells | Cancer Research | 67, 227–236, 2007 |
| Kikuchi Y, <u>Tamai K, Kaneda Y</u> | Cutaneous gene delivery | J Dermatol Sci | 2007 |
| Suwanasuthi, S., <u>Tamai, K. and Kaneda, Y</u> | Rapid transport of plasmid DNA into the nucleolus via actin depolymerization using HVJ envelope vector | J. Gene Medicine | 9, 55-62, 2007 |
| Jayawardana ST, Ushio H, Niyonsaba F, Gondokaryono SP, Takenaka H, <u>Ikeda S, Okumura K, Ogawa H</u> | Monomeric IgE and lipopolysaccharide synergistically prevent mast-cell apoptosis | Biochem Biophys Res Commun | 2007 Oct 30 |
| Hasegawa T, Suga Y, Mizoguchi M, Muramatsu S, Mizuno Y, Haruna K, <u>Ikeda S, Kuroyanagi Y, Ogawa H</u> | Intractable venous leg ulcer treated successfully with allogeneic cultured dermal substitute | Scand J Plast Reconstr Surg Hand Surg | 1-3,2007 |
| Zheng Y, Niyonsaba F, Ushio H, Nagaoka I, <u>Ikeda S, Okumura K, Ogawa H</u> | Cathelicidin LL-37 induces the generation of reactive oxygen species and release of human alpha-defensins from neutrophils | Br J Dermatol | 2007 Oct 4 |
| Ng W, Nishiyama C, Mizoguchi M, Nakano N, Suga Y, <u>Ikeda S, Itoh S, Kinoshita K, Okumura K, Ogawa H</u> | Human umbilical cord epithelial cells express Notch1: Implications for its epidermal-like differentiation | J Dermatol Sci | 2007 Sep 10 |
| Okuma K, Mitsuishi K, Hasegawa T, Tsuchihashi H, Ogawa H, <u>Ikeda S</u> | A Case Report of Steroid and Immunosuppressant-resistant Pyoderma Gangrenosum Successfully Treated by Granulocytapheresis | Ther Apher Dial | 11(5):387-90,2007 |
| Hasegawa T, Matsukura T, Suga Y, Muramatsu S, Mizuno Y, Tsuchihashi H, Haruna K, Ogawa H, <u>Ikeda S</u> | Case of acne conglobata successfully treated by CO(2) laser combined with topical tretinoin therapy | J Dermatol | 34(8):583-5,2007 |
| Jiang J, Kuhara T, Ueki R, Zheng Y, Suto H, <u>Ikeda S, Ogawa H</u> | Inhibitory effects of paroxetine on the development of atopic dermatitis-like lesions in NC/Nga mice | J Dermatol Sci | 47(3):244-7,2007 |
| Takagi A, Nishiyama C, Maeda K, Tokura T, Kawada H, Kanada S, Niwa Y, Nakano N, Mayuzumi N, Nishiyama M, <u>Ikeda S, Okumura K, Ogawa H</u> | Role of Sp1 in Transcription of Human ATP2A2 Gene in Keratinocytes | J Invest Dermatol | 2007 Jun 28 |
| Hasegawa T, Suga Y, Ikejima A, Muramatsu S, Mizuno Y, Tsuchihashi H, Haruna K, Ogawa H, <u>Ikeda S</u> | Suction blister grafting with CO(2) laser resurfacing of the graft recipient site for vitiligo | J Dermatol | 34(7):490-2,2007 |

| | | | |
|--|--|--------------------------|------------------------|
| Gondokaryono SP, Ushio H, Niyonsaba F, Hara M, Takenaka H, Jayawardana ST, <u>Ikeda S</u> , Okumura K, Ogawa H | The extra domain A of fibronectin stimulates murine mast cells via toll-like receptor 4 | J Leukoc Biol | 82(3):657-65,2007 |
| Hasegawa T, Sumiyoshi K, Tsuchihashi H, <u>Ikeda S</u> , Nakao A, Ogawa H | FK506 inhibits the enhancing effects of TGF-beta on wound healing in a rabbit dermal ulcer model | J Dermatol Sci | 47(1):37-40,2007 |
| Hasegawa T, Mizoguchi M, Haruna K, Mizuno Y, Muramatsu S, Suga Y, Ogawa H, <u>Ikeda S</u> | Amnia for intractable skin ulcers with recessive dystrophic epidermolysis bullosa: report of three cases | J Dermatol | 34(5):328-32,2007 |
| Ng W, Kikuchi Y, Chen X, Hira K, Ogawa H, <u>Ikeda S</u> | Pseudocysts of the auricle in a young adult with facial and ear atopic dermatitis | J Am Acad Dermatol | 56(5):858-61,2007 |
| Haruna K, Suga Y, Mizuno Y, Kourou K, Muramatsu S, Hasegawa T, Inoue A, Ogawa H, <u>Ikeda S</u> , Lakhmiri M, Seto A | A Moroccan patient with Sjogren-Larsson syndrome | J Dermatol | 34(2):153-5,2007 |
| Chen X, Niyonsaba F, Ushio H, Hara M, Yokoi H, Matsumoto K, Saito H, Nagaoka I, <u>Ikeda S</u> , Okumura K, Ogawa H | Antimicrobial peptides human beta-defensin (hBD)-3 and hBD-4 activate mast cells and increase skin vascular permeability | Eur J Immunol | 37(2):434-44,2007 |
| Tsuchihashi H, Hasegawa T, Sumiyoshi K, Ogawa H, <u>Ikeda S</u> , Ando T, Nakao A, Okumura K | TWEAK inhibits TGF-beta-induced contraction of normal and keloid fibroblast-embedded collagen gel | J Dermatol Sci | 45(3):216-8,2007 |
| Haruna K, Suga Y, Mizuno Y, Hasegawa T, Kourou K, Matsuba S, Muramatsu S, <u>Ikeda S</u> | R156C mutation of keratin 10 causes mild form of epidermolytic hyperkeratosis | J Dermatol | 34(8):545-8,2007 |
| Kohroh K, Suga Y, Mizuno Y, Ishii N, Hashimoto T, <u>Ikeda S</u> | Case of localized bullous pemphigoid with unique clinical manifestations in the lower legs | J Dermatol | 34(7):482-5,2007 |
| <u>山本明美</u> | Cornified cell envelope | 日本皮膚科学会雑誌 | 117 : 137-141,2007 |
| <u>山本明美</u> | 免疫組織染色・蛍光抗体法所見の読み方 | Visual Dermatology | 6(8), 834-839, 2007 |
| Ishida-Yamamoto A, Kishibe M, Takahashi H, Iizuka H | Rab11 is associated with epidermal lamellar granules | J Invest Dermatol | 127(9):2166-70. 2007 |
| Magnúsdóttir E, Kalachikov S, Mizukoshi K, Savitsky D, <u>Ishida-Yamamoto A</u> | Panteleyev AA, Calame K. Epidermal terminal differentiation depends on B lymphocyte-induced maturation protein-1 | Proc Natl Acad Sci U S A | 104(38):14988-93. 2007 |

| | | | |
|--|--|----------------------|-----------------------|
| Zeeuwen PL, <u>Ishida-Yamamoto A</u> , van Vlijmen-Willems IM, Cheng T, Bergers M, Iizuka H, Schalkwijk J | Colocalization of cystatin M/E and cathepsin V in lamellar granules and corneodesmosomes suggests a functional role in epidermal differentiation | J Invest Dermatol | 127(1):120-8. 2007 |
| Basel-Vanagaite L, Attia R, <u>Ishida-Yamamoto A</u> , Rainshtein L, Ben Amitai D, Lurie R, Pasmanik-Chor M, Indelman M, Zvulunov A, Saban S, Magal N, Sprecher E, Shohat M | Autosomal recessive ichthyosis with hypotrichosis caused by a mutation in ST14, encoding type II transmembrane serine protease matriptase | Am J Hum Genet | 80(3):467-77. 2007 |
| Kishibe M, Bando Y, Terayama R, Namikawa K, Takahashi H, Hashimoto Y, <u>Ishida-Yamamoto A</u> , Jiang YP, Mitrovic B, Perez D, Iizuka H, Yoshida S | Kallikrein 8 is involved in skin desquamation in cooperation with other kallikreins | J Biol Chem | 282(8):5834-41. 2007 |
| Lin Y, Kikuchi S, Tamakoshi A, Yagyu K, Obata Y, Inaba Y, <u>Kurosawa M</u> , et al | Obesity, physical activity and the risk of pancreatic cancer in a large Japanese cohort | Int J Cancer | 120(12):2665-71,2007 |
| Uchiyama T, <u>Kurosawa M</u> , Inaba Y | MMR-Vaccine and Regression in Autism Spectrum Disorders: Negative Results Presented from Japan | Autism Dev Disord | 37(2):210-7,2007 |
| Fukumoto T, Shiroyama Y, <u>Niizeki H</u> , et al | Paraneoplastic pemphigus presenting as erythrodermic lichenoid dermatitis with concomitant features of pemphigus foliaceus | J Dermatol | 34: 645-9, 2007 |
| 宮川幸子、 <u>新関寛徳</u> 、榎本美生、吉田さやか | 特集「小児の膠原病」 新生児エリテマトーデス | 皮膚病診療 | 29:538-542,2007 |
| 北村華奈、 <u>新関寛徳</u> 、古林郁乃、福本隆也、浅田秀夫、宮川幸子、村上伸介、玉置伸二、木村弘 | 間質性肺炎を伴ったAmyopathic Dermatomyositisの1例 | 皮膚の科学 | 6(3): 255-260, 2007 |
| <u>新関寛徳</u> | 水疱症の診断と治療 自己免疫性水疱症 2) 炎症の強い天疱瘡: 疱疹状天疱瘡、増殖性天疱瘡、腫瘍随伴性天疱瘡の診断と治療 | Monthly Derma | in press |
| Saito-Katsuragi M, Asada H, <u>Niizeki H</u> , Katoh F, Masuzawa M, Tsutsumi M, Kuniyasu H, Ito A, Nojima H, Miyagawa S | Role for connexin 26 in metastasis of human malignant melanoma: communication between melanoma and endothelial cells via connexin 26 | Cancer | 110(5): 1162-72, 2007 |
| 馬淵智生, 梅澤慶紀, <u>小澤 明</u> | 乾癬 | 皮膚病診療 | 29: 801-804, 2007 |
| T. Mabuchi, A. Oka, M. Iizuka, Y. Umezawa, T. Matsuyama, Y. Kubota, J. Nakayama, M. Ozawa, T. Terui, S. Yasumoto, T. Hashimoto, S. Ikeda, Y. Matsumoto, H. Sueki, M. Iijima, J.K. Kulski and <u>A. Ozawa</u> | Fine mapping of psoriasis-susceptibility locus within the HLA class II region by using microsatellite markers in an association study of Japanese cases and controls | Tokai J Exp Clin Med | 32: 6-13, 2007 |

| | | | |
|---|---|----------------------------------|----------------------|
| Okuyama R, Ogawa E, Nagoshi H, Yabuki M, Kurihara A, Terui T, Aiba S, Obinata M, Tagami H, Ikawa S | p53 homologue, p51/p63, maintains the immaturity of keratinocyte stem cells by inhibiting Notch1 activity | Oncogene | 2007 Jan 22 |
| Kusakari Y, Okuyama R, Hashimoto A, Ichinohasama R, Terui T, Tagami H, Aiba S | Recurrent classic Kaposi's sarcoma in a Japanese man: detection of human herpesvirus 8 infection by PCR and immunostaining | J Eur Acad Dermatol Venereol | 21(1):112-3,2007 |
| Nakajima K, Tamai K, Yamazaki T, Toyomaki Y, Nakano H, Uitto J, Sawamura D | Identification of Skn-1n, a Splice Variant Induced by High Calcium Concentration and Specifically Expressed in Normal Human Keratinocytes | J Invest Dermatol | in press |
| Yamamoto C, Tamai K, Nakano H, Matsuzaki Y, Kaneko T, Sawamura D | Vitamin D(3) inhibits expression of bullous pemphigoid antigen 1 through post-transcriptional mechanism without new protein synthesis | J Dermatol Sci | 2008 |
| 玉井克人 | 骨髓由来末梢循環細胞による組織再生 | 実験医学 | 2007 |
| Nakano H, Toyomaki Y, Ohashi S, Nakano A, Jin H, Munakata T, Akita N, Tamai K, Mitsuhashi Y | Novel COL7A1 mutations in a Japanese family with transient bullous dermolysis of the newborn associated with pseudosyndactyly | Br J Dermatol | 157(1):179-82,2007 |
| Mabuchi E, Umegaki N, Murota H, Nakamura T, Tamai K and Katayama I | Oral steroid improves bullous pemphigoid-like clinical manifestations in non-Herlitz junctional epidermolysis bullosa with COL17A1 mutation | Br J Dermatol | 157(3):596-8,2007 |
| 米田耕造、窪田泰夫 | 表皮細胞：疾患とユビキチン化 | 日本皮膚科学会雑誌 | in press |
| Yoneda K, Demitsu T, Matsuda Y and Kubota Y | Possible molecular mechanisms for sebaceous hyperplasia overlying dermatofibroma | Br J Dermatol | in press |
| Demitsu T, Yoneda K, Umemoto N, Azuma R, Kakurai M, Nishida J, Sadahira C and Kubota Y | Agminated mollusca contagiosa with attenuated ubiquitination of molluscum bodies as an initial sign of malignant lymphoma | J Eur Acad Dermatol and Venereol | 21: 691- 692, 2007 |
| Matsuoka Y, Yoneda K, Katsuura J, Moriue T, Nakai K, Sadahira C, Yokoi I, Nibu N, Demitsu T and Kubota Y | Successful treatment of follicular cutaneous T-cell lymphoma without mucinosis treated with narrow-band UVB irradiation | J Eur Acad Dermatol and Venereol | 21: 1121-1122, 2007 |
| Yoneda K, Bandoh S, Kanaji N, Moriue T, Katsuura J, Matsuoka Y, Nakai K, Demitsu T, Ishida T and Kubota Y | Eosinophilic pustular folliculitis associated with pulmonary infiltration with eosinophilia | J Eur Acad Dermatol and Venereol | 21: 1122- 1124, 2007 |
| Moriue T, Yoneda K, Katsuura J, Matsuoka Y, Nakai K, Yokoi I, Matsuda Y, Nibu N, Miyamoto I, Demitsu T and Kubota Y | Planar xanthomas due to cholestasis in graft versus host disease | Br J Dermatol | 156: 1374-1376, 2007 |

| | | | |
|---|---|----------------------------------|-------------------|
| <u>Yoneda K</u> , Katsuura J, Matsuoka Y, Moriawaki S, Moriue T, Nakai K, Yokoi I, Nibu N, Demitsu T and Kubota Y | Xeroderma pigmentosum group G in association with malignant melanoma | Eur J Dermatol | 17: 540-541, 2007 |
| Demitsu T, Kakurai M, Iida E, Takada M, Hiratsuka Y, Azuma R and <u>Yoneda K</u> | A case of nodular cystic fat necrosis with systemic lupus erythematosus presenting the multiple subcutaneous nodules on the extremities | J Eur Acad Dermatol and Venereol | in press |
| Demitsu T, Kakurai M, Azuma R, Haratsuka Y, Umemoto N, Ozaki K and <u>Yoneda K</u> | Long-lasting, urticaria-like erythema associated with IgA monoclonal gammopathy of undetermined significance | J Dermatol | in press |
| Nakai K, <u>Yoneda K</u> , Moriue T, Matsuoka Y, Yokoi I, Nibu N and Kubota Y | Fabry disease masquerading as multiple leg ulcers | J Eur Acad Dermatol and Venereol | 22:382-383,2008 |
| Demitsu T, <u>Yoneda K</u> , Iida E, Takada M, Azuma R, Umemoto N, Hiratsuka Y, Yamada T and Kakurai M | Urticarial vasculitis with hemorrhagic vesicles successfully treated with reserpine: report of a case. | J Eur Acad Dermatol and Venereol | in press |
| <u>Yoneda K</u> , Demitsu T, Matsuda Y and Kubota Y | Possible molecular mechanisms for sebaceous hyperplasia overlying dermatofibroma | Br J Dermatol | in press |
| Kubota Y, <u>Yoneda K</u> , Nakai K, Moriue T, Moriue J and Yokoi I | DRESS in a patient taking a dietary supplement containing phenobarbital purchased by internet shopping | Eur J Dermatol | in press |
| Moriue T, <u>Yoneda K</u> , Moriue J, Matsuoka Y, Nakai K, Yokoi I, Matsuda Y, Nibu N, Miyamoto I and Kubota Y | Simple therapeutic strategy for ingrown toenail with super elastic wire | Dermatol Surg | in press |
| Demitsu T, Kakurai M, Azuma R, Hiratsuka Y, Yamada T and <u>Yoneda K</u> | Recalcitrant pemphigus foliaceus with Kaposi's varicelliform eruption: report of a fatal case | Clin Exp Dermatol | in press |
| <u>Yoneda K</u> , Demitsu T, Matsuoka Y, Moriue T, Nakai K, Kushida Y, Haba R and Kubota Y | Subcellular activation site of caspase 3 in apoptotic keratinocytes observed in lichenoid tissue reaction | Br J Dermatol | in press |
| Kubota Y, <u>Yoneda K</u> , Nakai K, Moriue T, Matsuoka Y, Yokoi I and Ohya Y | Open-label pilot study of topical sequential applications with corticosteroids and tacrolimus of pediatric atopic dermatitis | J Am Acad Dermatol | in press |
| Nakai K, <u>Yoneda K</u> , Moriue T, Kosaka H and Kubota Y | Angiotensin II enhanced cell proliferation and EGF receptor expression levels via ROS formation in HaCaT cells | J Dermatol Sci | in press |
| Moriue T, Igarashi J, <u>Yoneda K</u> , Nakai K, Kosaka H and Kubota Y | Sphingosine-1-phosphate attenuates H ₂ O ₂ -induced apoptosis in cultured vascular endothelial cells: roles of caspase-3 and p38 MAP kinase | Biochem Biophys Res Commun | 368:852-857,2008 |

| | | | |
|--|---|----------------|----------------------|
| 米田耕造、窪田泰夫 | 角化症をどのように診るか | Clinical Derma | 8: 3-6, 2007 |
| 米田耕造、窪田泰夫 | 表皮細胞:疾患とユビキチン化 | 日本皮膚科学会雑誌 | 117: 2422-2424, 2007 |
| 中井浩三、米田耕造、横井郁美、丹生名都子、宮本 泉、松田保史、松岡由恵、森上徹也、串田吉生、羽場礼次、窪田泰夫 | アトピー性皮膚炎に合併したアミロイド苔癬の1例 | 皮膚科の臨床 | 49:1528-1529, 2007 |
| 宮本 泉、米田耕造、松田保史、丹生名都子、松岡由恵、横井郁美、中井浩三、森上徹也、窪田泰夫 | インフリキシマブ投与により改善したアトピー性皮膚炎の1例 | 皮膚科の臨床 | 49:984-985, 2007 |
| 横井郁美、米田耕造、窪田泰夫 | Zosteriform lichen planusの1例 | 皮膚科の臨床 | 49:986-987, 2007 |
| 丹生名都子、横井郁美、森上徹也、森上純子、松岡由恵、中井浩三、定平知江子、宮本 泉、松田保史、米田耕造、窪田泰夫 | 前治療抵抗性アトピー性皮膚炎に対するTh2サイトカイン阻害薬の有用性の検討-QOL評価を加えて- | 皮膚の科学 | 6: 33-37, 2007 |
| 窪田泰夫、森上徹也、中井浩三、松岡由恵、横井郁美、丹生名都子、宮本 泉、松田保史、米田耕造 | 患者主体の評価を基にしたベシル酸ベボタスチン(タリオン錠)の慢性蕁麻疹患者に対する臨床的有用性と患者QOLに及ぼす影響-慢性蕁麻疹の症状はどのように改善されるか- | 西日本皮膚科 | 69: 554-560, 2007 |
| 窪田泰夫、森上徹也、森上純子、松岡由恵、中井浩三、横井郁美、丹生名都子、宮本 泉、松田保史、米田耕造 | そう痒性皮膚疾患に対するロラタジン口腔内速溶錠(クラリチンレディタブ錠)の服用性及び有用性の検討-患者主体の評価を中心に- | 西日本皮膚科 | 69: 653-659, 2007 |
| 松田保史、松岡由恵、宮本 泉、米田耕造、窪田泰夫 | 脂腺の過形成を認めた皮膚線維腫の1例 | 皮膚科の臨床 | in press |
| 窪田泰夫、横井郁美、米田耕造 | 糖尿病合併皮膚潰瘍に対するLipo PGE1 製剤の有用性と患者QOLに及ぼす影響 | 診療と新薬 | in press |
| 奥田容子 新田悠紀子 小池文美香 大野俊之 黒木のぞみ 倉橋直子 青山裕美 | 線状IgA IgG水疱症の男児例 | 臨床皮膚科 | in press |

2. 単行本

| 著者名 | 論文題目 | 書籍名 | 頁、西暦年号 |
|----------------------|--|---|--------------|
| Kitajima Y, Aoyama Y | Desmosome assembly, disassembly and signaling in keratinocytes : Lessons from pemphigus, a desmosome-remodeling disease | Pathophysiology of Desmosome (RESEACH SIGNPOST) | in press |
| 北島康雄 (分担) | 1. 水疱症2. 非感染性膿皮症 | 図説臨床看護医学 (同朋舎メディアプラン) | in press |
| 北島康雄 (分担) | 角化異常症・汗孔角化症・急性痘瘡状苔癬状秕糠疹・黒色表皮腫・ジベルバラ色・秕糠疹・尖圭紅色苔癬・ダリエー病・頭部秕糠疹・連環状秕糠疹 (遠山)・播種性・汗孔角化症・反転性毛包角化症・砒素角化症・ヘブラ紅色秕糠疹・扁平苔癬・滴状・類乾癬毛孔性紅色秕糠疹・類乾癬 (17項目) | 医学大辞典第2版 (医学書院) | in press |
| 市來善郎、北島康雄 (分担) | 皮膚のしくみ | 皮膚の事典 (朝倉書店) | in press |
| 橋本 隆 | 自己免疫性皮膚疾患 | 目で見るアレルギー性皮膚疾患 (南山堂) | 250-261、2007 |
| 橋本 隆 | 尋常性天疱瘡、増殖性天疱瘡 | 目で見るアレルギー性皮膚疾患 (南山堂) | 262-266、2007 |
| 橋本 隆 | CD20抗体による自己免疫性水疱瘡の治療 | 目で見るアレルギー性皮膚疾患 (南山堂) | 436-438、2007 |
| 橋本 隆 | 天疱瘡 | 看護のための最新医学講座[第2版] 19皮膚疾患 (中山書店) | 96-102、2007 |
| 岩月啓氏、山本剛伸、辻和英 | EBウイルスとT/NK細胞増殖症 | 新しい診断と治療のABC/血液5 潜伏感染ウイルスによる血液疾患 (最新医学社) | 75-81、2007 |
| 岩月啓氏、辻 和英 | シックハウス症候群 | 目で見るアレルギー性皮膚疾患 (南山堂) | 354-364、2007 |
| 小宮根真弓 | ここまでわかった乾癬の病態 | WHAT' S NEW IN 皮膚科学2008-2009(メディカルレビュー社) | 2007 |
| 小宮根真弓 | 女性と乾癬 | P-PAT Letter 第12号 | 2008 |
| 小宮根真弓 | ジベルバラ色ひこう疹 | 看護のための最新医学講座第二版第19巻皮膚科疾患(中山書店) | 125-127,2007 |
| 小宮根真弓 | 毛孔性紅色ひこう疹 | 看護のための最新医学講座第二版第20巻皮膚科疾患(中山書店) | 123-125,2007 |
| 山本明美 | 毛孔性苔癬、顔面毛包性紅斑黒皮症 | 皮膚疾患最新の治療2007-2008 (南江堂) | 138、2007 |
| 新関寛徳、宮川幸子 | 皮膚筋炎 | 皮膚科診療カラーアトラス、第8巻 内臓疾患と皮膚 (講談社) | in press |

| | | | |
|-----------|-------------|---------------------------------------|---------------|
| 新関寛徳 | 乾癬性紅皮症 | 皮膚科診療カラーアトラス、第2巻 角 化異常症（非遺伝性）（講談社） | in press |
| 米田耕造、窪田泰夫 | ケラチン病-最近の進歩 | WHAT' S NEW in 皮膚科学(メジカル ビュー社) | 30-31, 2007 |
| 米田耕造、窪田泰夫 | 乾皮症 | スキンケア最前線(メジカルビュー社) | in press |
| 青山裕美（分担） | 落葉状天疱瘡 | 目で見えるアレルギー性皮膚疾患（南山 堂） | 267-275, 2007 |

厚生労働科学研究費補助金難治性疾患克服研究事業

稀少難治性皮膚疾患に関する調査研究班


平成19年度第1回総会プログラム

主任研究者 北島康雄

日時：平成19年8月3日（金）10時00分～17時00分

場所：KKRホテル東京11階 朱鷺の間

（東京都千代田区大手町1-4-1 ☎03-3287-2932）



連絡先：岐阜大学大学院医学研究科 病態制御学講座 皮膚病態学

☎501-1194 岐阜市柳戸1-1 ☎058-230-6394 FAX 058-230-6396

稀少難治性皮膚疾患に関する調査研究班

主任研究者 北島康雄

事務担当

青山裕美 ymaoyama@gifu-u.ac.jp

8. 天疱瘡の病変部には IL-17 産生リンパ球が存在する
大日輝記、荒川正崇、橋本隆 (久留米大)
9. 天疱瘡における HLA-DRB1 遺伝子プロフィールの検討 (腫瘍随伴性、病型移行例、尋常性天疱瘡関連 HLA-DRB1 アレル陰性例について)
新関寛徳、浅田秀夫、宮川幸子 (奈良医大)、青山裕美、北島康雄 (岐阜大)、清水宏 (北海道大)、橋本隆(久留米大)、池田志孝 (順天堂大)、高橋勇人、天谷雅行 (慶應大)、桑名正隆 (同 内科)、白方裕司、橋本公二 (愛媛大)、山本哲也、岩月啓氏 (岡山大)、岡晃、成瀬妙子、猪子英俊 (東海大分子生命科学)
10. 天疱瘡の H15 年度臨床調査個人票データを用いた最近 1 年間の治療状況
黒沢美智子(順天堂大・医・衛生)、池田志孝(順天堂大・医・皮膚)、天谷雅行(慶應大・皮膚科)、北島康雄(岐阜大・医・皮膚)、永井正規(埼玉医大・公衛)

—連絡会議 昼食—

12:20~13:00

先天性魚鱗癬様紅皮症

座長 池田志孝

13:00~14:00

- 1 1. 水疱型先天性魚鱗癬様紅皮症における過角化機構の解析。Rab11 と層板顆粒の関係
山本明美 (旭川医大 皮膚科)
- 1 2. 魚鱗癬の病態解明に関する研究
米田耕造¹、成清富貴子¹、出光俊郎²、窪田泰夫¹ (香川大医学部皮膚科¹、自治医科大学大宮医療センター皮膚科²)
- 1 3. 葉状魚鱗癬表皮におけるS-S結合の分布について
水野優起、須賀康、池田志孝 (順天堂大 皮膚科)

- 1 4. 水疱型先天性魚鱗癬様紅皮症のケラチン K1 変異による表皮細胞への影響
小宮根真弓 (東京大学)

膿疱性乾癬

座長 岩月啓氏

14:00~14:50

- 1 5. ゲノムワイドな遺伝的相関解析による乾癬感受性遺伝子の同定
馬淵智生, 梅澤慶紀, 小澤 明 (東海大), 岡 晃, 猪子英俊 (同分生)
- 1 6. 膿疱性乾癬の病態解明とその対策に向けて
—S100A11 によるヒト表皮角化細胞の増殖制御機構
許 南浩 (岡山大学医歯薬学総合研究科細胞生物学分野)
- 1 7. 膿疱性乾癬の新規バイオマーカーとしての血清 S 1 0 0 A 8 / A 9 の検討 : 重症度・合併症との関連について
中西元¹、辻和英¹、青地聖子¹、許南浩²、津田達也³、山西清文³、小宮根真弓⁴、岩月啓氏¹ (¹岡山大学大学院医歯薬学総合研究科 皮膚・粘膜・結合織学、²岡山大学大学院医歯薬学総合研究科 細胞生物学、³兵庫医科大学皮膚科、⁴東京大学皮膚科)
- 1 8. 膿疱性乾癬の病因と治療—表皮角化細胞の増殖・分化と脂質代謝
照井 正、横山 愛 (日本大)、青山裕美 (岐阜大)、槇島 眞 (日本大生化学)

—休憩— 10分

治療指針作成に関する発表

座長 北島康雄

15:00~16:50

北島康雄：治療指針作成に関する提案
天谷雅行：天疱瘡
橋本隆：類天疱瘡
岩月啓氏：膿疱性乾癬
清水宏：表皮水疱症
池田志孝：水疱型先天性魚鱗癬様紅皮症

閉会の挨拶

主任研究者 北島康雄

[V]

平成19年度総会プログラム

厚生労働科学研究費補助金難治性疾患克服研究事業


稀少難治性皮膚疾患に関する調査研究班

平成19年度第2回総会プログラム

主任研究者 北島康雄

日時：平成20年1月11日(金) 午前10時00分～午後17時00分

場所：KKR ホテル東京11階 丹頂の間
(東京都千代田区大手町1-4-1 ☎03-3287-2932)



連絡先：岐阜大学大学院医学研究科 病態制御学講座 皮膚病態学

☎501-1194 岐阜市柳戸1-1 ☎ 058-230-6394 FAX 058-230-6396

稀少難治性皮膚疾患に関する調査研究班

主任研究者 北島康雄

事務担当 青山裕美 ymaoyama@gmail.com cc.gifu-u.ac.jp