

20. Kaplan G, Racaniello VR. Construction and characterization of poliovirus subgenomic replicons. *J Virol* 1988;62:1687–96.
21. Liljestrom P, Garoff H. A new generation of animal cell expression vectors based on the Semliki Forest virus replicon. *Biotechnology (NY)* 1991;9:1356–61.
22. Bredenbeek PJ, Frolov I, Rice CM, Schlesinger S. Sindbis virus expression vectors: packaging of RNA replicons by using defective helper RNAs. *J Virol* 1993;67:6439–46.
23. Johanning FW, Conry RM, LoBuglio AF, Wright M, Sumerel LA, Pike MJ, et al. A Sindbis virus mRNA polynucleotide vector achieves prolonged and high level heterologous gene expression in vivo. *Nucleic Acids Res* 1995;23:1495–501.
24. Kamrud KI, Powers AM, Higgs S, Olson KE, Blair CD, Carlson JO, et al. The expression of chloramphenicol acetyltransferase in mosquitoes and mosquito cells using a packaged Sindbis replicon system. *Exp Parasitol* 1995;81:394–403.
25. Xiong C, Levis R, Shen P, Schlesinger S, Rice CM, Huang HV. Sindbis virus: an efficient, broad host range vector for gene expression in animal cells. *Science* 1989;243:1188–91.
26. Khromykh AA, Westaway EG. Subgenomic replicons of the flavivirus Kunjin: construction and applications. *J Virol* 1997;71:1497–505.
27. McKnight KL, Lemon SM. Capsid coding sequence is required for efficient replication of human rhinovirus 14 RNA. *J Virol* 1996;70:1941–52.
28. Behrens SE, Grassmann CW, Thiel HJ, Meyers G, Tautz N. Characterization of an autonomous subgenomic pestivirus RNA replicon. *J Virol* 1998;72:2364–72.
29. Blight KJ, Kolykhalov AA, Rice CM. Efficient initiation of HCV RNA replication in cell culture. *Science* 2000;290:1972–4.
30. Krieger N, Lohmann V, Bartenschlager R. Enhancement of hepatitis C virus RNA replication by cell culture-adaptive mutations. *J Virol* 2001;75:4614–24.
31. Lohmann V, Korner F, Dobierzewska A, Bartenschlager R. Mutations in hepatitis C virus RNAs conferring cell culture adaptation. *J Virol* 2001;75:1437–49.
32. Evans MJ, Rice CM, Goff SP. Phosphorylation of hepatitis C virus nonstructural protein 5A modulates its protein interactions and viral RNA replication. *Proc Natl Acad Sci USA* 2004;101:13038–43.
33. Appel N, Pietschmann T, Bartenschlager R. Mutational analysis of hepatitis C virus nonstructural protein 5A: potential role of differential phosphorylation in RNA replication and identification of a genetically flexible domain. *J Virol* 2005;79:3187–94.
34. Lohmann V, Hoffmann S, Herian U, Penin F, Bartenschlager R. Viral and cellular determinants of hepatitis C virus RNA replication in cell culture. *J Virol* 2003;77:3007–19.
35. Yi M, Lemon SM. Adaptive mutations producing efficient replication of genotype 1a hepatitis C virus RNA in normal Huh7 cells. *J Virol* 2004;78:7904–15.
36. Bartenschlager R. Hepatitis C virus molecular clones: from cDNA to infectious virus particles in cell culture. *Curr Opin Microbiol* 2006;9:416–22.
37. Brass V, Moradpour D, Blum HE. Molecular virology of hepatitis C virus (HCV): 2006 update. *Int J Med Sci* 2006;3:29–34.
38. Bartenschlager R. The hepatitis C virus replicon system: from basic research to clinical application. *J Hepatol* 2005;43:210–6.
39. Blight KJ, McKeating JA, Rice CM. Highly permissive cell lines for subgenomic and genomic hepatitis C virus RNA replication. *J Virol* 2002;76:13001–14.
40. Friebe P, Boudet J, Simorre JP, Bartenschlager R. Kissing-loop interaction in the 3' end of the hepatitis C virus genome essential for RNA replication. *J Virol* 2005;79:380–92.
41. Kato T, Furusaka A, Miyamoto M, Date T, Yasui K, Hiramoto J, et al. Sequence analysis of hepatitis C virus isolated from a fulminant hepatitis patient. *J Med Virol* 2001;64:334–9.
42. Kato T, Date T, Miyamoto M, Furusaka A, Tokushige K, Mizokami M, et al. Efficient replication of the genotype 2a hepatitis C virus subgenomic replicon. *Gastroenterology* 2003;125:1808–17.
43. Date T, Kato T, Miyamoto M, Zhao Z, Yasui K, Mizokami M, et al. Genotype 2a hepatitis C virus subgenomic replicon can replicate in HepG2 and IMY-N9 cells. *J Biol Chem* 2004;279:22371–6.
44. Kato T, Date T, Miyamoto M, Zhao Z, Mizokami M, Wakita T. Nonhepatic cell lines HeLa and 293 support efficient replication of the hepatitis C virus genotype 2a subgenomic replicon. *J Virol* 2005;79:592–6.
45. Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, et al. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med* 2005;11:791–6.
46. Zhong J, Gastaminza P, Cheng G, Kapadia S, Kato T, Burton DR, et al. Robust hepatitis C virus infection in vitro. *Proc Natl Acad Sci USA* 2005;102:9294–9.
47. Lindenbach BD, Evans MJ, Syder AJ, Wolk B, Tellinghuisen TL, Liu CC, et al. Complete replication of hepatitis C virus in cell culture. *Science* 2005;309:623–6.
48. Compans RW. Virus entry and release in polarized epithelial cells. *Curr Top Microbiol Immunol* 1995;202:209–19.
49. Garoff H, Hewson R, Opstelten DJ. Virus maturation by budding. *Microbiol Mol Biol Rev* 1998;62:1171–90.
50. Schmitt AP, Lamb RA. Escaping from the cell: assembly and budding of negative-strand RNA viruses. *Curr Top Microbiol Immunol* 2004;283:145–96.
51. Takimoto T, Portner A. Molecular mechanism of paramyxovirus budding. *Virus Res* 2004;106:133–45.
52. Murakami K, Ishii K, Ishihara Y, Yoshizaki S, Tanaka K, Gotoh Y, et al. Production of infectious hepatitis C virus particles in three-dimensional cultures of the cell line carrying the genome-length dicistronic viral RNA of genotype 1b. *Virology* 2006;351:381–92.
53. Kawada M, Nagamori S, Aizaki H, Fukaya K, Niiya M, Matsuura T, et al. Massive culture of human liver cancer cells in a newly developed radial flow bioreactor system: ultrafine structure of functionally enhanced hepatocarcinoma cell lines. *In Vitro Cell Dev Biol Anim* 1998;34:109–15.
54. Matsuura T, Kawada M, Hasumura S, Nagamori S, Obata T, Yamaguchi M, et al. High density culture of immortalized liver endothelial cells in the radial-flow bioreactor in the development of an artificial liver. *Int J Artif Organs* 1998;21:229–34.
55. Aizaki H, Nagamori S, Matsuda M, Kawakami H, Hashimoto O, Ishiko H, et al. Production and release of infectious hepatitis C virus from human liver cell cultures in the three-dimensional radial-flow bioreactor. *Virology* 2003;314:16–25.
56. Iwahori T, Matsuura T, Maehashi H, Sugo K, Saito M, Hosokawa M, et al. CYP3A4 inducible model for in vitro analysis of human drug metabolism using a bioartificial liver. *Hepatology* 2003;37:665–73.
57. Yoshioka H, Mikami M, Mori Y, Tsuchida E. A synthetic hydrogel with thermoreversible gelation. *J Macromol Sci* 1994; A31:113–20.
58. Bukh J, Purcell RH, Miller RH. Sequence analysis of the 5' noncoding region of hepatitis C virus. *Proc Natl Acad Sci USA* 1992;89:4942–6.
59. Brown EA, Zhang H, Ping LH, Lemon SM. Secondary structure of the 5' untranslated regions of hepatitis C virus and pestivirus genomic RNAs. *Nucleic Acids Res* 1992;20:5041–5.
60. Honda M, Beard MR, Ping LH, Lemon SM. A phylogenetically conserved stem-loop structure at the 5' border of the internal ribosome entry site of hepatitis C virus is required for cap-independent viral translation. *J Virol* 1999;73:1165–74.
61. Honda M, Brown EA, Lemon SM. Stability of a stem-loop involving the initiator AUG controls the efficiency of internal initiation of translation on hepatitis C virus RNA. *RNA* 1996;2:955–68.

62. Tsukiyama-Kohara K, Iizuka N, Kohara M, Nomoto A. Internal ribosome entry site within hepatitis C virus RNA. *J Virol* 1992; 66:1476–83.
63. Wang C, Sarnow P, Siddiqui A. Translation of human hepatitis C virus RNA in cultured cells is mediated by an internal ribosome-binding mechanism. *J Virol* 1993;67:3338–44.
64. Hellen CU, Pestova TV. Translation of hepatitis C virus RNA. *J Viral Hepat* 1999;6:79–87.
65. Lu HH, Wimmer E. Poliovirus chimeras replicating under the translational control of genetic elements of hepatitis C virus reveal unusual properties of the internal ribosomal entry site of hepatitis C virus. *Proc Natl Acad Sci USA* 1996;93:1412–7.
66. Reynolds JE, Kaminski A, Kettinen HJ, Grace K, Clarke BE, Carroll AR, et al. Unique features of internal initiation of hepatitis C virus RNA translation. *EMBO J* 1995;14:6010–20.
67. Rijnbrand R, Bredenbeek P, van der Straaten T, Whetter L, Inchauspe G, Lemon S, et al. Almost the entire 5' non-translated region of hepatitis C virus is required for cap-independent translation. *FEBS Lett* 1995;365:115–9.
68. Yoo BJ, Spaete RR, Geballe AP, Selby M, Houghton M, Han JH. 5' end-dependent translation initiation of hepatitis C viral RNA and the presence of putative positive and negative translational control elements within the 5' untranslated region. *Virology* 1992;191:889–99.
69. Lemon S, Honda M. Internal ribosome entry sites within the RNA genomes of hepatitis C virus and other flaviviruses. *Semin Virol* 1997;8:274–88.
70. Buratti E, Tisminetzky S, Zotti M, Baralle FE. Functional analysis of the interaction between HCV 5'UTR and putative subunits of eukaryotic translation initiation factor eIF3. *Nucleic Acids Res* 1998;26:3179–87.
71. Kieft JS, Zhou K, Jubin R, Doudna JA. Mechanism of ribosome recruitment by hepatitis C IRES RNA. *RNA* 2001;7:194–206.
72. Sizova DV, Kolupaeva VG, Pestova TV, Shatsky IN, Hellen CU. Specific interaction of eukaryotic translation initiation factor 3 with the 5' nontranslated regions of hepatitis C virus and classical swine fever virus RNAs. *J Virol* 1998;72:4775–82.
73. Ali N, Pruijn GJ, Kenan DJ, Keene JD, Siddiqui A. Human La antigen is required for the hepatitis C virus internal ribosome entry site-mediated translation. *J Biol Chem* 2000;275:27531–40.
74. Ali N, Siddiqui A. The La antigen binds 5' noncoding region of the hepatitis C virus RNA in the context of the initiator AUG codon and stimulates internal ribosome entry site-mediated translation. *Proc Natl Acad Sci USA* 1997;94:2249–54.
75. Isoyama T, Kamoshita N, Yasui K, Iwai A, Shiroki K, Toyoda H, et al. Lower concentration of La protein required for internal ribosome entry on hepatitis C virus RNA than on poliovirus RNA. *J Gen Virol* 1999;80:2319–27.
76. Hahm B, Kim YK, Kim JH, Kim TY, Jang SK. Heterogeneous nuclear ribonucleoprotein L interacts with the 3' border of the internal ribosomal entry site of hepatitis C virus. *J Virol* 1998; 72:8782–8.
77. Fukushi S, Okada M, Kageyama T, Hoshino FB, Nagai K, Katayama K. Interaction of poly(rC)-binding protein 2 with the 5'-terminal stem loop of the hepatitis C-virus genome. *Virus Res* 2001;73:67–79.
78. Anwar A, Ali N, Tanveer R, Siddiqui A. Demonstration of functional requirement of polypyrimidine tract-binding protein by SELEX RNA during hepatitis C virus internal ribosome entry site-mediated translation initiation. *J Biol Chem* 2000;275: 34231–5.
79. Shimoike T, Mimori S, Tani H, Matsuura Y, Miyamura T. Interaction of hepatitis C virus core protein with viral sense RNA and suppression of its translation. *J Virol* 1999;73:9718–25.
80. Tanaka Y, Shimoike T, Ishii K, Suzuki R, Suzuki T, Ushijima H, et al. Selective binding of hepatitis C virus core protein to synthetic oligonucleotides corresponding to the 5' untranslated region of the viral genome. *Virology* 2000;270:229–36.
81. Shimoike T, Koyama C, Murakami K, Suzuki R, Matsuura Y, Miyamura T, et al. Down-regulation of the internal ribosome entry site (IRES)-mediated translation of the hepatitis C virus: critical role of binding of the stem-loop IIIId domain of IRES and the viral core protein. *Virology* 2006;345:434–45.
82. Wang TH, Rijnbrand RC, Lemon SM. Core protein-coding sequence, but not core protein, modulates the efficiency of cap-independent translation directed by the internal ribosome entry site of hepatitis C virus. *J Virol* 2000;74:11347–58.
83. Zhang J, Yamada O, Yoshida H, Iwai T, Araki H. Autogenous translational inhibition of core protein: implication for switch from translation to RNA replication in hepatitis C virus. *Virology* 2002;293:141–50.
84. Li D, Takyar ST, Lott WB, Gowans EJ. Amino acids 1–20 of the hepatitis C virus (HCV) core protein specifically inhibit HCV IRES-dependent translation in HepG2 cells, and inhibit both HCV IRES- and cap-dependent translation in HuH7 and CV-1 cells. *J Gen Virol* 2003;84:815–25.
85. Friebe P, Lohmann V, Krieger N, Bartenschlager R. Sequences in the 5' nontranslated region of hepatitis C virus required for RNA replication. *J Virol* 2001;75:12047–57.
86. Hüsey P, Langen H, Mous J, Jacobsen H. Hepatitis C virus core protein: carboxy-terminal boundaries of two processed species suggest cleavage by a signal peptide peptidase. *Virology* 1996;224: 93–104.
87. McLauchlan J, Lemberg MK, Hope G, Martoglio B. Intramembrane proteolysis promotes trafficking of hepatitis C virus core protein to lipid droplets. *EMBO J* 2002;21:3980–8.
88. Lemberg MK, Martoglio B. Requirements for signal peptide peptidase-catalyzed intramembrane proteolysis. *Mol Cell* 2002; 10:735–44.
89. Okamoto K, Moriishi K, Miyamura T, Matsuura Y. Intramembrane proteolysis and endoplasmic reticulum retention of hepatitis C virus core protein. *J Virol* 2004;78:6370–80.
90. Weihofen A, Binns K, Lemberg MK, Ashman K, Martoglio B. Identification of signal peptide peptidase, a presenilin-type aspartic protease. *Science* 2002;296:2215–8.
91. Pallaoro M, Lahm A, Biasiol G, Brunetti M, Nardella C, Orsatti L, et al. Characterization of the hepatitis C virus NS2/3 processing reaction by using a purified precursor protein. *J Virol* 2001;75: 9939–46.
92. Thibeault D, Maurice R, Pilote L, Lamarre D, Pause A. In vitro characterization of a purified NS2/3 protease variant of hepatitis C virus. *J Biol Chem* 2001;276:46678–84.
93. Bartenschlager R, Ahlborn-Laake L, Mous J, Jacobsen H. Kinetic and structural analyses of hepatitis C virus polyprotein processing. *J Virol* 1994;68:5045–55.
94. Failla C, Tomei L, De Francesco R. An amino-terminal domain of the hepatitis C virus NS3 protease is essential for interaction with NS4A. *J Virol* 1995;69:1769–77.
95. Lin C, Pragai BM, Grakoui A, Xu J, Rice CM. Hepatitis C virus NS3 serine proteinase: *trans*-cleavage requirements and processing kinetics. *J Virol* 1994;68:8147–57.
96. Tanji Y, Hijikata M, Hirowatari Y, Shimotohno K. Hepatitis C virus polyprotein processing: kinetics and mutagenic analysis of serine proteinase-dependent cleavage. *J Virol* 1994;68:8418–22.
97. Jin L, Peterson DL. Expression, isolation, and characterization of the hepatitis C virus ATPase/RNA helicase. *Arch Biochem Biophys* 1995;323:47–53.
98. Kim DW, Gwack Y, Han JH, Choe J. C-terminal domain of the hepatitis C virus NS3 protein contains an RNA helicase activity. *Biochem Biophys Res Commun* 1995;215:160–6.
99. Suzich JA, Tamura JK, Palmer-Hill F, Warrenner P, Grakoui A, Rice CM, et al. Hepatitis C virus NS3 protein polynucleotide-stimulated nucleoside triphosphatase and comparison with the related pestivirus and flavivirus enzymes. *J Virol* 1993;67: 6152–8.

100. Tai CL, Chi WK, Chen DS, Hwang LH. The helicase activity associated with hepatitis C virus nonstructural protein 3 (NS3). *J Virol* 1996;70:8477–84.
101. Tai CL, Pan WC, Liaw SH, Yang UC, Hwang LH, Chen DS. Structure-based mutational analysis of the hepatitis C virus NS3 helicase. *J Virol* 2001;75:8289–97.
102. Wolk B, Sansonno D, Krausslich HG, Dammacco F, Rice CM, Blum HE, et al. Subcellular localization, stability, and trans-cleavage competence of the hepatitis C virus NS3–NS4A complex expressed in tetracycline-regulated cell lines. *J Virol* 2000;74:2293–304.
103. Barbato G, Cicero DO, Nardi MC, Steinkuhler C, Cortese R, De Francesco R, et al. The solution structure of the N-terminal proteinase domain of the hepatitis C virus (HCV) NS3 protein provides new insights into its activation and catalytic mechanism. *J Mol Biol* 1999;289:371–84.
104. Tanji Y, Hijikata M, Satoh S, Kaneko T, Shimotohno K. Hepatitis C virus-encoded nonstructural protein NS4A has versatile functions in viral protein processing. *J Virol* 1995;69:1575–81.
105. Restrepo-Hartwig MA, Ahlquist P. Brome mosaic virus helicase- and polymerase-like proteins colocalize on the endoplasmic reticulum at sites of viral RNA synthesis. *J Virol* 1996;70:8908–16.
106. Schaad MC, Jensen PE, Carrington JC. Formation of plant RNA virus replication complexes on membranes: role of an endoplasmic reticulum-targeted viral protein. *EMBO J* 1997;16:4049–59.
107. van der Meer Y, van Tol H, Locker JK, Snijder EJ. ORF1a-encoded replicase subunits are involved in the membrane association of the arterivirus replication complex. *J Virol* 1998;72:6689–98.
108. Shi ST, Schiller JJ, Kanjanahaluethai A, Baker SC, Oh JW, Lai MM. Colocalization and membrane association of murine hepatitis virus gene 1 products and de novo-synthesized viral RNA in infected cells. *J Virol* 1999;73:5957–69.
109. Froshauer S, Kartenbeck J, Helenius A. Alphavirus RNA replicase is located on the cytoplasmic surface of endosomes and lysosomes. *J Cell Biol* 1988;107:2075–86.
110. Egger D, Wolk B, Gosert R, Bianchi L, Blum HE, Moradpour D, et al. Expression of hepatitis C virus proteins induces distinct membrane alterations including a candidate viral replication complex. *J Virol* 2002;76:5974–84.
111. Gosert R, Egger D, Lohmann V, Bartenschlager R, Blum HE, Bienz K, et al. Identification of the hepatitis C virus RNA replication complex in Huh-7 cells harboring subgenomic replicons. *J Virol* 2003;77:5487–92.
112. Shi ST, Lee KJ, Aizaki H, Hwang SB, Lai MM. Hepatitis C virus RNA replication occurs on a detergent-resistant membrane that cofractionates with caveolin-2. *J Virol* 2003;77:4160–8.
113. Piccininni S, Varaklioti A, Nardelli M, Dave B, Raney KD, McCarthy JE. Modulation of the hepatitis C virus RNA-dependent RNA polymerase activity by the non-structural (NS) 3 helicase and the NS4B membrane protein. *J Biol Chem* 2002;277:45670–9.
114. Gao L, Aizaki H, He JW, Lai MM. Interactions between viral nonstructural proteins and host protein hVAP-33 mediate the formation of hepatitis C virus RNA replication complex on lipid raft. *J Virol* 2004;78:3480–8.
115. Brass V, Bieck E, Montserret R, Wolk B, Hellings JA, Blum HE, et al. An amino-terminal amphipathic alpha-helix mediates membrane association of the hepatitis C virus nonstructural protein 5A. *J Biol Chem* 2002;277:8130–9.
116. Shimakami T, Hijikata M, Luo H, Ma YY, Kaneko S, Shimotohno K, et al. Effect of interaction between hepatitis C virus NS5A and NS5B on hepatitis C virus RNA replication with the hepatitis C virus replicon. *J Virol* 2004;78:2738–48.
117. Hamamoto I, Nishimura Y, Okamoto T, Aizaki H, Liu M, Mori Y, et al. Human VAP-B is involved in hepatitis C virus replication through interaction with NS5A and NS5B. *J Virol* 2005;79:13473–82.
118. Ito T, Lai MM. An internal polypyrimidine-tract-binding protein-binding site in the hepatitis C virus RNA attenuates translation, which is relieved by the 3'-untranslated sequence. *Virology* 1999;254:288–96.
119. Kolykhalov AA, Agapov EV, Blight KJ, Mihalik K, Feinstone SM, Rice CM. Transmission of hepatitis C by intrahepatic inoculation with transcribed RNA. *Science* 1997;277:570–4.
120. Tanaka T, Kato N, Cho M-J, Shimotohno K. A novel sequence found at the 3' terminus of hepatitis C virus genome. *Biochem Biophys Res Commun* 1995;215:744–9.
121. Yi M, Lemon SM. 3' nontranslated RNA signals required for replication of hepatitis C virus RNA. *J Virol* 2003;77:3557–68.
122. Friebe P, Bartenschlager R. Genetic analysis of sequences in the 3' nontranslated region of hepatitis C virus that are important for RNA replication. *J Virol* 2002;76:5326–38.
123. Aizaki H, Lee KJ, Sung VM, Ishiko H, Lai MM. Characterization of the hepatitis C virus RNA replication complex associated with lipid rafts. *Virology* 2004;324:450–61.
124. Ali N, Tardif KD, Siddiqui A. Cell-free replication of the hepatitis C virus subgenomic replicon. *J Virol* 2002;76:12001–7.
125. Lai VC, Dempsey S, Lau JY, Hong Z, Zhong W. In vitro RNA replication directed by replicase complexes isolated from the subgenomic replicon cells of hepatitis C virus. *J Virol* 2003;77:2295–300.
126. Miyazaki Y, Hijikata M, Yamaji M, Hosaka M, Takahashi H, Shimotohno K. Hepatitis C virus non-structural proteins in the probable membranous compartment function in viral genome replication. *J Biol Chem* 2003;278:50301–8.
127. Hardy RW, Marcotrigiano J, Blight KJ, Majors JE, Rice CM. Hepatitis C virus RNA synthesis in a cell-free system isolated from replicon-containing hepatoma cells. *J Virol* 2003;77:2029–37.
128. Waris G, Sarker S, Siddiqui A. Two-step affinity purification of the hepatitis C virus ribonucleoprotein complex. *RNA* 2004;10:321–9.
129. Simons K, Ikonen E. Functional rafts in cell membranes. *Nature* 1997;387:569–72.
130. Simons K, Ikonen E. How cells handle cholesterol. *Science* 2000;290:1721–6.
131. Simons K, Toomre D. Lipid rafts and signal transduction. *Nat Rev Mol Cell Biol* 2000;1:31–9.
132. Barman S, Ali A, Hui EK, Adhikary L, Nayak DP. Transport of viral proteins to the apical membranes and interaction of matrix protein with glycoproteins in the assembly of influenza viruses. *Virus Res* 2001;77:61–9.
133. Scheiffele P, Rietveld A, Wilk T, Simons K. Influenza viruses select ordered lipid domains during budding from the plasma membrane. *J Biol Chem* 1999;274:2038–44.
134. Zhang J, Pekosz A, Lamb RA. Influenza virus assembly and lipid raft microdomains: a role for the cytoplasmic tails of the spike glycoproteins. *J Virol* 2000;74:4634–44.
135. Ding L, Derdowski A, Wang JJ, Spearman P. Independent segregation of human immunodeficiency virus type 1 Gag protein complexes and lipid rafts. *J Virol* 2003;77:1916–26.
136. Ono A, Freed EO. Plasma membrane rafts play a critical role in HIV-1 assembly and release. *Proc Natl Acad Sci USA* 2001;98:13925–30.
137. Bavari S, Bosio CM, Wiegand E, Ruthel G, Will AB, Geisbert TW, et al. Lipid raft microdomains: a gateway for compartmentalized trafficking of Ebola and Marburg viruses. *J Exp Med* 2002;195:593–602.
138. Stuart AD, Eustace HE, McKee TA, Brown TD. A novel cell entry pathway for a DAF-using human enterovirus is dependent on lipid rafts. *J Virol* 2002;76:9307–22.
139. Narayan S, Barnard RJ, Young JA. Two retroviral entry pathways distinguished by lipid raft association of the viral receptor and differences in viral infectivity. *J Virol* 2003;77:1977–83.

140. Ashbourne Excoffon KJ, Moninger T, Zabner J. The Coxsackie B virus and adenovirus receptor resides in a distinct membrane microdomain. *J Virol* 2003;77:2559–67.
141. Brown DA, Rose JK. Sorting of GPI-anchored proteins to glycolipid-enriched membrane subdomains during transport to the apical cell surface. *Cell* 1992;68:533–44.
142. Hijikata M, Mizushima H, Tanji Y, Komoda Y, Hirowatari Y, Akagi T, et al. Proteolytic processing and membrane association of putative nonstructural proteins of hepatitis C virus. *Proc Natl Acad Sci USA* 1993;90:10773–7.
143. Mottola G, Cardinali G, Ceccacci A, Trozzi C, Bartholomew L, Torrisi MR, et al. Hepatitis C virus nonstructural proteins are localized in a modified endoplasmic reticulum of cells expressing viral subgenomic replicons. *Virology* 2002;293:31–43.
144. Tu H, Gao L, Shi ST, Taylor DR, Yang T, Mircheff AK, et al. Hepatitis C virus RNA polymerase and NSSA complex with a SNARE-like protein. *Virology* 1999;263:30–41.
145. Pietschmann T, Lohmann V, Rutter G, Kurpanek K, Bartenschlager R. Characterization of cell lines carrying self-replicating hepatitis C virus RNAs. *J Virol* 2001;75:1252–64.
146. Choi J, Lee KJ, Zheng Y, Yamaga AK, Lai MM, Ou JH. Reactive oxygen species suppress hepatitis C virus RNA replication in human hepatoma cells. *Hepatology* 2004;39:81–9.
147. Kaito M, Watanabe S, Tsukiyama-Kohara K, Yamaguchi K, Kobayashi Y, Konishi M, et al. Hepatitis C virus particle detected by immunoelectron microscopic study. *J Gen Virol* 1994;75:1755–60.
148. Shimizu YK, Feinstone SM, Kohara M, Purcell RH, Yoshikura H. Hepatitis C virus: detection of intracellular virus particles by electron microscopy. *Hepatology* 1996;23:205–9.
149. Maillard P, Krawczynski K, Nitkiewicz J, Bronnert C, Sidorkiewicz M, Gounon P, et al. Nonenveloped nucleocapsids of hepatitis C virus in the serum of infected patients. *J Virol* 2001;75:8240–50.
150. Andre P, Komurian-Pradel F, Deforges S, Perret M, Berland JL, Sodoyer M, et al. Characterization of low- and very-low-density hepatitis C virus RNA-containing particles. *J Virol* 2002;76:6919–28.
151. Moradpour D, Wakita T, Tokushige K, Carlson RI, Krawczynski K, Wands JR. Characterization of three novel monoclonal antibodies against hepatitis C virus core protein. *J Med Virol* 1996;48:234–41.
152. Barba G, Harper F, Harada T, Kohara M, Goulinet S, Matsuura Y, et al. Hepatitis C virus core protein shows a cytoplasmic localization and associates to cellular lipid storage droplets. *Proc Natl Acad Sci USA* 1997;94:1200–5.
153. Moriya K, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, Ishibashi K, et al. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065–7.
154. Hope RG, Murphy DJ, McLauchlan J. The domains required to direct core proteins of hepatitis C virus and GB virus-B to lipid droplets share common features with plant oleosin proteins. *J Biol Chem* 2002;277:4261–70.
155. Suzuki R, Sakamoto S, Tsutsumi T, Rikimaru A, Tanaka K, Shimoike T, et al. Molecular determinants for subcellular localization of hepatitis C virus core protein. *J Virol* 2005;79:1271–81.
156. Suzuki T, Suzuki R. Maturation and assembly of hepatitis C virus core protein. In: Kalitzky M, Borowski P, editors. *Molecular biology of the Flavivirus*. Norfolk, UK: Horizon Bioscience; 2006. p. 295–311.
157. Shintani Y, Fujie H, Miyoshi H, Tsutsumi T, Tsukamoto K, Kimura S, et al. Hepatitis C virus infection and diabetes: direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;126:840–8.
158. Moriishi K, Mochizuki R, Moriya K, Miyamoto H, Mori Y, Abe T, et al. Critical role of PA28 $\gamma$  in hepatitis C virus-associated steatogenesis and hepatocarcinogenesis. *Proc Natl Acad Sci U S A* 2007;104:1661–6.
159. Miyamoto H, Moriishi K, Moriya K, Murata S, Tanaka K, Suzuki T, et al. Involvement of the PA28 $\gamma$ -dependent pathway in insulin resistance induced by hepatitis C virus core protein. *J Virol* 2007;81:1727–35.
160. Nakai K, Okamoto T, Kimura-Someya T, Ishii K, Lim CK, Tani H, et al. Oligomerization of hepatitis C virus core protein is crucial for interaction with the cytoplasmic domain of E1 envelope protein. *J Virol* 2006;80:11265–73.
161. Baumert TF, Ito S, Wong DT, Liang TJ. Hepatitis C virus structural proteins assemble into viruslike particles in insect cells. *J Virol* 1998;72:3827–36.
162. Falcon V, Garcia C, de la Rosa MC, Menendez I, Seoane J, Grillo JM. Ultrastructural and immunocytochemical evidences of core-particle formation in the methylotrophic *Pichia pastoris* yeast when expressing HCV structural proteins (core-E1). *Tissue Cell* 1999;31:117–25.
163. Kunkel M, Lorinczi M, Rijnbrand R, Lemon SM, Watowich SJ. Self-assembly of nucleocapsid-like particles from recombinant hepatitis C virus core protein. *J Virol* 2001;75:2119–29.
164. Lorenzo LJ, Duenas-Carrera S, Falcon V, Acosta-Rivero N, Gonzalez E, de la Rosa MC, et al. Assembly of truncated HCV core antigen into virus-like particles in *Escherichia coli*. *Biochem Biophys Res Commun* 2001;281:962–5.
165. Acosta-Rivero N, Aguilar JC, Musacchio A, Falcon V, Vina A, de la Rosa MC, et al. Characterization of the HCV core virus-like particles produced in the methylotrophic yeast *Pichia pastoris*. *Biochem Biophys Res Commun* 2001;287:122–5.
166. Kunkel M, Watowich SJ. Conformational changes accompanying self-assembly of the hepatitis C virus core protein. *Virology* 2002;294:239–45.
167. Acosta-Rivero N, Falcon V, Alvarez C, Musacchio A, Chinae G, Cristina de la Rosa M, et al. Structured HCV nucleocapsids composed of P21 core protein assemble primary in the nucleus of *Pichia pastoris* yeast. *Biochem Biophys Res Commun* 2003;310:48–53.
168. Blanchard E, Hourieux C, Brand D, Ait-Goughoulte M, Moreau A, Trassard S, et al. Hepatitis C virus-like particle budding: role of the core protein and importance of its Asp111. *J Virol* 2003;77:10131–8.
169. Majeau N, Gagne V, Boivin A, Bolduc M, Majeau JA, Ouellet D, et al. The N-terminal half of the core protein of hepatitis C virus is sufficient for nucleocapsid formation. *J Gen Virol* 2004;85:971–81.
170. Klein KC, Polyak SJ, Lingappa JR. Unique features of hepatitis C virus capsid formation revealed by de novo cell-free assembly. *J Virol* 2004;78:9257–69.
171. Kunkel M, Watowich SJ. Biophysical characterization of hepatitis C virus core protein: implications for interactions within the virus and host. *FEBS Lett* 2004;557:174–80.
172. Matsumoto M, Hwang SB, Jeng KS, Zhu N, Lai MM. Homotypic interaction and multimerization of hepatitis C virus core protein. *Virology* 1996;218:43–51.
173. Nolandt O, Kern V, Muller H, Pfaff E, Theilmann L, Welker R, et al. Analysis of hepatitis C virus core protein interaction domains. *J Gen Virol* 1997;78(Pt 6):1331–40.
174. Yan BS, Tam MH, Syu WJ. Self-association of the C-terminal domain of the hepatitis-C virus core protein. *Eur J Biochem* 1998;258:100–6.
175. Ezelle HJ, Markovic D, Barber GN. Generation of hepatitis C virus-like particles by use of a recombinant vesicular stomatitis virus vector. *J Virol* 2002;76:12325–34.
176. Clayton RF, Owsianka A, Aitken J, Graham S, Bhella D, Patel AH. Analysis of antigenicity and topology of E2 glycoprotein present on recombinant hepatitis C virus-like particles. *J Virol* 2002;76:7672–82.
177. Lo S-Y, Selby MJ, Ou J-H. Interaction between hepatitis C virus core protein and E1 envelope protein. *J Virol* 1996;70:5177–82.

178. Ma HC, Ke CH, Hsieh TY, Lo SY. The first hydrophobic domain of the hepatitis C virus E1 protein is important for interaction with the capsid protein. *J Gen Virol* 2002;83:3085–92.
179. Hershko A, Ciechanover A. The ubiquitin system. *Annu Rev Biochem* 1998;67:425–79.
180. Finley D, Ciechanover A, Varshavsky A. Ubiquitin as a central cellular regulator. *Cell* 2004;116:S29–32, 2 p following S.
181. Suzuki R, Tamura K, Li J, Ishii K, Matsuura Y, Miyamura T, et al. Ubiquitin-mediated degradation of hepatitis C virus core protein is regulated by processing at its carboxyl terminus. *Virology* 2001;280:301–9.
182. Moriishi K, Okabayashi T, Nakai K, Moriya K, Koike K, Murata S, et al. Proteasome activator PA28 $\gamma$ -dependent nuclear retention and degradation of hepatitis C virus core protein. *J Virol* 2003;77:10237–49.
183. Shirakura M, Murakami K, Ichimura T, Suzuki R, Shimoji T, Fukuda K, et al. E6AP ubiquitin ligase mediates ubiquitylation and degradation of hepatitis C virus core protein. *J Virol* 2007; 81:1174–85.
184. Huijbregtse JM, Scheffner M, Beaudenon S, Howley PM. A family of proteins structurally and functionally related to the E6-AP ubiquitin–protein ligase. *Proc Natl Acad Sci USA* 1995;92: 2563–7.
185. Huijbregtse JM, Scheffner M, Howley PM. Cloning and expression of the cDNA for E6-AP, a protein that mediates the interaction of the human papillomavirus E6 oncoprotein with p53. *Mol Cell Biol* 1993;13:775–84.
186. Scheffner M, Huijbregtse JM, Vierstra RD, Howley PM. The HPV-16 E6 and E6-AP complex functions as a ubiquitin-protein ligase in the ubiquitination of p53. *Cell* 1993;75:495–505.
187. Tanahashi N, Yokota K, Ahn JY, Chung CH, Fujiwara T, Tanahashi E, et al. Molecular properties of the proteasome activator PA28 family proteins and gamma-interferon regulation. *Genes Cells* 1997;2:195–211.
188. Realini C, Jensen CC, Zhang Z, Johnston SC, Knowlton JR, Hill CP, et al. Characterization of recombinant REGalpha, REGbeta, and REGgamma proteasome activators. *J Biol Chem* 1997;272:25483–92.
189. Polyak SJ, Klein KC, Shoji I, Miyamura T, Lingappa JR. Assemble and interact pleiotropic functions of the HCV core protein. In: Tan S-L, editor. *Hepatitis C viruses: genomes and molecular biology*. Norwich, UK: Horizon Bioscience; 2006. p. 89–119.



# Hepatitis C viral life cycle <sup>☆</sup>

Tetsuro Suzuki <sup>\*</sup>, Koji Ishii, Hideki Aizaki, Takaji Wakita

*Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan*

Received 11 April 2007; accepted 11 April 2007

Available online 9 August 2007

## Abstract

Hepatitis C virus (HCV) has been recognized as a major cause of chronic liver diseases worldwide. Molecular studies of the virus became possible with the successful cloning of its genome in 1989. Although much work remains to be done regarding early and late stages of the HCV life cycle, significant progress has been made with respect to the molecular biology of HCV, especially the viral protein processing and the genome replication. This review summarizes our current understanding of genomic organization of HCV, features of the viral protein characteristics, and the viral life cycle.

© 2007 Elsevier B.V. All rights reserved.

*Keywords:* Hepatitis C virus; Translation; Polyprotein processing; RNA replication; Viral assembly

## Contents

1. Introduction . . . . .	1201
2. Genomic organization . . . . .	1201
3. Features of the viral proteins . . . . .	1202
3.1. Core protein . . . . .	1202
3.2. E1 and E2 envelope proteins. . . . .	1202
3.3. p7 protein . . . . .	1202
3.4. NS2 protein . . . . .	1203
3.5. NS3–4A complex . . . . .	1203
3.6. NS4B protein. . . . .	1203
3.7. NS5A protein. . . . .	1203
3.8. NS5B protein. . . . .	1203
4. HCV life cycle . . . . .	1204
4.1. Attachment and entry . . . . .	1204
4.2. Translation . . . . .	1204
4.3. Polyprotein processing . . . . .	1205
4.4. RNA replication . . . . .	1205
4.5. Viral assembly . . . . .	1206
5. Perspectives . . . . .	1207
Acknowledgements . . . . .	1207
References . . . . .	1207

<sup>☆</sup> This review is part of the *Advanced Drug Delivery Reviews* theme issue on “Toward Evidence Based Control of Hepatitis C Virus Infection”.

<sup>\*</sup> Corresponding author. Tel.: +81 3 5285 1111; fax: +81 3 5285 1161.

*E-mail address:* [tesuzuki@nih.go.jp](mailto:tesuzuki@nih.go.jp) (T. Suzuki).

## 1. Introduction

Since its discovery in 1989, representing a turning-point in the search for infectious agents associated with post-transfusion non-A, non-B hepatitis, hepatitis C virus (HCV) has been recognized as a major cause of chronic liver disease and affects approximately 200 million people worldwide at the present time [1–3]. Persistent infection with HCV is associated with the development of chronic hepatitis, hepatic steatosis, cirrhosis, and hepatocellular carcinoma [3–8]. In general, people with chronic hepatitis C are relatively asymptomatic and have few, if any, clinical manifestations prior to the development of cirrhosis.

HCV is a small, enveloped RNA virus belonging to the *Hepacivirus* genus of the *Flaviviridae* family, which also includes several classical flaviviruses, including dengue virus and yellow fever virus, as well as pestiviruses, such as bovine viral diarrhea virus and the unassigned GB viruses [9,10]. This review summarizes our current understanding of genomic organization of HCV, as well as features of the viral protein characteristics, and the viral life cycle.

## 2. Genomic organization

The HCV genome consists of a single-stranded positive-sense RNA of approximately 9.6 kb, which contains an open reading frame (ORF) encoding a polyprotein precursor of approximately 3000 residues flanked by untranslated regions (UTRs) at both ends [11]. The precursor is cleaved into at least 10 different proteins: the structural proteins Core, E1, E2 and p7, as well as the non-structural proteins NS2, NS3, NS4A, NS4B, NS5A and NS5B (Fig. 1).

An important feature of the HCV genome is its high degree of genetic variability [12,13]. Mutation rates, however, vary in different regions. The E1 and E2 regions are the most variable, while the 5'UTR and terminal segment of the 3'UTR have the highest degree of sequence conservation among various isolates. The 5' UTR, which is ~341 nucleotide (nt) in length, contains an

internal ribosomal entry site (IRES), which is essential for cap-independent translation of viral RNA, from which four highly structured domains (domains I–IV) are produced (Fig. 1) [14–19]. These are largely conserved among HCV and related viruses [15,16]. As with other RNA viruses with IRES-mediated expression, the HCV 5'NTR is thought to contain determinants for translation, as well as cis-acting elements for RNA replication. It has been shown that (i) the sequence upstream of the IRES is essential for viral RNA replication, (ii) sequences within the IRES are required for high-level HCV replication, and (iii) the stem-loop domain II of the IRES is crucial for replication [20]. A recent study has revealed that the 5'UTR is capable of binding to a liver-specific microRNA, miR-122, resulting in enhanced HCV RNA replication [21]. (Fig. 2).

The 3'UTR varies between 200 and 235 nt in length, including a short variable region, a poly(U/UC) tract with an average length of 80 nt, and a virtually invariant 98-nt X-tail region [22–24]. The X region forms three stable stem-loop structures that are highly conserved among all genotypes and, as a result, the HCV genome likely ends with a double-strand stem structure. It appears that the 3'X region, as well as the 52 nt upstream of the poly(U/C) tract, are crucial for RNA replication, while the remainder of the 3'UTR plays a role in enhancement of replication [25,26].

To date, hepaciviruses are divided into six principal genotypes of HCV that differ in their nucleotide sequences by 31–34%, and in their amino acid sequences by ~30%. HCV, like many other RNA viruses, circulates in infected individuals as a population of diverse but closely related variants referred to as quasispecies [12]. HCV heterogeneity is primarily due to a high error rate of the RNA-dependent RNA polymerase encoded by the NS5B gene. The existence of different quasispecies of the HCV genome appears to contribute to viral persistence. It has been shown that patients with chronic hepatitis C have greater genetic complexity in terms of the population of quasispecies they possess than patients with spontaneous clearance [13]. During the course of chronic infection, random genetic drift steadily induces the development of quasispecies primarily due to changes in the




Fig. 1.




Fig. 2.

hypervariable region 1, involving the 27 N-terminal 27 residues of the E2 envelope protein [27–29].

### 3. Features of the viral proteins

#### 3.1. Core protein

The HCV core protein, which is derived from the N-terminus of the polyprotein, most likely forms the viral nucleocapsid given similarities between its position and that of sequences encoding viral nucleocapsids in other flavivirus genomes. The amino acid sequence of the core protein is highly conserved among different HCV strains, compared with other HCV proteins. HCV core protein has been extensively used in a number of serologic assays since anti-core antibodies are highly prevalent among HCV-infected individuals. Although several core proteins of varying molecular weights have been identified [30–33], the core protein is released as a 191-residue precursor of 23 kDa and further processing yields the predominant form of 21 kDa. The N-terminal domain of the core protein is highly basic, while its C-terminus is hydrophobic. Several groups have reported a complex intracellular localization of the core protein [30,33–42]. The core protein is primarily detected in the cytoplasm, in association with the endoplasmic reticulum (ER), lipid droplets, and mitochondria. In some studies, a fraction of the core protein has also been found in the nucleus.

The ubiquitin–proteasome pathway, a major route by which selective protein degradation occurs in eukaryotic cells, is involved in post-translational modification of the core protein [32,43–45]. An initial report indicated that processing at the carboxyl-terminal hydrophobic domain of the core protein produced efficient polyubiquitylation and proteasomal degradation [32]. Recently, ubiquitin ligase E6AP has been identified as an HCV core-binding protein that enhances ubiquitylation and degradation of mature, as well as carboxyl-terminus truncated-core protein, and it has been suggested that E6AP-dependent degradation of the core protein is common to a

variety of HCV isolates and plays a critical role in the HCV life cycle [45].

The core protein is likely multifunctional and essential for viral replication, maturation, and pathogenesis. It is involved not only in formation of the HCV virion, but also has a number of regulatory functions, including modulation of signaling pathways, cellular and viral gene expression, cell transformation, apoptosis, and lipid metabolism [reviewed in 46].

#### 3.2. E1 and E2 envelope proteins

The E1 and E2 proteins are essential components of the virion envelope and are necessary for viral entry. These glycosylated proteins extend from aa 192–383 (E1) and from aa 384–746 (E2) of the polyprotein, and have molecular weights of 33–35 and 70–72 kDa, respectively [47]. Along the precursor polyprotein, it has been suggested that the C-terminal transmembrane domains of E1 and E2 form hairpin structures that pass through the membrane twice, thereby allowing processing by a signal peptide in the ER lumen [48]. Upon signal peptidase cleavage, the C-termini are thought to translocate into the cytoplasm in order to generate the type I membrane topology of mature E1 and E2. Mature E1 and E2 remain noncovalently associated, interacting in part through their C-terminal transmembrane domains, which also mediates retention of the E1–E2 complex in the ER. It has recently been demonstrated that, in addition to this conventional type I membrane topology, E1 protein also adopts a polytopic topology, in which the protein twice spans the ER membrane with an intervening cytoplasmic loop spanning aa 288–360 [49].

#### 3.3. p7 protein

The p7 protein is a small (63 aa) hydrophobic polypeptide that adopts a double membrane-spanning topology. This protein is essential for the production of infectious virions *in vivo* [50] and may belong to a small protein family of viroporins, which are known to enhance membrane permeability. It has been revealed



that p7 protein forms an ion channel in artificial lipid bilayers, suggesting it may function as a viroporin [51,52].

### 3.4. NS2 protein

The NS2 protein is a transmembrane protein of 21–23 kDa, with 96 highly hydrophobic N-terminal residues, forming either three or four transmembrane helices that insert into the ER membrane. The C-terminal part of NS2 presumably resides in the cytoplasm enabling zinc-stimulated NS2/3 autoprotease activity together with the N-terminal domain of NS3. Efficient cleavage at the NS2/3 site requires the 130 C-terminal residues and first 180 aa of the NS3. Site-directed mutagenesis has revealed that His-952, Glu-972, and Cys-993 may comprise the active site for proteolytic activity [53,54]. Deletion of NS2 from the nonstructural polyprotein has not been observed to abolish HCV RNA replication in cell cultures, indicating that NS2 is not essential for viral RNA replication [55,56]. However, the NS2 protein is essential for completion of the viral replication cycle *in vitro* and *in vivo* [57,58]. A recent report regarding the crystal structure of the C-terminus of NS2 suggests that the cytoplasmic domain of NS2 forms a dimeric cysteine protease with two composite active sites, in which His-952 and Glu-972 comprise the active site of one monomer, and Cys-993 contributes to the active site of the other [59].

### 3.5. NS3–4A complex

NS3–4A is a complex bifunctional molecule essential for viral polyprotein processing and RNA replication. NS3 is a fairly hydrophobic protein of 69 kDa with a serine protease encoded by its N-terminal one-third region that non-covalently binds the NS4A cofactor, which is a 54-aa polypeptide [reviewed in 60]. The catalytic triad is formed by residues His-1083, Asp-1107 and Ser-1165 of NS3. The central portion of NS4A is important for efficient processing of the nonstructural proteins by NS3. It has been suggested that the N-terminus of NS4A might form a transmembrane helix that anchors the NS3–4A complex to the cellular membrane [61]. Crystal structural analyses of the NS3/4A complex have demonstrated structural similarities between the NS3 serine protease and trypsin, with two large domains primarily composed of six-stranded beta barrels separated by a cleft containing the active site and substrate binding pocket [62–64]. Of note, NS4A forms an integral part of this structure and interacts with the extreme N-terminal residues of NS3 to form two additional anti-parallel beta-strands. The NS3–4A complex has a shallow substrate-binding pocket, thus requiring extended interaction sites with the substrate.

The final 442 aa of the C-terminal of NS3 comprise the helicase-NTPase domain, which is a member of the superfamily-2 DexH/D-box helicase, which unwind RNA-RNA substrates in a 3'-to -5' direction [reviewed in 65]. This is supported by crystal structure analysis indicating the presence of NTPase domains and RNA binding within the protein [66]. During RNA replication, the NS3 helicase is believed to translocate along the nucleic acid substrate by changing its protein conformation, utilizing the energy of NTP hydrolysis [67–69]. Its helicase activity is

positively modulated by the NS3 protease domain and NS4A [70].

### 3.6. NS4B protein

NS4B is an integral membrane protein of 27 kDa, which is predicted to contain at least four transmembrane domains and an N-terminal amphipathic helix that is responsible for membrane association [71]. NS4B has the ability to induce the formation of a specialized membrane compartment, a sort of membranous web where viral RNA replication may take place [72,73].

### 3.7. NS5A protein

NS5A is a membrane-anchored phosphoprotein that is observed in basally phosphorylated (56 kDa) and hyperphosphorylated (58 kDa) forms. Based on the results of a comparative sequence analysis following limited proteolysis of purified protein, NS5A is predicted to contain three domains: domain 1 (aa 1–213), domain 2 (aa 250–342) and domain 3 (aa 356–447) [74]. A recent structural study has demonstrated that domain 1 immediately follows the membrane-anchoring alpha-helix and forms a dimeric structure with an unconventional zinc-coordinating motif [75]. Thus, it may interact with viral and cellular proteins, as well as membranes and RNA.

While its function has not fully been elucidated, NS5A is believed to be important in viral replication. A large number of cell culture-adaptive mutations mapped to the NS5A have been shown to enhance RNA replication [76–78]. These adaptive mutations often affect hyperphosphorylation of NS5A, suggesting that the phosphorylation status of NS5A might influence replication efficiency. NS5A has been reported to interact with other HCV nonstructural proteins [79–81]. In addition, several cellular proteins interact with NS5A, resulting in assembly of the viral replication complex and/or regulation of RNA replication, as described below.

### 3.8. NS5B protein

NS5B is a 68-kDa protein with a conserved sequence motif characteristic of viral RNA-dependent RNA polymerase (RdRp), including a hallmark GDD motif that produces catalytic activity. NS5B is a tail-anchored protein and its C-terminal 21-aa region forms an alpha-helical transmembrane domain, which is dispensable for polymerase activity *in vitro* but is responsible for post-translational targeting to the cytoplasmic side of the ER [82,83]. Analysis of the crystal structure of NS5B has revealed that the HCV RdRp resembles a right hand and contains fingers, palm, and thumb subdomains, similar to other template-dependent polymerases [84–86]. Unlike the more open structures of other template-dependent DNA polymerases, such as the Klenow Fragment and the human immunodeficiency virus 1 reverse transcriptase, the HCV RdRp has a fully encircled active site through extensive interactions between the fingers and thumb subdomains, resulting in a protein that predominantly exists in a “closed” conformation. HCV RdRp also has an unusual hairpin loop that protrudes into the active site and helps position the 3'-

end of the RNA template for proper initiation of RNA synthesis and inhibits extension from a primed template [87].

## 4. HCV life cycle

### 4.1. Attachment and entry

Attachment of the virus to a cell followed by viral entry is the first step in the virus life cycle. In order to enter the host cell, the virus must first bind to a receptor on the cell surface. The specific interaction between a host cell receptor and viral attachment proteins on the surface of the virion determines tissue tropism and host range.

Low levels of HCV replication in cultured cells hindered study of the HCV life cycle. In order to overcome this, alternative models have been developed to study viral attachment and entry using recombinant HCV envelope proteins, including virus-like particles produced by baculovirus [88,89], vesicular stomatitis virus and retrovirus pseudotypes (HCVpp) [90–93], as well as infectious particles derived from a JFH-1 isolate (HCVcc) [94–96].

By using soluble E2 as a probe to identify cell-surface proteins potentially involved in HCV entry, CD81 was first identified as a putative HCV receptor [97]. CD81, a widely expressed 25-kDa cell surface protein, belongs to a family of tetraspanins and is involved in a number of activities, including cell adhesion, motility, metastasis, cell activation and signal transduction [98]. CD81 has a small and large extracellular loop, which mediate binding to recombinant E2 [97,99]. Studies with HCVpp and HCVcc confirm the involvement of CD81 in HCV entry. HCVpp shows a restricted tropism for human hepatic cell lines expressing CD81 [92,93,100–102]. Although necessary, CD81 expression alone is not sufficient for cell entry of HCVpp. Of note, HepG2, which does not express CD81 on its cell surface, is resistant to HCVpp infection, but over-expression of CD81 renders the HepG2 permissive to HCVpp infection [102–105]. Significant infection of CD81-negative cell lines with HCVpp has not been reported. However, as mentioned, not all CD81-positive cell lines can be infected [93,101,103]. Expression of CD81 in host cells is also required for infectivity of HCVcc. Recombinant CD81 and antibodies to CD81 have been observed to neutralize infection [94–96]. Thus, CD81 may function as a post-attachment entry co-receptor and may play a role after binding of the virion to another receptor.

The human scavenger receptor class B type I (SR-BI) has been identified as another putative receptor for HCV [106]. SR-BI is an 82-kDa glycoprotein with two C- and N-terminal cytoplasmic domains separated by a large extracellular domain involved in cellular lipometabolism. SR-BI is expressed in a wide variety of mammalian tissues and cell types [100,107,108], with particularly high levels of expression in the liver and steroidogenic tissue [107,109,110]. SR-BI recognition by soluble E2 requires the HVR1 of E2 [103,111]. A role of SR-BI in HCV cell entry has been confirmed using HCVpp in receptor competition assays using polyclonal anti-SR-BI serum, which has been observed to specifically inhibit HCVpp entry efficiently in a dose-dependent manner [103]. Recent reports have demonstrated that serum factors, especially high-density

lipoprotein (HDL), a ligand to SR-BI, enhance the infectivity of HCVpp [105,112–115]. These results suggest that SR-BI modulates HCV entry.

Several human cell lines co-expressing CD81 and SR-BI are non-permissive for HCVpp infection [100,102,103], suggesting that another cell surface molecule(s) may be required for HCV entry. C-type (calcium-dependent) lectins, such as L-SIGN, DC-SIGN, and the asialoglycoprotein receptor, have also been investigated as potential HCV receptors based on their affinity for recombinant HCV envelope proteins [116–119]. However, L-SIGN and DC-SIGN are not expressed on hepatocytes and therefore cannot be receptors for HCV entry. A possible role of L-SIGN and DC-SIGN involves the capture and transfer of HCV to hepatocytes [120,121]. The LDL receptor is another candidate receptor based on the finding that HCV particles associate with lipoproteins in serum and their infectivity correlates with lipoprotein association. The LDL receptor has been shown to mediate HCV internalization by binding to virion-associated LDL particles [122]. However, a role for the LDL receptor in virus entry has not been confirmed using HCVpp [93], likely since the binding is mediated by lipoproteins rather than viral components.

Recently, a tight junction component claudin-1 has been identified as a co-receptor of HCV [123]. Claudin-1 appears to be critical for HCV entry into hepatic cells and is thought to act during the late stages of viral entry.

### 4.2. Translation

As opposed to cellular capped mRNA molecules which are translated via a cap-dependent scanning mechanism, the naturally uncapped RNA molecules of viruses such as flaviviruses and picornaviruses are translated via a cap-independent IRES-mediated process, in which viral protein expression is regulated by direct recruitment of each ribosome to the start site of translation [18,19]. The first 40 nt of the 5'UTR, which include a single stem-loop (domain I), are not essential for translation. Of note, the 5' border of the IRES was mapped between nt 38 and 46 [17,124,125]. Other domains in the 5'UTR are more complex: domain II consisting of a stem with several internal loops, domain III consisting of a pseudoknot connected to a four-helix junction, as well as stem-loop III<sub>d</sub> and domain IV, a small hairpin containing the AUG start codon at nt 342. It has been suggested that the first 12 to 40 nt downstream of the start codon are also important for IRES activity [126–128].

Structural analysis of the HCV IRES indicates that all of the RNA elements adopt tertiary structures capable of binding to the translation initiation complex with high affinity [129]. IRES-mediated translation of HCV RNA is initiated by direct binding of a vacant 40S ribosomal subunit to the IRES. The 40S subunit appears to interact with the viral RNA at multiple sites including stems, loops, pseudoknots, as well as the start codon. This binary complex then binds to eukaryotic initiation factor (eIF) 3, as well as the ternary complex eIF-2: Met-tRNA<sub>i</sub>: GTP to form a 48S-like complex dependent upon both the basal domain III and the start codon. Subsequent formation of the 80S complex, which is the rate-limiting step, is dependent upon GTP

hydrolysis and attachment of the 60S subunit, after which the first peptide bond is formed [130].

In addition to the requirements described above, additional factors modulate IRES activity. Cellular factors such as the La autoantigen [131–133], heterogeneous ribonucleoprotein L [134], poly-C binding protein [135], and pyrimidine tract-binding protein (PTB)[136], have also been shown to bind to the IRES element and modulate HCV translation. HCV translation is also regulated through various interactions with viral proteins and the IRES.

We have found that HCV core protein expression inhibits HCV translation, possibly through binding to domain III<sub>d</sub>, particularly a GGG triplet within the hairpin loop structure of the domain [137–139]. We therefore propose a model in which competitive binding of the core protein for the IRES and 40S subunit regulates HCV translation. Although there is an increasing body of evidence to suggest involvement of the core protein in translational regulation, there is conflicting data regarding the mechanism by which this occurs. In contrast to studies describing modulation of initiation of HCV translation by expression of the core protein [137,139–141], another study suggest that the core protein sequence, and not the core protein itself, modulates HCV IRES function through a long-range RNA-RNA interaction [142].

In addition to the 10 known viral proteins (Fig. 1), the core-coding region of HCV has also been observed to express low levels of a 16–17 kDa protein [143–145]. This protein, which has been named the F protein, is thought to be produced by a +1 translational frameshift by ribosomes initiating translation at the start codon during synthesis of the HCV polyprotein, which shifts the reading frame between codons 9 and 11 of the polyprotein.

#### 4.3. Polyprotein processing

The main translation product of the HCV genome is a large precursor polyprotein that is subsequently processed by cellular and viral proteases into mature structural and nonstructural proteins (Fig. 1). As deduced from the hydrophobicity profile and dependence on microsomal membranes, junctions at core/E1, E1/E2, E2/p7, p7/NS2 are processed by host signal peptidases. Secondary structure analysis of the core protein has revealed that all major alpha helices are located in the C-terminal half of the protein. A predicted alpha helix encoded by aa 174–191 is extremely hydrophobic and resembles typical signal peptide sequences. Further post-translational cleavage close to the C terminus of the core protein takes place, removing the E1 signal sequence by a signal peptide peptidase [146–149]. This peptidase has been identified as a presenilin-type aspartic protease [150] and shown to exhibit protease activity within cellular membranes, resulting in cleavage of peptide bonds in the plane of lipid bilayers.

As described above, HCV nonstructural proteins are processed by two viral proteases: cleavage between NS2 and NS3 is a rapid intramolecular reaction mediated by a NS2-3 protease spanning NS2 and the N-terminal domain of NS3, whereas the remaining four junctions are processed by a serine protease located within the 180 N-terminal residues of the NS3 protein. The NS3-NS5B

region is presumably processed by sequential cleavage: NS3/4A → NS5A/5B → NS4A/4B → NS4B/5A [151–154]. Processing at the NS3/4A site is intramolecular, whereas cleavage at the other sites occurs intermolecularly.

#### 4.4. RNA replication

As with other positive-strand RNA viruses, HCV replication is assumed to start with synthesis of a complementary negative-strand RNA using the genome as a template, after which genomic positive-strand RNA is produced from a negative-strand RNA template, both steps of which are catalyzed by the NS5B RdRp. The positive-strand RNA progeny are transcribed at a level 5- to 10-fold that of negative-strand RNA. Recombinant NS5B protein demonstrates RdRp activity *in vitro*, however, appears to lack strict template specificity and fidelity, which are essential for viral RNA synthesis. Thus, other viral and/or host factors are believed to be responsible for RNA replication and formation of the replication complex (RC), together with NS5B, which is required for catalyzing HCV RNA synthesis during replication. Several research groups have demonstrated HCV RC-mediated replication *in vitro* in crude membrane fractions of cells harboring subgenomic replicons [155–158]. Studies of cell-free replication systems, which provide a useful source of viral RCs have revealed that RNA synthesis can be initiated in the absence of additional negative-strand template RNA, suggesting that pre-initiated template RNA co-purifies with viral RCs [156–159].

Co-precipitation and immunostaining studies have revealed that newly synthesized HCV RNA exists as distinct specks of material, while all of the viral nonstructural proteins coexist [160]. These distinct structures may be equivalent to a membranous web, as described above. Expression of all structural and non-structural proteins in the context of the entire HCV polyprotein has been observed to induce similar membrane changes [72]. It is of interest that morphologically similar structures, termed sponge-like inclusions [161], have been identified by electron microscopy within the hepatocytes of HCV-infected chimpanzees. Thus, HCV RC may exist in the context of a membranous web in infected cells. Because all nonstructural proteins of HCV are associated with the ER membrane in cells harboring subgenomic replicon RNA molecules [162,163], and since the membranous web is frequently observed in close proximity with the ER membrane, it is likely that the membranous web in HCV-infected cells is derived from the ER membrane.

On the other hand, there is accumulating evidence to support an association between HCV RNA replication and detergent-insoluble membrane domains or lipid rafts, which are microdomains rich in cholesterol and sphingolipids. Membrane flotation analysis and replication assays have shown that viral RNA and proteins exist within detergent-resistant, lipid-raft membranes, and that RNA replication occurs even after treatment with detergent [155,164]. Inhibitors of *de novo* sphingolipid synthesis have been shown to inhibit HCV replication, presumably by disrupting the association of viral nonstructural proteins with lipid rafts [165,166]. It is now accepted that HCV nonstructural proteins synthesized at the ER localize to lipid raft membranes when they are actively engaged in RNA replication.

Membrane separation analysis has demonstrated that HCV nonstructural proteins exist both in the ER and the Golgi apparatus, but that viral RNA replication primarily occurs in the Golgi fraction [155]. Further studies to elucidate the cellular processes involved in HCV RC formation and replication of the HCV genome in infected cells are needed.

Studies of RNA replicons have demonstrated the greatest viral RNA levels during the growth phase of the cells, after which a significant drop is observed as the cells reach confluence, suggesting that HCV replication and/or translation is tightly linked to host cell metabolism [163]. Huh-7 cells, in which adapted replicons are cured by treatment with IFN, yield cell populations that are more permissive for the replicon tested. Thus, it is likely that some interplay between the cellular environment and specific adaptive mutations of viral RNA contributes to efficient RNA replication of HCV.

Several cellular proteins capable of interacting with NS5A, such as vesicle-associated membrane protein-associated protein (VAP) subtypes A and B (VAP-A and -B) [73,167], FKBP8 [168], FBL2 [169,170], growth factor receptor-bound protein 2 adaptor protein [171], SRCAP [172], and karyopherin b3 [173], as well as Raf-1 kinase [174], have been identified. VAP-A and -B and SNARE-like proteins are known to localize within the ER and Golgi apparatus and are essential for HCV replication by binding with both NS5A and NS5B. VAP-A interacts with VAP-B through its transmembrane domain. Thus, VAP-A and -B are thought to be involved in the formation of functional HCV RCs. FKBP8, a member of the FK506-binding protein family, and Hsp90 form a complex with NS5A, further contributing to viral RNA replication. Statins that decrease the production of mevalonate by inhibiting 3-hydroxy-3-methylglutaryl CoA reductase have been shown to inhibit HCV RNA replication [170,175], which can be reversed by adding geranylgeraniol, suggesting that viral replication requires geranylgeranylated proteins. A geranylgeranylated protein, FBL2, which contains an F-box motif and is therefore likely involved in protein degradation, has been identified as a NS5A-binding protein.

Host factors that interact with NS5B and might participate in HCV replication include cyclophilin B [176], p68 [177], nucleolin [178,179], and hnRNP A1 [180]. Cyclophilin B, a cellular peptidyl-prolyl cis-trans isomerase, interacts with the C-terminal region of NS5B to directly stimulate its RNA binding activity, and thereby contributes to efficient replication of HCV RNA. Redistribution of p68, an RNA helicase, from the nucleus to the cytoplasm occurs through its binding to NS5B, and the p68-NS5B interaction may further serve to mediate HCV replication. Nucleolin, a representative nucleolar marker, interacts with NS5B through two independent regions of NS5B and may be essential for HCV replication. hnRNP A1, a heterogeneous nuclear ribonucleoprotein, also interacts with septin 6, as well as the 5'-UTR and 3'-UTR of HCV RNA, and contains the cis-acting elements required for replication. Thus, hnRNP A1 and septin 6 play important roles in HCV replication through RNA-protein and protein-protein interactions. Other cellular components that bind to HCV RNA, such as PTB, may also be involved in viral replication. PTB has been observed to modulate HCV IRES

activity by binding to several sites within the viral genome [22,181–184]. Recent studies have shown that PTB also forms part of the HCV RC and participates in viral RNA synthesis [185].

#### 4.5. Viral assembly

Little is known about the assembly of HCV or its virion structure since efficient production of authentic HCV particles has only recently been achieved. As with related viruses, the mature HCV virion likely consists of a nucleocapsid and outer envelope composed of a lipid membrane and envelope proteins. Various forms of HCV have been reported to circulate in the sera of infected hosts, including (i) free mature virions, (ii) virions bound to low-density lipoproteins and very-low-density lipoproteins, (iii) virions bound to immunoglobulins, as well as (iv) non-enveloped nucleocapsids, which exhibit different physicochemical and antigenic properties [186–189].

Several expression systems have been used to investigate HCV capsid assembly using lysate from mammalian cells, insects, yeast, bacteria, and reticulocytes, as well as purified recombinant protein [88,89,190–195]. The results suggest that the immunogenic nucleocapsid-like particles of HCV are variable in size ranging from 30 to 80 nm in diameter. The N-terminal half of the core protein is important for nucleocapsid assembly [190,194,195]. HCV capsid formation occurs in the presence or absence of ER-derived membrane, which supports cleavage of the signal peptide at the C-terminus [195].

Nucleocapsid assembly generally involves oligomerization of the capsid protein and encapsidation of genomic RNA. This process is thought to occur upon interaction of the core protein with viral RNA, and the core-RNA interaction may be critical for switching from RNA replication to packaging. In fact, HCV core protein can bind to positive-strand HCV RNA through stem-loop domains I, III and nt 24–41 [138]. Two-hybrid systems have identified a potential homotypic interaction domain within the N-terminal region of the core protein (aa 1–115 or -122), with particular emphasis on the region encompassing aa 82–102 [196,197]. Using purified HCV core protein, a C-terminally truncated core protein (aa 1–124) and structured RNA have been implicated in nucleocapsid formation to produce homogenous spherical HCV particles. When core protein containing the C terminus up to aa 174 is similarly examined, a heterogeneous array of irregularly shaped particles is observed, suggesting that the C-terminus of the core protein influences self-assembly. Furthermore, Pro substitution within the C-terminal region has been observed to abolish core protein self-interaction [198]. Circular dichroism spectroscopy has further shown that a Trp-rich region spanning aa 76–113 is largely solvent-exposed and unlikely to play a role in multimerization. Recently, our group has demonstrated that self-oligomerization of the core protein is promoted by aa 72 to 91 of the core protein [49].

Once a HCV nucleocapsid is formed in the cytoplasm, it acquires an envelope as it buds through an intracellular membrane. Interactions between the core and E1/E2 proteins are thought to determine viral morphology. Expression of HCV structural proteins using recombinant virus vectors has succeeded in generating virus-like particles with similar ultrastructural

properties to HCV virions. Packaging of these HCV-like particles into intracellular vesicles as a result of budding from the ER has been noted [88,199,200]. Mapping studies to determine the nature of interaction between core and E1 proteins have demonstrated the importance of C-terminal regions in this interaction [201,202]. Since corresponding sequences are not well conserved among various HCV isolates, interactions between core and E1 proteins might depend more on hydrophobicity than specific sequences. In contrast, it has been shown that the interaction between self-oligomerized HCV core protein and the E1 glycoprotein is mediated through a cytoplasmic loop of the polytopic form of the E1 protein [49].

It is believed that HCV particles are released from the cell through the secretory pathway. HCV structural proteins have been observed both in the ER and Golgi apparatus [203]. In addition, complex N-linked glycans, which transit through the Golgi apparatus, have been detected on the surface of HCV particles isolated from patient sera [204].

## 5. Perspectives

Since the discovery of HCV, which is a major cause of liver disease worldwide, significant progress has been made regarding the molecular biology of this virus. However, details regarding early and late stages of the HCV life cycle, including cell entry, genome packaging, assembly and release, remain unclear. In addition, the role of some viral proteins and their importance to replication remains unclear, as well as the role of certain host factors in regulation of the HCV life cycle.

## Acknowledgements

The authors are grateful to all of their co-workers who contributed to the studies cited here. We also thank T. Mizoguchi for her secretarial work. This work was supported in part by a grant for Research on Health Sciences focusing on Drug Innovation from the Japan Health Sciences Foundation, and by grants-in-aid from the Ministry of Health, Labor, and Welfare, Japan.

## References

- [1] G. Kuo, Q.-L. Choo, H.J. Alter, G.L. Gitnick, A.G. Redeker, R.H. Purcell, T. Miyamura, J.L. Dienstag, M.J. Alter, C.E. Stevens, G.E. Tegtmeier, F. Bonino, M. Colombo, W.-S. Lee, C. Kuo, K. Berger, F.R. Shuster, L.R. Overby, D.W. Bradley, M. Houghton, An assay for circulating antibodies to a major etiologic virus of human non-A, non-B hepatitis, *Science* 244 (1989) 362–364.
- [2] A. Grakoui, H.L. Hanson, C.M. Rice, Bad time for Bonzo? Experimental models of hepatitis C virus infection, replication, and pathogenesis, *Hepatology* 33 (3) (2001) 489–495.
- [3] G.M. Lauer, B.D. Walker, Hepatitis C virus infection, *N. Engl. J. Med.* 345 (1) (2001) 41–52.
- [4] I. Saito, T. Miyamura, A. Ohbayashi, H. Harada, T. Katayama, S. Kikuchi, Y. Watanabe, S. Koi, M. Onji, Y. Ohta, et al., Hepatitis C virus infection is associated with the development of hepatocellular carcinoma, *Proc. Natl. Acad. Sci. U. S. A.* 87 (17) (1990) 6547–6549.
- [5] M.J. Alter, Epidemiology of hepatitis C in the West, *Semin. Liver Dis.* 15 (1) (1995) 5–14.
- [6] M. Di Bisceglie, Hepatitis C and hepatocellular carcinoma, *Hepatology* 26 (3 Suppl 1) (1997) 34S–38S.
- [7] T. Poynard, M.F. Yuen, V. Ratziu, C.L. Lai, Viral hepatitis C, *Lancet* 362 (9401) (2003) 2095–2100.
- [8] J.M. Pawlotsky, Pathophysiology of hepatitis C virus infection and related liver disease, *Trends Microbiol.* 12 (2) (2004) 96–102.
- [9] M. Houghton, A. Weiner, J. Han, G. Kuo, Q.-L. Choo, Molecular biology of the hepatitis C viruses: implications for diagnosis, development and control of viral disease, *Hepatology* 14 (1991) 381–388.
- [10] B. Robertson, G. Myers, C. Howard, T. Bretin, J. Bukh, B. Gaschen, T. Gojobori, G. Maertens, M. Mizokami, O. Nainan, S. Netesov, K. Nishioka, T. Shin i, P. Simmonds, D. Smith, L. Stuyver, A. Weiner, Classification, nomenclature, and database development for hepatitis C virus (HCV) and related viruses: proposals for standardization. International Committee on Virus Taxonomy, *Arch. Virol.* 143 (12) (1998) 2493–2503.
- [11] Q.-L. Choo, G. Kuo, A.J. Weiner, L.R. Overby, D.W. Bradley, M. Houghton, Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome, *Science* 244 (1989) 359–362.
- [12] M. Martell, J.I. Esteban, J. Quer, J. Genesca, A. Weiner, R. Esteban, J. Guardia, J. Gomez, Hepatitis C virus (HCV) circulates as a population of different but closely related genomes: quasispecies nature of HCV genome distribution, *J. Virol.* 66 (1992) 3225–3229.
- [13] J.M. Pawlotsky, Hepatitis C virus population dynamics during infection, *Curr. Top. Microbiol. Immunol.* 299 (2006) 261–284.
- [14] J. Bukh, R.H. Purcell, R.H. Miller, Sequence analysis of the 5′ noncoding region of hepatitis C virus, *Proc. Natl. Acad. Sci. U. S. A.* 89 (11) (1992) 4942–4946.
- [15] E.A. Brown, H. Zhang, L.H. Ping, S.M. Lemon, Secondary structure of the 5′ nontranslated regions of hepatitis C virus and pestivirus genomic RNAs, *Nucleic Acids Res.* 20 (19) (1992) 5041–5045.
- [16] M. Honda, M.R. Beard, L.H. Ping, S.M. Lemon, A phylogenetically conserved stem-loop structure at the 5′ border of the internal ribosome entry site of hepatitis C virus is required for cap-independent viral translation, *J. Virol.* 73 (2) (1999) 1165–1174.
- [17] M. Honda, E.A. Brown, S.M. Lemon, Stability of a stem-loop involving the initiator AUG controls the efficiency of internal initiation of translation on hepatitis C virus RNA, *RNA* 2 (10) (1996) 955–968.
- [18] K. Tsukiyama-Kohara, N. Iizuka, M. Kohara, A. Nomoto, Internal ribosome entry site within hepatitis C virus RNA, *J. Virol.* 66 (1992) 1476–1483.
- [19] C. Wang, P. Sarnow, A. Siddiqui, Translation of human hepatitis C virus RNA in cultured cells is mediated by an internal ribosome-binding mechanism, *J. Virol.* 67 (1993) 3338–3344.
- [20] P. Friebe, V. Lohmann, N. Krieger, R. Bartenschlager, Sequences in the 5′ nontranslated region of hepatitis C virus required for RNA replication, *J. Virol.* 75 (24) (2001) 12047–12057.
- [21] L. Jopling, M. Yi, A.M. Lancaster, S.M. Lemon, P. Sarnow, Modulation of hepatitis C virus RNA abundance by a liver-specific MicroRNA, *Science* 309 (5740) (2005) 1577–1581.
- [22] T. Ito, M.M. Lai, An internal polypyrimidine-tract-binding protein-binding site in the hepatitis C virus RNA attenuates translation, which is relieved by the 3′-untranslated sequence, *Virology* 254 (2) (1999) 288–296.
- [23] A. Kolykhalov, E.V. Agapov, K.J. Blight, K. Mihalik, S.M. Feinstone, C.M. Rice, Transmission of hepatitis C by intrahepatic inoculation with transcribed RNA, *Science* 277 (1997) 570–574.
- [24] T. Tanaka, N. Kato, M.-J. Cho, K. Shimotohno, A novel sequence found at the 3′ terminus of hepatitis C virus genome, *Biochem. Biophys. Res. Commun.* 215 (1995) 744–749.
- [25] M. Yi, S.M. Lemon, 3′ nontranslated RNA signals required for replication of hepatitis C virus RNA, *J. Virol.* 77 (6) (2003) 3557–3568.
- [26] P. Friebe, R. Bartenschlager, Genetic analysis of sequences in the 3′ nontranslated region of hepatitis C virus that are important for RNA replication, *J. Virol.* 76 (11) (2002) 5326–5338.
- [27] M. Hijikata, N. Kato, Y. Ootsuyama, M. Nakagawa, S. Ohkoshi, K. Shimotohno, Hypervariable regions in the putative glycoprotein of hepatitis C virus, *Biochem. Biophys. Res. Commun.* 175 (1991) 220–228.

- [28] J. Weiner, M.J. Brauer, J. Rosenblatt, K.H. Richman, J. Tung, K. Crawford, F. Bonino, G. Saracco, Q.-L. Choo, M. Houghton, Variable and hypervariable domains are found in the regions of HCV corresponding to the flavivirus envelope and NS1 proteins and the pestivirus envelope glycoproteins, *Virology* 180 (1991) 842–848.
- [29] N. Kato, Y. Ootsuyama, T. Tanaka, M. Nakagawa, T. Nakazawa, K. Muraiso, S. Ohkoshi, M. Hijikata, K. Shimotohno, Marked sequence diversity in the putative envelope proteins of hepatitis C viruses, *Virus Res.* 22 (1992) 107–123.
- [30] S. Harada, Y. Watanabe, K. Takeuchi, T. Suzuki, T. Katayama, Y. Takebe, I. Saito, T. Miyamura, Expression of processed core protein of hepatitis C virus in mammalian cells, *J. Virol.* 65 (1991) 3015–3021.
- [31] Q. Liu, C. Tackney, R.A. Bhat, A.M. Prince, P. Zhang, Regulated processing of hepatitis C virus core protein is linked to subcellular localization, *J. Virol.* 71 (1) (1997) 657–662.
- [32] R. Suzuki, K. Tamura, J. Li, K. Ishii, Y. Matsuura, T. Miyamura, T. Suzuki, Ubiquitin-mediated degradation of hepatitis C virus core protein is regulated by processing at its carboxyl terminus, *Virology* 280 (2) (2001) 301–309.
- [33] K. Yasui, T. Wakita, K. Tsukiyama-Kohara, S.I. Funahashi, M. Ichikawa, T. Kajita, D. Moradpour, J.R. Wands, M. Kohara, The native form and maturation process of hepatitis C virus core protein, *J. Virol.* 72 (7) (1998) 6048–6055.
- [34] J. Selby, Q.-L. Choo, K. Berger, G. Kuo, E. Glazer, M. Eckart, C. Lee, D. Chien, C. Kuo, M. Houghton, Expression, identification and subcellular localization of the proteins encoded by the hepatitis C viral genome, *J. Gen. Virol.* 74 (1993) 1103–1113.
- [35] R. Suzuki, Y. Matsuura, T. Suzuki, A. Ando, J. Chiba, S. Harada, I. Saito, T. Miyamura, Nuclear localization of the truncated hepatitis C virus core protein with its hydrophobic C terminus deleted, *J. Gen. Virol.* 76 (1995) 53–61.
- [36] D. Moradpour, T. Wakita, K. Tokushige, R.I. Carlson, K. Krawczynski, J.R. Wands, Characterization of three novel monoclonal antibodies against hepatitis C virus core protein, *J. Med. Virol.* 48 (3) (1996) 234–241.
- [37] G. Barba, F. Harper, T. Harada, M. Kohara, S. Goulinet, Y. Matsuura, G. Eder, Z. Schaff, M.J. Chapman, T. Miyamura, C. Brechot, Hepatitis C virus core protein shows a cytoplasmic localization and associates to cellular lipid storage droplets, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 1200–1205.
- [38] K. Moriya, H. Fujie, Y. Shintani, H. Yotsuyanagi, T. Tsutsumi, K. Ishibashi, Y. Matsuura, S. Kimura, The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice, *Nat. Med.* 4 (1998) 1065–1067.
- [39] R.G. Hope, J. McLauchlan, Sequence motifs required for lipid droplet association and protein stability are unique to the hepatitis C virus core protein, *J. Gen. Virol.* 81 (Pt 8) (2000) 1913–1925.
- [40] M. Okuda, K. Li, M.R. Beard, L.A. Showalter, F. Scholle, S.M. Lemon, S.A. Weinman, Mitochondrial injury, oxidative stress, and antioxidant gene expression are induced by hepatitis C virus core protein, *Gastroenterology* 122 (2) (2002) 366–375.
- [41] B. Schwer, S. Ren, T. Pietschmann, J. Kartenbeck, K. Kaehlcke, R. Bartenschlager, T.S. Yen, M. Ott, Targeting of hepatitis C virus core protein to mitochondria through a novel C-terminal localization motif, *J. Virol.* 78 (15) (2004) 7958–7968.
- [42] R. Suzuki, S. Sakamoto, T. Tsutsumi, A. Rikimaru, K. Tanaka, T. Shimoike, K. Moriishi, T. Iwasaki, K. Mizumoto, Y. Matsuura, T. Miyamura, T. Suzuki, Molecular determinants for subcellular localization of hepatitis C virus core protein, *J. Virol.* 79 (2) (2005) 1271–1281.
- [43] K. Moriishi, R. Mochizuki, K. Moriya, H. Miyamoto, Y. Mori, T. Abe, S. Murata, K. Tanaka, T. Miyamura, T. Suzuki, K. Koike, Y. Matsuura, Critical role of PA28 $\{\gamma\}$  in hepatitis C virus-associated steatogenesis and hepatocarcinogenesis, *Proc. Natl. Acad. Sci. U. S. A.* 104 (5) (2007) 1661–1666.
- [44] K. Moriishi, T. Okabayashi, K. Nakai, K. Moriya, K. Koike, S. Murata, T. Chiba, K. Tanaka, R. Suzuki, T. Suzuki, T. Miyamura, Y. Matsuura, Proteasome activator PA28 $\gamma$ -dependent nuclear retention and degradation of hepatitis C virus core protein, *J. Virol.* 77 (19) (2003) 10237–10249.
- [45] M. Shirakura, K. Murakami, T. Ichimura, R. Suzuki, T. Shimoji, K. Fukuda, K. Abe, S. Sato, M. Fukasawa, Y. Yamakawa, M. Nishijima, K. Moriishi, Y. Matsuura, T. Wakita, T. Suzuki, P.M. Howley, T. Miyamura, I. Shoji, E6AP ubiquitin ligase mediates ubiquitylation and degradation of hepatitis C virus core protein, *J. Virol.* 81 (3) (2007) 1174–1185.
- [46] T. Suzuki, R. Suzuki, Maturation and assembly of hepatitis C virus core protein, in: M. Kalitzky, P. Borowski (Eds.), *Molecular Biology of the Flavivirus*, Horizon Bioscience, Norfolk, U.K., 2006, pp. 295–311.
- [47] A. Op De Beeck, L. Cocquerel, J. Dubuisson, Biogenesis of hepatitis C virus envelope glycoproteins, *J. Gen. Virol.* 82 (Pt 11) (2001) 2589–2595.
- [48] L. Cocquerel, A. Op de Beeck, M. Lambot, J. Roussel, D. Delgrange, A. Pillez, C. Wychowski, F. Penin, J. Dubuisson, Topological changes in the transmembrane domains of hepatitis C virus envelope glycoproteins, *EMBO J.* 21 (12) (2002) 2893–2902.
- [49] K. Nakai, T. Okamoto, T. Kimura-Someya, K. Ishii, C.K. Lim, H. Tani, E. Matsuo, T. Abe, Y. Mori, T. Suzuki, T. Miyamura, J.H. Nunberg, K. Moriishi, Y. Matsuura, Oligomerization of hepatitis C virus core protein is crucial for interaction with the cytoplasmic domain of E1 envelope protein, *J. Virol.* 80 (22) (2006) 11265–11273.
- [50] A. Sakai, M.S. Claire, K. Faulk, S. Govindarajan, S.U. Emerson, R.H. Purcell, J. Bukh, The p7 polypeptide of hepatitis C virus is critical for infectivity and contains functionally important genotype-specific sequences, *Proc. Natl. Acad. Sci. U. S. A.* 100 (20) (2003) 11646–11651.
- [51] D. Pavlovic, D.C. Neville, O. Argaud, B. Blumberg, R.A. Dwek, W.B. Fischer, N. Zitzmann, The hepatitis C virus p7 protein forms an ion channel that is inhibited by long-alkyl-chain iminosugar derivatives, *Proc. Natl. Acad. Sci. U. S. A.* 100 (10) (2003) 6104–6108.
- [52] S.D. Griffin, L.P. Beales, D.S. Clarke, O. Worsfold, S.D. Evans, J. Jaeger, M.P. Harris, D.J. Rowlands, The p7 protein of hepatitis C virus forms an ion channel that is blocked by the antiviral drug, Amantadine, *FEBS Lett.* 535 (1–3) (2003) 34–38.
- [53] A. Grakoui, D.W. McCourt, C. Wychowski, S.M. Feinstone, C.M. Rice, A second hepatitis C virus-encoded proteinase, *Proc. Natl. Acad. Sci. U. S. A.* 90 (1993) 10583–10587.
- [54] M. Hijikata, H. Mizushima, T. Akagi, S. Mori, N. Kakiuchi, N. Kato, T. Tanaka, K. Kimura, K. Shimotohno, Two distinct proteinase activities required for the processing of a putative nonstructural precursor protein of hepatitis C virus, *J. Virol.* 67 (1993) 4665–4675.
- [55] V. Lohmann, F. Komer, J. Koch, U. Herian, L. Theilmann, R. Bartenschlager, Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line, *Science* 285 (1999) 110–113.
- [56] A. Khromykh, E.G. Westaway, Subgenomic replicons of the flavivirus Kunjin: construction and applications, *J. Virol.* 71 (2) (1997) 1497–1505.
- [57] T. Pietschmann, A. Kaul, G. Koutsoudakis, A. Shavinskaya, S. Kallis, E. Steinmann, K. Abid, F. Negro, M. Dreux, F.L. Cosset, R. Bartenschlager, Construction and characterization of infectious intragenotypic and intergenotypic hepatitis C virus chimeras, *Proc. Natl. Acad. Sci. U. S. A.* 103 (19) (2006) 7408–7413.
- [58] A. Kolykhalov, K. Mihalik, S.M. Feinstone, C.M. Rice, Hepatitis C virus-encoded enzymatic activities and conserved RNA elements in the 3' nontranslated region are essential for virus replication *in vivo*, *J. Virol.* 74 (2000) 2046–2051.
- [59] C. Lorenz, J. Marcotrigiano, T.G. Dentzer, C.M. Rice, Structure of the catalytic domain of the hepatitis C virus NS2-3 protease, *Nature* 442 (7104) (2006) 831–835.
- [60] R. De Francesco, G. Migliaccio, Challenges and successes in developing new therapies for hepatitis C, *Nature* 436 (7053) (2005) 953–960.
- [61] B. Wolk, D. Sansonno, H.G. Krausslich, F. Dammacco, C.M. Rice, H.E. Blum, D. Moradpour, Subcellular localization, stability, and trans-cleavage competence of the hepatitis C virus NS3-NS4A complex expressed in tetracycline-regulated cell lines, *J. Virol.* 74 (5) (2000) 2293–2304.
- [62] L. Kim, K.A. Morgenstern, C. Lin, T. Fox, M.D. Dwyer, J.A. Landro, S.P. Chambers, W. Markland, C.A. Lepre, E.T. O'Malley, S.L. Harbeson, C.M. Rice, M.A. Murcko, P.R. Caron, J.A. Thomson, Crystal structure of the hepatitis C virus NS3 protease domain complexed with a synthetic NS4A cofactor peptide, *Cell* 87 (1996) 343–355.



- [63] R.A. Love, H.E. Parge, J.A. Wickersham, Z. Hostomsky, N. Habuka, E.W. Moomaw, T. Adachi, Z. Hostomska, The crystal structure of hepatitis C virus NS3 proteinase reveals a trypsin-like fold and a structural zinc binding site, *Cell* 87 (1996) 331–342.
- [64] Y. Yan, Y. Li, S. Munshi, V. Sardana, J.L. Cole, M. Sardana, C. Steinkuehler, L. Tomei, R. De Francesco, L.C. Kuo, Z. Chen, Complex of NS3 protease and NS4A peptide of BK strain hepatitis C virus: a 2.2 Å resolution structure in a hexagonal crystal form, *Protein Sci.* 7 (4) (1998) 837–847.
- [65] D. Kwong, J.L. Kim, C. Lin, Structure and function of hepatitis C virus NS3 helicase, *Curr. Top. Microbiol. Immunol.* 242 (2000) 171–196.
- [66] L. Kim, K.A. Morgenstern, J.P. Griffith, M.D. Dwyer, J.A. Thomson, M.A. Murcko, C. Lin, P.R. Caron, Hepatitis C virus NS3 RNA helicase domain with a bound oligonucleotide: the crystal structure provides insights into the mode of unwinding, *Structure* 6 (1) (1998) 89–100.
- [67] V. Serebrov, A.M. Pyle, Periodic cycles of RNA unwinding and pausing by hepatitis C virus NS3 helicase, *Nature* 430 (6998) (2004) 476–480.
- [68] M.K. Levin, M. Gurjar, S.S. Patel, A Brownian motor mechanism of translocation and strand separation by hepatitis C virus helicase, *Nat. Struct. Mol. Biol.* 12 (5) (2005) 429–435.
- [69] S. Dumont, W. Cheng, V. Serebrov, R.K. Beran, I. Tinoco Jr., A.M. Pyle, C. Bustamante, RNA translocation and unwinding mechanism of HCV NS3 helicase and its coordination by ATP, *Nature* 439 (7072) (2006) 105–108.
- [70] D.N. Frick, R.S. Rypma, A.M. Lam, B. Gu, The nonstructural protein 3 protease/helicase requires an intact protease domain to unwind duplex RNA efficiently, *J. Biol. Chem.* 279 (2) (2004) 1269–1280.
- [71] M. Lundin, M. Monne, A. Widell, G. Von Heijne, M.A. Persson, Topology of the membrane-associated hepatitis C virus protein NS4B, *J. Virol.* 77 (9) (2003) 5428–5438.
- [72] D. Egger, B. Wolk, R. Gosert, L. Bianchi, H.E. Blum, D. Moradpour, K. Bienz, Expression of hepatitis C virus proteins induces distinct membrane alterations including a candidate viral replication complex, *J. Virol.* 76 (12) (2002) 5974–5984.
- [73] L. Gao, H. Aizaki, J.W. He, M.M. Lai, Interactions between viral nonstructural proteins and host protein hVAP-33 mediate the formation of hepatitis C virus RNA replication complex on lipid raft, *J. Virol.* 78 (7) (2004) 3480–3488.
- [74] T.L. Tellinghuisen, J. Marcotrigiano, A.E. Gorbalenya, C.M. Rice, The NS5A protein of hepatitis C virus is a zinc metalloprotein, *J. Biol. Chem.* 279 (47) (2004) 48576–48587.
- [75] T.L. Tellinghuisen, J. Marcotrigiano, C.M. Rice, Structure of the zinc-binding domain of an essential component of the hepatitis C virus replicase, *Nature* 435 (7040) (2005) 374–379.
- [76] K.J. Blight, A.A. Kolykhalov, C.M. Rice, Efficient initiation of HCV RNA replication in cell culture, *Science* 290 (2000) 1972–1974.
- [77] N. Krieger, V. Lohmann, R. Bartenschlager, Enhancement of hepatitis C virus RNA replication by cell culture-adaptive mutations, *J. Virol.* 75 (10) (2001) 4614–4624.
- [78] V. Lohmann, F. Korner, A. Dobierzewska, R. Bartenschlager, Mutations in hepatitis C virus RNAs conferring cell culture adaptation, *J. Virol.* 75 (3) (2001) 1437–1449.
- [79] T. Shimakami, M. Hijikata, H. Luo, Y.Y. Ma, S. Kaneko, K. Shimotohno, S. Murakami, Effect of interaction between hepatitis C virus NS5A and NS5B on hepatitis C virus RNA replication with the hepatitis C virus replicon, *J. Virol.* 78 (6) (2004) 2738–2748.
- [80] Y. Shiota, H. Luo, W. Qin, S. Kaneko, T. Yamashita, K. Kobayashi, S. Murakami, Hepatitis C virus (HCV) NS5A binds RNA-dependent RNA polymerase (RdRP) NS5B and modulates RNA-dependent RNA polymerase activity, *J. Biol. Chem.* 277 (13) (2002) 11149–11155.
- [81] M. Dimitrova, I. Imbert, M.P. Kienny, C. Schuster, Protein–protein interactions between hepatitis C virus nonstructural proteins, *J. Virol.* 77 (9) (2003) 5401–5414.
- [82] D. Moradpour, V. Brass, E. Bieck, P. Friebe, R. Gosert, H.E. Blum, R. Bartenschlager, F. Penin, V. Lohmann, Membrane association of the RNA-dependent RNA polymerase is essential for hepatitis C virus RNA replication, *J. Virol.* 78 (23) (2004) 13278–13284.
- [83] J. Schmidt-Mende, E. Bieck, T. Hogle, F. Penin, C.M. Rice, H.E. Blum, D. Moradpour, Determinants for membrane association of the hepatitis C virus RNA-dependent RNA polymerase, *J. Biol. Chem.* 276 (47) (2001) 44052–44063.
- [84] H. Ago, T. Adachi, A. Yoshida, M. Yamamoto, N. Habuka, K. Yatsunami, M. Miyano, Crystal structure of the RNA-dependent RNA polymerase of hepatitis C virus, *Struct. Fold. Des.* 7 (11) (1999) 1417–1426.
- [85] A. Lesburg, M.B. Cable, E. Ferrari, Z. Hong, A.F. Mannarino, P.C. Weber, Crystal structure of the RNA-dependent RNA polymerase from hepatitis C virus reveals a fully encircled active site, *Nat. Struct. Mol. Biol.* 6 (10) (1999) 937–943.
- [86] S. Bressanelli, L. Tomei, F.A. Rey, R. De Francesco, Structural analysis of the hepatitis C virus RNA polymerase in complex with ribonucleotides, *J. Virol.* 76 (7) (2002) 3482–3492.
- [87] S.J. Butcher, J.M. Grimes, E.V. Makeyev, D.H. Bamford, D.I. Stuart, A mechanism for initiating RNA-dependent RNA polymerization, *Nature* 410 (6825) (2001) 235–240.
- [88] T.F. Baumert, S. Ito, D.T. Wong, T.J. Liang, Hepatitis C virus structural proteins assemble into viruslike particles in insect cells, *J. Virol.* 72 (5) (1998) 3827–3836.
- [89] E. Matsuo, H. Tani, C. Lim, Y. Komoda, T. Okamoto, H. Miyamoto, K. Moriishi, S. Yagi, A.H. Patel, T. Miyamura, Y. Matsuura, Characterization of HCV-like particles produced in a human hepatoma cell line by a recombinant baculovirus, *Biochem. Biophys. Res. Commun.* 340 (1) (2006) 200–208.
- [90] L.M. Lagging, K.M. Randall, J. Owens, R. Ray, Functional Role of Hepatitis C Virus Chimeric Glycoproteins in the Infectivity of Pseudotyped Virus, *J. Virol.* 72 (1998) 3539–3546.
- [91] Y. Matsuura, H. Tani, K. Suzuki, T. Kimura-Someya, R. Suzuki, H. Aizaki, K. Ishii, K. Moriishi, C.S. Robison, M.A. Whitt, T. Miyamura, Characterization of pseudotyped VSV possessing HCV envelope proteins, *Virology* 286 (2) (2001) 263–275.
- [92] B. Bartosch, J. Bukh, J.C. Meunier, C. Granier, R.E. Engle, W.C. Blackwelder, S.U. Emerson, F.L. Cosset, R.H. Purcell, In vitro assay for neutralizing antibody to hepatitis C virus: evidence for broadly conserved neutralization epitopes, *Proc. Natl. Acad. Sci. U. S. A.* 100 (24) (2003) 14199–14204.
- [93] M. Hsu, J. Zhang, M. Flint, C. Logvinoff, C. Cheng-Mayer, C.M. Rice, J.A. McKeating, Hepatitis C virus glycoproteins mediate pH-dependent cell entry of pseudotyped retroviral particles, *Proc. Natl. Acad. Sci. U. S. A.* 100 (12) (2003) 7271–7276.
- [94] T. Wakita, T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H.G. Krausslich, M. Mizokami, R. Bartenschlager, T.J. Liang, Production of infectious hepatitis C virus in tissue culture from a cloned viral genome, *Nat. Med.* 11 (7) (2005) 791–796.
- [95] J. Zhong, P. Gastaminza, G. Cheng, S. Kapadia, T. Kato, D.R. Burton, S.F. Wieland, S.L. Uprichard, T. Wakita, F.V. Chisari, Robust hepatitis C virus infection in vitro, *Proc. Natl. Acad. Sci. U. S. A.* 102 (26) (2005) 9294–9299.
- [96] D. Lindenbach, M.J. Evans, A.J. Syder, B. Wolk, T.L. Tellinghuisen, C.C. Liu, T. Maruyama, R.O. Hynes, D.R. Burton, J.A. McKeating, C.M. Rice, Complete replication of hepatitis C virus in cell culture, *Science* 309 (5734) (2005) 623–626.
- [97] P. Pileri, Y. Uematsu, S. Campagnoli, G. Galli, F. Falugi, R. Petracca, A.J. Weiner, M. Houghton, D. Rosa, G. Grandi, S. Abrignani, Binding of hepatitis C virus to CD81, *Science* 282 (1998) 938–941.
- [98] S. Levy, S.C. Todd, H.T. Maecker, CD81 (TAPA-1): a molecule involved in signal transduction and cell adhesion in the immune system, *Annu. Rev. Immunol.* 16 (1998) 89–109.
- [99] R. Petracca, F. Falugi, G. Galli, N. Norais, D. Rosa, S. Campagnoli, V. Burgio, E. Di Stasio, B. Giardina, M. Houghton, S. Abrignani, G. Grandi, Structure–function analysis of hepatitis C virus envelope-CD81 binding, *J. Virol.* 74 (10) (2000) 4824–4830.
- [100] B. Bartosch, J. Dubuisson, F.L. Cosset, Infectious hepatitis C virus pseudo-particles containing functional E1–E2 envelope protein complexes, *J. Exp. Med.* 197 (5) (2003) 633–642.
- [101] E.G. Cormier, F. Tsamis, F. Kajumo, R.J. Durso, J.P. Gardner, T. Dragic, CD81 is an entry coreceptor for hepatitis C virus, *Proc. Natl. Acad. Sci. U. S. A.* 101 (19) (2004) 7270–7274.

- [102] J. Zhang, G. Randall, A. Higginbottom, P. Monk, C.M. Rice, J.A. McKeating, CD81 is required for hepatitis C virus glycoprotein-mediated viral infection, *J. Virol.* 78 (3) (2004) 1448–1455.
- [103] B. Bartosch, A. Vitelli, C. Granier, C. Goujon, J. Dubuisson, S. Pascale, E. Scarselli, R. Cortese, A. Nicosia, F.L. Cosset, Cell entry of hepatitis C virus requires a set of co-receptors that include the CD81 tetraspanin and the SR-B1 scavenger receptor, *J. Biol. Chem.* 278 (43) (2003) 41624–41630.
- [104] A. McKeating, L.Q. Zhang, C. Logvinoff, M. Flint, J. Zhang, J. Yu, D. Butera, D.D. Ho, L.B. Dustin, C.M. Rice, P. Balfe, Diverse hepatitis C virus glycoproteins mediate viral infection in a CD81-dependent manner, *J. Virol.* 78 (16) (2004) 8496–8505.
- [105] D. Lavillette, A.W. Tarr, C. Voisset, P. Donot, B. Bartosch, C. Bain, A.H. Patel, J. Dubuisson, J.K. Ball, F.L. Cosset, Characterization of host-range and cell entry properties of the major genotypes and subtypes of hepatitis C virus, *Hepatology* 41 (2) (2005) 265–274.
- [106] E. Scarselli, H. Ansuini, R. Cerino, R.M. Roccasecca, S. Acali, G. Filocamo, C. Traboni, A. Nicosia, R. Cortese, A. Vitelli, The human scavenger receptor class B type I is a novel candidate receptor for the hepatitis C virus, *EMBO J.* 21 (19) (2002) 5017–5025.
- [107] D. Rhoads, L. Brissette, The role of scavenger receptor class B type I (SR-B1) in lipid trafficking. defining the rules for lipid traders, *Int. J. Biochem. Cell Biol.* 36 (1) (2004) 39–77.
- [108] E. Yamada, M. Montoya, C.G. Schuettler, T.P. Hickling, A.W. Tarr, A. Vitelli, J. Dubuisson, A.H. Patel, J.K. Ball, P. Borrow, Analysis of the binding of hepatitis C virus genotype 1a and 1b E2 glycoproteins to peripheral blood mononuclear cell subsets, *J. Gen. Virol.* 86 (Pt 9) (2005) 2507–2512.
- [109] S. Acton, A. Rigotti, K.T. Landschulz, S. Xu, H.H. Hobbs, M. Krieger, Identification of scavenger receptor SR-B1 as a high density lipoprotein receptor, *Science* 271 (5248) (1996) 518–520.
- [110] K.T. Landschulz, R.K. Pathak, A. Rigotti, M. Krieger, H.H. Hobbs, Regulation of scavenger receptor, class B, type I, a high density lipoprotein receptor, in liver and steroidogenic tissues of the rat, *J. Clin. Invest.* 98 (4) (1996) 984–995.
- [111] H. Barth, R. Cerino, M. Arcuri, M. Hoffmann, P. Schurmann, M.I. Adah, B. Gissler, X. Zhao, V. Ghisetti, B. Lavezzo, H.E. Blum, F. von Weizsacker, A. Vitelli, E. Scarselli, T.F. Baumert, Scavenger receptor class B type I and hepatitis C virus infection of primary tupaia hepatocytes, *J. Virol.* 79 (9) (2005) 5774–5785.
- [112] B. Bartosch, G. Verney, M. Dreux, P. Donot, Y. Morice, F. Penin, J.M. Pawlotsky, D. Lavillette, F.L. Cosset, An interplay between hypervariable region 1 of the hepatitis C virus E2 glycoprotein, the scavenger receptor BI, and high-density lipoprotein promotes both enhancement of infection and protection against neutralizing antibodies, *J. Virol.* 79 (13) (2005) 8217–8229.
- [113] D.D. Lavillette, Y. Morice, G. Germanidis, P. Donot, A. Soulier, E. Pagkalos, G. Sakellariou, L. Intrator, B. Bartosch, J.M. Pawlotsky, F.L. Cosset, Human serum facilitates hepatitis C virus infection, and neutralizing responses inversely correlate with viral replication kinetics at the acute phase of hepatitis C virus infection, *J. Virol.* 79 (10) (2005) 6023–6034.
- [114] J.C. Meunier, R.E. Engle, K. Faulk, M. Zhao, B. Bartosch, H. Alter, S.U. Emerson, F.L. Cosset, R.H. Purcell, J. Bukh, Evidence for cross-genotype neutralization of hepatitis C virus pseudo-particles and enhancement of infectivity by apolipoprotein C1, *Proc. Natl. Acad. Sci. U. S. A.* 102 (12) (2005) 4560–4565.
- [115] C. Voisset, N. Callens, E. Blanchard, A. Op De Beeck, J. Dubuisson, N. Vu-Dac, High density lipoproteins facilitate hepatitis C virus entry through the scavenger receptor class B type I, *J. Biol. Chem.* 280 (9) (2005) 7793–7799.
- [116] J.P. Gardner, R.J. Durso, R.R. Arrigale, G.P. Donovan, P.J. Maddon, T. Dragic, W.C. Olson, L-SIGN (CD 209L) is a liver-specific capture receptor for hepatitis C virus, *Proc. Natl. Acad. Sci. U. S. A.* 100 (8) (2003) 4498–4503.
- [117] Y. Lozach, H. Lortat-Jacob, A. de Lacroix de Lavalette, I. Staropoli, S. Foung, A. Amara, C. Houles, F. Fieschi, O. Schwartz, J.L. Virelizier, F. Arenzana-Seisdedos, R. Altmeyer, DC-SIGN and L-SIGN are high affinity binding receptors for hepatitis C virus glycoprotein E2, *J. Biol. Chem.* 278 (22) (2003) 20358–20366.
- [118] S. Pohlmann, J. Zhang, F. Baribaud, Z. Chen, G.J. Leslie, G. Lin, A. Granelli-Piperno, R.W. Doms, C.M. Rice, J.A. McKeating, Hepatitis C virus glycoproteins interact with DC-SIGN and DC-SIGNR, *J. Virol.* 77 (7) (2003) 4070–4080.
- [119] B. Saunier, M. Triyatni, L. Ulianich, P. Maruvada, P. Yen, L.D. Kohn, Role of the asialoglycoprotein receptor in binding and entry of hepatitis C virus structural proteins in cultured human hepatocytes, *J. Virol.* 77 (1) (2003) 546–559.
- [120] E.G. Cormier, R.J. Durso, F. Tsamis, L. Boussemart, C. Manix, W.C. Olson, J.P. Gardner, T. Dragic, L-SIGN (CD209L) and DC-SIGN (CD209) mediate transinfection of liver cells by hepatitis C virus, *Proc. Natl. Acad. Sci. U. S. A.* 101 (39) (2004) 14067–14072.
- [121] Y. Lozach, A. Amara, B. Bartosch, J.L. Virelizier, F. Arenzana-Seisdedos, F.L. Cosset, R. Altmeyer, C-type lectins L-SIGN and DC-SIGN capture and transmit infectious hepatitis C virus pseudotype particles, *J. Biol. Chem.* 279 (31) (2004) 32035–32045.
- [122] V. Agnello, G. Abel, M. Elfahal, G.B. Knight, Q.X. Zhang, Hepatitis C virus and other flaviviridae viruses enter cells via low density lipoprotein receptor, *Proc. Natl. Acad. Sci. U. S. A.* 96 (22) (1999) 12766–12771.
- [123] M.J. Evans, T. von Hahn, D.M. Tschernie, A.J. Syder, M. Panis, B. Wolk, T. Hatzioannou, J.A. McKeating, P.D. Bieniasz, C.M. Rice, Claudin-1 is a hepatitis C virus co-receptor required for a late step in entry, *Nature* 446 (7137) (2007) 801–805.
- [124] R. Rijnbrand, P. Bredenbeek, T. van der Straaten, L. Whetter, G. Inchauspe, S. Lemon, W. Spaan, Almost the entire 5' non-translated region of hepatitis C virus is required for cap-independent translation, *FEBS Lett.* 365 (2-3) (1995) 115–119.
- [125] B.J. Yoo, R.R. Spaete, A.P. Geballe, M. Selby, M. Houghton, J.H. Han, 5' end-dependent translation initiation of hepatitis C viral RNA and the presence of putative positive and negative translational control elements within the 5' untranslated region, *Virology* 191 (2) (1992) 889–899.
- [126] U. Hellen, T.V. Pestova, Translation of hepatitis C virus RNA, *J. Viral Hepatitis* 6 (2) (1999) 79–87.
- [127] H.H. Lu, E. Wimmer, Poliovirus chimeras replicating under the translational control of genetic elements of hepatitis C virus reveal unusual properties of the internal ribosomal entry site of hepatitis C virus, *Proc. Natl. Acad. Sci. U. S. A.* 93 (4) (1996) 1412–1417.
- [128] J.E. Reynolds, A. Kaminski, H.J. Kettinen, K. Grace, B.E. Clarke, A.R. Carroll, D.J. Rowlands, R.J. Jackson, Unique features of internal initiation of hepatitis C virus RNA translation, *EMBO J.* 14 (23) (1995) 6010–6020.
- [129] M. Spahn, J.S. Kieft, R.A. Grassucci, P.A. Penczek, K. Zhou, J.A. Doudna, J. Frank, Hepatitis C virus IRES RNA-induced changes in the conformation of the 40s ribosomal subunit, *Science* 291 (5510) (2001) 1959–1962.
- [130] A. Otto, J.D. Puglisi, The pathway of HCV IRES-mediated translation initiation, *Cell* 119 (3) (2004) 369–380.
- [131] N. Ali, G.J. Pruijn, D.J. Kenan, J.D. Keene, A. Siddiqui, Human La antigen is required for the hepatitis C virus internal ribosome entry site-mediated translation, *J. Biol. Chem.* 275 (36) (2000) 27531–27540.
- [132] N. Ali, A. Siddiqui, The La antigen binds 5' noncoding region of the hepatitis C virus RNA in the context of the initiator AUG codon and stimulates internal ribosome entry site-mediated translation, *Proc. Natl. Acad. Sci. U. S. A.* 94 (6) (1997) 2249–2254.
- [133] T. Isoyama, N. Kamoshita, K. Yasui, A. Iwai, K. Shiroki, H. Toyoda, A. Yamada, Y. Takasaki, A. Nomoto, Lower concentration of La protein required for internal ribosome entry on hepatitis C virus RNA than on poliovirus RNA, *J. Gen. Virol.* 80 (Pt 9) (1999) 2319–2327.
- [134] B. Hahm, Y.K. Kim, J.H. Kim, T.Y. Kim, S.K. Jang, Heterogeneous nuclear ribonucleoprotein L interacts with the 3' border of the internal ribosomal entry site of hepatitis C virus, *J. Virol.* 72 (11) (1998) 8782–8788.
- [135] I.S. Fukushi, M. Okada, T. Kageyama, F.B. Hoshino, K. Nagai, K. Katayama, Interaction of poly(rC)-binding protein 2 with the 5'-terminal stem loop of the hepatitis C-virus genome, *Virus Res.* 73 (1) (2001) 67–79.



- [136] A. Anwar, N. Ali, R. Tanveer, A. Siddiqui, Demonstration of functional requirement of polypyrimidine tract-binding protein by SELEX RNA during hepatitis C virus internal ribosome entry site-mediated translation initiation, *J. Biol. Chem.* 275 (44) (2000) 34231–34235.
- [137] T. Shimoike, S. Mimori, H. Tani, Y. Matsuura, T. Miyamura, Interaction of hepatitis C virus core protein with viral sense RNA and suppression of its translation, *J. Virol.* 73 (12) (1999) 9718–9725.
- [138] Y. Tanaka, T. Shimoike, K. Ishii, R. Suzuki, T. Suzuki, H. Ushijima, Y. Matsuura, T. Miyamura, Selective binding of hepatitis C virus core protein to synthetic oligonucleotides corresponding to the 5' untranslated region of the viral genome, *Virology* 270 (1) (2000) 229–236.
- [139] T. Shimoike, C. Koyama, K. Murakami, R. Suzuki, Y. Matsuura, T. Miyamura, T. Suzuki, Down-regulation of the internal ribosome entry site (IRES)-mediated translation of the hepatitis C virus: critical role of binding of the stem-loop III domain of IRES and the viral core protein, *Virology* 345 (2) (2006) 434–445.
- [140] J. Zhang, O. Yamada, H. Yoshida, T. Iwai, H. Araki, Autogenous translational inhibition of core protein: implication for switch from translation to RNA replication in hepatitis C virus, *Virology* 293 (1) (2002) 141–150.
- [141] D. Li, S.T. Takyar, W.B. Lott, E.J. Gowans, Amino acids 1–20 of the hepatitis C virus (HCV) core protein specifically inhibit HCV IRES-dependent translation in HepG2 cells, and inhibit both HCV IRES- and cap-dependent translation in HuH7 and CV-1 cells, *J. Gen. Virol.* 84 (Pt 4) (2003) 815–825.
- [142] T.H. Wang, R.C. Rijnbrand, S.M. Lemon, Core protein-coding sequence, but not core protein, modulates the efficiency of cap-independent translation directed by the internal ribosome entry site of hepatitis C virus, *J. Virol.* 74 (23) (2000) 11347–11358.
- [143] J.L. Walewski, T.R. Keller, D.D. Stump, A.D. Branch, Evidence for a new hepatitis C virus antigen encoded in an overlapping reading frame, *RNA* 7 (5) (2001) 710–721.
- [144] Z. Xu, J. Choi, T.S. Yen, W. Lu, A. Strohecker, S. Govindarajan, D. Chien, M.J. Selby, J. Ou, Synthesis of a novel hepatitis C virus protein by ribosomal frameshift, *EMBO J.* 20 (14) (2001) 3840–3848.
- [145] A. Varaklioti, N. Vassilaki, U. Georgopoulou, P. Mavromara, Alternate translation occurs within the core coding region of the hepatitis C viral genome, *J. Biol. Chem.* 277 (20) (2002) 17713–17721.
- [146] P. Hüseyin, H. Langen, J. Mous, H. Jacobsen, Hepatitis C virus core protein: carboxy-terminal boundaries of two processed species suggest cleavage by a signal peptide peptidase, *Virology* 224 (1996) 93–104.
- [147] J. McLauchlan, M.K. Lemberg, G. Hope, B. Martoglio, Intramembrane proteolysis promotes trafficking of hepatitis C virus core protein to lipid droplets, *EMBO J.* 21 (15) (2002) 3980–3988.
- [148] M.K. Lemberg, B. Martoglio, Requirements for signal peptide peptidase-catalyzed intramembrane proteolysis, *Mol. Cell* 10 (4) (2002) 735–744.
- [149] K. Okamoto, K. Moriishi, T. Miyamura, Y. Matsuura, Intramembrane proteolysis and endoplasmic reticulum retention of hepatitis C virus core protein, *J. Virol.* 78 (12) (2004) 6370–6380.
- [150] A. Weihofen, K. Binns, M.K. Lemberg, K. Ashman, B. Martoglio, Identification of signal peptide peptidase, a presenilin-type aspartic protease, *Science* 296 (5576) (2002) 2215–2218.
- [151] R. Bartenschlager, L. Ahlborn-Laake, J. Mous, H. Jacobsen, Kinetic and structural analyses of hepatitis C virus polyprotein processing, *J. Virol.* 68 (1994) 5045–5055.
- [152] C. Failla, L. Tomei, R. De Francesco, An amino-terminal domain of the hepatitis C virus NS3 protease is essential for interaction with NS4A, *J. Virol.* 69 (3) (1995) 1769–1777.
- [153] C. Lin, B.M. Pragai, A. Grakoui, J. Xu, C.M. Rice, Hepatitis C virus NS3 serine proteinase: *trans*-cleavage requirements and processing kinetics, *J. Virol.* 68 (1994) 8147–8157.
- [154] Y. Tanji, M. Hijikata, Y. Hirowatari, K. Shimotohno, Hepatitis C virus polyprotein processing: kinetics and mutagenic analysis of serine proteinase-dependent cleavage, *J. Virol.* 68 (1994) 8418–8422.
- [155] H. Aizaki, K.J. Lee, V.M. Sung, H. Ishiko, M.M. Lai, Characterization of the hepatitis C virus RNA replication complex associated with lipid rafts, *Virology* 324 (2) (2004) 450–461.
- [156] N. Ali, K.D. Tardif, A. Siddiqui, Cell-free replication of the hepatitis C virus subgenomic replicon, *J. Virol.* 76 (23) (2002) 12001–12007.
- [157] V.C. Lai, S. Dempsey, J.Y. Lau, Z. Hong, W. Zhong, In vitro RNA replication directed by replicase complexes isolated from the subgenomic replicon cells of hepatitis C virus, *J. Virol.* 77 (3) (2003) 2295–2300.
- [158] Y. Miyanari, M. Hijikata, M. Yamaji, M. Hosaka, H. Takahashi, K. Shimotohno, Hepatitis C virus non-structural proteins in the probable membranous compartment function in viral genome replication, *J. Biol. Chem.* 278 (50) (2003) 50301–50308.
- [159] R.W. Hardy, J. Marcotrigiano, K.J. Blight, J.E. Majors, C.M. Rice, Hepatitis C virus RNA synthesis in a cell-free system isolated from replicon-containing hepatoma cells, *J. Virol.* 77 (3) (2003) 2029–2037.
- [160] S.T. Shi, K.J. Lee, H. Aizaki, S.B. Hwang, M.M. Lai, Hepatitis C virus RNA replication occurs on a detergent-resistant membrane that cofractionates with caveolin-2, *J. Virol.* 77 (7) (2003) 4160–4168.
- [161] U. Pfeifer, R. Thomssen, K. Legler, U. Bottcher, W. Gerlich, E. Weinmann, O. Klinge, Experimental non-A, non-B hepatitis: four types of cytoplasmic alteration in hepatocytes of infected chimpanzees, *Virchows Arch., B Cell Pathol. Incl. Mol. Pathol.* 33 (3) (1980) 233–243.
- [162] G. Mottola, G. Cardinali, A. Ceccacci, C. Trozzi, L. Bartholomew, M.R. Torrisi, E. Pedrazzini, S. Bonatti, G. Migliaccio, Hepatitis C virus nonstructural proteins are localized in a modified endoplasmic reticulum of cells expressing viral subgenomic replicons, *Virology* 293 (1) (2002) 31–43.
- [163] T. Pietschmann, V. Lohmann, G. Rutter, K. Kurpanek, R. Bartenschlager, Characterization of cell lines carrying self-replicating hepatitis C virus RNAs, *J. Virol.* 75 (3) (2001) 1252–1264.
- [164] G. Waris, S. Sarker, A. Siddiqui, Two-step affinity purification of the hepatitis C virus ribonucleoprotein complex, *RNA* 10 (2) (2004) 321–329.
- [165] T. Umehara, M. Sudoh, F. Yasui, C. Matsuda, Y. Hayashi, K. Chayama, M. Kohara, Serine palmitoyltransferase inhibitor suppresses HCV replication in a mouse model, *Biochem. Biophys. Res. Commun.* 346 (1) (2006) 67–73.
- [166] H. Sakamoto, K. Okamoto, M. Aoki, H. Kato, A. Katsume, A. Ohta, T. Tsukuda, N. Shimma, Y. Aoki, M. Arisawa, M. Kohara, M. Sudoh, Host sphingolipid biosynthesis as a target for hepatitis C virus therapy, *Nat. Chem. Biol.* 1 (6) (2005) 333–337.
- [167] I. Hamamoto, Y. Nishimura, T. Okamoto, H. Aizaki, M. Liu, Y. Mori, T. Abe, T. Suzuki, M.M. Lai, T. Miyamura, K. Moriishi, Y. Matsuura, Human VAP-B is involved in hepatitis C virus replication through interaction with NSSA and NS5B, *J. Virol.* 79 (21) (2005) 13473–13482.
- [168] T. Okamoto, Y. Nishimura, T. Ichimura, K. Suzuki, T. Miyamura, T. Suzuki, K. Moriishi, Y. Matsuura, Hepatitis C virus RNA replication is regulated by FKBP8 and Hsp90, *EMBO J.* 25 (20) (2006) 5015–5025.
- [169] J. Ye, C. Wang, R. Sumpter Jr., M.S. Brown, J.L. Goldstein, M. Gale Jr., Disruption of hepatitis C virus RNA replication through inhibition of host protein geranylgeranylation, *Proc. Natl. Acad. Sci. U. S. A.* 100 (26) (2003) 15865–15870.
- [170] S.B. Kapadia, F.V. Chisari, Hepatitis C virus RNA replication is regulated by host geranylgeranylation and fatty acids, *Proc. Natl. Acad. Sci. U. S. A.* 102 (7) (2005) 2561–2566.
- [171] T.L. Tan, H. Nakao, Y. He, S. Vijaysri, P. Neddermann, B.L. Jacobs, B.J. Mayer, M.G. Katze, NS5A, a nonstructural protein of hepatitis C virus, binds growth factor receptor-bound protein 2 adaptor protein in a Src homology 3 domain/ligand-dependent manner and perturbs mitogenic signaling, *Proc. Natl. Acad. Sci. U. S. A.* 96 (10) (1999) 5533–5538.
- [172] K. Ghosh, M. Majumder, R. Steele, P. Yaciuk, J. Chrivia, R. Ray, R.B. Ray, Hepatitis C virus NS5A protein modulates transcription through a novel cellular transcription factor SRCAP, *J. Biol. Chem.* 275 (10) (2000) 7184–7188.
- [173] K.M. Chung, J. Lee, J.E. Kim, O.K. Song, S. Cho, J. Lim, M. Seedorf, B. Hahn, S.K. Jang, Nonstructural protein 5A of hepatitis C virus inhibits the function of karyopherin beta3, *J. Virol.* 74 (11) (2000) 5233–5241.
- [174] T. Burckstummer, M. Kriegs, J. Lupberger, E.K. Pauli, S. Schmittl, E. Hildt, Raf-1 kinase associates with Hepatitis C virus NS5A and regulates viral replication, *FEBS Lett.* 580 (2) (2006) 575–580.
- [175] M. Ikeda, K. Abe, M. Yamada, H. Dansako, K. Naka, N. Kato, Different anti-HCV profiles of statins and their potential for combination therapy with interferon, *Hepatology* 44 (1) (2006) 117–125.

- [176] K. Watahi, N. Ishii, M. Hijikata, D. Inoue, T. Murata, Y. Miyanari, K. Shimotohno, Cyclophilin B is a functional regulator of hepatitis C virus RNA polymerase, *Mol. Cell* 19 (1) (2005) 111–122.
- [177] P.Y. Goh, Y.J. Tan, S.P. Lim, Y.H. Tan, S.G. Lim, F. Fuller-Pace, W. Hong, Cellular RNA helicase p68 relocalization and interaction with the hepatitis C virus (HCV) NS5B protein and the potential role of p68 in HCV RNA replication, *J. Virol.* 78 (10) (2004) 5288–5298.
- [178] M. Hirano, S. Kaneko, T. Yamashita, H. Luo, W. Qin, Y. Shirota, T. Nomura, K. Kobayashi, S. Murakami, Direct interaction between nucleolin and hepatitis C virus NS5B, *J. Biol. Chem.* 278 (7) (2003) 5109–5115.
- [179] T. Shimakami, M. Honda, T. Kusakawa, T. Murata, K. Shimotohno, S. Kaneko, S. Murakami, Effect of hepatitis C virus (HCV) NS5B-nucleolin interaction on HCV replication with HCV subgenomic replicon, *J. Virol.* 80 (7) (2006) 3332–3340.
- [180] S. Kim, S.K. Seol, O.K. Song, J.H. Park, S.K. Jang, An RNA-binding protein, hnRNP A1, and a scaffold protein, Septin 6, facilitate hepatitis C virus replication, *J. Virol.* 81 (8) (2007) 3852–3865.
- [181] N. Ali, A. Siddiqui, Interaction of polypyrimidine tract-binding protein with the 5' noncoding region of the hepatitis C virus RNA genome and its functional requirement in internal initiation of translation, *J. Virol.* 69 (10) (1995) 6367–6375.
- [182] T. Ito, M.M. Lai, Determination of the secondary structure of and cellular protein binding to the 3'-untranslated region of the hepatitis C virus RNA genome, *J. Virol.* 71 (11) (1997) 8698–8706.
- [183] K. Murakami, M. Abe, T. Kageyama, N. Kamoshita, A. Nomoto, Down-regulation of translation driven by hepatitis C virus internal ribosomal entry site by the 3' untranslated region of RNA, *Arch. Virol.* 146 (4) (2001) 729–741.
- [184] K. Tsuchihara, T. Tanaka, M. Hijikata, S. Kuge, H. Toyoda, A. Nomoto, N. Yamamoto, K. Shimotohno, Specific interaction of polypyrimidine tract-binding protein with the extreme 3'-terminal structure of the hepatitis C virus genome, the 3'X, *J. Virol.* 71 (9) (1997) 6720–6726.
- [185] H. Aizaki, K.S. Choi, M. Liu, Y.J. Li, M.M. Lai, Polypyrimidine-tract-binding protein is a component of the HCV RNA replication complex and necessary for RNA synthesis, *J. Biomed. Sci.* 13 (4) (2006) 469–480.
- [186] M. Kaito, S. Watanabe, K. Tsukiyama-Kohara, K. Yamaguchi, Y. Kobayashi, M. Konishi, M. Yokoi, S. Ishida, S. Suzuki, M. Kohara, Hepatitis C virus particle detected by immunoelectron microscopic study, *J. Gen. Virol.* 75 (1994) 1755–1760.
- [187] Y.K. Shimizu, S.M. Feinstone, M. Kohara, R.H. Purcell, H. Yoshikura, Hepatitis C virus: detection of intracellular virus particles by electron microscopy, *Hepatology* 23 (1996) 205–209.
- [188] P. Maillard, K. Krawczynski, J. Nitkiewicz, C. Bronnert, M. Sidorkiewicz, P. Gounon, J. Dubuisson, G. Faure, R. Cranic, A. Budkowska, Nonenveloped nucleocapsids of hepatitis C virus in the serum of infected patients, *J. Virol.* 75 (17) (2001) 8240–8250.
- [189] P. Andre, F. Komurian-Pradel, S. Deforges, M. Perret, J.L. Berland, M. Sodoyer, S. Pol, C. Brechot, G. Paranhos-Baccala, V. Lotteau, Characterization of low- and very-low-density hepatitis C virus RNA-containing particles, *J. Virol.* 76 (14) (2002) 6919–6928.
- [190] M. Kunkel, M. Lorinczi, R. Rijnbrand, S.M. Lemon, S.J. Watowich, Self-assembly of nucleocapsid-like particles from recombinant hepatitis C virus core protein, *J. Virol.* 75 (5) (2001) 2119–2129.
- [191] L.J. Lorenzo, S. Duenas-Carrera, V. Falcon, N. Acosta-Rivero, E. Gonzalez, M.C. de la Rosa, I. Menendez, J. Morales, Assembly of truncated HCV core antigen into virus-like particles in *Escherichia coli*, *Biochem. Biophys. Res. Commun.* 281 (4) (2001) 962–965.
- [192] N. Acosta-Rivero, J.C. Aguilar, A. Musacchio, V. Falcon, A. Vina, M.C. de la Rosa, J. Morales, Characterization of the HCV core virus-like particles produced in the methylotrophic yeast *Pichia pastoris*, *Biochem. Biophys. Res. Commun.* 287 (1) (2001) 122–125.
- [193] E. Blanchard, C. Hourieux, D. Brand, M. Ait-Goughoulte, A. Moreau, S. Trassard, P.Y. Sizaret, F. Dubois, P. Roingard, Hepatitis C virus-like particle budding: role of the core protein and importance of its Asp111, *J. Virol.* 77 (18) (2003) 10131–10138.
- [194] N. Majeau, V. Gagne, A. Boivin, M. Bolduc, J.A. Majeau, D. Ouellet, D. Leclerc, The N-terminal half of the core protein of hepatitis C virus is sufficient for nucleocapsid formation, *J. Gen. Virol.* 85 (Pt 4) (2004) 971–981.
- [195] C. Klein, S.J. Polyak, J.R. Lingappa, Unique features of hepatitis C virus capsid formation revealed by de novo cell-free assembly, *J. Virol.* 78 (17) (2004) 9257–9269.
- [196] M. Matsumoto, S.B. Hwang, K.S. Jeng, N. Zhu, M.M. Lai, Homotypic interaction and multimerization of hepatitis C virus core protein, *Virology* 218 (1) (1996) 43–51.
- [197] O. Nolandt, V. Kern, H. Muller, E. Pfaff, L. Theilmann, R. Welker, H.G. Krausslich, Analysis of hepatitis C virus core protein interaction domains, *J. Gen. Virol.* 78 (Pt 6) (1997) 1331–1340.
- [198] S. Yan, M.H. Tam, W.J. Syu, Self-association of the C-terminal domain of the hepatitis-C virus core protein, *Eur. J. Biochem.* 258 (1) (1998) 100–106.
- [199] H.J. Ezelle, D. Markovic, G.N. Barber, Generation of hepatitis C virus-like particles by use of a recombinant vesicular stomatitis virus vector, *J. Virol.* 76 (23) (2002) 12325–12334.
- [200] K. Murakami, K. Ishii, Y. Ishihara, S. Yoshizaki, K. Tanaka, Y. Gotoh, H. Aizaki, M. Kohara, H. Yoshioka, Y. Mori, N. Manabe, I. Shoji, T. Sata, R. Bartenschlager, Y. Matsuura, T. Miyamura, T. Suzuki, Production of infectious hepatitis C virus particles in three-dimensional cultures of the cell line carrying the genome-length dicistronic viral RNA of genotype 1b, *Virology* 351 (2) (2006) 381–392.
- [201] S.-Y. Lo, M.J. Selby, J.-H. Ou, Interaction between hepatitis C virus core protein and E1 envelope protein, *J. Virol.* 70 (1996) 5177–5182.
- [202] H.C. Ma, C.H. Ke, T.Y. Hsieh, S.Y. Lo, The first hydrophobic domain of the hepatitis C virus E1 protein is important for interaction with the capsid protein, *J. Gen. Virol.* 83 (Pt 12) (2002) 3085–3092.
- [203] A. Serafino, M.B. Valli, F. Andreola, A. Crema, G. Ravagnan, L. Bertolini, G. Carloni, Suggested role of the Golgi apparatus and endoplasmic reticulum for crucial sites of hepatitis C virus replication in human lymphoblastoid cells infected in vitro, *J. Med. Virol.* 70 (1) (2003) 31–41.
- [204] K. Sato, H. Okamoto, S. Aihara, Y. Hoshi, T. Tanaka, S. Mishiro, Demonstration of sugar moiety on the surface of hepatitis C virions recovered from the circulation of infected humans, *Virology* 196 (1993) 354–357.