

Imai K, Ogai Y, Nishizawa D, Kasai S, Ikeda K, Koga H.

A novel SNP detection technique utilizing a multiple primer extension (MPEX) on a phospholipid polymer-coated surface. *Mol Biosyst*,3,547-553, 2007

Mouri, A., Noda, Y., Noda, A., Nakamura, T., Tokura, T., Yura, Y., Nitta, A., Furukawa, H. and Nabeshima, T.

Involvement of a dysfunctional dopamine-D1/NMDA-NR1 and CaMK II pathway in the impairment of latent learning in a model of schizophrenia induced by phencyclidine. *Mol. Pharmacol.*,71,1598-1609, 2007

Cen XB, Nitta A., Ibi D, Zhao Y., Niwa M, Taguchi K., Hamada M, Ito Yoshihisa., Ito Yasumoto, Wang L., Nabeshima T.,:

Identification of Piccolo as a regulator of behavioral plasticity and dopamine transporter internalization : Piccolo as a regulator of behavioral plasticity and DAT internalization. *Mol. Psychiat.*, in press.

Shin EJ, Nah SY, Chae JS, Bing G., Shin SW, Yen T., Baek IH, Kim WK, Maurice, T., Nabeshima T. and Kim HC.:

Dextromethorphan attenuates trimethyltin-induced neurotoxicity via $\sigma 1$ receptor activation in rats. *Neurochem. Int.*, 50,791-799,2007

Mouri, A., Noda, Y., Enomoto, T. and Nabeshima, T.:

Phencyclidine animal models of schizophrenia: approaches from abnormality of glutamatergic neurotransmission and neurodevelopment. *Neurochem. Int.*, 51, 173-184, 2007

Fujihira T, Kanematsu S, Umino A, et al

Selective increase in the extracellular D-serine contents by D-cycloserine in the rat medial frontal cortex. *Neurochem. Int.*,51,233-236, 2007

Yamamoto H, Kamegaya E, Hagino Y, Imai K, Fujikawa A, Tamura K, Enokiya T, Yamamoto T, Takeshima T, Koga H, Uhl GR, Ikeda K, Sora I.

Genetic deletion of vesicular monoamine transporter-2 (VMAT2) reduces dopamine transporter activity in mesencephalic neurons in primary culture. *Neurochem Int*,51,237-244, 2007

Wang, D., Noda, Y., Zhou, Y., Nitta, A., Furukawa H. and Nabeshima, T.

Synergistic effect of galantamine with risperidone on impairment of social interaction in phencyclidine-treated mice as a schizophrenic animal model. *Neuropharmacology*, 52, 1179-1187, 2007

- Wang, D., Noda, Y., Zhou, Y., Nitta, A., Furukawa, H. and Nabeshima, T.:
Synergistic effect of combined treatment with risperidone and galantamine on phencyclidine-induced impairment of latent visuospatial learning and memory: role of nAChR activation-dependent increase of dopamine D1 receptor-mediated neurotransmission. *Neuropharmacology*, 53, 379-389, 2007
- Hashimoto K, Fujita Y, Iyo M.
Phencyclidine-induced cognitive deficits in mice are improved by subsequent subchronic administration of fluvoxamine: role of sigma-1 receptors. *Neuropsychopharmacology*,32(3),514-521, 2007
- Kaneko Y, Kashiwa A, Ito T, et al
Selective serotonin reuptake inhibitors, fluoxetine and paroxetine, attenuate the expression of the established behavioral sensitization induced by methamphetamine. *Neuropsychopharmacology*, 32, 658-664, 2007
- Wang, D., Noda, Y., Zhou, Y., Mouri, A., Mizoguchi, H., Nitta, A., Chen, W. and Nabeshima, T
The allosteric potentiation of nicotinic acetylcholine receptors by galantamine ameliorates the cognitive dysfunction in beta amyloid25-35 i.c.v.-injected mice: involvement of dopaminergic systems. *Neuropsychopharmacology*,32,1261-1271, 2007
- Zhang L, Shirayama Y, Iyo M, Hashimoto K.
Minocycline attenuates hyperlocomotion and prepulse inhibition deficits in mice after administration of the NMDA receptor antagonist dizocilpine. *Neuropsychopharmacology*,32 (9),2001-2010, 2007
- Harburg, GC., Hall, FS., Harrist, AV., Sora, I., Uhl, GR. and Eisch, AJ.
Knockout of the mu opioid receptor enhances the survival of adult-generated hippocampal granule cell neurons. *Neuroscience*,144(1),77-87, 2007
- Yan, Y., Yamada, K., Mizoguchi, H., Noda, Y., Nagai, T., Nitta, A. and Nabeshima, T.
Reinforcing effects of morphine are reduced in tissue plasminogen activator (tPA)-knockout mice. *Neuroscience*,146,50-59, 2007
- Numachi, Y., Shen, H., Yoshida, S., Fujiyama, K., Toda, S., Matsuoka, H., Sora, I. and Sato, M.
Methamphetamine alters expression of DNA methyltransferase 1 mRNA in rat brain. *Neurosci. Lett.*, 414,213–217, 2007

Numachi Y, Yoshida S, Yamashita M, et al

Altered EphA5 mRNA expression in rat brain with a single methamphetamine treatment.
Neurosci. Lett.,424,116-121, 2007

Waga C, Ikeda K, Iwahashi K

The relationship between alcoholism and DBI gene polymorphism in Japanese -genotyping of the +529A/T in DBI gene polymorphism based on PCR-. Nihon Arukoru Yakubutsu Igakkai Zasshi,42(6),629-634, 2007

Hritcu, L., Clicinschi, M. and Nabeshima, T.: . ., (2007)

Brain serotonin depletion impairs short-term memory, but not long-term memory in rats
Physiol. Behav,91, 652-657, 2007

Shimizu E, Watanabe H, Kojima T, Hagiwara H, Fujisaki M, Miyatake R, Hashimoto K, Iyo M.

Combined intoxication with methylone and 5-MeO-MIPT. Prog Neuropsychopharmacol Biol Psychiatry.,31(1),288-291., 2007

Shimizu E, Imai M, Fujisaki M, Shinoda N, Handa S, Watanabe H, Nakazato M, Hashimoto K, Iyo M.

Maintenance electroconvulsive therapy (ECT) for treatment-resistant disorganized schizophrenia.
Prog Neuropsychopharmacol Biol Psychiatry,31(2),571-573, 2007

Okamura N, Hashimoto K, Iyo M, Shimizu E,

Dempfle A, Friedel S, Reinscheid RK. Gender-specific association of a functional coding polymorphism in the neuropeptide S receptor gene with panic disorder but not with schizophrenia or attention-deficit/hyperactivity disorder. Prog Neuropsychopharmacol Biol Psychiatry.,31(7),1444-1448, 2007

Fujita Y, Ishima T, Kunitachi S, Hagiwara H, Zhang L, Iyo M, Hashimoto K.

Phencyclidine-induced cognitive deficits in mice are improved by subsequent subchronic administration of the antibiotic drug minocycline. Prog Neuropsychopharmacol Biol Psychiatry, in press.

Hashimoto T, Shimizu E, Koike K, Orita Y, Suzuki T, Kanahara N, Fukami G, Miyatake R, Shinoda N, Fujisaki M, Shirayama Y, Hashimoto K, Iyo M.

Deficits in auditory P50 inhibition in obsessive-compulsive disorder. Prog Neuropsychopharmacol Biol Psychiatry., in press

Iyo M, Shirayama Y, Watanabe H, Fujisaki M, Miyatake R, Fukami G, Akihiro Shiina A, Nakazato M, Shiraishi T, Ookami T, Hashimoto K

Fluvoxamine as a sigma-1 receptor agonist improved cognitive impairments in a patient with schizophrenia Prog Neuropsychopharmacol Biol Psychiatry., in press

Takeuchi T, Furuta K, Hirasawa T, Masaki H, Yukizane T, Atsuta H, Nishikawa T.

Perospirone in the treatment of patients with delirium. Psychiatry Clin Neurosci.,61,67-70, 2007

Arai, M., Obata, N., Kockelkorn, TT., Yamada, K., Toyota, T., Haga, S., Yoshida, Y., Ujike, H., Sora, I., Ikeda, K., Yoshikawa, T. and Itokawa, M.

Lack of association between polymorphisms in the 5' upstream region of the DISC1 gene and mood disorders. Psychiatric Genetics.,17(6),357, 2007

Fukushima S, Shen H, Ikeda K, Hata H, Ohara A, Ohmi K, Kobayashi H, Numachi Y, Hall F, Uhl GR, Sora I.

Methamphetamine-induced locomotor activity and sensitization in dopamine transporter and vesicular monoamine transporter 2 double mutant mice. Psychopharmacology,193(1),55-62, 2007

Nagai, T., Takuma, K., Dohniwa, M., Ibi, D., Mizoguchi, H., Kamei, H., Nabeshima, T. and Yamada, K.:

Repeated methamphetamine treatment impairs spatial working memory in rats: reversal by clozapine but not haloperidol. Psychopharmacology (Berl.), 194, 21-32, 2007

Nakajima, M., Yamada, K., Hattori, E., Iwayama, Y., Toyota, T., Iwata, Y., Tsuchiya, K.J., Sugihara, G., Hashimoto, K., Watanabe, H., Iyo, M., Kikuchi, M., Okazaki, Y., Yoshikawa, T.

Genetic examination of the PLXNA2 gene in Japanese and Chinese schizophrenics. Schizophrenia Res., in press

Kobayashi T, Nishizawa D, Iwamura T, Ikeda K.

Inhibition by cocaine of G protein-activated inwardly rectifying K⁺ channels expressed in Xenopus oocytes. Toxicol In Vitro,21,656-664, 2007

熱田英範, 横溝美緒, 筒井啓太 他

Imipramine によって音楽性幻聴が生じた反復性うつ病の1症例. 精神科,10,518-523, 2007

曾良一郎, 猪狩もえ, 山本秀子, 池田和隆

依存性薬物の分子標的としてのモリアミントランスポーター 日本薬理学雑誌,130,450-454, 2007

柏 淳, 金子雄二郎, 伊藤 卓 他

統合失調症の薬理学再発防止への展望 —セロトニン作動性薬物による逆耐性消失から
脳と精神の医学,18,197-203, 2007

林田眞和, 福田謙一, 池田和隆

ゲノムペインクリニック時代へ向けて-その2- ペインクリニック,28,1451-1452, 2007

桂 昌司, 大熊誠太郎

【薬理学・毒性学】ベンゾジアゼピン系薬物依存の新知見. 医学のあゆみ,223,187,2007

桂 昌司, 大熊誠太郎

【アルコール医学・医療の最前線】アルコール依存症、アルコール依存症病態発現の
神経薬理的機序. 医学のあゆみ,222,677-684, 2007

曾良一郎

薬物依存の分子病態. 実験医学,25(13),179-184, 2007

井手聡一郎, 笠井慎也, 池田和隆

医療用麻薬の鎮痛効果の個人差 神経精神薬理学雑誌, in press

山本直樹, 島津 奈, 海野麻未 他

グルタミン酸-D-セリン系に作用する統合失調症の新規治療薬開発に関する研究-
精神薬療研究年報,39,22-27, 2007

桂 昌司, 芝崎真裕, 大熊誠太郎

細胞内カルシウム動態変化からみたアルコール精神および身体依存形成. 日本アル
コール・薬物医学会雑誌,42,59-66, 2007

桂 昌司, 芝崎真裕, 黒川和宏, 大熊誠太郎

依存性薬物による精神依存および身体依存形成機序の相違. 日本アルコール・薬物
医学会雑誌,42,481-486, 2007

山本経之

行動薬理学的側面からみた動物モデルの意義・役割・問題点 日本薬理学雑誌, 130, 94-96,
2007

笠井慎也, 池田和隆

オピオイド受容体～ミューオピオイド受容体の機能を中心に～ 日本薬理学雑誌, 130,
235-237, 2007

曾良一郎, 猪狩もえ, 山本秀子, 池田和隆

依存性薬物の分子標的としてのモノアミントランスポーター. 日本薬理学雑誌, 130
(6), 450-454, 2007

岩橋和彦, 加賀央子, 青木淳, 吉原英児, 村山洋, 渡辺全朗, 池田和隆, 石郷岡純

オランザピンの副作用による体重増加・血糖上昇と俟約遺伝子やセロトニン系遺伝子
との関連—統合失調症 16 例による予備的検討—. 脳と精神の医学, 18(2), 143-148, 2007

曾良一郎, 福島攝, 山本 秀子, 池田 和隆

ADHD 動物モデルとしてのドーパミントランスポーター(DAT) 欠損マウス. 脳と精神
の医学, 18(4), 279-283, 2007

山本経之, 縄田陽子, 當原真奈美

脳内 endocannabinoid の異常性に基づくマウスの知的機能障害 分子精神医学, 7, 83-84,
2007

正木秀和, 西川 徹.

統合失調症の治療薬開発研究: 特集「新しい時代の統合失調症—研究から治療へ—」.
臨床精神医学, 36, 43-51, 2007