

INTRODUCTION

Ultraviolet (UV) absorbers are added to plastics to prevent polymer degradation due to UV rays, such as loss of strength, reduced flexibility and electric properties, discoloration, scratching, and loss of gloss (Commerce Online, 2007; Tenkazai.com, 2007). Currently, many kinds of UV absorbers are used: benzotriazoles, benzophenones, salicylates, cyanoacrylates, nickels, triazines, etc. Among them, benzotriazole UV absorbers are known to have the most excellent absorption capacity with a full spectrum of UV absorption and are, therefore, used in a variety of polymers.

2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole (CAS No. 3846-71-7; HDBB) is a benzotriazole UV absorber added at ~0.02%–2% mainly to unsaturated polyester resin, polycarbonate, vinyl chloride resin, polyacrylic acid ester, polyacetal, polyolefin, polymethacrylic acid ester, and polyamide (METI, 2006). From these resins, plastic resin products, such as building materials and automobile components, are manufactured. In addition, HDBB is also used in printing or sensitive materials and coating compounds, all intended for UV absorption.

In spite of such widespread use, no reliable data were available on the toxicity of HDBB; therefore, this chemical was selected as an object substance in an existing chemical testing program by the Japanese Government (MHLW, 2003; 2006). Previously, we reported the result of a 28-day repeated dose toxicity study of HDBB conducted under this program (Hirata-Koizumi et al., 2007). In this study, CD(SD)IGS rats were administered HDBB by gavage at a dose of 0.5, 2.5, 12.5, or 62.5 mg/kg/day. As a result, adverse effects, mainly on the liver and heart, were found at all doses in males and at 12.5 mg/kg and higher in females. Anemic changes and histopathological changes in the kidneys and thyroids were also observed at the higher dose. These changes remained after the 14-day recovery period. The no observed adverse effect level (NOAEL) for females was concluded to be 2.5 mg/kg/day based on the induction of hypertrophy and increased mitosis of hepatocytes, and the degeneration and hypertrophy of the myocardium at 12.5 mg/kg. On the other hand, the NOAEL for males could not be determined because hypertrophy and decreased incidence of fatty change of hepatocytes and bile duct proliferation were noted at the lowest dose of 0.5 mg/kg. Considering the toxic effects observed at a relatively low dose and the incomplete recovery, more severe damage induced by longer exposure was a concern; therefore, a chronic toxicity study was performed under the Japanese existing chemical testing program. We here report the details of the results of a 52-week repeated dose toxicity study in rats.

MATERIALS AND METHODS

This study was performed in compliance with the OECD Guideline 452 "Chronic Toxicity Studies" (OECD, 1981) and in accordance with the principles

for Good Laboratory Practice (OECD, 1998; EA, MHW and MITI, 2000) at the Safety Assessment Laboratory, Panapharm Laboratories Co., Ltd. (Kumamoto, Japan).

Chemicals

HDBB was obtained from Shipro Kasei Kaisha, Ltd. (Osaka, Japan). The HDBB (Lot no. S4-034-1) used in this study was 100% pure, based on analysis using liquid chromatography, and it was kept at room temperature. The purity and stability during the study were verified by analysis before and after animal experiments. HDBB was dissolved in corn oil once or twice a week and kept in a dark, cool place until dosing since stability under these conditions was confirmed for up to eight days. The concentrations of formulations were confirmed to be 98.0%–102.0% of the target by analysis using high-performance liquid chromatography (HPLC). All other reagents used in this study were of specific purity grade.

Animals

Crj: CD (SD) IGS rats (SPF, five weeks old) were purchased from Atsugi Breeding Center, Charles River Laboratories Japan, Inc. (Yokohama, Japan). After a seven- or eight-day acclimation, they were subjected to treatment at six weeks of age. Rats found to be in good health were selected and assigned to four groups of 20 males and 20 females by stratified random sampling based on body weight.

All animals were maintained in an air-conditioned room at 21–27°C, with a relative humidity of 47%–60%, a 12-h light/dark cycle, and ventilation with 13–15 air changes/h. They were housed individually, except during the acclimation period, in stainless steel hanger cages. A basal diet (CRF-1; Oriental Yeast Co., Ltd., Tokyo, Japan) and sodium-hypochlorite-added well water were provided *ad libitum*.

This experiment was approved by the Ethical Committee for Animal Experiments of Panapharm Laboratories, Co., Ltd. and performed in accordance with the Guidance for Animal Experiments of Panapharm Laboratories, Co., Ltd.

Experimental Design

Male and female rats were given HDBB once-daily by gavage for 52 weeks at 0 (vehicle control), 0.1, 0.5, or 2.5 mg/kg/day and at 0, 0.5, 2.5, or 12.5 mg/kg/day, respectively. The dosage levels were determined based on the results of our previous 28-day repeated dose toxicity study in rats given HDBB by gavage at 0.5, 2.5, 12.5, or 62.5 mg/kg/day, in which adverse effects, mainly on the liver and hearts, were found at all doses in males, and at 12.5 mg/kg and more in females (Hirata-Koizumi et al., 2007). The volume of each dose was

adjusted to 5 mL/kg of body weight, based on the latest body weight. At the end of the 13-week administration period, 10 males and 10 females from each group were euthanized for the assessment of hematology, blood biochemistry, organ weights, and macroscopic and microscopic findings. The remaining animals in all groups (10 rats/sex/dose) were fully examined at the completion of the 52-week administration period.

All animals were observed daily before and after dosing for clinical signs of toxicity. Body weight and food consumption were recorded weekly for the first 13 weeks of the administration period, and once every four weeks for the remainder of the dosing period. At weeks 13 and 52 of the dosing period, fresh urine was collected. It was examined microscopically for urinary sediment and analyzed for dipstick parameters, such as occult blood, pH, protein, glucose, ketone bodies, bilirubin, and urobilinogen. In addition, a 24-h urine sample was also collected for the determination of sodium, potassium, and chlorine levels, color, specific gravity, osmotic pressure, and volume of urine.

Prior to necropsy at the end of the 13- and 52-week dosing periods, blood was collected from the caudal vena cava in the abdomen under deep anesthesia by the intraperitoneal (i.p.) injection of pentobarbital sodium after overnight starvation. One portion of the blood was treated with ethylenediaminetetraacetic acid (EDTA)-2K and examined for hematological parameters, such as red blood cell count, hemoglobin, hematocrit, mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), white blood cell count, platelet count, reticulocyte count, and differential leukocyte count. Prothrombin time (PT) and activated partial thromboplastin time (APTT) were measured using plasma separated from another blood sample treated with 3.8% sodium citrate. Serum from the remaining portions of blood was analyzed for blood biochemistry (total protein, protein fraction ratio, albumin-globulin (A/G) ratio, glucose, total cholesterol, triglycerides, phospholipid, total bilirubin, urea nitrogen (BUN), creatinine, aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase (ALP), calcium, inorganic phosphorus, sodium, potassium, and chlorine).

Following the collection of blood, all animals were sacrificed by exsanguination, and organs and tissues of the entire body were macroscopically observed. The brain, pituitary, thymus, thyroids (including parathyroids), heart, lungs (including bronchus), liver, spleen, kidneys, adrenals, testes, epididymides, ovaries, and uterus were then excised and weighed. The trachea, pancreas, lymph nodes (mandibular and mesenteric), tongue, sublingual gland, submandibular gland, parotid gland, esophagus, stomach, duodenum, jejunum, ileum, cecum, colon, rectum, urinary bladder, eyeballs, optic nerve, Harderian gland, spinal cord (pectoral and lumbar part), sciatic nerve, seminal vesicles, prostates, vagina, mammary gland, aorta (thoracic), bone (sternum and femur including bone marrow), skeletal muscle (biceps femoris

muscle), and skin (hypogastric) as well as the above organs were fixed in 10% neutral-buffered formalin solution (following Bouin's fixation for testes and epididymides, and 2.5% glutaraldehyde fixation for eyeballs, optic nerve, and Harderian gland). Histopathological examination of these organs was conducted for all animals found dead or moribund, and for scheduled-sacrifice animals in the control and highest dose groups. In addition, the livers of males in the lowest dose group and of both sexes in the middle-dose group were examined, since test substance-related changes were found in the higher group. Paraffin sections for microscopic examination were routinely prepared and stained with hematoxylin and eosin.

Data Analysis

Parametric data, such as body weight, food consumption, urinalysis findings (sodium, potassium, chlorine, specific gravity, osmotic pressure, and volume), hematological and blood biochemical findings, and organ weights were analyzed by Bartlett's test (Bartlett, 1937) for homogeneity of distribution. When homogeneity was recognized, Dunnett's test (Dunnett, 1964) was conducted for comparison between control and individual treatment groups. If not homogenous, data were analyzed using Steel's multiple comparison test (Steel, 1959). For dipstick parameters, color, and sediment of urine, the grades were converted into numeric values, for which Steel's multiple comparison test (Steel, 1959) was conducted. Macroscopic and histopathological findings were analyzed using Fisher's exact test (Fisher, 1973) and Mann-Whitney's U test (Mann and Whitney, 1947), respectively. These analyses were all conducted by a two-tailed test with a significance level of 1% and 5%.

RESULTS

One male at 2.5 mg/kg was found dead on day 54 of the administration period. Two males at 0.1 mg/kg were also found dead on days 231 or 357 of the administration period. In addition, one female at 12.5 mg/kg was found moribund and was, therefore, euthanized on day 354 of the administration period.

In animals surviving to completion of the 13- or 52-week administration period, no substance-related clinical signs of toxicity were observed; however, body weight was significantly lowered from day 36 to the end of the 52-week dosing period at 2.5 mg/kg in males. A significant increase in food consumption was also detected on days 120, 204–288, and 364 of the dosing period in this group of males.

Examination at Completion of the 13-Week Administration Period

With urine analysis, a significant increase in osmotic pressure and specific gravity was detected at 2.5 mg/kg in males. No changes were noted in other parameters of urinalysis in any HDBB-treated groups (data not shown).

In hematological examination, a significant decrease in hemoglobin and hematocrit at 0.5 mg/kg and higher, decrease in red blood cell count, and increase in platelet count at 2.5 mg/kg was found in males (Table 1). In females, a significant decrease in hematocrit and MCV was noted at 12.5 mg/kg (Table 2). Blood biochemical examination revealed a significant increase in serum levels of glucose, BUN, and ALP at 0.5 mg/kg and higher in males (Table 3) and of total protein at 12.5 mg/kg in females (Table 4). A significant change in the serum protein fraction, such as an increase in albumin and decrease in α_2 - and β -globulin at 0.5 mg/kg and higher in males, and at 12.5 mg/kg in females, and a decrease in α_1 -globulin at 0.5 mg/kg and higher in males, was also found with a significant increase in the A/G ratio at 0.5 mg/kg and higher in males, and at 12.5 mg/kg in females. There were no substance-related changes in other blood biochemical parameters, including total bilirubin level (data not shown).

At necropsy, enlargement of the liver was observed in five of nine males at 2.5 mg/kg and in one of ten females at 12.5 mg/kg, and the absolute and relative liver weight was significantly increased at 0.5 mg/kg and higher in males, and at 12.5 mg/kg in females (Tables 5 and 6). A significant increase in

Table 1: Hematological findings in male rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.1	0.5	2.5
At completion of the 13-week administration period				
No. of animals	10	10	10	9
Red blood cells ($10^4/\mu\text{L}$)	855 \pm 27	870 \pm 29	828 \pm 43	807 \pm 22**
Hemoglobin (g/dL)	15.6 \pm 0.4	15.5 \pm 0.5	15.0 \pm 0.6*	14.3 \pm 0.6**
Hematocrit (%)	43.3 \pm 1.6	43.2 \pm 1.2	41.6 \pm 1.8*	40.0 \pm 1.3**
MCV (fL)	50.6 \pm 1.6	49.6 \pm 1.1	50.3 \pm 0.9	49.6 \pm 2.6
MCH (pg)	18.3 \pm 0.5	17.8 \pm 0.5	18.1 \pm 0.5	17.7 \pm 1.0
MCHC (g/dL)	36.1 \pm 0.7	35.9 \pm 0.4	36.0 \pm 0.7	35.7 \pm 0.5
Reticulocyte ($10^4/\mu\text{L}$)	15.8 \pm 2.8	16.3 \pm 1.8	16.2 \pm 3.2	14.8 \pm 3.6
Platelet count ($10^4/\mu\text{L}$)	103.4 \pm 11.2	108.7 \pm 8.2	112.9 \pm 16.0	130.5 \pm 27.1*
PT (s)	15.2 \pm 2.7	14.9 \pm 1.1	15.1 \pm 1.5	14.3 \pm 1.4
APTT (s)	24.6 \pm 1.9	24.1 \pm 1.9	23.0 \pm 1.5	23.5 \pm 3.1
At completion of the 52-week administration period				
No. of animals	10	8	10	10
Red blood cells ($10^4/\mu\text{L}$)	840 \pm 68	780 \pm 145	754 \pm 133*	778 \pm 66*
Hemoglobin (g/dL)	14.0 \pm 1.1	13.1 \pm 2.7	12.7 \pm 2.1	12.9 \pm 1.1
Hematocrit (%)	44.2 \pm 2.9	41.3 \pm 7.4	40.3 \pm 5.7	40.7 \pm 3.6*
MCV (fL)	52.7 \pm 2.1	53.1 \pm 1.2	53.9 \pm 4.5	52.3 \pm 2.3
MCH (pg)	16.7 \pm 0.8	16.7 \pm 0.7	16.9 \pm 1.0	16.6 \pm 0.7
MCHC (g/dL)	31.7 \pm 0.8	31.5 \pm 1.5	31.3 \pm 1.1	31.8 \pm 0.3
Reticulocyte ($10^4/\mu\text{L}$)	18.2 \pm 8.4	20.1 \pm 9.8	27.1 \pm 20.4	15.7 \pm 3.3
Platelet count ($10^4/\mu\text{L}$)	106.5 \pm 12.6	110.2 \pm 28.5	123.7 \pm 28.5	140.1 \pm 13.6**
PT (s)	13.5 \pm 1.0	13.8 \pm 1.0	14.5 \pm 1.9	21.8 \pm 9.0**
APTT (s)	21.5 \pm 1.5	20.9 \pm 2.7	21.2 \pm 2.6	29.5 \pm 9.3

Values are expressed as the mean \pm SD.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

Table 2: Hematological findings in female rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.5	2.5	12.5
At completion of the 13-week administration period				
No. of animals	10	10	10	10
Red blood cells ($10^4/\mu\text{L}$)	768 \pm 38	793 \pm 40	762 \pm 23	753 \pm 25
Hemoglobin (g/dL)	13.9 \pm 0.5	14.1 \pm 0.6	13.8 \pm 0.4	13.4 \pm 0.5
Hematocrit (%)	40.1 \pm 1.7	40.7 \pm 2.2	39.5 \pm 0.9	38.1 \pm 1.2*
MCV (fL)	52.2 \pm 1.1	51.3 \pm 0.7	51.9 \pm 1.3	50.6 \pm 1.0**
MCH (pg)	18.1 \pm 0.4	17.7 \pm 0.4	18.1 \pm 0.5	17.7 \pm 0.5
MCHC (g/dL)	34.6 \pm 0.5	34.6 \pm 0.6	35.0 \pm 0.3	35.1 \pm 0.5*
Reticulocyte ($10^4/\mu\text{L}$)	16.5 \pm 3.4	13.9 \pm 1.9	14.8 \pm 3.4	13.7 \pm 1.6
Platelet count ($10^4/\mu\text{L}$)	106.1 \pm 12.1	110.4 \pm 6.8	117.4 \pm 11.6	106.2 \pm 9.9
PT (s)	11.7 \pm 0.5	11.7 \pm 0.3	11.7 \pm 0.3	11.8 \pm 0.4
APTT (s)	19.2 \pm 1.5	19.7 \pm 0.9	19.0 \pm 1.6	19.2 \pm 1.5
At completion of the 52-week administration period				
No. of animals	10	10	10	9
Red blood cells ($10^4/\mu\text{L}$)	707 \pm 100	708 \pm 62	730 \pm 55	673 \pm 115
Hemoglobin (g/dL)	13.2 \pm 1.4	13.5 \pm 0.8	13.5 \pm 1.0	12.3 \pm 1.5
Hematocrit (%)	40.3 \pm 3.8	41.0 \pm 2.5	41.3 \pm 3.0	37.3 \pm 4.4
MCV (fL)	57.5 \pm 4.3	58.1 \pm 2.3	56.6 \pm 2.4	56.1 \pm 4.8
MCH (pg)	18.8 \pm 1.0	19.1 \pm 0.7	18.5 \pm 0.8	18.4 \pm 1.4
MCHC (g/dL)	32.7 \pm 0.9	33.0 \pm 0.5	32.7 \pm 0.6	32.9 \pm 0.4
Reticulocyte ($10^4/\mu\text{L}$)	14.9 \pm 8.9	16.4 \pm 9.6	13.9 \pm 5.8	17.1 \pm 15.1
Platelet count ($10^4/\mu\text{L}$)	90.2 \pm 10.0	94.2 \pm 14.7	101.5 \pm 13.9	105.6 \pm 11.9*
PT (s)	12.3 \pm 0.8	12.9 \pm 0.7	12.5 \pm 0.5	12.1 \pm 0.5
APTT (s)	18.4 \pm 0.9	18.5 \pm 0.9	17.7 \pm 1.4	17.7 \pm 1.2

Values are expressed as the mean \pm SD.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

the relative weight of the brain, heart, kidneys, and testes was also found at 2.5 mg/kg in males, but the absolute weight was not significantly changed. On histopathology, centrilobular hypertrophy of hepatocytes, accompanied with eosinophilic granular cytoplasm, was observed in the liver (Tables 7 and 8). The incidence was significantly increased at 2.5 mg/kg in males and at 12.5 mg/kg in females.

Examination at Completion of the 52-Week Administration Period

Urinalysis revealed a significant increase in osmotic pressure at 0.5 mg/kg and higher in males, while it was significantly decreased at 12.5 mg/kg in females. A significant increase in urine volume was also detected at 12.5 mg/kg in females (data not shown).

On hematological examination, a significant decrease in the red blood cell count at 0.5 mg/kg and higher, and in hematocrit at 2.5 mg/kg in males, and increase in platelet count at 2.5 mg/kg in males, and at 12.5 mg/kg in females was found (Tables 1 and 2). In addition, PT was significantly prolonged at 2.5 mg/kg in males. In the blood biochemical examination, a significant

Table 3: Blood biochemical findings in male rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.1	0.5	2.5
At completion of the 13-week administration period				
No. of animals	10	10	10	9
Total protein (g/dL)	5.8 ± 0.3	5.8 ± 0.2	5.7 ± 0.5	5.8 ± 0.5
A/G ratio	1.22 ± 0.12	1.30 ± 0.09	1.67 ± 0.23**	2.09 ± 0.27**
Protein fraction ratio				
α ₁ -Globulin (%)	18.7 ± 1.6	17.9 ± 1.6	15.6 ± 1.3**	12.1 ± 2.4**
α ₂ -Globulin (%)	7.1 ± 0.7	6.8 ± 0.6	5.9 ± 0.6**	5.6 ± 0.6**
β-Globulin (%)	15.2 ± 0.8	14.4 ± 0.6	11.5 ± 1.0**	9.9 ± 0.7**
γ-Globulin (%)	4.2 ± 0.5	4.3 ± 0.6	4.6 ± 0.8	5.0 ± 1.4
Albumin (%)	54.8 ± 2.3	56.6 ± 1.6	62.4 ± 2.9**	67.4 ± 3.0**
ALP (IU/L)	164 ± 23	216 ± 57	373 ± 60**	619 ± 115**
Glucose (mg/dL)	121 ± 9	120 ± 7	154 ± 13**	151 ± 9**
BUN (mg/dL)	12.3 ± 1.1	11.8 ± 1.7	14.2 ± 1.7*	14.8 ± 1.8**
At completion of the 52-week administration period				
No. of animals	10	8	10	10
Total protein (g/dL)	5.8 ± 0.2	5.8 ± 0.3	5.8 ± 0.5	5.8 ± 0.2
A/G ratio	1.01 ± 0.21	1.01 ± 0.29	1.42 ± 0.31**	1.75 ± 0.30**
Protein fraction ratio				
α ₁ -Globulin (%)	19.2 ± 2.2	18.2 ± 1.8	15.2 ± 2.4**	13.4 ± 2.0**
α ₂ -Globulin (%)	7.5 ± 0.5	7.1 ± 1.4	6.1 ± 1.3*	5.0 ± 1.1**
β-Globulin (%)	17.9 ± 2.3	18.5 ± 4.5	15.3 ± 3.0	12.7 ± 2.2**
γ-Globulin (%)	5.7 ± 2.3	6.9 ± 3.1	5.2 ± 1.7	5.8 ± 1.2
Albumin (%)	49.7 ± 5.4	49.3 ± 8.4	58.1 ± 5.4**	63.2 ± 4.7**
ALP (IU/L)	141 ± 42	165 ± 56	364 ± 87**	565 ± 137**
Glucose (mg/dL)	125 ± 27	115 ± 11	139 ± 17	125 ± 16
BUN (mg/dL)	9.1 ± 1.5	8.8 ± 0.9	10.4 ± 1.9	12.8 ± 1.5**

Values are expressed as the mean ± SD.

*Significantly different from the control, $p < 0.05$; **Significantly different from the control, $p < 0.01$.

increase in the levels of ALP at 0.5 mg/kg and higher in males, and at 12.5 mg/kg in females, of BUN at 2.5 mg/kg in males, and of glucose at 12.5 mg/kg in females was found (Tables 3 and 4). For the serum protein fraction ratio, a significant increase in albumin and decrease in α₁- and α₂-globulin at 0.5 mg/kg and higher, and a decrease in β-globulin at 2.5 mg/kg was detected in males. The A/G ratio was significantly increased at 0.5 mg/kg and higher in males. No substance-related changes were found in other blood biochemical parameters, including total bilirubin level (data not shown).

At necropsy, enlarged liver was observed in seven of ten males at 0.5 mg/kg, nine of ten males at 2.5 mg/kg, and five of nine females at 12.5 mg/kg, and light gray macules were grossly detected in the liver of two of ten males at 2.5 mg/kg and of one of nine females at 12.5 mg/kg. Absolute and relative liver weight was significantly increased at 0.5 mg/kg and higher in males, and at 12.5 mg/kg in females (Tables 5 and 6). A significant increase in the relative weight of the brain, pituitary, thyroids, lungs, heart, kidneys, testes, and epididymides at 2.5 mg/kg in males was also found, but no statistically significant

Table 4: Blood biochemical findings in female rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.5	2.5	12.5
At completion of the 13-week administration period				
No. of animals	10	10	10	10
Total protein (g/dL)	6.2 ± 0.4	6.3 ± 0.2	6.4 ± 0.4	6.7 ± 0.5*
A/G ratio	1.78 ± 0.16	1.87 ± 0.22	1.93 ± 0.19	2.24 ± 0.31**
Protein fraction ratio				
α ₁ -Globulin (%)	13.8 ± 1.0	12.9 ± 1.7	12.6 ± 1.6	12.9 ± 1.8
α ₂ -Globulin (%)	5.6 ± 0.8	5.6 ± 0.2	5.5 ± 0.6	4.7 ± 0.5*
β-Globulin (%)	12.6 ± 0.9	12.4 ± 1.2	12.1 ± 1.4	9.9 ± 0.8**
γ-Globulin (%)	3.9 ± 0.8	4.3 ± 1.0	4.2 ± 1.0	3.6 ± 1.1
Albumin (%)	64.0 ± 2.0	64.9 ± 2.8	65.7 ± 2.2	68.9 ± 2.9**
ALP (IU/L)	92 ± 30	107 ± 25	101 ± 31	136 ± 81
Glucose (mg/dL)	119 ± 13	117 ± 10	118 ± 15	130 ± 10
BUN (mg/dL)	14.5 ± 1.7	14.3 ± 1.7	13.6 ± 1.1	14.1 ± 1.8
At completion of the 52-week administration period				
No. of animals	10	10	10	9
Total protein (g/dL)	6.4 ± 0.3	6.7 ± 0.2	6.7 ± 0.3	6.5 ± 0.5
A/G ratio	1.79 ± 0.25	1.69 ± 0.17	1.73 ± 0.17	2.00 ± 0.19
Protein fraction ratio				
α ₁ -Globulin (%)	13.5 ± 1.6	14.2 ± 1.6	12.8 ± 1.4	12.1 ± 1.0
α ₂ -Globulin (%)	4.8 ± 0.6	4.8 ± 0.5	5.0 ± 0.9	4.1 ± 0.4
β-Globulin (%)	13.2 ± 1.5	13.5 ± 0.7	13.6 ± 1.6	12.2 ± 1.2
γ-Globulin (%)	4.6 ± 0.9	4.9 ± 1.2	5.4 ± 1.2	5.0 ± 1.2
Albumin (%)	63.9 ± 3.1	62.6 ± 2.5	63.3 ± 2.3	66.5 ± 2.1
ALP (IU/L)	57 ± 26	59 ± 16	57 ± 14	86 ± 20**
Glucose (mg/dL)	103 ± 9	110 ± 9	106 ± 16	119 ± 16*
BUN (mg/dL)	13.4 ± 2.7	12.6 ± 2.8	12.7 ± 3.1	12.1 ± 2.0

Values are expressed as the mean ± SD.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

change was noted in the absolute weight. As observed at the end of the 13-week administration period, centrilobular hypertrophy of hepatocytes with eosinophilic granular cytoplasm was observed on histopathological examination, and the incidence was significantly increased at 0.5 mg/kg and higher in males, and at 12.5 mg/kg in females (Tables 7 and 8). In addition, significant increases in the incidence of cystic degeneration and lipofuscin deposition in hepatocytes at 2.5 mg/kg, and of altered hepatocellular foci (clear cell foci) at 0.5 mg/kg and higher were found in the liver of males.

DISCUSSION

In the present study, one male receiving the highest dose of 2.5 mg/kg died early in the dosing period. Although the cause of death was not identified on histopathological examination, it is unlikely that this death was due to treatment with HDBB because no deaths in this group occurred during the remaining dosing period. Other the deaths of two males at 0.1 mg/kg and the

Table 5: Relative organ weight in male rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.1	0.5	2.5
At completion of the 13-week administration period				
No. of animals	10	10	10	9
Body weight ^a	530.1 ± 32.1	566.3 ± 42.2	546.5 ± 40.3	450.1 ± 27.8**
Brain ^b	0.42 ± 0.02	0.40 ± 0.03	0.42 ± 0.03	0.49 ± 0.03**
Pituitary ^c	2.7 ± 0.3	2.5 ± 0.2	2.6 ± 0.2	2.8 ± 0.2
Thyroids ^c	3.8 ± 1.0	4.7 ± 0.8	4.5 ± 1.1	4.1 ± 0.7
Heart ^b	0.29 ± 0.03	0.29 ± 0.02	0.30 ± 0.02	0.33 ± 0.02**
Lungs ^b	0.29 ± 0.02	0.28 ± 0.03	0.30 ± 0.02	0.31 ± 0.03
Liver ^b	2.75 ± 0.10	2.82 ± 0.23	3.71 ± 0.21**	5.12 ± 0.72**
Kidneys ^b	0.62 ± 0.04	0.62 ± 0.02	0.67 ± 0.06	0.70 ± 0.07*
Testes ^b	0.65 ± 0.07	0.62 ± 0.07	0.61 ± 0.06	0.81 ± 0.07**
Epididymides ^b	0.26 ± 0.02	0.25 ± 0.02	0.23 ± 0.02*	0.28 ± 0.03
At completion of the 52-week administration period				
No. of animals	10	8	10	10
Body weight ^a	819.9 ± 145.4	792.5 ± 140.4	842.4 ± 136.3	614.2 ± 97.3**
Brain ^b	0.30 ± 0.04	0.31 ± 0.07	0.29 ± 0.04	0.39 ± 0.05**
Pituitary ^c	2.0 ± 0.2	2.0 ± 0.5	1.9 ± 0.3	2.8 ± 0.3**
Thyroids ^c	3.8 ± 0.9	3.9 ± 1.0	4.1 ± 0.8	4.9 ± 0.9*
Heart ^b	0.23 ± 0.02	0.25 ± 0.04	0.25 ± 0.03	0.31 ± 0.03**
Lungs ^b	0.23 ± 0.02	0.24 ± 0.05	0.23 ± 0.02	0.29 ± 0.03**
Liver ^b	2.22 ± 0.25	2.26 ± 0.20	2.95 ± 0.47**	4.13 ± 0.62**
Kidneys ^b	0.47 ± 0.05	0.48 ± 0.08	0.51 ± 0.06	0.68 ± 0.09**
Testes ^b	0.45 ± 0.06	0.47 ± 0.10	0.46 ± 0.07	0.61 ± 0.15**
Epididymides ^b	0.16 ± 0.03	0.18 ± 0.04	0.17 ± 0.02	0.22 ± 0.06*

Values are expressed as the mean ± SD.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

^aBody weight after overnight starvation following the last dosing (g).

^bg/100 g body weight.

^cmg/100 g body weight.

moribund sacrifice of one female at 12.5 mg/kg were related to pituitary, renal, or muscular disorders, which was not observed in scheduled-sacrifice animals, and were considered incidental.

In scheduled-sacrifice animals, a lowered body weight was found at 2.5 mg/kg in males. This change was not observed even at the highest dose of 62.5 mg/kg in the previous 28-day repeated dose toxicity study of HDBB (Hirata-Koizumi et al., 2007). Since increased food consumption, blood glucose, and A/G ratio were noted in both previous 28-day and present 52-week studies, prolonged disturbance of metabolic homeostasis might affect body weight gain. Increased relative weight of the brain, heart, kidneys, testes, etc., without changes in the absolute weight, which was noted at 2.5 mg/kg in males in the present study, is considered to be due to the lowering of body weight.

Anemic changes, such as decreased red blood cell count, hematocrit, and hemoglobin, were also found at 0.5 mg/kg and higher in males in the current study. In females, slight changes indicative of anemia, such as decreased hematocrit and MCV, were noted at 12.5 mg/kg at the end of the 13-week

Table 6: Relative organ weight in female rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.5	2.5	12.5
At completion of the 13-week administration period				
No. of animals	10	10	10	10
Body weight ^a	304.1 ± 26.9	303.0 ± 31.0	297.0 ± 17.5	299.8 ± 23.1
Brain ^b	0.68 ± 0.06	0.69 ± 0.05	0.70 ± 0.03	0.70 ± 0.05
Pituitary ^c	5.6 ± 0.5	6.1 ± 0.7	6.4 ± 1.0*	6.2 ± 0.8
Thyroids ^c	5.5 ± 1.1	5.9 ± 0.8	6.5 ± 1.1*	6.2 ± 0.7
Heart ^b	0.32 ± 0.02	0.30 ± 0.02	0.32 ± 0.02	0.32 ± 0.03
Lungs ^b	0.37 ± 0.03	0.37 ± 0.02	0.36 ± 0.02	0.38 ± 0.03
Liver ^b	2.63 ± 0.14	2.63 ± 0.18	2.80 ± 0.18	3.88 ± 0.50**
Kidneys ^b	0.70 ± 0.25	0.64 ± 0.07	0.64 ± 0.05	0.66 ± 0.06
Ovaries ^c	26.1 ± 4.0	26.5 ± 3.2	26.9 ± 4.6	27.0 ± 4.0
Uterus ^b	0.19 ± 0.03	0.22 ± 0.04	0.19 ± 0.03	0.21 ± 0.03
At completion of the 52-week administration period				
No. of animals	10	10	10	9
Body weight ^a	423.2 ± 87.2	441.8 ± 71.4	481.0 ± 104.7	425.8 ± 71.4
Brain ^b	0.54 ± 0.12	0.51 ± 0.07	0.47 ± 0.10	0.52 ± 0.08
Pituitary ^c	6.6 ± 2.3	7.0 ± 3.8	7.1 ± 3.2	7.4 ± 2.4
Thyroids ^c	5.7 ± 1.1	5.5 ± 1.3	5.9 ± 1.1	6.4 ± 1.4
Heart ^b	0.28 ± 0.04	0.28 ± 0.04	0.26 ± 0.04	0.29 ± 0.03
Lungs ^b	0.33 ± 0.07	0.30 ± 0.05	0.29 ± 0.07	0.32 ± 0.05
Liver ^b	2.48 ± 0.39	2.42 ± 0.14	2.45 ± 0.32	3.54 ± 0.41**
Kidneys ^b	0.55 ± 0.08	0.54 ± 0.06	0.52 ± 0.13	0.63 ± 0.09
Ovaries ^c	16.0 ± 3.3	14.3 ± 4.4	13.5 ± 5.5	14.3 ± 2.5
Uterus ^b	0.24 ± 0.08	0.22 ± 0.06	0.22 ± 0.09	0.25 ± 0.08

Values are expressed as the mean ± SD.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

^aBody weight after overnight starvation following the last dosing (g).

^bg/100 g body weight.

^cmg/100 g body weight.

administration period but not at the completion of the 52-week administration period. The previous 28-day study also showed anemic effects of HDBB at 2.5 mg/kg and higher in males (Hirata-Koizumi et al., 2007). Since no change in the serum bilirubin level or hemosiderin accumulation in the liver, spleen, or kidneys were found in both the present 52-week and previous 28-day studies, anemic changes seem at least not to come from the hemolytic action of HDBB. In order to clarify the mechanisms, further study is required.

In the previous 28-day study, histopathological changes in the liver and heart were observed at 0.5 mg/kg and higher in males, and at 12.5 mg/kg and higher in females (Hirata-Koizumi et al., 2007). At higher doses, histopathological changes were also found in the kidneys and thyroids. In the current study, histopathological changes were observed in the liver. At the end of the 13-week administration, the incidence of centrilobular hypertrophy of hepatocytes was increased at 2.5 mg/kg in males and at 12.5 mg/kg in females, and this change was accompanied with eosinophilic granular cytoplasm. In addition to these changes, increased incidences of altered

Table 7: Histopathological findings in the liver of male rats given HDBB by gavage.

	Grade	Dose (mg/kg/day)				
		0	0.1	0.5	2.5	
At completion of the 13-week administration period						
No. of animals		10	10	10	9	
Centrilobular hypertrophy of hepatocytes ^a	+	0	0	3	6] **
	++	0	0	0	3	
Focal necrosis	+	1	0	1	2	
At completion of the 52-week administration period						
No. of animals		10	8	10	10	
Centrilobular hypertrophy of hepatocytes ^a	+	0	0	5*	7] **
	++	0	0	0	1	
Focal necrosis	+	1	0	3	4	
Lipofuscin deposition in hepatocytes ^b	+	0	0	0	6**	
Altered hepatocellular foci	+	0	1	7**	6] **
	++	0	0	0	1	
Cystic degeneration of hepatocytes	+	0	2	2	4*	

Values represent the number of animals with findings.

+ = mild; ++ = moderate.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

^aAccompanied with eosinophilic granular cytoplasm.

^bIdentified by the Schmorl method, Berlin blue staining, and the Hall method.

Table 8: Histopathological findings in the liver of female rats given HDBB by gavage.

	Grade	Dose (mg/kg/day)			
		0	0.5	2.5	12.5
At completion of the 13-week administration period					
No. of animals		10	10	10	10
Centrilobular hypertrophy of hepatocytes ^a	+	0	-	0	6**
Focal necrosis	+	0	-	1	0
At completion of the 52-week administration period					
No. of animals		10	10	10	9
Centrilobular hypertrophy of hepatocytes ^a	+	0	-	0	4*
Focal necrosis	+	2	-	0	0
Lipofuscin deposition in hepatocytes ^b	+	0	-	0	2

Values represent the number of animals with findings.

+ = mild; - = not examined.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

^aAccompanied with eosinophilic granular cytoplasm.

^bIdentified by the Schmorl method, Berlin blue staining, and the Hall method.

hepatocellular foci (clear cell foci) at 0.5 mg/kg and higher and of cystic degeneration and lipofuscin deposition in hepatocytes at 2.5 mg/kg were found in males at the completion of the 52-week administration.

Centrilobular hypertrophy of hepatocytes with eosinophilic granular cytoplasm is known to be a characteristic change observed in rodents administered with peroxisome proliferators, such as fibrate hypolipidemic drugs and phthalate plasticizers (Cattley and Popp, 2002). Prolonged exposure to these substances has been shown in many studies to induce liver tumors in rats and mice (IARC, 1995), and preferential growth of altered hepatocytes, as detected in the present 52-week study, could be observed in the developmental process. In addition, increased mitosis of hepatocytes, indicating hepatocellular proliferation, was observed in the higher dose group in the previous 28-day study of HDBB (Hirata-Koizumi et al., 2007). Further longer-term studies are needed to precisely evaluate whether HDBB induces liver tumors in rats. In the current study, lipofuscin deposition in hepatocytes was also apparent at the completion of the 52-week administration. While lipofuscin accumulates in hepatocytes with aging, increased amounts of lipofuscin have also been reported in the liver of rats treated for long periods with peroxisome proliferators (IARC, 1995; Cattley and Popp, 2002). Based on these findings, HDBB might exert an effect on the liver via the mechanism of peroxisome proliferation, although the ultrastructure or peroxisome-associated enzyme was not analyzed in the current study. The hepatic changes caused by this mechanism are considered not to be significant for human risk assessment (Hasegawa et al., 2004) because primates are much less sensitive to peroxisome proliferators than rodents (Elcombe and Mitchell, 1986; Blaauboer et al., 1990). For HDBB, however, the incidence of cystic degeneration of hepatocytes was increased at the end of the current 52-week study, and increased incidence of focal necrosis, vacuolar degeneration of hepatocytes, and bile duct proliferation in the liver was found in the previous 28-day study (Hirata-Koizumi et al., 2007). These changes may not be necessarily associated with the mechanism of peroxisome proliferation. Considering the possible induction of neoplastic change in the liver by mechanisms relevant to humans, further study is required.

In the current study, histopathological changes in the heart were not detected even at the highest dose of 2.5 mg/kg in males and 12.5 mg/kg in females, at which degeneration and hypertrophy of the myocardium or cell infiltration were found in the previous 28-day study (Hirata-Koizumi et al., 2007). Although the cause of this difference between studies is not clear, the borderline dose of HDBB for affecting the heart is considered to be around 2.5 mg/kg in males and 12.5 mg/kg in females. As functional parameters are considered to be more sensitive than histopathological changes in the heart (Glaister, 1992), further studies are required to clarify the adverse effects of HDBB on cardiac function. Histopathological changes in the kidneys and

thyroids, detected in the previous 28-day study (Hirata-Koizumi et al., 2007), were also not observed in the present study, which would be due to the low dosage administered; however, changes in osmotic pressure, specific gravity, or volume of urine, and/or increase in the levels of BUN, noted at 0.5 mg/kg and higher in males and at 12.5 mg/kg in females, suggest renal effects of HDBB.

Based on these findings in the current study, the NOAEL for chronic toxicity of HDBB was concluded to be 0.1 mg/kg/day in male rats and 2.5 mg/kg/day in female rats based on the induction of altered hepatocellular foci and/or hypertrophy of hepatocytes. This result showed that male rats are nearly 25 times more susceptible to HDBB toxicity than female rats, which is consistent with the results of the previous 28-day study (Hirata-Koizumi et al., 2007). Since male rats showed higher susceptibility to various effects of HDBB (on the liver, blood, etc.) consistently, sex-related variations in toxicokinetic determinants, such as metabolism and elimination, might increase the blood concentration of causative substances (i.e., HDBB or its metabolites) in males. In order to clarify the cause of the sexual differences in the HDBB toxicity, we are planning a toxicokinetic study of HDBB, which would include the identification of metabolites and the related metabolic enzyme as well as measurement of the blood concentration of HDBB both after single and repeated administration of HDBB to rats.

Gender-related differences in toxic susceptibility have been documented for other substances. For example, a recent subchronic toxicity study using F344 rats showed that fluoranthene, a polycyclic aromatic hydrocarbon, had greater effects on males than females, especially in the kidneys (Knuckles et al., 2004). In contrast, it was reported that female rats exhibited a greater susceptibility to hypothermic effects and inhibition of hypothalamic cholinesterase by a carbamate cholinesterase inhibitor, rivastigmine (Wang et al., 2001). For such gender differences, sexual hormones must play an important role. In fact, Wang et al. (2001) reported that orchidectomy completely abolished the above-mentioned sex differences in hypothalamic cholinesterase inhibition induced by rivastigmine. Since testosterone decreased cholinesterase inhibition in gonadectomized males and females, it is apparent that testosterone interferes with the effects of rivastigmine. It is interesting to investigate the role of sex steroids in the mediation of sex differences in toxic susceptibility to HDBB; therefore, we are currently performing a repeated-dose toxicity study of HDBB using male and female castrated rats.

CONCLUSIONS

The current results showed that the oral administration of HDBB for 52 weeks principally affected the liver. The NOAEL for chronic toxicity was concluded to be 0.1 mg/kg/day in male rats and 2.5 mg/kg/day in female rats.

ACKNOWLEDGMENTS


This study was supported by the Ministry of Health, Labour and Welfare, Tokyo, Japan.

REFERENCES

- Bartlett, M. S. (1937). Properties of sufficiency and statistical tests. *Proc. R. Soc. Lond. Ser. A* 160:268–282.
- Blaauboer, B. J., van Holsteijn, C. W., Bleumink, R., Mennes, W. C., van Pelt, F. N., Yap, S. H., van Pelt, J. F., van Iersel, A. A., Timmerman, A., Schmid, B. P. (1990). The effect of becloric acid and clofibrilic acid on peroxisomal β -oxidation and peroxisome proliferation in primary cultures of rat, monkey, and human hepatocytes. *Biochem. Pharmacol.* 40:521–528.
- Cattley, R. C., Popp, J. A. (2002). Liver. In Haschek, W. M., Rousseaux, C. G., Wallig, M.A., (eds.), *Handbook of Toxicologic Pathology*, 2nd ed. Vol. 2. San Diego: Academic Press, pp. 187–225.
- Commerce Online. (2007). Product Keywords on Wujiang Dongfeng Chemical Co., Ltd. Accessed on April 25, 2007 from http://www.commerce.com.tw/company_inside.php?ID=C0013309.
- Dunnnett, C. W. (1964). New tables for multiple comparisons with a control. *Biometrics* 20:482–491.
- EA, MHW and MITI (Environment Agency, Ministry of Health and Welfare, and Ministry of International Trade and Industry of Japan). (2000). “Testing Facility Provided in the Article 4 in the Ordinance Prescribing Test Relating to New Chemical Substances and Toxicity Research of Designated Chemical Substances,” Planning and Coordination Bureau, Environment Agency No. 41 and Environmental Health Bureau, Ministry of Health and Welfare No. 268, dated March 1, 2000, and Basic Industries Bureaus, Ministry of International Trade and Industry No. 1, dated February 14, 2000.
- Elcombe, C. R., Mitchell, A. M. (1986). Peroxisome proliferation due to di(2-ethylhexyl) phthalate (DEHP): species differences and possible mechanisms. *Environ. Health Perspect.* 70:211–219.
- Fisher, R. A. (1973). *Statistical Methods of Research Workers*, 14th ed. New York: Hapner Publishing Company, p. 6.
- Glaister, J. R. (1992). Histopathology of target organs—cardiovascular. In *Principles of Toxicological Pathology (Japanese version supervised by Takahashi, M.)*, Tokyo: Soft Science Inc., pp. 135–142.
- Hasegawa, R., Koizumi, M., Hirose, A. (2004). Principles of risk assessment for determining the safety of chemical: recent assessment of residual solvents in drugs and di(2-ethylhexyl)phthalate. *Congenit. Anom. Kyoto* 44:51–59.
- Hirata-Koizumi, M., Watari, N., Mukai, D., Imai, T., Hirose, A., Kamata, E., Ema, M. (2007) A 28-day repeated dose toxicity study of ultraviolet absorber 2-(2'-hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole in rats. *Drug Chem. Toxicol.* 30:327–341.
- IARC (International Agency for Research on Cancer). (1995). *Peroxisome Proliferation and its Role in Carcinogenesis (Technical Report no. 24)*. Lyon: IARC Press.
- Knuckles, M. E., Inyang, F., Ramesh, A. (2004). Acute and subchronic oral toxicity of fluoranthene in F-344 rats. *Ecotoxicol. Environ. Saf.* 59:102–108.

- Mann, H. B., Whitney, D. R. (1947). On a test of whether one of two random variables is stochastically larger than the other. *Ann. Math. Stat.* 18:50–60.
- METI (Ministry of Economy, Trade and Industry of Japan). (2006). 2-(2H-1,2,3-Benzotriazole-2-yl)-4,6-di-*tert*-butylphenol, document distributed in Committee on Safety of Chemical Substances, Chemical Substances Council, 30 June 2006. Accessed on April 25, 2007 from <http://www.meti.go.jp/committee/materials/g60705aj.html>.
- MHLW (Ministry of Health, Labour and Welfare of Japan). (2003). 2-(2'-Hydroxy-3',5'-de-*tert*-butylphenyl)benzotriazole. In *Toxicity Testing Reports of Environmental Chemicals (Ministry of Health, Labour and Welfare, ed.)*, Vol. 10. Tokyo: Chemicals Investigation Promoting Council, pp. 215–247.
- MHLW (Ministry of Health, Labour and Welfare of Japan). (2006). 2-(2'-Hydroxy-3',5'-de-*tert*-butylphenyl)benzotriazole. In *Toxicity Testing Reports of Environmental Chemicals (Ministry of Health, Labour and Welfare, ed.)*, Vol. 13, Tokyo: Chemicals Investigation Promoting Council, pp. 187–202.
- OECD (Organization for Economic Co-operation and Development). (1981). Guideline 452, Chronic Toxicity Studies (adopted 12th May 1981), OECD Guidelines for the Testing of Chemicals, section 5, OECD, Paris.
- OECD (Organization for Economic Co-operation and Development). (1998). Principles on Good Laboratory Practice (as revised in 1997), OECD Series on Principles of Good Laboratory Practice and Compliance Monitoring, No. 1, OECD, Paris.
- Steel, R. D. (1959). A multiple comparison rank sum test: treatment versus control. *Biometrics* 15:560–572.
- Tenkazai.com. (2007). Market Trend of Resin Additives “Light Stabilizer.” Accessed on April 25, 2007 from <http://www.tenkazai.com/market.html>.
- Wang, R. H., Schorer-Apelbaum, D., Weinstock, M. (2001). Testosterone mediates sex difference in hypothermia and cholinesterase inhibition by rivastigmine. *Eur. J. Pharmacol.* 433:73–79.

This article was downloaded by:[Ema, Makoto]
On: 28 December 2007
Access Details: [subscription number 789019941]
Publisher: Informa Healthcare
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Drug and Chemical Toxicology

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title~content=t713597244>

A 52-Week Repeated Dose Toxicity Study of Ultraviolet Absorber

2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole in Rats

Mutsuko Hirata-Koizumi ^a; Hidehiro Ogata ^b; Toshio Imai ^a; Akihiko Hirose ^a; Eiichi Kamata ^a; Makoto Ema ^a

^a Biological Safety Research Center, National Institute of Health Sciences, Tokyo, Japan

^b Safety Assessment Laboratory, Panapharm Laboratories Co., Ltd., Kumamoto, Japan

Online Publication Date: 01 January 2008

To cite this Article: Hirata-Koizumi, Mutsuko, Ogata, Hidehiro, Imai, Toshio, Hirose, Akihiko, Kamata, Eiichi and Ema, Makoto (2008) 'A 52-Week Repeated Dose Toxicity Study of Ultraviolet Absorber

2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole in Rats', *Drug and Chemical Toxicology*, 31:1, 81 - 96

To link to this article: DOI: 10.1080/01480540701688758

URL: <http://dx.doi.org/10.1080/01480540701688758>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Gonadal Influence on the Toxicity of 2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole in Rats

Mutsuko Hirata-Koizumi,¹ Takashi Matsuyama,² Toshio Imai,¹
Akihiko Hirose,¹ Eiichi Kamata,¹ and Makoto Ema¹

¹Biological Safety Research Center, National Institute of Health Sciences, Tokyo, Japan

²Shin Nippon Biomedical Laboratories, Ltd., Drug Safety Research Laboratories (SNBL DSR), Kagoshima, Japan

Previously, we showed that susceptibility of male rats to the toxicity of an ultraviolet absorber, 2-(2'-hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole (HDBB), was nearly 25 times higher than that of females. In the current study, we investigated the role of sex steroids in the mediation of the gender-related difference using castrated rats. Male and female castrated CD(SD) rats were given HDBB by gavage at 0, 0.5, 2.5, or 12.5 mg/kg/day for 28 days. No deaths, clinical signs of toxicity, or changes in body weight or food consumption were found at any doses. Blood biochemical changes suggestive of hepatic damage, such as increased levels of aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase, and lactate dehydrogenase, were detected at 12.5 mg/kg/day in males. Absolute and relative liver weight increased at 0.5 mg/kg/day and above in males and at 12.5 mg/kg/day in females. In the liver, histopathological changes, such as nucleolar enlargement, increased mitosis, hypertrophy in hepatocytes, and/or focal necrosis were observed at 0.5 mg/kg/day and above in males, and at 2.5 mg/kg/day and above in females. These findings indicate that castration markedly reduced the gender-related differences in toxicity of HDBB in rats.

Keywords Benzotriazole UV absorber, Castration, Gender-related difference, Rats.

INTRODUCTION

2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole (CAS No. 3846-71-7; HDBB) is an ultraviolet (UV) absorber used in plastic resin products, such as

Address correspondence to Makoto Ema, Division of Risk Assessment, Biological Safety Research Center, National Institute of Health Sciences, 1-18-1, Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan; E-mail: ema@nihs.go.jp

building materials and automobile components (METI, 2006). Previously, we showed a marked difference in the susceptibility of male and female rats to the toxicity of HDBB in 28-day and 52-week repeated oral dose toxicity studies (Hirata-Koizumi et al., 2007; 2008). In the 28-day study, toxic effects in the liver, heart, kidneys, thyroids, and blood were observed. The no observed adverse effect level (NOAEL) for females was 2.5 mg/kg/day based on histopathological changes in the liver and heart detected at 12.5 mg/kg/day. However, the NOAEL for males could not be determined because hepatic changes were noted even at the lowest dose of 0.5 mg/kg/day. In the 52-week study, the NOAEL was concluded to be 0.1 mg/kg/day in males and 2.5 mg/kg/day in females based on histopathological changes in the liver. These findings consistently showed that male rats have a nearly 25 times higher susceptibility to HDBB toxicity than female rats.

Gender-related differences in susceptibility to toxicity have been documented for other substances; for example, a subchronic toxicity study in rats showed that fluoranthene, a polycyclic aromatic hydrocarbon, had greater effects on males than females, especially on the kidneys (Knuckles et al., 2004). In contrast, female rats exhibited greater susceptibility to hypothalamic cholinesterase inhibitory and hypothermic effects of a carbamate cholinesterase inhibitor, rivastigmine (Wang et al., 2001). Such gender-related variations are also reported in humans, mostly for medicines. Examples include the more severe adverse effects, but with greater improvement in response, to antipsychotic drugs such as chlorpromazine and fluspirilene in women (Harris et al., 1995).

Sex hormones are likely to play an important role in gender differences in toxicity responses. In fact, Wang et al. (2001) reported that orchidectomy completely abolished the above-mentioned sex differences in hypothalamic cholinesterase inhibition induced by rivastigmine, and testosterone treatment to gonadectomized males and females decreased the cholinesterase inhibitory effects of rivastigmine; therefore, it is apparent that testosterone interferes with the effects of rivastigmine. On the other hand, estrogen has been shown to act as a dopamine antagonist (Harris et al., 1995), which is considered to contribute, at least in part, to sex differences in response to antipsychotic drugs. The role of sex hormones in differences between sexes in toxicity responses seems to vary from case to case.

In the present study, we performed a repeated dose toxicity study of HDBB using male and female castrated rats to investigate the role of sex steroids in the mediation of sex difference in the susceptibility of rats to the toxicity of HDBB. Administration was conducted in the same way as the previous 28-day study using intact animals (Hirata-Koizumi et al., 2007) for comparison, and effects on the liver and heart, which were principally affected in the previous study of HDBB, were examined.

MATERIALS AND METHODS

This study was performed at Shin Nippon Biomedical Laboratories, Ltd., Drug Safety Research Laboratories (SNBL DSR; Kagoshima, Japan). The experiment was approved by the Institutional Animal Care and Use Committee of SNBL DSR and was performed in accordance with the ethics criteria contained in the bylaws of the Committee of SNBL DSR.

Chemicals

HDBB (Lot no. AY11) was obtained from Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan). The HDBB used in this study was 100% pure and was kept in a light-resistant and airtight container at room temperature. Test solutions were prepared as suspensions in corn oil twice a week and kept cool in a light-resistant and airtight container until dosing. Stability under refrigerated conditions was confirmed for seven days in the previous 28-day repeated dose toxicity study using intact animals (Hirata-Koizumi et al., 2007). All other reagents used in this study were of specific purity grade.

Animals

CrI:CD(SD) rats (SPF, three weeks old) were purchased from Hino Breeding Center, Charles River Laboratories Japan, Inc. (Yokohama, Japan). All animals were maintained in an air-conditioned room at 21.8–22.8°C, with a relative humidity of 45%–55%, a 12-h light/dark cycle, and ventilation with 15 air changes/h. Animals were housed individually in stainless cages suspended over a cage board. A basal diet (CE-2; CLEA Japan, Inc., Tokyo, Japan) and water, which meets the drinking water standard under the Water Works Law of Japan, were provided *ad libitum*.

Male and female rats were castrated under ether anesthesia between five and eight days after purchase. After a two-week acclimation, they were subjected to treatment at six weeks of age. Rats found to be in good health were selected and assigned to four groups of 10 males and 10 females by stratified random sampling based on body weight. One female in the highest dose group was excluded from the present study because remnants of the left ovary were confirmed at necropsy.

Experimental Design

Male and female castrated rats were given HDBB once-daily at 0 (vehicle control), 0.5, 2.5, and 12.5 mg/kg/day by gavage for 28 days. The dosage levels were determined based on the results of our previous 28-day study using intact rats given HDBB by gavage at 0.5, 2.5, 12.5, or 62.5 mg/kg/day, at which adverse effects, mainly on the liver and heart, were found at all doses in males and at

12.5 mg/kg/day and above in females (Hirata-Koizumi et al., 2007). The volume of each dose was adjusted to 10 mL/kg based on the latest body weight.

All animals were observed daily before and one to two hours after dosing for clinical signs of toxicity. Body weight was measured on days 0, 3, 7, 10, 14, 17, 21, 24, and 28 of the dosing period, and food consumption was recorded on days 0, 3, 7, 10, 14, 17, 21, 24, and 27 of the dosing period.

On the day after the last dosing, blood was drawn from the caudal vena cava in the abdomen with a heparin-added syringe under ether anesthesia and centrifuged to obtain plasma. The plasma was examined for biochemical parameters, such as total protein, albumin, glucose, total cholesterol, triglycerides, total bilirubin, urea nitrogen (BUN), creatinine, aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), creatinine phosphokinase, calcium, inorganic phosphorus, sodium, potassium, and chlorine. Following the collection of blood, all animals were euthanized by exsanguination, and the surface of the body, and organs and tissues of the entire body, were examined macroscopically. The liver and heart were then removed and weighed. Both organs were fixed in 10% neutral-buffered formalin, processed routinely for embedding in paraffin, and sections were prepared for staining with hematoxylin and eosin. Histopathological observation was performed for all groups.

Data Analysis

Parametric data, such as body weight, food consumption, blood biochemical parameters, and organ weights, were analyzed by Bartlett's test (Bartlett, 1937) for homogeneity of distribution ($p < 0.05$). When homogeneity was recognized, Dunnett's test (Dunnett, 1964) was conducted to compare control and individual treatment groups ($p < 0.01$ or 0.05). If not homogenous, the data were analyzed using a Dunnett-type mean rank test ($p < 0.01$ or 0.05) (Hollander and Wolfe, 1973).

RESULTS

No deaths or clinical signs of toxicity were found in any groups. There was no significant difference in body weight between the control and HDBB-treated groups (Fig. 1). Food consumption was also not significantly changed, except for a transient increase on day 21 of the administration period at 12.5 mg/kg/day and on day 27 of the administration period at 2.5 mg/kg/day in males (data not shown).

Blood biochemical examination revealed significant increases in the level of albumin at 0.5 mg/kg/day and above in males and at 2.5 mg/kg/day and above in females, total protein at 0.5 mg/kg/day and above in females, glucose at 12.5 mg/kg/day in males, and BUN at 12.5 mg/kg/day in both sexes (Table 1).