

Table 3-1. Effects of maternal administration of BPA on neurotransmitter contents of 6-wk-old female offspring (medulla oblongata and cerebellum)


		NE	DA	DOPAC	HVA	DOPAC/DA	HVA/DA
Medulla oblongata	Control	100 ± 5.9	100 ± 6.3	100 ± 5.0	100 ± 8.7	100 ± 6.1	100 ± 5.3
	4 mg/kg	96.1 ± 11.2	96.2 ± 11.9	88.8 ± 9.1	97.7 ± 8.2	92.2 ± 3.1	100.5 ± 13.9
	40 mg/kg	91.9 ± 3.2	105.6 ± 13.8	91.6 ± 13.2	96.2 ± 7.5	85.4 ± 3.8	92.0 ± 14.3
	A100	3.38	0.492	0.119	0.281	0.244	0.600
Cerebellum	Control	100 ± 4.4	100 ± 5.4	100 ± 18.2	100 ± 12.2	100 ± 19.1	100 ± 16.6
	4 mg/kg	104.4 ± 10.8	107.9 ± 10.4	69.3 ± 8.6	102.5 ± 7.4	64.7 ± 7.8	92.9 ± 4.5
	40 mg/kg	97.3 ± 3.5	142.5 ± 25.9	86.4 ± 16.6	118.0 ± 14.3	65.8 ± 7.8	92.9 ± 18.8
	A100	1.23	0.0636	0.0204	0.181	0.307	2.94

Results are shown as means ± SEM (%). A100, Absolute values for 100% (N = 4–5 for medulla oblongata and N = 3–5 for cerebellum, nmoles/g tissue).

Table 3-2. Effects of maternal administration of BPA on neurotransmitter contents of 6-wk-old female offspring (medulla oblongata and cerebellum)

		5HT	5HIAA	5HIAA/5HT	ACh	Ch	ACh/Ch
Medulla oblongata	Control	100 ± 12.5	100 ± 14.1	100 ± 9.8	100 ± 7.5	100 ± 17.8	100 ± 13.1
	4 mg/kg	115.3 ± 21.8	116.4 ± 11.1	121.2 ± 17.0	90.4 ± 8.6	182.0 ± 30.0	50.3* ± 10.8
	40 mg/kg	63.1 ± 5.2	95.3 ± 7.0	172.6* ± 18.3	90.8 ± 5.5	147.5 ± 34.6	68.1 ± 14.6
	A100	31.5	7.10	0.202	19.3	30.5	0.693
Cerebellum	Control	100 ± 29.3	100 ± 26.8	100 ± 25.1	100 ± 13.0	100 ± 21.3	100 ± 16.6
	4 mg/kg	90.4 ± 4.9	77.7 ± 17.9	86.3 ± 21.9	108.8 ± 16.9	152.0 ± 26.0	66.5 ± 11.6
	40 mg/kg	114.5 ± 13.9	103.9 ± 15.3	93.3 ± 15.7	97.6 ± 14.3	128.8 ± 17.0	70.9 ± 13.1
	A100	1.35	0.251	0.188	3.47	12.3	0.326

Results are shown as means ± SEM (%). A100, Absolute values for 100% (N = 4–5, nmoles/g tissue). *: p < 0.05 by Dunnett's multiple t-test.

**Fig. 3-1. Effects of perinatal administration of BPA on the neurotransmitter contents of forebrain in 9-wk-old female offspring.**

Results are shown as means ± SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 0.722 for NE, 6.44 for DA, 0.945 for DOPAC, 1.32 for HVA, 12.0 for 5HT, and 1.35 for 5HIAA; absolute values of ratios for 100% were as follows: 0.148 for DOPAC/DA, 0.210 for HVA/DA, and 0.111 for 5HIAA/5HT. N = 5. *: p < 0.05 by Dunnett's multiple t-test.




Fig. 3-2. Effects of perinatal administration of BPA on the neurotransmitter contents of hindbrain in 9-wk-old female offspring.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 1.65 for NE, 1.64 for DA, 0.203 for DOPAC, 0.570 for HVA, 9.58 for 5HT, and 3.87 for SHIAA; absolute values of ratios for 100% were as follows: 0.126 for DOPAC/DA, 0.361 for HVA/DA, and 0.420 for SHIAA/5HT. $N = 5$.




Fig. 3-3. Effects of perinatal administration of BPA on the neurotransmitter contents of medulla oblongata in 9-wk-old female offspring.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 1.98 for NE, 0.193 for DA, 0.0881 for DOPAC, 0.695 for HVA, 2.47 for 5HT, and 3.30 for SHIAA; absolute values of ratios for 100% were as follows: 0.479 for DOPAC/DA, 3.76 for HVA/DA, and 1.41 for SHIAA/5HT. $N = 5$. *: $p < 0.05$ by Dunnett's multiple t -test.

Discussion

The reproductive effects of BPA have been studied in detail²³, but little is understood about the effects of BPA on the nervous system. Both positive and negative effects of BPA on the reproductive and other functions of offspring after perinatal exposure have been reported^{7, 10, 24, 25}. In our present study, changes in monoamine and metabolite levels due to BPA treatment were not observed in the brains of 1-wk-old female rat pups. Increases in DA metabolite, DOPAC and HVA, were observed in the female rat pups at 3 wk of age, and these increases were statistically significant in the

forebrain, DOPAC in the 40 mg/kg group and HVA in the 4 mg/kg group. HVA contents in the hindbrain and medulla oblongata of the 40 mg/kg group were greater than the control. The HVA/DA ratio was significantly high in the frontal cortex of the 4 mg/kg group and it was also high in the occipital cortex and medulla oblongata of the 40 mg/kg group, but the differences were not significant. These results mean that the turnover of DA was accelerated in the BPA-treated groups, and suggest that the release of DA from nerve endings was increased in these groups. Significant increases in 5HT and SHIAA were observed in the forebrain, and in 5HT in the medulla oblongata of the 4 mg/kg group. The 5HT level




Fig. 3-4. Effects of perinatal administration of BPA on the neurotransmitter contents of cerebellum in 9-wk-old female offspring.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 0.635 for NE, 0.0355 for DA, 0.411 for HVA, 0.346 for 5HT, and 0.464 for SHIAA; absolute values of ratios for 100% were as follows: 11.7 for HVA/DA, and 1.36 for SHIAA/5HT. N = 5. *: $p < 0.05$ by Dunnett's multiple *t*-test.




Fig. 4-1. Effects of perinatal administration of BPA on the neurotransmitter contents of frontal cortex in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 1.65 for NE, 4.74 for DA, 1.11 for DOPAC, 0.771 for HVA, 13.8 for 5HT, and 1.70 for SHIAA; absolute values of ratios for 100% were as follows: 0.232 for DOPAC/DA, 0.166 for HVA/DA, and 0.123 for SHIAA/5HT. N = 5-6. *: $p < 0.05$ by Dunnett's multiple *t*-test.

in the medulla oblongata of the 40 mg/kg group was higher than the control; however, no changes were observed in 5HT and SHIAA in the forebrain of the 40 mg/kg group. The 5HT and SHIAA levels in the hindbrain of the BPA-treated groups were less than control, although the degree of the decrease was small. Though 5HT and SHIAA increased in the forebrain and medulla oblongata, changes in 5HT and SHIAA seemed to be dependent on brain area and dose of BPA. In 6-wk-old offspring, increases in Ch levels were observed in all of the eight brain areas of the BPA-treated groups, but there were no accompanying changes in ACh levels. Synthesis or uptake into the synaptosome of Ch seems

to have been accelerated in the BPA-treated groups. Among changes in catecholamine, serotonin, and their metabolites, DA and DOPAC in the hippocampus, and DA in the cerebellum increased in the 40 mg/kg group by 40 to 50% compared with the control. In 9-wk-old offspring, significant changes in monoamines and metabolites were observed in the forebrain, medulla oblongata, and cerebellum of the BPA-treated groups, however, these changes were scattered and not great.

BPA treatment affected the monoamine and metabolite contents of the brain of dams. Large and dose dependent increases in HVA and in the HVA/DA ratio occurred in the




Fig. 4-2. Effects of perinatal administration of BPA on the neurotransmitter contents of occipital cortex in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 1.59 for NE, 3.44 for DA, 0.609 for DOPAC, 0.0749 for HVA, 2.83 for 5HT, and 1.59 for 5HIAA; absolute values of ratios for 100% were as follows: 0.179 for DOPAC/DA, 0.0221 for HVA/DA, and 0.576 for 5HIAA/5HT. N = 5–6.




Fig. 4-3. Effects of perinatal administration of BPA on the neurotransmitter contents of hippocampus in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 1.37 for NE, 0.123 for DA, 0.0673 for DOPAC, 2.06 for 5HT, and 1.97 for 5HIAA; absolute values of ratios for 100% were as follows: 0.564 for DOPAC/DA, and 1.02 for 5HIAA/5HT. N = 5–6. **: $p < 0.01$ by Dunnett's multiple *t*-test.

occipital cortex of the BPA-treated groups, although they were not statistically significant. A dose dependent increase in the HVA/DA ratio was also observed in the frontal cortex. These findings suggest that DA turnover was accelerated in specific brain areas following BPA treatment. 5HT and 5HIAA increased in the frontal cortex of the 4 mg/kg group and 5HIAA increased in the striatum of the 40 mg/kg group. Everitt *et al.* reported that the serotonin turnover was accelerated in female rats by estrogen administration²⁶⁾. According to Shimizu and Bray, estradiol administration increased the ratio of DOPAC/DA but decreased the 5HIAA/5HT ratio in the nucleus accumbens of ovariectomized rats

when measured by microdialysis²⁷⁾. Our findings are consistent with the changes found by Everitt *et al.* following estrogen treatment, because 5HIAA in the frontal cortex and striatum increased in BPA-treated dams. BPA possesses very weak estrogenic activity and the 5HIAA increase observed in our experiment may be related to the estrogenic activity of BPA. Our experimental conditions were very different from those of Shimizu and Bray, therefore, it is difficult to compare our findings with their results. They measured extracellular neurotransmitters and metabolites in a microdialysis study, whereas, we measured them following the homogenization of the brain, in which both intracellular




Fig. 4-4. Effects of perinatal administration of BPA on the neurotransmitter contents of striatum in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 0.669 for NE, 48.2 for DA, 7.16 for DOPAC, 3.75 for HVA, 1.39 for 5HT, and 2.05 for SHIAA; absolute values of ratios for 100% were as follows: 0.150 for DOPAC/DA, 0.0819 for HVA/DA, and 1.85 for SHIAA/5HT. N = 5-6. *: $p < 0.05$ by Dunnett's multiple *t*-test.




Fig. 4-5. Effects of perinatal administration of BPA on the neurotransmitter contents of midbrain in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 6.87 for NE, 4.75 for DA, 1.44 for DOPAC, 3.11 for 5HT, and 3.54 for SHIAA; absolute values of ratios for 100% were as follows: 0.304 for DOPAC/DA, and 1.41 for SHIAA/5HT. N = 4-6.

and extracellular substances were included. At present, it is not clear whether the changes in monoamine turnover observed in the dams in our experiments were due to the estrogenic activity of BPA. An effect of BPA on prolactin secretion has been reported^{28, 29}. DA inhibits the secretion of prolactin in the anterior pituitary gland. Male rats were exposed to BPA from postnatal days 22 to 32²⁹. During this period, BPA stimulated prolactin secretion in the same manner as pimozide (a dopamine antagonist) and 17 β -estradiol. Steinmetz *et al.* reported that BPA induces hyperprolactinemia in F344 rats with an efficacy similar to that of estradiol²⁸. On the assumption that such effects of BPA on prolactin secretion are via the inhibition of

dopaminergic activity in the anterior pituitary gland, BPA would inhibit the activity of DA neurons. In our experiment, DA in the hippocampus significantly increased in the 4 mg/kg group. HVA levels and HVA/DA ratios in the occipital cortex of dams treated with BPA at 4 or 40 mg/kg were higher than the control, although without statistical significance. These results from female CD (SD) IGS rats are inconsistent with previous findings. This may be due to sex differences. Alternatively, the stimulation of prolactin secretion by BPA might be due to activities of BPA other than dopamine-mediated action. Changes in DA, 5HT, and their metabolites were observed in some brain regions of dams dosed with 4 or 40 mg/kg of BPA in our present study. Although the




Fig. 4-6. Effects of perinatal administration of BPA on the neurotransmitter contents of hypothalamus in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 9.56 for NE, 2.07 for DA, 0.498 for DOPAC, 0.192 for HVA, 4.91 for 5HT, and 2.55 for SHIAA; absolute values of ratios for 100% were as follows: 0.240 for DOPAC/DA, 0.113 for HVA/DA, and 0.497 for SHIAA/5HT. N = 4–6.




Fig. 4-7. Effects of perinatal administration of BPA on the neurotransmitter contents of medulla oblongata in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 2.64 for NE, 0.463 for DA, 0.232 for DOPAC, 2.36 for 5HT, and 2.62 for SHIAA; absolute values of ratios for 100% were as follows: 0.502 for DOPAC/DA, and 1.11 for SHIAA/5HT. N = 5–6.

effects of BPA on GABA (A) and nicotinic receptors have been reported^{30, 31}), those of BPA on dopaminergic and serotonergic neurons have not been described. Our results suggest that the metabolism of DA and 5HT was accelerated in BPA-treated female rats. We postulate that BPA may affect some DA- and 5HT-related brain functions.

The assay of brain substances of the male offspring sacrificed in the same series of experiments is now underway in our laboratory. Kubo *et al.* reported that sexual differentiation of the brain locus coeruleus is disrupted in rats perinatally exposed to BPA³²). In that study, maternal rats received BPA at 1.5 mg/kg per day. We found that levels of NE in the forebrain of 9-wk-old offspring were dose-

dependently increased and reached significance in the 40 mg/kg group; a similar increase was also observed in the hindbrain. Cell bodies of NE neurons are dense in the locus coeruleus and NE neurons might be altered by BPA in this disruption of sexual differentiation, although the size of the locus coeruleus is much smaller than the forebrain and hindbrain. According to Farabollini *et al.*, the maternal administration of BPA during the critical period of fetal brain organization produces different effects on the behavior of male and female offspring rats³³). A comparison of data regarding neurotransmitters obtained from female and male offspring may explain such sexually differentiated behavior effects of BPA.




Fig. 4-8. Effects of perinatal administration of BPA on the neurotransmitter contents of cerebellum in maternal rats.

Results are shown as means \pm SEM (%). Absolute values (nmoles/g tissue) for 100% were as follows: 0.785 for NE, 0.0527 for DA, 0.0520 for DOPAC, 0.0447 for HVA, 0.455 for 5HT, and 0.366 for 5HIAA; absolute values of ratios for 100% were as follows: 0.987 for DOPAC/DA, 0.896 for HVA/DA, and 0.939 for 5HIAA/5HT. N = 5–6.

We found that dams given BPA at 400 mg/kg weighed significantly less than controls¹². The 40 mg/kg group weighed somewhat less than controls, but BPA at 4 mg/kg did not affect the body weight of dams. The weight of the 40 mg/kg group recovered to the control level during lactation. The body weight of female offspring did not statistically differ between control and BPA-treated groups at 1 to 9 wk of age. No differences were statistically significant in the weights of the liver and kidneys among groups. These results show that the neurochemical alterations in the brains of dams and offspring after BPA exposure were not caused by differences in somatic growth. Anogenital distances in female offspring were not significantly affected by BPA at 1, 3 or 9 wk of age as observed in the same rats in this study¹². Though the anogenital distance is not always sensitive to the reproductive effects of chemicals, neurotransmitters in the brain might be more sensitive to BPA than reproductive organ sensitivity to the estrogenic action of BPA. At present we have no data to explain the reason why the changes in monoamines and metabolites occurred in pups as well as dams. These changes were observed in specific brain areas, except the increase in Ch. Levels of Ch were higher than control in all of the eight brain areas of 6-wk-old pups of the 4 and 40 mg/kg groups. Unfortunately, we have no data for Ch in pups at ages other than 6 wk.

Acknowledgements

The authors are grateful to Ms. N. Ebara for excellent

technical assistance and Ms. J. Kuwahara for statistical treatment. This investigation is part of a contract for research study with the Ministry of Health, Labour and Welfare, and is supported in part by the Ministry of Environment, Japan.

References

- 1) Krishnan AV, Stathis P, Permuth SF, Tokes L, Feldman D (1993) Bisphenol-A: An estrogenic substance is released from polycarbonate flasks during autoclaving. *Endocrinol* **132**, 2279–86.
- 2) Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT (1998) Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinol* **139**, 4252–63.
- 3) Nagel SC, vom Saal FS, Thayer KA, Dhar MG, Boehler M, Welshons WV (1997) Relative binding affinity-serum modified access (RBA-SMA) assay predicts the relative in vivo bioactivity of the xenoestrogens bisphenol A and octylphenol. *Environ Health Perspect* **105**, 70–6.
- 4) Howdeshell KL, Hotchkiss AK, Thayer KA, Vandenberg JG, vom Saal FS (1999) Exposure to bisphenol A advances puberty. *Nature* **401**, 763–4.
- 5) Welshons WV, Nagel SC, Thayer KA, Judy BM, vom Saal FS (1999) Low-dose bioactivity of xenoestrogens in animals: fetal exposure to low doses of methoxychlor and other xenoestrogens increases adult prostate size in mice. *Toxicol Ind Health* **15**, 12–25.
- 6) Honma S, Suzuki A, Buchanan DL, Katsu Y, Watanabe H, Iguchi T (2002) Low dose effect of in utero exposure to bisphenol A and diethylstilbestrol on female mouse reproduction. *Reprod Toxicol* **16**, 117–22.
- 7) Schonfelder G, Flick B, Mayr E, Talsness C, Paul M, Chahoud

- I (2002) In utero exposure to low doses of bisphenol A lead to long-term deleterious effects in the vagina. *Neoplasia* **4**, 98–102.
- 8) Jacobson JL, Jacobson SW (1996) Intellectual impairment in children exposed to polychlorinated biphenyls in utero. *N Engl J Med* **335**, 783–9.
 - 9) Faroon O, Jones D, de Rosa C (2001) Effects of polychlorinated biphenyls on the nervous system. *Toxicol Ind Health* **16**, 305–33.
 - 10) vom Saal FS, Cooke PS, Buchanan DL, Palanza P, Thayer KA, Nagel SC, Parmigiani S, Welshons WV (1998) A physiologically based approach to the study of bisphenol A, and other estrogenic chemicals on the size of reproductive organs, daily sperm production, and behavior. *Toxicol Ind Health* **14**, 239–60.
 - 11) Laws SC, Carey SA, Ferrell JM, Bodman GJ, Cooper RL (2000) Estrogenic activity of octylphenol, nonylphenol, bisphenol A and methoxychlor in rats. *Toxicol Sci* **54**, 154–67.
 - 12) Kobayashi K, Miyagawa M, Wang RS, Sekiguchi S, Suda M, Honma T (2002) Effects of in utero and lactational exposure to bisphenol A on somatic growth and anogenital distance in F1 rat offspring. *Ind Health* **40**, 375–81.
 - 13) Watanabe S, Wang RS, Miyagawa M, Kobayashi K, Suda M, Sekiguchi S, Honma T (2003) Imbalance of testosterone level in male offspring rats perinatally exposed to bisphenol A. *Ind Health* **41**, 338–41.
 - 14) Cooper JR, Bloom FE, Roth RH (2003) *The Biochemical Basis of Neuropharmacology*. Oxford University Press, Oxford.
 - 15) Honma T (1992) Brain microdialysis study of the effects of hazardous chemicals on the central nervous system. 1. Changes in monoamine metabolites induced by cerebral methyl bromide administration measured by two-probe microdialysis (TPMD) method. *Ind Health* **30**, 47–60.
 - 16) Tsuga H, Haga T, Honma T (2002) Effects of toluene exposure on signal transduction: toluene reduced the signaling via stimulation of human muscarinic acetylcholine receptor m2 subtypes in CHO cells. *Jpn J Pharmacol* **89**, 282–9.
 - 17) Kwon S, Stedman DB, Elswick RC, Cattey RC, Welsh F (2000) Pubertal development and reproductive functions of Crl: CD BR Sprague-Dawley rats exposed to bisphenol A during prenatal and postnatal development. *Toxicol Sci* **55**, 399–406.
 - 18) Tsuga H, Honma T (2000) Effects of short-term toluene exposure on ligand binding to muscarinic acetylcholine receptors in the rat frontal cortex and hippocampus. *Neurotoxicol Teratol* **22**, 603–6.
 - 19) Glowinski J, Iversen LL (1966) Regional studies of catecholamines in the rat brain. I. The disposition of ³Hnorepinephrine, ³Hdopamine and ³Hdopa in various regions of the brain. *J Neurochem* **13**, 655–69.
 - 20) Honma T, Miyagawa M, Sato M (1987) Methyl bromide alters catecholamine and metabolites concentrations in rat brain. *Neurotoxicol Teratol* **9**, 369–75.
 - 21) Honma T, Miyagawa M, Sato M (1991) Inhibition of tyrosine hydroxylase activity by methyl bromide exposure. *Neurotoxicol Teratol* **13**, 1–4.
 - 22) Honma T, Suda M (2004) Brain microdialysis study of the effects of hazardous chemicals on the central nervous system. 2. Toluene exposure and cerebral acetylcholine. *Ind Health* **42**, 336–47.
 - 23) Witorsch RJ (2002) Low-dose in utero effects of xenoestrogens in mice and their relevance to humans: an analytical review of the literature. *Food Chem Toxicol* **40**, 905–12.
 - 24) Tinwell H, Haseman J, Lefevre PA, Wallis N, Ashby J (2002) Normal sexual development of two strains of rat exposed in utero to low doses of bisphenol A. *Toxicol Sci* **68**, 339–48.
 - 25) Yoshino H, Ichihara T, Kawabe M, Imai N, Hagiwara A, Asamoto M, Shirai T (2002) Lack of significant alteration in the prostate or testis of F344 rat offspring after transplacental and lactational exposure to bisphenol A. *J Toxicol Sci* **27**, 433–9.
 - 26) Everitt BJ, Fuxe K, Hokfelt FT, Jonsson G (1975) Role of monoamines in the control by hormones of sexual receptivity in the female rat. *J Comp Physiol Psychol* **89**, 556–72.
 - 27) Shimizu H, Bray GA (1993) Effects of castration, estrogen replacement and estrus cycle on monoamine metabolism in the nucleus accumbens, measured by microdialysis. *Brain Res* **621**, 200–6.
 - 28) Steinmetz R, Brown NG, Allen DL, Bigsby RM, Ben-Jonathan N (1997) The environmental estrogen bisphenol A stimulates prolactin release in vitro and in vivo. *Endocrinol* **138**, 1780–6.
 - 29) Stoker TE, Robinette CL, Britt BH, Laws SC, Cooper RL (1999) Prepubertal exposure to compounds that increase prolactin secretion in the male rat: effects on the adult prostate. *Biol Reprod* **61**, 1636–43.
 - 30) Aoshima H, Hossain SJ, Imamura H, Shingai R (2001) Effects of bisphenol A and its derivatives on the response of GABA (A) receptors expressed in *Xenopus* oocytes. *Biosci Biotechnol Biochem* **65**, 2070–7.
 - 31) Nakazawa K, Ohno Y (2001) Modulation by estrogens and xenoestrogens of recombinant human neuronal nicotinic receptors. *Eur J Pharmacol* **430**, 175–83.
 - 32) Kubo K, Arai O, Ogata R, Omura M, Hori T, Aou S (2001) Exposure to bisphenol A during the fetal and suckling periods disrupts sexual differentiation of the locus coeruleus and of behavior in the rat. *Neurosci Lett* **304**, 73–6.
 - 33) Farabolini F, Porrini S, Dessi-Fulgherit F (1999) Perinatal exposure to the estrogenic pollutant bisphenol A affects behavior in male and female rats. *Pharmacol Biochem Behav* **64**, 687–94.

Effects of in Utero and Lactational Exposure to Di(2-ethylhexyl)phthalate on Somatic and Physical Development in Rat Offspring

Kenichi KOBAYASHI*, Muneyuki MIYAGAWA, Rui-Sheng WANG, Megumi SUDA, Soichiro SEKIGUCHI and Takeshi HONMA

National Institute of Occupational Safety and Health, 21-1, Nagao 6-chome, Tama-Ku, Kawasaki 214-8585, Japan

Received March 27, 2006 and accepted July 3, 2006

Abstract: Di(2-ethylhexyl)phthalate (DEHP) has been reported to act as an antiandrogen and to affect the reproductive organs and accessory genital glands. Thus, to assess the reproductive toxicity of DEHP it is important to examine both its adverse effects on the development of offspring following maternal exposure and its effects on sexual function and fertility. In the present study, we examined whether in utero and lactational exposure to DEHP affects postnatal somatic growth of offspring in the rat. Pregnant females were orally administered various doses of DEHP (0, 25, 100 or 400 mg/kg body weight/day) from gestational day (GD) 6 through postnatal day (PND) 20. There were no significant changes in body weight, body length, tail length, or the weight of individual organs between the control and DEHP-treated groups. Somatic hormonal parameters were the same for all DEHP doses. These findings suggest that in utero and lactational exposure to various concentrations of DEHP has very little effect on postnatal development or endocrine and physical status of male and female rat offspring under the experimental conditions of the present study.

Key words: Di(2-ethylhexyl)phthalate, Postnatal development, In utero and lactational exposure, Offspring, Rat

Introduction

To date, several compounds have been suspected of exerting endocrine-disturbing effects even at ultra-low concentrations. Phthalates have been produced and used in the manufacture of chemically derived materials and products. Di(2-ethylhexyl)phthalate (DEHP) has been most widely used in polyvinyl chloride to impart structural flexibility, and it is used as a plasticizer in products such as food packaging, children's products (toys and crib bumpers) and medical devices. Significantly, DEHP has been detected in plasma samples¹⁾. Mono(2-ethylhexyl)phthalate (MEHP), which is an active and the predominant DEHP metabolite, is also considered as a testicular toxicant²⁾. It has been estimated that mean DEHP intake is 8.2 $\mu\text{g}/\text{kg}$ body weight per day for adults³⁾. During recent years, DEHP has been

excluded from many products to avoid consumer exposure. However, recent heightened public concerns about environmental exposure to high concentrations of DEHP have raised new questions about its possible occupational and medical health hazards.

Developmental toxicity studies of DEHP have been conducted in laboratory mice⁴⁻⁸⁾ and rats⁸⁻¹⁰⁾. These reports suggest that in utero exposure to high doses of DEHP induces embryotoxicity and/or teratogenicity. Animal reproductive toxicity studies of DEHP have also been reported. In a study of adult male rats, testicular defects such as atrophy of the seminiferous tubules, loss of spermatogenesis and vacuolation of Sertoli cells were observed after 90 days of dietary exposure to DEHP at 500 and 5,000 ppm (equivalent to 37.6 and 375.2 mg/kg/day, respectively)¹¹⁾. Perinatal exposure to DEHP in rats from gestational day (GD) 14 through postnatal day (PND) 3 reduced anogenital distance, testis weight or the weight of androgen-dependent tissues¹²⁾.

*To whom correspondence should be addressed.

Dietary exposure of adult male rats given 0, 320, 1,250, 5,000, and 20,000 ppm DEHP (equivalent to 0, 17.5, 69.2, 284.1 and 1156.4 mg/kg/day, respectively) for 60 days, when mated with untreated adult females, did not affect the rate of neonatal death, initial pup weight or growth (up to PND 7), whereas the average litter size decreased in rats fed 20,000 ppm DEHP¹³. Inhalation exposure of adult male Wistar rats to 25 mg/m³ for 6 h/day for 8 wk increased plasma testosterone level and seminal vesicle weight in a dose-dependent manner¹⁴. In a study of adult female rats, DEHP induced prolonged estrous cycles and suppressed plasma concentrations of estradiol and subsequent ovulation¹⁵.

Several studies have shown that in utero and lactational exposure to DEHP leads to abnormalities in the hypothalamus-pituitary-testicular axis. Sprague-Dawley rats were orally dosed with DEHP (0–1,500 mg/kg/day) from GD 3 through PND 21, and dose-related effects in the male offspring included several parameters involved in sexual development¹⁶. Oral exposure of pregnant female Long-Evans rats to 100 mg/kg/day DEHP from GD 12–21 induced significantly increased levels of testosterone and luteinizing hormone in male offspring on PND 21 and PND 35, but by PND 90 the levels were comparable between treated and untreated animals¹⁷, indicating that the magnitude of DEHP toxicity on reproductive function is influenced by the stage of development.

Thus, DEHP toxicity studies in laboratory animals have focused on embryotoxicity, teratogenicity and reproductive toxicological effects in addition to some developmental effects in the early postnatal period, yet extensive toxicity information for long-term development after DEHP exposure is still lacking. The purpose of the present study was to evaluate postnatal growth and physical development following in utero and lactational exposure to DEHP in male and female rat offspring until the post-pubertal period. We examined the effects of DEHP on pubertal development, and doses of DEHP were chosen based on the levels that caused no overt maternal toxicity. Additionally, the exposure period was extended to examine the effects of lactational exposure in addition to the effects of in utero exposure, to complement previous studies^{4–10}. Thus, we administered several doses of DEHP orally by gavage to pregnant rats using an experimental schedule identical to one used previously¹⁸, and we examined the effects on postnatal somatic and organ growth, as assessed by body weight, body length, tail length and main organ weights, including reproductive organs, in male and female offspring. In addition, to better assess physical status following DEHP exposure, we evaluated the levels of several plasma

hormonal landmarks with regard to postnatal somatic growth.

Materials and Methods

Chemicals and experimental animals

DEHP (purity >99.9%, Cat# 289-10442) and corn oil were obtained from Wako Pure Chemical Industries, Ltd., Osaka, Japan. A total of 52 pregnant (GD 3) female rats (Crj: CD (SD) IGS strain, 9 wk of age) were purchased from Charles River Japan, Inc. (Tsukuba, Japan). The presence of a copulatory plug defined GD 0. They were acclimated on GD 3–6 and housed individually in plastic cages with sterilized wood chips (Soft chip, Japan Sic Inc., Shizuoka, Japan) for bedding and were maintained under controlled temperature ($23 \pm 1^\circ\text{C}$) and humidity ($55 \pm 5\%$) and with a 12-h light-dark cycle (08:00–20:00) throughout the study. A standard laboratory diet (CE-2, Clea Japan, Inc., Tokyo, Japan) and drinking water were available ad libitum.

Dose range-finding evaluation

Dams were randomly divided into five groups (four pregnant females per group). The DEHP-exposed groups were orally administered 500, 1,000, 1,500, or 2,000 mg DEHP/kg/day in corn oil vehicle (10 ml/kg of body weight); DEHP was given between 08:30 and 09:30 for five consecutive days each week (Monday–Friday) from GD 6 through GD 20, and the control group was given the same amount of corn oil during the same period. During the exposure period, we recorded maternal body weights and noted any clinical signs or abnormal behavior that may have resulted from toxic effects. These results were used to determine the range of the DEHP dose for the main study.

Main study

Dams were randomly divided into four groups (eight pregnant females per group) and weighed once daily from GD 3 through PND 20 (except for GD 4 and 5). The DEHP-exposed groups were orally administered 25, 100 or 400 mg DEHP/kg/day in corn oil vehicle (10 ml/kg of body weight); DEHP was given between 08:30 and 09:30 from GD 6 through PND 20, and the control group was given the same amount of corn oil during the same period. Maternal data were recorded as described above. For each dam, the gestational duration was recorded, and weight gain during gestation and lactation was measured. Dams were checked for birth until 10:00 on each day; the day on which pups were first observed was designated PND 0. The number of

live births and the weight of each live pup on PND 1 were recorded. The litter size was standardized to 10 (five males and five females when possible) between 10:00 and 11:00 on PND 7 (1 wk of age). Litters with a total of nine or fewer pups were not culled regardless of the sex ratio. Culled pups were used for the analysis at 1 wk of age. On PND 21, the remaining offspring were weaned, and thereafter males and females were housed in separate stainless steel cages by litter. Body weights were recorded with an electric balance (Shimadzu, Kyoto, Japan). Body length and tail length (millimeters) were measured with a digital caliper (Mitutoyo, Kanagawa, Japan). The nose-anus length was considered the body length. One male and one female offspring from each dam were dissected at 3 and 9 wk of age when possible. While the rat was under ether anesthesia, liver, kidneys and testes, prostate and seminal vesicles or ovaries and uterus were carefully removed and weighed.

Hormone determinations

For hormone determinations, blood samples were collected from the postcaval vein following euthanasia by ether inhalation at 9 wk of age. Plasma samples were obtained by centrifugation at 4°C and stored at -20°C until the analysis. Concentrations of the plasma thyroid hormones thyroxine (T₄) and tri-iodothyronine (T₃) were determined by a time-resolved fluoroimmunoassay (DELFLIA T₄ Reagents and DELFLIA T₃ Reagents, respectively, PerkinElmer Life and Analytical Sciences, Inc., MA, USA). Plasma growth hormone (GH) concentrations were determined by enzyme immunoassay (EIA) (Rat GH EIA Biotrak system, GE Healthcare Bio-Sciences Corp., NJ, USA). Plasma insulin-like growth factor-I (IGF-I) concentrations were also measured by EIA (ACTIVE mouse/rat IGF-I EIA kit, Diagnostic Systems Laboratories, Inc., TX, USA). Time-resolved fluorescence and absorbance were measured by a multilabel counter (VICTOR², PerkinElmer Life and Analytical Sciences, Inc.). All hormones were assayed according to the manufacturer's instructions.

Statistical analysis

The differences from the corresponding control group were statistically analyzed by an analysis of variance followed by Dunnett's test (significance at $p < 0.05$).

Results

Dose range-finding evaluation

In the 1,000 mg/kg/day and higher DEHP groups, maternal toxicity was clearly manifested as greatly suppressed weight

gain during gestation, which led us to discontinue subsequent dosing by GD 17 of this preliminary study. In the 500 mg/kg/day group, mean body weights decreased slightly at later stages of gestation compared with the control group (data not shown). Based on these observations, we set the highest dose at 400 mg/kg/day to exclude the influence of maternal toxicity and observe the effect of DEHP on the offspring. The lowest dose and the middle dose were set at 25 mg/kg/day and 100 mg/kg/day, respectively.

Main study

Dams

Table 1 shows the number of dams and their offspring used for examinations in each group. Weight gain did not differ between dams from the control group and the DEHP groups from GD 6 through GD 21. In the 400 mg/kg/day group, one dam was found dead on GD 23, and thus the dam was excluded from the analysis. No significant differences were observed between the control group and the DEHP groups with regard to gestational duration or the number of live births per litter on PND 1.

Figure 1 shows maternal body changes during gestation (left panel) and lactation (right panel). There were no statistically significant differences among groups with regard to maternal body weight during the gestation and lactation periods, although the 25 mg/kg/day group showed a transient but not significant weight reduction during early lactation.

Offspring

The number of offspring examined is shown in Table 2. In male and female offspring, there were no statistically significant differences in body weight, body length or tail length between the control and DEHP-exposed groups at 1, 3 or 9 wk of age (Figs. 2, 3 and 4). There were no statistically significant effects on liver or kidney weights in males or females at 1, 3 or 9 wk of age (Table 3, 4). In male offspring, testis weights did not differ among the control group and DEHP groups at 3 or 9 wk of age (Table 3). Prostate and seminal vesicle weights did not differ among the control group and DEHP groups at 9 wk of age (Table 3). In female offspring, ovary and uterus weights did not differ among the groups at 3 or 9 wk of age (Table 4).

Physical status of offspring

In male offspring, no statistically significant differences in plasma concentrations of T₄, T₃, GH or IGF-I were observed among the control group and the DEHP groups at 9 wk of age (Table 5). In female offspring, no statistically significant differences in plasma concentrations of T₄, T₃,

Table 1. Dams and litter data

	DEHP dose (mg/kg/day)			
	0	25	100	400
Females (n)	8	8	8	8
Pregnant females (n)	8	8	8	8
Dam weight gain (GD 6-21)	130 ± 7 ^a	127 ± 5	135 ± 4	133 ± 5
Gestational period (days)	21.1 ± 0.1	21.4 ± 0.2	21.3 ± 0.2	21.3 ± 0.2
Live births/litter on PND 1	11.8 ± 0.7	13.6 ± 0.6	13.5 ± 0.5	11.7 ± 0.5 (7) ^b

^aValues are mean ± SEM.

^bThe number in parentheses represents dams per dose group. One dam was found dead on GD 23, and thus the dam was excluded from the analysis.




Fig. 1. Effects of exposure to di(2-ethylhexyl)phthalate (DEHP) on maternal body weight during gestation (left panel) and lactation (right panel).
Each point represents the mean.

GH or IGF-I were observed between the control group and the DEHP groups at 9 wk of age (Table 6).

Discussion

In recent years, the issue of endocrine-disrupting chemicals has been the topic of much discussion. Nagel *et al.*¹⁹⁾ and vom Saal *et al.*²⁰⁾ reported that *in utero* exposure to low doses of bisphenol A (2 and/or 20 µg/kg/day) affects prostate and preputial gland weight and decreases daily sperm production efficiency in mouse offspring; moreover, their results indicated that exposure to low doses of xenoestrogens during a critical period can affect the reproductive organ systems of male offspring. On the other hand, other investigators have failed to find such effects in mouse offspring when using an identical experimental design^{21,22)}. Thus, the issue of low-dose exposure to these potential endocrine-disrupting chemicals remains a matter of debate among investigators. Hence, as more refined analytical methods become available, risk assessment for previously characterized chemical

Table 2. Number of subjects examined

Group	DEHP dose (mg/kg/day)	No. of offspring examined			
		Age (wk)	1	3	9
Control	0	Male	8	8	8
		Female	6	8	8
DEHP	25	Male	10	7	7
		Female	11	7	7
DEHP	100	Male	13	8	7
		Female	9	8	8
DEHP	400	Male	9	7	6
		Female	7	7	7

substances should be repeated.

Embryo-fetotoxicity and teratotoxicity of DEHP have been studied in mice⁴⁻⁸⁾ and rats⁸⁻¹⁰⁾. These studies were conducted to elucidate whether *in utero* exposure to high doses of DEHP induces embryotoxicity and/or teratogenicity. The doses used in these previous studies were far in excess of human




Fig. 2. Effects of maternal exposure to DEHP on postnatal body weight of offspring.

Body weights of male (left panel) and female (right panel) offspring are shown at 1, 3 and 9 wk of age. Each column and vertical bar represent the mean and SEM, respectively. There were no significant differences among groups.




Fig. 3. Effects of maternal exposure to DEHP on postnatal body length of offspring.

Body lengths (nose to anus) of males (left panel) and females (right panel) are shown at 1, 3 and 9 wk of age. Each column and vertical bar represent the mean and SEM, respectively. There were no significant differences among groups.




Fig. 4. Effects of maternal exposure to DEHP on postnatal tail length of offspring.

Tail lengths of males (left panel) and females (right panel) are shown at 1, 3 and 9 wk of age. Each column and vertical bar represent the mean and SEM, respectively. There were no significant differences among groups.

Table 3. Organ weights in male offspring

Organ	Group	DEHP dose (mg/kg/day)	Age (wk)		
			1	3	9
Liver (g)	Control	0	0.372 ± 0.011 ^a	1.974 ± 0.090	15.55 ± 0.439
	DEHP	25	0.367 ± 0.024	1.984 ± 0.156	16.73 ± 0.560
	DEHP	100	0.334 ± 0.016	1.936 ± 0.138	14.78 ± 0.735
	DEHP	400	0.372 ± 0.037	2.276 ± 0.122	15.83 ± 0.691
Kidneys (g)	Control	0	0.191 ± 0.004	0.618 ± 0.018	2.951 ± 0.093
	DEHP	25	0.188 ± 0.008	0.585 ± 0.037	3.049 ± 0.124
	DEHP	100	0.164 ± 0.007	0.582 ± 0.042	2.842 ± 0.078
	DEHP	400	0.163 ± 0.015	0.632 ± 0.024	3.071 ± 0.092
Testes (g)	Control	0	- ^b	0.222 ± 0.009	3.065 ± 0.095
	DEHP	25	-	0.225 ± 0.014	2.999 ± 0.102
	DEHP	100	-	0.213 ± 0.011	2.834 ± 0.050
	DEHP	400	-	0.241 ± 0.012	3.070 ± 0.092
Prostate (g)	Control	0	-	-	0.443 ± 0.026
	DEHP	25	-	-	0.428 ± 0.033
	DEHP	100	-	-	0.372 ± 0.032
	DEHP	400	-	-	0.358 ± 0.026
Seminal vesicles (g)	Control	0	-	-	1.109 ± 0.057
	DEHP	25	-	-	1.064 ± 0.060
	DEHP	100	-	-	0.979 ± 0.034
	DEHP	400	-	-	1.014 ± 0.096

^aValues are mean ± SEM. ^b -, not examined.

environmental exposure, and the duration of dosing was limited to the period of gestation. The present study was thus designed to investigate whether in utero and lactational exposure to DEHP affects the development of the next generation. For the main study, we set the highest dose at 400 mg/kg/day to avoid the influence of maternal toxicity and observe the effect of DEHP on the offspring. The exposure period was prolonged to examine the effects of lactational exposure in addition to the effects of gestational exposure. The offspring of dams in which no overt toxicity was observed (0, 25, 100 and 400 mg/kg/day), as determined by body weight and general behavior during gestation and lactation, were used in our study.

In recent years, certain studies have focused on the effects of DEHP and its antiandrogenic action on the hypothalamus-pituitary-gonadal axis^{16, 17, 29}; very few studies, however, have reported the effect of DEHP on longer term postnatal development. Hence, it is important to examine the developmental toxicity of DEHP from birth until puberty. In this regard, our study was performed to evaluate the effects of in utero and lactational exposure to DEHP in rat offspring with a special focus on postnatal growth and physical status. We found that somatic and tissue growth and related endocrine landmarks were not affected by DEHP exposure.

Liver weights were slightly increased in the 400 mg/kg/day group for both male and female offspring at 3 wk of age, but no significant differences were observed among treatment groups. DEHP and other phthalates, such as di(2-ethylhexyl) adipate (DEHA) and butylbenzyl phthalate, are peroxisome proliferators that activate peroxisome proliferator-activated receptors and cause liver enlargement²³. Induction of peroxisome proliferator-activated receptors could result in liver enlargement following DEHP exposure (Table 3, 4). This phenomenon could be an adaptive response following consecutive exposures to DEHP. However, this trend was no longer apparent at 9 wk of age. Since the DEHP groups were not exposed to the compound after 3 wk of age, body burden might be decreased because of metabolic clearance.

In a study of reproductive and accessory organ development following DEHP exposure, dose-dependent reductions in ventral, dorsolateral and/or anterior prostate weight were reported in rat offspring on PND 21 and PND 63 in response to oral administration of DEHP (0, 375, 750 and 1,500 mg/kg/day, GD3-PND21)¹⁶. This study also showed that DEHP significantly reduced testis weight on PND 21 and PND 63 in a dose-dependent manner. In the present study, on the other hand, testis weights were not

Table 4. Organ weights in female offspring

Organ	Group	DEHP dose (mg/kg/day)	Age (wk)		
			1	3	9
Liver (g)	Control	0	0.338 ± 0.007 ^a	1.899 ± 0.117	9.665 ± 0.573
	DEHP	25	0.322 ± 0.015	1.886 ± 0.103	9.279 ± 0.511
	DEHP	100	0.349 ± 0.014	1.808 ± 0.105	9.760 ± 0.505
	DEHP	400	0.367 ± 0.030	2.046 ± 0.092	9.643 ± 0.441
Kidneys (g)	Control	0	0.176 ± 0.006	0.605 ± 0.026	2.039 ± 0.078
	DEHP	25	0.177 ± 0.006	0.593 ± 0.025	1.849 ± 0.091
	DEHP	100	0.179 ± 0.007	0.583 ± 0.023	1.983 ± 0.055
	DEHP	400	0.171 ± 0.007	0.583 ± 0.020	1.959 ± 0.039
Ovaries (mg)	Control	0	- ^b	18.95 ± 0.76	79.57 ± 4.08
	DEHP	25	-	17.80 ± 1.98	74.28 ± 8.14
	DEHP	100	-	14.83 ± 1.83	71.00 ± 4.26
	DEHP	400	-	16.67 ± 0.82	73.42 ± 3.29
Uterus (mg)	Control	0	-	26.03 ± 1.91	327.4 ± 25.3
	DEHP	25	-	30.72 ± 3.95	300.7 ± 14.2
	DEHP	100	-	31.96 ± 2.37	376.3 ± 30.9
	DEHP	400	-	27.82 ± 2.15	340.5 ± 16.1

^aValues are mean ± SEM. ^b-, not examined.

Table 5. Hormone determinations in male offspring at 9 wk of age

Parameter	DEHP dose (mg/kg/day)			
	0	25	100	400
T ₄ (ng/ml)	83.1 ± 6.9 ^a	74.1 ± 3.7	73.2 ± 4.7	81.2 ± 7.5
T ₃ (ng/ml)	1.74 ± 0.05	1.70 ± 0.06	1.63 ± 0.07	1.81 ± 0.09
GH (ng/ml)	140.0 ± 35.3	137.3 ± 30.2	130.5 ± 16.3	96.5 ± 19.5
IGF-I (ng/ml)	669.6 ± 49.0	641.7 ± 57.8	758.6 ± 49.6	743.5 ± 23.8

^aValues are mean ± SEM.

Table 6. Hormone determinations in female offspring at 9 wk of age

Parameter	DEHP dose (mg/kg/day)			
	0	25	100	400
T ₄ (ng/ml)	70.0 ± 7.4 ^a	70.7 ± 5.4	67.7 ± 4.8	69.1 ± 6.4
T ₃ (ng/ml)	1.88 ± 0.11	1.91 ± 0.06	1.76 ± 0.06	1.79 ± 0.10
GH (ng/ml)	98.4 ± 9.6	99.5 ± 19.6	121.3 ± 22.4	109.4 ± 19.4
IGF-I (ng/ml)	499.0 ± 34.4	574.0 ± 34.6	528.6 ± 42.5	632.6 ± 66.0

^aValues are mean ± SEM.

significantly different between the control and DEHP groups. No significant differences in prostate weights were observed among the groups, although they were reduced in a dose-dependent manner (Table 3). The outcomes of the present study at the highest dose (400 mg/kg/day) were in accordance with those of Moore *et al.*, who conducted a study that used 375 mg/kg/day as the lowest dose¹⁶⁾. The magnitude of DEHP

effects in the present study was much smaller than that found in the study by Moore *et al.*¹⁶⁾; this discrepancy could be explained by the large difference in dosage range.

Thyroid hormones play pivotal roles in normal growth, neuronal development and metabolism in animals. Endocrine disturbance following chemical exposure is suspected to occur at the embryonic and/or neonatal stage rather than at

the adult stage. An epidemiological study has suggested that toxicants such as polychlorinated biphenyls and dioxins, which are persistent and cumulative compounds in the environment, may affect growth and development through thyroid impairment²⁴. Animal studies have reported that 2,3,7,8-tetrachlorodibenzo-*p*-dioxin disrupts thyroid homeostasis²⁵ and causes developmental defects²⁶ and bone growth deficits²⁷. Thyroid hormones are hormonal regulators of bone growth. The principal hormonal regulators during postnatal development are GH and IGF-I, and these hormones, which are regulated by thyroid hormones, are considered biomarkers for longitudinal somatic growth²⁸. In the present study, hormonal parameters regarding developmental somatic growth were determined in the offspring to better assess the physical status following DEHP exposure. There were no significant differences in any parameters in male and female rat offspring (Table 5, 6). The fact that normal hormonal parameters were observed in rat offspring following exposure of dams to DEHP (even at high doses) leads us to conclude that postnatal development remains intact in the offspring.

The level of DEHP exposure used in the present study was much greater (~1,000-fold higher) than the estimated intake due to either medical exposure or consumer exposure in adult humans³. It was recently suggested that the magnitude of testicular toxicity after DEHP exposure is associated with the duration and/or the route of exposure^{14, 29}. Inhalation of DEHP caused an elevation of plasma testosterone without affecting gonadotropin and several steroid enzymes that are involved in testosterone synthesis in male prepubertal rats¹⁴. These findings suggest that levels of DEHP that cause hormonal disturbance when inhaled may not have the same effect if consumed orally.

In conclusion, our results suggest that prenatal and postnatal exposure to DEHP does not affect postnatal somatic growth or endocrine and physical status of either males or females under the experimental conditions we used. The effects of DEHP exposure, however, remain uncertain and must be clarified using a wider dosage range, an extended exposure period, a side-by-side comparison of different exposure routes and a larger number of animals.

Acknowledgements

The authors thank Mr. T. Murase, Ms. S. Watanabe and Mr. S. Numajiri for their help throughout this study. This study was conducted as a part of contract research with the Ministry of Health, Labour and Welfare, which was supported by funds from the Ministry of the Environment.

References

- 1) NTP-CERHR (2000) CERHR Expert panel report on di(2-ethylhexyl) phthalate. National Toxicology Program, U.S. Department of Health and Humane Services.
- 2) EHC 131. Diethylhexyl phthalate. Environmental Health Criteria 131, The International Programme on Chemical Safety, WHO, Geneva. <http://www.inchem.org/documents/ehc/ehc/ehc131.htm>. Accessed June 5, 2006.
- 3) Clark K, Cousins I, Mackay D (2003) Assessment of Critical Exposure Pathways. In: The handbook of environmental chemistry, Vol. 3, Part Q, Phthalate esters, Staples CA (Ed.), 227–62, Springer-Verlag, Berlin.
- 4) Shiota K, Chou MJ, Nishimura H (1980) Embryotoxic effects of di-2-ethylhexyl phthalate (DEHP) and di-n-butyl phthalate (DBP) in mice. *Environ Res* **22**, 245–53.
- 5) Shiota K, Nishimura H (1982) Teratogenicity of di(2-ethylhexyl) phthalate (DEHP) and di-n-butyl phthalate (DBP) in mice. *Environ Health Perspect* **45**, 65–70.
- 6) Shiota K, Miwa S (1985) Assessment of the teratogenicity of di (2-ethylhexyl) phthalate and mono (2-ethylhexyl) phthalate in mice. *Arch Toxicol* **56**, 263–6.
- 7) Tomita I, Nakamura Y, Yagi Y, Tutikawa K (1982) Teratogenicity/fetotoxicity of DEHP in mice. *Environ Health Perspect* **45**, 71–5.
- 8) Tyl RW, Price CJ, Marr MC, Kimmel CA (1988) Developmental toxicity evaluation of dietary di(2-ethylhexyl)phthalate in Fischer 344 rats and CD-1 mice. *Fundam Appl Toxicol* **10**, 395–412.
- 9) Singh WR, Lawrence WH, Autian J (1972) Teratology of phthalate esters in rat. *J Pharm Sci* **61**, 51–5.
- 10) Lewandowski M, Fernandes J, Chen TS (1980) Assessment of the teratogenic potential of plasma-soluble extracts of diethylhexyl phthalate plasticized polyvinyl chloride plastics. *Toxicol Appl Pharmacol* **54**, 141–7.
- 11) Poon R, Lecavalier P, Mueller R, Valli VE, Procter BG, Chu I (1997) Subchronic oral toxicity of di-n-octyl phthalate and di (2-ethylhexyl) phthalate in the rat. *Food Chem Toxicol* **35**, 225–39.
- 12) Gray LE Jr, Ostby J, Furr J, Price M, Veeramachaneni DN, Parks L (2000) Perinatal exposure to the phthalates DEHP, BBP, and DINP, but not DEP, DMP, or DOTP, alters sexual differentiation of the male rat. *Toxicol Sci* **58**, 350–65.
- 13) Agarwal DK, Eustis S, Lamb JC 4th, Reel JR, Kluwe WM (1986) Effects of di(2-ethylhexyl) phthalate on the gonadal pathophysiology, sperm morphology, and reproductive performance of male rats. *Environ Health Perspect* **65**, 343–50.
- 14) Kurahashi N, Kondo T, Omura M, Umemura T, Ma M, Kishi R (2005) The effects of subacute inhalation of di (2-ethylhexyl) phthalate (DEHP) on the testes of prepubertal Wistar rats. *J Occup Health* **47**, 437–44.
- 15) Davis BJ, Maronpot RR, Heindel JJ (1994) Di-(2-ethylhexyl) phthalate suppresses estradiol and ovulation in cycling rats. *Toxicol Appl Pharmacol* **128**, 216–23.
- 16) Moore RW, Rudy TA, Lin T-M, Ko K, Peterson RE (2001)

- Abnormalities of sexual development in male rats with in utero and lactational exposure to the antiandrogenic plasticizer di(2-ethylhexyl)phthalate. *Environ Health Perspect* **109**, 229–37.
- 17) Akingbemi BT, Youker RT, Sottas CM, Ge R, Katz E, Klinefelter GR, Zirkin BR, Hardy MP (2001) Modulation of rat Leydig cell steroidogenic function by di(2-ethylhexyl)phthalate. *Biol Reprod* **65**, 1252–9.
 - 18) Kobayashi K, Miyagawa M, Wang RS, Sekiguchi S, Suda M, Honma T (2002) Effects of *in utero* and lactational exposure to bisphenol A on somatic growth and anogenital distance in F₁ rat offspring. *Ind Health* **40**, 375–81.
 - 19) Nagel SC, vom Saal FS, Thayer KA, Dhar MG, Boechler M, Welshons WV (1997) Relative binding affinity-serum modified access (RBA-SMA) assay predicts the relative *in vivo* bioactivity of the xenoestrogens bisphenol A and octylphenol. *Environ Health Perspect* **105**, 70–6.
 - 20) vom Saal FS, Cooke PS, Buchanan DL, Palanza P, Thayer KA, Nagel SC, Parmigiani S, Welshons WV (1998) A physiologically based approach to the study of bisphenol A and other estrogenic chemicals on the size reproductive organs, daily sperm production, and behavior. *Toxicol Ind Health* **14**, 239–60.
 - 21) Ashby J, Tinwell H, Haseman J (1999) Lack of effects for low dose levels of bisphenol A and diethylstilbestrol on the prostate gland of CF1 mice exposed *in utero*. *Regul Toxicol Pharmacol* **30**, 156–66.
 - 22) Cagen SZ, Waechter JM Jr, Dimond SS, Breslin WJ, Butala JH, Jekat FW, Joiner RL, Shiotsuka RN, Veenstra GE, Harris LR (1999) Normal reproductive organ development in CF-1 mice following prenatal exposure to bisphenol A. *Toxicol Sci* **50**, 36–44.
 - 23) Kersten S, Wahli W (2000) Peroxisome proliferator activated receptor agonists. *EXS* **89**, 141–51.
 - 24) Koopman-Esseboom C, Morse DC, Weisglas-Kuperus N, Lutkeschipholt IJ, Van der Paauw CG, Tuinstra LG, Brouwer A, Sauer PJ (1994) Effects of dioxins and polychlorinated biphenyls on thyroid hormone status of pregnant women and their infants. *Pediatr Res* **36**, 468–73.
 - 25) Nishimura N, Yonemoto J, Miyabara Y, Sato M, Tohyama C (2003) Rat thyroid hyperplasia induced by gestational and lactational exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Endocrinology* **144**, 2075–83.
 - 26) Pohjanvirta R, Tuomisto J (1994) Short-term toxicity of 2,3,7,8-tetrachlorodibenzo-p-dioxin in laboratory animals: effects, mechanisms, and animal models. *Pharmacol Rev* **46**, 483–549.
 - 27) Miettinen HM, Pulkkinen P, Jamsa T, Koistinen J, Simanainen U, Tuomisto J, Tuukkanen J, Viluksela M (2005) Effects of *in utero* and lactational TCDD exposure on bone development in differentially sensitive rat lines. *Toxicol Sci* **85**, 1003–12.
 - 28) Ohlsson C, Bengtsson BA, Isaksson OG, Andreassen TT, Słotweg MC (1998) Growth hormone and bone. *Endocr Rev* **19**, 55–79.
 - 29) Akingbemi BT, Ge R, Klinefelter GR, Zirkin BR, Hardy MP (2004) Phthalate-induced Leydig cell hyperplasia is associated with multiple endocrine disturbances. *Proc Natl Acad Sci USA* **101**, 775–80.

Effects of *in Utero* and Lactational Exposure to Bisphenol A on Thyroid Status in F₁ Rat Offspring

Kenichi KOBAYASHI*, Muneyuki MIYAGAWA, Rui-Sheng WANG,
Megumi SUDA, Soichiro SEKIGUCHI and Takeshi HONMA

Department of Health Effects Research, National Institute of Industrial Health, 6-21-1, Nagao, Tama-Ku, Kawasaki 214-8585, Japan

Received December 21, 2004 and accepted August 24, 2005

Abstract: Bisphenol A (BPA), a xenoestrogen, has been reported to mimic the actions of estrogen or to affect the endocrine glands *in vivo* and *in vitro*. In this study, we examined whether *in utero* and lactational exposure to BPA alters thyroid status in rat F₁ offspring. Dams were orally administered various doses of BPA (0, 4 or 40 mg/kg body weight per day) from gestation day (GD) 6 through postnatal day (PND) 20. The BPA and control groups did not differ significantly with respect to plasma thyroxine (T₄) concentration. The thyroid glands from the BPA groups had normal T₄ responses to exogenous thyroid-stimulating hormone *in vivo*. These results suggest that *in utero* and lactational exposure (indirect exposure) to BPA (4–40 mg/kg/day, GD 6 - PND 20) does not affect thyroid functions in the F₁ generation of male and female rats.

Key words: Bisphenol A, Thyroid, Thyroxine, Thyroid-stimulating hormone, Offspring, Rat

Introduction

Bisphenol A (BPA) is very widely used in the manufacture of polycarbonate and epoxy resins, dental sealants and other chemically derived products. BPA released from lacquer coatings has been detected in food cans¹, and it has also been found in saliva collected from subjects treated with dental sealants². Krishnan *et al.* have reported the weak estrogenic action of BPA eluted from a polycarbonate bottle into medium during an autoclaving procedure. They showed that BPA increases the number of progesterone receptors and promotes proliferation of a cultured cell line that originated from a human breast cancer (MCF-7)³. It has been estimated that total daily intake is 0.48 µg/kg body weight (BW) per day for 60-kg adults⁴.

Reproductive toxicity of BPA has been reported in mice and rats, and low-dose effects of BPA *in vivo* have been observed in mice. BPA increased prostate and preputial gland weight and decreased daily sperm production efficiency in

male mice offspring prenatally exposed to BPA at 2 or 20 µg/kg/day from gestation day (GD) 11 through GD 17^{5, 6}. Other investigators, however, failed to confirm such effects in mouse offspring using identical experimental designs^{7, 8}. Cagen *et al.* reported normal reproductive development in Wistar rat offspring born from mothers supplied with BPA in drinking water at a concentration range of 0.01 to 10 ppm (equivalent to approximately 0.001–4.022 mg/kg/day) for 10 weeks, from the pre-mating day (at 9 weeks old) to the weaning day⁹. In addition, multigeneration reproductive toxicity studies of BPA have been conducted in rats^{10, 11}. Both of these studies confirmed that there were no BPA-related effects on developmental and reproductive parameters at low doses of exposure.

Although the reported effects of BPA are not entirely consistent, it has received a great deal of attention with regard to its possible effects on reproductive glands and accessory genital glands, which are due primarily to its estrogenic activity. It has been known that BPA exerts weak estrogenic activity *in vivo*¹² and *in vitro*¹³. BPA may also disrupt thyroid homeostasis because of its structural similarity to thyroid

*To whom correspondence should be addressed.

Table 1. Treatment design and number of subjects examined

Measurement	Group	Dose (mg/kg/day)	No. of offspring tested			
			Age (wks)	1	3	9
Plasma T ₄	Control	0	Male	6	5	5
			Female	8	5	5
	BPA	4	Male	9	6	3
			Female	8	6	5
	BPA	40	Male	1	5	5
			Female	9	5	5
TSH stimulation test	Control	0	Male	— ^a	—	4
			Female	—	—	7
	BPA	4	Male	—	—	2
			Female	—	—	4
	BPA	40	Male	—	—	4
			Female	—	—	4

^aNot examined.

hormone. *In vitro* studies have demonstrated that BPA binds weakly to the thyroid hormone receptor and suppresses transcriptional activity that is stimulated by tri-iodothyronine (T₃)¹⁴. On the other hand, BPA does not inhibit the binding of T₃ to the thyroid hormone receptor and does not inhibit the hormonal activity of T₃ to induce growth and growth hormone (GH) production in the rat pituitary cell line GH3¹⁵. Although BPA is suspected to mimic thyroid hormone by modulating the thyroid hormone receptor, the *in vitro* effect of BPA on thyroid function/action is controversial, and the *in vivo* effect of BPA on thyroid status is unknown.

In the present study, we administered relatively high oral doses of BPA to pregnant rats and measured the effects on thyroid parameters including plasma thyroxine (T₄) levels at 1, 3 and 9 weeks of age in male and female offspring. To better assess thyroid function, we also measured *in vivo* T₄ responsiveness of thyroid glands to exogenous thyroid-stimulating hormone (TSH) at 9 weeks of age.

Materials and Methods

Animals

BPA (purity >99.8%, Cat#: 280-08561, Lot#: HCE9312) and corn oil (Cat#: 034-17015) were obtained from Wako Pure Chemical Industries, Ltd., Osaka, Japan. A total of 24 pregnant female rats (Crj: CD (SD) IGS strain, 9 weeks of age) at GD 3 were purchased from Charles River Japan, Inc. (Tsukuba) and housed separately and maintained under controlled temperature (23 ± 1°C), humidity (55 ± 5%) and a 12-h light-dark cycle (08:00–20:00) conditions throughout the study. The presence of a copulatory plug defined GD 0.

A standard laboratory diet (CE-2, Clea Japan, Inc., Tokyo, Japan) and drinking water were available *ad libitum*. Dams were randomly divided into four groups (6 pregnant rats per group) and weighed once a day from GD 3 through PND 20 (except for GD 4–5). The BPA-exposed groups were dosed by oral gavage with 4, 40 or 400 mg/kg BW per day of BPA in corn oil vehicle (10 ml/kg BW) once daily between 08:30 and 09:30 from GD 6 through PND 20, and the control group was given the same amount of corn oil during the same period. The litter size was standardized to ten (male:female = 5:5, if possible) between 10:00 and 11:00 on PND 7 (1 week of age). Examinations were performed as soon as possible after offspring were culled. On PND 21, the remaining offspring were weaned, and thereafter males and females were housed separately per litter. A pair of male and female offspring from each dam was dissected at 3 and 9 weeks of age. The remaining offspring were used for brain function and behavioral effects (not reported here). Table 1 shows the treatment design and number of subjects examined. The 400 mg/kg/day group was excluded from further analysis because of its excessive maternal toxicity¹⁶.

Hormone assay

Plasma samples were prepared as described previously¹⁶. Plasma T₄ levels were determined by a chemiluminescence immunoassay (ACS; Centaur; Chiron Corp., Emeryville, CA). The plasma T₄ levels were also used as basal values for the TSH stimulation test (see below).

TSH stimulation test

The BPA-exposed and control groups of both sexes at 9




Fig. 1. Effects of maternal exposure to BPA on plasma T₄ levels in male and female offspring.

Each column and vertical bar represents the mean and SEM, respectively. There were no significant differences among groups.

weeks of age were injected with bovine TSH (bTSH; Sigma-Aldrich Corp., St. Louis, MO) intraperitoneally at 25.0 mIU/5 μ l/g BW and intramuscularly at 12.5 mIU/5 μ l/g BW. Blood samples were collected from the postcaval vein under euthanasia by ether inhalation at 6 h after bTSH administration and then stored at -20°C until the analysis. Plasma T₄ levels were measured as described above and are presented as the percentage of corresponding basal values.

Statistical analysis

For plasma T₄ determinations, the differences from the corresponding control group were statistically analyzed by analysis of variance followed by Dunnett's test (significance at $p < 0.05$). In the TSH stimulation tests, the difference from the corresponding basal value was statistically analyzed using the Student's or Welch's *t*-test (significance at $p < 0.05$).

Results

In male and female offspring, no statistically significant differences in plasma T₄ levels were observed between the control and BPA groups at 1, 3 and 9 weeks of age (Fig. 1). Plasma T₄ levels after TSH administration were significantly elevated compared with the corresponding basal values in the control and the BPA groups. No treatment-related differences were measured in the T₄ levels or in the relative T₄ increase in response to exogenous TSH in all groups of both sexes (Figs. 2, 3). In male offspring, plasma T₄ levels in the control, 4 and 40 mg/kg/day groups increased by 243%, 206% and 330%, respectively, compared with the corresponding basal values. Likewise, in female offspring, plasma T₄ levels increased by 285%, 311% and 275%, respectively (Fig. 3).

Discussion

Reproductive and developmental toxicity studies have been conducted using higher doses of BPA. Although maternal toxicity (reduction in maternal weight gain during gestation) was noted in rats exposed to BPA by gastric intubation at 160, 320 and 640 mg/kg/day from GD 6 through GD 15, there were no toxic effects on the development of their pups¹⁷. Kwon *et al.* showed that exposure at 320 mg/kg/day from GD 11 through PND 20 resulted in no apparent change in male and female pubertal development and reproductive function in SD rats¹⁸. We previously reported that exposure at 4 or 40 mg/kg/day from GD 6 through PND 20 did not cause changes in somatic growth or anogenital distance in rat offspring of both sexes¹⁶, but these doses did affect testosterone homeostasis in the male testis¹⁹. Thus, much of the interest in BPA toxicity has focused on its putative effects on reproductive organs and genital glands. In the present study, we investigated whether *in utero* and lactational exposure to BPA affects endocrine status in the thyroid gland of rat offspring. Plasma T₄ levels were unaffected in the BPA groups (Fig. 1). A TSH stimulation test was then performed to examine thyroid function. Plasma T₄ levels were significantly elevated after exogenous TSH administration in the BPA-exposed groups, similar to the elevation seen in the control group (Fig. 2), suggesting that BPA exposure does not affect the synthesis and release of thyroid hormone in offspring *in vivo*. To our knowledge, this study represents the first attempt to better examine the effects of BPA on thyroid function *in vivo* in rat offspring exposed to relatively high levels of BPA.

Thyroid hormones play important roles in normal growth, neuronal development and metabolism in animals. During