

- Mann, H. B., Whitney, D. R. (1947). On a test of whether one of two random variables is stochastically larger than the other. *Ann. Math. Stat.* 18:50–60.
- METI (Ministry of Economy, Trade and Industry of Japan). (2006). 2-(2H-1,2,3-Benzotriazole-2-yl)-4,6-di-*tert*-butylphenol, document distributed in Committee on Safety of Chemical Substances, Chemical Substances Council, 30 June 2006. Accessed on April 25, 2007 from <http://www.meti.go.jp/committee/materials/g60705aj.html>.
- MHLW (Ministry of Health, Labour and Welfare of Japan). (2003). 2-(2'-Hydroxy-3',5'-de-*tert*-butylphenyl)benzotriazole. In *Toxicity Testing Reports of Environmental Chemicals (Ministry of Health, Labour and Welfare, ed.)*, Vol. 10. Tokyo: Chemicals Investigation Promoting Council, pp. 215–247.
- MHLW (Ministry of Health, Labour and Welfare of Japan). (2006). 2-(2'-Hydroxy-3',5'-de-*tert*-butylphenyl)benzotriazole. In *Toxicity Testing Reports of Environmental Chemicals (Ministry of Health, Labour and Welfare, ed.)*, Vol. 13, Tokyo: Chemicals Investigation Promoting Council, pp. 187–202.
- OECD (Organization for Economic Co-operation and Development). (1981). Guideline 452, Chronic Toxicity Studies (adopted 12th May 1981), OECD Guidelines for the Testing of Chemicals, section 5, OECD, Paris.
- OECD (Organization for Economic Co-operation and Development). (1998). Principles on Good Laboratory Practice (as revised in 1997), OECD Series on Principles of Good Laboratory Practice and Compliance Monitoring, No. 1, OECD, Paris.
- Steel, R. D. (1959). A multiple comparison rank sum test: treatment versus control. *Biometrics* 15:560–572.
- Tenkazai.com. (2007). Market Trend of Resin Additives “Light Stabilizer.” Accessed on April 25, 2007 from <http://www.tenkazai.com/market.html>.
- Wang, R. H., Schorer-Apelbaum, D., Weinstock, M. (2001). Testosterone mediates sex difference in hypothermia and cholinesterase inhibition by rivastigmine. *Eur. J. Pharmacol.* 433:73–79.

Gonadal Influence on the Toxicity of 2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole in Rats

Mutsuko Hirata-Koizumi,¹ Takashi Matsuyama,² Toshio Imai,¹ Akihiko Hirose,¹ Eiichi Kamata,¹ and Makoto Ema¹

¹Biological Safety Research Center, National Institute of Health Sciences, Tokyo, Japan

²Shin Nippon Biomedical Laboratories, Ltd., Drug Safety Research Laboratories (SNBL DSR), Kagoshima, Japan

Previously, we showed that susceptibility of male rats to the toxicity of an ultraviolet absorber, 2-(2'-hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole (HDBB), was nearly 25 times higher than that of females. In the current study, we investigated the role of sex steroids in the mediation of the gender-related difference using castrated rats. Male and female castrated CD(SD) rats were given HDBB by gavage at 0, 0.5, 2.5, or 12.5 mg/kg/day for 28 days. No deaths, clinical signs of toxicity, or changes in body weight or food consumption were found at any doses. Blood biochemical changes suggestive of hepatic damage, such as increased levels of aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase, and lactate dehydrogenase, were detected at 12.5 mg/kg/day in males. Absolute and relative liver weight increased at 0.5 mg/kg/day and above in males and at 12.5 mg/kg/day in females. In the liver, histopathological changes, such as nucleolar enlargement, increased mitosis, hypertrophy in hepatocytes, and/or focal necrosis were observed at 0.5 mg/kg/day and above in males, and at 2.5 mg/kg/day and above in females. These findings indicate that castration markedly reduced the gender-related differences in toxicity of HDBB in rats.

Keywords Benzotriazole UV absorber, Castration, Gender-related difference, Rats.

INTRODUCTION

2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole (CAS No. 3846-71-7; HDBB) is an ultraviolet (UV) absorber used in plastic resin products, such as

Address correspondence to Makoto Ema, Division of Risk Assessment, Biological Safety Research Center, National Institute of Health Sciences, 1-18-1, Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan; E-mail: ema@nihs.go.jp

building materials and automobile components (METI, 2006). Previously, we showed a marked difference in the susceptibility of male and female rats to the toxicity of HDBB in 28-day and 52-week repeated oral dose toxicity studies (Hirata-Koizumi et al., 2007; 2008). In the 28-day study, toxic effects in the liver, heart, kidneys, thyroids, and blood were observed. The no observed adverse effect level (NOAEL) for females was 2.5 mg/kg/day based on histopathological changes in the liver and heart detected at 12.5 mg/kg/day. However, the NOAEL for males could not be determined because hepatic changes were noted even at the lowest dose of 0.5 mg/kg/day. In the 52-week study, the NOAEL was concluded to be 0.1 mg/kg/day in males and 2.5 mg/kg/day in females based on histopathological changes in the liver. These findings consistently showed that male rats have a nearly 25 times higher susceptibility to HDBB toxicity than female rats.

Gender-related differences in susceptibility to toxicity have been documented for other substances; for example, a subchronic toxicity study in rats showed that fluoranthene, a polycyclic aromatic hydrocarbon, had greater effects on males than females, especially on the kidneys (Knuckles et al., 2004). In contrast, female rats exhibited greater susceptibility to hypothalamic cholinesterase inhibitory and hypothermic effects of a carbamate cholinesterase inhibitor, rivastigmine (Wang et al., 2001). Such gender-related variations are also reported in humans, mostly for medicines. Examples include the more severe adverse effects, but with greater improvement in response, to antipsychotic drugs such as chlorpromazine and fluspirilene in women (Harris et al., 1995).

Sex hormones are likely to play an important role in gender differences in toxicity responses. In fact, Wang et al. (2001) reported that orchidectomy completely abolished the above-mentioned sex differences in hypothalamic cholinesterase inhibition induced by rivastigmine, and testosterone treatment to gonadectomized males and females decreased the cholinesterase inhibitory effects of rivastigmine; therefore, it is apparent that testosterone interferes with the effects of rivastigmine. On the other hand, estrogen has been shown to act as a dopamine antagonist (Harris et al., 1995), which is considered to contribute, at least in part, to sex differences in response to antipsychotic drugs. The role of sex hormones in differences between sexes in toxicity responses seems to vary from case to case.

In the present study, we performed a repeated dose toxicity study of HDBB using male and female castrated rats to investigate the role of sex steroids in the mediation of sex difference in the susceptibility of rats to the toxicity of HDBB. Administration was conducted in the same way as the previous 28-day study using intact animals (Hirata-Koizumi et al., 2007) for comparison, and effects on the liver and heart, which were principally affected in the previous study of HDBB, were examined.

MATERIALS AND METHODS

This study was performed at Shin Nippon Biomedical Laboratories, Ltd., Drug Safety Research Laboratories (SNBL DSR; Kagoshima, Japan). The experiment was approved by the Institutional Animal Care and Use Committee of SNBL DSR and was performed in accordance with the ethics criteria contained in the bylaws of the Committee of SNBL DSR.

Chemicals

HDBB (Lot no. AY11) was obtained from Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan). The HDBB used in this study was 100% pure and was kept in a light-resistant and airtight container at room temperature. Test solutions were prepared as suspensions in corn oil twice a week and kept cool in a light-resistant and airtight container until dosing. Stability under refrigerated conditions was confirmed for seven days in the previous 28-day repeated dose toxicity study using intact animals (Hirata-Koizumi et al., 2007). All other reagents used in this study were of specific purity grade.

Animals

CrI:CD(SD) rats (SPF, three weeks old) were purchased from Hino Breeding Center, Charles River Laboratories Japan, Inc. (Yokohama, Japan). All animals were maintained in an air-conditioned room at 21.8–22.8°C, with a relative humidity of 45%–55%, a 12-h light/dark cycle, and ventilation with 15 air changes/h. Animals were housed individually in stainless cages suspended over a cage board. A basal diet (CE-2; CLEA Japan, Inc., Tokyo, Japan) and water, which meets the drinking water standard under the Water Works Law of Japan, were provided *ad libitum*.

Male and female rats were castrated under ether anesthesia between five and eight days after purchase. After a two-week acclimation, they were subjected to treatment at six weeks of age. Rats found to be in good health were selected and assigned to four groups of 10 males and 10 females by stratified random sampling based on body weight. One female in the highest dose group was excluded from the present study because remnants of the left ovary were confirmed at necropsy.

Experimental Design

Male and female castrated rats were given HDBB once-daily at 0 (vehicle control), 0.5, 2.5, and 12.5 mg/kg/day by gavage for 28 days. The dosage levels were determined based on the results of our previous 28-day study using intact rats given HDBB by gavage at 0.5, 2.5, 12.5, or 62.5 mg/kg/day, at which adverse effects, mainly on the liver and heart, were found at all doses in males and at

12.5 mg/kg/day and above in females (Hirata-Koizumi et al., 2007). The volume of each dose was adjusted to 10 mL/kg based on the latest body weight.

All animals were observed daily before and one to two hours after dosing for clinical signs of toxicity. Body weight was measured on days 0, 3, 7, 10, 14, 17, 21, 24, and 28 of the dosing period, and food consumption was recorded on days 0, 3, 7, 10, 14, 17, 21, 24, and 27 of the dosing period.

On the day after the last dosing, blood was drawn from the caudal vena cava in the abdomen with a heparin-added syringe under ether anesthesia and centrifuged to obtain plasma. The plasma was examined for biochemical parameters, such as total protein, albumin, glucose, total cholesterol, triglycerides, total bilirubin, urea nitrogen (BUN), creatinine, aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), creatinine phosphokinase, calcium, inorganic phosphorus, sodium, potassium, and chlorine. Following the collection of blood, all animals were euthanized by exsanguination, and the surface of the body, and organs and tissues of the entire body, were examined macroscopically. The liver and heart were then removed and weighed. Both organs were fixed in 10% neutral-buffered formalin, processed routinely for embedding in paraffin, and sections were prepared for staining with hematoxylin and eosin. Histopathological observation was performed for all groups.

Data Analysis

Parametric data, such as body weight, food consumption, blood biochemical parameters, and organ weights, were analyzed by Bartlett's test (Bartlett, 1937) for homogeneity of distribution ($p < 0.05$). When homogeneity was recognized, Dunnett's test (Dunnett, 1964) was conducted to compare control and individual treatment groups ($p < 0.01$ or 0.05). If not homogenous, the data were analyzed using a Dunnett-type mean rank test ($p < 0.01$ or 0.05) (Hollander and Wolfe, 1973).

RESULTS

No deaths or clinical signs of toxicity were found in any groups. There was no significant difference in body weight between the control and HDBB-treated groups (Fig. 1). Food consumption was also not significantly changed, except for a transient increase on day 21 of the administration period at 12.5 mg/kg/day and on day 27 of the administration period at 2.5 mg/kg/day in males (data not shown).

Blood biochemical examination revealed significant increases in the level of albumin at 0.5 mg/kg/day and above in males and at 2.5 mg/kg/day and above in females, total protein at 0.5 mg/kg/day and above in females, glucose at 12.5 mg/kg/day in males, and BUN at 12.5 mg/kg/day in both sexes (Table 1).




Figure 1: Body weight of male and female castrated rats given HDBB by gavage for 28 days.

Table 1: Blood biochemical findings in male and female castrated rats given HDBB by gavage for 28 days.

Dose (mg/kg/day)	0	0.5	2.5	12.5
Male				
No. of animals	10	10	10	10
Total protein (g/dL)	6.19 ± 0.32	6.44 ± 0.23	6.45 ± 0.40	6.26 ± 0.31
Albumin (g/dL)	4.43 ± 0.18	4.90 ± 0.17**	4.99 ± 0.25**	5.03 ± 0.18**
AST (IU/L)	61.0 ± 6.2	54.4 ± 3.5	63.6 ± 8.0	91.4 ± 24.0**
ALT (IU/L)	40.2 ± 8.9	37.9 ± 4.2	46.2 ± 8.6	55.5 ± 7.2**
ALP (IU/L)	868 ± 200	995 ± 267	989 ± 344	1552 ± 538**
LDH (IU/L)	112 ± 28	129 ± 18	173 ± 30*	403 ± 189**
Glucose (mg/dL)	176 ± 12	199 ± 13**	176 ± 7	196 ± 22*
BUN (mg/dL)	15.8 ± 2.0	15.3 ± 2.1	16.0 ± 1.8	19.7 ± 1.6**
Creatinine (mg/dL)	0.208 ± 0.020	0.174 ± 0.022**	0.176 ± 0.027**	0.175 ± 0.016**
Na (mEq/L)	145 ± 1	145 ± 1	145 ± 1	142 ± 1**
Cl (mEq/L)	107 ± 1	106 ± 2	106 ± 2	104 ± 2**
Female				
No. of animals	10	10	10	9 ^a
Total protein (g/dL)	5.81 ± 0.21	6.17 ± 0.26**	6.15 ± 0.18*	6.41 ± 0.34**
Albumin (g/dL)	4.19 ± 0.12	4.39 ± 0.22	4.55 ± 0.19**	5.14 ± 0.32**
AST (IU/L)	54.8 ± 3.5	62.4 ± 5.1*	57.4 ± 6.2	58.4 ± 10.0
ALT (IU/L)	39.1 ± 4.6	43.2 ± 7.8	39.5 ± 5.9	45.8 ± 8.7
ALP (IU/L)	727 ± 164	742 ± 122	703 ± 199	1026 ± 217**
LDH (IU/L)	138 ± 44	254 ± 27**	209 ± 44*	235 ± 116*
Glucose (mg/dL)	202 ± 25	181 ± 13	182 ± 10	216 ± 16
BUN (mg/dL)	20.0 ± 1.6	20.2 ± 2.1	18.2 ± 2.6	23.2 ± 2.2**
Creatinine (mg/dL)	0.230 ± 0.022	0.229 ± 0.025	0.196 ± 0.022*	0.208 ± 0.030
Na (mEq/L)	142 ± 1	143 ± 1	144 ± 1**	141 ± 1
Cl (mEq/L)	104 ± 1	105 ± 2	106 ± 1**	102 ± 2*

Values are expressed as the mean ± SD.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

^aOne female was excluded because left ovary remnants were found at autopsy.

The levels of LDH at 2.5 mg/kg/day and above in males and at 0.5 mg/kg/day and above in females, ALP at 12.5 mg/kg/day in both sexes, and AST and ALT at 12.5 mg/kg/day in males were also significantly increased. In addition, significant decreases in the levels of creatinine at 0.5 mg/kg/day and above, of sodium at 12.5 mg/kg/day in males, and of chloride at 12.5 mg/kg/day in both sexes were detected.

At necropsy, no gross abnormality was found at any dose. Absolute and relative liver weight was significantly increased at 0.5 mg/kg/day and above in males and at 12.5 mg/kg/day in females (Table 2). No significant change was found in the absolute and relative heart weight.

Histopathological findings in the liver are summarized in Table 3. Diffuse hypertrophy of hepatocytes were observed at 0.5 mg/kg/day and above in males and at 2.5 mg/kg/day and above in females. The cytoplasm of the hepatocytes was slightly eosinophilic. At these doses, anisokaryosis, nucleolar enlargement, and decreased glycogen in hepatocytes were also found. In addition, focal coagulative necrosis at 12.5 mg/kg/day in males and at 2.5 mg/kg/day and above in females, and increased mitosis of hepatocytes at 2.5 mg/kg/day and above and mononuclear cell infiltration at 12.5 mg/kg/day in males, were detected. No substance-related histopathological findings were detected in the heart.

DISCUSSION

The current study was designed to investigate the role of sex steroids in the mediation of gender-related differences in HDBB toxicity. The dosage of HDBB used in the present study was sufficiently high to be expected to induce

Table 2: Organ weight of the heart and liver in male and female castrated rats given HDBB by gavage for 28 days.

Dose (mg/kg/day)	0	0.5	2.5	12.5
Male				
No. of animals	10	10	10	10
Heart (g)	1.30 ± 0.07 (0.352 ± 0.022) ^a	1.25 ± 0.09 (0.331 ± 0.028)	1.35 ± 0.12 (0.362 ± 0.020)	1.37 ± 0.12 (0.373 ± 0.030)
Liver (g)	15.5 ± 1.5 (4.18 ± 0.27)	18.2 ± 2.7* (4.78 ± 0.47*)	21.6 ± 3.0** (5.76 ± 0.61**)	26.9 ± 1.9** (7.32 ± 0.40**)
Female				
No. of animals	10	10	10	9 ^b
Heart (g)	1.14 ± 0.07 (0.342 ± 0.027)	1.11 ± 0.09 (0.322 ± 0.027)	1.15 ± 0.10 (0.329 ± 0.024)	1.25 ± 0.14 (0.352 ± 0.035)
Liver (g)	14.5 ± 1.9 (4.33 ± 0.34)	14.8 ± 1.4 (4.28 ± 0.19)	16.2 ± 2.5 (4.63 ± 0.32)	27.0 ± 3.3** (7.63 ± 0.87**)

Values are expressed as the mean ± SD.

*Significantly different from the control, $p < 0.05$; **significantly different from the control, $p < 0.01$.

^aRelative organ weight (g/100 g body weight).

^bOne female was excluded because left ovary remnants were found at autopsy.

Table 3: Histopathological findings in the liver of male and female castrated rats given HDBB by gavage for 28 days.

	Grade	Dose (mg/kg/day)			
		0	0.5	2.5	12.5
Male					
No. of animals		10	10	10	10
Anisokaryosis of hepatocytes	±	0	1	8	3
	+	0	0	0	7
Nucleolar enlargement in hepatocytes	±	0	1	10	5
	+	0	0	0	5
Increased mitosis of hepatocytes	±	0	0	1	4
Hypertrophy of hepatocytes	±	0	4	10	10
Decreased glycogen in hepatocytes	±	0	1	6	8
	+	0	0	0	2
Focal necrosis	±	0	0	0	3
Mononuclear cell infiltration	±	1	0	0	5
Female					
No. of animals		10	10	10	9 ^a
Anisokaryosis of hepatocytes	±	0	0	5	8
Nucleolar enlargement in hepatocytes	±	0	0	5	9
Hypertrophy of hepatocytes	±	0	0	2	9
Decreased glycogen in hepatocytes	±	0	0	2	8
Focal necrosis	±	0	0	3	2
Mononuclear cell infiltration	±	1	1	1	0

Values represent the number of animals with the findings.

± = very slight; + = slight.

^aOne female was excluded because left ovary remnants were found at autopsy.

toxicological effects on the liver, based on the results of the previous 28-day and 52-week repeated dose toxicity study using intact rats (Hirata-Koizumi et al., 2007; 2008). As expected, absolute and relative liver weight increased at 0.5 mg/kg/day and above in males and at 12.5 mg/kg/day in females, and histopathological changes in the liver, including anisokaryosis, nucleolar enlargement, increased mitosis, hypertrophy and decreased glycogen in hepatocytes, focal necrosis, and/or mononuclear cell infiltration, were observed at 0.5 mg/kg/day and above in males and at 2.5 mg/kg/day and above in females. Blood biochemical changes, such as increases in the level of total protein, albumin, AST, ALT, ALP, and/or LDH, were also found at all doses in both sexes. Although these changes in blood biochemical parameters were mostly slight and lacked dose dependence in some cases, simultaneous increase in hepatic enzymes (AST, ALT, ALP, and LDH) at 12.5 mg/kg/day in males is considered to be related to hepatic damage caused by HDBB.

A previous 28-day study using intact rats showed the cardiac toxicity of HDBB; degeneration and hypertrophy of the myocardium or cell infiltration were found at 2.5 mg/kg/day and above in males and at 12.5 mg/kg/day and above in females (Hirata-Koizumi et al., 2007). In the present study, using castrated rats, however, histopathological changes in the heart were not

detected even at the highest dose of 12.5 mg/kg/day. Considering that histopathological effects on the heart were also not found at the highest dose of 2.5 mg/kg/day in males and 12.5 mg/kg/day in females in the previous 52-week study using intact rats (Hirata-Koizumi et al., 2008), the present results would not necessarily mean that castration caused a change in the cardiac effect of HDBB. Although the cause of the difference in the cardiac toxicity of HDBB in our studies is not clear, further study is required to investigate the toxicological effects of HDBB on the heart in more detail, including the effect on cardiac function (e.g., electrocardiographic parameters, blood pressure, etc.).

In the previous 28-day study, male and female intact rats were given HDBB by gavage at 0.5, 2.5, 12.5, or 62.5 mg/kg/day (Hirata-Koizumi et al., in press). Histopathological findings similar to those observed in the present study were detected in the liver at all doses in males and at 12.5 mg/kg/day and above in females. The changes were accompanied with an increase in the absolute and/or relative liver weight. Serum levels of hepatic enzymes increased at 12.5 mg/kg/day and above in males and slightly at 62.5 mg/kg/day in females. When comparing the sensitive endpoint for hepatotoxicity of HDBB, histopathological changes in the liver, between sexes, the changes detected at 0.5 mg/kg/day in male rats were comparable in severity and incidence to those at 12.5 mg/kg/day in females. Thus, it was considered that male rats showed a nearly 25 times higher susceptibility to the hepatotoxicity of HDBB than females. In the present study, using castrated rats, histopathological findings in the liver were detected in males but not in females at the lowest dose of 0.5 mg/kg/day. The hepatic changes at 0.5 mg/kg/day in males were slightly milder than those at 2.5 mg/kg/day in females, showing that the difference in the susceptibility of male and female castrated rats was less than five times. Thus, castration markedly reduced gender-related differences in the hepatotoxicity of HDBB. As shown in Figure 2, a comparison of the rate of changes in the relative liver weight provided a more clear description of a nearly 25 times difference in the susceptibility of male and female intact rats to HDBB hepatotoxicity and the marked reduction by castration.

When comparing the histopathological findings of the liver from the previous 28-day study using intact rats and the present study using castrated rats, those in males were approximately equivalent at the same dose. On the other hand, for females, hepatic changes were observed at 12.5 mg/kg/day and above in intact rats, but clear changes in the histopathology of the liver were detected in castrated rats at a lower dose of 2.5 mg/kg/day. Therefore, castration of female rats enhanced the adverse effects of HDBB on the liver, suggesting suppressive effects of estrogen on HDBB hepatotoxicity in rats. Comparison of the relative liver weight change (Fig. 2) showed decreased male susceptibility as well as increased female susceptibility by castration. Androgen might have an enhancing effect on the hepatotoxicity of HDBB.




Figure 2: Comparison of change in relative liver weight of male and female intact and castrated rats given HDBB by gavage.

^aThe result of the previous 28-day repeated dose toxicity study, in which male and female intact rats were given HDBB once-daily at 0 (vehicle control), 0.5, 2.5, 12.5, and 62.5 mg/kg/day by gavage (Hirata-Koizumi et al., 2007)

The current study showed that the gender-related difference in susceptibility to HDBB hepatotoxicity was reduced, but not abolished, by castration. Sexual differences found in the present study were considered to be due to exposure to sexual hormones before four weeks of age, when castration was conducted. In female rats, serum estradiol concentration during the first three weeks after birth is as high as or higher than the level during the proestrus stage in young adults (Döhler and Wuttke, 1975); however, because serum estradiol concentration is similarly high during this preweaning period in male rats, it is unlikely that exposure to estradiol during this period contributes to the sexual difference in susceptibility of rats to the toxicity of HDBB. On the other hand, serum androgen levels before four weeks of age are much higher in male than female rats (Döhler and Wuttke, 1975). Ketelslegers et al. (1978) reported that plasma testosterone level in male rats was as high as 50 ng/100 mL two days after birth and it remained at the same level until day 8. The progressive decline occurred from days 8–24, and the testosterone level remained low, at the limit of detection of the assay (18 ng/100 mL), until day 30. There is a possibility that neonatal exposure to testosterone plays some role in the different susceptibility of male and female rats to the toxicity of HDBB. In fact, neonatal exposure to androgen irreversibly programs brain centers involved in the hypothalamo-pituitary control of hepatic sex-dependent metabolism (Gustafsson et al., 1981). We are currently in the process of performing a repeated dose toxicity study of HDBB using preweaning rats to clarify when gender-related differences in susceptibility to the toxicity of HDBB develop.

As in the case of HDBB, the male-predominant induction of toxicity in rats has been reported for many other substances, such as adenine (Ogirima et al.,

2006), acetaminophen (Raheja et al., 1983), dapsone (Coleman et al., 1990), flu-oranthene (Knuckles et al., 2004), 3-nitropropionic acid (Nishino et al., 1998), and mercuric chloride (Muraoka and Itoh, 1980). Various causes of such gender-related differences are indicated mainly for toxicokinetic determinants. It is well known that hepatic metabolism differs between the sexes, with male rats generally having higher activity than females (Gad, 2006). Furthermore, gender differences in membrane transport in various organs, including the kidneys, liver, intestine, and brain, have emerged relatively recently (Morris et al., 2003). In the case of HDBB, male rats consistently showed greater susceptibility to various effects of HDBB (e.g., on the liver, blood, etc.) in the previous 28-day and 52-week studies (Hirata-Koizumi et al., 2007; 2008); therefore, such differences in metabolism or transport between the sexes might increase the blood concentration of causative substances (i.e., HDBB or its metabolites) in males.

For gender-related variations in toxicokinetic determinants, many mechanistic studies on the metabolic enzyme cytochrome P450 have been reported (Waxman and Chang, 2005). In rats, a subset of P450s is expressed in a sex-dependent fashion and is subject to endocrine control. Whereas testosterone has a major positive regulatory influence on male-specific P450 forms, estrogen plays a somewhat lesser role in the expression of the female-specific/predominant liver P450 enzymes. If the male-specific/predominant metabolic enzymes have an intimate involvement in the toxic activation of HDBB, our results, showing the higher susceptibility of male rats to HDBB toxicity than females and decreased susceptibility by castration of male rats, could be explained. Interestingly, it was reported that estradiol suppressed the expression of male-specific/predominant P450 enzymes (Waxman and Chang, 2005). This is consistent with our results that female susceptibility to the hepatotoxicity of HDBB was increased by castration, given that the male-specific/predominant P450 enzymes activate HDBB. Since the expression of female-specific/predominant P450 enzymes is reduced by testosterone treatment as well as by castration of females (Waxman and Chang, 2005), there is also the possibility that these enzymes might be involved in the detoxication of HDBB. In order to clarify the cause of the sexual differences in susceptibility of rats to the toxicity of HDBB, we are planning a toxicokinetic study of HDBB, which would include the identification of metabolites and the related metabolic enzyme as well as measurement of the blood concentration of HDBB both after a single and repeated administration of HDBB to rats.

CONCLUSIONS

The current results showed that an oral administration of HDBB to castrated rats for 28 days caused hepatotoxicity at 0.5 mg/kg/day and above in males and at 2.5 mg/kg/day and above in females. Castration markedly reduced gender-related differences in the toxicity of HDBB in male and female rats.

ACKNOWLEDGMENT

This study was supported by the Ministry of Health, Labour and Welfare, Tokyo, Japan.

REFERENCES

- Bartlett, M. S. (1937). Properties of sufficiency and statistical tests. *Proc. R. Soc. Lond. Ser. A* 160:268–282.
- Coleman, M. D., Tingle, M. D., Winn, M. J., Park, B. K. (1990). Gonadal influence on the metabolism and haematological toxicity of dapsone in the rat. *J. Pharm. Pharmacol.* 42:698–703.
- Döhler, K. D., Wuttke, W. (1975). Changes with age in levels of serum gonadotropins, prolactin, and gonadal steroids in prepubertal male and female rats. *Endocrinology* 97:898–907.
- Dunnett, C. W. (1964). New tables for multiple comparisons with a control. *Biometrics* 20:482–491.
- Gad, S. C. (2006). Metabolism. In Gad, S. C., (ed.), *Animal Models in Toxicology*, 2nd ed. Florida, USA: CRC Press, pp. 217–247.
- Gustafsson, J. A., Eneroth, P., Hokfelt, T., Mode, A., Norstedt, G., Skett, P. (1981). Role of the hypothalamo-pituitary-liver axis in sex differences in susceptibility of the liver to toxic agents. *Environ. Health Perspect.* 38:129–141.
- Harris, R. Z., Benet, L. Z., Schwartz, J. B. (1995). Gender effects in pharmacokinetics and pharmacodynamics. *Drugs* 50:222–239.
- Hirata-Koizumi, M., Ogata, H., Imai, T., Hirose, A., Kamata, E., Ema, M. (2008). A 52-week repeated dose toxicity study of ultraviolet absorber 2-(2'-hydroxy-3',5'-di-tert-butylphenyl)benzotriazole in rats. *Drug Chem. Toxicol.* 31:In press.
- Hirata-Koizumi, M., Watari, N., Mukai, D., Imai, T., Hirose, A., Kamata, E., Ema, M. (2007). A 28-day repeated dose toxicity study of ultraviolet absorber 2-(2'-hydroxy-3',5'-di-tert-butylphenyl) benzotriazole in rats. *Drug Chem. Toxicol.* 30:327–341.
- Hollander, M., Wolfe, D. A. (1973). *Nonparametric Statistical Methods*. New York: John Wiley & Sons.
- Ketelslegers, J. M., Hetzel, W. D., Sherins, R. J., Catt, K. J. (1978). Developmental changes in testicular gonadotropin receptors: plasma gonadotropins and plasma testosterone in the rat. *Endocrinology* 103:212–222.
- Knuckles, M. E., Inyang, F., Ramesh, A. (2004). Acute and subchronic oral toxicity of fluoranthene in F-344 rats. *Ecotoxicol. Environ. Saf.* 59:102–108.
- METI (Ministry of Economy, Trade and Industry of Japan). (2006). 2-(2H-1,2,3-Benzotriazole-2-yl)-4,6-di-tert-butylphenol, document distributed in Committee on Safety of Chemical Substances, Chemical Substances Council, 30 June 2006. Accessed on May 18, 2007 from <http://www.meti.go.jp/committee/materials/g60705aj.html>.
- Morris, M. E., Lee, H. J., Predko, L. M. (2003). Gender differences in the membrane transport of endogenous and exogenous compounds. *Pharmacol. Rev.* 55:229–240.
- Muraoka, Y., Itoh, F. (1980). Sex difference of mercuric chloride-induced renal tubular necrosis in rats—from the aspect of sex differences in renal mercury concentration and sulfhydryl levels. *J. Toxicol. Sci.* 5:203–214.

- Nishino, H., Nakajima, K., Kumazaki, M., Fukuda, A., Muramatsu, K., Deshpande, S. B., Inubushi, T., Morikawa, S., Borlongan, C. V., Sanberg, P. R. (1998). Estrogen protects against while testosterone exacerbates vulnerability of the lateral striatal artery to chemical hypoxia by 3-nitropropionic acid. *Neurosci. Res.* 30:303–312.
- Ogirima, T., Tano, K., Kanehara, M., Gao, M., Wang, X., Guo, Y., Zhang, Y., Guo, L., Ishida, T. (2006). Sex difference of adenine effects in rats: renal function, bone mineral density, and sex steroidogenesis. *Endocr. J.* 53:407–413.
- Raheja, K. L., Linscheer, W. G., Cho, C. (1983). Hepatotoxicity and metabolism of acetaminophen in male and female rats. *J. Toxicol. Environ. Health* 12:143–158.
- Wang, R. H., Schorer-Apelbaum, D., Weinstock, M. (2001). Testosterone mediates sex difference in hypothermia and cholinesterase inhibition by rivastigmine. *Eur. J. Pharmacol.* 433:73–79.
- Waxman, D. J., Chang, T. K. (2005). Hormonal regulation of liver cytochrome P450 enzymes. In Ortiz de Montellano, P. R. (ed.), *Cytochrome P450 – Structure, Mechanism, and Biochemistry*, 3rd ed. New York: Kluwer Academic/ Plenum Publishers, pp. 347–376

Lack of Gender-Related Difference in the Toxicity of 2-(2'-Hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole in Prewaning Rats

Mutsuko Hirata-Koizumi,¹ Takashi Matsuyama,² Toshio Imai,¹ Akihiko Hirose,¹ Eiichi Kamata,¹ and Makoto Ema¹

¹Division of Risk Assessment, Biological Safety Research Center, National Institute of Health Sciences, Tokyo, Japan

²Drug Safety Research Laboratories, Shin Nippon Biomedical Laboratories, Ltd. (SNBL DSR), Kagoshima, Japan

In our previous toxicity studies using young rats, we showed that an ultraviolet absorber, 2-(2'-hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole (HDBB), principally affected the liver, and male rats had nearly 25 times higher susceptibility to the toxic effects than females. In the present study, the toxicity of HDBB was investigated in preweaning rats. HDBB was administered by gavage to male and female CD(SD) rats from postnatal days 4 to 21 at a dose of 0, 0.1, 0.5, 2.5, or 12.5 mg/kg/day. No substance-related deaths, clinical signs of toxicity, or body-weight changes were observed. Increased levels of albumin, AST and ALP in both sexes, BUN in males, and LDH in females were found at 12.5 mg/kg. Liver weights increased at 2.5 mg/kg and above in both sexes. Histopathologically, hepatocellular findings, such as nucleolar enlargement, anisokaryosis, increased mitosis, and/or hypertrophy, were observed at 2.5 mg/kg and above in both sexes. These results indicate no gender-related differences in the susceptibility to the toxic effects of HDBB in preweaning rats.

Keywords Benzotriazole UV absorber, Prewaning rat, Gender-related difference, Hepatotoxicity.

Address correspondence to Makoto Ema, Division of Risk Assessment, Biological Safety Research Center, National Institute of Health Sciences, 1-18-1, Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan; E-mail: ema@nihs.go.jp

INTRODUCTION

A number of reports have been published on gender-related differences in the toxic effects of chemicals in rats (Agarwal et al., 1982; Coleman et al., 1990; McGovren et al., 1981; Muraoka and Itoh, 1980; Nishino et al., 1998; Ogirima et al., 2006; Raheja et al., 1983). For example, fluoranthene, a polycyclic aromatic hydrocarbon, showed greater effects on male rats than females, especially on the kidneys, in a subchronic toxicity study (Knuckles et al., 2004). In contrast, female rats exhibited greater susceptibility to hypothalamic cholinesterase inhibitory and hypothermic effects of a carbamate cholinesterase inhibitor, rivastigmine (Wang et al., 2001). Such gender-related variations are also reported in humans, mostly for medicines (Harris et al., 1995). Examples include more severe adverse effects, but with greater improvement in response, to antipsychotic drugs such as chlorpromazine and fluspirilene in women.

Previously, we reported that male and female rats showed markedly different susceptibilities to the toxicity of 2-(2'-hydroxy-3',5'-di-*tert*-butylphenyl)benzotriazole (HDBB), which is an ultraviolet absorber used in plastic resin products, such as building materials and automobile components (METI, 2006). In a 28-day repeated-dose toxicity study, male and female rats were administered HDBB by gavage, and adverse effects on the liver, heart, blood, kidneys, and thyroids were found (Hirata-Koizumi et al., 2007). The no observed adverse effect level (NOAEL) for females was 2.5 mg/kg/day based on histopathological changes in the liver and heart detected at 12.5 mg/kg, but the NOAEL for males could not be determined because hepatic changes were noted even at the lowest dose of 0.5 mg/kg. In the 52-week repeated-dose toxicity study, chronic oral administration of HDBB principally affected the liver, and the NOAEL was concluded to be 0.1 mg/kg/day in males and 2.5 mg/kg/day in females (Hirata-Koizumi et al., 2008a), showing that male rats have approximately 25 times higher susceptibility to HDBB toxicity than females.

For such gender differences in toxic responses, sexual hormones are likely to play important roles. In fact, Wang et al. (2001) reported that orchidectomy completely abolished the above-mentioned sex differences in hypothalamic cholinesterase inhibition induced by rivastigmine, and testosterone treatment to gonadectomized males and females decreased the cholinesterase inhibitory effects of rivastigmine; therefore, it is apparent that testosterone interferes with the effects of rivastigmine. On the other hand, estrogen has been shown to act as a dopamine antagonist (Harris et al., 1995), which is considered to contribute, at least in part, to sex differences in response to antipsychotic drugs.

In order to investigate the role of sex steroids in the mediation of sex differences in the susceptibility to the toxic effects of HDBB, we recently performed a 28-day repeated-dose toxicity study using male and female

castrated rats (Hirata-Koizumi et al., 2008b). As expected, castration markedly reduced the sexual variation in HDBB toxicity, but some difference, less than five times, remained between male and female castrated rats. It is speculated that the determinants of susceptibility to HDBB toxicity are already differentiated between sexes by four weeks of age, when the castration was performed; therefore, in the present study, we determined the sexual difference in the susceptibility to HDBB toxicity in preweaning rats.

MATERIALS AND METHODS

This study was performed at Shin Nippon Biomedical Laboratories, Ltd., Drug Safety Research Laboratories (SNBL DSR; Kagoshima, Japan) in 2006–2007. The experiment was approved by the Institutional Animal Care and Use Committee of SNBL DSR and was performed in accordance with the ethics criteria contained in the bylaws of the Committee.

Animals and Housing Conditions

Eleven-week-old male and 10-week-old female Crl:CD(SD) rats were purchased from Hino Breeding Center, Charles River Laboratories Japan, Inc. (Yokohama, Japan) and individually housed in stainless steel cages suspended over a cage board. After a seven-day acclimation, females were cohabited overnight with one male each. Females with vaginal plugs were regarded as pregnant, and this day was designated as Day 0 of gestation. On gestation day 20, the pregnant females were transferred to aluminum cages with wooden chips as bedding (White Flake; Charles River Laboratories Japan, Inc.) and allowed to deliver spontaneously and rear their pups. The day of birth was defined as postnatal day (PND) 0. On PND 4, the sex of the pups was determined, and the litters were adjusted randomly to four males and four females. Five litters were selected and randomly assigned to each of five dose groups, including control groups; the initial number of pups for treatment was 20/sex/group.

Throughout the study, the animals were maintained in an air-conditioned room at 21.5–22.4°C, with a relative humidity of 43–55%, a 12-h light/dark cycle, and ventilation with 15 air changes/hour. A basal diet (CE-2; CLEA Japan, Inc., Tokyo, Japan) and water, which met the drinking water standard under the Water Works Law of Japan, were provided *ad libitum*.

Chemicals and Doses

HDBB (CAS No. 3846-71-7, Lot no. AY11) was 100% pure and was obtained from Tokyo Chemical Industry Co., Ltd. (Tokyo, Japan); it was kept in a dark place at room temperature under airtight conditions. Dosing

solutions were prepared as a suspension in corn oil (Wako Pure Chemical Industries, Ltd., Osaka, Japan) once or twice a week and kept cool in a dark place under airtight conditions until dosing. Stability under refrigerated conditions was confirmed for seven days in the previous 28-day repeated-dose toxicity study using young animals (Hirata-Koizumi et al., 2007).

Male and female preweaning rats were given HDBB by gavage once-daily from PNDs 4 to 21. Control rats received the vehicle only. A nutrient catheter (Type 3Fr; Atom Medical Corporation, Tokyo, Japan), attached to a disposable syringe, was used for dosing. The volume of each dose was adjusted to 10 mL/kg of body weight, based on the latest body weight.

The dosage levels of HDBB were determined to be 0.1, 0.5, 2.5, or 12.5 mg/kg/day, based on the results of our previous 28-day repeated-dose toxicity study using young rats (Hirata-Koizumi et al., 2007). In this previous study, male and female young rats were given HDBB by gavage at 0.5, 2.5, 12.5, or 62.5 mg/kg/day, and adverse effects, mainly on the liver and heart, were found at all doses in males and at 12.5 mg/kg and above in females.

Observations

All dams were observed daily for clinical signs of toxicity, and body weight was recorded on Days 0, 10, and 20 of pregnancy and on Days 0, 10, 20, and 22 after delivery. On Day 22 after delivery, they were euthanized by exsanguination under deep ether anesthesia, and the surface, organs, and tissues of the entire body were macroscopically observed.

All pups were observed daily before and three to four hours after dosing for clinical signs of toxicity. Body weight was recorded on PNDs 0, 4, 6, 8, 10, 12, 14, 16, 18, 21, and 22. On PND 22, blood was collected from the caudal vena cava in the abdomen of two male and two female pups per litter under deep ether anesthesia. Plasma separated from the blood by centrifugation was examined for total protein, albumin, glucose, total cholesterol, triglycerides, total bilirubin, blood urea nitrogen (BUN), creatinine, aspartate aminotransferase (AST), alanine aminotransferase (ALT), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), creatine phosphokinase, calcium, inorganic phosphorus, sodium, potassium, and chlorine. Following the collection of blood, all pups (four males and four females per litter) were euthanized by exsanguination under deep ether anesthesia, and the surface, organs, and tissues of the entire body were macroscopically observed. The heart, lungs, liver, spleen, kidneys, and adrenals were then collected and weighed. The liver and heart were histopathologically examined in one male and one female per litter. The organs were fixed in 10% neutral-buffered formalin, and paraffin sections for microscopic examination were routinely prepared and stained with hematoxylin-eosin.

Data Analysis

Body weight, blood biochemical parameters, and organ weights of pups were analyzed by Bartlett's test (Bartlett, 1937) for homogeneity of distribution ($p < 0.01$). When homogeneity was recognized, Dunnett's test (Dunnett, 1964) was conducted to compare between control and individual treatment groups ($p < 0.01$ or 0.05). If not homogenous, data were analyzed using the mean rank test of Dunnett's type (Hollander and Wolfe, 1973) ($p < 0.01$ or 0.05). Histopathological findings were analyzed using Wilcoxon's rank sum test (Wilcoxon, 1945) ($p < 0.01$ or 0.05).

RESULTS

HDBB, orally administered to pups from PNDs 4 to 21, did not induce any clinical signs of toxicity or affect the body weight of maternal rats (data not shown). At necropsy, no gross abnormality was found in the dams.

One male pup each at 0 or 0.5 mg/kg and one female pup each at 0, 0.5, or 12.5 mg/kg died, which was confirmed to be due to gavage error. No substance-related clinical signs of toxicity were found in pups of any groups. There were also no significant changes in the body weight of male and female pups, as shown in Figure 1.

Principle blood biochemical values are summarized in Table 1. In males, the levels of albumin, AST, ALP, and BUN were significantly increased at 12.5 mg/kg. In females, significant increases in the levels of albumin, AST, ALP, and LDH were found at the same dose. There were no substance-related changes in other blood biochemical parameters.




Figure 1: Body weight curves of male and female preweaning rats given HDBB by gavage.

Table 1: Principle blood biochemical values in male and female preweaning rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.1	0.5	2.5	12.5
No. of males	10	10	10	10	10
Total protein (g/dL)	4.49 ± 0.28	4.53 ± 0.22	4.48 ± 0.26	4.43 ± 0.17	4.42 ± 0.18
Albumin (g/dL)	3.62 ± 0.24	3.60 ± 0.24	3.59 ± 0.21	3.74 ± 0.27	4.04 ± 0.17**
BUN (mg/dL)	11.4 ± 1.5	14.1 ± 2.6	13.7 ± 5.3	12.9 ± 1.8	14.7 ± 2.3**
AST (IU/L)	91.4 ± 15.9	85.2 ± 4.8	88.7 ± 5.2	91.6 ± 12.2	100.2 ± 8.5*
ALT (IU/L)	34.8 ± 5.7	34.0 ± 6.3	29.4 ± 5.3	30.7 ± 5.5	35.9 ± 6.1
ALP (IU/L)	1557 ± 203	1529 ± 240	1412 ± 279	1286 ± 249	2054 ± 444**
LDH (IU/L)	198 ± 123	165 ± 16	184 ± 40	236 ± 170	326 ± 221
No. of females	10	10	10	10	10
Total protein (g/dL)	4.49 ± 0.24	4.54 ± 0.24	4.53 ± 0.28	4.55 ± 0.18	4.50 ± 0.14
Albumin (g/dL)	3.59 ± 0.28	3.66 ± 0.24	3.70 ± 0.26	3.80 ± 0.25	4.04 ± 0.16**
BUN (mg/dL)	12.5 ± 2.0	15.4 ± 1.5	13.5 ± 4.0	14.1 ± 4.1	15.5 ± 3.3
AST (IU/L)	87.3 ± 9.4	85.1 ± 8.2	86.5 ± 6.3	85.2 ± 6.6	101.3 ± 9.2**
ALT (IU/L)	30.7 ± 5.9	30.7 ± 3.6	27.1 ± 5.5	27.1 ± 4.5	35.9 ± 4.2
ALP (IU/L)	1470 ± 136	1394 ± 215	1287 ± 105	1339 ± 183	1872 ± 259**
LDH (IU/L)	175 ± 52	176 ± 36	179 ± 35	139 ± 28	370 ± 295*

Values are expressed as the mean ± SD.

BUN, blood urea nitrogen; AST, aspartate aminotransferase; ALT, alanine aminotransferase; ALP, alkaline phosphatase; LDH, lactate dehydrogenase.

*Significantly different from the control group ($p < 0.05$).

**Significantly different from the control group ($p < 0.01$).

At necropsy, no gross abnormality was observed. Absolute and relative organ weights of scheduled sacrifice animals are shown in Table 2. In males, absolute liver weight at 12.5 mg/kg and relative weight at 2.5 mg/kg and above were significantly increased. In addition, absolute and relative weights of the lungs and spleen were significantly decreased at 12.5 mg/kg. In females, significant increases in absolute liver weight at 12.5 mg/kg and relative liver weight at 2.5 mg/kg and above, and decreases in relative spleen weight and absolute and relative adrenal weight at 12.5 mg/kg, were found. No substance-related changes were detected in other organ weights.

Histopathological findings in the liver are presented in Table 3. In males, nucleolar enlargement, anisokaryosis, and increased mitosis of hepatocytes were observed at 2.5 mg/kg and above. In the 12.5 mg/kg group, hypertrophy of hepatocytes accompanied with eosinophilic granular changes was also observed. Further, increased incidence and/or severity of decreased glycogen in hepatocytes was found at 2.5 mg/kg and above. Similarly, in females, nucleolar enlargement, anisokaryosis, and increased mitosis of hepatocytes at 2.5 mg/kg and above, and hypertrophy and eosinophilic granular change of hepatocytes at 12.5 mg/kg were detected, and the incidence and/or severity of decreased glycogen in hepatocytes was higher at 12.5 mg/kg. No substance-related histopathological changes were detected in the heart in both sexes.

Table 2: Organ weights of male and female preweaning rats given HDBB by gavage.

Dose (mg/kg/day)	0	0.1	0.5	2.5	12.5
No. of males	19	20	19	20	20
Body weight (g)	67.2 ± 7.3	71.3 ± 6.9	67.3 ± 5.8	66.2 ± 9.6	66.2 ± 5.0
Heart (g)	0.37 ± 0.04 (0.55 ± 0.04) 0.58 ± 0.07 (0.87 ± 0.07)	0.37 ± 0.04 (0.52 ± 0.04) 0.58 ± 0.04 (0.82 ± 0.09)	0.36 ± 0.05 (0.53 ± 0.03) 0.53 ± 0.03* (0.80 ± 0.06*)	0.36 ± 0.05 (0.54 ± 0.03) 0.59 ± 0.08 (0.90 ± 0.09)	0.35 ± 0.04 (0.53 ± 0.04) 0.53 ± 0.04* (0.80 ± 0.06*)
Lung (g)	2.83 ± 0.47 (4.19 ± 0.36)	2.88 ± 0.34 (4.04 ± 0.26)	2.75 ± 0.44 (4.07 ± 0.42)	3.24 ± 0.68 (4.87 ± 0.40**)	4.54 ± 0.61** (6.84 ± 0.53**)
Liver (g)	0.37 ± 0.09 (0.55 ± 0.10)	0.40 ± 0.05 (0.57 ± 0.06)	0.34 ± 0.08 (0.51 ± 0.10)	0.38 ± 0.77 (0.57 ± 0.08)	0.29 ± 0.05** (0.44 ± 0.06**)
Spleen (g)	0.72 ± 0.09 (1.07 ± 0.07)	0.74 ± 0.06 (1.04 ± 0.07)	0.72 ± 0.08 (1.07 ± 0.08)	0.68 ± 0.10 (1.03 ± 0.05)	0.71 ± 0.07 (1.07 ± 0.08)
Kidneys (g)	17.5 ± 3.7 (26.2 ± 5.1)	19.3 ± 3.7 (27.3 ± 5.8)	18.1 ± 3.3 (27.4 ± 5.8)	21.5 ± 5.2* (32.4 ± 6.8**)	17.0 ± 2.4 (25.6 ± 3.3)
Adrenals (mg)	19	20	19	20	19
No. of females	19	20	19	20	19
Body weight (g)	64.0 ± 7.1	68.6 ± 7.5	63.6 ± 4.7	63.6 ± 8.9	65.3 ± 4.1
Heart (g)	0.35 ± 0.05 (0.54 ± 0.04)	0.35 ± 0.05 (0.51 ± 0.05)	0.33 ± 0.03 (0.52 ± 0.06)	0.34 ± 0.05 (0.53 ± 0.04)	0.35 ± 0.04 (0.53 ± 0.04)
Lung (g)	0.54 ± 0.08 (0.85 ± 0.11)	0.54 ± 0.06 (0.80 ± 0.09)	0.55 ± 0.06 (0.86 ± 0.10)	0.57 ± 0.09 (0.90 ± 0.12)	0.51 ± 0.05 (0.78 ± 0.06)
Liver (g)	2.72 ± 0.47 (4.23 ± 0.43)	2.77 ± 0.41 (4.02 ± 0.24)	2.62 ± 0.38 (4.12 ± 0.44)	3.01 ± 0.54 (4.71 ± 0.27*)	4.47 ± 0.39** (6.84 ± 0.41**)
Spleen (g)	0.36 ± 0.12 (0.55 ± 0.15)	0.37 ± 0.06 (0.53 ± 0.07)	0.32 ± 0.07 (0.50 ± 0.10)	0.33 ± 0.06 (0.52 ± 0.08)	0.28 ± 0.07 (0.43 ± 0.09*)
Kidneys (g)	0.70 ± 0.07 (1.09 ± 0.05)	0.71 ± 0.07 (1.04 ± 0.04**)	0.67 ± 0.06 (1.05 ± 0.05)	0.66 ± 0.09 (1.04 ± 0.05*)	0.72 ± 0.07 (1.10 ± 0.07)
Adrenals (mg)	19.2 ± 3.7 (29.9 ± 4.6)	18.8 ± 4.5 (27.5 ± 6.8)	16.9 ± 2.3 (26.8 ± 4.2)	19.9 ± 3.7 (31.4 ± 5.2)	15.4 ± 3.5** (23.5 ± 4.8**)

Values are expressed as the mean ± SD.
 Values in parentheses are relative organ weights (g or mg/100 g body weight).
 *Significantly different from the control group (p < 0.05).
 **Significantly different from the control group (p < 0.01).