

Fig. 1. Isaji, T., et al.




Fig.2. Isaji, T., et al.




Fig. 3. Isaji, T., et al.




Fig. 4. Isaji, T., et al.




Fig. 5. Isaji, T., et al.




Fig.6. Isaji, T., et al.





Fig.7. Isaji, T., et al.



An *N*-glycosylation site on the β -propeller domain of the integrin $\alpha 5$ subunit plays key roles in both its function and site-specific modification by $\beta 1,4$ -*N*-acetylglucosaminyltransferase III*

Yuya Sato[‡], Tomoya Isaji[‡], Michiko Tajiri[§], Shumi Yoshida-Yamamoto^{§,*}, Tsuyoshi Yoshinaka[†], Toshiaki Somehara[†], Tomohiko Fukuda[‡], Yoshinao Wada[§] and Jianguo Gu^{‡,§}
From the [‡] Division of Regulatory Glycobiology, Institute of Molecular Biomembrane and Glycobiology, Tohoku Pharmaceutical University, 4-4-1 Komatsushima, Aoba-ku, Sendai, Miyagi 981-8558, Japan; [§] Department of Molecular Medicine, Osaka Medical Center and Research Institute for Maternal and Child Health, 840 Murodo-cho Izumi, Osaka 594-1101, Japan; *present address. Department of Food Science and Nutrition, Mukogawa Women's University, Nishinomiya, Hyogo 663-8558, Japan; [†] Laboratory Chemicals Division, Wako Pure Chemical Industries, Ltd., 1-2, Doshomachi 3-chome, Chuo-ku, Osaka 540-8605, Japan.

Running title: *N*-glycosylation site-4 on $\alpha 5$ is important for its functional regulation

[†]To whom all correspondence should be addressed. Division of Regulatory Glycobiology, Tohoku Pharmaceutical University, 4-4-1 Komatsushima, Aobaku, Sendai, Miyagi 981-8558, Japan
Tel: 81-2-727-0216; Fax: 81-2-727-0078; E-mail: jgu@tohoku-pharm.ac.jp

Recently, we reported that *N*-glycans on the β -propeller domain of the integrin $\alpha 5$ subunit (S-3,4,5) are essential for $\alpha 5\beta 1$ heterodimerization, expression and cell adhesion. Herein, to further investigate which *N*-glycosylation site is the most important for the biological function and regulation, we characterized the S-3,4,5 mutants in detail. We found that site-4 is a key site that can be specifically modified by *N*-acetylglucosaminyltransferase III (GnT-III). The introduction of bisecting GlcNAc into the S-3,4,5 mutant catalyzed by GnT-III, decreased cell adhesion and migration on fibronectin, whereas overexpression of

N-acetylglucosaminyltransferase V (GnT-V) promoted cell migration. The phenomenon is similar to previous observations that the functions of the wild-type $\alpha 5$ subunit were positively and negatively regulated by GnT-V and GnT-III, respectively, suggesting that the $\alpha 5$ subunit could be duplicated by the S-3,4,5 mutant. Interestingly, GnT-III specifically modified the S-4,5 mutant, but not the S-3,5 mutant. This result was confirmed by E4-PHA lectin blot analysis. The reduction in cell adhesion was consistently observed in the S-4,5 mutant, but not in the S-3,5 mutant cells. Furthermore, mutation of site-4 alone

ASBMB
THE JOURNAL OF BIOLOGICAL CHEMISTRY

resulted in a substantial decrease in E4-PHA lectin staining and suppression of cell spread induced by GnT-III, compared with that of either the site-3 single mutant or wild-type $\alpha 5$. These results, taken together, strongly suggest that *N*-glycosylation of site-4 on the $\alpha 5$ subunit is the most important site for its biological functions. To our knowledge, this is the first demonstration that site-specific modification of *N*-glycans by a glycosyltransferase results in functional regulation.

Glycosylation is a crucial post-translational modification of most secreted and cell surface proteins (1). Glycosylation is involved in a variety of physiological and pathological events, including cell growth, migration, differentiation and tumor invasion. It is well known that glycans play important roles in cell-cell communication, intracellular signal transduction, protein folding and stability (2,3).

Integrins comprise a family of receptors that are important for cell adhesion. The major function of integrins is to connect cells to the extracellular-matrix, activate intracellular signaling pathways, and regulate cytoskeletal formation (4). Integrin $\alpha 5 \beta 1$ is well known as a fibronectin (FN) receptor. The interaction between integrin $\alpha 5$ and FN is essential for cell migration, cell survival and development (5-8). In addition, integrins are *N*-glycan carrier proteins. For example,

$\alpha 5 \beta 1$ integrin contains 14 and 12 putative *N*-glycosylation sites on the $\alpha 5$ and $\beta 1$ subunits, respectively. Several studies suggest that *N*-glycosylation is essential for functional integrin $\alpha 5 \beta 1$. When human fibroblasts were cultured in the presence of 1-deoxymannojirimycin, which prevents *N*-linked oligosaccharide processing, immature $\alpha 5 \beta 1$ integrin appeared on the cell surface, and FN-dependent adhesion was greatly reduced (9). Treatment of purified integrin $\alpha 5 \beta 1$ with *N*-glycosidase F, which cleaves between the innermost *N*-acetylglucosamine (GlcNAc) and asparagine *N*-glycan residues of *N*-linked glycoproteins, prevented the inherent association between subunits and blocked $\alpha 5 \beta 1$ -binding to FN (10).

A growing body of evidence indicates that the presence of the appropriate oligosaccharide can modulate integrin activation. *N*-acetylglucosaminyltransferase III (GnT-III) catalyzes the addition of GlcNAc to mannose that is $\beta 1,4$ -linked to an underlying *N*-acetylglucosamine, producing what is known as a "bisecting" GlcNAc linkage, as shown in Fig.1B. GnT-III is generally regarded as a key glycosyltransferase in *N*-glycan biosynthetic pathways and contributes to inhibition of metastasis. The introduction of a bisecting GlcNAc catalyzed by GnT-III suppresses additional processing and elongation of *N*-glycans. These reactions, which are catalyzed *in vitro* by other glycosyltransferases, such as

N-acetylglucosaminyltransferase V (GnT-V), which catalyzes the formation of β 1,6 GlcNAc branching structures (Fig.1B) and plays important roles in tumor metastasis, do not proceed because the enzymes cannot utilize the bisected *N*-glycans as a substrate. Introduction of the bisecting GlcNAc to integrin α 5 by overexpression of GnT-III, resulted in decreased in ligand binding and down-regulation of cell-adhesion and migration (11-13). Contrary to the functions of GnT-III, overexpression of GnT-V promoted integrin α 5 β 1-mediated cell migration on FN (14). These observations clearly demonstrate that the alteration of *N*-glycan structure affected the biological functions of integrin α 5 β 1. Similarly, characterization of the carbohydrate moieties in integrin α 3 β 1 from non-metastatic and metastatic human melanoma cell lines showed that expression of β 1,6 GlcNAc branched structures was higher in metastatic cells compared with non-metastatic cells, confirming the notion that the β 1,6 GlcNAc branched structure confers invasive and metastatic properties to cancer cells. In fact, Partridge et al (15) reported that GnT-V-modified *N*-glycans containing poly-*N*-acetylglucosamine, the preferred ligand for galectin-3, on surface receptors oppose their constitutive endocytosis, promoting intracellular signaling, and consequently cell migration and tumor metastasis.

In addition, sialylation on the non-reducing terminus of *N*-glycans of α 5 β 1 integrin plays

an important role in cell adhesion. Colon adenocarcinomas express elevated levels of α 2,6 sialylation and increased activity of ST6Gall sialyltransferase. Elevated ST6Gall positively correlated with metastasis and poor survival. Therefore, ST6Gall-mediated hypersialylation likely plays a role in colorectal tumor invasion (16,17). In fact, oncogenic ras up-regulated ST6Gall and, in turn, increased sialylation of β 1 integrin adhesion receptors in colon epithelial cells (18). However, this is not always the case. The expression of hyposialylated integrin α 5 β 1 was induced by phorbol ester-stimulated differentiation in myeloid cells, in which the expression of the ST6Gall was down-regulated by the treatment, increasing FN binding (19). A similar phenomenon was also observed in hematopoietic or other epithelial cells. In these cells, the increased sialylation of the β 1 integrin subunit was correlated with reduced adhesiveness and metastatic potential (20-22). In contrast, the enzymatic removal of α 2,8-linked oligosialic acids from the α 5 integrin subunit inhibited cell adhesion to FN (23). Collectively, these findings suggest that the interaction of integrin α 5 β 1 with FN is dependent on its *N*-glycosylation and the processing status of *N*-glycans.

Because integrin α 5 β 1 contains multi-potential *N*-glycosylation sites, it is important to determine the sites that are crucial for its biological function and regulation. Recently, we found that *N*-glycans on the β -propeller domain (site-3,4

and 5) of the integrin $\alpha 5$ subunit are essential for $\alpha 5\beta 1$ heterodimerization, cell surface expression, and biological function (24). In this study, to further investigate the underlying molecular mechanism of GnT-III-regulated biological functions, we characterized the *N*-glycans on the $\alpha 5$ subunit in detail using genetic and biochemical approaches, and found that site-4 is a key site that can be specifically modified by GnT-III.

EXPERIMENTAL PROCEDURES

Reagents and Antibodies – A monoclonal antibody (mAb) against human integrin $\alpha 5$ subunit (clone1) for Western blot analysis was obtained from BD Biosciences (Franklin Lakes, NJ). For immunoprecipitation, the agarose-conjugated anti-green fluorescent protein (GFP) antibody (RQ2) was obtained from Medical & Biological Laboratories Co. LTD. (Nagoya, Japan). Peroxidase-conjugated anti-mouse IgG was obtained from Cell Signaling Technology, Inc. (Danvers, MA). A VECTASTAIN ABC Kit was purchased from Vector Laboratories, Inc. (Burlingame, CA). Antibodies against GnT-III (33A8) and GnT-V (24B11) were obtained from FUJIREBIO Inc. (Tokyo, Japan). Biotinylated erythroagglutinating phytohemagglutinin (E4-PHA), biotinylated leucoagglutinating phytohemagglutinin (L4-PHA), biotinylated Datura stramonium (DSA) lectin were purchased from Seikagaku Corp. (Tokyo, Japan). For

fluorescence activated cell sorting (FACS) analysis, mouse anti-human $\alpha 5\beta 1$ integrin mAb (HA5, MAB1999) was purchased from Chemicon (Temecula, CA).

Cells and Cell culture – The integrin $\alpha 5$ subunit-deficient CHO K1 cell line (CHO-B2) was a gift from Dr. Rudolf Juliano (School of Medicine, University of North Carolina, Chapel Hill) (25). The CHO-B2 stable expression cells containing various integrin $\alpha 5$ with altered *N*-glycosylation sites were established in our laboratory (24). As shown in Fig.1A, WT indicates CHO-B2 expressing wild type (full *N*-glycosylation sites) integrin $\alpha 5$; S-3,4,5, S-3,5, and S-4,5 show that all *N*-glycosylation sites were removed with site-directed mutagenesis except indicated sites; and D-3 or D-4 represent single mutations at the indicated site. A HeLa cell line was purchased from RIKEN BioResource Center (Tsukuba, Japan). The stable expression of S-3,5 and S-4,5 mutants in HeLa cells were obtained by viral expression vector as mentioned below. These mutants and cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum (FBS), non-essential amino acids (Invitrogen), penicillin (100 units/ml) and streptomycin (100 μ g/ml) (Nacalai Tesque, Inc., Kyoto, Japan) under a humidified atmosphere containing 5% CO₂.

GnT-III, GnT-V and $\alpha 5$ mutants (S-3,5 and

S-4,5) expression with viral vectors - The cDNAs encoding human GnT-III and GnT-V were amplified for cloning into pENTR-D-Topo for Gateway Conversion System (Invitrogen) according to the manufacturer's protocol. The cloned genes were inserted into the virus expression vector, pBABE-puro (Addgene, Inc. Cambridge, MA) accommodated into the Gateway Conversion System using LR clonase reaction. The GnT-III and GnT-V constructs were transfected into Phoenix-Ampho cells with Lipofectamine 2000 (Invitrogen) for production of viral supernatants. The various $\alpha 5$ integrin mutants were infected with the resulting viral supernatant containing 10 $\mu\text{g/ml}$ polybrene (Sigma-Aldrich, St. Louis, MO), and selected with 13 $\mu\text{g/ml}$ of puromycin for two weeks. In case of HeLa cells expressed of S-3,5 and S-4,5 mutants, after virus infection, the infected cells were selected with 2.5 $\mu\text{g/ml}$ of puromycin. For mock transfection, the same protocol was performed using the empty virus expression vector only.

Cell Adhesion assay using 96-well plate - 96-well plates (Corning Inc., PA) were coated with 3 $\mu\text{g/ml}$ of FN at 37 °C for 1 h and blocked with 1% bovine serum albumin (BSA) in DMEM at 37 °C for 1 h. The cells were detached with trypsin containing 1 mM EDTA, resuspended with 0.5 mg/ml trypsin inhibitor (Nacalai Tesque) in DMEM. The suspended cells were centrifuged at 1,000 rpm for 3 min, and diluted to 4×10^5 or $8 \times$

10^5 cells/ml with assay medium, 0.1% BSA in DMEM. One hundred-microliter aliquots of cell suspension were added to each well and the plates were incubated at 37 °C for 20-25 min. After incubation, attached cells were fixed with 25% glutaraldehyde (Nacalai Tesque), and stained with 0.5% crystal violet. The absorbance at 590 nm was measured using an automated micro-titer plate spectrometer, Powerscan® HT (Dainippon Sumitomo Pharma Co., Ltd. Osaka, Japan) operated with Microplate Data Analysis Software, KC4™ (BIO-TEC Instruments, Inc., Winooski, VT). Cell spreading assays were performed as described previously (12,24). After a 20-min incubation, representative fields were observed using a phase-contrast microscopy, and spread cells were counted. The rounded cells were not considered as spread cells.

Cell adhesion kinetics assay using the RT-CES system - The cell adhesion kinetics assay was performed using a real-time cell electronic sensing (RT-CES™) system (ACEA Biosciences, Inc., CA) (26). Briefly, ACEA electro-sensing 16-well plates were coated with 50 μL of 10 $\mu\text{g/ml}$ FN (Sigma) at 37 °C for 1 h and then blocked with 1% BSA in DMEM at 37 °C for 1 h. The cells were detached with trypsin containing 1 mM EDTA, resuspended with 0.5 mg/ml trypsin inhibitor (Nacalai Tesque) in DMEM. The suspended cells were centrifuged at $1,000 \times g$ for 3 min, and diluted to 1×10^5 or 8×10^5 cells/ml with assay medium (0.1% BSA in

DMEM). Fifty-microliter aliquots of the cell suspension were added to each well and the assay was performed using RT-CES SP software. The program was set up such that the cell index was measured every 2 min for 3 h.

Cell migration - Transwells (BD BioCoat™ Control Inserts, 8.0- μ m inserts; BD Biosciences) were coated only on the bottom side with 10 μ g/ml of FN at 37 °C for 1 h. Cells were starved in serum-free medium for 4 h, trypsinized and suspended with 0.5 mg/ml trypsin inhibitor (Nacalai Tesque) in DMEM. Suspended cells were centrifuged and supernatants were removed. The cell pellets were resuspended with assay medium (0.1% BSA in DMEM containing 1% FBS) and diluted to 4×10^5 cells/ml. One-hundred-microliter aliquots of the cell suspension were added to each FN-coated transwell; the cells were then incubated at 37 °C for 3 h. After incubation, cells on the upper side were removed by scraping with a cotton swab. The membranes in the transwells were fixed with 4% paraformaldehyde and stained with 0.5% crystal violet for 30 min. Cells that had migrated to the lower side were counted using a phase-contrast microscope.

Immunoprecipitation, Western Blot and Lectin Blot - Subconfluent cells were washed with phosphate buffered saline (PBS) twice and lysed with ice-cold lysis buffer (1% Triton in Tris-buffered saline (TBS)

containing protease inhibitor cocktail (Nacalai Tesque)). The cell lysates were centrifuged at $15,000 \times g$ for 10 min at 4 °C. The supernatants were obtained and protein concentrations were determined using a BCA™ Protein Assay Kit (PIERCE, Rockford, IL). Equal amounts of protein were incubated with 10 μ l of agarose-conjugated anti-GFP antibody and 15 μ l of Sepharose™ 4B at 4 °C for 1 h. The immunocomplexes were washed twice with ice-cold lysis buffer and then were eluted with SDS sample buffer and boiled for 5 min. The immunoprecipitates were subjected to 6.0% or 7.5% SDS-PAGE, and then were transferred to nitrocellulose membranes. The membranes were blocked with 5% non-fat milk or 5% BSA in TBS for Western blot and Lectin blot, respectively. After blocking, the membranes were incubated with either primary antibody or lectin for 1 h. For Western Blot, membranes were incubated with the secondary antibody conjugated with peroxidase for 1 h and the immunoreactive proteins were visualized using an ECL kit (GE Healthcare Life Sciences). For Lectin Blot, the lectin binding proteins were detected using a VECTASTAIN ABC kit and an ECL kit.

Flow Cytometry - Flow cytometry was performed as described previously (24). Briefly, cells were detached by trypsinizing, incubated with mouse anti-human $\alpha 5\beta 1$ integrin mAb (HA5), followed by Alexa Fluor 647 anti-mouse IgG. Negative controls

underwent the same procedure without primary antibody. The analyses were performed using a FACS Calibur instrument (BD Biosciences) equipped with CELLQuestPro software.

RESULTS

Comparison of N-Glycosylation Patterns on S-3,4,5 α 5 Subunit Mutant in GnT-III and GnT-V Transfectants.

N-Glycosylation is essential for integrin α 5 β 1 heterodimer formation, and, therefore, plays an important role in the biological function of integrin. GnT-III-modified integrin α 5 β 1 decreased cell adhesion and cell migration on FN (12). In contrast to GnT-III, GnT-V specifically modified only the β 1 subunit and up-regulated integrin α 5 β 1-mediated cell migration (14). Recently, we found that three N-glycosylation sites—sites 3, 4 and 5 from the N-terminus of the α 5 subunit—were essential for the biological functions of integrin, such as cell adhesion and migration on FN, and heterodimerization.

The purpose of the present study was to determine whether the S-3,4,5 mutant, which contained only three potential N-glycosylation sites (i.e., sites 3, 4 and 5) had characteristics similar to those of the wild-type α 5 subunit, such as modification by GnT-III and GnT-V as described above. Various α 5 subunit mutants were used in this study as shown in Fig.1A. First, the expression levels of GnT-III and GnT-V in

S-3,4,5 mutant cells that had been transfected with a retrovirus system, were examined by Western blotting (Fig. 2A). Their products were detected by E4-PHA lectin, which specifically recognizes bisecting GlcNAc, and by L4-PHA lectin, which selectively recognizes β 1,6- branching GlcNAc, blots (Fig. 2B) (27,28). As expected, bands corresponding to GnT-III and GnT-V, as well as lectin reactivities of E4-PHA and L4-PHA, were increased in the GnT-III and GnT-V transfectants, respectively (Fig.2A, B). Equal amounts of protein (20 μ g) were loaded in each lane and α -tubulin was used as the loading control. Next, we immunoprecipitated α 5 and detected N-glycans using E4-PHA and L4-PHA lectin blotting (Fig.2 C). The E4-PHA reactivities were much stronger in GnT-III transfectants than those in mock or GnT-V transfectants. This result indicates that both the α 5 and β 1 subunits are targets of GnT-III. In contrast, results of L4-PHA lectin staining indicated that only the β 1 subunit could be modified by GnT-V, consistent with a previous study (14). The reactivities of DSA lectin, which preferentially reacts with branched sugar chains (more than triantennary) (29), consistently showed a significant increase in the β 1 subunit of GnT-V transfectants. A significant decrease in the α 5 subunit of the GnT-III transfectants, further supports the notion that introduction of GnT-III suppresses additional processing and branching formation of N-glycans catalyzed by other endogenous glycosyltransferases,

such as GnT-V and GnT-IV. Although it is not clear if the modification levels of branched *N*-glycans on $\alpha 5$ subunits were enhanced in GnT-V transfectants compared with the levels in mock transfectants, it could be argued that intrinsic GnT-V-mediated glycosylation is enough for occupation of some *N*-glycosylation sites on the $\alpha 5$ subunit, which may be specifically and exclusively modified by GnT-V.

Effects of GnT-III and GnT-V on Integrin-mediated Cell Adhesion and Migration in S-3,4,5 Transfectants.

It is well known that wild-type integrin $\alpha 5 \beta 1$ -mediated cell migration can be positively and negatively regulated by GnT-V and GnT-III, respectively. Therefore, we determined whether modifications of S-3,4,5 mutants could mimic wild-type $\alpha 5$ to affect its biological functions, such as cell adhesion and cell migration. As shown in Fig.3A, cell adhesion on FN was down-regulated in GnT-III transfectants compared with mock and GnT-V transfectants. The cell adhesion kinetics assay using RT-CES also showed the same tendency (Fig.3B). On the other hand, cell migration was determined using the transwell assay as described in "Experimental Procedures." Interestingly, overexpression of GnT-III significantly inhibited cell migration on FN, while GnT-V promoted cell migration, relative to the mock transfectants (Fig. 3C). These results, taken together, suggest that the $\alpha 5$ subunit could be duplicated by the S-3,4,5

mutant. The up-regulation of cell migration in GnT-V transfectants could be ascribed to the *N*-glycans of the $\beta 1$ subunit modified by GnT-V.

To determine whether overexpression of GnT-III or GnT-V affected integrin $\alpha 5 \beta 1$ expression on cell surface, we performed FCAS analysis using anti $\alpha 5 \beta 1$ integrin antibody. As shown in Fig.3D, there were no significant differences in the levels of cell surface expression among the three cell types, indicating that the functional alterations shown in Fig. 3, were due to *N*-glycosylation of the integrin modified by GnT-III or GnT-V.

GnT-III Selectively Modifies N-glycosylation Site-4 on the $\alpha 5$ Subunit.

As described above, the characteristics of the S-3,4,5 mutant are similar to those of wild-type $\alpha 5$. We therefore determined if GnT-III could specifically modify the *N*-glycosylation site among site-3, site-4, and site-5. Because the *N*-glycosylation site-5 is essential for its expression on the cell surface, the mutant did not exhibit biological function, such as cell adhesion (24). Thus, we chose the S-3,5 and S-4,5 mutants for use in future studies.

First, GnT-III was overexpressed in both transfectants. The expression levels of GnT-III were almost same in S-3,5 as in S-4,5 transfectants, which were examined by Western blot using anti-GnT-III antibody (Fig4.A). It is of particular interest that the mutant S-4,5, but not the S-3,5 mutant, was

clearly detected using E4-PHA lectin blot. The intensity of the lectin staining was comparable to that of S-3,4,5 (Fig. 4B). These results, taken together, suggest that the site-3 may not be modified by GnT-III. Because introduction of bisecting GlcNAc into the $\alpha 5$ subunit down-regulates cell adhesion as described above, we checked whether this phenomenon occurred in these mutants. Overexpression of GnT-III in S-4,5 cells consistently inhibited cell adhesion on FN, while the inhibition of cell adhesion was not observed in S-3,5 cells overexpressing GnT-III (Fig. 4C,D). It should be noted that cell adhesion activities of the S-3,5 was similar to that of S-4,5 mutant, since CHO-B2 cells do not express enough endogenous GnT-III to modify integrin as shown in Fig. 2.

To confirm that whether the site-specific modification also happens in endogenous conditions, we introduced the S-3,5 and S-4,5 mutants into HeLa cells that express a relatively higher endogenous GnT-III, to examine the products of GnT-III, as confirmed by E4-PHA lectin blot. To be consistent with the results in overexpressing GnT-III, the E4-PHA lectin staining was clearly detected in S-4,5, but not in S-3,5 of $\alpha 5$ subunit transfectants (Fig.4E). Taken together, these results strongly suggest that GnT-III may specifically modify site-4 on the $\alpha 5$ subunit, which down-regulates its biological functions.

To further elucidate the importance of site-4 for GnT-III modification, we compared

the E4-PHA staining patterns of wild-type (WT) with single mutants, such as site-3 (D-3) or site-4 (D-4), as shown in Fig.1A. These three expression plasmids were co-transfected with GnT-III, and the $\alpha 5$ integrins were immunoprecipitated using an anti-GFP antibody. As shown in Fig.5A, there were no significant differences in the GnT-III expression levels among the three transfectants. The intensity of E4-PHA staining in D-3 cells was less than that in WT cells, but they were comparable. However, the intensity of E4-PHA staining in D-4 cells was substantially less than that in WT or D-3 cells (Fig. 5B). The ratios of E4-PHA staining versus total $\alpha 5$ staining purified from WT, D3 and D4 cells, were 1.0, 0.6 and 0.2, respectively. Furthermore, to directly examine the effects of GnT-III on Site-4 for cell adhesion, we compared cell spreading of the D4 mutant with WT of $\alpha 5$ integrin. As expected, GnT-III significantly down regulated cell spreading on FN in WT transfectants, whereas the deletion of site-4 abolished the suppression of cell spread induced by GnT-III in D4 transfectants (Fig. 5C). Taken together, these results clearly show that *N*-glycosylation of site-4 is critical and effective for GnT-III modification.

DISCUSSION

In the present study, we intensively investigated the effects of *N*-glycosylation on the β -propeller of the integrin $\alpha 5$ subunit on

its biological functions such as cell adhesion and cell migration, and found that site-4 is essential and effective for GnT-III modification among 14 potential *N*-glycosylation sites. To our knowledge, this is the first report to clearly demonstrate that a glycosyltransferase of *N*-glycosylation can specifically modify a *N*-glycosylation site among multiple potential sites, and effectively regulates its biological functions.

Integrins can be activated by inside-out signaling mechanisms that trigger global conformational changes, which ultimately modulate integrin ligand-affinity. It is apparent that integrin activity can be regulated by other mechanisms, such as posttranscriptional modification, *N*-glycosylation. Altered integrin glycosylation has been associated with tumorigenesis, autoimmune disease, chronic inflammation, and cell adhesion events (11,30). In particular, *N*-glycosylation of the integrin $\alpha 5$ and $\beta 1$ subunits appears to be important for both structure and function. It has been reported that *N*-glycosylation of both the $\alpha 5$ and $\beta 1$ subunits is necessary for $\alpha 5\beta 1$ heterodimerization, and its binding to FN. Moreover, changes in integrin glycan composition, resulting from forced expression of selected glycosyltransferases, i.e., "remodeling," reportedly modulate integrin functions as described above. However, most of these earlier studies examined only total changes, but without individual information. Therefore, the exact molecular mechanisms by which

N-glycosylation of site(s) or glycan(s) occurs remain unknown. Recently, we used site-directed mutagenesis to determine that *N*-glycosylation site-5 on the β -propeller plays an important role in the assembly of the integrin for its expression on the cell surface (24). These observations prompted us to determine whether there are specific *N*-glycosylation sites that regulate its biological functions. Here, we clearly showed that site-4 is a key *N*-glycosylation site for the biological function of $\alpha 5$ subunit that is effectively modified by GnT-III. Taken together these results indicate that individual *N*-glycosylation sites may have unique functions.

Although the molecular mechanism by which bisecting GlcNAc is introduced into site-4, inhibiting its biological function remain unknown, we speculate that the effect of altered glycosylation of site-4 may be related to conformational changes in the key functional regions of the β -propeller domain of the $\alpha 5$ subunit that are critical for integrin activation. In fact, the β -propeller domain has been postulated to be required for effective interaction between $\alpha 5\beta 1$ integrin and its ligand (31). In contrast, the crystal structure of integrin $\alpha V\beta 3$ has been successfully determined, and the main contact between the αV and $\beta 3$ subunits is the β -propeller on the α and A domain on $\beta 3$ with hydrophobic, ionic, and mixed contacts (32,33). Because the $\alpha 5$ subunit has 47% homology to αV , Mould et al. made a homologous modeling structure of $\alpha 5\beta 1$ (34).

Based on the model, the $\alpha 5$ subunit seems to be surrounded by *N*-glycans, which explains the dissociation of the $\alpha\beta$ heterodimer that occurs when $\alpha 5\beta 1$ is deglycosylated by treatment with PNGase F or removal of *N*-glycans on the β -propeller. Very recently, Liu et al. used a molecular modeling approach to study the effects of altered glycosylation on the I-like domain of the $\beta 1$ subunit, which is the partner of the β -propeller of the α subunit. These researchers found that $\alpha 2,6$ sialic acid affected the interactions between *N*-glycans and the I-like domain, which in turn altered the accessibility of the loop that determines specificity of ligand binding (35). In fact, the remodeling of *N*-glycans by GnT-III affects either the branching formations catalyzed by GnT-V and GnT-IV, or the sialylation on the terminus of the *N*-glycans (11,36). Therefore, a possible mechanism by which *N*-glycans are involved in the $\alpha\beta$ interaction or conformational arrangement is that an unknown lectin domain may exist on the α or β subunit. The lectin domain of $\alpha M\beta 2$ integrin is associated with GlcNAc on the non-reducing terminal of sugar chains on platelets, facilitating their phagocytosis (37,38). These studies further support the observation that modification of bisecting GlcNAc on site-4 of the β -propeller may be critical for the regulation of its biological functions, which may shed light on the structural studies.

It is of interest to understand why GnT-III specifically and effectively modifies site-4 of

the 14 putative *N*-glycosylation sites. There is currently no detailed information available regarding this observation, but several explanations have been proposed. First, *N*-glycosylation occurs on site-4 because it provides the easiest access for GnT-III. Because the integrin $\alpha 5$ crystal structure is currently unavailable this hypothesis cannot be proven. Second, GnT-III may associate with some other molecules, which define the specificities for protein or peptide substrates. Reportedly, protein *O*-mannosyltransferase 1 (POMT1) and its homolog POMT2 are responsible for catalyzing the first step in *O*-mannosyl glycan, which is important for muscle and brain development (39). Interestingly, Manya et al., reported that formation of a POMT1-POMT2 complex is essential for POMT activity (40). Only two peptides derived from the mucin domain of α -dystroglycan are highly *O*-mannosylated by POMT, but no *O*-mannosylation occurs in mucin tandem repeat peptides (41). Similarly, complex formation is also important for T-synthase (core 1 $\beta 1,3$ -GalT) activity. Ju et al. reported that Tn syndrome, a rare autoimmune disease, in which subpopulations of blood cells of all lineages carry an incompletely glycosylated membrane glycoprotein, known as the Tn antigen, is associated with a somatic mutation in *Cosmc*, a gene on the X chromosome that encodes a molecular 'chaperone' that is required for the proper folding and hence full activity of T-synthase (42,43). Indeed, it has been reported that caveolin-1 may co-localize

with GnT-III to regulate its localization and activity (44). Those results, taken together, suggest that glycosyltransferase complex formation may play a crucial role in determination of both activity and substrate specificity. The detailed molecular mechanism remains further study.

This study specifically focused on *N*-glycosylation of the integrin $\alpha 5$ subunit.

To fully understand the effects of the *N*-glycans on integrin structure and function, it will be necessary, in future studies, to investigate the interaction of glycans with glycans or peptides of integrin. The current study has also implications for engineering $\alpha 5$ that contains the glycans necessary for its activation, which may facilitate the study of its crystal structure.

REFERENCES

1. Apweiler, R., Hermjakob, H., and Sharon, N. (1999) *Biochim Biophys Acta* **1473**(1), 4-8
2. Dwek, R. A. (1995) *Biochem Soc Trans* **23**(1), 1-25
3. Saxon, E., and Bertozzi, C. R. (2001) *Annu Rev Cell Dev Biol* **17**, 1-23
4. Hynes, R. O. (2002) *Cell* **110**(6), 673-687
5. George, E. L., Georges-Labouesse, E. N., Patel-King, R. S., Rayburn, H., and Hynes, R. O. (1993) *Development* **119**(4), 1079-1091
6. Goh, K. L., Yang, J. T., and Hynes, R. O. (1997) *Development* **124**(21), 4309-4319
7. Watt, F. M., and Hodivala, K. J. (1994) *Curr Biol* **4**(3), 270-272
8. Yang, J. T., Rayburn, H., and Hynes, R. O. (1993) *Development* **119**(4), 1093-1105
9. Akiyama, S. K., Yamada, S. S., and Yamada, K. M. (1989) *J Biol Chem* **264**(30), 18011-18018
10. Zheng, M., Fang, H., and Hakomori, S. (1994) *J Biol Chem* **269**(16), 12325-12331
11. Gu, J., and Taniguchi, N. (2004) *Glycoconj J* **21**(1-2), 9-15
12. Isaji, T., Gu, J., Nishiuchi, R., Zhao, Y., Takahashi, M., Miyoshi, E., Honke, K., Sekiguchi, K., and Taniguchi, N. (2004) *J Biol Chem* **279**(19), 19747-19754
13. Zhao, Y., Nakagawa, T., Itoh, S., Inamori, K., Isaji, T., Kariya, Y., Kondo, A., Miyoshi, E., Miyazaki, K., Kawasaki, N., Taniguchi, N., and Gu, J. (2006) *J Biol Chem* **281**(43), 32122-32130
14. Guo, H. B., Lee, I., Kamar, M., Akiyama, S. K., and Pierce, M. (2002) *Cancer Res* **62**(23), 6837-6845
15. Partridge, E. A., Le Roy, C., Di Guglielmo, G. M., Pawling, J., Cheung, P., Granovsky, M., Nabi, I. R., Wrana, J. L., and Dennis, J. W. (2004) *Science* **306**(5693), 120-124
16. Seales, E. C., Jurado, G. A., Brunson, B. A., Wakefield, J. K., Frost, A. R., and Bellis, S.

- L. (2005) *Cancer Res* **65**(11), 4645-4652
17. Shaikh, F. M., Seales, E. C., Clem, W. C., Hennessy, K. M., Zhuo, Y., and Bellis, S. L. (2008) *Exp Cell Res*
 18. Seales, E. C., Jurado, G. A., Singhal, A., and Bellis, S. L. (2003) *Oncogene* **22**(46), 7137-7145
 19. Semel, A. C., Seales, E. C., Singhal, A., Eklund, E. A., Colley, K. J., and Bellis, S. L. (2002) *J Biol Chem* **277**(36), 32830-32836
 20. Pretzlaff, R. K., Xue, V. W., and Rowin, M. E. (2000) *Cell Adhes Commun* **7**(6), 491-500
 21. Kawano, T., Takasaki, S., Tao, T. W., and Kobata, A. (1993) *Int J Cancer* **53**(1), 91-96
 22. Dennis, J., Waller, C., Timpl, R., and Schirmacher, V. (1982) *Nature* **300**(5889), 274-276
 23. Nadanaka, S., Sato, C., Kitajima, K., Katagiri, K., Irie, S., and Yamagata, T. (2001) *J Biol Chem* **276**(36), 33657-33664
 24. Isaji, T., Sato, Y., Zhao, Y., Miyoshi, E., Wada, Y., Taniguchi, N., and Gu, J. (2006) *J Biol Chem* **281**(44), 33258-33267
 25. Schreiner, C. L., Bauer, J. S., Danilov, Y. N., Hussein, S., Sczekan, M. M., and Juliano, R. L. (1989) *J Cell Biol* **109**(6 Pt 1), 3157-3167
 26. Solly, K., Wang, X., Xu, X., Strulovici, B., and Zheng, W. (2004) *Assay Drug Dev Technol* **2**(4), 363-372
 27. Cummings, R., and Kornfeld, S. (1982) *J Biol Chem* **257**(19), 11230-11234
 28. Yamashita, K., Hitoi, A., and Kobata, A. (1983) *J Biol Chem* **258**(24), 14753-14755
 29. Yamashita, K., Totani, K., Ohkura, T., Takasaki, S., Goldstein, I. J., and Kobata, A. (1987) *J Biol Chem* **262**(4), 1602-1607
 30. Bellis, S. L. (2004) *Biochim Biophys Acta* **1663**(1-2), 52-60
 31. Mould, A. P., Askari, J. A., and Humphries, M. J. (2000) *J Biol Chem* **275**(27), 20324-20336
 32. Xiong, J. P., Stehle, T., Zhang, R., Joachimiak, A., Frech, M., Goodman, S. L., and Arnaout, M. A. (2002) *Science* **296**(5565), 151-155
 33. Xiong, J. P., Stehle, T., Diefenbach, B., Zhang, R., Dunker, R., Scott, D. L., Joachimiak, A., Goodman, S. L., and Arnaout, M. A. (2001) *Science* **294**(5541), 339-345
 34. Mould, A. P., Symonds, E. J., Buckley, P. A., Grossmann, J. G., McEwan, P. A., Barton, S. J., Askari, J. A., Craig, S. E., Bella, J., and Humphries, M. J. (2003) *J Biol Chem* **278**(41), 39993-39999
 35. Liu, Y., Pan, D., Bellis, S. L., and Song, Y. (2008) *Proteins* **73**(4): 989-1000
 36. Koyota, S., Ikeda, Y., Miyagawa, S., Ihara, H., Koma, M., Honke, K., Shirakura, R.,