

Highly-purified *Helicobacter pylori* LPS preparations induce weak inflammatory reactions and utilize Toll-like receptor 2 complex but not Toll-like receptor 4 complex

Shin-ichi Yokota¹, Takahiro Ohnishi², Masashi Muroi², Ken-ichi Tanamoto³, Nobuhiro Fujii¹ & Ken-ichi Amano⁴

¹Department of Microbiology, Sapporo Medical University School of Medicine, Chuo-ku, Sapporo, Japan; ²Division of Microbiology, National Institute of Health Sciences, Setagaya-ku, Tokyo, Japan; ³Division of Food Additives, National Institute of Health Sciences, Setagaya-ku, Tokyo, Japan; and

⁴Bioscience Education and Research Center, Akita University, Akita, Japan

Correspondence: Ken-ichi Amano, Bioscience Education and Research Center, Akita University, 1-1-1, Hondo, Akita 010-8543, Japan. Tel.: +81 18 884 6452; fax: +81 18 884 6190; e-mail: amanocr@med.akita-u.ac.jp

Received 25 October 2006; revised 21 May 2007; accepted 22 May 2007.
First published online 20 July 2007.

DOI: 10.1111/j.1574-695X.2007.00288.x

Editor: Artur Ulmer

Keywords

Helicobacter pylori; LPS; Toll-like receptor; NF- κ B; interleukin-8.

Introduction

Helicobacter pylori is a micro-aerophilic and spiral Gram-negative bacterium that can colonize gastric epithelial cells or gastric mucin. *Helicobacter pylori* is now recognized as a causative factor of chronic gastritis, gastroduodenal ulcer, gastric cancer and mucosa-associated lymphatic tissue lymphoma (Cover & Blaser, 1996; Parsonnet, 1998). Cell injury and inflammation caused by long-term chronic *H. pylori* infection is believed to underlie these diseases. Several bacterial products probably mediate cytotoxicity and the inflammatory reaction. Among these are vacuolating toxin (VacA), ammonium ions produced by urease, cytotoxin-associated gene A (CagA) protein, and the type IV secretion system, reviewed in Covacci *et al.* (1999) and Kusters *et al.* (2006). In general, Gram-negative bacterial LPSs are key inflammation inducers. However, *H. pylori* LPSs show

Abstract

Helicobacter pylori is recognized as an etiologic agent of gastroduodenal diseases. Among toxic substances produced by *H. pylori*, LPS exhibits extremely low endotoxic activity as compared to the typical LPSs, such as that produced by *Escherichia coli*. We found that the LPS-low-responder stomach cancer cell line MKN28, which expresses Toll-like receptor 4 (TLR4) at extremely low levels, showed similar levels of interleukin-8 (IL-8) induction by *H. pylori* or *E. coli* LPS preparations. Weak IL-8 induction by *H. pylori* LPS preparations was suppressed by expression of a dominant negative mutant of TLR2 but not of TLR4. Data from luciferase reporter analysis indicated that cotransfection of TLR2–TLR1 or TLR2–TLR6 was required for the activation induced by *H. pylori* LPS preparations. In conclusion, the *H. pylori* LPS preparations significantly induce an inflammatory reaction via the receptor complex containing TLR2–TLR1 or TLR2–TLR6 but not that containing TLR4. The TLR2–TLR1 complex was preferentially recognized by the *H. pylori* LPS preparations over the TLR2–TLR6 complex. Whereas the magnitude of response to *H. pylori* LPS preparation was markedly less than that to *E. coli* LPS preparation in LPS-high-responder cells strongly expressing TLR4, it was comparable to that of *E. coli* LPS in low-responder cells expressing negligible amount of TLR4.

extremely low endotoxic activity in comparison to typical Gram-negative bacterial LPSs, such as those from *Escherichia coli* (Nielsen *et al.*, 1994; Perez-Perez *et al.*, 1995; Yokota *et al.*, 2003; Hynes *et al.*, 2004).

Various substances produced by microorganisms are recognized by host receptors and trigger an innate immune response. Toll-like receptors (TLRs) have been shown to play important roles in the pattern recognition of microbial components, so the microbial substances recognized by TLRs are called 'pathogen-associated molecular pattern' (PAMP). At least 13 TLR family members in vertebrates and 10 TLRs in humans have been reported (Akira *et al.*, 2001; Barton & Medzhitov, 2002). These proteins recognize diverse PAMPs. Typical LPSs are recognized by a TLR4 complex consisting of TLR4, CD14 and MD2 expressed on host cells. Currently, it is controversial which TLR contributes to signal transduction by *H. pylori* LPS. Some reports

suggest that *H. pylori* LPS transduces signals via a TLR4 system. Kawahara *et al.* (2001) reported that TLR4 is required for induction of mitogen oxidase 1 by *H. pylori* LPS. Ishihara *et al.* (2004) reported that *H. pylori* infection upregulated MD2 expression, and upregulated MD2 increased cell surface expression of the MD2-TLR4 complex and the subsequent response to *H. pylori* LPS. Ogawa *et al.* (1997) reported that chemically synthesized *H. pylori* lipid A activates cytokine production in macrophages, and that the induction is inhibited by anti-TLR4 antibody. In contrast, Smith *et al.* (2003) and Lepper *et al.* (2005) reported that TLR2 is required for signal transduction induced by *H. pylori* LPS. Notably, Lepper *et al.* (2005) suggested that LPSs derived from some *H. pylori* strains antagonize TLR4 signaling activated by a typical LPS. Here, we investigated which TLR responded to *H. pylori* LPS in stomach adenocarcinoma cell lines, whose response to LPS is very weak and comparable in the case of both *E. coli* and *H. pylori* LPS.

Materials and methods

Bacterial strains and preparation of LPS

Helicobacter pylori clinical strains were isolated from the biopsy specimens obtained from patients with chronic gastritis (CG), gastric ulcer (GU), duodenal ulcer (DU), and gastric cancer (CA). After fewer than three passages of laboratory culture, these bacteria were grown on brucella broth (BBL, Cockysville, MD) supplemented with 10% (v/v) horse serum at 37 °C for 2 days in micro-aerophilic conditions, using the Campyack system (BBL). The organisms were collected by centrifugation (10 000 g, 30 min), washed with de-ionized water twice, and lyophilized. The conventional LPS preparation was obtained from the lyophilized cells by the hot phenol/water method (Yokota *et al.*, 2000). In some experiments, a highly purified LPS preparation, which was further purified in several steps as follows, was used, due to the potential influence of contaminants. The conventional LPS preparation was treated with 10 µg mL⁻¹ DNase I (Roche Diagnostic, Tokyo, Japan), 10 µg mL⁻¹ RNase A (Qiagen, Hilden, Germany), 2 µg mL⁻¹ lipoprotein lipase from bovine milk (Sigma-Aldrich, St Louis, MO) and 10 µg mL⁻¹ lipoprotein lipase from *Pseudomonas* sp. (Sigma-Aldrich) at 37 °C for 12 h in 50 mM sodium phosphate buffer (pH 7.2). Finally, proteinase K (Wako Junyaku, Tokyo, Japan) was added to the reaction mixture at a concentration of 20 µg mL⁻¹, and the mixture was incubated at 60 °C for 4 h. The resulting material was subjected to Octyl-Sepharose column chromatography (GE Healthcare Bio-Science, Piscataway, NJ). The column was equilibrated with 0.5 M NaCl in 0.1 M sodium/acetate buffer (pH 4.7). The column was washed with 30% propanol, and then LPS was eluted with a linear gradient of 30–60% propanol.

Fractions were analyzed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) with silver staining as previously described (Hitchcock & Brown, 1983). For buffer exchange, the pooled LPS fraction was applied to a PD-10 desalting column (GE Healthcare Bio-Science) equilibrated with 0.2% triethylamine. The material eluted at void volume was pooled, lyophilized and used as a highly purified LPS preparation.

Campylobacter jejuni strains 81–176 and OH4832, which were isolated from a gastroenteritis patient and a Guillain-Barré syndrome patient, respectively, were provided by Dr Misawa (Miyazaki University, Miyazaki, Japan). *Campylobacter jejuni* LPS was prepared by phenol/water extraction (Yokota *et al.*, 2000). LPSs derived from *E. coli* O111:B4 (wild type) and D31m4 (Re mutant) were purchased from LIST Biochemical Laboratories (Campbell, CA).

Cell lines

The human gastric carcinoma cell lines MKN28 and MKN45 were obtained from the Japanese Collection of Research Biosources (JCRB, Ibaraki, Japan). The human uroepithelial cell line T24 and the human monocytic cell line THP-1 were obtained from the American Type Culture Collection (ATCC, Manassas, VA). The human embryonic kidney 293 cell line was obtained from the Human Science Research Resource Bank (Tokyo, Japan). MKN28, MKN45, T24 and 293 were routinely cultured in Dulbecco's modified minimum essential medium supplemented with 10% (v/v) heat-inactivated fetal bovine serum (FBS), penicillin (100 U mL⁻¹), and streptomycin (100 µg mL⁻¹). THP-1 was cultured in RPMI-1640 containing 10% (v/v) FBS, penicillin (100 U mL⁻¹), and streptomycin (100 µg mL⁻¹).

Expression plasmids

Human cDNAs encoding TLR2 and MD2 were prepared from RNA derived from THP-1 cells by reverse transcription (RT)-PCR. Human TLR4 cDNA was prepared from human spleen total RNA (OriGene Technologies, Rockville, MD) by RT-PCR. Human CD14 cDNA was kindly provided by Dr Shunsuke Yamamoto (Medical College of Oita, Oita, Japan). cDNAs encoding human TLR2, TLR4, MD2 and CD14 minus signal peptide sequences were subcloned into the downstream portion of a modified pcDNA3 vector (Invitrogen, Carlsbad, CA). The product generated from this vector contains the coding sequence for the preprotrypsin signal peptide sequence preceding the N-terminal equine infectious anemia virus tag epitope. Expression plasmids encoding TLR5, dominant negative TLR2 and dominant negative TLR4 were kindly donated by Dr M.F. Smith (University of Virginia) (Smith *et al.*, 2003). Dominant negative mutants had deletions of the C-terminal portion of the molecules

(TLR2 truncated at amino acid 670, and TLR4 at amino acid 700). Expression plasmids encoding TLR1 and TLR6 were kindly donated by Dr Ken-ichiro Shibata (Hokkaido University, Sapporo, Japan) (Fujita et al., 2003).

Enzyme-linked immunosorbent assay (ELISA)

The amounts of interleukin-8 (IL-8) in culture supernatants were determined with an ELISA Development kit for human IL-8 (R&D Systems, Minneapolis, MN).

Western blotting analysis

SDS-PAGE and Western blotting were carried out as described previously (Yokota et al., 2000). A mouse anti-CD14 monoclonal antibody (UCH-M1) and a rabbit anti-TLR2 polyclonal antibody (H-175) were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Rabbit anti-TLR1 polyclonal antibodies and mouse anti-TLR6 monoclonal antibodies were purchased from Abcam (Cambridge, UK). Alkaline phosphatase-labeled goat anti-mouse or anti-rabbit IgG antibodies were purchased from BioSource International (Camarillo, CA), and used as secondary antibodies. Specific binding was detected using tetrazolium bromochloroindolylphosphate/Nitro Blue Tetrazolium as a developing substrate.

Real-time RT-PCR

Total RNA was isolated from cells using an RNeasy Mini kit (Qiagen). The cDNA was prepared from total RNA with a Reverse Transcription System kit (Promega, Madison, WI) using random primers (GE Healthcare Bio-Science). Real-time PCR was performed using a TaqMan Gene Expression Assay kit (Applied Biosystems, Foster City, CA) with a 7300 Fast Real-Time PCR system (Applied Biosystems). mRNA levels of TLR2, TLR4, CD14, MD2 and glyceraldehyde-3-phosphate dehydrogenase were determined. The mRNA levels were normalized with the mRNA levels of glyceraldehyde-3-phosphate dehydrogenase.

Flow cytometry

Flow cytometry was performed using a FACSCalibur (Beckton-Dickinson, San Jose, CA). Phycoerythrin-labeled mouse monoclonal antibodies against TLR2 (clone TL2.1), TLR4 (HTA125) and CD14 (61D3), and their isotype controls, phycoerythrin-labeled mouse monoclonal IgG₁ and IgG_{2a}, were purchased from eBioscience (San Diego, CA).

Luciferase reporter gene assay

A luciferase reporter plasmid carrying the promoter region of E-selectin (pELAM-L) was constructed by inserting a PCR fragment (-730 to +52) of the E-selectin promoter

into the SacI-HindIII site of the pGL3 Basic vector (Promega, Madison, WI). After plating of 293 cells in 12-well plates (2×10^5 well⁻¹), cells were transfected by the calcium phosphate precipitation method with the indicated expression plasmids (0.02 µg of TLR1, TLR2, and TLR6), together with 0.02 µg of CD14 expression plasmid, 0.1 µg of pELAM-L luciferase reporter plasmid, and 0.0025 µg of pRL-TK (Promega) for normalization. In some experiments, CD14 expression vector was eliminated, as described in 'Results'. After 24 h, cells were stimulated with LPS for 6 h. Reporter activity was measured according to the manufacturer's (Promega) recommendations.

Pam₃CSK₄ (as a TLR2-TLR1 agonist), MALP-2 (as a TLR2-TLR6 agonist) and recombinant flagellin protein of *Salmonella muenchen* (as a TLR5 agonist) were purchased from Bachem (Bubendorf, Switzerland), Alexis (Lausen, Switzerland), and Calbiochem (San Diego, CA), respectively.


Results

Relationship of TLR4 expression levels to LPS responsiveness in various cell lines

We examined responsiveness to *H. pylori* LPS preparations as compared with *E. coli* LPS preparations of stomach adenocarcinoma cell lines (MKN28 and MKN45) and a uroepithelial cell line (T24). The *H. pylori* LPSs used in this study were from six strains. LPS preparations derived from strains GU2 and DU2 are smooth types carrying the highly antigenic epitope. LPS preparations derived from strains CA2 and CA6 are smooth types carrying the weakly antigenic epitope, and are frequently found in strains derived from gastric cancer patients. LPS preparations derived from CG10 and DU8 are rough types lacking the O-polysaccharide chain (Yokota et al., 1997; Yokota et al., 2000).

The LPS-high-responder cell line T24 produced high levels of IL-8 after stimulation with an *E. coli* wild-type LPS preparation (Fig. 1a). The *E. coli* rough LPS preparation was less active than the smooth LPS preparation. *Helicobacter pylori* LPS preparations from all strains showed very weak (500-fold less than *E. coli* smooth LPS) IL-8 induction. The weak response to the *H. pylori* LPS preparation was consistent with previous reports (Nielsen et al., 1994; Perez-Perez et al., 1995; Yokota et al., 2003; Hynes et al., 2004). MKN28 showed a weak response to the *E. coli* LPS preparation (Fig. 1c). Interestingly, similar levels of IL-8 induction were observed with *E. coli* and *H. pylori* LPS preparations. MKN45 cells showed responses between those of T24 and MKN28 cells (Fig. 1b). Here, we also examined *H. pylori* LPS from which probable contaminants were extensively removed. Such contaminants were destroyed by treatment with proteinase K, DNase, RNase and lipoprotein lipases,

Fig. 1. IL-8 induction by LPS derived from *Helicobacter pylori* and *Escherichia coli* in the human uroepithelial cell line T24 and in human gastric carcinoma lines MKN28 and MKN45. Cells were treated with various concentrations of LPS for 24 h. Concentrations of IL-8 in the resulting culture supernatant were determined by ELISA. The LPSs used were derived from *Helicobacter pylori* strain GU2 (○), DU2 (●), CA2 (△), CA6 (▲), CG10 (□), and DU8 (■), and *Escherichia coli* O111 : B4 wild-type (✱) and Re mutant (+) strains. The data represent results for the conventional LPS preparations. Similar results were obtained when the highly purified LPS preparation was used.



and the LPSs were further subjected to hydrophobic interaction chromatography. The lipoprotein lipase derived from *Pseudomonas* treatment decreased IL-8 induction by MALP-2 and Pam₃CSK₄ by < 5% in THP-1 cells. Lipoprotein lipase derived from bovine milk treatment decreased IL-8 induction by Pam₃CSK₄ by < 2% (data not shown). Hashimoto *et al.* (2004) showed that lipoprotein lipase could reduce the activity of contaminated lipoprotein in a preparation of *Pseudomonas gingivalis* LPS isolated with a conventional method. We further performed hydrophobic interaction chromatography on an Octyl-Sepharose column. This chromatographic method has been shown to be suitable for removal of contaminants in the conventional LPS preparation (Hashimoto *et al.*, 2004; Lepper *et al.*, 2005). After enzyme treatment and hydrophobic interaction chromatography, our LPS preparation did not show significantly altered reactivity (data not shown). This suggests that the activity of our *H. pylori* LPS preparation obtained by the conventional method is due to LPS itself and not to contaminants. *Helicobacter pylori* LPS preparations from GU2 and DU2 showed weaker IL-8 induction than did the *H. pylori* LPS preparation from other strains in all cells tested (Fig. 1).

We analyzed the expression of LPS receptor components and other TLRs in these cell lines. Cell surface expression of TLR4, CD14 and MD2 is required for recognition and signal transduction of a typical LPS, such as *E. coli* LPS, by host cells. TLR2 and MD2 mRNA levels were comparable in all three cell lines. TLR4 and CD14 mRNA levels were in the following order, from higher to lower: T24 > MKN45 > MKN28 (Fig. 2a). Cell surface protein expression levels as determined by flow cytometry were consistent with mRNA levels (Fig. 2b). This expression level of TLR4 and CD14 was positively correlated with responsiveness to the *E. coli* LPS preparation. TLR1 and TLR6 mRNA levels were in the following order, from higher to lower: MKN45 > T24 > MKN28. In particular, TLR6 mRNA was not detectable in MKN28 cells.

Effect of dominant negative mutant of TLR4 on IL-8 production induced by *E. coli* and *H. pylori* LPS preparations

To examine the contribution of TLR4 to *E. coli* or *H. pylori* LPS-induced IL-8 production, we overexpressed dominant negative mutants of TLR4 or TLR2 in LPS-high-responder T24 cells (Fig. 3). *Escherichia coli* LPS-induced IL-8 production was markedly suppressed by expression of dominant negative TLR4 but not by dominant negative TLR2. On the other hand, IL-8 production induced by the *H. pylori* LPS preparation was significantly suppressed by dominant negative TLR2 but not by dominant negative TLR4. These results indicate that TLR2, but not TLR4, is necessary for IL-8 induction by the highly purified *H. pylori* LPS preparations.

Effect of *E. coli*, *C. jejuni* and *H. pylori* LPS preparations on nuclear factor- κ B (NF- κ B) activation in 293 cells expressing TLRs and related molecules

Next, we examined NF- κ B activation induced by *H. pylori*, *C. jejuni* and *E. coli* LPS preparations in 293 cells coexpressing TLR4, CD14 and MD2 as determined by a luciferase reporter assay (Fig. 4). LPS preparations derived from *C. jejuni*, which, like *Helicobacter*, belongs to the *Campylobacteriales* family, activated NF- κ B to a similar extent as *E. coli* LPS preparations. *Helicobacter pylori* LPS preparations had little effect on NF- κ B activation. The results indicate that the *H. pylori* LPS preparation does not transduce cellular signaling via TLR4, in contrast to other Gram-negative bacterial LPSs.

To address the contribution of other TLRs to *H. pylori* LPS signal transduction, we expressed various TLRs and related molecules and undertook similar assays in 293 cells, which lack TLRs (Fig. 5). 293 cells expressing TLR2, TLR4-MD2 or TLR5, plus CD14 did not respond to *H. pylori* LPS preparations, even at a concentration of 10 μ g mL⁻¹. Next, we coexpressed TLR1 or TLR6 with




Fig. 2. Expression of (a) mRNA levels (by real-time RT-PCR) of TLR2, TLR4, TLR1, TLR6, CD14 and MD2, and (b) cell surface protein levels (by flow cytometry) of TLR2, TLR4 and CD14 in MKN28, MKN45 and T24 cells. THP-1 cells were used as positive controls. The relative amounts of mRNA levels in relation to THP-1 cells (expression levels in THP-1 cells taken as 1) were calculated. (b) Hatched lines indicate cells incubated with isotype-matched control antibodies.


TLR2 plus CD14. CD14 expression is required for recognition of ligands for TLR2–TLR1 and TLR2–TLR6 (Schroder *et al.*, 2004). We also found that coexpression of CD14 significantly elevated TLR2–TLR1- and TLR2–TLR6-mediated responses (data not shown). We examined the expression levels of TLR2, TLR1, TLR6 and CD14 in each transfectant by Western blotting (Fig. 5b). The expression levels of each of these molecules was not affected by the coexpression of other TLRs and CD14. This suggested that differences in responsiveness of these TLR systems were not due to different expression levels of TLRs and CD14. Significant NF- κ B activation in response to *H. pylori* LPS was observed in 293 cells expressing TLR2–TLR1 or TLR2–TLR6. The TLR2–TLR1 system showed stronger activation than the TLR2–TLR6 system. Both TLR2–TLR1 and TLR2–TLR6 agonists, as positive controls, responded well. The results suggested that the *H. pylori* LPS preparations preferentially transduced signals via TLR2–TLR1 rather than TLR2–TLR6. Significant activation via the TLR2–TLR1 system was observed for LPS preparations derived from CA2 and GU2 strains at concentrations of 1 and 10 $\mu\text{g mL}^{-1}$, respectively. The CA2 LPS preparation was

a more potent NF- κ B activator (Fig. 5) and IL-8 inducer (Fig. 1) than the GU2 LPS preparation. However, the magnitude of NF- κ B activation by *H. pylori* LPS preparations was markedly less than that of a conventional agonist (Fig. 5). Thus, the degree of NF- κ B activation by the *H. pylori* LPS preparation appeared to correlate with the degree of IL-8 induction. These results suggest that the *H. pylori* LPS preparation weakly activates NF- κ B via hetero-oligomers containing TLR2 but not TLR4, and transduce a signal preferentially via TLR2–TLR1 rather than TLR2–TLR6.

Discussion

Helicobacter pylori LPS exhibits extremely weak endotoxic activities, such as NF- κ B-mediated inflammatory reactions. LPS preparations from several bacteria, such as *Porphyromonas gingivalis* (Ogawa *et al.*, 1997), *Bacteroides fragilis* (Lindberg *et al.*, 1990), *Chlamydia trachomatis* (Heine *et al.*, 2003) and *Rhodobacter* species (Rose *et al.*, 1995; Weckesser & Mayer, 1988) also show weak endotoxic activity. Whereas the typical LPS transduces signals via the TLR4 complex containing MD2 and CD14, the identity of signal-


Fig. 3. Effect of overexpression of dominant negative mutants of TLR2 or TLR4 on LPS-induced IL-8 induction in T24 cells. Expression plasmids encoding dominant negative mutants of TLR2 (\blacktriangle) or TLR4 (\blacksquare), or empty plasmid (\bullet), were transfected into T24 cells. After 24 h of transfection, cells were treated with *Helicobacter pylori* LPS derived from strain GU2 (a), strain CA2 (b) or strain CG10 (c), or LPS from *Escherichia coli* O111:B4 (d) or *Escherichia coli* Re mutant (e). Concentrations of IL-8 in the resulting culture supernatants were determined by ELISA. Experiments were done in triplicate. The results were expressed as fold induction (mean values \pm SD), relative to the value obtained from control experiments without LPS stimulation. * $P < 0.01$ compared to LPS-untreated control. The data represent results for the highly purified LPS preparations. Similar results were obtained when the conventional LPS preparations were used.



transducing receptor(s) of weak endotoxic LPS is controversial. For example, Erridge *et al.* (2004) have reported that LPS with the low endotoxic activity shown above commonly transduces signals via TLR2. In contrast, Hashimoto *et al.* (2004) indicated that contaminating lipopeptide in a *P. gingivalis* LPS preparation is an agonist of TLR2, and LPS is a weak agonist of a TLR4-containing complex.

In the case of *H. pylori*, bacterial components other than LPS can induce NF- κ B-mediated inflammatory reactions. As a TLR-independent pathway, the type IV secretion system, consisting of proteins encoded by genes located in

Fig. 4. Reporter gene assay of NF- κ B activation by *Helicobacter pylori*, *Campylobacter jejuni* and *Escherichia coli* LPS using 293 cells expressing TLR4, CD14 and MD2. Expression plasmids of TLR4, CD14 and MD2 and reporter plasmids of pELAM-L and pHR-TK were cotransfected into 293 cells. After 24 h, cells were stimulated with the conventional LPS preparation derived from *Helicobacter pylori* (strains GU2, CA2, and CG10), *Campylobacter jejuni* (strains OH4832 and 81-176) or *Escherichia coli* O111:B4 for 6 h. The reporter activity was measured. The experiments were done in triplicate. The results were expressed as fold induction (mean values \pm SD), relative to the value obtained from control experiments without LPS stimulation.






Fig. 5. Reporter gene assay of NF- κ B activation by *Helicobacter pylori* LPS using 293 cells expressing various TLR systems. (a) Expression plasmids indicated at the top of each graph plus expression plasmid of CD14, and reporter plasmids (pELAM-L and pRL-TK), were cotransfected into 293 cells. After 24 h, cells were stimulated with the conventional LPS preparation derived from *Helicobacter pylori* strain GU2 (\bullet) or strain CA2 (\blacktriangle) for 6 h. Positive control experiments were carried out with Pam₂CSK₄ (for TLR2 and TLR2-TLR1), MALP-2 (for TLR2-TLR6), LPS derived from *Escherichia coli* O111:B4 (for TLR4-MD2) or flagella derived from *Salmonella* (for TLR5) as a stimulant. The reporter activity was measured. The experiments were done in triplicate. The results were expressed as fold induction (mean values \pm SD), relative to the value obtained from control experiments without LPS stimulation. * $P < 0.01$ compared to LPS-untreated control. (b) Protein expression levels of TLR2, TLR1, TLR6 and CD14 in transfectants with plural expression plasmids. 293 cells were transfected with the expression plasmids indicated at the top of the figure. After 24 h of transfection, cells were lysed and assessed by Western blotting.

the *cag* pathogenicity island, including CagE, contributes to NF- κ B activation (Glocker *et al.*, 1998; Naumann *et al.*, 1999). Viable bacterial cells are required for type IV secretion system-dependent NF- κ B activation. Although flagella derived from various bacteria activate NF- κ B via the TLR5-dependent pathway, *H. pylori* flagella cannot activate such systems (Andersen-Nissen *et al.*, 2005). Other protein components, such as heat shock protein 60 (Takenaka *et al.*, 2004) and urease B (Beswick *et al.*, 2006), act as NF- κ B activators and are suggested to be TLR agonists.

Helicobacter pylori LPS is also believed to be a TLR agonist. It is still controversial which receptor is recognized by *H. pylori* LPS. Lepper *et al.* (2005) and Smith *et al.* (2003) reported that TLR2 is essential for signal transduction activated by *H. pylori* LPS. On the other hand, Kawahara *et al.* (2001), Ishihara *et al.* (2004) and Hashimoto *et al.* (2004) reported that TLR4 plays an essential role in LPS-induced inflammatory responses. Maeda *et al.* (2001) indicated that mechanisms of NF- κ B activation differ among cell types. Monocyte-macrophage (THP-1) cells respond to *H. pylori* in a TLR4-CD14 system-dependent manner. In contrast, epithelial (MKN45) cells show TLR-independent and *H. pylori* *cag*-dependent responses to *H. pylori*. Maeda *et al.* concluded that monocyte-macrophage cells are acti-

vated by *H. pylori* via the TLR4 system, and this might be dependent on LPS. In contrast, epithelial cells are mainly activated by viable *H. pylori* cells, and this is dependent on the type IV secretion system but not on LPS.

There might be several reasons for such contradictions. The first might be contamination of LPS preparations by lipoproteins, proteins, nucleic acids and other bacterial LPSs. Lipoproteins and other bacterial LPSs may stimulate TLR2 and TLR4 signals, respectively. Furthermore, bacterial nucleic acids may stimulate other TLRs, such as TLR9 (Akira *et al.*, 2001; Barton & Medzhitov, 2002). The second reason might be differences in the experimental systems used, such as different cell lines and strategies used to identify each TLR. For example, the strategies are transfection of dominant negative mutants, suppression by neutralizing antibodies, and transfection of TLR-expression plasmids into TLR-deficient cell lines, such as 293 cells. The third, and possibly most important, reason is heterogeneity of the structures of *H. pylori* LPS molecules within a given preparation. This could result from strain differences and/or bacterial culture condition. The chemical structure of lipid A, e.g. the number and chain length of fatty acid residues and the presence of phosphate groups, derived from these bacteria differs from that of a typical enterobacterial LPS (Netea *et al.*,

2002). Tran *et al.* (2005) reported that the lipid A portion of *H. pylori* LPS undergoes several structural modifications through the action of specific modifying enzymes. The 'remodeling' of LPS should generate structural heterogeneity in lipid A among strains and culture conditions.

Here, we used highly purified preparations of *H. pylori* LPS, from which contaminants were removed by extensive purification. Such contaminants were removed by proteinase K, DNase, RNase and lipoprotein lipases, and the LPS was further subjected to hydrophobic interaction chromatography. After enzyme treatment and hydrophobic interaction chromatography, our LPS preparation did not show altered reactivity, indicating that the activity is due to LPS itself and not to contaminants. Our results clearly suggest that inflammatory signal transduction mediated by *H. pylori* LPS in our highly purified preparation requires TLR2 but not TLR4. Complexes containing TLR2 and TLR1 were more efficient signal-transducing receptors for *H. pylori* LPS than complexes containing TLR2 and TLR6. The similar activities were observed for six strains, which had various characters, such as smooth/rough phenotypes and antigenicity of the polysaccharide portion. In the present study, we cannot completely rule out the possibility that very small amounts of TLR agonists were present in the highly purified LPS preparations. Furthermore, we have not examined different LPS preparations from cells grown under various culture conditions, and determined the chemical structures of lipid A portions. However, the TLR2-dependent activity that we observed seemed to be common to LPS preparations from various strains of *H. pylori*. The structural requirements for the TLR2-dependent activity need to be clarified.

The magnitude of response to a typical LPS, such as that from *E. coli*, was correlated with expression levels of typical LPS receptor complexes, which contain TLR4, CD14 and MD2. Higher expression levels of TLR4 and CD14 were seen in LPS-high-responder cells, such as T24. On the other hand, TLR2 expression levels were similar among all cell lines used in this study. In conclusion, *H. pylori* LPS is a weak but significant inducer of inflammatory reactions. The magnitude of the inflammatory response stimulated by *H. pylori* LPS is comparable to that of typical LPS in LPS-low-responder cells (e.g. MKN28), which express TLR4 at very low levels.

Acknowledgements

We thank Dr Michael F. Smith and Dr Ken-ichiro Shibata for providing expression plasmids.

References

Akira S, Takeda K & Kaisho T (2001) Toll-like receptors: critical proteins linking innate and acquired immunity. *Nat Immunol* 2: 675–680.

- Andersen-Nissen E, Smith KD, Strobe KL, Barrett SL, Cookson BT, Logan SM & Aderem A (2005) Evasion of Toll-like receptor 5 by flagellated bacteria. *Proc Natl Acad Sci USA* 102: 9247–9252.
- Barton GM & Medzhitov R (2002) Toll-like receptors and their ligands. *Curr Top Microbiol Immunol* 270: 81–92.
- Beswick EJ, Pinchuk IV, Minch K, Suarez G, Sierra JC, Yamaoka Y & Reyes VE (2006) The *Helicobacter pylori* urease B subunit binds to CD74 on gastric epithelial cells and induces NF- κ B activation and interleukin-8 production. *Infect Immun* 74: 1148–1155.
- Covacci A, Telford JL, Del Giudice G, Parsonnet J & Rappuoli R (1999) *Helicobacter pylori* virulence and genetic geography. *Science* 284: 1328–1333.
- Cover TL & Blaser MJ (1996) *Helicobacter pylori* infection, a paradigm for chronic mucosal inflammation: pathogenesis and implications for eradication and prevention. *Adv Intern Med* 41: 85–117.
- Erridge C, Pridmore A, Eley A, Stewart J & Poxton IR (2004) Lipopolysaccharides of *Bacteroides fragilis*, *Chlamydia trachomatis* and *Pseudomonas aeruginosa* signal via toll-like receptor 2. *J Med Microbiol* 53: 735–740.
- Fujita M, Into T, Yasuda M, Okusawa T, Hamahira S, Kuroki Y, Eto A, Nisizawa T, Morita M & Shibata K (2003) Involvement of leucine residues at positions 107, 112, and 115 in a leucine-rich repeat motif of human Toll-like receptor 2 in the recognition of diacylated lipoproteins and lipopeptides and *Staphylococcus aureus* peptidoglycans. *J Immunol* 171: 3675–3683.
- Glocker E, Lange C, Covacci A, Bereswill S, Kist M & Pahl HL (1998) Proteins encoded by the *cag* pathogenicity island of *Helicobacter pylori* are required for NF- κ B activation. *Infect Immun* 66: 2346–2348.
- Hashimoto M, Asai Y & Ogawa T (2004) Separation and structural analysis of lipoprotein in a lipopolysaccharide preparation from *Porphyromonas gingivalis*. *Int Immunol* 16: 1431–1437.
- Heine H, Muller-Loennies S, Brade L, Lindner B & Brade H (2003) Endotoxic activity and chemical structure of lipopolysaccharides from *Chlamydia trachomatis* serotypes E and L2 and *Chlamydia psittaci* 6BC. *Eur J Biochem* 270: 440–450.
- Hitchcock P & Brown TM (1983) Morphological heterogeneity among *Salmonella* lipopolysaccharide chemotypes in silver-stained polyacrylamide gels. *J Bacteriol* 154: 269–277.
- Hynes SO, Ferris JA, Szponar B *et al.* (2004) Comparative chemical and biological characterization of the lipopolysaccharides of gastric and enterohepatic helicobacters. *Helicobacter* 9: 313–323.
- Ishihara S, Rumi MA, Kadowaki Y *et al.* (2004) Essential role of MD-2 in TLR4-dependent signaling during *Helicobacter pylori*-associated gastritis. *J Immunol* 173: 1406–1416.
- Kawahara T, Teshima S, Oka A, Sugiyama T, Kishi K & Rokutan K (2001) Type I *Helicobacter pylori* lipopolysaccharide stimulates

- toll-like receptor 4 and activates mitogen oxidase 1 in gastric pit cells. *Infect Immun* 69: 4382–4389.
- Kusters JG, van Vliet AH & Kuipers EJ (2006) Pathogenesis of *Helicobacter pylori* infection. *Clin Microbiol Rev* 19: 449–490.
- Lepper PM, Triantafyllou M, Schumann C, Schneider EM & Triantafyllou K (2005) Lipopolysaccharides from *Helicobacter pylori* can act as antagonists for Toll-like receptor 4. *Cell Microbiol* 7: 519–528.
- Lindberg AA, Weintraub A, Zahringer U & Rietschel ET (1990) Structure–activity relationships in lipopolysaccharides of *Bacteroides fragilis*. *Rev Infect Dis* 12 (suppl 2), S133–S141.
- Maeda S, Akanuma M, Mitsuno Y, Hirata Y, Ogura K, Yoshida H, Shiratori Y & Omata M (2001) Distinct mechanism of *Helicobacter pylori*-mediated NF- κ B activation between gastric cancer cells and monocytic cells. *J Biol Chem* 276: 44856–44864.
- Naumann M, Wessler S, Bartsch C, Wieland B, Covacci A, Haas R & Meyer TF (1999) Activation of activator protein 1 and stress response kinases in epithelial cells colonized by *Helicobacter pylori* encoding the *cag* pathogenicity island. *J Biol Chem* 274: 31655–31662.
- Netea MG, van Deuren M, Kullberg BJ, Cavaillon JM & Van der Meer JW (2002) Does the shape of lipid A determine the interaction of LPS with Toll-like receptors? *Trends Immunol* 23: 135–139.
- Nielsen H, Birkholz S, Andersen LP & Moran AP (1994) Neutrophil activation by *Helicobacter pylori* lipopolysaccharides. *J Infect Dis* 170: 135–139.
- Ogawa T, Suda Y, Kashiwara W, Hayashi T, Shimoyama T, Kusumoto S & Tamura T (1997) Immunobiological activities of chemically defined lipid A from *Helicobacter pylori* LPS in comparison with *Porphyromonas gingivalis* lipid A and *Escherichia coli*-type synthetic lipid A (compound 506). *Vaccine* 15: 1598–1605.
- Parsonnet J (1998) *Helicobacter pylori*: the size of the problem. *Gut* 43 Suppl 1, S6–S9.
- Perez-Perez GI, Shepherd VL, Morrow JD & Blaser MJ (1995) Activation of human THP-1 cells and rat bone marrow-derived macrophages by *Helicobacter pylori* lipopolysaccharide. *Infect Immun* 63: 1183–1187.
- Rose JR, Christ WJ, Bristol JR, Kawata T & Rossignol DP (1995) Agonistic and antagonistic activities of bacterially derived *Rhodobacter sphaeroides* lipid A: comparison with activities of synthetic material of the proposed structure and analogs. *Infect Immun* 63: 833–839.
- Schroder NW, Heine H, Alexander C, Manukya M, Ecker J, Hamann L, Gobel U & Schumann RR (2004) Lipopolysaccharide binding protein binds to triacylated and diacylated lipopeptides and mediates innate immunity. *J Immunol* 173: 2683–2691.
- Smith MF Jr, Mitchell A, Li G, Ding S, Fitzmaurice AM, Ryan K, Crowe S & Goldberg JB (2003) Toll-like receptor (TLR) 2 and TLR5, but not TLR4, are required for *Helicobacter pylori*-induced NF- κ B activation and chemokine expression by epithelial cells. *J Biol Chem* 278: 32552–32560.
- Takenaka R, Yokota K, Ayada K et al. (2004) *Helicobacter pylori* heat-shock protein 60 induces inflammatory responses through the Toll-like receptor-triggered pathway in cultured human gastric epithelial cells. *Microbiology* 150: 3913–3922.
- Tran AX, Stead CM & Trent MS (2005) Remodeling of *Helicobacter pylori* lipopolysaccharide. *J Endotoxin Res* 11: 161–166.
- Weckesser J & Mayer H (1988) Different lipid A types in lipopolysaccharides of phototrophic and related non-phototrophic bacteria. *FEMS Microbiol Rev* 4: 143–153.
- Yokota S, Amano K, Hayashi S & Fujii N (1997) Low antigenicity of the polysaccharide region of *Helicobacter pylori* lipopolysaccharides derived from tumors of patients with gastric cancer. *Infect Immun* 65: 3509–3512.
- Yokota S, Amano K, Shibata Y, Nakajima M, Suzuki M, Hayashi S, Fujii N & Yokochi T (2000) Two distinct antigenic types of the polysaccharide chains of *Helicobacter pylori* lipopolysaccharides characterized by reactivity with sera from humans with natural infection. *Infect Immun* 68: 151–159.
- Yokota S, Amano K, Chiba S & Fujii N (2003) Structures, biological activities and antigenic properties of *Helicobacter pylori* lipopolysaccharide. *Recent Res Dev Microbiol* 7: 251–267.

Quantitative nuclear magnetic resonance spectroscopic determination of the oxyethylene group content of polysorbates

NAOKI SUGIMOTO¹, RYO KOIKE², NORIKO FURUSHO¹, MAKOTO TANNO²,
CHIKAKO YOMOTA¹, KYOKO SATO¹, TAKESHI YAMAZAKI¹, &
KENICHI TANAMOTO¹

¹National Institute of Health Sciences: 1-18-1, Kamiyoga, Setagaya, Tokyo 158-8501, Japan and ²Kao Corporation, Analytical Research Center, 1334, Minato, Wakayama-shi, Wakayama 640-8580, Japan

(Received 30 November 2006; revised 5 February 2007; accepted 13 February 2007)

Abstract

Guidelines for the oxyethylene group (EO) content of polysorbates are set by the Food and Agriculture Organization/World Health Organization Joint Expert Committee on Food Additives. However, the classical titration method for EO determination is difficult and time-consuming. Here, we show that quantitative ¹H-nuclear magnetic resonance spectroscopy can determine the EO contents of polysorbates rapidly and simply. The EO signals were identified through comparisons with sorbitan monolaurate and poly(ethylene glycol) distearate. Potassium hydrogen phthalate was used as an internal standard. The EO contents were estimated from the ratio of the signal intensities of EO to the internal standard. Two nuclear magnetic resonance systems were used to validate the proposed method. The EO content of commercial polysorbates 20, 60, 65, and 80 was determined to be within the recommended limits using this technique. Our approach thus represents an additional or alternative method of determining the EO contents of polysorbates.

Keywords: Analytical method, food additive, oxyethylene, polysorbate, quantitative nuclear magnetic resonance


Introduction

Polysorbates are non-ionic surfactants that are widely used as emulsifiers, dispersants, and stabilizers in food processing. Polysorbates consist of a mixture of fatty-acid partial esters of sorbitol and condensed sorbitol anhydrides, and contain approximately 20 moles of ethylene oxide (comprising the oxyethylene unit [EO] –OC₂H₄–) for each mole of sorbitol, along with its monohydrides and dianhydrides. The main fatty acids of polysorbates 20, 60, 65, and 80 are monolauric acid, monostearic acid, tristearic acid, and monooleic acid, respectively. The typical structures of these polysorbates are shown in Figure 1.

Guidelines for the EO contents of polysorbates are set by the Food and Agriculture Organization (FAO)/World Health Organization (WHO) Joint Expert Committee on Food Additives (JECFA). To comply with the JECFA standards, the quality

and composition of commercially synthesized polysorbates must be monitored and regulated. The standard method of measuring EO as described in “section VI. Methods for fats and related substances in the guide to specification” is as follows: “The oxyethylene groups are converted to ethylene and ethyl iodide which can be determined by titration. By utilizing a conversion factor determined on a reference sample, it is possible to compute the polyoxyethylene ester content” (JECFA [internet]). However, this classical titration method requires a complicated apparatus and involves several time-consuming steps. Alternative methods for determining the EO contents of polysorbates have not previously been reported, because these complex compounds are mixtures of isomers that are non-selectively substituted with EOs and fatty acids.

The quantitative nuclear magnetic resonance (qNMR) approach is based upon the International



Compound		Formula (MW)	EO(%) in molecule
Polysorbate 20 (polyoxyethylene (20) sorbitan monolaurate)	$R_1 = \text{H}_3\text{C}-(\text{CH}_2)_5-\text{C}(=\text{O})-\text{CH}_2-$ $R_2 = R_3 = \text{H}$	$C_{58}H_{114}O_{26}$ (MW1227.5)	EO(%) = 71.8
Polysorbate 60 (polyoxyethylene (20) sorbitan monostearate)	$R_1 = \text{H}_3\text{C}-(\text{CH}_2)_7-\text{C}(=\text{O})-\text{CH}_2-$ $R_2 = R_3 = \text{H}$	$C_{64}H_{126}O_{26}$ (MW1311.7)	EO(%) = 67.2
Polysorbate 65 (polyoxyethylene (20) sorbitan tristearate)	$R_1 = R_2 = R_3 = \text{H}_3\text{C}-(\text{CH}_2)_7-\text{C}(=\text{O})-\text{CH}_2-$	$C_{100}H_{194}O_{28}$ (MW1844.6)	EO(%) = 47.8
Polysorbate 80 (polyoxyethylene (20) sorbitan monooleate)	$R_1 = \text{H}_3\text{C}-(\text{CH}_2)_6-\text{CH}=\text{CH}-(\text{CH}_2)_6-\text{C}(=\text{O})-\text{CH}_2-$ $R_2 = R_3 = \text{H}$	$C_{63}H_{122}O_{26}$ (MW1295.6)	EO(%) = 68.0

Figure 1. Typical structures of polysorbates 20, 60, 65, and 80. The formulae and EO (%) were estimated based on the assumption that there were 20 moles of EO per molecule.

system of units (SI units). This valuable technique meets the requirements of a primary ratio analytical method (Jancke 1998). The use of qNMR to determine the ethanol content of deuterium oxide solution was previously reported as a part of an intercomparison study organized by the Comité Consultatif pour la Quantité de Matière (CCQM). The results showed that the accuracy of qNMR was equivalent to that of gas chromatography with a flame ionization detector (GC-FID) (Saito et al. 2003). qNMR exploits the fact that the signal intensities of a given NMR resonance are directly proportional to the molar amount of the nucleus within the sample. qNMR can determine the quantity of a compound, its substituent contents, or its absolute quality if the whole sample weight is known. This technique has several advantages for the analysis of organic compounds: it is non-destructive, it provides both quantitative data and structural information about a compound, and high-throughput spectral-acquisition instruments are commercially available. The main drawback of the qNMR approach is that manual spectral assignment is required; however, this can easily be rectified by applying current NMR technical experiments such as total correlated spectroscopy (TOCSY), heteronuclear multiple quantum correlation (HMQC), heteronuclear multiple bond coherence (HMBC), etc.

Based on these features of qNMR, we predicted that the method could be used to determine the EO

contents of polysorbates. In the current paper, we detail the application of qNMR along with an internal standard for the direct determination of the EO contents of polysorbates.

Materials and methods

Materials

Samples of reagent-grade polysorbates 20, 60, 65, 80, and sorbitan monolaurate (Span 20) were purchased from Wako Pure Chemical Industries, Ltd (Osaka, Japan). Poly(ethylene glycol) distearate was purchased from Sigma-Aldrich Japan KK (Tokyo). Commercial samples of polysorbates were obtained from companies A-E via the Japan Food Additives Association. The NMR solvents, methanol- d_4 and acetone- d_6 with 0.03% tetramethylsilane (TMS), were purchased from Isotec Inc. (Miamisburg, OH). Potassium hydrogen phthalate (PHP), which was standard grade for volumetric analysis according to Japanese Industrial Standard (JIS) K8005, was purchased from Wako Pure Chemical Industries, Ltd.

Instrumentation

NMR spectra were recorded on JNM-ECA (500 MHz; JEOL, Tokyo) and MERCURY (400 MHz; VARIAN, Palo Alto, CA) pulsed Fourier-transform (FT) spectrometers, equipped with 5 mm $^1\text{H}\{\text{X}\}$ inverse detection gradient

Table I. Instruments and acquisition parameters.

	MERCURY400 (VARIAN) and ECA500 (JEOL)
Spectrometer	
Probe	5 mm indirect detection probe
Spectral width	2.5–12.5 ppm
Data points	64 000
Flip angle	45°
Pulse delay	30 s (>5 * T ₁)
Scan times	8
Sample spin	15 Hz
Probe temperature	25°C
Solvent	Mixture of methanol-d ₄ and acetone-d ₆ (1:1)
Internal standard	Potassium hydrogen phthalate (PHP)
Range of integral signal	Oxyethylene group (EO) = 3.40–3.85 ppm 4 protons of PHP = 7.46–7.66 ppm + 8.18–8.38 ppm

probes, with methanol-d₄:acetone-d₆ (1:1) and 0.3% (w/v) PHP as an NMR solvent. The spectra were referenced internally to TMS by ¹H-NMR. The samples and internal standard were weighed on a LIBROR AEG-80SM (Shimadzu, Kyoto, Japan) electronic balance to an accuracy of ±0.01 mg.

Preparation of samples and NMR measurement conditions

The polysorbate samples were prepared as follows. PHP was crushed into a powder in a mortar and dried for 1 h at 120°C. After cooling in a desiccator, the powder (300 mg) was dissolved in 100 ml of methanol-d₄:acetone-d₆ (1:1) with ultrasonic agitation for 30 min. This stock solution was used as the NMR solvent and included an internal standard. A 50-mg polysorbate sample was then dissolved in 3 ml of the NMR solvent described above, and 0.6 ml of the sample solution was placed into a 5-mm NMR tube (Kusano Science Co. Ltd, Tokyo). The ¹H-NMR spectra were recorded on MERCURY400 and ECA500 spectrometers operating at 400 and 500 MHz, respectively. Typical ¹H-NMR parameters for the quantitative analyses are listed in Table I. The free induction decay (FID) signals of the samples from the MERCURY400 and ECA500 spectrometers were loaded onto a Windows XP-based personal computer (PC) equipped with the Alice 2 Version 5 (JEOL) NMR data-processing and analytical software. Fourier transformations of the FID signals were carried out with this software using the default parameters; window function = exponential, BF = 0.12 Hz, zero filling = 1, T₁ = T₂ = 0%, T₃ = 90%, T₄ = 100%. After phase adjustments and baseline corrections of the NMR spectra were performed using the same algorithms in the automatic mode of Alice 2, the signal intensities

of the EOs and internal standard protons were measured, respectively.

Results and discussion

Identification of EO signals in polysorbates

Polysorbate molecules contain approximately 20 moles of EO according to the JECFA definition. However, recently reported matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) spectra showed that polysorbates include numerous other chemical species, including polyethylenes, unesterified, monoesterified, and diesterified polyoxyethylene sorbitans, and isosorbides (Frison-Norrie and Sporns 2001). Furthermore, analysis by liquid chromatography (LC)-mass spectrometry (MS) confirmed that polysorbates contain not only polyoxyethylene sorbitan fatty acid esters but also numerous intermediates, such as polyoxyethylene sorbitan and isosorbitan, and the monoesters and diesters of fatty acids (Vu Dang et al. 2006). These studies have confirmed that polysorbates comprise many types of chemical isomers. This molecular diversity makes it difficult to determine the EO contents of polysorbates. However, we hypothesized that the EO contents of polysorbates could be measured rapidly and simply by qNMR if the signals could be identified on ¹H-NMR spectra, regardless of whether they contained numerous chemical isomers.

Thus, in order to identify the EO signals in polysorbates, we compared the ¹H-NMR spectra of polysorbate 20, sorbitan monolaurate, and poly(ethylene glycol) distearate. The partial structures of sorbitan monolaurate and poly(ethylene glycol) distearate, which comprised a sorbitol anhydride core and poly(ethylene glycol), were similar to those of polysorbate 20 (Figures 2 and 3). The sorbitan monolaurate and poly(ethylene glycol) distearate spectra revealed fatty-acid moiety signals with δ_{H} values ranging from 0.9 to 2.4 ppm, similar to those of polysorbate 20. The triplet signal at δ_{H} c. 0.9 ppm, the major broad signal and multiplet signal at δ_{H} c. 1.3 ppm and 1.6 ppm, and the triplet signal at δ_{H} c. 2.4 ppm were identified as the terminal CH₃-, -CH₂-, and -CH₂C=O- groups of the fatty acids, respectively. Most of the EO signals in poly(ethylene glycol) distearate were observed between δ_{H} values of 3.40 and 3.85 ppm. One of the -CH₂O- groups appeared to have been shifted downfield to δ_{H} c. 4.2 ppm, near to the residual proton of methanol-d₄ at δ_{H} c. 4.4 ppm. A HMBIC experiment revealed that the proton at δ_{H} c. 4.2 ppm was correlated to the carbonyl carbon of the fatty acid at δ_{C} 173.4 ppm. Thus, the proton signal was assigned




Figure 2. Structures of sorbitan monolaurate and poly(ethylene glycol) distearate.




Figure 3. Comparison of NMR spectra of polysorbate 20, sorbitan monolaurate and poly(ethylene glycol) distearate. ¹H-NMR spectra were obtained using the ECA500 system (500 MHz; JEOL) under the conditions shown in Table I.

to the $-\text{CH}_2\text{O}-$ group adjacent to the fatty acid side chain. In the sorbitan monolaurate spectrum, various minor proton signals were observed from δ_{H} values of *c.* 3.4–5.0 ppm; these were attributed to the sorbitan moiety in sorbitan monolaurate, which consists of a mixture of cyclic sorbitol-derived ethers (such as sorbitan, isosorbite, and other isomers). These signals were also observed on the spectrum of polysorbate 20. However, the signals were broad and negligibly smaller than that of sorbitan monolaurate, as polysorbate 20 has the diversity of molecule more than sorbitan monolaurate. The polysorbate 20 signals ranging from δ_{H} 0.9 to 2.4 ppm that were attributed to the fatty-acid moiety were similar to those of sorbitan monolaurate and poly(ethylene glycol) distearate. Polysorbate 60, 65, and 80 also showed the signals of fatty acid as same as sorbitan monolaurate, but

the olefinic protons were only observed at δ_{H} 5.3 ppm on the spectrum of polysorbate 80 consisting of an unsaturated fatty acid (data not shown). The EO signals were assigned to a large envelop between δ_{H} 3.40 and 3.85 ppm, and at δ_{H} 4.20 ppm, which overlapped with the negligible small broad signals seen for the mixture of sorbitan, isosorbite, and other isomers moieties between δ_{H} values of *c.* 3.4 and 5.0 ppm. The EO signals of polysorbates 60, 65, and 80 also appeared within these ranges (data not shown). This was due to the fact that polysorbates basically comprise the same units: sorbitol anhydrides core, EO chains, and fatty acids. Although proton signals of the $-\text{CH}_2\text{O}-$ group adjacent to the fatty acid at δ_{H} *c.* 4.20 ppm were observed, and the signals of the sorbitol anhydrides core were overlapped on EO signals, they were negligible and did not effect the




Figure 4. $^1\text{H-NMR}$ spectrum of polysorbate 20. The spectrum was obtained using the ECA500 system (500 MHz; JEOL). PHP was added as an internal standard. Signals of the four protons on the benzene ring of PHP were observed at δ_{H} values of 7.46–7.66 ppm and 8.18–8.38 ppm. Most of the EO signals of polysorbate 20 were observed in a large envelope between δ_{H} 3.40 and 3.85 ppm.

determination of the EO contents. Thus, in the current research, we used the EO signals between δ_{H} 3.40 and 3.85 ppm to determine the EO contents of polysorbates by NMR.

Determination of EO contents in polysorbates 20, 60, 65, and 80

Several reports have described the applications of qNMR to determine specific types of chemical compound, such as natural products, impurities, and polymers (Stefanova et al. 1988; Paula 2001; Jake et al. 2002; Wells et al. 2002; Paula et al. 2005). Recently, a practical set of parameters for qNMR has been discussed (Saito et al. 2004). Furthermore, qNMR using an internal standard has been suggested as a new way of determining the contents of surfactants with a relatively high throughput (Koike et al. 2004a, 2004b, 2005). To minimize quantitative errors, we used the qNMR conditions described by Koike and colleagues, as listed in Table I. In particular, the flip angle was set to 45° , and the spectral width was set at a value sufficient for the peak of interest to fall within 80% of its centre, because the signal intensities decreased towards both edges of the spectral window. The number of data points was set at 64 000 to enhance the resolution. The pulse delay was set at up to 30 s, as high-precision NMR can only be achieved when the pulse delay time is greater than the quintuple spin-lattice relaxation time ($>5 \times T_1$). As qNMR is based on the fact that the signal intensities of a given resonance are directly proportional to the molar quantity of the nucleus within the sample, the EO signal intensity of polysorbates and four protons on the benzene ring of PHP were used to determine the EO contents. The total time taken to obtain one FID using these parameters was <10 min.

The weight percentage of the EO groups was calculated according to Equation 1.

$$\text{EO(w/w\%)} = \frac{(I_{\text{EO}}/H_{\text{EO}} \times M_{\text{EO}}/W_{\text{sample}})}{(I_{\text{standard}}/H_{\text{standard}} \times M_{\text{standard}}/W_{\text{standard}})} \times 100. \quad (1)$$

Here, I_{EO} is the signal intensity of the EO group; H_{EO} is the number of protons of the EO group (four); M_{EO} is the partial molecular weight of the EO group (44); W_{sample} is the weight (mg) of the sample in 3 ml of NMR solvent including PHP as an internal standard; I_{standard} is the total signal intensity of PHP; H_{standard} is the number of protons on the benzene ring of PHP (four); M_{standard} is the molecular weight of PHP (204); and W_{standard} is the weight (mg) of PHP in 3 ml of NMR solvent.

We initially confirmed that the qNMR showed linearity between the intensity of the EO signal and the amount of polysorbate 20. Various amounts of the reagent-grade polysorbate 20 sample were analysed by $^1\text{H-NMR}$ under the conditions described in the Materials and methods and Table I. The NMR spectrum of polysorbate 20 with the internal standard is shown in Figure 4. The four protons of the PHP benzene ring were observed as two double-doublet signals at δ_{H} values of 7.46–7.66 ppm and 8.18–8.38 ppm, respectively. The ratio of the EO signal intensity was calculated as follows: intensity of EO/total intensities of four protons on PHP benzene ring. The relationship between EO/PHP and the amount of polysorbate 20 was linear ($R^2 = 0.9996$) in the range of 12.5–100 mg of polysorbate 20 in 3 ml of NMR solvent. Based on these results, we concluded

Table II. Determination of EO contents in polysorbates by qNMR.^a

Sample name	MERCURY (400 MHz, VARIAN)			ECA500 (500 MHz, JEOL)		
	Entry	EO (%)	SD	Entry	EO (%)	SD
Polysorbate 20 (polyoxyethylene (20) sorbitan monolaurate)	1	73.0		1	72.2	
	2	71.8		2	71.8	
	3	73.2		3	72.3	
	4	71.7		4	72.5	
	5	71.9		5	71.6	
				6	72.9	
				7	72.0	
				8	72.7	
				9	73.7	
		AV	72.3	0.7	AV	72.4
Polysorbate 60 (polyoxyethylene (20) sorbitan monostearate)	1	67.7		1	67.4	
	2	65.3		2	67.7	
	3	68.9		3	67.5	
	4	67.8		4	67.9	
	5	66.9		5	68.6	
	AV	67.3	1.3	AV	67.8	0.5
Polysorbate 65 (polyoxyethylene(20) sorbitan tristearate)	1	49.1		1	49.8	
	2	49.8				
	3	49.5				
	4	49.8				
	5	48.7				
	AV	49.4	0.5			
Polysorbate 80 (polyoxyethylene (20) sorbitan monooleate)	1	65.0		1	67.0	
	2	65.5				
	3	66.2				
	4	64.8				
	5	65.1				
	AV	65.3	0.6			

^aReagent-grade polysorbates were purchased from Wako Pure Chemical Industries, Ltd. "Entry" means that the same sample was measured repeatedly on different days.

that qNMR could quantitatively determine the EO contents of polysorbates.

In order to verify whether qNMR could accurately determine the EO contents of polysorbates, two different NMR instruments (MERCURY and ECA500, with magnetic field strengths of 400 and 500 MHz, respectively) were used to repeatedly measure the EO contents of reagent grade polysorbates 20, 60, 65, and 80, which are generally used as standards. The results are shown in Table II. Reproducible results were obtained from each sample using the MERCURY system. Furthermore, the results obtained by the two NMR instruments did not differ significantly (standard deviations = 0.5–1.3%). These findings confirmed that it was possible to determine the EO contents of polysorbates using this approach regardless of the NMR instrument employed.

Finally, to confirm the validity of qNMR, we determined the EO contents of the commercially synthesized polysorbates 20, 60, 65, and 80, which

met the specifications of the JECFA. All of the EO contents of the polysorbates were within the limits described in the *Compendium of Food Additive and Flavoring Agent Specifications* (JECFA [internet]) (Table III). The qNMR method for determining the EO contents of polysorbates demonstrated in this paper thus represents a simple and rapid alternative to the classic titration method recommended by the JECFA, which does not require specific chemical reactions or sophisticated apparatus. Moreover, the qNMR method made it possible to distinguish between Polysorbates 60 and 80, which have the same stipulated value, by comparison with the ¹H-NMR spectra, as polysorbate 80 consisting of an unsaturated fatty acid only showed the signals of olefinic protons at δ_{H} 5.3 ppm. It is theoretically possible to determine the ratio of a substituted group in any molecule, or the quality of any compound, using the proposed qNMR method with an internal standard, provided that the target proton signals can be separated from those

Table III. EO contents in commercial polysorbates determined using qNMR.*

Name	Stipulated value	Brand	EO (%)	SD
Polysorbate 20 (polyoxyethylene (20) sorbitan monolaurate)	70.0-74.0%	A	71.2	1.0
		B	73.0	
		C	70.3	
		D	71.0	
		E	71.5	
		AV	71.4	
Polysorbate 60 (polyoxyethylene (20) sorbitan monostearate)	65.0-69.5%	A	66.9	1.1
		B	65.4	
		C	68.0	
		D	68.1	
		E	67.2	
		AV	67.1	
Polysorbate 65 (polyoxyethylene (20) sorbitan tristearate)	46.0-50.0%	A	48.3	1.1
		B	46.0	
		C	-	
		D	47.2	
		E	48.1	
		AV	47.4	
Polysorbate 80 (polyoxyethylene (20) sorbitan monooleate)	65.0-69.5%	A	67.4	1.6
		B	65.1	
		C	69.3	
		D	66.7	
		E	68.0	
		AV	67.1	

*Brands A-E were purchased from five manufacturers. Brand C does not supply polysorbate 65.

of non-target groups and impurities. We are currently investigating the potential for this technique to determine various other compounds and polymers.

Conclusions

This research demonstrated that the EO contents of commercial polysorbates 20, 60, 65, and 80 could be readily determined using qNMR with an internal standard. Clear NMR data for the polysorbates were obtained from simple sample preparations. Two different NMR instruments validated the proposed method, and no significant differences were observed among the results. Moreover, the data obtained for commercial polysorbates 20, 60, 65, and 80 were in good agreement with the JECFA guidelines.

It is generally difficult to determine the amounts of substituted groups within polymers owing to their great diversity in molecular weights and structures. Classical methods require time-consuming preparation to set up the apparatus, and technically skilled operators. Furthermore, as there are no alternative methods to validate the results, they have to be accepted without verification. Our proposed qNMR

is a rapid and simple analysis that provides the structural information of target compounds together. These advantages will reduce dramatically the time and manpower cost required, even if the NMR spectrometer and the solvents are expensive. qNMR is thus a valuable additional and/or alternative method, with a broad range of applications in quantitative analysis.

Acknowledgements

This work was supported by a Research on Food Safety in Health and Labour Sciences Research Grants from the Ministry of Health, Labor, and Welfare of Japan.

References

- Frison-Norrie S, Sporns P. 2001. Investigating the molecular heterogeneity of polysorbate emulsifiers by MALDI-TOF MS. *Journal of Agricultural and Food Chemistry* 49:3335-3340.
- Jake B, Stücher O, Veit M, Fröhlich R, Paili GF. 2002. Evaluation of glucoiberin reference material from *Iberis amara* by spectroscopic fingerprinting. *Journal of Natural Products* 65:517-522.
- Jancke H. 1998. NMR spectroscopy as a primary analytical method. Comité Consultatif pour la Quantité de Matière Report 98: 1-12.

- Joint FAO/WHO Expert Committee on Food Additives (JECFA). Available: http://www.fao.org/ag/agn/jecfa/archive_en.stm. Accessed 20 November 2006.
- Koike R, Jo S, Azuma M, Wakisaka T. 2004a. Precise and rapid determination of anionic and cationic surfactants by ^1H nuclear magnetic resonance using an internal standard. *Bunseki Kagaku* (The Japan Society for Analytical Chemistry) 53:1125-1131.
- Koike R, Jo S, Azuma M, Wakisaka T. 2004b. Precise and rapid determination of amphoteric and nonionic surfactants by ^1H nuclear magnetic resonance using an internal standard. *Bunseki Kagaku* (The Japan Society for Analytical Chemistry) 53:1133-1138.
- Koike R, Jo S, Azuma M, Wakisaka T. 2005. Precise and simultaneous determination of surfactants by ^1H nuclear magnetic resonance using an internal standard. *Bunseki Kagaku* (The Japan Society for Analytical Chemistry) 54:715-722.
- Paula GF. 2001. qNMR—a versatile concept for the validation of natural product reference compounds. *Phytochemical Analysis* 12:28-42.
- Paula GF, Jake BU, Lankin DC. 2005. Quantitative ^1H NMR: Development and potential of a method for natural products analysis. *Journal of Natural Products* 68:133-149.
- Saito T, Ihara T, Sato H, Jancke H, Kinugasa S. 2003. International comparison on the determination of an ethanol aqueous solution by ^1H nuclear magnetic resonance. *Bunseki Kagaku* (The Japan Society for Analytical Chemistry) 52:1029-1036.
- Saito T, Nakaie S, Kinoshita M, Ihara T, Kinugasa S, Nomura A, Maeda T. 2004. Practical guide for accurate quantitative solution state NMR analysis. *Metrologia* 41:213-218.
- Stefanova R, Rankoff D, Panayotova S, Spasov SL. 1988. Quantitative proton NMR determination of linoleic acid mono- and diester of polyethyleneglycols via reaction with trichloroacetyl isocyanate. *Journal of American Oil Chemists' Society* 65:1516-1518.
- Vu Dang HV, Gray AJ, Watson D, Bates CD, Scholes P, Gillian GM. 2006. Composition analysis of two batches of polysorbate 60 using MS and NMR techniques. *Journal of Pharmaceutical and Biomedical Analysis* 40:1155-1165.
- Wells RJ, Hook JM, Al-Deen TS, Hibbert DB. 2002. Quantitative nuclear magnetic resonance (QNMR) spectroscopy for assessing the purity of technical grade agrochemicals: 2,4-Dichlorophenoxyacetic acid (2,4-D) and sodium 2,2-dichloropropionate (dalapon sodium). *Journal of Agricultural and Food Chemistry* 50:3366-3374.

Neocrocetin A: a Novel Crocetin Glycoside with a Unique System for Binding Sugars Isolated from Gardenia Yellow

Yoshinori UEKUSA,^{a,1)} Naoki SUGIMOTO,^{*,b} Kyoko SATO,^b Young Sook YUN,^a Akira KUNUGI,^a Takeshi YAMAZAKI,^b and Ken-ichi TANAMOTO^b

^aSchool of Life Science, Tokyo University of Pharmacy and Life Science; 1432-1 Horinouchi, Hachioji, Tokyo 192-0392, Japan; and ^bNational Institute of Health Sciences; 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan.

Received June 20, 2007; accepted August 29, 2007; published online September 3, 2007

A novel crocetin glycosyl ester, neocrocetin A (**2**), was isolated from gardenia yellow. The structure of **2** was elucidated as that of an all-*trans*-crocetin β -D-gentiobiosyl β -D-glucopyranosyl-(1 \rightarrow 6)-D-2-deoxy-glucopyranos-2-yl diester based on chemical and spectral data. The findings provide evidence that the binding system of crocetin glycosides is not limited to the anomeric position.

Key words *Gardenia jasminoides*; gardenia yellow; neocrocetin A; crocetin glycoside; crocin

Crocetin (**1**)²⁾ is a digentiobiosyl all-*trans*-crocetin (8,8'-dipapocrotene-8,8'-dioic acid) ester that is the major yellow pigment in gardenia yellow and saffron, which are extracts of *Gardenia jasminoides* fruits and *Crocus sativus* stigmas, respectively.^{3,4)} Gardenia yellow and saffron consist of many minor pigments as well as some relatively abundant pigments which have been characterized as all-*trans*- and 13-*cis*-crocetin monoglucosyl ester,²⁾ diglucosyl ester,²⁾ monogentiobiosyl ester,^{2,5)} glucosyl gentiobiosyl ester²⁾ and gentiobiosyl neapolitanosyl ester.⁵⁾ However, the structures of the other minor pigments have so far remained uncertain, and the binding systems for sugars have not previously been confirmed by detailed spectroscopic investigations, such as NMR analysis, after isolation. Here we report on the isolation and structural elucidation of a novel crocetin glycoside, neocrocetin A (**2**), which has a unique binding system for sugars, based on spectral data and chemical derivatization (Fig. 1).

Results and Discussion

Gardenia yellow extracted from dried gardenia fruits was fractionated on a Diaion HP-20 column. The 60–70% methanol eluate was then concentrated and the residue was loaded into a preparative LC-MS system,⁶⁾ led to the isolation of crocin (**1**) and neocrocetin A (**2**).

Neocrocetin A (**2**) was isolated as a red amorphous powder. The molecular formula of **2** was established as C₄₄H₆₄O₂₄, which was as the same as that of **1**, according to HR-ESI-MS (*m/z* 999.3707, [M+Na]⁺, Calcd 999.3685), and the IR spectrum and UV/Vis absorption were similar to those of **1**. All-*trans*-crocetin dimethyl ester (**3**) and D-glucose were ob-

tained, respectively, after the methanolysis and hydrolysis⁷⁾ of **2**. These observations implied that **2** had the same carotenoid moiety, all-*trans*-crocetin, as the chromophore group, but that the binding system for glucoses differed from that of **1**. The ¹H- and ¹³C-NMR spectra of **1** showed simple subduplet signals, because **1** had C₂ structural symmetry. The ¹H-NMR spectra of **2** were similar to, but more complex than, those of **1**. Based on this comparison, we predicted that the C₂ structural symmetry was disrupted in **2** by the different binding system for glucoses at each end of crocetin, and that equilibrated isomerization could occur readily in the NMR solvent. The ¹H-NMR spectrum of **2** revealed a crocetin moiety (δ_{H} 6.49–7.31), anomeric doublets of β configuration (δ_{H} 5.38 (d, *J*=7.8 Hz), 4.53 (d, *J*=6.9 Hz), 4.13 (d, *J*=7.8 Hz), 4.15 (d, *J*=7.8 Hz)) and an anomeric doublet of α configuration (δ_{H} 5.06 (d, *J*=3.6 Hz)). Two anomeric protons (δ_{H} 4.53, 5.06) were shifted to a high magnetic field, and two oxymethines on H-2 (δ_{H} 4.39 (dd, *J*=3.7, 10.1 Hz), 4.53 (t, *J*=6.9 Hz)) were shifted to a low magnetic field, in comparison to those of **1**. This observation indicated the existence of a free hydroxyl group at an anomeric position on the glycosyl groups. Furthermore, HMBC correlations were observed between the H-2 of the α -glucoside and β -glucoside (glucose C) at δ_{H} 5.06 and δ_{H} 4.53, and the carbonyl carbons on the crocetin moiety at δ_{C} 167.92 and δ_{C} 167.11, respectively. Based on these chemical and spectral data, **2** was determined to be an all-*trans*-crocetin β -D-gentiobiosyl β -D-glucopyranosyl-(1 \rightarrow 6)-D-2-deoxy-glucopyranos-2-yl ester. The NMR spectral data for **1** and **2** are summarized in Table 1.

Furthermore, to confirm the binding system for the sugars of neocrocetin A (**2**), we firstly carried out peracetylation of **2**. However, the sufficient quantity of peracetylated **2** was not obtained for the structure determination, because **2** was unstable more than crocin (**1**). Hence, after peracetylation of gardenia yellow, we isolated peracetylated crocin (**1a**) and an isomer of neocrocetin A (**2a**). The other isomer of peracetylated **2** could not be isolated using preparative LC-MS because of overlapping with the peak of **1a**. The HR-ESI-MS spectra indicated that the molecular formula of both **1a** and **2a** was C₇₂H₉₂O₃₈. To compare the ¹H- and ¹³C-NMR data between **2a** and **1a**, detailed 2D-NMR experiments were performed and the coupling constants were assigned using 1D-TOCSY. The chemical shifts and coupling constants of glu-




Fig. 1. Chemical Structures of Crocin (**1**) and Neocrocetin A (**2**)

* To whom correspondence should be addressed. e-mail: nsugimot@nihs.go.jp

Table 1. ^1H (800 MHz)- and ^{13}C (200 MHz)-NMR Spectral Data^a of Crocin (1) and Neocrocin A (2) (in $\text{DMSO}-d_6/\text{D}_2\text{O}$ (9:1))

Crocin (1) ^b						Neocrocin A (2) ^b					
α isomer			β isomer			α isomer			β isomer		
No.	δ_{H}	δ_{C}	No.	δ_{H}	δ_{C}	No.	δ_{H}	δ_{C}	No.	δ_{H}	δ_{C}
8, 8'	166.80	166.72	8	167.92	167.92	8			8		
9, 9'	125.70	125.77	9	126.63	126.63	9			9		
10, 10'	140.50	140.44	10	139.42	139.42	10	7.31 (d, 11.4)	7.31 (d, 11.4)	10	7.20 (d, 11.0)	7.20 (d, 11.0)
11, 11'	6.62 (br t, 14.7)	124.40	11	6.63 (br t, 13.1)	124.52	11	6.63 (br t, 13.1)	6.63 (br t, 11.7)	11	6.62 (br t, 13.1)	6.62 (br t, 13.1)
12, 12'	6.77 (d, 14.7)	145.20	12	6.78 (d, 14.6)	144.40	12	6.78 (d, 14.6)	6.78 (d, 14.6)	12	6.72 (d, 15.1)	6.72 (d, 15.1)
13, 13'	137.40	137.45	13	137.35	137.35	13			13		
14, 14'	6.50 (br d, 7.8)	136.60	14	6.49 (br d, 8.2)	136.15	14	6.49 (br d, 8.2)	6.49 (br d, 8.2)	14	6.49 (br d, 8.2)	6.49 (br d, 8.2)
15, 15'	6.81 (br d, 7.8)	132.50	15	6.82 (br d, 8.2)	132.32	15	6.82 (br d, 8.2)	6.82 (br d, 8.2)	15	6.82 (br d, 8.2)	6.82 (br d, 8.2)
19, 19'	1.92 (s)	13.16	19	1.92—1.95 ^c	13.08—13.48 ^d	19	1.92—1.95 ^c	1.92—1.95 ^c	19	1.92—1.95 ^c	1.92—1.95 ^c
20, 20'	1.94 (s)	13.04	20		13.08—13.48 ^d	20		13.08—13.48 ^d	20		13.08—13.48 ^d
Glucose A, C			Glucose C			Glucose A			Glucose C		
1	5.37 (d, 7.8)	95.00	1	5.06 (d, 3.6)	89.58	1	5.38 (d, 7.8)	95.05	1	4.53 (d, 6.9)	94.80
2	3.18 (t, 7.8)	72.78	2	4.39 (dd, 3.7, 10.1)	74.87	2	4.39 (dd, 3.7, 10.1)	74.87	2	4.53 (t, 6.9)	75.87
3	3.23 (t, 8.0)	76.51	3			3			3		
4	3.20 (t, 8.0)	69.57	4	2.90—4.10 ^c	68.00—77.00 ^d	4	2.90—4.10 ^c	68.00—77.00 ^d	4	2.90—4.10 ^c	68.00—77.00 ^d
5	3.38 (m)	76.70	5			5			5		
6	3.54 (m)	68.36	6			6			6		
	3.94 (d, 10.1)										
Glucose B, D			Glucose D			Glucose B			Glucose D		
1	4.13 (d, 7.8)	103.49	1	4.13 (d, 7.8)	103.70	1	4.13 (d, 7.8)	103.60	1	4.15 (d, 7.8)	103.70
2	2.91 (t, 8.0)	71.88	2			2			2		
3	3.08 (t, 8.0)	77.00	3			3			3		
4	2.99 (t, 8.0)	70.37	4	2.90—4.10 ^c	61.00—77.00 ^d	4	2.90—4.10 ^c	61.00—77.00 ^d	4	2.92—4.10 ^c	61.00—77.00 ^d
5	3.01 (m)	77.23	5			5			5		
6	3.38 (m)	61.38	6			6			6		
	3.60 (d, 10.5)										

a) Chemical shifts are in ppm and coupling constants in Hz are in parentheses. b) Assignments were made using 1D-TOCSY, COSY, HMQC and HMBC. c) Signals were overlapped and could not be assigned.

Table 2. ^1H (800 MHz)- and ^{13}C (200 MHz)-NMR Spectral Data^{a)} of Peracetylated Crocin (1a) and Neocrocins A (2a) (in CDCl_3)

Peracetylated crocin (1a) ^{b)}			Peracetylated neocrocins A (2a) (α isomer) ^{b)}					
No.	δ_{H}	δ_{C}	No.	δ_{H}	δ_{C}	No.	δ_{H}	δ_{C}
8, 8'		166.09	8		166.07	8'		166.92
9, 9'		124.57	9		124.55	9'		125.02
10, 10'	7.33 (d, 12.4)	141.45	10	7.32 (d, 11.4)	141.45	10'	7.20 (d, 12.4)	140.46
11, 11'	6.54 (dd, 11.7, 14.9)	123.66	11'	6.53 (dd, 11.5, 15.1)	123.71	11'	6.50 (dd, 11.7, 14.9)	123.63
12, 12'	6.68 (d, 14.5)	145.42	12	6.68 (d, 15.2)	145.42	12'	6.60 (d, 15.1)	144.85
13, 13'		138.00	13		136.96	13'		136.85
14, 14'	6.42 (d, 10.1)	136.95	14	6.41 (d, 9.7)	136.33	14'	6.39 (d, 10.1)	136.02
15, 15'	6.73 (dd, 2.1, 8.0)	131.83	15	6.71 (brdd)	131.84	15'	6.72 (brdd)	131.67
19, 19'	1.98 (s)	12.87	19	1.97 (s)	12.70	19'	1.91 (s)	12.80
20, 20'	1.99 (s)	12.73	20	1.98 (s)	12.90	20'	1.98 (s)	12.90
Glucose A, C			Glucose A		Glucose C			
1	5.74 (d, 8.2)	92.08	1	5.74 (d, 7.8)	92.08	1	6.73 (d, 3.7)	88.99
2	5.20 (t, 9.0)	70.29	2	5.20 (t, 9.0)	70.29	2	5.07 (dd, 3.9, 10.3)	69.63
3	5.27 (t, 9.4)	72.90	3	5.26 (t, 9.6)	72.90	3	5.55 (t, 9.9)	70.08
4	5.01 (t, 9.6)	68.42	4	5.01 (t, 9.8)	68.38	4	5.03 (t, 10.1)	68.47
5	3.83 (ddd, 2.4, 6.4, 10.5)	74.01	5	3.82 (m)	74.01	5	4.05 (m)	70.99
6	3.59 (dd, 5.8, 11.7)	67.68	6	3.59 (dd, 5.8, 11.7)	67.68	6	3.53 (dd, 6.0, 11.0)	67.96
	3.93 (dd, 2.3, 11.4)			3.92 (1H, brd)			3.93 (brd)	
Glucose B, D			Glucose B'		Glucose D			
1	4.55 (d, 8.2)	100.65	1	4.55 (d, 8.2)	100.65	1	4.53 (d, 7.8)	100.85
2	4.98 (dd, 7.8, 9.6)	70.99	2	4.97 (t, 8.9)	70.99	2	4.97 (t, 8.9)	70.99
3	5.17 (t, 9.4)	72.90	3	5.17 (t, 9.4)	72.90	3	5.19 (t, 9.4)	72.76
4	5.05 (t, 10.2)	68.42	4	5.05 (t, 9.4)	68.43	4	5.07 (t, 9.9)	69.03
5	3.63 (ddd, 2.3, 4.6, 10.1)	71.98	5	3.63 (m)	71.98	5	3.67 (ddd, 2.4, 4.5, 10.0)	72.02
6	4.09 (dd, 2.3, 12.4)	61.88	6	4.09 (dd, 1.8, 12.4)	61.88	6	4.11 (dd, 2.1, 12.2)	61.88
	4.24 (dd, 4.6, 12.4)			4.23 (dd, 5.0, 12.3)			4.26 (dd, 4.6, 12.4)	
Acetyl	2.00–2.07 (s)	20.65–20.80		1.98–2.06 (s)	20.65–20.94			
		169.45–170.73			169.36–170.79			

a) Chemical shifts are in ppm and coupling constants in Hz are in parentheses. b) Assignments were made using 1D-TOCSY, COSY, HMQC and HMBC.

crocin A and B of **2a** were identical to those of **1a**, and the HMBC correlations of the H-1 (δ_{H} 5.74) of glucose A to C-8 (δ_{C} 166.07) of crocetin, and the H-1 (δ_{H} 4.55) of glucose B to C-6 (δ_{C} 67.68) of glucose A, suggested the existence of a gentiobiosyl moiety connected to crocetin at the anomeric position, similar to in **1a**. By contrast, the chemical shifts and coupling constants of glucose C suggested that the anomeric position of glucose C was in the α configuration, and the HMBC correlations of the H-2 (δ_{H} 5.07) of glucose C to C-8' (δ_{C} 166.07) of crocetin, and the H-1 (δ_{H} 4.53) of glucose D to C-6 (δ_{C} 67.96) of glucose C, supported the notion that another gentiobiosyl moiety was connected to the crocetin *via* the number 2 position on glucose C. The ^1H - and ^{13}C -NMR data for **1a** and **2a** are summarized in Table 2 and key correlations of **2a** are shown in Fig. 2. These observations provided evidence that neocrocins A (**2**) possessed a unique binding system for sugars.

As neocrocins A (**2**) isolated from gardenia yellow was a mixture of isomers along with crocin (**1**), we initially suspected that neocrocins A (**2**) was an artefact formed by crocin (**1**) *via* the rearrangement of the acyl moiety during the isolation process. However, the LC profile of gardenia yellow, especially the peak height of neocrocins A (**2**), remained unchanged after dissolving again with ethanol and water under acidic and basic conditions. We thus concluded that neocrocins A (**2**) was an original constituent of the *G. jasminoides* fruits. To our knowledge, this is the first published evidence that the binding system for the sugars of a crocetin glycosyl ester is not limited to the anomeric position




Fig. 2. Key Correlations of Peracetylated Neocrocins A (**2a**)

in nature. We are further investigating the possibility that other types of crocetin glycoside might also exist in this fruit through an ongoing study.

Experimental

General IR spectra were recorded on a JASCO FT/IR-4100. UV/Vis spectra were recorded on a JASCO UV-560. FAB-MS spectra were measured with JEOL JMS-700 system and ESI-MS spectra were measured with JEOL JMS-T100LC "AccuTOF". NMR spectra were recorded on JEOL JNM-ECA (800 MHz) and JNM-ECA (500 MHz). Diaion HP-20 was used for open-column chromatography. The preparative LC-MS system (Waters FractionLynxTMMS auto purification system), which is described in our previous report,⁹ was used for isolation of compounds on this research. The