

effect on the expansion of human CB stem/progenitor cells using this coculture system.

Materials and Methods

HbVs

Hemoglobin vesicles were prepared under sterile conditions in a similar manner as previously described.^{22,23} In brief, the Hb was purified from outdated donated blood provided by the Japanese Red Cross Society (Tokyo, Japan). The encapsulated Hb solution (38 g/dL) contained 14.7 mM pyridoxal 5'-phosphate (PLP) at a molar ratio of [PLP]/[Hb] = 2.5 as an allosteric effector. The lipid bilayer was composed of 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, cholesterol, 1,5-O-dihexadecyl-N-succinyl-L-glutamate (Nippon Fine Chemical Co. Ltd, Osaka, Japan), and 1,2-distearoyl-sn-glycero-3-phosphatidylethanol-amine-N-[poly(ethylene glycol) (5,000)] (NOF Corp., Tokyo, Japan), at a molar ratio of 5:5:1:0.033. The physicochemical parameters were P_{50} , 27 Torr; 262 ± 77 nm particle diameter; and <3% MetHb content. The concentration of Hb in the HbVs dispersion was adjusted to 10 g/dl. The concentration of HbVs in this study was set at about 3 vol/vol%, based on the following rationale. Intravenously injected HbVs are eventually captured by phagocytes in the reticuloendothelial system, including the spleen, liver, and bone marrow. The circulation half-life of HbVs in the elimination phase in humans has been estimated to be approximately 66 h by the study of circulation kinetics using rats and rabbits,⁸ and the percentage of the infused dose of HbVs of bone marrow in humans was estimated to be 6.4% at 48 h after 25% top-loading of HbVs, in studies of the organ distribution of HbVs in rats and rabbits.⁸ Based on this estimation, the distribution of HbVs in the human bone marrow at 48 h after infusion at 25 vol/vol% (1225 ml of HbVs) of the blood volume (4.9 L, 70 ml/kg, body weight) in a 70-kg individual could be expected to be 78.4 ml (6.4 vol/vol% of the infused dose of HbVs). The volume of the bone marrow space has been estimated as 2.6–4 L in an average-sized human (70 kg).²⁴ From these values, the amount of HbVs in the human bone marrow can be calculated to be about 2 to 3 vol/vol%.

Preparation of Human CB

Experiments using human umbilical CB were approved by the Committee of Hokkaido CB Bank. Cord blood was obtained during normal full-term deliveries. Cord blood CD34⁺ cells were prepared as described previously.²⁵ Briefly, after the sedimentation of red blood cells by incubating CB samples with the same volume of 6% (w/v) hydroxyethyl starch dissolved in Ringer's solution (Veen-D Inj., Nikken Chemical, Tokyo, Japan) at room temperature for 30 min, low-density (<1.077 g/ml) mononuclear cells were collected with Ficoll-Paque (Pharmacia biotech, Uppsala, Sweden). The cells were then enriched further with CD34⁺ cells using a MACS CD34 Progenitor Isolation Kit (Miltenyi Biotec, Bergish-Gladbach, Germany) according to the manufacturer's instructions. The purity of CD34⁺ cells was >85%.

Coculture of CD34⁺ Cells and hTERT-Transfected Stromal Cells

The coculture of CD34⁺ cells and hTERT-transformed cells was carried out as previously described.²⁵ Briefly, hTERT-

transfected human stromal cells were cultured in 25 cm² flasks at 37°C in 5% CO₂ in a humidified atmosphere using a culture medium comprising Iscove's modified Dulbecco's medium (IMDM, Invitrogen, Carlsbad, Ca), 12.5% FCS (Equitech Bio Inc., Igram, TX), 12.5% horse serum [Invitrogen], 1 μM hydrocortisone (BSA; Sigma Chemical, St Louis, MO), and 50 μM β-mercaptoethanol. The cells were cultured until confluent. On the day of the coculture, the stromal cells were washed several times with phosphate buffered saline before the addition of CB CD34⁺ cells. Some 5,000 CB CD34⁺ cells were seeded on a monolayer of hTERT-stromal cells that had been preestablished in 4 ml of serum-free medium, X-VIVO 10 (BioWhittaker, Walkersville, MD), supplemented with 50 ng/ml human thrombopoietin (TPO: a gift from Kirin Brewery Co LTD, Tokyo, Japan), 10 ng/ml stem cell factor (SCF), and 100 ng/ml human Flt-ligand (FL: a gift from Immunex Corporation, Seattle, WA), in the presence of HbVs (final concentration, up to 3%). After 1 week of coculture, 4 ml of fresh complete medium containing these same concentrations of cytokines without HbVs were added, and the coculture was continued for 1 week. At this point, each concentration of HbVs was halved. At the end of the second week, both expanded hematopoietic cells that had not attached to the stromal cells and those that had attached weakly to them were collected by gentle pipetting. Cells attached strongly to stromal cells were also recovered by treatment with ethylenediamine-tetra-acetic acid and trypsin after the removal of non-adherent cells. Stromal cells were removed by culturing in a tissue culture flask for 1 h. A mixture of the hematopoietic cells was washed to remove HbVs and the population of CD34⁺ cells was determined with a flow cytometer. Subsequently, expanded cells (300 CD34⁺ cells/dish) were subjected to a clonal assay.

In the experiment for examining the effect of the exposure period to HbVs, HbVs (0%, 1.5%, or 3%) were added to the coculture for either 3 days or 7 days, at the end of which both hematopoietic cells that had not attached to the stromal cells and those that had attached weakly to them were collected by gentle pipetting, washed by repeated centrifugation to remove HbVs, and resuspended in complete culture medium without HbVs. The monolayers of h-TERT stromal cells were washed with phosphate-buffered saline twice to remove the residual HbVs, and were cultured with the resuspended hematopoietic cells as described above up to a total of 14 days.

Clonal Cell Culture

The methylcellulose clonal culture was carried out as described previously, except for absence of HbVs. The population of CD34⁺ cells among the mononuclear cells was determined by flow cytometry, and CB-derived mononuclear cells were seeded at 300 CD34⁺ cells/dish. Burst-forming units of erythrocyte (BFU-E), colony-forming units of granulocytes/macrophages (CFU-GM), CFU-mix, and colony forming units in culture (CFC) were scored with an inverted microscope after 14 days of incubation. Densely packed colonies that reached >1 mm in size were scored as high-proliferative potential colonies (HPP-CFC) after 28 days of incubation.




Figure 1. Effects of hemoglobin vesicles (HbVs) on expansion of stem/progenitor cells *in vitro*. Five thousand CD34⁺ cells derived from (cord blood) CB were cultured on a monolayer of h-TERT-transfected human stromal cells in the presence of thrombopoietin (TPO), stem cell factor (SCF), and Flt-ligand (FL). Various concentrations of HbVs were added to the coculture system. Two weeks later, the total number of expanded cells, and the number of expanded CD34⁺ cells were determined. Data represent the mean \pm SD of three experiments performed on three separate CB donors. * $p < 0.01$ versus HbVs (0%).

Statistical Analysis

Results are expressed as the mean \pm standard deviation (SD). A two-way nonrepeated analysis of variance followed by Bonferroni's test was used for multiple comparisons between the control (HbVs; 0%) group and a treatment group. For analysis of the difference between HbVs and empty liposomes, paired t test was used. Values of $p < 0.05$ were considered statistically significant.

Results

Effects of HbVs on Expansion of Hematopoietic Stem/Progenitor Cells

The coculture of hematopoietic stem/progenitor cells with bone marrow-derived stromal cells mimics the physiological hematopoiesis in bone marrow more closely than does a liquid culture. CD34⁺ cells were cocultured for 2 weeks with h-TERT stromal cells in the presence of different concentrations of HbVs. The total cell numbers obtained without HbVs were $1.28 \times 10^6 \pm 2.7 \times 10^5$, showing an approximately 260-fold increase of cells. The continuous presence of HbVs at 1.5% for 1 week and then at 0.75% for an additional week, caused a




Figure 2. Effects of empty liposomes on the expansion of stem/progenitor cells. Experimental conditions were the same as those of Figure 1 except that either 3% (vol/vol) empty liposomes (closed column) or hemoglobin vesicles (HbVs) (open column) were added to the coculture for 14 d. * $p < 0.05$, ** $p < 0.01$, empty liposomes vs HbVs; paired Student's t test.

significant reduction in the proliferation of cells. Furthermore, the presence of HbVs at 3% for 1 week and then at 1.5% for an additional week, caused a marked reduction (Figure 1). The CD34⁺ cells obtained without HbVs were $9.3 \times 10^4 \pm 3.9 \times 10^4$, showing an approximately 20-fold expansion. Again, the presence of HbVs for 14 days dose-dependently inhibited the expansion of CD34⁺ cells (Figure 1).

A colony assay of the CD34⁺ cells expanded with or without HbVs was carried out (Table 1). The number of CD34⁺ cells expanded in the presence of 3% HbVs was 17.7% of those without HbVs. Similarly, the numbers of BFU-E, CFU-GM, CFU-mix, CFC, and primitive HPP-CFC obtained from the CD34⁺ cells expanded in the presence of HbVs were 49.9%, 15.8%, 18.7%, 18.2%, and 12.3%, respectively, of those without HbVs (Table 1).

Effects of Empty Liposomes on the Expansion of Hematopoietic Stem/Progenitor Cells

We then tested the question whether the inhibitory effect of HbVs on the expansion of total cells as well as CD34⁺ cells themselves was due to either the liposomes themselves or Hb in HbVs. To do this, CD34⁺ cells were cocultured with h-TERT stromal cells in the presence of either HbVs or empty liposomes for 14 days. As shown in Figure 2, the 14-day exposure of CD34⁺ cells to empty liposomes reduced expansion of the

Table 1. Clonogenic Assay of Expanded Cells

HbV	CD34 ⁺ Cells	BFU-E	CFU-GM	CFU-Mix	CFC	HPP-CFC
0%	$9.6 \times 10^4 \pm 6.4 \times 10^3$	1370 ± 1150	15500 ± 2570	1270 ± 1360	18240 ± 3440	2530 ± 1260
3%	$1.7 \times 10^4 \pm 9.5 \times 10^3$ (17.7%)	683 ± 76 (49.9%)	2450 ± 700 (15.8%)	237 ± 70 (18.7%)	3320 ± 715 (18.2%)	310 ± 90 (12.3%)

Five thousand CD34⁺ cells derived from CB were cultured without or with HbV as shown in the legend of Figure 1. After 14 d of coculture, the total hematopoietic cells were counted, and the population of CD34⁺ cells was determined with flow cytometry. Subsequently, expanded hematopoietic mononuclear cells were seeded at 300 CD34⁺ cells/dish. The numbers of colonies shown in the Table were calculated by the following equation: (the numbers of colony formed) \times (total CD34⁺ cells)/300.

Parentheses show the percentage of the control (HbV 0%).

Data represent the mean \pm SD of 3 experiments performed on 3 separate CB donors.

HbV, Hemoglobin vesicles; BFU-E, burst-forming units of erythrocyte; CFU-GM, colony-forming units of granulocytes/macrophages; CFC, colony forming units in culture; HPP-CFC, high-proliferative potential colonies.




Figure 3. Effects of short exposure to hemoglobin vesicles (HbVs) on the expansion of stem/progenitor cells derived from cord blood (CB). Cord blood-derived CD34⁺ cells were exposed to HbVs (0%, 1.5%, or 3%) for 3 d (open circle), 7 d (open square), or 14 d (closed circle). After culture for a total of 14 d, the total number of expanded cells and the number of expanded CD34⁺ cells were determined, respectively. Data represent the mean \pm SD of three experiments performed on three separate CB donors. A two-way paired analysis of variance (ANOVA) followed by Bonferroni's test was used for comparisons of multiple HbV-treated groups with the control (HbVs 0%) group. * $p < 0.05$, ** $p < 0.01$ versus HbVs (0%).

total cells. However, the liposomal inhibitory effects were much less than those of HbVs. Similarly, the 14-day exposure of CD34⁺ cells to empty liposomes only modestly inhibited the expansion of CD34⁺ cells themselves as compared with those by HbVs (Figure 2).

Effects of Short Exposure to HbVs on the Expansion of Hematopoietic Stem/Progenitor Cells

Next, the effects of the short exposure of CD34⁺ cells to HbVs on the expansion of both total cells and CD34⁺ cells were examined. Exposure to HbVs even at 3% for 3 days only slightly inhibited the increase of the number of the total cells and did not affect expansion of CD34⁺ cells (Figure 3). In the case of 7-day exposure, the expansion of total cells and CD34⁺ cells was reduced by a similar extent as was observed in the case of continuous exposure.

Discussion

We previously examined the biocompatibility of HbVs with CB hematopoietic progenitor cells by testing whether HbVs might affect the clonogenic activity of progenitor cells in a semisolid culture, and the proliferation and differentiation of erythroid and myeloid lineages from progenitor cells in a liquid culture. In the study, we showed that a short exposure (1–3 days) to HbVs had no adverse effects on progenitor activity, although a longer exposure (14 days) suppressed progenitor activity.¹⁴ Under these two culture conditions, hematopoietic stem/progenitor activity could not be tested.

In the present study, therefore, we used the coculture system with hTERT-transfected human stromal cells and CD34⁺ cells, which mimics the physiological conditions of hematopoiesis in the bone marrow in terms of the interaction between hematopoietic stem/progenitor cells and stromal cells, and importantly supports *in vitro* expansion of hematopoietic stem/progenitor cells.²¹ Under this system, we found that 14-day exposure to HbVs significantly suppressed expansion of hema-

topoietic stem/progenitor cells, although the effect of 3-day exposure to HbVs was minimal. This observation is in agreement with our previous result.¹⁴

It has been shown that heme (the ferric chloride salt of heme) exerts a proliferative effect on erythroid progenitor cells. The administration of heme to mice increases the number of BFU-E within the bone marrow *in vivo*.²⁶ Heme was also shown to stimulate a two-fold increase in a murine erythroid colony compared with those stimulated by Epo alone in the *in vitro* culture.²⁷ Along with heme's effect on erythroid progenitor cells, purified and cross-linked hemoglobin has also been shown to have erythropoietic activity.²⁸ Purified and other chemically modified hemoglobin, as a blood substitute, has been shown to support expansion of erythroid progenitor cells from CB-derived CD34⁺/CD38⁻ and CD34⁺/CD33⁻ stem/progenitor cells in a liquid culture.²⁹ In addition, under a human CD34⁺ bone marrow cell liquid culture system, a cross-linked human recombinant hemoglobin alone did not affect the proliferation of erythroid and myeloid lineage cells, but protected erythroid lineage cells from 3'-azido-3'-deoxythymidine (AZT)-mediated inhibition.³⁰ The protection of this recombinant hemoglobin from AZT toxicity was also achieved in animal models.³¹ The authors speculated that free heme could be released from hemoglobin before its uptake by erythroid progenitor cells or, alternatively, intact hemoglobin could be taken up intracellularly by the cells before the release of heme. As a result, heme stimulates proliferation and differentiation of erythroid progenitor cells. These erythropoietic activities observed in an acellular type of hemoglobin substitute were not observed in HbVs, cellular-type artificial oxygen carriers, in our culture conditions. HbVs might degrade during long-term incubation, leading to the release of Hb. However, our previous study showed that the released Hb level in the liquid culture in the presence of 3% HbVs was low enough not to affect the proliferation of erythroid and myeloid lineage cells.¹⁴ Another possible explanation is that the uptake of HbVs may not occur in erythroid progenitor cells as efficiently as that of modified hemoglobin.

The suppression of expansion by 14-day exposure to HbVs appeared to be mostly attributed to Hb itself in the HbVs, because the effect of empty liposomes was much less than that of HbVs. The modest inhibitory effect observed in the liposomes might be caused by physical suppression of the interaction of CD34⁺ cells and stromal cells.

There are several explanations for the observed suppression by 14-day exposure to HbVs. Higher dissolved oxygenations in the culture medium were theoretically more expected in the presence of HbVs than in the absence of HbVs, which may be involved in the inhibition of expansion of hematopoietic stem/progenitor cells after the prolonged exposure to HbVs. In addition, Hb in HbVs might autoxidize during 14 days in culture, producing reactive oxygen species, which in turn cause lipid peroxidation of liposomes, resulting in the generation of lipid radicals. The production of lipid peroxides and their by-products ultimately causes the loss of membrane function and integrity of cells including hematopoietic stem/progenitor cells.^{32,33} In fact, conversion of Hb to Met-Hb within HbVs occurs.³⁴ In addition, concentrated Hb within HbVs interacted with exogenously added H₂O₂, and subsequently Met-Hb and ferrylHb were formed inside the HbVs.³⁵ However, there was no peroxidation of lipids comprising the vesi-

cles,³⁵ due to the saturated lipid membrane of HBVs. Furthermore, when vesicles composed of egg yolk lecithin as unsaturated lipids were added to the mixture of HBVs and H₂O₂, no lipid peroxidation was observed in vesicles composed of egg yolk lecithin.³⁵ This is due to the stable encapsulation within HBVs of the products of Met-HbV and ferrylHb. Thus, the possibility that lipid peroxidation may be involved in a suppressive effect associated with longer exposure is less likely.

We found exposure time-dependent suppression of stem/progenitor activity by HBVs in this *in vitro* study. In contrast, our animal studies demonstrated a lack of any suppressive effects of HBVs on hematopoiesis. For instance, rats and beagle dogs after receiving an acute 40% exchange-transfusion with HBVs, showed elevated erythropoietic activity and completely recovered from the decreased hematocrit within 7 days.^{36,37} In addition, the number of red blood cells, leukocytes, and platelets were maintained constantly for 1 week after the infusion of HBVs into rats at 20% of the whole blood volume.³⁸ In the animal model, HBVs accumulated transiently in the bone marrow by 1–3 days and significantly decreased at 7 days, and then became undetectable at 14 days after infusion of HBVs.³⁶ The HBVs distributed in the bone marrow were accumulating in bone marrow macrophages, and the HBVs were degraded in the phagosomes in the reticuloendothelial system.³⁹ However, an *in vitro* coculture system lacks any reticuloendothelial system in addition to any shielding by the vasculature and blood stream, and these closed systems might magnify the direct contact between HBVs and CD34⁺ cells, resulting in the different outcome (*i.e.*, the exposure time-dependent suppression of stem/progenitor activity).

In conclusion, although the absence of adverse effects on hematopoiesis by HBVs in animal models suggests that HBVs, when transfused into humans, may have minimal side effects on hematopoiesis, the present data raise some concern regarding hematopoiesis, and one should therefore pay attention to the effect of HBVs on various hematological aspects in future clinical setting.

Acknowledgment

Supported, in part, by Health and Labor Sciences Research Grants (Research on Regulatory Science of Pharmaceuticals and Medical Devices), Ministry of Health, Labor and Welfare, Japan, and by Oxygen Inc., Tokyo, Japan.

References

- Djordjevic L, Miller IF: Synthetic erythrocytes from lipid encapsulated hemoglobin. *Exp Hematol* 8: 584–592, 1980.
- Phillips WT, Klipper RW, Awasthi VD, et al: Polyethylene glycol-modified liposome-encapsulated hemoglobin: A long circulating red cell substitute. *J Pharmacol Exp Ther* 288: 665–670, 1999.
- Chang TM: Hemoglobin-based red blood cell substitutes. *Artif Organs* 28: 789–794, 2004.
- Sakai H, Takeoka S, Park SI, et al: Surface modification of hemoglobin vesicles with poly(ethylene glycol) and effects on aggregation, viscosity, and blood flow during 90% exchange transfusion in anesthetized rats. *Bioconjug Chem* 8: 23–30, 1997.
- Sakai H, Masada Y, Horinouchi H, et al: Hemoglobin-vesicles suspended in recombinant human serum albumin for resuscitation from hemorrhagic shock in anesthetized rats. *Crit Care Med* 32: 539–545, 2004.
- Cabrales P, Sakai H, Tsai AG, et al: Oxygen transport by low and normal oxygen affinity hemoglobin vesicles in extreme hemodilution. *Am J Physiol Heart Circ Physiol* 288: 1885–1892, 2005.
- Yoshizu A, Izumi Y, Park S, et al: Hemorrhagic shock resuscitation with an artificial oxygen carrier, hemoglobin vesicle, maintains intestinal perfusion and suppresses the increase in plasma tumor necrosis factor- α . *ASAIO J* 50: 458–463, 2004.
- Sou K, Klipper R, Goins B, et al: Circulation kinetics and organ distribution of Hb-vesicles developed as a red blood cell substitute. *J Pharmacol Exp Ther* 312: 702–709, 2005.
- Shizuru JA, Negrin RS, Weissman IL: Hematopoietic stem and progenitor cells: Clinical and preclinical regeneration of the hematolymphoid system. *Annu Rev Med* 56: 509–538, 2005.
- Lutton JD, Jiang S, Drummond GS, et al: Pharmacologic effects of the red blood cell substitutes cross-linked and non-cross-linked hemoglobins on hematopoiesis in rabbits. *Pharmacology* 58: 319–324, 1999.
- Pessina A, Malerba I, Gribaldo L: Hematotoxicity testing by cell clonogenic assay in drug development and preclinical trials. *Curr Pharm Des* 11: 1055–1065, 2005.
- Giarratana MC, Kobari L, Lapillonne H, et al: Ex vivo generation of fully mature human red blood cells from hematopoietic stem cells. *Nat Biotechnol* 23: 69–74, 2005.
- Unverzagt KL, Bender JG, Loudovaris M, et al: Characterization of a culture-derived CD15+CD11b- promyelocytic population from CD34+ peripheral blood cells. *J Leukoc Biol* 62: 480–484, 1997.
- Yamaguchi M, Fujihara M, Wakamoto S, et al: Influence of hemoglobin vesicles, cellular-type artificial oxygen carriers, on human umbilical cord blood hematopoietic progenitor cells in vitro. *J Biomed Mater Res A* 88: 34–42, 2009.
- Petzer AL, Zandstra PW, Piret JW, Eaves CJ: Differential cytokine effects on primitive (CD34+CD38-) human hematopoietic cells: Novel responses to Flt3-ligand and thrombopoietin. *J Exp Med* 183: 2551–2558, 1996.
- McNiece I, Kubegov D, Kerzic P, et al: Increased expansion and differentiation of cord blood products using a two-step expansion culture. *Exp Hematol* 28: 1181–1186, 2000.
- Ueda T, Tsuji K, Yoshino H, et al: Expansion of human NOD/SCID-repopulating cells by stem cell factor, FLK2/Flt3 ligand, thrombopoietin, IL-6, and soluble IL-6 receptor. *J Clin Invest* 105: 1013–1021, 2000.
- Xu MJ, Tsuji K, Ueda T, et al: Stimulation of mouse and human primitive hematopoiesis by murine embryonic aorta-gonad-mesonephros-derived stromal cell lines. *Blood* 92: 2032–2040, 1998.
- Breems DA, Blokland EA, Siebel KE, et al: Stroma-contact prevents loss of hematopoietic stem cell quality during ex vivo expansion of CD34+ mobilized peripheral blood stem cells. *Blood* 91: 111–117, 1998.
- Bennaceur-Grisicelli A, Tourino C, Izac B, et al: Murine stromal cells counteract the loss of long-term culture-initiating cell potential induced by cytokines in CD34(+)/CD38(low/neg) human bone marrow cells. *Blood* 94: 529–538, 1999.
- Kawano Y, Kobune M, Yamaguchi M, et al: Ex vivo expansion of human umbilical cord hematopoietic progenitor cells using a coculture system with human telomerase catalytic subunit (hTERT)-transfected human stromal cells. *Blood* 101: 532–540, 2003.
- Sakai H, Yuasa M, Onuma H, et al: Synthesis and physicochemical characterization of a series of hemoglobin-based oxygen carriers: Objective comparison between cellular and acellular types. *Bioconjug Chem* 11: 56–64, 2000.
- Sou K, Naito Y, Endo T, et al: Effective encapsulation of proteins into size-controlled phospholipid vesicles using freeze-thawing and extrusion. *Biotechnol Prog* 19: 1547–1552, 2003.
- Lipton JM, Nathan DG: The anatomy and physiology of hematopoiesis, in Nathan DG, Oski FA (eds), *Hematology of Infancy and Childhood*, Philadelphia, WB Saunders, 1987, pp. 128–158.
- Ohkawara JJ, Ikebuchi K, Fujihara M, et al: Culture system for extensive production of CD19+IgM+ cells by human cord blood CD34+ progenitors. *Leukemia* 12: 764–771, 1998.

26. Monette FC, Holden SA, Sheehy MJ, Matzinger EA: Specificity of hemin action in vivo at early stages of hematopoietic cell differentiation. *Exp Hematol* 12: 782-787, 1984.
27. Porter PN, Meints RH, Mesner K: Enhancement of erythroid colony growth in culture by hemin. *Exp Hematol* 7: 11-16, 1979.
28. Rosenthal GJ, Gerber MJ: Method of stimulating hematopoiesis with hemoglobin. US Patent 5631219, 1997.
29. Bell D, Mueller SG: Enhanced stimulation of erythropoiesis. EP Patent 099644, 2004.
30. Fowler DA, Rosenthal GJ, Sommadossi JP: Effect of recombinant human hemoglobin on human bone marrow progenitor cells: Protection and reversal of 3'-azido-3'-deoxythymidine-induced toxicity. *Toxicol Lett* 85: 55-62, 1996.
31. Moqattash S, Lutton JD, Rosenthal G, et al: Effect of blood substitute, recombinant hematology, on in vivo hematopoietic recovery from AZT toxicity. *Acta Haematol* 98: 76-82, 1997.
32. Peled-Kamar M, Loten J, Okon E, et al: Thymic abnormalities and enhanced apoptosis of thymocytes and bone marrow cells in transgenic mice overexpressing Cu/Zn-superoxide dismutase: Implications for Down syndrome. *EMBO J* 14: 4985-4993, 1995.
33. Shao J, White CC, Dabrowski MJ, et al: The role of mitochondria and oxidative injury in BDE47 toxicity to human fetal liver hematopoietic stem cells. *Toxicol Sci* 101: 81-90, 2008.
34. Teramura Y, Kanazawa H, Sakai H, et al: Prolonged oxygen-carrying ability of hemoglobin vesicles by coencapsulation of catalase in vivo. *Bioconjug Chem* 14: 1171-1176, 2003.
35. Takeoka S, Teramura Y, Atoji T, Tsuchida E: Effect of Hb-encapsulation with vesicles on H₂O₂ reaction and lipid peroxidation. *Bioconjugate Chem* 13: 1302-1308, 2002.
36. Sakai H, Horinouchi H, Yamamoto M, et al: Acute 40 percent exchange-transfusion with hemoglobin-vesicles (HbVs) suspended in recombinant human serum albumin solution: Degradation of HbVs and erythropoiesis in a rat spleen for 2 weeks. *Transfusion* 46: 339-347, 2006.
37. Ikeda T, Horinouchi H, Kohno M, et al: Resuscitation effect and long term effect of Hb vesicles on organ function in beagle dog. *Artif Cells Blood Substit Immobil Biotechnol* 36: 214, 2008.
38. Abe H, Azuma H, Yamaguchi M, et al: Effects of hemoglobin vesicles, a liposomal artificial oxygen carrier, on hematological responses, complement and anaphylactic reactions in rats. *Artif Cells Blood Substit Immobil Biotechnol* 35: 157-172, 2007.
39. Sakai H, Horinouchi H, Tomiyama K, et al: Hemoglobin-vesicles as oxygen carriers: influence on phagocytic activity and histopathological changes in reticuloendothelial system. *Am J Pathol* 159: 1079-1188, 2001.

AQ: 1

ヘモグロビン小胞体の *in vitro* におけるヒト血液細胞および血漿タンパクへの適合性

Biocompatibility of Hemoglobin Vesicles, a Cellular-type Artificial Oxygen Carrier, on Human Blood Cells and Plasma Proteins *In Vitro*

藤原 満博, 東 寛, 池田 久實

Mitsuhiro Fujihara, Hiroshi Azuma, Hisami Ikeda

和文抄録

ヘモグロビン小胞体 (hemoglobin vesicles: HbV) は、脂質二重膜からなるリポソームの中にヘモグロビンを内包させたセル型の人工酸素運搬体である。HbVはいわゆる「人工赤血球」として血管内に輸注されるため、その臨床応用に向けた安全性の評価には、生体適合性、とくに血液構成成分に対する適合性をみるのが重要である。そこで我々はヒト血液細胞およびヒト血漿タンパク系を用いた *in vitro* の評価に焦点をあてた検討をおこなってきた。止血や炎症反応に関与する血小板に対し、HbV 自体には活性化作用はみられず、またアゴニストによる活性化への影響もみられなかった。自然免疫に重要な働きをする好中球の機能 (走化能、脱顆粒、活性酸素産生) の活性化に対しても、HbV は影響を与えなかった。造血機能に対しては、HbV は *in vitro* での共存が 3 日間までの短期間であれば造血前駆細胞活性を損なうことはないことが示された。またリポソームの特性から補体の活性化が懸念されたが、現行の HbV では活性化はみられず、さらに凝固系、カリクレイン・キニン系にも影響はみられなかった。以上のことから、HbV はヒト血液細胞ならびに血漿タンパクに対し高い適合性を持つことが示唆される。

Abstract

Hemoglobin vesicles (HbV), a cellular-type artificial oxygen carrier, are composed of human hemoglobin encapsulated within a phospholipid bilayer membrane. As HbV are injected intravenously, biocompatibility of the HbV with blood components is very important to ensure safety of this material for clinical use. We have evaluated this biocompatibility by focusing on the influence of HbV on human blood cells as well as plasma proteins *in vitro*. 1) As to the influences to platelets which are involved not only in the hemostasis but also in inflammation, HbV themselves did not activate platelets, and had no aberrant effect on agonist-induced platelet activation. 2) HbV did not affect on agonist-induced activation of neutrophil functions (chemotaxis, degranulation, and production of superoxide) which play important roles on innate immunity. 3) HbV had no effect on the hematopoietic progenitor activity, if the exposure period is brief. 4) The present HbV (containing DHSG) did not activate complement system, although the old-type HbV (containing DPPG, no PEG modification) did markedly. 5) The coagulation as well as kallikrein-kinin cascades were not affected by the present HbV. Thus, our *in vitro* studies show that HbV are highly biocompatible with human blood cells and human plasma proteins.

Keywords

artificial oxygen carrier, biocompatibility, platelets, neutrophils, hematopoietic progenitor cells, complement, coagulation, kallikrein-kinin

はじめに

赤血球製剤の人工代替物として人工酸素運搬体の開発が進められてきている。人工酸素運搬体の臨床応用における利点は、

輸血時の血液型判定の必要がないこと、製剤の長期保存が可能となること、同種抗原感作や細菌感染およびウイルス感染の伝播を回避できること等があげられる¹⁾。これらの利点から、災

北海道赤十字血液センター 研究部 〒063-0002 札幌市西区山の手2条2丁目 Hokkaido Red Cross Blood Center, Yamanote 2-2, Nishi-ku, Sapporo 063-0002, Japan

論文受付 2008年11月12日 論文受理 2008年12月3日

害時の緊急事態において大量の血液製剤が必要とされる時、また血液型検査および適合した血液製剤の供給を待機する間、人工酸素運搬体を応急措置として用いることが想定される。現行の同種血輸血や自己血輸血との組み合わせにより、人工酸素運搬体は合理的な輸血システムを構築することに寄与することが期待される。

ヘモグロビン小胞体 (hemoglobin vesicles; HbV) は、粒子径約250nmのリポソーム包埋型ヘモグロビン¹⁾。細胞膜を模擬し人工的に作成したリン脂質/コレステロール/陰性荷電脂質/PEG脂質からなる脂質二重膜のリポソームの中にヘモグロビンを内包させたセル型的人工酸素運搬体である。HbVの人工赤血球としての酸素運搬能に関する有効性、組織分布、代謝等についてはこれまで詳細に検討されている¹⁾。一方、HbVの臨床応用を進めていく上では、その生体適合性を評価することが重要である。生体適合性を評価すべき項目は多岐にわたる²⁾。その中で我々は、HbVが輸注された際に相互作用を起しうる血液構成成分、即ち、血液細胞および血漿タンパクに対しHbVがどのような影響をあたえるかについて焦点をあてた検討をおこなってきた。本稿においてはヒト血液細胞として血小板、好中球、造血前駆細胞を、ヒト血漿タンパクとして補体系、血液凝固系、カリクレイン-キニン系をとりあげ、HbVのin vitroにおけるそれぞれへの影響について我々の検討結果を概説し、HbVの生体適合性について言及したい。

1. ヘモグロビン小胞体の血小板活性化に対する影響

げっ歯類の動物モデルにおいては、リポソーム包埋型ヘモグロビン投与による副作用が知られ、その症状の一つとして一過性の急性血小板減少が報告されている³⁾。同様な一過性の急性血小板減少は、陰性荷電のリポソームの投与においてもみられている⁴⁾。この急性血小板減少の発生原因を解明するため、in vitroにおいてリポソームとげっ歯類の血小板との結合に関して研究がなされた結果⁵⁾、リポソームの構成成分が負電荷の脂質である場合、ラット多血小板血漿とのインキュベーションによりリポソームは血漿中の補体成分C3bに覆われ、C3bを介したリポソーム-C3b-血小板上CR1 receptorの結合による凝集塊が形成されることが示された。そしてin vivoにおいては、この凝集塊が速やかに網内系によって処理されることが急性血小板減少の原因と考えられている⁶⁾。これに対しヒト血小板上にはCR1 receptorが存在しないため、リポソームと血小板との凝集塊は観察されない⁷⁾。従ってヒトにおいてはリポソームの構成成分が負電荷の脂質であったとしても、動物モデルで観察されるリポソーム輸注後の急性血小板減少は発症しないと推測される。

一方、血小板はその顆粒中に種々の炎症性生理活性物質を含有していることや、膜表面の接着分子を介して白血球と結合することから、止血血栓の形成のみならず炎症反応にも関与していると考えられている⁸⁾。HbVの投与により急性の血小板減少症が起らないとしても、HbVが生体内で血小板を活性化したり、あるいはアゴニストによる反応を助長させたりするよ

うな影響を及ぼし、血小板由来生理活性物質の放出が起こるといふ事態は、生体にとって好ましいことではない。

このような背景のもと、われわれはリポソームの構成成分としてDPPG (1,2-dipalmitoyl-*sn*-glycero-3-phosphatidyl glycerol) を含有するHbVの血小板活性化への影響について、RANTESの放出反応を指標として検討した⁹⁾。RANTESは血小板の α 顆粒に存在する炎症性ケモカインで、好塩基球、好酸球、単球およびリンパ球に対する強力な走化作用を有する¹⁰⁾。そこでヒト多血小板血漿と最大20% (vol/vol) までの濃度のHbVを予めインキュベーション後、アゴニストであるコラーゲンで刺激した反応上清中のRANTESの濃度の増加を測定した。その結果、コラーゲンの濃度に依存したRANTESの放出がみとめられたが、20% HbVと予めインキュベーションさせた場合においては、RANTESの放出への影響はみられなかった (Table 1)。またアゴニスト刺激なしの場合においても、RANTESの放出に関して血小板とHbVを予めインキュベーションした影響はならみとめられないことから、HbVが血小板を活性化する作用もないと考えられた。

次に、リポソームの構成成分としてDPPGをカルボン酸脂質DHSG (1,5-*O*-dihexadecyl-*N*-succinyl-L-glutamate) に置き換えた現行のHbVの血小板への影響について、HbV濃度最大40%まで検討した¹¹⁾。血小板活性化の評価は、RANTESに加え同じく α 顆粒に存在する炎症性ケモカイン β -thromboglobulin (β -TG)、濃染顆粒に存在するセロトニンの放出反応、ポジティブフィードバックによって血小板活性化を増幅するThromboxane A_2 (TXA₂) の産生、さらに α 顆粒由来のCD62Pの膜表面への発現、および血小板膜糖タンパク α IIb β 3の構造変化を認識する抗体PAC-1の結合を指標とした。なお、TXA₂は半減期が短いためその代謝産物のThromboxane B₂ (TXB₂) を測定した (Table 1)。

Table 1. Effect of HbV on human platelets.

Index	Stimulant	Type of HbV	Conc. of HbV	Effect
RANTES release	Collagen (+) (-)	DPPG-HbV	≤ 20%	No effect
		DPPG-HbV	≤ 20%	No effect
RANTES release	Collagen (+) (-)	DHSG-HbV	≤ 40%	No effect
		DHSG-HbV	≤ 40%	Marginal reduction
β -TG release	Collagen (+) (-)	DHSG-HbV	≤ 40%	No effect
		DHSG-HbV	≤ 40%	No effect
Serotonin release	Collagen (+) (-)	DHSG-HbV	≤ 40%	No effect
		DHSG-HbV	≤ 40%	No effect
TXB ₂ production	Collagen (+) (-)	DHSG-HbV	≤ 40%	No effect
		DHSG-HbV	≤ 40%	No effect
CD62 expression	ADP (+) (-)	DHSG-HbV	≤ 40%	No effect
		DHSG-HbV	≤ 40%	No effect
PAC-1 binding	ADP (+) (-)	DHSG-HbV	≤ 40%	Slight potentiation
		DHSG-HbV	≤ 40%	No effect

セロトニンは血小板の濃染顆粒に存在する生理活性アミンで、血管収縮作用をはじめ繊維芽細胞増殖作用、マクロファージの活性酸素産生促進作用、NK細胞活性化促進作用を有する¹¹⁾。CD62Pはセレクチンファミリーに属する接着分子で¹²⁾、血小板

活性化に伴って α 顆粒から膜表面に分布するため、血小板活性化マーカーとして汎用されている¹¹⁾。PAC-1は血小板上の α IIb β 3 (GPIIb-IIIa)が血小板活性化にともなって構造変化を起こした時に発現するエピトープを認識するモノクローナル抗体である^{11,12)}。低濃度のADP刺激によっても発現の増加がみられることから、高感度でかつ特異的な血小板活性化マーカーとして有用であるとされている。 α IIb β 3はインテグリンファミリーに属する接着分子であり、血小板においてフィブリノーゲンやvon Willbrand因子のレセプターとして機能し、血小板凝集や粘着の重要な分子である¹³⁾。

ヒト多血小板血漿に20%および40%の濃度のHbVを1時間予めインキュベーション後、アゴニスト刺激の無い場合は、血小板反応上清中のRANTES, セロトニン, β -TG, TXB₂および血小板表面CD62Pのレベルの上昇はみられないことから、HbVに血小板活性化作用は無いと考えられた。この際、RANTESレベルはむしろ減少したが、この程度の変化での臨床的意義は少ないと推察された。アゴニスト刺激により血小板の脱顆粒を惹起させると、いずれの濃度のHbV処理においてもRANTES, セロトニン, β -TGの放出, TXB₂の産生, およびCD62P発現が引き起こされ、そのレベルはHbV未処理のものと同程度はみられなかった (Table 1)。一方、PAC-1の発現はHbV未処理では影響がないものの、HbV処理により低濃度のADP刺激血小板において僅かな亢進がみられた。PAC-1の発現の亢進は空の小胞体処理においても観察されたため、ヘモグロビンの有無にはよらず、小胞体それ自体または小胞体の構成成分に起因する可能性が考えられた。以上のことから、HbVは低濃度のアゴニスト存在下で α IIb β 3の構造変化を僅かに促進するものの、CD62Pの膜への発現, RANTES, セロトニン, β -TGの放出, TXB₂の産生にはアゴニスト刺激の有無にかかわらず影響を与えないことから、ヒト血小板に対し高い生体適合性があると結論された (Table 1)。

2. ヘモグロビン小胞体の好中球活性化に対する影響

HbVがヒトに投与される状況として大量出血が想定される。その際に、生体防御のファーストラインを担う好中球の機能に対し、HbVが抑制的に働けば、感染症などを合併する危険性が懸念される。循環血中の好中球は活性化を受け、炎症部位に遊走し、生体を脅かす細菌を効率的に殺菌する必要がある。好中球は、ゼイモザンや細菌由来のペプチド (*N*-formyl-methionyl-leucyl-phenylalanine (fMLP)) によって活性化を受け、多岐にわたる機能を発揮することが知られているが、PEG-distearoylphosphatidylethanolamine (PEG-DSPE) 修飾を受けたある種のリボソームは、これらの刺激に対する好中球の反応を阻害することが報告されている¹⁴⁾。逆に、フォスファチジルコリンやフォスファチジルセリンからなるリボソームが、マウスのアレルギーモデルにおいて、好中球の肺への集積を促進させたという報告がなされている¹⁵⁾。そこで、HbV自体がPEG-DSPE修飾をうけていることもあり、fMLP刺激によって好中球が発揮する機能に対し、どのような影響を及ぼすかに

Table 2. Effect of DPPP-HbV on fMLP-induced neutrophil functions.

Function	Conc. of HbV	Effect
Chemotaxis	≤ 0.6%	No effect
Upregulation of CD11b expression	≤ 0.6%	No effect
Degranulation of Gelatinase B	≤ 6%	No effect
Superoxide production	≤ 6%	No effect

ついて検討を加えた¹⁶⁾。

この検討においては、リボソームの構成成分としてDPPPを含有するHbVを用いた。fMLP刺激による好中球の機能評価として、(1)好中球の走化能、(2) β ₂インテグリンであるCD11bの膜発現、(3)脱顆粒にともなうgelatinase B (matrix metalloproteinase-9)の放出、(4)活性酸素の産生をとりあげた。Table 2にこれらの結果を示した。

走化性(ケモタキシス)は、細胞が化学物質の濃度勾配に従い、一定の方向に運動性を示すことであり、ヒト単離好中球を異なる濃度(最大0.6%)のHbVと37℃にて30分間予めインキュベーション(プレインキュベーション)後、ケモタキシス・チャンパー法によって評価をおこなった。fMLP (1 μ M)刺激によって好中球の走化能が活性化されたが、これに対するHbVの影響はみとめられなかった。CD11bはCD18と二量体を形成して好中球の遊走能における接着因子Mac-1として機能する分子であるが、HbVとプレインキュベーションさせた好中球において、fMLP刺激によるCD11bの発現亢進は、HbVを作用させなかった好中球の場合と同等であった。gelatinase Bは、好中球の組織浸潤に関与する酵素であるが、fMLP刺激による好中球からの放出の経時変化をHbVとのプレインキュベーションの有無で比較したが、両者には違いはみとめられなかった。また好中球の殺菌能に関与する活性酸素産生についても、HbVとプレインキュベーションした好中球からのfMLPによる産生量は、HbVを作用させなかった好中球の場合と同等であった。以上の結果より、fMLP刺激によって発揮される好中球の少なくとも4つの機能において、用いたHbVの濃度ではなら影響はみられず、HbVの好中球への高い適合性が示唆された。

PEG-DSPE修飾したリボソームがfMLP刺激による好中球の走化能活性化を阻害するという報告¹⁷⁾を上述したが、リン脂質濃度にして大過剰の濃度を用いたにもかかわらず、HbVでは阻害がみられないことの原因は不明である。リボソームの構成成分であるリン脂質の種類、脂肪酸の飽和度の程度、脂肪酸の側鎖の数が、fMLPやゼイモザンによる好中球の走化能に影響することが報告されている¹⁸⁾。前述のPEG-DSPE修飾したリボソームとHbVとの間では、これらのパラメーターを始め、PEG-DSPEとリボソームのモル比が異なっており、これらの違いが異なる結果を得た理由となっている可能性が考えられる。

以上、HbVの好中球への高い適合性が示されたが、より高濃度即ち臨床で用いられる20%および40%での検討をみる必要性は残されている。

3. ヘモグロビン小胞体の造血前駆細胞に対する影響

血管内投与したりボソーム製剤は、漸次、肝臓のクッパー細胞、脾臓および骨髄のマクロファージなどの細網内皮系において捕捉される²¹⁾。HbVにおいても、アイソトープラベルを用いた追跡によって組織分布が詳細に検討され、肝臓、脾臓、骨髄に主として分布していることが示されている²²⁾。他の組織学的検討においても、HbVは細網内皮系において捕捉されることが報告されている²³⁾。骨髄にHbVが移行することから、骨髄を主たる場とする造血に対し少なからず影響するのではないかと考えられた。40%脱血しHbVを輸血したラットさらにビーグル犬の場合、脱血による赤血球造血の亢進によってヘマトクリットは1週間以内には回復し、特に造血系への影響は認められていない²⁴⁾。さらに、20%のHbVをtoploadさせたラットの場合においても、末梢血の血球数は投与後1週間観察しても一定に保たれていた²⁵⁾。とはいえヒトへの投与を考えた場合、その造血前駆細胞の増殖・分化への影響について、in vitroの系において検討する必要があると考えられ検討をおこなった²¹⁾。

造血のin vitroのアッセイ系としては、コロニー形成細胞の数を評価するコロニーアッセイと造血前駆細胞の増殖の度合いを評価する液体培養系を用いた。そして臍帯血から分離したCD34陽性細胞を含む単核球を、造血を支持する各種サイトカインを添加したメチルセルロースからなる半固形培地に播種し、一定期間培養後に形成したコロニーの数をカウントした。赤芽球バーストとよばれるBFU-E (Burst forming unit of erythrocyte)、顆粒球・マクロファージ系コロニーとよばれるCFU-GM (Colony forming unit of granulocyte and macrophage)、両者が混在したCFU-mix。これらのコロニーの総数であるColony-forming cells (CFC)は培養2週間後に、増殖能が高くコロニーのサイズが1 mm以上になったものをHPP-CFC (High proliferative potential colony forming cell)と定義し、培養4週間後にそれぞれカウントした。用いたHbVはPEG修飾されたDHSGを構成成分とするHbVで、その濃度は最大3%と設定した²¹⁾。

コロニーアッセイの培地にHbVを添加し、14日間共存させると、それぞれのコロニーの数およびサイズがHbVの濃度に依存して減少する傾向がみられた。さらに14日間HbVと共存後にコロニー数を測定するHPP-CFCの数は著しく減少した (Fig. 1A)。一方、HbVにCD34陽性細胞を含む単核球を20時間だけ曝してコロニーアッセイした場合は、HPP-CFCの数に影響はなく、3日間曝した場合でも、HPP-CFCの数の減少の程度は少なかった (Fig. 1B)。次に液体培地に血清と造血支持サイトカインを添加して、赤芽球系または顆粒球系の細胞へと増殖させる系にHbVを添加させ影響を評価した。10日間の培養期間、HbVが継続して存在する場合には赤芽球系または顆粒球系への細胞の増殖が抑制された。これに対し、コロニーア




Fig. 1. Effects of HbVs on the formation of high proliferative potential colony forming cells (HPP-CFC) in the clonogenic assay of cord blood-derived hematopoietic cells. Cord blood-derived mononuclear cells were seeded at 300 at CD34⁺ cells per dish in complete methylcellulose medium for human clonal culture assays. HPP-CFC was scored after 28 days incubation. (A) HbVs (0 to 3%) were added to the cells until the end of culture. (B) Cord blood-derived mononuclear cells were incubated for 20 h (open column) or for 3 days (shaded column) with different concentrations of HbVs in IMDM containing FCS, BSA, and cytokines. Subsequently, they were recovered, washed to remove the HbVs, and subjected to the clonal assay. Data are expressed as the mean \pm SD of the percentage of control (HbV 0%) performed on three separate cord blood donors in (A) and (B).

ッセイの場合と同様に、HbVに20時間または3日間曝しただけの場合は、赤芽球系または顆粒球系への細胞の増殖は影響されなかった (Table 3)。

以上HbVは、in vitroの閉鎖系において、長期間にわたり共存することにより造血前駆細胞の活性を阻害するという結果が、2つの評価系から明らかとなった。しかしながら、動物モデルでHbVを投与した場合、骨髄において1-3日にHbVの蓄積が認められ、以後1週間では激減することが報告されている²⁴⁾。

Table 3. Effect of DHSG-HbV on the proliferation of erythroid and myeloid lineage cells in liquid culture.

Exposure period to HbV	CD235a ⁺ cells			CD15 ⁺ cells		
	HbV conc.(%)			HbV conc.(%)		
	0.75	1.5	3.0	0.75	1.5	3.0
20h	93.7 \pm 10.0	94.9 \pm 1.2	92.2 \pm 8.8	100.8 \pm 14.3	96.3 \pm 7.9	96.6 \pm 13.3
3 days	85.2 \pm 22.3	92.6 \pm 11.5	89.0 \pm 14.5	92.9 \pm 6.1	95.8 \pm 5.4	91.7 \pm 4.7
10 days	47.5 \pm 16.6**	46.6 \pm 18.2**	27.3 \pm 23.4**	85.2 \pm 20.3*	55.2 \pm 18.6*	37.2 \pm 22.8*

Various concentrations of HbVs were added to the medium containing the cord blood-derived CD34⁺ cells. After 10 days' incubation, CD235a⁺ cells for erythroid lineage and CD15⁺ cells for myeloid lineage, respectively, were analyzed by flow cytometry. The number of CD235a⁺ cells or CD15⁺ cells at each concentration of DHSG-HbV is expressed as a percentage of the number in the control (HbV 0%). Data are represented as the mean \pm SD from three experiments performed on three separate cord blood donors. *p < 0.05, **p < 0.01 versus HbV (0%).

よって、2週間連続してHbVに造血幹・前駆細胞が曝されるという状況はヒトの臨床例では想定できないし、むしろ1-3日間曝されるというのが想定できる期間である。さらにin vivoでは、HbVはマクロファージ内に蓄積しており、in vitroのようにHbVと造血幹・前駆細胞が直接相互作用をすることは極めて少ないと考えられる。従って造血幹・前駆細胞が直接1-3日間HbVに曝される場合において、造血前駆細胞の活性に大きな影響がなかったという結果は、HbVをヒトに投与した場合、造血に大きな悪影響を及ぼす可能性が極めて少ないことを支持するものと捉えることができる。

国外では、ヘモグロビンを基盤として化学修飾させたセルフリ一型の人工酸素運搬体の開発が進められている。これらの修飾ヘモグロビンや分子内架橋を行ったヘモグロビンは、我々と同じようにヒト臍帯血から調製したCD34陽性細胞を用いた液体培養系において、赤芽球前駆細胞の増殖の亢進を促すと報告されている²⁹。また、AIDSの治療に用いられる薬剤3'-azido-3'-deoxythymidine (AZT)による造血抑制を、ヘモグロビンおよび化学修飾ヘモグロビンがin vitro, in vivoの両方で回復させる(但し赤血球造血に特異的)ことが示され^{29,30}。これらの赤血球造血作用は酸素運搬能以外の付加価値と位置付けられている。修飾ヘモグロビンのこれらのin vitroでの作用は、赤芽球系前駆細胞が修飾ヘモグロビンを取り込み、赤血球造血作用のあるhemeに分解するためと推測されている²⁹。HbVの場合は、赤芽球系前駆細胞を含めて造血前駆細胞の増殖亢進作用は現時点で明らかでない。少なくとも赤芽球系前駆細胞は、HbVを修飾ヘモグロビンのように容易には取り込まないと考えられる。

4. ヘモグロビン小胞体の補体系、凝固系、カリクレイン-キニン系に対する影響

リポソームは細胞膜と同様リン脂質を構成成分としているが、生体にとっては異物であり、血液細胞のみならず血漿タンパク質と相互作用する³¹。この作用は、リポソームの大きさ、荷電、構成脂質、さらにリポソーム自体への化学修飾により異なってくる。リポソームの中で陰性荷電のものは、ラット、モルモット、そしてヒトの補体を活性化することが広く知られている³²⁻³⁴。その結果、オプソニン化したリポソームは、細網内皮系により循環血液中から速やかに排除される。また陰性荷電リポソームは凝固系第7因子を活性化させ、内因系の凝固カスケードならびにカリクレイン・キニン系活性のトリガーとなる^{35,36}。リポソームの構成成分のコレステロール含量も補体の活性化に影響することが知られ、この活性化は血液中の自然抗体によると考えられている^{31,37,38}。これまで述べてきたリポソーム包埋型ヘモグロビンにおいても、ヒトの血液中の脂質に対する自然抗体によって、古典経路と第二経路の両方を介して補体の活性化が惹起される³⁹。さらに、ブタモデル実験においては、ある種のリポソームによって偽アレルギー反応が起こり、補体活性化との関連が考えられている^{40,41}。一方、リポソーム

をPEG修飾することで、これらの反応を防ぐことができることが報告されている^{19,42}。しかしながら、リポソーム包埋型ヘモグロビンのみならず抗ガン剤doxorubicinを封入したPEG修飾リポソームDoxil/Caelyxにおいても、顔面紅潮、呼吸困難、紅斑、胸痛、背面痛、血圧低下または血圧上昇が報告されている^{43,44}。またこれらの症状はdoxorubicin自体では起こらないため、リポソームによるものと考えられている⁴⁵。したがってHbVの生体適合性を評価するうえで、ヒトの補体系、凝固系、カリクレイン-キニン系に対する影響を検討することの意義は大きいと考えられた。

この検討においては、DHSGをそのリポソーム構成成分としPEG修飾された現行のHbV (DHSG-HbV)、DPPGを構成成分としPEG修飾されたHbV (DPPG-HbV)、DPPGを構成成分とするがPEG未修飾のHbV (DPPG-HbV (no PEG))、強陰性荷電PEG未修飾リポソーム (EL-A)を用いた³⁹。なお、EL-Aの脂質構成はDPPC:CHOL:DPPG (30:40:30) (mol%)からなり、その生理食塩水中でのゼータ電位は-47.2mVであった³⁹。

HbVを健康人ヒト血清と20%あるいは40%で混合し、37℃にて24時間インキュベーション後、遠心により各種HbVを除去し、その上清の補体価(CH50)を測定した。Table 4に示すように、DHSG-HbVを混合することによって血清が希釈されるためCH50は低下した。強陰性荷電PEG未修飾リポソーム(EL-A)を添加した群では著しいCH50の低下がみられるのに対し、DHSG-HbVを20%あるいは40%で添加した群では、それぞれ生食を添加した場合のCH50より減少することはない。さらに3種のHbVと比較してみると(Table 5)、DHSG-HbVとDPPG-HbVは、補体消費を起こさなかったのに対し、DPPG-HbV (no PEG)は40%の添加によってほぼ完全に補体が消費された。以上の結果から、現行のDHSG-HbVおよびPEG修飾されたDPPG-HbVは、ヒト血清補体系を活性化しないと考えられた。

Table 4. Consumption of complement by HbV and Liposome.

Additive	CH50 (U/mL)	
	(additive:serum)	
	20:80	40:60
Saline	33.4 ± 2.8	21.4 ± 1.7
DHSG-HbV	33.5 ± 2.9	22.9 ± 2.4
EL-A	25.1 ± 2.7*	5.9 ± 0.7*

The complement titer (CH50) was measured using a 50% hemolysis assay with a commercial kit. DHSG-HbV, saline or Coatsome EL-A (a negative charged liposome) were mixed with serum as indicated ratio (v/v) at 37°C for 24 hr. The lipid composition (mol%) of Coatsome EL-A was DPPC:CHOL:DPPG=30:40:30. Data are represented as the mean ± SD using serum from five individuals. The CH50 of 100% serum was 38 ± 3.2 U/mL. *p < 0.05 vs. saline.

Table 5. Consumption of complement by various types of HbV.

Additive	CH50 (U/mL)	
	(additive:serum)	
	20:80	40:60
Saline	36.4	27.9
DHSG-HbV	37.6	31.4
DPPG-HbV	35.9	28.4
DPPG-HbV(no PEG)	29.9	Under detection limit

The complement titer (CH50) was measured using a 50% hemolysis assay with a commercial kit. DHSG-HbV, DPPG-HbV, DPPG-HbV (no PEGylation) or saline were mixed with serum as indicated ratio (V/V) at 37°C for 24 hr. The CH50 of 100% serum was 45.1 U/mL.

次に凝固系に及ぼす影響をこれら3種のリボソーム構成成分の異なるHbVを用い、プロトロンビン時間(PT)および活性化部分トロンボプラスチン時間(APTT)について比較した。PTは健康人ヒト血漿との混合比率が上昇するにつれて凝固時間が遅延した。血漿に対し生食の比が20%および60%においては、これら3種のHbVはいずれも生食添加の場合のPT時間を有意に短縮させたが、その差は1秒以内であった。APTT時間に対し、DPPG-HbVとDPPG-HbV(no PEG)は、生食添加の場合のAPTT時間を2秒以内ではあるが有意に短縮させたのに対し、DHSG-HbVはどの血漿比率の場合においても生食の場合と差はみられなかった。

カリクレイン-キニン系に対する影響は、血漿にそれぞれこれら3種のHbVまたは生食を添加し、37°Cにて24時間インキュベーション後、血漿中高分子キニノーゲンの分解による低分子化された高分子キニノーゲンの出現をイムノブロットング法にて検出した。血漿に対し40%あるいは60%にてDPPG-HbVまたはDPPG-HbV(no PEG)を添加させた場合、intactな高分子キニノーゲンが減少し、低分子化された高分子キニノーゲンの増加が顕著に観察される。これに対して、DHSG-HbVではいずれの血漿比率の場合においても生食の場合と差はみられず、DHSG-HbVは血漿中高分子キニノーゲンの分解を引き起こさないことが示された(Fig. 2)。

以上の検討から、陰性荷電のDPPGを構成成分とするHbVではPEG修飾がなければ補体の活性化が惹起されるのに対し、PEG修飾することで補体の活性化が抑えられることが明らかとなった。生理食塩水中でゼータ電位を測定するとDPPG-HbV(no PEG)が-14.5 mV、DPPG-HbVが-3.4 mVであり³⁰⁾、表面荷電の影響を裏付けていた。しかしながら、ここに示したごとくDPPG-HbVでは、わずかではあるが、凝固時間、特にAPTT時間の短縮およびカリクレイン-キニン系の活性化を引き起こした。これに対し、DHSG-HbVは、補体系、APTT時間、カリクレイン-キニン系のいずれに対しても影響が無いことが示された。DHSG-HbVのゼータ電位は-2.6 mVでDPPG-




Fig. 2. Activation of kallikrein-kinin cascade by HbVs. HbVs or saline were mixed with plasma at 40% or 60% at 37°C for 24 hr. Appearance of digested S-HMWK was detected using western blot analysis as a result of kallikrein activation. Arrowhead (▶) indicates digested S-HMWK.

HbVのそれとほとんど変わらないことから、単に表面荷電だけでなく、構成脂質の親水性頭部(hydrophilic head group)が作用の有無に影響を及ぼすものと考えられる。

いずれにしても、今後の臨床試験に用いられる現行のHbVは、血漿中の補体系に影響を及ぼさないということから、“補体活性化関連偽アレルギー反応”と呼ばれるリボソーム投与直後の心血管系の急性副作用³¹⁾に対する懸念はほとんど無いと推測される。加えて、凝固系やカリクレイン-キニン系への影響も無いことは、ヒト血漿タンパクへの高い適合性を示している。

第1章の血小板に対するHbVの影響に関して、げっ歯類においては血小板上に補体成分C3bが結合するCRIレセプターを有しているが、ヒトを含む霊長類では有さないことを上述した。これに対し、ヒトを含む霊長類では赤血球がCRIレセプターを有しているため、補体活性化に伴って免疫複合体の赤血球への結合が報告されている³²⁾。そのため、補体活性化を惹起する化合物および医薬品のヒト血液を対象としたin vitroにおける適合試験や臨床試験においては、赤血球も相互作用する可能性のある血液構成成分の一つとして重要であり、そのため、赤血球をモニターする意義や結果の解釈にも影響する可能性を考慮しておく必要がある。しかしながら、少なくとも今後の臨床試験に用いられる現行のHbVは、in vitroの評価試験において血漿中の補体系に影響を及ぼさないため、補体成分を介してHbVが赤血球と結合する可能性は少ないと推測される。

おわりに

HbVが臨床応用に向けて、前臨床試験でその有効性、安全性について検討しなければならない事項の中に、免疫系および血液構成成分に対する影響があげられている。ヒト血液細胞およびヒト血漿タンパク系を用いた評価により、止血また炎症反

応にも関与する血小板に対し、HbV自体による活性化作用はみられないこと、またアゴニストによる活性化を亢進するような作用はないことが示された。自然免疫に重要な働きをする好中球の機能（走化能、脱顆粒、活性酸素産生）にも影響を与えなかった。造血機能への影響としては、短期間であれば造血前駆細胞活性を損なうことはないことが示された。またリポソームの特性から補体の活性化が懸念されたが、現行のHbVでは活性化はみられず、また、凝固系、カリクレイン・キニン系にも影響はみられなかった。しかしながら *in vitro* では評価できない抗体産生、投与後の補体価の変動、リンパ球の抗原刺激による増殖反応等については現在ラットを用いて検討をおこなっている。本稿にては詳細は割愛したが、それらの影響は、投与直後に一過性の抑制がみられるものの1週間では回復することをみだしている^{28,30}。以上のことから、免疫系、および血液構成成分に対してHbVが高い適合性を持つことを示すことができた。

最近、Natansonら³¹は、ヘモグロビンを基盤とする非セル型の人工酸素運搬体を対象とした無作為化された前臨床試験および臨床試験の中から、一定の基準を満たした16トリアルにおいて、死亡のリスクが30%増、心筋梗塞のリスクが2.7倍増加しているというメタ解析の結果を報告した。この報告に伴って、ヘモグロビンを基盤とする非セル型の人工酸素運搬体の安全性を見直し、リスクの増加の原因を明らかにすべきとの指摘がなされている³²。このような状況下において、我が国だけが開発を進めているセル型の人工酸素運搬体の一つであるHbVに、大きな期待がよせられる。

謝辞

本研究は、厚生科学研究費補助金（高度先端医療研究事業）「酸素運搬能を有する人工赤血球の創製とその評価に関する研究（平成9年～11年）」「臨床応用可能な人工赤血球の創製に関する研究（平成12年～14年）」、厚生科学研究費補助金（高度先端医療研究事業）「人工赤血球の安全性向上に関する研究（平成15年～17年）」、「人工酸素運搬体の臨床応用に関する研究（平成18年～20年）」によって行われた。記して謝意を表する。

参考文献

1. Sakai H, Sou K, Horinouchi H, Kobayashi K, Tsuchida E. Haemoglobin-vesicles as artificial oxygen carriers: present situation and future visions. *J Intern Med* 2008;263:4-15.
2. 高折益彦: 人工血液としての条件. *人工血液* 2002;10:28-35.
3. Rabinovici R, Rudolph AS, Feuerstein G: Characterization of hemodynamic, hematologic, and biochemical responses to administration of liposome-encapsulated hemoglobin in the conscious, freely moving rat. *Circ Shock* 1989;29:115-32.
4. Rabinovici R, Rudolph AS, Yue T, Feuerstein G: Biological responses to administration of liposome-encapsulated hemoglobin (LEH) are improved by a PAF antagonist.

- Circ Shock* 1990;31:431-5.
5. Phillips WT, Klipper R, Fresne D, Rudolph AS, Javors M, Goins B: Platelet reactivity with liposome-encapsulated hemoglobin in the rat. *Exp Hematol* 1997;25:1347-56.
6. Reinish LW, Bally MB, Loughrey HC, Cullis PR: Interactions of liposomes and platelets. *Thromb Haemost* 1988;60:518-523.
7. Doerschuk CM, Gie RP, Bally MB, Cullis PR, Reinish L: Platelet distribution in rabbits following infusion of liposomes. *Thromb Haemost* 1989;61:392-6.
8. Loughrey HC, Bally MB, Reinish LW, Cullis PR: The binding of phosphatidylglycerol (PG) liposomes to rat platelets is mediated by complement. *Thromb Haemost* 1990;64:172-6.
9. Klinger MHF: Platelets and inflammation. *Anat Embryol* 1997;196:1-11.
10. Gawaz M, Langer H, May AE: Platelets in inflammation and atherogenesis. *J Clin Invest* 2005;115:3378-84.
11. Wakamoto S, Fujihara M, Abe H, Sakai H, Takeoka S, Tsuchida E, Ikeda H, Ikebuchi K: Effects of poly(ethyleneglycol)-modified hemoglobin vesicles on agonist-induced platelet aggregation and RANTES release *in vitro*. *Artif Cells Blood Substit Immobil Biotechnol* 2001;29:191-201.
12. Baggolini M, Dahinden CA: CC chemokines in allergic inflammation. *Immunol Today* 1994;15:127-33.
13. Wakamoto S, Fujihara M, Abe H, Yamaguchi M, Azuma H, Ikeda H, Takeoka S, Tsuchida E: Effects of hemoglobin vesicles on resting and agonist-stimulated human platelets *in vitro*. *Artif Cells Blood Substit Biotechnol* 2005;33:101-11.
14. Hagberg IA, Lyberg T: Evaluation of circulating platelet-leukocyte conjugates: a sensitive flow cytometric assay well suited for clinical studies. *Platelets* 2000;11:151-60.
15. 幸村 近, 林 由紀子, 池田久實: フローサイトメトリーを用いた活性化血小板の検出法. *臨床病理* 1999;47:447-52.
16. Hynes RO: Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 1992; 69:11-25.
17. Hatipoglu U, Gao XP, Verral S, Sejourne F, Pitrak D, Alkan-Onyuksel H, Rubinstein I: Sterically stabilized phospholipids attenuate human neutrophils chemotaxis *in vitro*. *Life Sci* 1998; 63:693-9.
18. Bellemare F, Israel-Assayag E, Cormier Y: PC:PS liposomes induce a recruitment of neutrophils and the release of TNF alpha in the lungs of mice sensitized with *Saccaropolyspora rectivirgula*. *Eur J Clin Invest* 1995;25:340-5.
19. Ito T, Fujihara M, Abe H, Yamaguchi M, Wakamoto S, Takeoka S, Sakai H, Tsuchida E, Ikeda H, Ikebuchi K: Effects of poly(ethyleneglycol)-modified hemoglobin

- vesicles on N-formyl-methionyl-leucyl-phenylalanine-induced responses of polymorphonuclear neutrophils in vitro. *Artif Cells Blood Substit Immobil Biotechnol* 2001;29:427-37.
20. Galdiero F, Carratelli CR, Bentivoglio C, Capasso C, Cioffi S, Folgore A, Gorga F, Ianniello R, Mattera S, Nuzzo I, Rizzo A, Tufano MA: Correlation between modification of membrane phospholipids and some biological activity of lymphocytes, neutrophils and macrophages. *Immunopharmacol Immunotoxicol* 1991;13:623-42.
 21. Torchilin VP: Recent advances with liposomes as pharmaceutical carriers. *Nat Rev Drug Discov* 2005;4:145-60.
 22. Sou K, Klipper R, Goins B, Tsuchida E, Phillips WT: Circulation kinetics and organ distribution of Hb-vesicles developed as a red blood cell substitute. *J Pharmacol Exp Ther* 2005;312:702-9.
 23. Sakai H, Horinouchi H, Tomiyama K, Ikeda E, Takeoka S, Kobayashi K, Tsuchida E: Hemoglobin-vesicles as oxygen carriers: Influence on phagocytic and histopathological changes in reticuloendothelial system. *Am J Pathol* 2001;159:1079-88.
 24. Sakai H, Horinouchi H, Yamamoto M, Ikeda E, Takeoka S, Takaori M, Tsuchida E, Kobayashi K: Acute 40 percent exchange-transfusion with hemoglobin-vesicles (HbV) suspended in recombinant human serum albumin solution: degradation of HbV and erythropoiesis in a rat spleen for 2 weeks. *Transfusion* 2006;46:339-47.
 25. Ikeda T, Horinouchi H, Kohno M, Izumi Y, Watanabe M, Sakai H, Sou K, Tsuchida E, Kobayashi K: Resuscitation effect and long term effect of Hb vesicles on organ function in beagle dog. *Artif Cells Blood Substitutes Biotechnol* 2008;36:214.
 26. Abe H, Azuma H, Yamaguchi M, Fujihara M, Ikeda H, Sakai H, Takeoka S, Tsuchida E: Effects of hemoglobin vesicles, a liposomal artificial oxygen carrier, on hematological responses, complement and anaphylactic reactions in rats. *Artif Cells Blood Substit Biotechnol* 2007;35:157-72.
 27. Yamaguchi M, Fujihara M, Wakamoto S, Sakai H, Takeoka S, Tsuchida E, Azuma H, Ikeda H: Influence of hemoglobin vesicles, cellular-type artificial oxygen carriers, on human umbilical cord blood hematopoietic progenitor cells in vitro. *J Biomed Mater Res A* 2008. (in press)
 28. Bell D, Mueller SG: Enhanced stimulation of erythropoiesis. Patent EP099644, 2004.
 29. Fowler DA, Rosenthal GJ, Sommadossi JP: Effect of recombinant human hemoglobin on human bone marrow progenitor cells: protection and reversal of 3'-azido-3'-deoxythymidine-induced toxicity. *Toxicology Letters* 1996;85:55-62.
 30. Moqattash S, Lutton JD, Rosenthal G, Abu-Hijleh MF, Abraham NG: Effect of blood substitute, recombinant hematology, on in vivo hematopoietic recovery from AZT toxicity. *Acta Haematol* 1997;98:76-82.
 31. 阿部秀樹, 藤原満博, 東 寛, 池田久實. リポソームと補体. *人工血液* 2003;11:151-9.
 32. Chonn A, Cullis PR, Devine DV: The role of surface charge in the activation of the classical and alternative pathways of complement by liposomes. *J Immunol* 1991;146:4234-41.
 33. Cunningham CM, Kingzette M, Richards RL, Alving CR, Lint TF, Gewurz H: Activation of human complement by liposomes: a model for membrane activation of the alternative pathway. *J Immunol* 1979;122:1237-42.
 34. Devine DV, Wong K, Serrano K, Chonn A, Cullis PR: Liposome-complement interactions in rat serum: implications for liposome survival studies. *Biochim Biophys Acta* 1994;1191:43-51.
 35. Griep MA, Fujikawa K, Nelsestuen GL: Binding and activation properties of human factor XII, prekallikrein, and derived peptides with acidic lipid vesicles. *Biochemistry* 1985;24:4124-30.
 36. Mitropoulos KA, Martin JC, Reeves BE, Esnouf MP: The activation of the contact phase of coagulation by physiologic surfaces in plasma: the effect of large negatively charged liposomal vesicles. *Blood* 1989;73:1525-33.
 37. Alving CR, Richards RL, Guirguis AA: Cholesterol-dependent human complement activation resulting in damage to liposomal model membranes. *J Immunol* 1977;118:342-7.
 38. Alving CR: Natural antibodies against phospholipids and liposomes in humans. *Biochem Soc Trans* 1984;12:342-4.
 39. Szebeni J, Wassef NM, Rudolph AS, Alving CR: Complement activation in human serum by liposome-encapsulated hemoglobin: the role of natural anti-phospholipid antibodies. *Biochim Biophys Acta* 1996;1285:127-30.
 40. Bradley AJ, Devine DV, Ansell SM, Janzen J, Brooks DE: Inhibition of liposome-induced complement activation by incorporated poly (ethylene glycol)-lipids. *Arch Biochem Biophys* 1998;357:185-94.
 41. Klibanov AL, Maruyama K, Torchilin VP, Huang L: Amphipathic polyethyleneglycols effectively prolong the circulation time of liposomes. *FEBS Lett* 1990;268:235-7.
 42. Woodle MC, Matthey KK, Newman MS, Hidayat JE,

- Collins LR, Redemann C, Martin FJ, Papahadjopoulos D: Versatility in lipid compositions showing prolonged circulation with sterically stabilized liposomes. *Biochim Biophys Acta* 1992;1105:193-200.
43. Laverman P, Boerman OC, Oyen WJG, Corstens FHM, Storm G: In vivo applications of PEG liposomes: unexpected observations. *Crit Rev Ther Drug Carrier Syst* 2001;18:551-66.
44. Szebeni J, Baranyi L, Savay S, Bodo M, Morse DS, Basta M, Stahl GL, Bünger R, Alving CR. Liposome-induced pulmonary hypertension: properties and mechanism of a complement-mediated pseudoallergic reaction. *Am J Physiol Heart Circ Physiol* 2000;279:H1319-28.
45. Wassef NM, Johnson SH, Graeber GM, Swartz GM Jr, Schultz CL, Hailey JR, Johnson AJ, Taylor DG, Ridgway RL, Alving CR: Anaphylactoid reactions mediated by autoantibodies to cholesterol in miniature pigs. *J Immunol* 1989;143:2990-5.
46. Uziely B, Jeffers S, Isacson R, Kutsch K, Wei-Tsao D, Yehoshua Z, Libson E, Muggia FM, Gabizon A. Liposomal doxorubicin: antitumor activity and unique toxicities during two complementary phase I studies. *J Clin Oncol* 1995;13:1777-85.
47. Koukourakis MI, Koukouraki S, Giatromanolaki A, Archimandritis SC, Skarlatos J, Beroukas K, Bizakis JG, Retalis G, Karkavitsas N, Helidonis ES. Liposomal doxorubicin and conventionally fractionated radiotherapy in the treatment of locally advanced non-small-cell lung cancer and head and neck cancer. *J Clin Oncol* 1999;17:3512-21.
48. Lyass O, Uziely B, Ben-Yosef R, Tzemach D, Heshing NI, Lotem M, Brufman G, Gabizon A. Correlation of toxicity with pharmacokinetics of pegylated liposomal doxorubicin (Doxil) in metastatic breast carcinoma. *Cancer* 2000;89:1037-47.
49. Chanan-Khan A, Szebeni J, Savay S, Liebes L, Rafique NM, Alving CR, Muggia FM. Complement activation following first exposure to pegylated liposomal doxorubicin (Doxil): possible role in hypersensitivity reactions. *Ann Oncology* 2003; 14:1430-7.
50. Abe H, Fujihara M, Azuma H, Ikeda H, Ikebuchi K, Takeoka S, Tsuchida E, Harashima H: Interaction of hemoglobin vesicles, a cellular-type artificial oxygen carrier, with human plasma: effects on coagulation, kallikrein-kinin, and complement systems. *Artif Cells Blood Substit Biotechnol* 2006;34:1-10.
51. Szebeni J, Fontana JL, Wassef NM, Mongan PD, Morse DS, Dobbins DE, Stahl GL, Bunge R, Alving CR. Hemodynamic changes induced by liposomes and liposome-encapsulated hemoglobin in pigs. A model for pseudoallergic cardiopulmonary reactions to liposomes: role of complement and inhibition by soluble CRI and anti-C5a antibody. *Circulation* 1999;99:2302-2309.
52. Cornacoff JB, Hebert LA, Smead WL, VanAman ME, Birmingham DJ, Waxman FJ: Primate erythrocyte-immune complex-clearing mechanism. *J Clin Invest* 1983;71:236-47.
53. 阿部秀樹, 山口美樹, 藤原満博, 酒井宏水, 武岡真司, 土田英俊, 東 寛, 池田久實: ヘモグロビン小胞体 (HbV) 投与がラット免疫系に及ぼす影響. *人工血液* 2004;12:55.
54. Natanson C, Kern SJ, Lurie P, Banks SM, Wolfe AM: Cell-free hemoglobin-based blood substitutes and risk of myocardial infarction and death: a meta-analysis. *JAMA* 2008; 299:304-12.
55. Furgusson DA, McIntyre L: The future of clinical trials evaluating blood substitutes. *JAMA* 2008;99:2324-6.



Contents lists available at ScienceDirect

Journal of Controlled Release

journal homepage: www.elsevier.com/locate/jconrel



Pharmacokinetics of single and repeated injection of hemoglobin-vesicles in hemorrhagic shock rat model

Kazuaki Taguchi^a, Toru Maruyama^{a,b}, Yasunori Iwao^a, Hiromi Sakai^c, Koichi Kobayashi^d, Hirohisa Horinouchi^d, Eishun Tsuchida^c, Toshiya Kai^{a,e}, Masaki Otagiri^{a,*}

^a Department of Biopharmaceutics, Graduate School of Pharmaceutical Sciences, Kumamoto University, 5-1 Oe-honmachi, Kumamoto 862-0973, Japan

^b Center for Clinical Pharmaceutical Sciences, Kumamoto University, 5-1 Oe-honmachi, Kumamoto 862-0973, Japan

^c Research Institute for Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku, Tokyo 169-8555, Japan

^d Department of Surgery, School of Medicine, Keio University, 35 Shinano Shinjyuku, Tokyo 160-8582, Japan

^e The Pharmaceutical Research Center, Nipro Corporation 3023 Najicho, Kusatsu, Shiga 525-0055, Japan

ARTICLE INFO

Article history:

Received 10 November 2008

Accepted 10 February 2009

Available online xxxxx

Keywords:

Liposome

Oxygen carrier

Hemorrhagic shock

Accelerated blood clearance (ABC)

phenomenon

Extrapolation

ABSTRACT

Hemoglobin-vesicles (HbV) are liposomal artificial oxygen carriers that may be useful as a resuscitation fluid during hemorrhagic shock (HS). It is well-known that the pharmacokinetic properties of liposome change in response to both pathological conditions and repeated administration. Therefore, we compared the pharmacokinetics of single versus repeated administration of HbV during HS. HS was induced by withdrawal of 40% of total blood volume. The normal (non-HS) and HS₁ group was received an injection of ¹²⁵I-labeled HbV (¹²⁵I-HbV). The HS₂ group was resuscitated with non-labeled HbV, and 1 h later, it received an injection of ¹²⁵I-HbV. The half-life was shorter in HS₁ rats, but it returned to non-HS levels after the second HbV injection. During 12 h after administration of HbV, tissue distribution of HbV was greatest in the HS₁ group; however, the HS₂ group had the greatest tissue distribution at subsequent time points. Excretion into urine, major elimination pathway, did not differ between non-HS and HS₁ rats, but was significantly reduced in the HS₂ group. Furthermore, the half-life of HbV in humans was estimated to be approximately 3–4 days using an allometric equation. This suggests that HbV may be a useful artificial oxygen carrier in HS based on HbV pharmacokinetics.

© 2009 Elsevier B.V. All rights reserved.

1. Introduction

Before new drugs are approved for clinical use, they must undergo many types of evaluation, including physicochemical tests, animal studies and clinical trials. Among these assessments, pre-clinical pharmacokinetic studies in various mammalian species are essential, as the results of such studies can be extrapolated to humans, allowing estimation of appropriate dosing regimens in humans. In fact, Kennedy et al. reported that inadequate pharmacokinetic data accounts for approximately 40% of the failed attempts to develop new drug-administration protocols during the clinical-trial stage of drug development [1].

To date, hemoglobin-based artificial oxygen carriers (HBOCs), which include cross-linked [2], polymerized [3] and polymer-conjugated Hbs [4], have been developed to overcome the problems associated with blood transfusion, such as cross-matching, blood-borne infection (human immunodeficiency virus, hepatitis virus), and shortage of donated blood. Several of these HBOCs are currently in the final stages of clinical evaluation. Unfortunately, however, some

HBOCs cause vasoconstriction, and consequently, hypertension, due to scavenging of nitric oxide (NO) by Hb. This phenomenon may occur because, unlike red blood cells (RBCs), HBOCs lack a cellular structure [5].

In contrast, hemoglobin vesicles (HbVs) are artificial oxygen carriers with a cellular structure similar to that of RBCs: highly concentrated Hb encapsulated in a phospholipid bilayer membrane. Thus, HbV does not cause the adverse effects that are associated with HBOCs [6,7], and HbV can control oxygen release to adjust the amount of allosteric effector [8]. Recently, Sakai et al. reported that the resuscitative effect of HbV is equivalent to that of autologous blood when injected into rats with hemorrhagic shock (HS) induced by withdrawal of 50% of total blood volume [9]. Yamasaki et al. also demonstrated that use of HbV as a cardiopulmonary bypass (CPB) priming solution improved neurologic and neurocognitive outcomes in a rat model of CPB [10].

Despite evidence for the effectiveness of HbV, the pharmacokinetics of HbV has not been well characterized. Thus far, pharmacokinetic studies of HbV have evaluated only changes in serum concentration and organ distribution in normal rats and rabbits administered ^{99m}Tc-labeled hexamethyl-propylenamine oxime (HMPAO) encapsulated in HbV [11]. However, resuscitation after

* Corresponding author. Tel.: +81 96 371 4150; fax: +81 96 362 7690.

E-mail address: otagirim@gpo.kumamoto-u.ac.jp (M. Otagiri).

massive hemorrhage resulting from clinically relevant causes, such as injury, accidental blood loss or major surgery, has not been studied in detail. Clinical conditions affect the pharmacokinetics of numerous drugs [12,13]. For example, clinical trials have shown that the pharmacokinetics of liposome-encapsulated amphotericin B differ between normal individuals and patients [14,15]. Therefore, clarifying the pharmacokinetics of HbV in animal models of HS should provide useful information, such as dosing regimens, for future clinical applications.

Moreover, clinical use of HbV in patients with HS would require repeated administration of HbV. Recently, PEGylated liposomes were shown to lose their long-circulation half-life after being administered twice to the same animals (referred to as “accelerated blood clearance (ABC) phenomenon”) [16]. Therefore, the pharmacokinetics of HbV, which has a liposomal structure, might be altered by repeated administration.

In the present study, we hypothesized that the pharmacokinetics of HbV would be altered by the presence of a pathological condition and by repeated administration. We directly labeled the internal Hb of HbV with iodine to create ^{125}I -labeled HbV (^{125}I -HbV) and subsequently investigated the changes in HbV pharmacokinetics resulting from a pathological condition and repeated administration using a rat model of HS induced by massive hemorrhage.

2. Materials and methods

2.1. Materials and animal

Hb solution, which was purified from outdated donated blood, was provided by the Japanese Red Cross Society (Tokyo, Japan). Pyridoxal 5'-phosphate (PLP) was purchased from Sigma Chemical Co. (St. Louis, MO). 1,2-dipalmitoyl-*sn*-glycero-3-phosphatidylcholine (DPPC), cholesterol, and 1,5-bis-*O*-hexadecyl-*N*-succinyl-L-glutamate (DHSG) were purchased from Nippon Fine Chemical Co. Ltd. (Osaka, Japan). 1,2-distearoyl-*sn*-glycero-3-phosphatidyl-ethanolamine-*N*-PEG (PEG-DSPE) was purchased from NOF Corp. (Tokyo, Japan). Recombinant human serum albumin (rHSA) was given by Nipro Corp. (Osaka, Japan).

Sprague-Dawley (SD) rats (180–210 g) were purchased from Kyudou Co. (Kumamoto, Japan). SD rats were maintained in a temperature-controlled room with a 12-hr dark/light cycle and *ad libitum* access to food and water. All animal experiments were performed according to the guidelines, principles, and procedures for the care and use of laboratory animals of Kumamoto University.

2.2. Preparation of HbV

HbVs were prepared under sterile conditions as previously reported [17]. Briefly, the encapsulated Hb (38 g/dl) contained 14.7 mM of PLP as an allosteric effector to regulate P_{50} to 25–28 Torr. The lipid bilayer was a mixture of DPPC, cholesterol, and DHSG at a molar ratio of 5/5/1, and PEG-DSPE (0.3 mol%). The HbVs were suspended in a physiological salt solution at [Hb] 10 g/dL, filter-sterilized (Dismic, Toyo-Roshi, Tokyo, Japan; pore size, 450 nm), and bubbled with N_2 for storage.

2.3. HbV labeling with ^{125}I (^{125}I -HbV) and preparation of HbV solution

^{125}I -HbV was prepared by incubation of HbV with Na^{125}I (Piscataway, NJ, USA) in an Iodo-Gen (1, 3, 4, 6-tetrachloro-3, 6-diphenylglycoluril) tube for 30 min at room temperature. Thereafter, ^{125}I -HbVs were isolated from free ^{125}I by passage through a PD-10 column (GE Healthcare Bio-Sciences AB). The ^{125}I -HbVs were then filter-sterilized (pore size, 450 nm) to remove aggregates. Over 97% of iodine was bound to internal Hb in HbV. Three different HbV suspensions were prepared: non-radiolabeled HbV and ^{125}I -HbV for HS model rats; and,




Fig. 1. Scheme illustrating (A) the method of preparing HS model and (B) the pharmacokinetic experimental protocol.

^{125}I -HbV for non-HS model rats. All suspensions were mixed with rHSA to adjust the albumin concentration of the vesicle-suspension medium to 5 g/dL. Under these conditions, the colloid osmotic pressure of the suspension is kept constant at approximately 20 mm Hg [17].

2.4. Preparation of hemorrhagic shock model rats

SD rats were anesthetized with pentobarbital. Subsequently, polyethylene catheters (PE 50 tubing, outer diameter equal to 0.965 mm, and inner diameter equal to 0.58 mm; Becton Dickinson and Co., Tokyo, Japan) containing saline and heparin were introduced into the left femoral artery for infusion and blood withdrawal, and into the right femoral artery to monitor mean arterial pressure (MAP), heart rate (HR) and pressure pulse. HS was induced by removal of 40% of total blood volume (22.4 mL/kg). Systemic blood volume was estimated to be 56 mL/kg [9]. Briefly, blood was removed as follows: constant withdrawal of 20% (1 mL/min); and, removal of 5% (0.5 mL/min) of total blood volume four times over a period of 5 min. After the last withdrawal of blood, animals were allowed to recover for 10 min at room temperature (Fig. 1A).

2.5. Measurement of cardiocirculatory dynamic parameters and plasma clinical chemistry

Ten HS rats were resuscitated by infusion of either isovolemic HbV (1400 mgHb/kg, 1 mL/min; $n = 5$) or saline (22.4 mL/kg, 1 mL/min; $n = 5$). After resuscitation, the polyethylene catheter of right femoral

artery was removed and the monitoring finished. Blood samples were collected immediately prior to blood withdrawal, after bleeding, and 7 days after resuscitation. Immediately after withdrawal, the pH of all blood samples was measured using a pH meter (HORIBA, Kyoto, Japan). An aliquot of blood was treated with perchloric acid to remove blood proteins for lactate analysis. The remaining blood was centrifuged (3000 g, 5 min) to obtain plasma for analysis of albumin, aspartate aminotransferase (AST), alanine aminotransferase (ALT), γ -glutamyltransferase (γ -GTP), amylase, lipase, total-cholesterol, free-cholesterol, HDL-cholesterol, phospholipids, urea nitrogen (BUN), creatinine (CRE) and iron (Fe). All plasma samples were stored at -80°C prior to analysis by a commercial clinical testing laboratory (SRL, Tokyo, Japan).

2.6. The pharmacokinetic experimental protocol

All rats were given water containing 5 mM sodium iodide (NaI) for the duration of the experiment to avoid specific accumulation in the glandula thyroidea. Ninety-six male SD rats were anesthetized with pentobarbital and polyethylene catheters were inserted into the left femoral artery. The rats were divided into three groups. Two groups (one-injection group (HS₁), $n=32$; two-injection group (HS₂), $n=32$) of rats underwent experimental HS. Both the non-HS (non-HS group, $n=32$) and HS₁ rats were injected with the ¹²⁵I-HbV suspension (1400 mg Hb/kg, 1 mL/min), 1 h after the first injection of non-labeled HbV suspension (1400 mg Hb/kg, 1 mL/min), the HS₂ rats received the ¹²⁵I-HbV suspension (1400 mg Hb/kg, 1 mL/min). The volume of the infused fluid was identical to the volume of blood withdrawn to induce HS (22.4 mL/kg) (Fig. 1B). In each group of rats, five rats were randomly selected to undergo the plasma concentration test. The rats were anesthetized using ether, blood samples were collected at multiple time points after the ¹²⁵I-HbV injection (3 min, 10 min, 30 min, 1 h, 6 h, 12 h, 24 h, 48 h, 72 h) and the plasma was separated by centrifugation (3000 g, 5 min). Degraded HbVs and free ¹²⁵I were removed from plasma by centrifugation in 1% bovine serum albumin and 40% trichloroacetic acid (TCA). Three rats in each treatment group ($n=27$ /group) were euthanized at each time point (3 min, 30 min, 1 h, 2 h, 6 h, 12 h, 24 h, 48 h, 72 h). The organs were excised (kidney, liver, spleen, lung, heart), rinsed with saline, and weighed. The levels of ¹²⁵I radiation in the excised organs, urine and feces, collected at fixed intervals (2 h, 6 h, 12 h, 24 h, 48 h, 72 h in each group; urine samples at 2 h and 6 h were not collected in the HS₁ group because of the reduction of renal function) using metabolic cages, were determined using a γ -counter (ARC-5000, Aloka, Tokyo, Japan). A two-compartment model was used to determine the pharmacokinetic parameters after HbV administration. Pharmacokinetic parameters were estimated by curve-fitting using MULTI, a normal least-squares program. Pharmacokinetic analysis was described in the supplementary materials and methods in detail.

2.7. Interspecies scaling of pharmacokinetic parameters

Allometric relationships between various pharmacokinetic parameters (P) and body weight (W) were plotted on a log-log scale. Linear regression of the logarithmic values was calculated using the least-squares method using Eq.(1) to obtain the coefficient (α) and exponent (β) values [18].

$$P = \alpha W^{\beta} \quad (1)$$

To calculate pharmacokinetic parameters, such as V_1 and CL for humans using Eq.(1), the total blood volume of rabbits and humans were assumed to be 5.7% and 7% of body weight, respectively [11]. The CL of rabbits and humans, which were previously reported by Sou et al., were calculated using V_1 and half-life (see supplementary methods Eq. (8)) [11]. The V_1 and CL of mice were 1.75 mL and 0.116 mL/h, respectively

(submitted in separate manuscript). After prediction of V_1 and CL for humans using Eq.(1), the half-life for human was estimated.

2.8. Data analysis

A two-compartment model was used in the pharmacokinetic analyses after HbV administration. Each parameter was calculated by fitting using MULTI, a normal least-squares program [19]. Data are means \pm SD for the indicated number of animals. Dunnett's test was used for comparison of baseline and subsequent values within each group. Significant differences among each group were examined using a Student's t -test. A probability value of $p < 0.05$ was considered to indicate statistical significance.

3. Results

3.1. Systemic response to hemorrhagic shock

MAP of the SD rats before induction of HS by bleeding was 79.9 ± 4.3 mm Hg; MAP declined significantly to 30.1 ± 1.8 mm Hg after induction of HS. During blood withdrawal to induce HS, MAP remained less than 40 mm Hg (hypotension) for 30 min. After resuscitation with HbV, MAP immediately increased to 81.4 ± 5.6 mm Hg, which was similar to MAP measured at baseline, and apparently HbV did not induce hypertension (Fig. 2A). Average HR before and after hemorrhage was 369 ± 44 and 290 ± 43 beats/min, respectively. However, during the interval between 10-min-post-bleeding and initiation of resuscitation, HR increased slightly (Fig. 2B). Pressure pulse, which reflects cardiac




Fig. 2. Change in (A) mean arterial pressure (MAP), (B) heart rate (HR), (C) pressure pulse from -5 min (5 min before bleeding) to 45 min (10 min after resuscitation). Each point represents the mean \pm SD ($n=5$). * <0.05 or ** <0.01 vs. 0 min.

Table 1

Plasma clinical chemistry analyses after withdrawal of 40% of total blood volume and 7 days after administration of HbV or saline.

	7 days after resuscitation			
	Baseline	After bleeding	HbV	Saline
Albumin, g/dL	3.38 ± 0.13	2.96 ± 0.15	3.34 ± 0.28	3.32 ± 0.26
Phospholipids, mg/dL	124.4 ± 15.4	103.6 ± 12.0	139.2 ± 12.9	129.4 ± 14.2
Total-cholesterol, mg/dL	75.0 ± 11.3	59.4 ± 8.8	80.8 ± 10.3	75.2 ± 9.5
Free-cholesterol, mg/dL	20.2 ± 2.0	16.2 ± 1.9	20.2 ± 2.8	18.8 ± 2.3
BUN, mg/dL	13.1 ± 1.9	13.7 ± 0.6	16.0 ± 1.0	15.6 ± 2.0
Creatinine, mg/dL	0.17 ± 0.01	0.29 ± 0.02 ^a	0.19 ± 0.02	0.19 ± 0.01
AST, IU/L	74.0 ± 9.3	65.6 ± 8.7	119.3 ± 28.8 ^a	101.8 ± 11.4 ^b
ALT, IU/L	42.4 ± 4.7	37.2 ± 2.4	57.0 ± 9.9	56.0 ± 9.7
γ-GTP, IU/L	<2.8	<2.8	<2.8	<2.8
HDL-cholesterol, mg/dL	23.0 ± 3.2	19.2 ± 2.6	28.2 ± 1.5	25.4 ± 2.9
Fe, μg/dL	153.5 ± 36.5	171.4 ± 54.7	148.7 ± 22.5	123.7 ± 66.3
Lipase, U/L	6.8 ± 0.4	6.8 ± 0.8	5.4 ± 0.5	5.8 ± 0.4
Amylase, U/L	669.8 ± 28.5	638.8 ± 34.0	604.0 ± 67.7	624.4 ± 66.0

The values are mean ± S.D. (n = 3–5). ^ap < 0.05, ^bp < 0.01 vs. baseline.

output, was 26.4 ± 3.2 and 15.3 ± 3.1 mm Hg before and after hemorrhage, respectively. After resuscitation, pressure pulse returned to baseline values (Fig. 2C). Acute bleeding resulted in a non-significant decrease in pH from 7.55 ± 0.06 to 7.51 ± 0.05, and a significant increase in lactate from 21.4 ± 4.0 to 62.8 ± 17.5 mg/dL (p < 0.01). In addition, skin pallor and hyperpnea developed during hypotension. These results indicate that acute blood withdrawal in rats is an adequate model of clinical HS.

3.2. Plasma clinical chemistry

All rats survived the complete experimental protocol. Table 1 summarizes the clinical chemistry parameters measured in plasma. Immediately after hemorrhage, the concentration of CRE in plasma, which reflects kidney function, was two-fold greater than the concentration at baseline (p < 0.01). In contrast, BUN, which is also a marker of kidney function, and other parameters were not affected by hemorrhage. We also performed clinical chemistry analyses in plasma that was collected from HS rats after 7 days of treatment with HbV or saline. Only AST increased significantly, reaching 119.3 ± 28.8 IU/L in the group given HbV (p < 0.01 vs. baseline) and 101.8 ± 11.4 IU/L in the group administered saline (p < 0.05 vs. baseline). CRE decreased from 0.29 ± 0.02 mg/dL to 0.19 ± 0.02 and 0.19 ± 0.01 mg/dL in the HbV and saline groups, respectively. Although HbV contains many lipids




Fig. 3. Relative plasma concentration of ¹²⁵I-HbV after administration of 1400 mgHb/kg via injection of non-HS, HS₁ or HS₂. Each point represents the mean ± SD (n = 5). * < 0.05 or ** < 0.01 vs. non-HS rats.

Table 2

Pharmacokinetic parameters of HbV in non-HS rats, HS₁ and HS₂ rats.

	Non-HS	HS ₁	HS ₂
t _{1/2α} , h	5.3 ± 3.9	0.58 ± 0.05 ^a	5.0 ± 1.7
t _{1/2β} , h	30.6 ± 4.0	18.1 ± 3.7 ^b	32.4 ± 1.1
C ₀ , % of dose/mL	9.1 ± 0.2	9.9 ± 0.5	9.8 ± 1.9
V ₁ , mL	10.9 ± 0.2	10.1 ± 0.5	10.4 ± 2.1
V ₂ , mL	5.5 ± 1.2	11.9 ± 1.6 ^b	5.9 ± 2.0
V _{ss} , mL	16.7 ± 1.1	21.6 ± 2.3 ^b	17.0 ± 3.1
ke, × 10 ⁻³ min ⁻¹	0.70 ± 0.06	1.4 ± 0.1 ^b	0.60 ± 0.07
k ₁₂ , × 10 ⁻³ min ⁻¹	0.41 ± 0.10	6.6 ± 2.3 ^b	7.4 ± 3.5 ^b
k ₂₁ , × 10 ⁻³ min ⁻¹	0.69 ± 0.23	5.3 ± 1.2 ^b	13.9 ± 4.7 ^b
AUC, % of dose/mL	210.3 ± 22.9	126.5 ± 12.8 ^b	259.0 ± 48.6
CL, mL/hr	0.47 ± 0.04	0.80 ± 0.08 ^b	0.40 ± 0.09
k ₁₂ /k ₂₁	0.60 ± 0.14	1.2 ± 0.1 ^b	0.55 ± 0.13

t_{1/2α}, the distribution-phase half-life; V_{ss}, the steady-state distribution volumes; k₁₂, the rate constants for the transfer from V₁ to V₂; k₂₁, the rate constants for the transfer from V₂ to V₁.

Values are means ± S.D. (n = 5). ^ap < 0.05, ^bp < 0.01 vs. non-HS rats.

and iron, lipid-related parameters, such as total-cholesterol, free-cholesterol, HDL-cholesterol, phospholipids, and Fe did not differ between the HbV- and saline-treated rats. These results show that neither organ function nor serum parameters were affected by HbV administration, despite the temporary disruption of circulatory function by HS.

3.3. Plasma concentration

The fate of the ¹²⁵I-HbV administered to non-HS, HS₁ and HS₂ rats was evaluated by determination of residual TCA-precipitable radioactivity in plasma. Fig. 3 shows the time course of the plasma concentration of ¹²⁵I-HbV in non-HS, HS₁ and HS₂ rats and Table 2 lists the pharmacokinetic parameters for these groups. The elimination-phase half-life (t_{1/2β}) of ¹²⁵I-HbV in HS₁ rats was reduced significantly—by 0.66-fold—compared with the t_{1/2β} of ¹²⁵I-HbV in non-HS rats. In addition, the AUC was decreased, while the elimination rate constant (ke) and clearance (CL) were increased in HS₁ rats compared with non-HS rats. Notably, the initial plasma concentration (C₀) and distribution volume of the central compartment of HbV (V₁) were identical between the two groups, whereas the distribution volume of the peripheral compartment (V₂) in HS₁ rats was nearly two-fold greater than non-HS rats. Interestingly, the plasma concentration of ¹²⁵I-HbV and pharmacokinetic parameters, except k₁₂ and k₂₁, of HS₂ rats were similar to the results observed in non-HS rats. Furthermore, V₁ (approximately 10 mL) was nearly equivalent to blood volume (56 mL/kg).

3.4. Organ distribution of HbV

Fig. 4 shows the time course of organ distribution of ¹²⁵I-HbV (% of ID) after ¹²⁵I-HbV administration. Shortly after ¹²⁵I-HbV administration, the amount of ¹²⁵I-HbV was significantly greater in the kidney, lung and heart of HS₁ rats compared with non-HS rats; however, there were no differences in distribution of ¹²⁵I-HbV in other organs between groups. As time passed, HbV was distributed to all observed organs (kidney, liver, spleen, lung and heart) of HS₁ rats, compared with the non-HS rats. In contrast, less ¹²⁵I-HbV was distributed in the liver, spleen, lung, and heart of the HS₂ rats compared with both non-HS and HS₁ rats. However, past 12 h after administration of HbV, HS₂ rats exhibited the greatest distribution of ¹²⁵I-HbV in kidney and liver. Furthermore, during the first 12 h after HbV administration, the total amount of ¹²⁵I-HbV distributed in all observed organs (kidney, liver, spleen, lung and heart) was greatest in HS₁ rats, while after 12 h post-administration of HbV, the greatest amount of total radioactivity was observed in the HS₂ rats.




Fig. 4. Distribution of HbV in kidney (a), liver (b), spleen (c), lung (d), heart (e), total organ (f) after injection of non-HS or HS₁ or HS₂ from the time of administration until 2 h post-administration and in kidney (A), liver (B), spleen (C), lung (D), heart (E), total observed organ (the sum of kidney, liver, spleen, lung and heart) (F) from the time of administration until 72 h post-administration. Each point represents the mean \pm SD ($n=3$). * <0.05 or ** <0.01 vs. non-HS rats. # <0.05 or ## <0.01 vs. HS₁.

The tissue distribution of ¹²⁵I-HbV in non-HS, HS₁ and HS₂ rats was also evaluated based on both percentage of the infused dose/organ weight (ID/g organ) and the tissue-to-plasma partition coefficient (*K_p*), which can be used to compare the kinetics of plasma and tissue HbV concentrations. Table 3 shows the ID/g organ and *K_p* data for the kidney, liver, spleen, lung and heart at 1 h, 12 h and 72 h after administration of HbV. The *K_p* data in spleen became more than 1 (mL/g of tissue), but that in other organ less than 1 (mL/g of tissue) at 1 h and 12 h. At 72 h post-administration of HbV, the % of ID and % ID/g organ values were

greatest, the *K_p* value was lowest, and there was an increase in the plasma concentration of HbV in the HS₂ group.

3.5. Urinary and fecal excretion of HbV

To identify the pathways by which HbV is excreted, the amount of HbV in urine and feces was measured in non-HS, HS₁ and HS₂ rats (Fig. 5). We found that HbV was excreted primarily in the urine of all groups (58.9 ± 5.6 , 65.5 ± 10.5 and $44.2 \pm 2.9\%$ of ID for non-HS, HS₁

Table 3

Dose/gram of organ weight (dose/g) and tissue-to-plasma partition coefficient (Kp) for non-HS rats, HS₁ and HS₂ rats.

		Dose/organ weight (dose/g)			Kp (mL/g of tissue)		
		Non-HS	HS ₁	HS ₂	Non-HS	HS ₁	HS ₂
1 h	Kidney	0.86 ± 0.18	1.32 ± 0.09 ^b	1.65 ± 0.14 ^{b,c}	0.14 ± 0.04	0.32 ± 0.02 ^b	0.37 ± 0.01 ^{b,c}
	Liver	1.15 ± 0.13	1.09 ± 0.14	0.93 ± 0.44 ^a	0.19 ± 0.01	0.26 ± 0.02 ^b	0.21 ± 0.03 ^c
	Spleen	7.87 ± 0.56	7.11 ± 0.42	9.30 ± 0.44 ^{a,d}	1.30 ± 0.09	1.72 ± 0.10 ^b	2.09 ± 0.19 ^{b,c}
	Lung	1.47 ± 0.42	1.48 ± 0.11	1.11 ± 0.16 ^c	0.24 ± 0.06	0.36 ± 0.06 ^a	0.25 ± 0.03 ^c
	Heart	0.90 ± 0.07	0.90 ± 0.17	0.52 ± 0.10 ^{b,c}	0.13 ± 0.01	0.22 ± 0.03 ^b	0.12 ± 0.02 ^d
12 h	Kidney	0.37 ± 0.05	0.64 ± 0.14 ^a	0.78 ± 0.09 ^b	0.16 ± 0.02	0.29 ± 0.07 ^a	0.25 ± 0.03 ^b
	Liver	0.87 ± 0.11	1.07 ± 0.31	1.29 ± 0.67	0.39 ± 0.05	0.47 ± 0.02 ^a	0.39 ± 0.16
	Spleen	7.23 ± 1.32	10.9 ± 2.69 ^a	6.49 ± 1.27 ^c	3.19 ± 0.61	4.89 ± 0.72 ^a	2.01 ± 0.08 ^{a,d}
	Lung	0.54 ± 0.08	0.71 ± 0.07 ^a	0.55 ± 0.10 ^c	0.24 ± 0.04	0.32 ± 0.11	0.17 ± 0.01 ^{a,c}
	Heart	0.24 ± 0.07	0.39 ± 0.09 ^a	0.30 ± 0.04	0.10 ± 0.03	0.18 ± 0.02 ^a	0.09 ± 0.01 ^d
72 h	Kidney	0.05 ± 0.01	0.08 ± 0.01 ^b	0.47 ± 0.15 ^{b,d}	0.21 ± 0.07	0.63 ± 0.01 ^b	0.52 ± 0.10 ^b
	Liver	0.33 ± 0.07	0.40 ± 0.07	0.75 ± 0.26 ^{a,c}	1.29 ± 0.28	3.12 ± 0.73 ^b	0.81 ± 0.15 ^{a,d}
	Spleen	1.52 ± 0.45	1.11 ± 0.16	1.75 ± 0.41 ^c	5.74 ± 0.23	7.06 ± 0.11 ^b	1.93 ± 0.39 ^{a,d}
	Lung	0.05 ± 0.01	0.05 ± 0.01	0.18 ± 0.07 ^d	0.18 ± 0.02	0.36 ± 0.09 ^a	0.19 ± 0.01 ^d
	Heart	0.03 ± 0.01	0.02 ± 0.01	0.09 ± 0.03 ^{b,d}	0.13 ± 0.02	0.19 ± 0.03 ^a	0.10 ± 0.01 ^d

Values are means ± S.D. (n = 3).

^ap < 0.05, ^bp < 0.01 vs. non-HS rats, ^cp < 0.05, ^dp < 0.01 vs. HS₁ rats.

and HS₂ rats, respectively), while radioactivity was virtually undetectable in feces (2.0 ± 0.5, 3.2 ± 1.3 and 2.8 ± 0.5% of ID, for non-HS, HS₁ and HS₂ rats, respectively). Urinary and fecal levels of radioactivity were not significantly different between the non-HS and HS₁ groups. Meanwhile, urinary radioactivity in HS₂ rats was significantly less than that in non-HS and HS₁ rats, and the half-life of HbV was significantly longer in HS₂ rats compared with the other groups.

3.6. Prediction of human pharmacokinetics

As mentioned in the "Plasma concentration" section, we found that V₁ was equivalent to arteriovenous blood flow. To predict the CL in humans, we examined the allometric relationship between V₁ and body weight in mice, rats, rabbits and humans. As shown in Fig. 6A, there was a good correlation between V₁ and body weight, with r² values of 0.99.

Using the relationship observed for both mice and rats, the CL of HbV in humans was predicted to be approximately 36.6 mL/h. In addition, we applied the rabbit [11] and human CL values for HbV to this relationship. We calculated the human CL (47.2 mL/h) using k_e, which is from Sou et al. [11], and our estimated V₁. These values fit our allometric equation well (Fig. 6B). The half-life of HbV in humans was extrapolated to be approximately 96 h.

4. Discussion

HbV is a cellular-type, artificial oxygen carrier that consists of a highly concentrated Hb solution coated with a phospholipid bilayer membrane. Recent findings have demonstrated that HbV restores tissue and plasma oxygenation [9], and improves survival [20] in HS animal models. To exert these pharmacological effects and enhance oxygen-carrying capacity, HbV, or any other RBC substitute, must have a long half-life in circulation (t_{1/2}; free Hb and PEG-Hb in rats for 1.5 and 10.9 h, respectively [21,22], while RBC in human for up to 60 days [23]).

In the present study, the elimination-phase half-life (t_{1/2β}) of ¹²⁵I-HbV in the HS₁ rats was shorter than that in non-HS rats. This result does not seem to support the therapeutic use of HbV, because the most important determinant of HbV efficacy is HbV-derived Hb concentration in blood. However, the shorter half-life in the HS₁ group may reflect transfer of HbV from arteriovenous blood to the organ capillary beds, thereby normalizing distribution of blood to the organs after bleeding. This scenario would result in an apparent reduction in

HbV in arteriovenous circulation. Consistent with this hypothesis, HbV has been shown to both reach microcirculation in critically ischemic peripheral tissue and promote recovery from capillary pathological hemodynamic conditions [24]. In addition, the results of the present study also support our hypothesis: 1) the initial plasma concentration of HbV (C₀) and the amount of HbV eliminated in HS₁ rats were similar to the values observed in non-HS rats, such that the amount of residual HbV in non-HS and HS₁ rats were similar; 2) the time course of HbV distribution in the kidney, liver, spleen, lung and heart in the HS₁ rats was significantly greater than that of the non-HS rats; and, 3) the plasma concentration of ¹²⁵I-HbV and pharmacokinetic parameters in the HS₂ rats were similar to the values observed in the non-HS rats.

For HbV to keep long circulating retention, the enhanced tissue distribution of HbV is a concern, as it may reflect both increased scavenging of HbV by cells of the mononuclear phagocyte system (MPS), such as Kupffer cells, red pulp zone splenocytes, and mesangial cells [7], and functional changes in MPS after HS. However, alterations in HbV, such as PEG-modification and controlled particle size (250 nm), suppress scavenging by the MPS relative to scavenging of Hb encapsulated in other liposomes [21,25]. Moreover, MPS scavenging of liposomal preparations is influenced by the dose administered [26], while mild hemorrhage does not affect phagocytic activation of Kupffer cells [27]. Therefore, because we used a high dose of HbV in




Fig. 5. Time-dependence of urinary and fecal radioactivity after administration of ¹²⁵I-HbV to non-HS or HS₁ or HS₂. The solid line represents urinary excretion, and the dotted line represents fecal excretion. Each point represents the mean ± SD (n = 3). * < 0.05 or ** < 0.01 vs. non-HS rats. ## < 0.01 vs. HS₁.