


3D). Similar results were obtained when *scnn1a* promoter was used for ChIP assay (data not shown). These findings strongly support the notion that protein levels of endogenous HEXIM1 might determine GR recruitment onto these promoters and subsequent glucocorticoid-responsive transcription of *atp1a1* and *scnn1a* in HepG2 cells.

### GR and HEXIM1 Stochastically Contribute to Hormone-Dependent Transcriptional Regulation of GR-Target Genes

To test relative contribution of GR and HEXIM1 in the expression of *atp1a1*, we overexpressed FLAG-GR alone or in combination with HEXIM1 in HepG2 cells



**Fig. 4.** GR and HEXIM1 Stochastically Contribute to Hormone-Dependent Transcriptional Regulation of GR-Target Genes


**A**, HepG2 cells were infected with the recombinant adenoviruses expressing FLAG-GR and HEXIM1 in phenol red-free Opti-MEM I at MOI of 5 for 24 h as indicated. Cells were lysed and subjected to Western blot analysis using the indicated antibodies. **B**, HepG2 cells were infected with the recombinant adenoviruses as described in panel A and treated with or without 100 nM DEX for 6 h. Total RNA was prepared, and mRNA for *atp1a1*, *sgk1*, and *gapdh* was measured with qRT-PCR. Samples were normalized to *gapdh*, and relative expression levels to vehicle-treated samples are presented as relative mRNA expression. Error bars represent SD values of at least three independent experiments. **C**, HepG2 cells were infected with the recombinant adenoviruses in phenol red-free Opti-MEM I for 24 h as indicated and treated with or without 1  $\mu$ M DEX for 20 min. ChIP assays were performed with anti-GR antibodies as described in *Materials and Methods*. IP, Immunoprecipitation.

using recombinant adenoviruses. Western blots showed that the protein levels of expressed FLAG-GR and HEXIM1 were approximately 5-fold and 8-fold compared with those of endogenous GR and HEXIM1, respectively (Fig. 4A). In HepG2 cells, overexpression of FLAG-GR restored glucocorticoid responsiveness and resulted in DEX-dependent induction of *atp1a1* mRNA by 1.8-fold, which was again canceled by coexpression of HEXIM1 (Fig. 4B). We may propose, therefore, that high-level expression of HEXIM1 relative to GR confers tissue-specific glucocorticoid resistance of *atp1a1* in HepG2 cells. This is also the case in *scnn1a* mRNA expression (data not shown). Interestingly, mRNA expression of *sgk1* was also negatively affected by exogenous expression of HEXIM1 (Fig. 4B). Our ChIP assay revealed that GR overexpression restored hormone-dependent recruitment of GR to *atp1a1* promoter in a dose-dependent manner, which was again canceled by overexpression of HEXIM1 (Fig. 4C). In the case of *sgk1* promoter as well, overexpression of GR further increased hormone-dependent GR recruitment, which was antagonized by exogenous HEXIM1 (Fig. 4C). These findings strongly support the notion that GR and HEXIM1 stochastically contribute to hormone-dependent transcriptional regulation of both *atp1a1* and *sgk1*, and that *atp1a1* promoter is more susceptible to HEXIM1.

### Knockdown of HEXIM1 Enhanced P-TEFb-Dependent Gene Expression


In addition to suppressing GR recruitment onto the target DNA, HEXIM1 is originally reported to inactivate kinase activity of P-TEFb, thereby suppressing transcription elongation (see *Introduction*). To investigate whether knockdown of HEXIM1 in our system affects P-TEFb-dependent mRNA expression as previously reported (51), we analyzed mRNA expression of several genes, expression of which has been reported to be critically regulated by P-TEFb at the step of transcription elongation.

mRNA level of *cyp1a1* was increased after stimulation with 1 h treatment of 10 nM 3-methylcholanthrene (3MC) by 3.7-fold, probably via activation of AhR and subsequent recruitment of P-TEFb onto *cyp1a1* promoter (38), and AdsiHEXIM1 further enhanced this 3MC effect by 5.0-fold (Fig. 5A). IL-6-mediated expression of *junB* mRNA (14-fold), which is mediated by STAT3 (52), was also enhanced by AdsiHEXIM1 (22-fold). Finally, heat shock-mediated amplification of mRNA expression of *hspa1a* and *fos* (11-fold and 8.1-fold, respectively), which is mediated by HSF1 (53, 54), was further enhanced by



**Fig. 5. Knockdown of HEXIM1 Enhanced P-TEFb-Dependent Gene Expression**

A, HepG2 cells were infected with AdsiCtrl or AdsiHEXIM1 as described in Fig. 3B and stimulated with 10 nM 3MC, 100 ng/ml IL-6, or culture at 42°C (heat shock) for 1 h as indicated. Endogenous mRNA for cytochrome P450, family 1, subfamily A, polypeptide 1 (*cyp1a1*), JunB (*junB*), heat shock 70-kDa protein 1A (*hspa1a*), Fos (*fos*), and *gapdh* was measured with qRT-PCR. Samples were normalized to *gapdh*, and relative expression levels to the AdsiCtrl-infected and unstimulated samples are presented as relative mRNA expression. Error bars represent SD values of at least three independent experiments. B, HepG2 cells were infected with the recombinant adenoviruses and stimulated as described in panel A. Nuclear extracts were prepared, and protein expression levels of endogenous AhR, STAT3, HSF1, CDK9, and  $\alpha$ -actinin were assessed by Western blotting.



**Fig. 6.** P-TEFb-Binding and GR Binding Are Separable for HEXIM1

**A.** Schematic illustration of wild-type (wt) and mutant HEXIM1 used in this study. BR encompassing 150–177 amino acids are depicted as a *solid box*. Numbers depict positions of amino acids. Nucleotide and amino acid sequences in the BR are shown. Substitutions of nucleotides are shown in *uppercase letters*. wt (SR) and 168–177A (SR) have nucleotide substitutions in the target nucleotide sequence for AdsiHEXIM1 without affecting original amino acid sequence. 159–167A (SR) and dBR+SV (SR) are resistant to AdsiHEXIM1 by nature. **B.** HeLa cells were cotransfected with empty vector or expression plasmids for indicated FLAG-tagged mutant HEXIM1. Whole-cell lysates were prepared and subjected to FLAG-affinity purification as described in *Materials and Methods*. Western blot analysis of input lysates (lanes 1–5) and affinity-purified fractions (lanes 6–10) were performed using anti-FLAG peptide, anti-CycT1, and anti-CDK9 antibodies. **C.** COS7 cells were cotransfected with empty vector or expression plasmids for indicated HA-tagged mutant HEXIM1 along with either FLAG-tagged GR (*middle panel*) or FLAG-tagged PPAR $\gamma$  (*lower panel*) expression plasmid. Cells were treated with 100 nM DEX (*middle panel*) or 100 nM TGZ for 2 h. Nuclear extracts were prepared and immunoprecipitated with anti-HEXIM1 antibodies. Western blot analysis of input extracts (lanes 1–5) and immunoprecipitated fractions (lanes 6–10) were performed using anti-HA peptide and anti-FLAG peptide antibodies. a.a., Amino acids; IP, immunoprecipitation; NE, nuclear extract; NLS, nuclear localization signal.

AdsiHEXIM1 (17-fold and 14-fold, respectively). Note that protein expression levels of AhR, STAT3, HSF1, and CDK9 were not significantly influenced by infection of AdsiHEXIM1 (Fig. 5B). We may conclude,

therefore, that P-TEFb activity was enhanced in HEXIM1-knocked down cells and that the P-TEFb-dependent elongation process was up-regulated in certain genes.

### P-TEFb Binding and GR Binding Are Separable for HEXIM1

To highlight the GR target gene-selective role of HEXIM1, we established an experimental system that enables us to clarify which function of HEXIM1 is important in regulation of GR-target gene expression, P-TEFb suppression or GR sequestration. In short, endogenous HEXIM1 was knocked down by infection of AdsiHEXIM1, and mutant HEXIM1, which lacks either P-TEFb-suppressing activity or direct interaction with GR, or both, was exogenously complemented. To obtain such a mutant HEXIM1, we focused on basic region (BR) of HEXIM1 and made alanine substitution and domain swap mutants, as schematically depicted in Fig. 6A, because we and others previously showed that BR is essential for nuclear localization, interaction with GR, and P-TEFb-inhibition (28, 30, 41, 55). Because siRNA against HEXIM1 in AdsiHEXIM1 was designed to target the region corresponding to amino acids 159–165 (Fig. 6A), the expression plasmids for siRNA-resistant wild-type (SR) and 168–177A (SR) were created with several nucleotide substitutions in HEXIM1 cDNA without affecting original amino acid sequence (Fig. 6A). In indirect immunofluorescence analysis, every mutant HEXIM1 protein was expressed in the nucleus in transfected cells (data not shown).

To verify the presence or absence of the interaction between P-TEFb and these mutant FLAG-tagged HEXIM1, we, after transfection of their expression plasmids into HeLa cells, immunoprecipitated cell ly-

sate with anti-FLAG monoclonal antibody, and blots were probed with the antibodies against major P-TEFb subunits CycT1 and CDK9. As expected, substitution of BR to the nuclear localization signal from simian virus (SV) 40 large T antigen, resulting in dBR+SV (SR), completely abolished binding of CycT1 and CDK9 (Fig. 6B). Alanine substitution of amino acids 159–167, which was shown to disrupt the interaction with 7SK snRNA (55), diminished consecutive recruitment of CycT1 and CDK9 (Fig. 6B), as seen in dBR+SV (SR). On the other hand, alteration of amino acids 168–177 to alanines did not affect binding of CycT1 or CDK9 (Fig. 6B).

Using these HEXIM1 mutants, we also studied the physical interaction between HEXIM1 and GR. For this purpose, hemagglutinin (HA)-tagged wild-type (SR) and mutant HEXIM1 were expressed in COS7 cells along with either FLAG-tagged GR or FLAG-tagged peroxisome proliferator-activated receptor  $\gamma$  (PPAR $\gamma$ ) as a control and immunoprecipitated with anti-HEXIM1 antibodies. As shown in Fig. 6C, GR bound not only wild-type (SR) but also HEXIM1 mutants with alanine substitution, but the swap mutant dBR+SV (SR) did not bind GR. These results may suggest that amino acids 159–177 of HEXIM1 are not critical for binding GR, but protein configuration of BR and its proximity is important for GR recognition. In contrast, PPAR $\gamma$  did not bind wild-type (SR) or any mutant HEXIM1 (Fig. 6C). To further confirm that these HEXIM1 mutants, especially 159–167A (SR), retain not P-TEFb-inhibition but GR suppression, we tested their





Fig. 7. Differential Functions of HEXIM1 on P-TEFb- and GR-Dependent Gene Expression

A, COS7 cells were cotransfected with empty vector or expression plasmids for the indicated HA-tagged mutant HEXIM1 along with GR expression plasmid and GRE reporter plasmid. Four hours later, media were replaced, further cultured for 20 h, and treated with vehicle or 100 nM DEX for 18 h as indicated. Cells were lysed and subjected to luciferase assay. Results are presented as relative light units (RLU) per microgram of protein in the lysates. Error bars represent SD values of at least three independent experiments. Protein expression levels of GR and HA-HEXIM1 were assessed in Western blotting. B, COS7 cells were cotransfected with empty vector or expression plasmids for indicated HA-tagged mutant HEXIM1 along with PPAR $\gamma$  expression plasmid and PPARRE reporter plasmid. Media were replaced 4 h later, further cultured for 20 h, and treated with vehicle or 100 nM TGZ for 18 h as indicated. Cells were lysed and subjected to luciferase assay. Results are presented as RLU per microgram of protein in the lysates. Error bars represent SD values of at least three independent experiments. Protein expression levels of PPAR $\gamma$  and HA-HEXIM1 were assessed by Western blotting.

functions in a GRE-luciferase reporter gene assay (Fig. 7A). PPAR $\gamma$ -dependent reporter gene assay served as a control (Fig. 7B), because neither wild-type (SR) nor any mutant HEXIM1 was capable of binding PPAR $\gamma$  (Fig. 6C). HEXIM1 dBR+SV (SR), which lacks binding activity to either P-TEFb or GR, did not significantly affect either reporter gene activity, as expected (Fig. 7, A and B, top). With respect to GR-driven reporter gene expression, any alanine-substituted HEXIM1 mutant suppressed ligand-dependent activation of the reporter gene as well as wild type (SR), indicating its functional interaction with GR (Fig. 7A, top). In clear contrast, PPAR $\gamma$ -mediated activation of the reporter gene was repressed solely by wild-type (SR) and 168–177A (SR) (Fig. 7B, top). In these experimental settings, protein expression of FLAG-tagged GR or FLAG-tagged PPAR $\gamma$  was not significantly affected by HEXIM1 mutants (Fig. 7, A and B, bottom). It is concluded, therefore, that these HEXIM1 mutants can serve an efficient tool for delineating mechanism of suppressing expression of particular genes by HEXIM1, i.e. P-TEFb suppression or GR binding.

### P-TEFb Is Not Involved in HEXIM1-Mediated Suppression of Glucocorticoid Responsiveness

Finally, we differentially evaluated the importance of P-TEFb-suppressing and GR-binding activities of HEXIM1 in regulating glucocorticoid sensitivity of glucocorticoid-inducible mRNA expression of *atp1a1* and *scnn1a*. HeLa cells were transfected with the expression plasmids for HEXIM1 (SR) mutants, infected with AdsiHEXIM1, and treated with the cognate ligands, after which RNA was isolated for qRT-PCR analyses. In HEXIM1 knocked-down cells, mRNA expression of *cyp11a1*, which is known to be P-TEFb dependent (38), was stimulated by 10-fold in response to 6 h treatment with 10 nM 3MC (Fig. 8A). Adding back of wild-type (SR) HEXIM1 repressed induction of mRNA expression of *cyp11a1*, to 3.3-fold, suggesting that ectopically expressed HEXIM1 (SR) functionally suppressed P-TEFb activity (Fig. 8A). However, neither 159–167A (SR) nor dBR+SV (SR) repressed *cyp11a1* expression, confirming that the suppression of P-TEFb activity may be critical for the repression (Fig. 8A). In support of this, 168–177A (SR), which binds P-TEFb



**Fig. 8.** P-TEFb Is Not Involved in HEXIM1-Mediated Suppression of Glucocorticoid Responsiveness

A, HeLa cells were transfected with 3  $\mu$ g of empty vector or expression plasmids for the indicated HA-tagged mutant HEXIM1. Cells were infected 4 h later with AdsiHEXIM1 in phenol red-free Opti-MEM I at MOI of 100 for 36 h, and treated with vehicle or 10 nM 3MC (top panel), or 100 nM DEX (middle and bottom panels) for 6 h. Endogenous mRNA levels for *cyp11a1*, *atp1a1*, *scnn1a*, and *gapdh* were measured with qRT-PCR. Samples were normalized to *gapdh* mRNA, and mRNA induction levels by cognate ligands are shown in fold induction. Error bars represent SD values of at least three independent experiments. B, HepG2 cells were infected with FLAG-GR-expressing adenovirus (MOI of 50) along with LacZ- or mutant HEXIM1-expressing adenoviruses (MOI of 40) in phenol red-free Opti-MEM I for 24 h, and the cells were treated with 1  $\mu$ M DEX for 20 min. ChIP assays were performed with polyclonal anti-GR antibodies as described in Materials and Methods. Error bars represent SD values of at least three independent experiments. IP, Immunoprecipitation.

(Figs. 6B and 7B), suppressed AhR-mediated transcription as well as wild type (SR) (Fig. 8A). These effects of wild-type (SR) and mutant HEXIM1 were also observed in the other P-TEFb-regulated genes depicted in Fig. 5A (data not shown). With respect to GR target genes, mRNA expression of *atp1a1* and *scnn1a* was stimulated by 1.8-fold and 2.1-fold, respectively, in response to 6 h treatment with 100 nM DEX in HEXIM1 knocked-down cells (Fig. 8A). Complementation of wild-type (SR) HEXIM1 or 168–177A (SR) significantly repressed induction of mRNA expression of *atp1a1* and *scnn1a* (Fig. 8A). In contrast to *cyp11a1*, 159–167A (SR) suppressed glucocorticoid-induced enhancement of mRNA expression of *atp1a1* and *scnn1a* comparable to that of wild type (SR) (Fig. 8A), indicating that P-TEFb-binding activity of HEXIM1 is dispensable but GR-binding activity is important for the suppression. Consistently, dBR+SV (SR), which does not bind GR, did not affect mRNA induction of *atp1a1* and *scnn1a* (Fig. 8A). The importance of GR binding of HEXIM1 was also confirmed in ChIP assay. Recombinant adenovirus-mediated expression of 159–167A in HepG2 cells suppressed DEX-dependent recruitment of FLAG-GR onto *atp1a1* promoter, whereas dBR+SV did not (Fig. 8B). Using *scnn1a*, we obtained identical results (data not shown). Taken together, we may conclude that direct interaction between GR and HEXIM1 is critical for HEXIM1-mediated glucocorticoid resistance of *atp1a1* and *scnn1a* in HepG2 cells.

## DISCUSSION

As described in the introductory section, HEXIM1 is currently considered to be a multifunctional protein, acting at a specific stage of gene expression. In the present study, we intended to characterize endogenous HEXIM1 function for modulation of GR-mediated transcriptional regulation. For that purpose, we focused on *atp1a1* and *scnn1a*, because expression of these genes is resistant in HEXIM1-rich HepG2 cells to treatment with DEX (Fig. 1A). Treatment with histone deacetylase inhibitor did not result in liberation of these genes in HepG2 cells (Fig. 1A), suggesting that the observed DEX resistance is not due to irreversible alteration in higher order chromatin structure or histone acetylation-related chromatin packaging. In support of this, these genes retain responsiveness to other extracellular stimuli in liver and HepG2 cells (Refs. 10 and 11 and data not shown). We showed that, in HepG2 cells, knockdown of HEXIM1 by siRNA not only canceled the DEX resistance but also rather enhanced DEX-responsive mRNA expression of these genes (Fig. 3C). Moreover, our ChIP assay clearly demonstrated that siRNA-mediated knockdown of HEXIM1 restored hormone-dependent GR recruitment onto the promoters of those genes in parallel with corresponding increase in RNAPII binding (Fig. 3D). Such effect of reduction in endogenous HEXIM1 level

was mimicked by exogenous overexpression of GR (Fig. 4, B and C), indicating that GR-HEXIM1 ratio could be a determinant of glucocorticoid resistance/sensitivity of those genes. As anticipated, overexpression of HEXIM1 turned those promoters more or less resistant to DEX (Fig. 4B).

Endogenous HEXIM1 seems to negatively modulate all GR target genes but not completely diminish DEX responsiveness of all of them in HepG2 cells (Fig. 4), indicating that efficiency of the suppression by HEXIM1 is dependent on gene context. Indeed, our previous DNA microarray analyses showed that the extent of reducing DEX responsiveness by overexpressed HEXIM1 was variable among different genes in HepG2 cells (30). It is also reported that GRE occupancy with GR in alveolar epithelial A549 cells is generally restricted to such genes that are actually regulated by glucocorticoids in those cells (21). This observation strongly supports the idea that gene-specific determination of GR recruitment to GRE is important in tissue-specific regulation of glucocorticoid-responsive gene expression at the level preceding transcription initiation. We recently demonstrated that HEXIM1 directly binds GR and that GR or other oxosteroid receptors are preferential partners of HEXIM1 (48). In this line, we might speculate that HEXIM1 sequesters GR in the nucleus and inhibits its access to target gene promoter, and such negative effect of HEXIM1 is, more or less, shared by many genes. Some GR-target genes, including *atp1a1* and *scnn1a*, therefore, might be particularly susceptible to HEXIM1 and resistant to glucocorticoids in HEXIM1-rich cells, i.e. HepG2 cells. Certain promoters, e.g. *sgk1* promoter, allow hormone-dependent GR binding in HepG2 cells, strongly supporting the notion that promoter recruitment of GR is determined in a gene context-dependent manner as previously predicted in other GR-regulated genes (56).

We also revealed that P-TEFb-suppressing activity of HEXIM1 is not prerequisite for glucocorticoid resistance of these genes (Fig. 8A). Furthermore, the fact that 159–167A binds GR and suppresses GR recruitment to the target gene (Figs. 6C and 8B) again emphasizes the importance of the suppression of GR recruitment through direct GR-HEXIM1 interaction in the mechanisms of glucocorticoid resistance by HEXIM1. These results highlighted the role of HEXIM1 in P-TEFb-independent and gene-selective suppression of mRNA expression. The bimodal roles of HEXIM1 may differentially contribute to suppressing mRNA expression in a gene context-dependent manner. In this line, it should be noted that other transcription factors, such as estrogen receptor (45) and CCAAT/enhancer binding protein  $\alpha$  (44), which were shown to directly interact with HEXIM1, may also be controlled by HEXIM1 through a P-TEFb-independent mechanism. Moreover, the interaction of HEXIM1 with these transcription factors has been shown to be a molecular basis for various physiological or pathological actions of HEXIM1 (44, 45).

Our previous observation revealed that HEXIM1/GR complexes reside in a distinct subnuclear area (30). Given this, HEXIM1 might prevent intranuclear GR from accessing to the promoter and decrease the amount of available GR for transcription. Since we revealed that the central and C-terminal regions of HEXIM1 are indispensable for its proper nuclear localization and GR repression (Refs. 41 and 48, and data not shown), HEXIM1 might anchor at as yet unknown but saturable subnuclear structure via these regions. Increasing evidence indicates that the C-terminal region of HEXIM1 possesses various functions, e.g. P-TEFb-binding (57, 58), self-oligomerization (59–61), and interaction with transcription factors (44, 51). Recently, nucleophosmin was shown to bind HEXIM1 via BR and promote its degradation (62). Taken together, it may be indicated that subnuclear localization and function of HEXIM1 might be tightly controlled via multimodal interactions among distinct HEXIM1 domains and various nuclear machineries to elicit fine tuning of transcriptional control of gene expression. In any case, an important question to be solved is how multiple functions of HEXIM1 are rationally regulated in a gene- or tissue-dependent manner.

Expression levels of HEXIM1 vary in different tissues and are modulated during differentiation and development as well as in response to extracellular stimuli (see "Introduction"). Disturbances of tissue-specific glucocorticoid responses have been implicated in pathophysiology of rheumatoid arthritis, osteoarthritis, Crohn's disease, ulcerative colitis, asthma, AIDS, osteoporosis, and metabolic syndromes (63). Numbers of proteins have been shown to affect GR activity at different steps of GR signaling pathway and indicated to be potentially involved in the pathogenesis of such diseases that have relations to disturbed glucocorticoid responses in particular tissues (63). HEXIM1-mediated repression of GR might be one of such mechanisms and play pathological roles in certain diseases. On the other hand, glucocorticoids are still indispensable in treatment for a numerous diseases (4, 64). However, the desired therapeutic effects are often accompanied by severe side effects. Pharmacological alteration of the expression levels of HEXIM1, if possible, might indirectly modulate glucocorticoid effects in a tissue-specific manner and enable selective expression of pharmacological actions of glucocorticoids in given tissues. Along with development of selective GR modulators (26), HEXIM1 might also be considered as a drug target for tissue-specific modulation of GR actions.

## MATERIALS AND METHODS

### Reagents and Antibodies

DEX, troglitazone (TGZ), 3MC, and TSA were purchased from Sigma-Aldrich (St. Louis, MO). Recombinant human IL-6 was from Peprotech (London, UK). Other reagents were from

Nacalai Tesque (Kyoto, Japan) unless otherwise specified. Polyclonal antibodies against CDK9, STAT3, HSF1, CycT1, PPAR $\gamma$ , GR, and HA-peptide were from Santa Cruz Biotechnology, Inc. (sc-484, sc-7179, sc-9144, sc-8127, sc-7196, sc-8992, and sc-805, respectively; Santa Cruz, CA). Polyclonal anti-AhR antibodies were from Biomol (SA-210; Plymouth Meeting, PA). Polyclonal anti-FLAG-peptide antibodies and monoclonal anti- $\alpha$ -actinin antibody were from Sigma-Aldrich (F7425 and A5044, respectively). Monoclonal anti-GR antibody was from BD Biosciences (San Jose, CA). Monoclonal anti-RNAPII antibody was from Covance Laboratories, Inc. (MMS-126R; Princeton, NJ). Rabbit antihuman HEXIM1 antiserum and rabbit antimouse HEXIM1 antiserum were generated against a peptide corresponding from 39–53 amino acids of human HEXIM1 (RVPEEDSRWQSRAPP) and 55–69 amino acids of mouse HEXIM1 (SGSRPGQEGEGGLKH-I), respectively. Polyclonal anti-HEXIM1 affinity-purified antibodies were obtained from antihuman HEXIM1 antiserum with immunogen-immobilized affinity matrix (Kitayama Labes, Ina, Japan).

### Cell Culture and Transfection

HepG2, 293, COS7, and HeLa cells were from RIKEN cell bank (Tsukuba, Japan) and maintained in DMEM supplemented with 10% fetal calf serum (Invitrogen, Carlsbad, CA) and antibiotics in a humidified atmosphere at 37 C with 5% CO $_2$ . Before transfection, cells were washed twice with PBS, and media were replaced with phenol red-free Opti-MEM 1 (Invitrogen). Transient transfection was performed using TransIt-LT1 transfection reagent (Panvera, Madison, WI) as described previously (65). Total amounts of plasmids to transfect were kept constant by adding empty vector.

### Western Blot Analysis

Whole-cell extracts or nuclear extracts were prepared as described previously (30), resolved in sodium dodecyl sulfate (SDS)-polyacrylamide gels, and blotted to polyvinylidene fluoride membranes. The membranes were incubated with Blocking One (Nacalai Tesque) at room temperature for 1 h, incubated with specific antibodies diluted in Blocking One (1:500 dilution for HA-peptide or 1:2000 for the others) at 4 C for 18 h, and then, washed three times with TBS-T (25 mM Tris-HCl, pH 8.0; 125 mM NaCl; 0.1% Tween 20), incubated with secondary antibodies conjugated to horseradish peroxidase (GE Healthcare, Buckinghamshire, UK) at room temperature for 30 min, washed three times with TBS-T, and detected with Chemi-Lumi One L (Nacalai Tesque) according to manufacturer's instruction.

### Recombinant DNA and Adenoviruses

Expression plasmids for FLAG-tagged HEXIM1 (wild-type and dBR+SV) were described previously (30). pFLAG-CMV2-derived mammalian expression plasmids for mutant FLAG-HEXIM1 (159–167A and 168–177A) were generous gifts from Dr. Q. Zhou (University of California, Berkeley, CA). cDNA fragments for wild-type (SR), 159–167A (SR), 168–177A (SR), and dBR+SV (SR) HEXIM1 were generated by a standard PCR protocol using custom-designed primers and subcloned into pCMV-HA (TaKaRa, Otsu, Japan) or pFLAG-CMV2 (Sigma-Aldrich) expression plasmid using blunt-ended EcoRI and XhoI sites. The expression plasmid for human PPAR $\gamma$ , pCMX-6His-PPAR $\gamma$ , was generated by cloning appropriate PCR fragments into pCMX-6His vector (65). The PPAR response element (PPARRE)-driven reporter plasmid p3xPPARRE-LUC was a kind gift from Dr. E. A. Jansson (Karolinska Institutet, Stockholm, Sweden). All plasmids constructed above were verified by DNA sequencing. Recombinant adenoviruses encoding double-stranded hairpin RNAs for siRNA

against HEXIM1, AdsiHEXIM1, or control siRNA, AdsiCtrl, were constructed by subcloning expression cassettes from pSilencer3.1-H1 neo-derived expression plasmids (30) into adenoviral genome using Adenovirus Expression Vector Kit (TaKaRa) according to the manufacturer's instruction. Recombinant adenoviruses prepared from 293 cells were purified with Virakit AdenoMini-24 (Virapur, San Diego, CA) and titrated using Adeno-X Rapid Titer Kit (TaKaRa).

#### qRT-PCR

Total RNA was prepared with Sepasol-RNA I super (Nacal Tesque), reverse-transcribed with oligo-dT primer using SuperScript First-Strand Synthesis System for RT-PCR (Invitrogen). qRT-PCR was performed with the LightCycler TaqMan Master, Universal ProbeLibrary Set, Human, and LightCycler ST300 systems (Roche, Indianapolis, IN) according to manufacturer's instructions. Expression levels of mRNA were calculated on the basis of standard curves generated for each gene. mRNA for *gapdh* was used as an internal control. Sequences of primers used in this study are shown below:

*atp1a1*: 5'-ccctggctgcttctctt-3' and 5'-ggcacagaaccacc-agga-3'  
*scn11a*: 5'-aacccaggtctctctgcaacc-3' and 5'-gaaagatag-cagttccatcacatcg-3'  
*sgk1*: 5'-cctgagcttgaatgccaac-3' and 5'-gccaaggtg-atttctgctgag-3'  
*adh1a*: 5'-aagggccatgaagttctgatt-3' and 5'-ccactggg-cactctgctg-3'  
*cyp11a1*: 5'-cccagctcagctcagctacct-3' and 5'-ggagattggg-aaaagcatga-3'  
*juno*: 5'-atacacagctacgggatacgg-3' and 5'-gctcgtttca-ggagttgt-3'  
*hspa1a*: 5'-ggagctcctacgcttcaaca-3' and 5'-ccagcactt-cttctgtctg-3'  
*fos*: 5'-ctaccactcaccocagact-3' and 5'-aggctcgtcaga-agtct-3'  
*gapdh*: 5'-agccatcgctcagaca-3' and 5'-gcccaatacacc-aaatcc-3'

#### ChIP

ChIP assay was performed with ChIP Assay Kit (Upstate Biotechnology Inc., Lake Placid, NY) according to the manufacturer's instructions with minor modification. First, HepG2 cells were cultured in phenol red-free Opti-MEM I for 24 h for hormone depletion. Then, the cells were treated with 1  $\mu$ M DEX or 0.1% ethanol (vehicle) for the indicated time periods. After treatment, the cells were cross-linked in 1% formaldehyde for 10 min at 37 C. Cross-linking was stopped with addition of glycine to medium to a final 125 mM for 5 min at 37 C, after which the cells were rinsed with ice-cold PBS twice and harvested. Cell pellets were collected and resuspended in SDS-lysis buffer (50 mM Tris, pH 8.0; 1% SDS; 10 mM EDTA; 1  $\mu$ M 4-(2-aminoethyl)-benzenesulfonyl fluoride hydrochloride; 800 nM aprotinin; 15  $\mu$ M E-64; 20  $\mu$ M leupeptin-hemisulfate; 50  $\mu$ M bestatin; and 10  $\mu$ M pepstatin A) for 10 min at 4 C. Chromatin was sheared to an average size of 500 bp by sonication of the lysate using a Bioruptor Ultrasonicator (Cosmo-Bio, Tokyo, Japan). Lysates corresponding to  $2 \times 10^6$  cells were diluted 10-fold in ChIP dilution buffer (0.01% SDS; 1.1% Triton X-100; 1.2 mM EDTA; 16.7 mM Tris, pH 8.1; and 167 mM NaCl) and precleared with Salmon Sperm DNA/Protein A Agarose beads (Upstate Biotechnology) at 4 C for 30 min. Supernatants were then collected and incubated with 5  $\mu$ g of anti-GR polyclonal antibodies or anti-RNAPIII monoclonal antibody at 4 C overnight. To collect immune complex, Salmon Sperm DNA/Protein A Agarose beads were added and further incubated at 4 C for 1 h. The beads were then washed twice each with Low-Salt Immune Complex Wash Buffer (0.1% SDS; 1% Triton X-100; 2 mM EDTA; 20 mM Tris, pH 8.1; and 150 mM NaCl), High-Salt

Immune Complex Wash Buffer (0.1% SDS; 1% Triton X-100; 2 mM EDTA; 20 mM Tris, pH 8.1; and 500 mM NaCl), LiCl Immune Complex Wash Buffer (0.25 M LiCl; 1% Nonidet P-40; 1% deoxycholate; 1 mM EDTA; and 10 mM Tris, pH 8.1), and Tris-EDTA buffer. Protein-chromatin complex was eluted with elution buffer (10 mM dithiothreitol, 1% SDS, and 0.1 M NaHCO<sub>3</sub>), and reversal of cross-link of eluates was performed in 200 mM NaCl at 65 C for 6 h, after which proteins were digested with proteinase K at 45 C for 1 h. Precipitated DNA fragments were recovered by QIAquick DNA purification kit (QIAGEN, Chatsworth, CA) and quantified with qRT-PCR using appropriate primer sets. Sequences of primers used in this study are shown below:

*sgk1* -1238: 5'-acctctcactcagctgttctgg-3' and *sgk1* -982: 5'-caagcaaggctgaaataac-3' for GR  
*sgk1* -173: 5'-cctctcaatgggggacagaac-3' and *sgk1* +85: 5'-ccttagcagcctcagctgttca-3' for RNAPII  
*atp1a1* -732: 5'-cgccctcagatctcatt-3' and *atp1a1* -447: 5'-ggactcagggatctgga-3' for GR  
*atp1a1* +158: 5'-ccctagctcctccacttg-3' and *atp1a1* +239: 5'-tcgtggagaatcagagaga-3' for RNAPII

#### FLAG-Affinity Purification

HeLa cells ( $2.5 \times 10^6$ ) were transfected with 4  $\mu$ g of pFLAG-CMV2-derived expression plasmids. After 4 h, media were replaced with DMEM supplemented with 10% fetal calf serum. After 32 h, cells were lysed in lysis buffer [50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1% (wt/vol) Nonidet P-40, 1  $\mu$ M dithiothreitol, 0.5  $\mu$ M phenylmethylsulfonyl fluoride], centrifuged at  $20,000 \times g$  for 20 min. Supernatant was diluted in FAR buffer [16.7 mM Tris-HCl (pH 8.0), 50 mM NaCl, 0.33% (wt/vol) Nonidet P-40, 0.33  $\mu$ M dithiothreitol, 0.17  $\mu$ M phenylmethylsulfonyl fluoride], applied to anti-FLAG M2-agarose beads (Sigma-Aldrich), incubated for 2 h at room temperature. The beads were washed three times with FAR buffer. Bound proteins were eluted with SDS-sample loading buffer and subjected to Western blot analysis using anti-FLAG peptide, anti-CycT1, and anti-CDK9 antibodies.

#### Luciferase Assay

COS7 cells ( $1 \times 10^6$ ) were transfected with 2  $\mu$ g of reporter plasmids (p2xGRE-LUC or p3xPPARRE-LUC), 2.5 ng of expression plasmids for the receptors (pCMX-6His-GR or pCMX-6His-PPAR $\gamma$ ), and pCMV-HA-derived HEXIM1 expression plasmids. After 4 h, media were replaced with fresh phenol red-free Opti-MEM I, and infected with recombinant adenoviruses at multiplicity of infection (MOI) of 100. After 20 h, cells were treated with 100 nM DEX, 100 nM TGZ or vehicle (0.1% ethanol), and further cultured for 18 h. Cells were lysed in Cell Culture Lysis Reagent (Promega Corp., Madison, WI), and cellular luciferase activity was measured by using Luciferase Assay System (Promega). Relative light units were normalized to the protein amounts determined with BCA Protein Assay Reagent (Pierce Chemical Co., Rockford, IL).

#### Acknowledgments

We thank Drs. Q. Zhou and E. A. Jansson for material transfer; Drs. H. Iba, S. Tokudome, and members of the Morimoto laboratory for help; Ms. Y. Nagafuji and Ms. J. Suzuki for clerical work; and Ms. T. Maruyama for diligent technical assistance.

Received March 28, 2008. Accepted September 12, 2008.

Address all correspondence and requests for reprints to: Hirotohi Tanaka, Division of Clinical Immunology, Advanced



Clinical Research Center, Institute of Medical Science, University of Tokyo, 4-6-1, Shirokanedai, Minato-ku, Tokyo 108-8639, Japan. E-mail: hirotnk@ims.u-tokyo.ac.jp.

This work was supported in part by Grants-in-Aid for Science Research and Creative Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology, the grant for Research on Intractable Diseases from the Ministry of Health, Labor and Welfare of Japan, and Nakatomi Foundation (to H.T.).

Disclosure Statement: The authors have nothing to disclose.

## REFERENCES

- Duma D, Jewell CM, Cidlowski JA 2006 Multiple glucocorticoid receptor isoforms and mechanisms of post-translational modification. *J Steroid Biochem Mol Biol* 102:11–21
- Beato M 1989 Gene regulation by steroid hormones. *Cell* 56:335–344
- Sapolsky RM, Romero LM, Munck AU 2000 How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev* 21:55–89
- Rhen T, Cidlowski JA 2005 Antiinflammatory action of glucocorticoids—new mechanisms for old drugs. *N Engl J Med* 353:1711–1723
- Lu NZ, Cidlowski JA 2004 The origin and functions of multiple human glucocorticoid receptor isoforms. *Ann NY Acad Sci* 1024:102–123
- Lingrel JB, Orlowski J, Shull MM, Price EM 1990 Molecular genetics of Na,K-ATPase. *Prog Nucleic Acid Res Mol Biol* 38:37–89
- Bens M, Chassin C, Vandewalle A 2006 Regulation of NaCl transport in the renal collecting duct: lessons from cultured cells. *Pflügers Arch* 453:133–146
- Garty H, Palmer LG 1997 Epithelial sodium channels: function, structure, and regulation. *Physiol Rev* 77:359–396
- Bonny O, Hummler E 2000 Dysfunction of epithelial sodium transport: from human to mouse. *Kidney Int* 57:1313–1318
- Gick GG, Ismail-Beigi F, Edelman IS 1988 Thyroidal regulation of rat renal and hepatic Na,K-ATPase gene expression. *J Biol Chem* 263:16610–16618
- Ismail-Beigi F, Pressley TA, Haber RS, Gick GG, Loeb JN, Edelman IS 1988 Kinetic analysis of Na,K-activated adenosine triphosphatase induced by low external K<sup>+</sup> in a rat liver cell line. *J Biol Chem* 263:8162–8167
- Mangelsdorf DJ, Thummel C, Beato M, Herrlich P, Schutz G, Umesono K, Blumberg B, Kastner P, Mark M, Chambon P, Evans RM 1995 The nuclear receptor superfamily: the second decade. *Cell* 83:835–839
- Beato M, Herrlich P, Schutz G 1995 Steroid hormone receptors: many actors in search of a plot. *Cell* 83:851–857
- Belandia B, Parker MG 2003 Nuclear receptors: a rendezvous for chromatin remodeling factors. *Cell* 114:277–280
- Lonard DM, O'Malley B W 2007 Nuclear receptor coregulators: judges, juries, and executioners of cellular regulation. *Mol Cell* 27:691–700
- Kumar R, Thompson EB 2005 Gene regulation by the glucocorticoid receptor: structure: function relationship. *J Steroid Biochem Mol Biol* 94:383–394
- Lu NZ, Wardell SE, Burnstein KL, Defranco D, Fuller PJ, Giguere V, Hochberg RB, McKay L, Rencio JM, Weigel NL, Wilson EM, McDonnell DP, Cidlowski JA 2006 International Union of Pharmacology. LXV. The pharmacology and classification of the nuclear receptor superfamily: glucocorticoid, mineralocorticoid, progesterone, and androgen receptors. *Pharmacol Rev* 58:782–797
- Woychik NA, Hampsey M 2002 The RNA polymerase II machinery: structure illuminates function. *Cell* 108:453–463
- Kadonaga JT 2004 Regulation of RNA polymerase II transcription by sequence-specific DNA binding factors. *Cell* 116:247–257
- Phuc Le P, Friedmann JR, Schug J, Brestelli JE, Parker JB, Bochkis IM, Kaestner KH 2005 Glucocorticoid receptor-dependent gene regulatory networks. *PLoS Genet* 1:e16
- So AY, Chaivorapol C, Bolton EC, Li H, Yamamoto KR 2007 Determinants of cell- and gene-specific transcriptional regulation by the glucocorticoid receptor. *PLoS Genet* 3:e94
- Wu W, Chaudhuri S, Brickley DR, Pang D, Karrison T, Conzen SD 2004 Microarray analysis reveals glucocorticoid-regulated survival genes that are associated with inhibition of apoptosis in breast epithelial cells. *Cancer Res* 64:1757–1764
- Rogatsky I, Wang JC, Derynck MK, Nonaka DF, Khodabakhsh DB, Ha CM, Darimont BD, Garabedian MJ, Yamamoto KR 2003 Target-specific utilization of transcriptional regulatory surfaces by the glucocorticoid receptor. *Proc Natl Acad Sci USA* 100:13845–13850
- Tomlinson JW, Walker EA, Bujalska IJ, Draper N, Lavery GG, Cooper MS, Hewison M, Stewart PM 2004 11 $\beta$ -Hydroxysteroid dehydrogenase type 1: a tissue-specific regulator of glucocorticoid response. *Endocr Rev* 25:831–866
- Lonard DM, O'Malley BW 2005 Expanding functional diversity of the coactivators. *Trends Biochem Sci* 30:126–132
- Lu NZ, Cidlowski JA 2006 Glucocorticoid receptor isoforms generate transcription specificity. *Trends Cell Biol* 16:301–307
- Kusuhara M, Nagasaki K, Kimura K, Maass N, Manabe T, Ishikawa S, Aikawa M, Miyazaki K, Yamaguchi K 1999 Cloning of hexamethylene-bis-acetamide-inducible transcript, HEXIM1, in human vascular smooth muscle cells. *Biomed Res* 20:273–279
- Michels AA, Nguyen VT, Fraidl A, Labas V, Edwards M, Bonnet F, Lania L, Bensaude O 2003 MAO1 and 7SK RNA interact with CDK9/cyclin T complexes in a transcription-dependent manner. *Mol Cell Biol* 23:4859–4869
- Yik JH, Chen R, Nishimura R, Jennings JL, Link AJ, Zhou Q 2003 Inhibition of P-TEFb (CDK9/Cyclin T) kinase and RNA polymerase II transcription by the coordinated actions of HEXIM1 and 7SK snRNA. *Mol Cell* 12:971–982
- Shimizu N, Ouchida R, Yoshikawa N, Hisada T, Watanabe H, Okamoto K, Kusuhara M, Handa H, Morimoto C, Tanaka H 2005 HEXIM1 forms a transcriptionally abortive complex with glucocorticoid receptor without involving 7SK RNA and positive transcription elongation factor b. *Proc Natl Acad Sci USA* 102:8555–8560
- Sims RJ, 3rd, Belotserkovskaya R, Reinberg D 2004 Elongation by RNA polymerase II: the short and long of it. *Genes Dev* 18:2437–2448
- Malik S, Roeder RG 2005 Dynamic regulation of pol II transcription by the mammalian Mediator complex. *Trends Biochem Sci* 30:256–263
- Peterlin BM, Price DH 2006 Controlling the elongation phase of transcription with P-TEFb. *Mol Cell* 23:297–305
- Karn J 1999 Tackling Tat. *J Mol Biol* 293:235–254
- Barboric M, Nissen RM, Kanazawa S, Jabrane-Ferrat N, Peterlin BM 2001 NF- $\kappa$ B binds P-TEFb to stimulate transcriptional elongation by RNA polymerase II. *Mol Cell* 8:327–337
- Giraud S, Hurlstone A, Avril S, Coqueret O 2004 Implication of BRG1 and cdk9 in the STAT3-mediated activation of the p21/waf1 gene. *Oncogene* 23:7391–7398

37. Boehm AK, Saunders A, Werner J, Lis JT 2003 Transcription factor and polymerase recruitment, modification, and movement on dhspp70 in vivo in the minutes following heat shock. *Mol Cell Biol* 23:7628–7637
38. Tian Y, Ke S, Chen M, Sheng T 2003 Interactions between the aryl hydrocarbon receptor and P-TEFb. Sequential recruitment of transcription factors and differential phosphorylation of C-terminal domain of RNA polymerase II at cyp1a1 promoter. *J Biol Chem* 278:44041–44048
39. Zhou Q, Yik JH 2006 The Yin and Yang of P-TEFb regulation: implications for human immunodeficiency virus gene expression and global control of cell growth and differentiation. *Microbiol Mol Biol Rev* 70:646–659
40. Dey A, Chao SH, Lane DP 2007 HEXIM1 and the control of transcription elongation: from cancer and inflammation to AIDS and cardiac hypertrophy. *Cell Cycle* 6:1856–1863
41. Ouchida R, Kusuhara M, Shimizu N, Hisada T, Makino Y, Morimoto C, Handa H, Ohsuzu F, Tanaka H 2003 Suppression of NF- $\kappa$ B-dependent gene expression by a hexamethylene bisacetamide-inducible protein HEXIM1 in human vascular smooth muscle cells. *Genes Cells* 8:95–107
42. Wittmann BM, Wang N, Montano MM 2003 Identification of a novel inhibitor of breast cell growth that is down-regulated by estrogens and decreased in breast tumors. *Cancer Res* 63:5151–5158
43. Haaland RE, Herrmann CH, Rice AP 2005 siRNA depletion of 7SK snRNA induces apoptosis but does not affect expression of the HIV-1 LTR or P-TEFb-dependent cellular genes. *J Cell Physiol* 205:463–470
44. Montano MM, Doughman YQ, Deng H, Chaplin L, Yang J, Wang N, Zhou Q, Ward NL, Watanabe M 2008 Mutation of the HEXIM1 gene results in defects during heart and vascular development partly through downregulation of vascular endothelial growth factor. *Circ Res* 102:415–422
45. Wittmann BM, Fujinaga K, Deng H, Ogba N, Montano MM 2005 The breast cell growth inhibitor, estrogen down regulated gene 1, modulates a novel functional interaction between estrogen receptor alpha and transcriptional elongation factor cyclin T1. *Oncogene* 24:5576–5588
46. Li Q, Cooper JJ, Altwerger GH, Feldkamp MD, Shea MA, Price DH 2007 HEXIM1 is a promiscuous double-stranded RNA-binding protein and interacts with RNAs in addition to 7SK in cultured cells. *Nucleic Acids Res* 35:2503–2512
47. Huang F, Wagner M, Siddiqui MA 2004 Ablation of the CLP-1 gene leads to down-regulation of the HAND1 gene and abnormality of the left ventricle of the heart and fetal death. *Mech Dev* 121:559–572
48. Yoshikawa N, Shimizu N, Sano M, Ohnuma K, Iwata S, Hosono O, Fukuda K, Morimoto C, Tanaka H 2008 Role of the hinge region of glucocorticoid receptor for HEXIM1-mediated transcriptional repression. *Biochem Biophys Res Commun* 371:44–49
49. Shull MM, Pugh DG, Lingrel JB 1990 The human Na, K-ATPase  $\alpha$  1 gene: characterization of the 5'-flanking region and identification of a restriction fragment length polymorphism. *Genomics* 6:451–460
50. Itani OA, Liu KZ, Cornish KL, Campbell JR, Thomas CP 2002 Glucocorticoids stimulate human sgk1 gene expression by activation of a GRE in its 5'-flanking region. *Am J Physiol Endocrinol Metab* 283:E971–E979
51. Kohoutek J, Blazek D, Peterlin BM 2006 Hexim1 sequesters positive transcription elongation factor b from the class II transactivator on MHC class II promoters. *Proc Natl Acad Sci USA* 103:17349–17354
52. Aida M, Chen Y, Nakajima K, Yamaguchi Y, Wada T, Handa H 2006 Transcriptional pausing caused by NELF plays a dual role in regulating immediate-early expression of the junB gene. *Mol Cell Biol* 26:6094–6104
53. Ishikawa T, Igarashi T, Hata K, Fujita T 1999 c-fos induction by heat, arsenite, and cadmium is mediated by a heat shock element in its promoter. *Biochem Biophys Res Commun* 254:566–571
54. Ni Z, Schwartz BE, Werner J, Suarez JR, Lis JT 2004 Coordination of transcription, RNA processing, and surveillance by P-TEFb kinase on heat shock genes. *Mol Cell* 13:55–65
55. Yik JH, Chen R, Pezda AC, Samford CS, Zhou Q 2004 A human immunodeficiency virus type 1 Tat-like arginine-rich RNA-binding domain is essential for HEXIM1 to inhibit RNA polymerase II transcription through 7SK snRNA-mediated inactivation of P-TEFb. *Mol Cell Biol* 24:5094–5105
56. Starr DB, Matsui W, Thomas JR, Yamamoto KR 1996 Intracellular receptors use a common mechanism to interpret signaling information at response elements. *Genes Dev* 10:1271–1283
57. Schulte A, Czudnochowski N, Barboric M, Schonichen A, Blazek D, Peterlin BM, Geyer M 2005 Identification of a cyclin T-binding domain in Hexim1 and biochemical analysis of its binding competition with HIV-1 Tat. *J Biol Chem* 280:24968–24977
58. Dames SA, Schonichen A, Schulte A, Barboric M, Peterlin BM, Grzesiek S, Geyer M 2007 Structure of the Cyclin T binding domain of Hexim1 and molecular basis for its recognition of P-TEFb. *Proc Natl Acad Sci USA* 104:14312–14317
59. Dulac C, Michels AA, Fraidi A, Bonnet F, Nguyen VT, Napolitano G, Lania L, Bensaude O 2005 Transcription-dependent association of multiple positive transcription elongation factor units to a HEXIM multimer. *J Biol Chem* 280:30619–30629
60. Barboric M, Kohoutek J, Price JP, Blazek D, Price DH, Peterlin BM 2005 Interplay between 7SK snRNA and oppositely charged regions in HEXIM1 direct the inhibition of P-TEFb. *EMBO J* 24:4291–4303
61. Blazek D, Barboric M, Kohoutek J, Oven I, Peterlin BM 2005 Oligomerization of HEXIM1 via 7SK snRNA and coiled-coil region directs the inhibition of P-TEFb. *Nucleic Acids Res* 33:7000–7010
62. Gurumurthy M, Tan CH, Ng R, Zeiger L, Lau J, Lee J, Dey A, Philp R, Li Q, Lim TM, Price DH, Lane DP, Chao SH 2008 Nucleophosmin interacts with HEXIM1 and regulates RNA polymerase II transcription. *J Mol Biol* 378:302–317
63. Kino T, De Martino MU, Charmandari E, Mirani M, Chrousos GP 2003 Tissue glucocorticoid resistance/hypersensitivity syndromes. *J Steroid Biochem Mol Biol* 85:457–467
64. Tanaka H, Yoshikawa N, Shimizu N, Morimoto C 2004 Selective modulation of glucocorticoid receptor toward development of novel anti-inflammation: lessons from a phenylpyrazolo-steroid cortivazol. *Mod Rheumatol* 14:347–355
65. Yoshikawa N, Yamamoto K, Shimizu N, Yamada S, Morimoto C, Tanaka H 2005 The distinct agonistic properties of the phenylpyrazolo-steroid cortivazol reveal inter-domain communication within the glucocorticoid receptor. *Mol Endocrinol* 19:1110–1124

*Molecular Endocrinology* is published monthly by The Endocrine Society (<http://www.endo-society.org>), the foremost professional society serving the endocrine community.

## Activation of Mitochondrial Biogenesis by Hormesis

Motoaki Sano, Keiichi Fukuda

Mitochondria play a major role in oxidative energy production, reduction-oxidation reaction (redox) control and calcium homeostasis. Although mitochondria contain DNA with mitochondrial-specific genes, most mitochondrial proteins are encoded by the nDNA, synthesized in the cytosol, and imported into mitochondria. The expression of nuclear genes that encode mitochondrial proteins that function in metabolic pathways such as the trichloroacetic acid cycle (TCA), oxidative phosphorylation, heme synthesis, and in mitochondrial DNA replication and transcription (eg, mitochondrial transcription factor A [Tfam]), is coordinately regulated by the transcriptional coactivators PPAR $\gamma$  coactivator (PGC)-1 $\alpha$  and PGC-1 $\beta$  through activation of nuclear respiratory factor (NRF)-1 and NRF-2.<sup>1</sup>

In their recent publications, Piantadosi et al provided insight into the mechanisms underlying the interaction between mitochondria-derived reactive oxygen species (ROS) signaling and mitochondrial biogenesis. First, lipid hydroperoxide regulates Tfam expression through phosphorylation of NRF-1 via Akt activation, which promotes nuclear translocation of NRF-1 and binding to the Tfam promoter.<sup>2</sup> Second, carbon monoxide (CO) induced mitochondrial biogenesis via activation of Akt/PKB and guanylate cyclase, which augmented gene and protein expression of NRF-1 and NRF-2, PGC-1 $\alpha$ , and TFAM.<sup>3</sup> CO-induced mitochondrial ROS result in the activation of AKT. Third, the anthracycline anticancer agent doxorubicin suppresses the nuclear program for mitochondrial biogenesis, and its associated intrinsic antiapoptosis proteins, leading to severe mitochondrial DNA (mtDNA) depletion and apoptosis. CO inhalation or heme oxygenase (Hmo) overexpression prevented doxorubicin-induced mtDNA depletion and apoptosis via activation of AKT and guanylate cyclase.<sup>4</sup> Lastly, new work in this issue of *Circulation Research*<sup>5</sup> sheds light on the role of NF-E2-related factor (Nrf2) as a key transcriptional regulator in mitochondrial ROS-dependent induction of NRF-1 mRNA.

There is increasing evidence to suggest that ROS may be a double-edged sword: although they can be toxic to cells, they may also play an important role in cell signaling involved in the antioxidant defense network. ROS are generated from many sources including the Nox family of NADPH oxidases,

xanthine oxidase, and mitochondria, where ROS are produced as a byproduct of oxidative energy production. ROS are very unstable and cannot penetrate lipid membranes; they are therefore retained within the compartment in which they are produced. However, ROS can attack neighboring polyunsaturated fatty acids of the membrane and trigger a chain reaction of lipid peroxidation, resulting in the generation of lipid hydroperoxides and  $\alpha$ ,  $\beta$ -unsaturated aldehydes, such as 4-hydroxy-2-nonenal (4-HNE) (Figure). They are highly electrophilic and react with biomolecules, such as proteins and nucleic acids, generating various adducts. By virtue of their increased chemical stability, these lipid peroxidation products can diffuse greater distances compared with their precursor ROS and can propagate and amplify oxidative injury. Thus, lipid peroxidation products have been implicated in the development and progression of a variety of pathological events such as oxidation of LDL, atherosclerosis, ischemia/reperfusion injury, Alzheimer's disease, cancers, and cell senescence.

However cells are able to sense macromolecular damage and counteract stress-induced damage to reestablish homeostasis. Electrophilic lipid peroxidation products can trigger a cascade of stress resistant pathways in both a tissue- and cell type-specific manner. The induction of stress-protective mechanisms by stress is referred to as "stress-response hormesis."<sup>6</sup> The principle of stress-response hormesis can be seen in many contexts. For example, the ninja, a group of spies and assassins in feudal Japan, were known to regularly take sublethal doses of poison to build their capacity to detoxify xenobiotics and thus protect themselves against assassination with poison. In cell culture, 4-HNE kills cells at a high dose, whereas pretreatment of cells with low-dose 4-HNE upregulates endogenous antioxidant and phase II enzymes, conferring greater tolerance against subsequent oxidative insult.<sup>7</sup> An effect of stress-response hormesis may also be seen in clinical studies that have tested antioxidant supplements for prevention of cardiovascular events<sup>8</sup> and cancers<sup>9</sup> based on the principle that they should prevent oxidative stress-induced macromolecular damage. In both clinical studies, antioxidant supplements may have failed to identify a beneficial effect because this inevitably attenuates the cell-signaling pathways necessary for protection against oxidative stress and reestablishment of redox homeostasis.<sup>8</sup>

Following the induction of oxidative stress, 2 basic leucine zipper transcription factors, Nrf2 and activating transcription factor (ATF)4, are activated at the posttranscriptional level and induce the expression of genes encoding proteins that function as antioxidants and enzymes involved in phase II detoxification and glutathione biosynthesis. Under non-stressed conditions, Nrf2 is tethered in the cytoplasm by Keap1. This complex directs Nrf2 polyubiquitination and degradation. On oxidative stress, Nrf2 is liberated from

The opinions expressed in this editorial are not necessarily those of the editors or of the American Heart Association.


From the Department of Regenerative Medicine and Advanced Cardiac Therapeutics, Keio University School of Medicine, Tokyo, Japan.

Correspondence to Motoaki Sano, Department of Regenerative Medicine and Advanced Cardiac Therapeutics, Keio University School of Medicine, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582. E-mail msano@sc.itc.keio.ac.jp

(*Circ Res*. 2008;103:1191-1193.)

© 2008 American Heart Association, Inc.

*Circulation Research* is available at <http://circres.ahajournals.org>  
DOI: 10.1161/CIRCRESAHA.108.189092



**Figure.** The generation of ROS and the subsequent oxidative modification of biomolecules are inevitable events in aerobic organisms. The polyunsaturated fatty acids in membrane lipids are particularly vulnerable to ROS attack and they undergo peroxidation. These lipid peroxidation products form protein and DNA adducts and have biphasic properties, in that high doses cause overt toxicity, whereas low doses interact with genetic signaling systems that upregulate gene expression to counteract stressor challenges and to re-establish homeostasis. LOOH indicates lipid hydroperoxides; LCHO, lipid aldehydes.

Keap1 and enters the nucleus, where it can form a heterodimer with the small Maf transcription factor Nrf2 to stimulate the expression of antioxidant response element-containing genes, including NAD(P)H:quinone oxidoreductase, heme-oxygenase 1,  $\gamma$ -glutamylcysteine synthetase, glutathione S-transferase, glutathione peroxidase, glutathione reductase, cysteine glutamate transporter, and multidrug resistance-associated protein 1.<sup>11</sup> Oxidative stress leads to the phosphorylation of the  $\alpha$  subunit of translation initiation factor 2 (eIF2 $\alpha$ ). Phosphorylation of eIF2 $\alpha$  inhibits general protein synthesis but specifically upregulates translation of ATF4. ATF4 forms homodimers and heterodimers with members of the AP-1 and C/EBP family of proteins to regulate the expression of genes involved in amino acid metabolism which provide the precursor amino acids necessary for glutathione biosynthesis, such as phosphoserine amino transferase, phosphoserine phosphatase, cystathione  $\gamma$ -lyase, and methylenetetrafolate dehydrogenase.<sup>12</sup> Thus, Nrf2 and ATF4 coordinately regulate glutathione biosynthesis and the glutathione redox cycle.

Intense muscular contractile activity by exercise results in oxidative stress, as indicated by altered muscle and blood glutathione concentrations and increases in protein, DNA, and lipid peroxidation. Interestingly, it was recently reported that excess vitamin C supplements decrease training efficiency via the reduction of the exercise-induced expression of PGC-1, NRF-1, and Tfam.<sup>13</sup> This observation further suggests that ROS cannot only be considered to be toxic byproducts; they also play an important role in the cell signaling that regulates expression of genes involved in mitochondrial biogenesis. Piantadosi et al first demonstrated a role for Nrf2 in ROS-mediated induction of NRF-1. The

NRF-1 promoter contains multiple antioxidant response element motifs and mitochondrial-derived ROS enhance Nrf2 binding to the NRF-1 promoter via AKT-mediated derepression of Nrf2 nuclear translocation. In the heart, however, the role of Nrf2 signaling in the basal expression, as well as the induction of antioxidants in pathological circumstances remains unclear.

Mitochondrial DNA copy number and mitochondrial gene expression are reduced in heart failure. Not surprisingly, concomitant downregulation of PGC-1 $\alpha$ , NRF-1, and Tfam in the failing heart is observed.<sup>14</sup> The mechanism by which pathophysiological cues downregulate PGC-1/NRF-1/Tfam expression have only begun to be resolved,<sup>15</sup> but it is tempting to speculate that rescue of PGC-1/NRF-1/Tfam expression may have beneficial effects on cardiac function. Indeed, transgenic overexpression of Tfam in the heart ameliorates the decrease in mitochondrial DNA copy number and attenuates left ventricular remodeling and failure after myocardial infarction.<sup>16</sup>

The novel concept that mitochondrial biogenesis seems to be triggered by mitochondrial ROS generation is intriguing. To move present knowledge toward more general applicability, the physiological and pathological relevance of mitochondrial ROS-mediated transcriptional and posttranscriptional activation of NRF-1 via AKT, in the setting of postneonatal normal growth, exercise-challenged, pressure-challenged, ischemic, and failing heart need to be clarified.

## Sources of Funding

This work was supported by a PRESTO (Metabolism and Cellular Function) grant from the Japanese Science and Technology Agency to M. Sano.

## Disclosures

None.

## References

- Kelly DP, Scarpulla RC. Transcriptional regulatory circuits controlling mitochondrial biogenesis and function. *Genes Dev*. 2004;18:357-368.
- Piantadosi CA, Suliman HB. Mitochondrial transcription factor A induction by redox activation of nuclear respiratory factor 1. *J Biol Chem*. 2006;281:324-333.
- Suliman HB, Carraway MS, Tatro LG, Piantadosi CA. A new activating role for CO in cardiac mitochondrial biogenesis. *J Cell Sci*. 2007;120:299-308.
- Suliman HB, Carraway MS, Ali AS, Reynolds CM, Wely-Wolf KE, Piantadosi CA. The CO/PO system reverses inhibition of mitochondrial biogenesis and prevents murine doxorubicin cardiomyopathy. *J Clin Invest*. 2007;117:3730-3741.
- Piantadosi CA, Carraway MS, Babiker A, Suliman HB. Heme oxygenase-1 regulates cardiac mitochondrial biogenesis via Nrf2-mediated transcriptional control of nuclear respiratory factor-1. *Circ Res*. 2008;103:1232-1240.
- Gems D, Partridge L. Stress-response hormesis and aging: "that which does not kill us makes us stronger." *Cell Metab*. 2008;7:200-203.
- Uchida K, Shiraishi M, Naito Y, Torii Y, Nakamura Y, Osawa T. Activation of stress signaling pathways by the end product of lipid peroxidation: 4-Hydroxy-2-nonenal is a potential inducer of intracellular peroxide production. *J Biol Chem*. 1999;274:2234-2242.
- Homarbakhti S, Schachter M. Vitamins and cardiovascular disease. *Br J Nutr*. 2008;1-19.
- Bjelakovic G, Nikolova D, Simonetti RG, Glud C. Antioxidant supplements for prevention of gastrointestinal cancers: a systematic review and meta-analysis. *Lancet*. 2004;364:1219-1228.

10. Hayashida K, Sano M, Ohsawa I, Shinmura K, Tamaki K, Kimura K, Endo J, Katayama T, Kawamura A, Kohsaka S, Makino S, Ohta S, Ogawa S, Fukuda K. Inhalation of hydrogen gas reduces infarct size in the rat model of myocardial ischemia-reperfusion injury. *Biochem Biophys Res Commun*. 2008;373:30-35.
11. Motohashi H, Yamamoto M. Nrf2-Keap1 defines a physiologically important stress response mechanism. *Trends Mol Med*. 2004;10:549-557.
12. Harding HP, Zhang Y, Zeng H, Novoa I, Lu PD, Calfon M, Sadri N, Yun C, Popko B, Paules R, Stojdl DF, Bell JC, Hettmann T, Leiden JM, Ron D. An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol Cell*. 2003;11:619-633.
13. Gomez-Cabrera MC, Domenech E, Romagnoli M, Arduini A, Borrás C, Pallardo FV, Sastre J, Vina J. Oral administration of vitamin C decreases muscle mitochondrial biogenesis and hampers training-induced adaptations in endurance performance. *Am J Clin Nutr*. 2008;87:142-149.
14. Garnier A, Fortin D, Delomenie C, Monken I, Veksler V, Ventura-Clapier R. Depressed mitochondrial transcription factors and oxidative capacity in rat failing cardiac and skeletal muscles. *J Physiol*. 2003;551:491-501.
15. Sano M, Wang SC, Shirai M, Scaglia F, Xie M, Sakai S, Tanaka T, Kulkarni PA, Barger PM, Youker KA, Taffet GE, Hamamori Y, Michael LH, Craigen WJ, Schneider MD. Activation of cardiac Cdk9 represses PGC-1 and confers a predisposition to heart failure. *EMBO J*. 2004;23:3559-3569.
16. Ikeuchi M, Matsusaka H, Kang D, Matsushima S, Ide T, Kubota T, Fujiwara T, Hamasaki N, Takeshita A, Sunagawa K, Tsutsui H. Overexpression of mitochondrial transcription factor  $\alpha$  ameliorates mitochondrial deficiencies and cardiac failure after myocardial infarction. *Circulation*. 2005;112:683-690.

KEY WORDS: lipid metabolites ■ mitochondria ■ oxidative stress ■ redox ■ signaling pathways ■ hormesis ■ stress response

**Ligand-based gene expression profiling reveals novel roles of glucocorticoid receptor in cardiac metabolism**

Noritada Yoshikawa<sup>1,2</sup>, Masao Nagasaki<sup>3</sup>, Motoaki Sano<sup>4,6</sup>, Satori Tokudome<sup>4</sup>, Kazuko Ueno<sup>3</sup>, Noriaki Shimizu<sup>1</sup>, Seiya Imoto<sup>3</sup>, Satoru Miyano<sup>3</sup>, Makoto Suematsu<sup>5</sup>, Keiichi Fukuda<sup>4</sup>, Chikao Morimoto<sup>1,2</sup>, Hirotohi Tanaka<sup>1,2</sup>

N.Y. and M.N. equally contributed to this work and both should be considered as first authors.

1, Division of Clinical Immunology, Advanced Clinical Research Center, 2, Research Hospital, 3, Laboratory of DNA Information Analysis, Human Genome Center, Institute of Medical Science, University of Tokyo, Tokyo, Japan, 4, Department of Regenerative Medicine and Advanced Cardiac Therapeutics, 5, Department of Biochemistry and Integrative Medical Biology, Keio University School of Medicine, Tokyo, Japan, 6, Precursory Research for Embryonic Science and Technology (PRESTO), Japan Science and Technology Agency, Saitama, Japan

**Running head: Novel Roles of GR in Cardiac Metabolism**

Address for correspondence to: Hirotohi Tanaka, M.D., Ph.D., Division of Clinical Immunology, Advanced Clinical Research Center, Institute of Medical Science, University of Tokyo, 4-6-1, Shirokanedai, Minato-ku, Tokyo 108-8639, JAPAN

Telephone: +81-3-5449-5547

Fax: +81-3-6409-2405

e-mail: hirotnk@ims.u-tokyo.ac.jp

**Abstract**

Recent studies have documented various roles of adrenal corticosteroid signaling in cardiac physiology and pathophysiology. It is known that glucocorticoids and aldosterone are able to bind glucocorticoid receptor (GR) and mineralocorticoid receptor (MR), and these ligand-receptor interactions are redundant. It, therefore, has been impossible to delineate how these nuclear receptors couple with corticosteroid ligands and differentially regulate gene expression for operation of their distinct functions in the heart. Here, to particularly define the role of GR in cardiac muscle cells, we applied ligand-based approach involving GR-specific agonist cortivazol (CVZ) and GR antagonist RU486, and performed microarray analysis using rat neonatal cardiomyocytes. We indicated that glucocorticoids appear to be a major determinant of GR-mediated gene expression when compared with aldosterone. Moreover, expression profiles of these genes highlighted numerous roles of glucocorticoids in various aspects of cardiac physiology. At first, we identified that glucocorticoids, via GR, induce mRNA and protein expression of a transcription factor Kruppel-like factor 15 and its downstream target genes including branched-chain aminotransferase 2, a key enzyme for amino acid catabolism in the muscle. CVZ treatment or overexpression of KLF15 decreased cellular branched-chain amino acid concentrations and introduction of siRNA against KLF15 cancelled these CVZ actions in cardiomyocytes. Second, glucocorticoid-GR signaling promoted gene expression of the enzymes involved in the prostaglandin biosynthesis including cyclooxygenase-2 and phospholipase A2 in cardiomyocytes. Together, we may conclude that GR signaling should have distinct roles for maintenance of cardiac function, for example, in amino acid catabolism and prostaglandin biosynthesis in the heart (248 words).

Key words: endocrinology, cardiovascular system, KLF15, COX-2, and PLA2

## Introduction

Glucocorticoid hormones are essential for homeostatic regulation and physiological maintenance of a variety of organ functions. Concerning the heart, numerous observations have suggested that glucocorticoids as well as aldosterone (ALD) have been shown to exert direct effects on cardiomyocytes and help maintain various cardiac functions. For example, it is shown that a synthetic glucocorticoid dexamethasone (DEX) significantly increases the L-type  $\text{Ca}^{2+}$  currents (51) and inhibits inducible nitric oxide synthase activity in rat cardiomyocytes (42). Moreover, DEX treatment enhances the development of contractile tension and increases contraction and relaxation velocities in cardiac muscle (35). The decrease in contractile force of rat papillary muscle induced by adrenalectomy is prevented by DEX treatment (27) by modulating membrane  $\text{Ca}^{2+}$  transport and  $\text{K}^{+}$  channels (33, 35, 50, 51). Short-term treatment with DEX has been shown to decrease resting heart rate in healthy human volunteers (5). It, thus, is apparent that glucocorticoids play essential roles in regulation of cardiac electrical and mechanical activities. On the other hand, numerous studies have documented the pathological consequences and deleterious effects of abnormal or excessive glucocorticoid signalings. Not only hypercortisolemia in patients with Cushing's syndrome but also the chronic therapeutic use of glucocorticoids is associated with several side effects, including adverse cardiovascular events, such as hypertension and left ventricular hypertrophy (48). Glucocorticoid excess also induces metabolic syndrome with hyperglycemia, dyslipidemia, and obesity, which is associated with early and progressive atherosclerosis, contributing to a cluster of cardiovascular risk factors, including heart failure (48). Moreover, several clinical studies have documented the distinct role of glucocorticoids in the prognosis of cardiac diseases; for example, rheumatoid factor-positive but not negative patients with rheumatoid arthritis were at increased risk of cardiovascular events following



exposure to glucocorticoids (8), and, in patients with chronic heart failure, higher serum levels of cortisol and ALD were independent predictors of increased mortality risk (16).

Glucocorticoids and mineralocorticoids bind to nuclear receptors-glucocorticoid receptor (GR) and mineralocorticoid receptor (MR)-both of which are transcription factors and expressed in cardiomyocytes. However, the role of these receptors in cardiac physiology remains elusive. Indeed, the ligand/receptor interactions are complex, as both ALD and glucocorticoids can activate cardiac MR, thereby directly affecting heart function (48). Some of the cardiac or peripheral effects of glucocorticoids may be mediated at least in part by MR activation. In "classical" aldosterone target cells (i.e., kidney and colon), MR is protected from illicit occupation by glucocorticoids due to the presence of 11 $\beta$ -hydroxysteroid dehydrogenase type II (11 $\beta$ HSD2), an enzyme that converts cortisol (human)/corticosterone (COR, rodents) into inactive metabolites. Cardiomyocytes belong to the so-called "nonclassical" ALD target tissues that express both GR and MR, but not 11 $\beta$ HSD2. In cardiomyocytes, thus, MR is not protected from occupancy by glucocorticoids and is not ALD selective. Taking into account that circulating cortisol/COR levels are at least 100-fold higher than those of ALD, and that MR has the same affinity for ALD and glucocorticoids, MR, as well as GR, may be permanently occupied by glucocorticoids and glucocorticoid effects could be mediated by both GR and MR (48). The recent advent of microarray and other technologies has facilitated the identification of a number of glucocorticoid-regulated genes (1, 20, 34, 36, 45) and it becomes apparent that the profile of those glucocorticoid-target genes differs according to the cell types and the mode of interaction with ligands (49). However, because of the redundancy of the ligand-receptor interaction, not a single study could clearly differentiate target genes for cardiac GR and MR. Recently a transgenic mouse model with conditionally inducible cardiac-specific expression of human GR was generated to preclude secondary effects due to general glucocorticoid-induced

alterations and to investigate the specific role of GR in cardiomyocytes, and electrophysiological phenotyping indicated that cardiac GR overexpression resulted in conduction defects, with high-degree atrio-ventricular block (39). These results strongly support such an idea that GR has as yet unknown but essential roles in the heart. It, therefore, is important to delineate how these nuclear receptors, especially GR, differentially couple with ligands and regulate gene expression for operation of their distinct functions in cardiomyocytes.

We previously reported that a synthetic glucocorticoid cortivazol (CVZ) could be extremely specific for GR and does not crossreact with MR (52, 53). Given this, we indicated that ligand-based approach involving CVZ and GR antagonist RU486 might be applied to define the role of GR in non-classical ALD target tissues. In the present study, we performed microarray analysis based on this ligand-based approach and differentially characterized corticosteroid target genes, and the distinct role of GR in cardiomyocytes was discussed.

## **Materials and Methods**

### **Reagents and antibodies**

CVZ was kindly gifted from Sanofi-Aventis (Paris, France). COR, ALD, IL-1 $\beta$ , lipopolysaccharide, estradiol, progesterone, and RU486 were purchased from Sigma (St. Louis, MO). MG-132 was purchased from Calbiochem (San Diego, CA). Other reagents were from Nacalai Tesque (Kyoto, Japan) unless otherwise specified. Anti-GR (sc-1004), anti-cyclooxygenase-2 (COX-2, sc-1747), and anti-KLF15 (sc-34827) antibodies were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Anti- $\alpha$ -actinin (A7811) and anti-FLAG (F1804) antibodies were obtained from Sigma.

**Plasmids, siRNA oligonucleotides, and recombinant adenoviruses**

To construct the expression plasmids for FLAG-tagged rat GR and MR, either full-length cDNAs for rat GR or MR were inserted into p3xFLAG-CMV10 vector (Sigma). The glucocorticoid response element (GRE)-driven reporter plasmid p2xGRE-LUC was described previously (52). siRNA oligonucleotides against rat GR (*Silencer*<sup>®</sup> Pre-designed siRNA ID: 199951) and control siRNA (*Silencer*<sup>®</sup> Negative control siRNA #1: 07606954A) were purchased from Ambion (Austin, TX). siRNA oligonucleotides against rat KLF15 (*Stealth*<sup>™</sup> Select RNAi RSS340443) was purchased from Invitrogen (Carlsbad, CA). Recombinant adenoviruses encoding FLAG-tagged rat KLF15 (Ad-KLF15) and Cre-recombinase (Ad-Cre) were generated by using Adenovirus Cre/loxP-regulated Expression Vector Set (TaKaRa, Otsu, Japan) as manufacturer's instructions and previously described (44). Recombinant adenoviruses prepared from 293 cells were purified with Virakit AdenoMini-24 (Virapur, San Diego, CA) and titrated using Adeno-X Rapid Titer Kit (TaKaRa).

**Cell culture**

COS7 cells were obtained from RIKEN Cell Bank (Tsukuba, Japan) and cultured in Dulbecco's modified Eagle's medium (DMEM, Invitrogen) supplemented with 10% fetal calf serum (FCS, SIGMA-ALDRICH, St. Louis, MO) and antibiotics in a humidified atmosphere at 37°C with 5% CO<sub>2</sub>. Primary cultures of cardiomyocytes were prepared as described previously (40). In brief, the ventricles of 1-day-old neonatal Wistar rats (CLEA Japan, Tokyo, Japan) were dissociated in 0.03% trypsin, 0.03% collagenase, and 20 µg/mL of DNase I. The cardiomyocytes and cardiac fibroblasts were separately prepared on the basis of their differential adhesiveness. Attached cells (mostly cardiac fibroblasts) were subcultured twice to deplete cardiomyocytes, and the third passage cells were used. Cardiomyocytes were

seeded at a density of  $1 \times 10^5$  cells/cm<sup>2</sup> on gelatin-coated dishes and grown in medium 199/DMEM (Invitrogen) supplemented with 10% FCS and antibiotics in a humidified atmosphere at 37°C with 5% CO<sub>2</sub>. Concerning animal experiments, all procedures and protocols were approved by the Animal Care and Use Committee of Keio University.

### **Immunofluorescence**

FLAG-tagged rat GR or MR expressing COS7 cells were plated onto glass coverslips in a 6-well plate. Fixed and permeabilized cells were blocked with blocking buffer (3% BSA and 0.1% Triton-X in Tris-buffered saline). The cells were then stained with primary antibodies against FLAG (1:500) for 1 h at room temperature, and then, secondary antibodies conjugated with Alexa Fluor 488 (1:500, Invitrogen) were applied for 1 h at room temperature. The stained cells were observed by confocal laser scanning microscopy (LSM510; Carl Zeiss, Jena, Germany) with appropriate emission filters..

### **Western blotting**

Whole cell extracts were prepared in NP-40 lysis buffer (150 mM NaCl, 1.0% NP-40, 50 mM Tris, pH 8.0, and protease inhibitor cocktail) and boiled in SDS sample buffer, analyzed by SDS-PAGE, and electrically transferred to a polyvinyl difluoride membrane (Millipore, Bedford, MA). Subsequently, immunoblotting was performed with anti-GR, anti- $\alpha$ -actinin, anti-KLF15, anti-FLAG, or anti-COX-2 antibodies diluted at 1:1000, followed by horseradish peroxidase-conjugated secondary antibodies (Amersham Biosciences, Buckinghamshire, UK) diluted at 1:2000. Antibody-protein complexes were visualized using the enhanced chemiluminescence method according to the manufacturer's protocol (Amersham Biosciences). Signal intensity of the band for GR relative to that for  $\alpha$ -actinin were quantified using the analysis soft ware from National Institutes of Health (NIH image