

FIG. 5. Changes in HMGB-1 protein expression in lung tissue upon administration of LPS in rats. Lung tissue was obtained from control, LPS, and LPS + landiolol group rats 12 h after LPS administration. The HMGB-1 expression was analyzed by immunohistochemistry in the three groups (hematoxylin-eosin, original magnification $\times 400$): control group (A), LPS group (B), and landiolol group (C). Western blot analysis (D) of HMGB-1 expression in lungs. β -Actin was used as a control for loading.

landiolol group compared with the LPS-only group (Fig. 2D). The wet-to-dry weight ratios obtained from the lungs of the LPS group animals were significantly elevated compared with those from control group animals (Fig. 3). Wet-to-dry weight ratios obtained from the LPS + landiolol group were comparable to the control group ratios, that is, significantly lower than those from LPS group animals.

Effects of landiolol on serum levels of TNF- α , IL-6, and HMGB-1

The cytokines IL-6 and TNF- α both showed marked peaks of expression in serum 3 h after LPS administration. Levels of both TNF- α and IL-6 were significantly lower in the LPS + landiolol group compared with LPS group, with the most marked effect seen at the 3-h time point (Fig. 4, A and B). Similarly, serum levels of HMGB-1 increased after the administration of LPS; however, the peak in expression did not occur until 12 h after treatment. This peak was significantly less prominent in LPS + landiolol group rats (Fig. 4C). Serum levels of IL-6, TNF- α , and HMGB-1 remained unchanged at all time points in the control group (data not shown).

Effects of landiolol on lung tissue

Immunohistochemical analysis revealed that the number of cells expressing HMGB-1 increased after LPS treatment alone (Fig. 5B). This increase was dramatically suppressed by landiolol cotreatment (Fig. 5C). Western blot analysis of lung tissue also showed the increase in HMGB-1 expression resulting from LPS treatment. This increase was less pronounced among LPS + landiolol group rats compared with the LPS group (Fig. 5D). We next examined NF- κ B signaling in lung tissue by determining the extent of DNA binding activity by the NF- κ B transcription factor p50/p65. Administration of LPS led to a robust increase in the DNA binding activity. However, cotreatment with landiolol significantly suppressed this increase (Fig. 6).

Effect of landiolol treatment on myocardial function in the LPS-induced systemic inflammation model

The administration of LPS was associated with a marked increase in LVEDP compared with values from control group animals. This increase was significantly suppressed in LPS + landiolol group animals (Fig. 7A). In contrast, the LPS + landiolol group showed markedly increased LVDP (Fig. 7B), LV +dP/dt_{max} (Fig. 7C), and LV -dP/dt_{min} (Fig. 7D) relative to the LPS group. Of particular note was the 12-h time point, at

which all indicated parameters were significantly improved by the cotreatment with landiolol.

DISCUSSION

In this study, we demonstrated that landiolol provided protection against lung injury and reduced the wet-to-dry weight ratio in the lung in a rat model of LPS-induced sepsis. Acute lung injury in acute respiratory distress syndrome is considered an acute inflammatory condition by many clinicians. Although the pathology of this syndrome is poorly defined, Zhang et al. (27) reported that β -adrenoceptor activation can attenuate the acute inflammatory response induced in the lung by LPS. In addition, β -blockers reduce lung injury (28). These results suggest a critical role for β -adrenoceptors in acute lung injury.

The inflammatory response leading to lung and cardiac dysfunction and failure continues to be the major problem after injury in many clinical conditions such as sepsis (29, 30). Our findings showed that serum levels of cytokines and HMGB-1 were significantly inhibited by the administration of landiolol. This is the first report of a landiolol-dependent suppression of HMGB-1 expression in the septic rat model. The results presented here build on the growing body of evidence suggesting that several cytokines, including TNF- α and IL-6, are released during the early phase of systemic inflammatory response syndrome (29, 30). TNF- α and IL-6 mediate cardiac inflammation and contractile dysfunction after sepsis (31, 32). In addition, Ueno et al. (14) reported that HMGB-1 is important in the initiation of acute lung injury.




FIG. 6. Effect of landiolol on the LPS-induced increase of specific binding of p50/p65 to DNA. Lung tissue cells were obtained from LPS and LPS + landiolol group rats 6 h after LPS administration. Nuclear fractions were prepared and subjected to a DNA binding activity assay specific for the NF- κ B (p50/p65) transcription factor. All data are expressed as the mean \pm SD. *Significant difference compared with LPS group ($P < 0.05$).




FIG. 7. Effects of landiolol treatment on heart function in rats from LPS-induced systemic inflammation model. Myocardial function in rats from the control (black bars), LPS (white bars), and LPS + landiolol (slashed bars) groups ($n = 10$ per group) was determined at the time points indicated. A, LVDP. B, LVDP. C, $+dP/dt_{max}$. D, $-dP/dt_{min}$. Data are presented as the mean \pm SD. *Significant difference compared with LPS group at 12 h ($P < 0.05$). *Significant difference compared with LPS group at 24 h ($P < 0.05$).

In this study, we analyzed the levels of HMGB-1 in serum and lung tissue in the rat model of LPS-induced lung injury. TNF- α is thought to mediate LPS-induced HMGB-1 release (21); a number of different pathways have been implicated in the inhibition of HMGB-1 secretion. The selective blockade of β 1-adrenoceptors has been shown to inhibit cytokine release in septic animals (33). We found that serum levels of TNF- α and IL-6 were significantly lower in animals cotreated with the β -blocker landiolol compared with levels found in rats treated with LPS alone.

In this study, we demonstrated that the release of inflammatory mediators can be inhibited by administration of the β 1-adrenoceptor blocker landiolol. On the other hand, previous studies indicated that β -adrenoceptor agonists inhibit the secretion of various cytokines in the LPS-induced inflammatory response (34, 35). Likewise, β 2-adrenoceptor stimulation reduces cytokine release and organ damage (36, 37). These results suggest that β 1- and β 2-adrenoceptors might play different roles in inflammation. Further studies will be required to elucidate the mechanisms of β 1- and β 2-adrenoceptor function in the regulation of inflammation.

Many signaling events lead to cytokine synthesis and release after LPS exposure. One such key event involves the transcription factor NF- κ B. Nuclear factor- κ B signaling controls tissue regeneration and early events in inflammation (38), and is a central regulator of septic shock (22). In addition, NF- κ B is markedly activated in peripheral mononuclear cells from patients with sepsis, and its activity is significantly higher in nonsurviving patients than in surviving patients (39, 40). Furthermore, numerous NF- κ B target genes are implicated in the pathogenesis of sepsis (22, 41). In this study, we demonstrated that NF- κ B activation is inhibited by administration of landiolol in the LPS-induced systemic inflammation model. The improvement of LPS-induced systemic inflammation might be attributed to the inhibition of NF- κ B activation. However, the mechanism by which landiolol administration inhibits NF- κ B remains unclear. One possibility involves free-radical scavenging. Further studies on the anti-inflammatory effects of landiolol are needed to elucidate the basis of its activity in this context.

Previous studies have demonstrated that β -blockers inhibit the activation of NF- κ B (42). We showed here that landiolol exerts such an effect in a rat model of LPS-induced lung injury. Nuclear factor- κ B is a major driver in the induction of several proinflammatory cytokine genes (22) as well as the secretion of HMGB-1 after LPS administration (43). Our

results here indicate that one mechanism by which landiolol administration suppresses cytokine and HMGB-1 secretion may be through the inhibition of NF- κ B, suggesting a critical role for β 1-adrenoceptor in NF- κ B activity in the LPS-induced shock model.

In our rat model of LPS-induced septic shock, cotreatment with landiolol prevented acute lung injury and improved cardiac function. In conclusion, our results suggest that landiolol may bring therapeutic benefit to patients with sepsis through its ability to inhibit the inflammatory response, especially by suppressing HMGB-1 expression, while improving lung injury and cardiac function. Further studies are needed to determine whether treatment with landiolol after the onset of sepsis is similarly efficacious. Additional studies are also required to better understand the mechanism of disease; our results point to the β -adrenoceptors being important new therapeutic targets for the treatment of LPS-induced acute lung injury.

ACKNOWLEDGMENTS

The authors thank Dr Tomohisa Uchida for giving helpful advice and for scoring lung injuries.

REFERENCES

- Martin GS, Mannino DM, Eaton S, Moss M: The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 348:1546-1554, 2003.
- Schlichting D, McCollam JS: Recognizing and managing severe sepsis: a common and deadly threat. *South Med J* 100:594-600, 2007.
- Alberti C, Brun-Buisson C, Burchardi H, Martin C, Goodman S, Artigas A, Sicignano A, Palazzo M, Moreno R, Boulmé R, et al.: Epidemiology of sepsis and infection in ICU patients from an international multicentre cohort study. *Intensive Care Med* 28:108-121, 2002.
- Rudiger A, Singer M: Mechanisms of sepsis-induced cardiac dysfunction. *Crit Care Med* 35(6):1599-1608, 2007.
- Costa EL, Schettino IA, Schettino GP: The lung in sepsis: guilty or innocent? *Endocr Metab Immune Disord Drug Targets* 6:213-216, 2006.
- McDonough KH, Vitrag JI: Sepsis-induced myocardial dysfunction and myocardial protection from ischemia/reperfusion injury. *Front Biosci* 11:23-32, 2006.
- Keasler W, Traeger T, Westerholt A, Neher F, Mikulcak M, Müller A, Maier S, Heidecke CD: The vagal nerve as a link between the nervous and immune system in the instance of polymicrobial sepsis. *Langenbecks Arch Surg* 391: 83-87, 2006.
- Iguchi S, Iwamura H, Nishizaki M, Hayashi A, Senokuchi K, Kobayashi K, Sakaki K, Hachiya K, Ichioka Y, Kawamura M: Development of a highly cardioselective ultra short-acting β -blocker, ONO-1101. *Chem Pharm Bull (Tokyo)* 40:1462-1469, 1992.
- Motomura S, Hagihara A, Narumi Y, Hashimoto K: Time course of a new ultrashort-acting β -adrenoceptor-blocking drug, ONO-1101: comparison with those of esmolol and propranolol by using the canine isolated, blood-perfused heart preparations. *J Cardiovasc Pharmacol* 31:431-440, 1998.
- Chi DS, Fitzgerald SM, Pitts S, Cantor K, King E, Lee SA, Huang SK, Krishnaswamy G: MAPK-dependent regulation of IL-1- and β -adrenoceptor-induced inflammatory cytokine production from mast cells: implications for the stress response. *BMC Immunol* 5:22, 2004.

11. Le Tulzo Y, Shenkar R, Kaneko D, Moine P, Fantuzzi G, Dinarello CA, Abraham E: Hemorrhage increases cytokine expression in lung mononuclear cells in mice: involvement of catecholamines in nuclear factor-kappaB regulation and cytokine expression. *J Clin Invest* 99:1516-1524, 1997.
12. Yu M, Wang H, Ding A, Golenbock DT, Latz E, Czura CJ, Fenton MJ, Tracey KJ, Yang H: HMGB1 signals through toll-like receptor (TLR) 4 and TLR2. *Shock* 26:174-179, 2006.
13. Park JS, Gamboni-Robertson F, He Q, Svetkauskaite D, Kim JY, Strassheim D, Sohn JW, Yamada S, Maruyama I, Banerjee A, et al.: High Mobility Group Box 1 protein (HMGB1) interacts with multiple Toll like receptors. *Am J Physiol Cell Physiol* 290:917-924, 2006.
14. Ueno H, Matsuda T, Hashimoto S, Amaya F, Kitamura Y, Tanaka M, Kobayashi A, Maruyama I, Yamada S, Hasegawa N, et al.: Contributions of high mobility group box protein in experimental and clinical acute lung injury. *Am J Respir Crit Care Med* 170:1310-1316, 2004.
15. Wang H, Bloom O, Zhang M, Vishnubhakt JM, Ombrellino M, Che J, Frazier A, Yang H, Ivanova S, Borovikova L, et al.: HMGB-1 as a late mediator of endotoxin lethality in mice. *Science* 285:248-251, 1999.
16. Hagiwara S, Iwasaka H, Uchino T, Noguchi T: High Mobility Group Box 1 induces a negative inotropic effect on the left ventricle in an isolated rat heart model of septic shock: a pilot study. *Circ J* 72:1012-1017, 2008.
17. Wang H, Yang H, Czura CJ, Sama AE, Tracey KJ: HMGB1 as a late mediator of lethal systemic inflammation. *Am J Respir Crit Care Med* 164:1768-1773, 2001.
18. Poltorak A, He X, Smirnova I, Liu MY, Van Huffel C, Du X, Birdwell D, Alejos E, Silva M, Galanos C, et al.: Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science* 282:2085-2088, 1998.
19. Li Q, Verma IM: NF-kappaB regulation in the immune system. *Nat Rev Immunol* 2:725-734, 2002.
20. Taguchi N, Kawahara K, Yone K, Hashiguchi T, Yamakuchi M, Goto M, Inoue K, Yamada S, Ijiri K, Matsunaga S, et al.: High mobility group box chromosomal protein 1 plays a role in the pathogenesis of rheumatoid arthritis as a novel cytokine. *Arthritis Rheum* 48:971-981, 2003.
21. Wang H, Vishnubhakt JM, Bloom O, Zhang M, Ombrellino M, Sama A, Tracey KJ: Proinflammatory cytokines (tumor necrosis factor and interleukin 1) stimulate release of high mobility group protein-1 by pituitary cells. *Surgery* 126:389-392, 1999.
22. Liu SF, Malik AB: NF-kappa B activation as a pathological mechanism of septic shock and inflammation. *Am J Physiol Lung Cell Mol Physiol* 290(4):L622-L645, 2006.
23. Murakami K, McGuire R, Cox RA, Jodojin JM, Bjertnaes LJ, Katahira J, Traber LD, Schmalstieg FC, Hawkins HK, Herndon DN, et al.: Heparin nebulization attenuates acute lung injury in sepsis following smoke inhalation in sheep. *Shock* 18:236-241, 2002.
24. Meng X, Brown JM, Ao L, Nordeen SK, Franklin W, Harken AH, Banerjee A: Endotoxin induces cardiac HSP70 and resistance to endotoxemic myocardial depression in rats. *Am J Physiol* 271(4 Pt 1):C1316-C1324, 1996.
25. Brown MA, Jones WK: NF-kappaB action in sepsis: the innate immune system and the heart. *Front Biosci* 9:1201-1217, 2004.
26. Meng X, Brown JM, Ao L, Banerjee A, Harken AH: Norepinephrine induces cardiac heat shock protein 70 and delayed cardioprotection in the rat through 1 adrenoceptors. *Cardiovasc Res* 32:374-383, 1996.
27. Zhang H, Kim YK, Govindarajan A, Baba A, Binnie M, Marco Ranieri V, Liu M, Slutsky AS: Effect of adrenoceptors on endotoxin-induced cytokines and lipid peroxidation in lung explants. *Am J Respir Crit Care Med* 160:1703-1710, 1999.
28. Takashima S, Schlidt SA, Koukoulis G, Sevala M, Egan TM: Isoproterenol reduces ischemia-reperfusion lung injury despite beta-blockade. *J Surg Res* 126(1):114-120, 2005.
29. Bhatia M, Mochhala S: Role of inflammatory mediators in the pathophysiology of acute respiratory distress syndrome. *J Pathol* 202:145-156, 2004.
30. Court O, Kumar A, Parrillo JE, Kumar A: Clinical review: myocardial depression in sepsis and septic shock. *Crit Care* 6:500-508, 2002.
31. Zhang H, Wang HY, Bassel-Duby R, Maass DL, Johnston WE, Horton JW, Tao W: Role of interleukin-6 in cardiac inflammation and dysfunction after burn complicated by sepsis. *Am J Physiol Heart Circ Physiol* 292:H2408-H2416, 2007.
32. Duncan DJ, Hopkins PM, Harrison SM: Negative inotropic effects of tumour necrosis factor-alpha and interleukin-1beta are ameliorated by alfentanil in rat ventricular myocytes. *Br J Pharmacol* 150:720-726, 2007.
33. Suzuki T, Morisaki H, Serita R, Yamamoto M, Kotake Y, Ishizaka A, Takeda J: Infusion of the beta-adrenergic blocker esmolol attenuates myocardial dysfunction in septic rats. *Crit Care Med* 33:2294-2301, 2005.
34. Izeboud CA, Monshouwer M, van Miert AS, Witkamp RF: The beta-adrenoceptor agonist clenbuterol is a potent inhibitor of the LPS-induced production of TNF-alpha and IL-6 *in vitro* and *in vivo*. *Inflamm Res* 48(9):497-502, 1999.
35. Izeboud CA, Mocking JA, Monshouwer M, van Miert AS, Witkamp RF: Participation of beta-adrenergic receptors on macrophages in modulation of LPS-induced cytokine release. *J Recept Signal Transduct Res* 19(1-4):191-202, 1999.
36. Izeboud CA, Hoebe KH, Grotenoster AF, Nijmeijer SM, van Miert AS, Witkamp RR, Rodenhoff RJ: Endotoxin-induced liver damage in rats is minimized by beta 2-adrenoceptor stimulation. *Inflamm Res* 53(3):93-99, 2004.
37. Nakamura A, Johns EJ, Imaizumi A, Yanagawa Y, Kohsaka T: Beta(2)-adrenoceptor agonist suppresses renal tumour necrosis factor and enhances interleukin-6 gene expression induced by endotoxin. *Nephrol Dial Transplant* 15(12):1928-1934, 2000.
38. Palumbo R, Galvez BG, Pusterla T, De Marchis F, Cossu G, Marcu KB, Bianchi ME: Cells migrating to sites of tissue damage in response to the danger signal HMGB1 require NF-kappaB activation. *J Cell Biol* 179:33-40, 2007.
39. Amalich F, Garcia-Palmero E, Lopez J, Jimenez M, Madero R, Renart J, Vazquez JJ, Montiel C: Predictive value of nuclear factor kappaB activity and plasma cytokine levels in patients with sepsis. *Infect Immun* 68:1942-1945, 2000.
40. Bohrer H, Qiu F, Zimmermann T, Zhang Y, Jilmer T, Mannel D, Bottiger BW, Stern DM, Waldherr R, Saeger HD, et al.: Role of NF-kappaB in the mortality of sepsis. *J Clin Invest* 100:972-985, 1997.
41. Pritts TA, Moon MR, Wang Q, Hungness ES, Salzman AL, Fischer JE, Hasselgren PO: Activation of NF-kappaB varies in different regions of the gastrointestinal tract during endotoxemia. *Shock* 14:118-122, 2000.
42. Yang SP, Ho LJ, Lin YL, Cheng SM, Tsao TP, Chang DM, Hsu YL, Shih CY, Juan TY, Lai JH: Carvedilol, a new antioxidative beta-blocker, blocks *in vitro* human peripheral blood T cell activation by downregulating NF-kappaB activity. *Cardiovasc Res* 59:776-787, 2003.
43. Dumitriu IE, Baruah P, Valentinis B, Voll RE, Herrmann M, Nawroth PP, Arnold B, Bianchi ME, Manfredi AA, Rovere-Querini P: Release of high mobility group box 1 by dendritic cells controls T cell activation via the receptor for advanced glycation end products. *J Immunol* 174:7506-7515, 2005.



Effects of hyperglycemia and insulin therapy on high mobility group box 1 in endotoxin-induced acute lung injury in a rat model*

Satoshi Hagiwara, MD, PhD; Hideo Iwasaka, MD, PhD; Akira Hasegawa, MD; Hironori Koga, MD; Takayuki Noguchi, MD, PhD

Objective: Hyperglycemia and insulin resistance are commonly seen in septic patients and are associated with increased morbidity and mortality. High mobility group box 1 (HMGB1) protein has been shown to play a key role as a significant factor in sepsis pathogenesis. This study investigated the increase in lung damage because of hyperglycemia and HMGB1 increase in a lipopolysaccharide-induced septic rat model and the potential for insulin therapy to reduce this lung damage by decreasing the serum level of HMGB1.

Design: Randomized, prospective animal study.

Setting: University medical center research laboratory.

Subjects: Male Wistar rats.

Interventions: Septic hyperglycemia was induced by infusion of glucose immediately after administration of lipopolysaccharide in rats.

Measurements and Main Results: Animals were monitored for blood glucose. Separate cohorts were killed at 12 and 24 hrs postlipopolysaccharide administration and analyzed for HMGB1

and lung damage. The effects of insulin treatment were also examined. Hyperglycemic septic animals had significantly higher blood glucose and enhanced lung damage. In addition, HMGB1 was increased in the serum of hyperglycemic rats. On the other hand, insulin treatment for hyperglycemia resulted in significantly lower blood glucose and decreased both the lung damage and the serum level of HMGB1. In an *in vitro* study, insulin treatment inhibited the activation of NF- κ B.

Conclusions: Hyperglycemia is associated with higher HMGB1 levels and lung damage in sepsis. Insulin therapy significantly reduced lung damage, suggesting that management of hyperglycemia with insulin might decrease HMGB1 levels in the serum and lung tissue. One of the mechanisms that could contribute to the inhibition of HMGB1 secretion might be related to the inhibition of NF- κ B. (Crit Care Med 2008; 36:2407–2413)

Key Words: lipopolysaccharide; lung injury; sepsis; hyperglycemia; insulin; high mobility group box 1

Sepsis is a significant cause of morbidity and mortality worldwide. This is partly due to complications, such as acute respiratory distress syndrome, which are difficult to treat (1–3). Hyperglycemia and insulin resistance are commonly observed in septic patients and have proinflammatory effects, including increased production of reactive oxygen species (4). Indeed, a study concluded that treating hyperglycemia with intensive insulin therapy reduced morbidity and mortality among critically ill patients in a surgical intensive care unit (5). However, a recent trial investigating the effects of intensive insulin therapy on patients with severe

sepsis was terminated early because the risk of adverse effects was significantly higher in the intensive-therapy group (6). Researchers and clinicians currently are debating the effects of intensive insulin therapy on patients with severe sepsis.

High mobility group box 1 (HMGB1) protein plays a key role in the pathogenesis of sepsis as a late-phase mediator of systemic inflammation (7). HMGB1 is an intranuclear protein that was originally identified as a regulator of chromatin structure (8). Once released from necrotic cells or secreted by activated monocytes/macrophages, HMGB1 can bind to receptors for advanced glycation end-products (RAGE). The proinflammatory effects of HMGB1 are mediated partially by signaling through RAGE (9, 10). HMGB1 contributes to lethality in septic patients and activates downstream cytokine release (11). Thereafter, release of HMGB1 to the extracellular space is controlled by cytokines (12). Extracellular HMGB1 seems to play a role in both clinical, acute respiratory distress syndrome, and experimental models. The concentrations of HMGB1 were elevated in the plasma and lungs of

patients with acute respiratory distress syndrome and in a mouse model of lipopolysaccharide (LPS)-induced lung injury (13).

Hyperglycemia up-regulates various inflammatory mediators (14). One reason for this enhanced inflammation is the activation of NF- κ B after glucose intake (15). On the other hand, insulin has an anti-inflammatory effect. Dandona et al. (16) reported that this effect includes the inhibition of NF- κ B. A recent study demonstrated that tight control of blood glucose by insulin therapy protected the endothelium of critically ill patients (17). Controlling hyperglycemia is acknowledged as an important component of sepsis treatment.

We hypothesized that hyperglycemia might increase the serum and tissue HMGB1 levels, thereby contributing to LPS-induced lung injury. Insulin therapy may be able to prevent LPS-induced lung injury by regulating hyperglycemia and inhibiting the levels of HMGB1 in the serum and tissue. To test this hypothesis, we investigated the effects of hyperglycemia and insulin therapy on LPS-induced lung injury in rats.

*See also p. 2475.

From the Department of Brain and Nerve Science, Anesthesiology, Oita University Faculty of Medicine, Oita, Japan.

The authors have not disclosed any potential conflicts of interest.

For information regarding this article, E-mail: saku@med.oita-u.ac.jp

Copyright © 2008 by the Society of Critical Care Medicine and Lippincott Williams & Wilkins

DOI: 10.1097/CCM.0b013e318180b3ba

METHODS

Materials. Insulin was obtained from Novo Nordisk A/S (Copenhagen, Denmark). LPS (O127:B8) was obtained from Sigma (St Louis, MO). All other reagents were of the highest available analytical grade. We used commercially available antibodies to HMGB1 (Shino-Test Corporation, Tokyo, Japan), phosphorylated IkappaB alpha (p-IkappaB alpha), IkappaB alpha (Cell Signaling Technology, Beverly, MA), and beta-actin (Abcam, Cambridge, UK).

Animals. All protocols conformed to the National Institute of Health guidelines and all animals received humane care in compliance with the Principles of Laboratory Animal Care. The study was approved by the Ethical Committee of Animal Research at the College of Medicine, Oita University, Oita, Japan. Male Wistar rats weighing 250–300 g (Kyudou, Saga, Japan) were used in all experiments. All animals were housed with free access to food and water *ad libitum*.

Animals were randomly assigned to one of five groups ($n = 10$ in each group): 1) saline group, rats intravenously received 0.9% NaCl solution (2.5 mL/kg/hr); 2) Glu group, rats received 20% glucose solution (2.5 mL/kg/hr) intravenously; 3) LPS group, rats received 0.9% NaCl solution (2.5 mL/kg/hr) intravenously after the intravenous injection of a bolus of LPS (2.5 mg/kg); 4) LPS + Glu group, rats received 20% glucose solution (2.5 mL/kg/hr) intravenously after the intravenous injection of a bolus of LPS (2.5 mg/kg); and 5) LPS + Ins group, rats received 20% glucose solution (2.5 mL/kg/hr) and insulin (20 IU/kg/hr) intravenously after the intravenous injection of a bolus of LPS (2.5 mg/kg). Serum samples of venous blood (0.1 mL) were obtained from the external jugular vein at various time points (0, 4, 6, 8, 10, 12, and 24 hrs). All animals were killed at 12 or 24 hrs and lung tissue specimens were quickly removed and processed as indicated below.

Blood Glucose Measurement. Blood samples ($n = 6$ in each group) were collected from the external jugular vein of rats for each time point (0, 4, 6, 8, 10, 12, and 24 hrs). The whole blood glucose levels of the rats were monitored using the glucose-oxidase method with a Glu-test-sensor blood glucose monitoring system from Sanwa Kagaku Kenkyosho (Nagoya, Japan).

Histologic Analysis. Left lungs ($n = 4$ in each group) were obtained from animals in the LPS, LPS + Glu, and LPS + Ins groups under sevoflurane anesthesia. The lung tissue specimens were fixed with 10% formalin and embedded in paraffin. The sections were stained with hematoxylin and eosin. A pathologist unaware of the group assignment analyzed the samples. An evaluation of lung injury was done according to Murakami's technique (18). Briefly, 24 areas in the lung parenchyma were graded on a scale of 0–4 (0, absent and appears normal; 1, light; 2, mod-

erate; 3, strong; 4, intense) for congestion, edema, inflammatory cells, and hemorrhaging. The mean score for each of the parameters was calculated.

Measurements of HMGB1. Serum samples ($n = 6$ in each group) were assayed using the sandwich enzyme-linked immunosorbent assay (ELISA) method. Ninety-six well plates were precoated with monoclonal antibodies specific to rat HMGB1 (Shino-Test, Tokyo, Japan). The samples, negative controls, and diluted HMGB1 standard markers were added into their respective wells. Detection of HMGB1 in the samples was performed according to the manufacturer's protocols. The A450 values were determined using an ELISA reader (Bio-Rad Laboratories, Hercules, CA).

Immunohistochemical Analysis. Lung tissue ($n = 4$ in each group) samples were fixed immediately in 4% paraformaldehyde at 4°C overnight and then embedded in O.C.T. Compound (Sakura Finetechnical, Tokyo, Japan). Sections were incubated with mouse anti-rat-HMGB1 polyclonal antibody (1:1000 dilution) overnight at 4°C. The sections were rinsed with phosphate buffered saline (PBS) and then incubated with peroxidase-conjugated anti-mouse-IgG (1:1000 dilution). After rinsing with PBS, the slides were stained using the LSAB2 kit (Dako, Carpinteria, CA), including the biotin-avidin-peroxidase complex system. After development, the slides were counterstained with Mayer's hematoxylin and mounted.

Cell Culture Study. The murine macrophage cell line RAW264.7 was maintained in Dulbecco's Modified Eagle's Medium containing 5% heat-inactivated fetal bovine calf serum and antibiotics at 37°C under 5% CO₂. The medium was removed and replaced with high glucose (450 mg/dL) or low glucose (100 mg/dL) Dulbecco's Modified Eagle's Medium containing 5% fetal bovine calf serum. Cells were randomly assigned to the following groups: 1) saline in low glucose medium; 2) LPS in normal glucose medium; 3) LPS in high glucose medium; and 4) LPS and insulin in high glucose medium. Cells were lysed and protein was extracted using nuclear and cytoplasmic extraction reagents (PIERCE Biotechnology, Rockford, IL). Cytoplasmic and nuclear protein fractions were used for Western blot analyses, as described below.

NF-kappaB Binding Assay. The DNA binding activity of NF-kappaB (p50/p65) was determined using an ELISA-based nonradioactive NF-kappaB p50/p65 transcription factor assay kit (Chemicon, Temecula, CA). Detection of NF-kappaB activity in the samples was performed according to the manufacturer's protocols. The A450 values were determined using an ELISA reader (Bio-Rad Laboratories).

Western Blotting. Proteins were subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis on 12% gels and then were transferred onto polyvinylidene difluoride sheets (Millipore, Bedford, MA). The membranes were incubated with the primary anti-

body at 1/1000 in PBS-T for 1 hr at room temperature. After three washes in PBS-T, peroxidase-labeled secondary antibody was added at 1/1000 in PBS-T and incubated for 1 hr at room temperature. Blots were developed using enhanced chemiluminescence (ECL) reagents and exposed to Hyperfilm ECL (Amersham, Buckinghamshire, UK).

Statistical Analysis. All data are presented as the mean \pm SE. The data were compared using nonparametric tests. The Kruskal-Wallis test for comparisons between all groups was used. A p value of <0.05 was considered to be statistically significant.

RESULTS

Effect of Glucose or Insulin Administration on Hyperglycemia

Figure 1 shows the changes in blood glucose concentrations in the five groups of rats during the infusions of saline (saline group), glucose (Glu group), saline after LPS injection (LPS group), glucose after LPS injection (LPS + Glu group), or glucose and insulin after LPS injection (LPS + Ins group). In the saline group, glucose levels were maintained at normal levels over 24 hrs and ranged from $158 \pm$




Figure 1. The changes in blood glucose concentration in lipopolysaccharide (LPS), LPS + glucose (Glu), LPS + insulin (Ins) groups after endotoxin challenge. Blood glucose concentration at the indicated hours is shown for each group ($n = 6$ for each group). Filled square (saline group), animals received saline (5 mL/kg/hr) infusion only; circles (Glu group), animals received glucose (5 mL/kg/hr) infusion only; triangles (LPS group), animals received saline infusion after a 2.5 mg/kg endotoxin injection; diamonds (LPS + Glu group), animals received glucose infusion after a 2.5 mg/kg endotoxin injection; open square (LPS + Ins group), animals received glucose and insulin infusions after a 2.5 mg/kg endotoxin injection. All data are expressed as the means \pm SE. * $p < 0.05$ vs. each point in saline only group.




Figure 2. Effects of high glucose on lung injury in lipopolysaccharide (LPS)-treated rats. Rats were given either saline (saline group), glucose (Glu group), saline and LPS (LPS group), glucose and LPS (LPS + Glu group), or glucose, insulin and LPS (LPS + Ins group). Lungs are from the saline (A, magnification $\times 40$), Glu (B, magnification $\times 40$), LPS (C, magnification $\times 40$), LPS + Glu (D, magnification $\times 40$), and LPS + Ins (E, magnification $\times 40$) groups, respectively. Sections shown are stained with hematoxylin and eosin.




Figure 3. Changes in lung histology score. The identified histologic changes included congestion, edema, inflammation, and hemorrhaging 24 hrs after the administration of lipopolysaccharide (LPS). White bars represent the saline-administrated sham animals (saline group), black bars represent animals injected with glucose (Glu group), gray bars represent animals injected with saline and LPS (LPS group), slashed bars represent animals injected with glucose and LPS (LPS + Glu), and checkered bars represent animals injected with glucose, insulin and LPS (LPS + Ins group). The data are expressed as the mean \pm SE. * $p < 0.05$ vs. LPS + Glu group.


8 to 192 ± 9 mg/dL. In the glucose group, glucose levels were slightly higher than normal levels over 24 hrs, with an average of around 200 mg/dL. In the LPS group, glucose levels were lowered below normal levels by 6 hrs after LPS administration. In the LPS + Glu group, after starting the infusion of glucose, blood glucose levels significantly increased and reached 312 ± 5 mg/dL over the first 6 hrs. In the LPS + Ins group, glucose levels were maintained at normal levels during the experimental period.

The Effect of Glucose or Insulin Administration on Lung Tissue Specimens

Histologic examinations of lung tissue sections were performed 24 hrs after LPS (2.5 mg/kg intravenous) administration. Observations of microscopic changes in the lung tissue specimens showed edema-like formation and inflammatory cell infiltration in the LPS and LPS + Glu groups compared with the saline and Glu groups (Fig. 2A–D). Interstitial edema and inflammatory cell infiltration were markedly reduced in the LPS + Ins group in comparison with the LPS and LPS + Glu groups (Fig. 2C–E). We observed no histologic changes after administering saline and glucose alone (Fig. 2A–B). The histology scores, based on the number of areas with congestion, edema, inflammation, and hemorrhaging (17) were all significantly higher after the administration of LPS than in the saline group. All of the scores were significantly higher in the LPS + Glu group compared with the saline, LPS, and LPS + Ins groups (Fig. 3).

Changes in HMGB1 Protein Expression in Lung Tissue

At 24 hrs after LPS administration, positive staining cells for HMGB1 were rarely observed in lung tissue from the saline (Fig. 4A) and Glu group (data not shown). Positive staining cells for HMGB1 were somewhat increased in lung tissue from the LPS group (Fig. 4B) and markedly increased in the LPS + Glu group (Fig. 4C). The number of cells expressing HMGB1 was substantially decreased in the LPS + Ins group (Fig. 4D) compared with the LPS + Glu group (Fig. 4C). At 12 hrs after LPS administration, Western blot analysis showed an increase in HMGB1 expression in the lung tissue from rats receiving glucose after LPS injection. This increase in HMGB1



LPS(2.5mg/kg)	-	+	+	+
Intravenous Injection	Saline	20% glucose	20% glucose	Saline
Insulin	-	-	+	+
Time(hour)	12	12	12	12

Figure 4. Changes in the high mobility group box 1 (HMGB1) protein expression in lung tissue specimens after lipopolysaccharide (LPS) administration in rats. The expression of HMGB1 protein 24 hrs after LPS administration was detected by immunohistochemistry in lung tissue specimens from rats in the saline group (A), LPS group (B), LPS + glucose group (C), or LPS + insulin group (D); arrows indicate cells that are positively stained for HMGB1. The expression of HMGB1 protein in lung tissue 12 hrs after LPS administration was detected by Western blotting (E); the rats were given either saline alone, LPS + glucose, LPS + glucose + insulin, or LPS + saline + insulin.

was not seen in lung tissue from rats that received insulin in addition to glucose after LPS injection (Fig. 4E).

Changes in HMGB1 Serum Concentrations

The serum level of HMGB1 increased in all groups at 12 and 24 hrs after the administration of LPS. HMGB1 serum concentration in the LPS + Glu group was significantly higher compared with the LPS group at both time points. On the other hand, in the LPS + Ins group, HMGB1 concentrations were not significantly different from the LPS group at

either time point (Fig. 5). The serum level of HMGB1 was below the detection level in all groups before the administration of LPS (data not shown).

Effect of High Glucose on Levels of HMGB1 Secreted by LPS-Stimulated Murine Macrophages

Secreted HMGB1 levels increased after LPS stimulation in high glucose medium compared with low glucose medium, but were inhibited by the addition of insulin in high glucose medium. In addition, the inhibition of HMGB1 by insulin

seemed to be dose-dependent, with minimal inhibition at a concentration of 1 μ M and the greatest effect observed at the highest dose (100 μ M) tested in this study (Fig. 6). Because in this experiment we observed the greatest inhibition of HMGB1 using 100 μ M insulin, we used this concentration for subsequent experiments.

Insulin Inhibits the I κ B Kinase Pathways and Modulates NF- κ B

The DNA binding activity of the transcription factor NF- κ B (p50/p65) in the nucleus of RAW264.7 cells increased 1 hr after administration of LPS in both low and high glucose medium. Activation of the NF- κ B transcription factor p50/p65 after the administration of LPS was higher in high glucose medium than in low glucose medium. However, administration of insulin suppressed this increase (Fig. 7).

Treatment with LPS resulted in the degradation of I κ B α in both the low and high glucose medium, which was inhibited by the addition of insulin to the high glucose medium. In addition, the phosphorylation of I κ B α in RAW264.7 cells increased after LPS administration in both the low glucose medium and high glucose medium, which was also inhibited by the addition of insulin to the high glucose medium (Fig. 8).

DISCUSSION

Our study showed that hyperglycemia induced histologic changes in our LPS-induced septic rat model. Hyperglycemia occurs frequently among critically ill patients, including those with sepsis (19). When hyperglycemia occurs as a complication, it has been shown to be associated with high mortality and poor clinical outcomes in subpopulations of clinically ill patients (20). One of the reasons for this is that hyperglycemia can amplify the inflammatory response (21). These results indicate that hyperglycemia aggravates the condition of a patient in septic shock by increasing the severity of lung injury.

HMGB1 has been shown to play an important role in various types of inflammation. In addition, HMGB1 is an important mediator of cell death, and its expression occurs at a relatively late stage after injury (7). Especially when it is released as a result of acute lung inflamma-




Figure 5. Temporal changes of high mobility group box 1 (HMGB1) serum concentration after lipopolysaccharide (LPS) administration in rats. Rats were given either saline and LPS (LPS group), glucose and LPS (LPS + Glu group), or glucose, insulin and LPS (LPS + Ins group). HMGB1 serum concentration at 12 and 24 hrs after LPS administration is shown for each group (n = 6 for each group). White bars represent LPS group animals, black bars represent LPS + Glu group animals, and slashed bars represent LPS + Ins group animals. All data are expressed as the means \pm SE. * p < 0.05 vs. LPS group at 12 hrs.




Figure 6. Effect of insulin in high glucose (HG) and low glucose (LG) medium on high mobility group box 1 (HMGB1) production by lipopolysaccharide (LPS)-stimulated murine macrophages. Murine macrophages treated with insulin were stimulated with LPS for 20 hrs. Conditioned media were collected and HMGB1 levels were determined by enzyme-linked immunosorbent assay. The checked bar represents values measured for the LG medium, the slashed bar represents values measured for the HG medium without insulin, the gray bar represents values measured for the HG medium with 1 μ M insulin, the black bar represents values measured for the HG medium with 10 μ M insulin, and the white bar represents values measured for the HG medium with 100 μ M insulin. All data are expressed as the means \pm SE. *Denotes a significant difference compared with LPS in HG medium (p < 0.05).

tion, HMGB1 can have various effects (13). In the present study, we demonstrated that the expression of HMGB1 in the serum and lung tissue increased in

hyperglycemic rats with LPS-induced lung injury. Moreover, hyperglycemia also increased histologic changes including interstitial and intra-alveolar inflam-

mation, edema, congestion, and hemorrhage. These findings are consistent with previous studies, which showed that HMGB1 plays a critical role in the development of LPS-induced lung injury (13, 22).

LPS stimulation of macrophages activates several intracellular signaling pathways, including the I κ B kinase (IKK)-NF- κ B pathway (23). In addition, the proinflammatory effects of glucose metabolism include an increase in IKK- α , IKK- β , and NF- κ B binding with a decrease in I κ B α (15). We found that insulin in high glucose medium suppressed the activation of NF- κ B and prevented the phosphorylation of a molecule central to the IKK pathway, I κ B. A previous study also demonstrated that insulin inhibits the activation of NF- κ B (16, 24). Accordingly, the inhibition of I κ B phosphorylation after insulin administration for hyperglycemia may lead to the inhibition of NF- κ B activation. The *in vitro* effect of insulin on NF- κ B/I κ B suggests that there is a direct mechanism that is not necessarily dependent on a decrease in glucose concentrations. When I κ B is phosphorylated after LPS stimulation, the activation of NF- κ B may be related to the secretion of HMGB1 (25). We demonstrated that insulin in the high glucose medium inhibits LPS-induced NF- κ B activation, which may participate in the inhibition of HMGB1 secretion in the LPS-induced lung injury model. However, in this study, we used two different species (rat *in vivo* study, and murine *in vitro* study) and acknowledge that there may be mechanistic differences between the two systems. Similar experiments using a mouse model of lung injury may clarify our results.

In the *in vitro* study, inhibiting HMGB1 secretion required a high concentration of insulin. Aljada et al. (24) has demonstrated the suppression of NF- κ B by insulin in human aortic endothelial cells at more physiologically relevant concentrations. However, we used human insulin in our murine *in vitro* study and acknowledge that there may be differences. This might be the reason we needed a very high concentration of insulin to observe the inhibition of HMGB1 secretion and NF- κ B activation.

We also demonstrated that insulin treatment significantly decreased the severity of lung injury and both serum and lung tissue HMGB1 levels in the LPS-induced septic rat model. Systemic in-




Figure 7. Effect of insulin on the lipopolysaccharide (LPS)-induced increase in the specific binding of p50/p65 to DNA. DNA binding activity of NF-kappaB p50/p65 is shown for nuclear fractions from LPS-stimulated RAW264.7 cells and low glucose (Lg) medium, high glucose (HG) medium, and HG medium with 100 μ M insulin. The dashed bar represents values measured for the LG medium, the black bar represents values for the HG medium without insulin and the white bar represents values for the HG medium with insulin. All data are expressed as the mean \pm SE. *Denotes a significant difference compared with LPS in HG medium at 1 hr ($p < 0.05$).




Figure 8. Effect of insulin on the lipopolysaccharide (LPS)-induced phosphorylation of IkappaB. Proteins were extracted from murine macrophages with or without insulin treatment 1 hr after stimulation with LPS (100 ng/mL). The cytoplasmic levels of phosphorylated IkappaB alpha, IkappaB alpha, and beta-actin were determined by Western blot analysis using antibodies specific for phosphorylated IkappaB alpha, IkappaB alpha, and beta-actin (1/1000 dilution). LG, low glucose; HG, high glucose.

inflammation can occur during hyperglycemia and persist after blood glucose levels have returned to normal. This can lead to an increase in the systemic inflammatory response to a subsequent moderate stress (14). An increase in systemic inflammation can lead to a greater risk of organ damage in septic shock patients. On the other hand, insulin treatment has been

demonstrated to reduce the inflammation in various diseases (26). Therefore, insulin therapy may benefit patients with conditions that are often complicated by severe systemic inflammation including septicemia, meningitis, and pneumonia (27). This anti-inflammatory effect of insulin has also been shown in rats injected with endotoxin (28). In this study, we

demonstrated that insulin treatment led to a decrease in the serum level of HMGB1 after LPS administration. These findings suggest that anti-inflammatory effect of insulin in the rat model of LPS-induced lung injury may be at least partially due to the ability of insulin to decrease HMGB1 levels in the serum and lung tissue.

This study only examined the short-term (up to 24 hrs) therapeutic effects of insulin treatment for hyperglycemia after the onset of lethal endotoxemia. However, recent clinical studies demonstrated that the anti-inflammatory effect of insulin was long-lived (17, 29). The results of this study cannot be extrapolated to chronic conditions in patients. Since long-term insulin treatment was effective in intensive care unit patients, we need further study of the long-term effects of insulin in a septic shock model.

In conclusion, hyperglycemia aggravated the pathophysiological findings in a rat model of LPS-induced lung injury and increased the serum level of HMGB1. The administration of insulin to treat hyperglycemia improved these findings. These observations indicated that insulin treatment for hyperglycemia could decrease the severity of lung injury by inhibiting the HMGB1 levels during the acute phase of LPS-induced lung injury in rats. As a caveat, in this study, we used small numbers of animals and the high levels of glucose might not be entirely relevant for clinical conditions in the intensive care unit.

ACKNOWLEDGMENT

We thank Dr. Tomohisa Uchida for giving us helpful advice and for scoring lung injuries.

REFERENCES

- Martin GS, Mannino DM, Eaton S, et al: The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 2003; 348:1546-1554
- Wyncoll DL, Evans TW: Acute respiratory distress syndrome. *Lancet* 1999; 354: 497-501
- Lehr HA, Bittinger F, Kirkpatrick CJ: Microcirculatory dysfunction in sepsis: A pathogenic basis for therapy? *J Pathol* 2000; 190: 373-386
- Marik PE, Raghavan M: Stress-hyperglycemia, insulin and immunomodulation in sepsis. *Intensive Care Med* 2004; 30:748-756
- van den Berghe G, Wouters P, Weekers F, et al: Intensive insulin therapy in the critically

- ill patients. *N Engl J Med* 2001; 345: 1359-1367
6. Brunkhorst FM, Engel C, Bloos F, et al: German Competence Network Sepsis (SepNet). Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med* 2008; 358:125-139
 7. Wang H, Yang H, Czura CJ, et al: HMGB1 as a late mediator of lethal systemic inflammation. *Am J Respir Crit Care Med* 2001; 164: 1768-1773
 8. Bustin M: Regulation of DNA-dependent activities by the functional motifs of the high-mobility-group chromosomal proteins. *Mol Cell Biol* 1999; 19:5237-5246
 9. Dumitriu IE, Baruah P, Valentini B, et al: Release of high mobility group box 1 by dendritic cells controls T cell activation via the receptor for advanced glycation end products. *J Immunol* 2005; 174:7506-7515
 10. Kokkola R, Andersson A, Mullins G, et al: RAGE is the major receptor for the proinflammatory activity of HMGB1 in rodent macrophages. *Scand J Immunol* 2005; 61:1-9
 11. Wang H, Bloom O, Zhang M, et al: HMG-1 as a late mediator of endotoxin lethality in mice. *Science* 1999; 285:248-251
 12. Wang H, Vishnubhakat JM, Bloom O, et al: Proinflammatory cytokines (tumor necrosis factor and interleukin 1) stimulate release of high mobility group protein-1 by pituitary cells. *Surgery* 1999; 126:389-392
 13. Ueno H, Matsuda T, Hashimoto S, et al: Contributions of high mobility group box protein in experimental and clinical acute lung injury. *Am J Respir Crit Care Med* 2004; 170: 1310-1316
 14. Ling PR, Smith RJ, Bistrian BR: Hyperglycemia enhances the cytokine production and oxidative responses to a low but not high dose of endotoxin in rats. *Crit Care Med* 2005; 33:1084-1089
 15. Aljada A, Friedman J, Ghanim H, et al: Glucose ingestion induces an increase in intranuclear nuclear factor kappaB, a fall in cellular inhibitor kappaB, and an increase in tumor necrosis factor alpha messenger RNA by mononuclear cells in healthy human subjects. *Metabolism* 2006; 55:1177-1185
 16. Dandona P, Aljada A, Mohanty P, et al: Insulin inhibits intranuclear nuclear factor kappaB and stimulates IkappaB in mononuclear cells in obese subjects: Evidence for an anti-inflammatory effect? *J Clin Endocrinol Metab* 2001; 86:3257-3265
 17. Langouche L, Vanhorebeek I, Vlasselaers D, et al: Intensive insulin therapy protects the endothelium of critically ill patients. *J Clin Invest* 2005; 115:2277-2286
 18. Murakami K, McGuire R, Cox RA, et al: Heparin nebulization attenuates acute lung injury in sepsis following smoke inhalation in sheep. *Shock* 2002; 18:236-241
 19. Frankenfield DC, Omert LA, Badellino MM, et al: Correlation between measured energy expenditure and clinically observed variables in trauma and sepsis patients. *JPEN J Parenter Enteral Nutr* 1994; 18:398-403
 20. Krinsley JS: Association between hyperglycemia and increased hospital mortality in a heterogeneous population of critically ill patients. *Mayo Clin Proc* 2003; 78:1471-1478
 21. Leonidou L, Mouzaki A, Michalaki M, et al: Cytokine production and hospital mortality in patients with sepsis-induced stress hyperglycemia. *J Infect* 2007; 55:340-346
 22. Wang H, Yang H, Tracey KJ: Extracellular role of HMGB1 in inflammation and sepsis. *J Intern Med* 2004; 255:320-331
 23. Schottelius AJ, Mayo MW, Sartor RB, et al: Interleukin-10 signaling blocks inhibitor of kappaB kinase activity and nuclear factor kappaB DNA binding. *J Biol Chem* 1999; 274: 31868-31874
 24. Aljada A, Ghanim H, Saadeh R, et al: Insulin inhibits NFkappaB and MCP-1 expression in human aortic endothelial cells. *J Clin Endocrinol Metab* 2001; 86:450-453
 25. Ulloa L, Ochani M, Yang H, et al: Ethyl pyruvate prevents lethality in mice with established lethal sepsis in the hospital setting: The case for improved control among non-dietics. *Ann Pharmacother* 2005; 39:492-501
 26. Conner TM, Flesner-Gurley KR, Barner JC: Hyperglycemia in the hospital setting: The case for improved control among non-diabetics. *Ann Pharmacother* 2005; 39: 492-501
 27. Dandona P, Mohanty P, Chaudhuri A, et al: Insulin infusion in acute illness. *J Clin Invest* 2005; 115:2069-2072
 28. Jesc MG, Klein D, Bolder U, et al: Insulin attenuates the systemic inflammatory response in endotoxemic rats. *Endocrinology* 2004; 145:4084-4093
 29. Chaudhuri A, Janicke D, Wilson MF, et al: Anti-inflammatory and profibrinolytic effect of insulin in acute ST-segment-elevation myocardial infarction. *Circulation* 2004; 109: 849-854

A Neutrophil Elastase Inhibitor, Sivelestat, Reduces Lung Injury Following Endotoxin-Induced Shock in Rats by Inhibiting HMGB1

Satoshi Hagiwara,^{1,2} Hideo Iwasaka,¹ Kazumi Togo,¹ and Takayuki Noguchi¹

Abstract—Neutrophil elastase (NE) plays an important role in the progression of acute lung injury (ALI). Sivelestat sodium hydrate (Sivelestat) is a highly specific synthetic inhibitor of NE. High mobility group box 1 (HMGB1) is one of the key mediators in the development of sepsis. The aim of this study was to evaluate the effect of sivelestat and to determine whether it can reduce lipopolysaccharide (LPS)-induced acute lung injury in rats. Rats were randomly divided into a negative control group, an LPS-induced sepsis group, and a group treated with sivelestat prior to LPS administration. Animals in the sivelestat group received a bolus of 10 mg/kg of sivelestat injected into the intraperitoneal cavity before the LPS treatment. Furthermore, rats were administered sivelestat at 0, 1, 3, and 6 h following LPS treatment. We measured cytokine and HMGB1 levels in the serum after the induction of sepsis. In addition, we observed histopathology, wet/dry weight ratio, inducible nitric oxide synthase and HMGB1 expression in the lung tissue. Lung histopathology was significantly improved in the sivelestat group compared to the LPS group. Serum and pulmonary HMGB1 levels were lower over time among sivelestat-treated animals. Furthermore, inhibition of NF- κ B activity was observed with the administration of sivelestat. These results suggest that sivelestat reduces LPS-induced lung injury at least partially by inhibiting inflammation and NF- κ B activity.

KEY WORDS: inflammation; sepsis; high mobility group box 1; cytokine; NF- κ B; nitric oxide.

INTRODUCTION

The high case fatality rate of sepsis is due to complications throughout the major organ systems that can occur by various mechanisms. In spite of advances in therapy, the mortality rate for severe sepsis remains high, reaching 80% to 90% for septic shock with multiple organ dysfunction [1]. Acute respiratory distress syndrome (ARDS) and acute lung injury (ALI) are

well defined clinical disorders that are a significant cause of morbidity and mortality in intensive care units (ICU). The most frequent cause of ARDS and ALI is sepsis [2]. Despite recent increases in our understanding of the molecular underpinnings of sepsis, most of these complications remain refractory to treatment [3]. The respiratory system in particular is severely affected and difficult to treat [4].

Neutrophil elastase (NE, also called leukocyte elastase) is a serine protease found primarily in the azurophilic granules of neutrophils. Intracellular NE is a key molecule of the innate immune system and is required for effective killing of phagocytosed bacterial and fungal pathogens [5, 6]. In addition, Moraes TJ *et al.* [7] reviewed the potentially pathologic role of NE in ARDS and ALI, including the ability of extracellular NE

¹Department of Brain and Nerve Science, Anesthesiology, Oita University Faculty of Medicine, 1-1 Idaigaoka-Hasamamachi-Yufu, City-Oita 879-5593, Japan

²To whom correspondence should be addressed at Department of Brain and Nerve Science, Anesthesiology, Oita University Faculty of Medicine, 1-1 Idaigaoka-Hasamamachi-Yufu, City-Oita 879-5593, Japan. E-mail: saku@med.oita-u.ac.jp

to cause endothelial damage and increase vascular permeability.

Sivelestat is a specific neutrophil elastase inhibitor that has no other known antiprotease effects [8]. Previous studies have reported that sivelestat can reduce lung injury associated with systemic inflammatory response syndrome (SIRS) in humans [9] and also decrease endotoxin-induced lung injury in animal models [10, 11]. These results suggested that sivelestat might be useful for the treatment of SIRS and ARDS/ALI. However, the protective mechanisms of sivelestat are not well understood.

HMGB1 was recognized as a late-phase mediator in the pathogenesis of sepsis [12] and was thought to contribute to the pathology and mortality of sepsis as an inflammatory mediator [13]. Recent studies indicate that HMGB1 plays a role in the enhancement of various inflammatory processes. Indeed, HMGB1 mediates cell-to-cell signaling by binding to Toll-like receptors (TLRs) [14, 15]. HMGB1 also acts as a procoagulant, thereby enhancing the inflammatory response in septic shock [16]. In addition, HMGB1 is present in the serum of sepsis patients, with higher levels associated with increased mortality [17, 18].

The release of HMGB1 from macrophages was first reported in response to stimulation by endotoxin, and increased serum levels of HMGB1 have been observed in sepsis [12, 19]. Endotoxin, also called lipopolysaccharide (LPS), is found in the outer membrane of Gram-negative bacteria and induces immune responses by interacting with TLR4 [20]. The release of HMGB1 from macrophages is regulated by various inflammatory mediators and activates signal transduction pathways, including the NF- κ B pathway. Specifically, nitric oxide and TNF- α induce HMGB1 release in response to LPS [21, 22].

We hypothesized that sivelestat prevents acute lung injury in a rat model by inhibiting serum and tissue HMGB1. To test this hypothesis, we investigated the impact of sivelestat administration on serum and lung levels of HMGB1, and lung histopathology in rats with LPS-induced lung injury. To further elucidate the mechanism of this effect, we assessed the effect of sivelestat by measuring the activity of NF- κ B in the lung tissue.

MATERIALS AND METHODS

Animals

Male Wistar rats weighing 250–300 g (Kyudou, Saga, Japan) were used in all experiments. All protocols

conformed to the National Institute of Health (NIH) guidelines and the animals received humane care in compliance with the Principles of Laboratory Animal Care. All animals were housed with access to food and water *ad libitum*.

Anesthesia was induced by 4% sevofluran. Animals were randomly assigned to one of three groups: (1) LPS group, (2) sivelestat group, or (3) negative control group. In the LPS group, the rats received an infusion of 0.9% NaCl solution (1.0 ml/kg) into the intraperitoneal cavity 30 min prior to LPS administration. When LPS (7.5 mg/kg) was administered, the rats simultaneously received 0.9% NaCl solution (1.0 ml/kg) into the vein. The rats received an infusion of 0.9% NaCl solution (1.0 ml/kg) into the intraperitoneal cavity 1, 3 and 6 h following LPS administration. In the sivelestat group, the rats received an infusion of sivelestat (10 mg/kg) into the intraperitoneal cavity 30 min prior to LPS administration. When LPS (7.5 mg/kg) was administered, the rats simultaneously received sivelestat (10 mg/kg) into the vein. The rats received an infusion of sivelestat (10 mg/kg) into the intraperitoneal cavity 1, 3 and 6 h following LPS administration. Sivelestat was donated by Ono Pharma Co. Ltd. (Osaka, Japan). In the negative control group, the rats received an infusion of 0.9% NaCl solution (1.0 ml/kg) into the intraperitoneal cavity a total of five times to correspond with injections in the other groups. Before and after surgery, animals in all groups had unlimited access to food and water.

Wet-to-dry Weight (W/D) Ratio

Animals were euthanized with an overdose of sodium pentobarbital 12 h following LPS treatment. After the administration of sodium pentobarbital, the chest was opened and the right ventricle was injected with 10 ml of phosphate-buffered saline (PBS) to wash blood cells from the pulmonary circulation. The lungs were removed, weighed, and then dried in an oven at 80°C for 48 h. The dried lungs were weighed again to calculate pulmonary wet/dry ratios.

Histological Analysis

Lung sections were stained with hematoxylin and eosin. A pathologist blinded to group assignment analyzed the samples and determined levels of lung injury according to Murakami's technique [23]. Briefly, 24 areas in the lung parenchyma were graded on a scale of 0–4 (0, absent and appears normal; 1, light; 2, moderate; 3, strong; 4, intense) for congestion, edema,

infiltration of inflammatory cells, and hemorrhaging. The mean score for each of the parameters was then used for analysis purposes.

Measurements of Cytokine and HMGB1 Secretions

HMGB1, IL-6 and TNF- α levels were determined using commercial enzyme-linked immunosorbent assay (ELISA) kits (HMGB1: Shino-Test Corporation, Tokyo, Japan; IL-6 and TNF- α : R&D Systems Inc, Minneapolis, MN). The absorbance at 450 nm was determined using an ELISA reader (Bio-Rad Laboratories, Hercules, CA, USA).

Nitrite/Nitrate Determination

In biological fluids NO is rapidly deactivated by oxidation to nitrite and nitrate. After collecting the blood samples by cardiac puncture, we detected NO via nitrite/nitrate. Serum nitrite/nitrate concentration in plasma was determined using commercial kits according to the manufacturers' protocols (R&D Systems Inc., Minneapolis, MN, USA). This kit measured nitrite/nitrate concentration using a modification of the Griess assay. The absorbance was measured at 540 nm. nitrite/nitrate concentration was calculated using a nitrite/nitrate standard curve and expressed as micromoles per liter.




Fig. 1. Effects of sivelestat on lung injury in LPS-treated rats and changes in the lung histology score. Lung tissue was obtained from rats in the negative control group (a: magnification $\times 40$), LPS group (b: magnification $\times 40$), or sivelestat group (c: magnification $\times 40$). Sections shown are stained with hematoxylin and eosin. d Histological changes were scored based on findings of congestion, edema, inflammation, and hemorrhage 12 h after treatment with LPS. White bars represent the negative control group, black bars represent animals administered LPS only, and the slashed bars represent animals injected with sivelestat before LPS treatment. The data are expressed as the mean \pm SE. Asterisk denotes a significant difference compared with the LPS group ($p < 0.05$).

Immunohistochemical Analysis

Tissue samples were fixed immediately in 4% paraformaldehyde and incubated at 4°C overnight, embedded in O.C.T. Compound (Sakura Finetechnical Co., Tokyo, Japan), and sectioned at 5 µm. Endogenous peroxidase activity was blocked with 0.3% H₂O₂ and sodium azide (1 mg/ml) for 10 min followed by blocking of non-specific protein binding by 10% sheep serum for 10 min. Blocked sections were incubated with mouse anti-HMGB1 (Shino-Test Corporation, Tokyo, Japan) polyclonal antibody (1:1,000 dilution) overnight at 4°C. The sections were rinsed with PBS three times for 5 min and then were incubated with peroxidase-conjugated anti-mouse IgG. After three PBS rinses, the slides were stained using the biotin avidin peroxidase complex system from an LSAB2 kit (Dako, Via Real Carpinteria, CA, USA). Following development, slides were counterstained with Mayer's hematoxylin and mounted.

Western Blotting

Proteins were subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) on a 12% gel and transferred onto polyvinylidene difluoride sheets (Millipore, Bedford, MA). The membranes were washed with PBS containing 0.1% Tween 20 (PBS-T) and 5% skim milk for 1 hr at room temperature. After three washes with PBS-T, the membranes were incubated with antibodies (1:1000 dilution) to iNOS (Santa Cruz Biotechnology, Inc, Santa Cruz, CA, USA), HMGB1 (Shino-Test Corporation, Tokyo, Japan) or beta-actin (Abcam, Cambridge, UK) for 1 h at room temperature. After three washes, peroxidase-labeled secondary antibodies (ZYMED, South San Francisco, CA) were added at 1/1,000 in PBS-T and incubated for 1 hr at room temperature. Blots were subsequently developed using an enhanced chemiluminescence detection kit (Amersham, Buckinghamshire, UK) and exposed on Hyperfilm ECL (Amersham, Buckinghamshire, UK).

NF-κB Binding Assay

Nuclear proteins were extracted from the lung tissue using NE-PER extraction reagents (PIERCE Biotechnology, Rockford, IL, USA). The DNA binding activity of NF-κB (p50/p65) was determined using an ELISA-based nonradioactive NF-κB p50/p65 transcription factor assay kit (Chemicon, Temecula, CA, USA). The absorbance at 450 nm was determined using an ELISA reader (Bio-Rad Laboratories, Hercules, CA, USA).

Statistical Analysis

All data were presented as the mean±standard error of the mean (SEM). Independent groups were compared using Mann-Whitney's *U*-test. Statistical analysis was performed using the Kruskal-Wallis test. A *p* value of <0.05 was considered to be statistically significant.

RESULTS

Effect of Sivelestat on the Lung in a Rat Model of LPS-Induced Septic Shock

In the negative control group, no pulmonary histological alterations were observed (Fig. 1a). Among the LPS affected rats, lung tissue specimens observed 12 h after the administration of LPS showed edema formation and interstitial infiltration by neutrophils and reduced the alveolar spaces (Fig. 1b). Sivelestat treatment significantly reduced the interstitial edema and inflammatory cell infiltration in comparison to the LPS group (Fig. 1c). Histology scores, based on the number of areas with congestion, edema, inflammation, and hemorrhaging, were all significantly higher in the LPS group than in the negative control group (Fig. 1d). All scores were lower in the sivelestat group compared to the LPS only group (Fig. 1d). At 12 h following LPS administration, the *W/D* ratios of the lungs in the LPS group were elevated (Fig. 2). The *W/D* ratios of the




Fig. 2. Effects of sivelestat on lung wet-to-dry weight (*W/D*) ratio of LPS-treated rats. The lung *W/D* ratio 6 h following LPS administration is shown for each group (*n*=6 for each group). The white bar represents animals in the negative control group, the black bar represents animals administered LPS only, and the slashed bar represents animals injected with sivelestat before LPS treatment. The data are expressed as the mean±SE. Asterisk denotes a significant difference compared with the LPS group (*p*<0.05).




Fig. 3. Changes in iNOS and HMGB1 protein expression in rat lung tissue following LPS treatment. Rats were given either saline (LPS group) or sivelestat (sivelestat group) prior to the administration of LPS. The expression of iNOS or HMGB1 in lung tissue 12 h following LPS treatment in both groups was detected by Western blot.

lungs in the sivelestat group were significantly lower than those in the LPS group and were comparable to the values of the control animals (Fig. 2).

Measurement of Serum Nitrite/Nitrate

Nitrite/nitrate are metabolic products of NO and often used as markers to indicate NO formation. Nitrite/nitrate concentration was measured in plasma 0, 3, 6, 9, 12 and 24 h following LPS treatment (Fig. 4). After LPS




Fig. 4. Effect of LPS administration on serum levels of nitrite/nitrate. Serum concentration of nitrite/nitrate was measured 0, 3, 6, 9, 12, and 24 h following LPS administration (7.5 mg/kg, I.V., n=6 for each group). The LPS group is represented by squares and the sivelestat group is represented by circles. All data are expressed as the mean±SE. Asterisk statistically significant (p<0.05) with respect to the LPS group.




Fig. 5. Temporal changes in serum TNF-alpha, IL-6, and HMGB1 concentrations following LPS treatment in rats. Rats were given either saline (LPS group), or sivelestat (sivelestat group) prior to the administration of LPS. Serum concentrations pre-treatment and 3, 6, 9, 12, and 24 h following LPS treatment are shown for TNF-alpha (a), IL-6 (b), and HMGB1 (c). The LPS group is shown with circle symbols (n=6) and the sivelestat group is shown with square symbols (n=6). All data are expressed as the mean±SE. Asterisk denotes a significant difference compared with the LPS group (p<0.05).



Fig. 6. Changes in HMGB1 protein expression in rat lung tissue following LPS treatment. Immunohistochemical analysis to detect HMGB1 in lungs from animals killed 12 h following intravenous saline administration; (a, magnification $\times 400$), 12 h following LPS treatment (b, magnification $\times 400$) or from animals treated with sivelestat and killed 12 h following LPS treatment (c, magnification $\times 400$). Arrows point to positive cells.

administration, the levels of serum nitrite/nitrate were increased. Sivelestat treatment significantly decreased the nitrite/nitrate levels compared to the LPS only group (Fig. 4).

Effects of Sivelestat on Serum Levels of IL-6, TNF-alpha, and HMGB1

Administration of sivelestat prior to LPS treatment in rats significantly decreased the concentration of TNF-alpha at 3 h post-treatment (Fig. 5a). Similarly, groups treated with sivelestat showed decreased serum levels of IL-6 at 6 h later (Fig. 5b). Serum levels of HMGB1 also increased after the administration of LPS. This increase was less prominent in rats treated with sivelestat compared to the LPS group (Fig. 5c). Serum levels of IL-6, TNF-alpha and HMGB1 remained constant in negative control group rats at all assay times (data not shown).

Effect of Sivelestat on the Lung Tissue

Western blot analysis showed that the expression of HMGB1 in the lung tissue increased 12 hr following LPS administration. This increase was less pronounced among sivelestat treated rats compared with the LPS group (Fig. 3). At the same time point, immunohistochemical analysis revealed that cells expressing HMGB1 increased following LPS treatment (Fig. 6b). In contrast, the percentage of cells expressing HMGB1 did not significantly increase in the rats receiving sivelestat prior to LPS treatment (Fig. 6c). Similar to the expression of HMGB1, the expression of iNOS in the lung tissue increased as a result of LPS treatment.

This increase was also less pronounced among sivelestat treated rats compared with the LPS group (Fig. 3). Administration of LPS led to a robust activation of the NF- κ B transcription factor p50/p65. However, administration of sivelestat partially suppressed this increase (Fig. 7).




Fig. 7. Effect of sivelestat on the LPS-induced increase in specific binding of p50 and p65 to DNA. Nuclear fractions were harvested 6 h following LPS treatment from rats with or without sivelestat administration ($n=6$ for each group). The DNA binding activity assay showed a marked decrease in the p50/p65 binding activity in nuclear fractions from lung tissue cells in the sivelestat group. All data are expressed as the mean \pm SE. Asterisk denotes a significant difference compared with LPS group ($p<0.05$).

DISCUSSION

In this study, we demonstrated that the administration of sivelestat significantly reduced the lung injury and wet/dry ratio of lung tissue following LPS treatment (Figs. 1 and 2). Acute lung injury in acute respiratory distress syndrome (ARDS) is considered an acute inflammatory condition by many clinicians. Although the pathology of this syndrome is poorly defined, recent evidence suggests that neutrophil elastase (NE) plays a role in the development of acute lung injury, which is characterized by increased permeability and resulting pulmonary edema [5, 24].

Inducible NOS (iNOS) is typically activated only in systemic inflammatory conditions, including septic shock. The up-regulation of iNOS is mediated by specific proinflammatory agents, such as TNF- α [25]. Increased expression of iNOS induces nitric oxide (NO) release, leading to organ damage. NO is believed to play a key role in inflammation, and has been implicated in a wide range of disease processes [25]. In this study, we demonstrated that iNOS and nitrate/nitrite were significantly increased in the LPS-induced lung injury model. Treatment with sivelestat, however, decreased these mediators (Figs. 3 and 4). These results suggest that improvement of LPS-induced lung injury by treatment with sivelestat might be related to a decrease in NO.

Our findings showed that serum levels of cytokines and HMGB1 were significantly inhibited by the administration of sivelestat prior to LPS treatment (Fig. 5a-c). Several cytokines, including TNF- α and IL-6, can cause neutrophil-mediated lung tissue injury [26]. The significant decrease in lung injury among rats treated with sivelestat, therefore, may be partially attributable to decreased levels of these pro-inflammatory cytokines. Our findings suggested that sivelestat may inhibit not only serum cytokine levels, but also levels of HMGB1 in both serum and lung tissue (Figs. 3 and 5c). This is the first report of sivelestat administration inhibiting HMGB1 in a rat model of sepsis. These results build on the growing body of evidence suggesting the importance of HMGB1 in the initiation of acute lung injury [13].

Various pathways have been identified as potential mechanisms behind the inhibition of HMGB1 secretion. Stimulation of multiple cytokines induces the release of HMGB1 [23]. In particular, evidence supports a potential role for TNF- α and NO in the regulation of LPS-induced HMGB1 release [23, 27]. In the present study,

the TNF- α and IL-6 levels in the serum were significantly lower in the sivelestat-treated group than in the LPS group. The decrease of TNF- α and NO in the sivelestat group suggests that the inhibition of cytokine and NO secretion by sivelestat in systemic inflammation mediates the inhibition of HMGB1.

We demonstrated that sivelestat inhibited activation of nuclear factor κ B (NF- κ B) in rat lung tissue in the LPS-induced lung injury model. In acute lung injury, multiple signaling pathways lead to cytokine synthesis and release of HMGB1. Systemic inflammation, such as septic shock, also results in activation of various signaling pathways [29, 30]. Significantly, these signaling pathways induce activation of the transcription factor NF- κ B, a central regulator of inflammation, cytokine synthesis and stress responses in many cell types, including macrophages [29, 31, 32]. In patients with septic lung injury, NF- κ B activity in alveolar macrophages has been shown to be significantly increased [28]. Because NE activates NF- κ B in LPS-induced lung injury [33], sivelestat might reduce NF- κ B activation in lung injury through inhibition of NE (Figs. 6 and 7).

In a rat model of LPS-induced lung injury, sivelestat administration at 10 mg/kg can prevent acute lung injury. Our results suggest that sivelestat might therapeutically benefit septic patients not only through inhibition of HMGB1, but also through inhibition of inflammatory responses in systemic inflammatory conditions. Further research will be necessary to assess the effect of delayed sivelestat administration in LPS-induced septic shock models.

ACKNOWLEDGMENTS

The authors wish to thank Dr. Tomohisa Uchida for giving us helpful advice and for scoring lung injuries.

REFERENCES

- Schlichting, D., and J. S. McCollam. 2007. Recognizing and managing severe sepsis: a common and deadly threat. *South Med. J.* 100(6):594-600.
- Costa, E. L., I. A. Schettino, and G. P. Schettino. 2006. The lung in sepsis: guilty or innocent? *Endocr. Metab. Immune. Disord. Drug Targets* 6(2):213-216.
- Andrews, P., E. Azoulay, M. Antonelli, L. Brochard, C. Brun-Buisson, G. Dobb, J. Y. Fagon, H. Gerlach, J. Groeneveld, J. Mancebo, P. Metnitz, S. Nava, J. Pugin, M. Pinsky, P. Radermacher, C. Richard, and R. Tasker. 2006. Year in review in intensive care medicine, 2005. II. Infection and sepsis, ventilator-associated pneumonia, ethics,

- haematology and haemostasis, ICU organisation and scoring, brain injury. *Intensive Care Med.* **32**:380-390.
4. Andrews, P., E. Azoulay, M. Antonelli, L. Brochard, C. Brun-Buisson, G. Dobb, J. Y. Fagon, H. Gerlach, J. Groeneveld, J. Mancebo, P. Metnitz, S. Nava, J. Pugin, M. Pinsky, P. Radermacher, C. Richard, R. Tasker, and B. Vallet. 2005. Year in review in intensive care medicine, 2004. I. Respiratory failure, infection, and sepsis. *Intensive Care Med.* **31**:28-40.
 5. Weiss, S. J. 1989. Tissue destruction by neutrophils. *N. Engl. J. Med.* **320**:365-376.
 6. Vender, R. L. 1996. Therapeutic potential of neutrophil-elastase inhibition in pulmonary disease. *J. Invest. Med.* **44**:531-539.
 7. Moreas, T. J., C. W. Chow, and G. P. Downey. 2003. Proteases and lung injury. *Crit. Care Med.* **31**:S189-S194.
 8. Kawabata, K., M. Suzuki, M. Sugitani, K. Imaki, M. Toda, and T. Miyamoto. 1991. ONO-5046, a novel inhibitor of human neutrophil elastase. *Biochem. Biophys. Res. Commun.* **177**:814-820.
 9. Bellomo, R., S. Uchino, T. Naka, and L. Wan. 2004. Hidden evidence to the West: multicentre, randomised, controlled trials in sepsis and systemic inflammatory response syndrome in Japanese journals. *Intensive Care Med.* **30**:911-917.
 10. Sakamaki, F., A. Ishizaka, T. Urano, K. Sayama, H. Nakamura, T. Terashima, Y. Waki, S. Tasaka, N. Hasegawa, K. Sato, N. Nakagawa, T. Obata, and M. Kanazawa. 1996. Effect of a specific neutrophil elastase inhibitor, ONO-5046, on endotoxin-induced acute lung injury. *Am. J. Respir. Crit. Care Med.* **153**(1):391-397.
 11. Wakayama, F., I. Fukuda, Y. Suzuki, and N. Kondo. 2007. Neutrophil elastase inhibitor, sivelestat, attenuates acute lung injury after cardiopulmonary bypass in the rabbit endotoxemia model. *Ann. Thorac. Surg.* **83**(1):153-160.
 12. Wang, H., O. Bloom, M. Zhang, J. M. Vishnubhakat, M. Ombrellino, J. Che, A. Frazier, H. Yang, S. Ivanova, L. Borovikova, K. R. Manogue, E. Faist, E. Abraham, J. Andersson, U. Andersson, P. E. Molina, N. N. Abumrad, A. Sama, and K. J. Tracey. 1999. HMG-1 as a late mediator of endotoxin lethality in mice. *Science* **285**:248-251.
 13. Ueno, H., T. Matsuda, S. Hashimoto, F. Amaya, Y. Kitamura, M. Tanaka, A. Kobayashi, I. Maruyama, S. Yamada, N. Hasegawa, J. Soejima, H. Koh, and A. Ishizaka. 2004. Contributions of high mobility group box protein in experimental and clinical acute lung injury. *Am. J. Respir. Crit. Care Med.* **170**:1310-1316.
 14. Yu, M., H. Wang, A. Ding, D. T. Golenbock, E. Latz, C. J. Czura, M. J. Fenton, K. J. Tracey, and H. Yang. 2006. HMGB1 signals through toll-like receptor (TLR) 4 and TLR2. *Shock* **26**:174-179.
 15. Park, J. S., F. Gamboni-Robertson, Q. He, D. Svetkauskaite, J. Y. Kim, D. Strassheim, J. W. Sohn, S. Yamada, I. Maruyama, A. Banerjee, A. Ishizaka, and E. Abraham. 2006. High Mobility Group Box 1 protein (HMGB1) interacts with multiple Toll like receptors. *Am. J. Physiol. Cell Physiol.* **290**:917-924.
 16. Sunden-Cullberg, J., A. Norrby-Teglund, and C. J. Treutiger. 2006. The role of high mobility group box-1 protein in severe sepsis. *Curr. Opin. Infect. Dis.* **19**:231-236.
 17. Hatada, T., H. Wada, T. Nobori, K. Okabayashi, K. Maruyama, Y. Abe, S. Uemoto, S. Yamada, and I. Maruyama. 2005. Plasma concentrations and importance of High Mobility Group Box protein in the prognosis of organ failure in patients with disseminated intravascular coagulation. *Thromb. Haemost.* **94**:975-979.
 18. Sunden-Cullberg, J., A. Norrby-Teglund, A. Rouhiainen, H. Rauvala, G. Herman, K. J. Tracey, M. L. Lee, J. Andersson, L. Tokics, and C. J. Treutiger. 2005. Persistent elevation of high mobility group box-1 protein (HMGB1) in patients with severe sepsis and septic shock. *Crit. Care Med.* **33**:564-573.
 19. Wang, H., H. Yang, C. J. Czura, A. E. Sama, and K. J. Tracey. 2001. HMGB1 as a late mediator of lethal systemic inflammation. *Am. J. Respir. Crit. Care Med.* **164**:1768-1773.
 20. Poltorak, A., X. He, I. Smirnova, M. Y. Liu, C. Van Huffel, X. Du, D. Birdwell, E. Alejos, M. Silva, C. Galanos, M. Freudenberg, P. Ricciardi-Castagnoli, B. Layton, and B. Beutler. 1998. Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in Tlr4 gene. *Science* **282**:2085-2088.
 21. Jiang, W., and D. S. Pisetsky. 2006. The role of IFN-alpha and nitric oxide in the release of HMGB1 by RAW 264.7 cells stimulated with polyinosinic-polycytidylic acid or lipopolysaccharide. *J. Immunol.* **177**(5):3337-3343.
 22. Wang, H., J. M. Vishnubhakat, O. Bloom, M. Zhang, M. Ombrellino, A. Sama, and K. J. Tracey. 1999. Proinflammatory cytokines (tumor necrosis factor and interleukin 1) stimulate release of high mobility group protein-1 by pituitary cells. *Surgery* **126**:389-392.
 23. Murakami, K., R. McGuire, R. A. Cox, J. M. Jodoin, L. J. Bjertnaes, J. Katahira, L. D. Traber, F. C. Schmalstieg, H. K. Hawkins, D. N. Herrndon, and D. L. Traber. 2002. Heparin nebulization attenuates acute lung injury in sepsis following smoke inhalation in sheep. *Shock* **18**:236-241.
 24. Tate, R. M., and J. E. Repine. 1983. Neutrophils and the adult respiratory distress syndrome. *Am. Rev. Respir. Dis.* **128**:552-559.
 25. Chandra, A., P. Enkhbaatar, Y. Nakano, L. D. Traber, and D. L. Traber. 2006. Sepsis: emerging role of nitric oxide and selectins. *Clinics* **61**:71-76.
 26. Bhatia, M., and S. Mochhala. 2004. Role of inflammatory mediators in the pathophysiology of acute respiratory distress syndrome. *J. Pathol.* **202**:145-156.
 27. Jiang, W., and D. S. Pisetsky. 2006. The role of IFN-alpha and nitric oxide in the release of HMGB1 by RAW 264.7 cells stimulated with polyinosinic-polycytidylic acid or lipopolysaccharide. *J. Immunol.* **177**(5):3337-3343.
 28. Schwartz, M. D., E. E. Moore, F. A. Moore, R. Shenkar, P. Moine, J. B. Haenel, and E. Abraham. 1996. Nuclear factor-kappa B is activated in alveolar macrophages from patients with acute respiratory distress syndrome. *Crit. Care Med.* **24**:1285-1292.
 29. Liu, S. F., and A. B. Malik. 2006. NF-kappa B activation as a pathological mechanism of septic shock and inflammation. *Am. J. Physiol. Lung Cell Mol. Physiol.* **290**(4):L622-L645.
 30. Pritts, T. A., M. R. Moon, Q. Wang, E. S. Hungness, A. L. Salzman, J. E. Fischer, and P. O. Hasselgren. 2000. Activation of NF-kappaB varies in different regions of the gastrointestinal tract during endotoxemia. *Shock* **14**:118-122.
 31. Kawai, T., O. Adachi, T. Ogawa, K. Takeda, and S. Akira. 1999. Unresponsiveness of MyD88-deficient mice to endotoxin. *Immunity* **11**(1):115-122.
 32. Adcock, I. M. 1997. Transcription factors as activators of gene transcription: AP-1 and NF-kappa B. *Monaldi Arch. Chest Dis.* **52**:178-186.
 33. Kuwahara, I., E. P. Lillehoj, W. Lu, I. S. Singh, Y. Isohama, T. Miyata, and K. C. Kim. 2006. Neutrophil elastase induces IL-8 gene transcription and protein release through p38/NF-(kappa)B activation via EGFR transactivation in a lung epithelial cell line. *Am. J. Physiol. Lung Cell Mol. Physiol.* **291**:L407-L416.

High Mobility Group Box 1 Induces a Negative Inotropic Effect on the Left Ventricle in an Isolated Rat Heart Model of Septic Shock

— A Pilot Study —

Satoshi Hagiwara, MD; Hideo Iwasaka, MD;
Tomoko Uchino, MD; Takayuki Noguchi, MD

Background Sepsis can be exacerbated by an inappropriate immune response and the severe impact of this disease on the cardiovascular system is well documented. High mobility group box 1 (HMGB1) protein is an important mediator in the pathogenesis of sepsis and its role in cardiovascular system dysfunction was investigated in an lipopolysaccharide (LPS)-induced rat model of sepsis.

Methods and Results Twelve hours after intravenous bolus injections of LPS (5 mg/kg), rats were killed and heart samples were harvested. Immunoblot analysis was performed to assess expression levels of HMGB1 in cardiac myocytes. Left ventricular developed pressure (LVDP) served as a measure of systolic function. LPS administration was associated with an increase in the expression of HMGB1 in cardiac myocytes and a decrease in cardiac function. Hearts from the LPS-treated rats were also perfused with recombinant HMGB1 and cardiac function measured. The dose-dependent effects observed with elevated HMGB1 included decreased LVDP, decreased left ventricular (LV) $+dP/dt_{max}$, decreased absolute value of LV $-dP/dt_{min}$, and increased LV end-diastolic pressure.

Conclusions HMGB1 stimulation produces a negative inotropic effect during septic shock, suggesting an important role for this molecule in cardiovascular system dysfunction during sepsis. (Circ J 2008; 72: 1012–1017)

Key Words: Cardiac function; HMGB1; Negative inotropic; Sepsis

Mortality associated with septic shock is thought to be a consequence of severe derangements in the cardiovascular system¹ with myocardial dysfunction playing a primary role². Indeed, reduced cardiac contractile function has been observed both in septic patients^{3,4} and experimental animal models of lipopolysaccharide (LPS)-induced sepsis⁵. The cellular and molecular mechanisms responsible for these effects, however, remain incompletely understood.

Recent studies in rat and mouse models of sepsis have partially elucidated these mechanisms. One key factor that has emerged is high mobility group box 1 (HMGB1) protein, which serves as a late-phase mediator in the pathogenesis of sepsis. HMGB1 is an intranuclear protein that was originally identified as an important gene regulator⁶. Following the activation of the inflammation cascade, HMGB1 is released from necrotic damaged cells and secreted by activated monocytes and macrophages. Its mechanism of action is presumably via its affinity for receptors for advanced glycation end-products, Toll-like receptor 2 and Toll-like

receptor 4. HMGB1 binding to its receptors activates intracellular signaling pathways, such as the NF- κ B pathway, which induce downstream cytokine release⁷, thereby enhancing the inflammatory response^{8,9}.

LPS, a cell membrane component shed from Gram-negative bacteria, is a key mediator of septicemia and its pathogenic effects on the heart¹⁰ but exactly how this endotoxin causes cardiac dysfunction remains largely unclear. We hypothesized that HMGB1 might be a key mediator in the pathogenesis of cardiac dysfunction and to test this, we exposed hearts from LPS-treated rats to recombinant HMGB1 protein.

Methods

Materials

Recombinant HMGB1 was purchased from Shino-Test Corporation (Tokyo, Japan) and LPS (O127:B8) was obtained from Sigma (St Louis, MO, USA). All reagents were of the highest available analytical grade.

Animals

Male Wistar rats weighing 250–300 g (Kyudou, Saga, Japan) were used in all experiments. All rats received humane care in compliance with the National Institute of Health (NIH) guidelines and the Principles of Laboratory Animal Care. All animals were housed with free access to food and water.

(Received November 26, 2007; revised manuscript received January 27, 2008; accepted January 30, 2008)

Department of Brain and Nerve Science, Anesthesiology, Oita University Faculty of Medicine, Yufu, Japan

Mailing address: Satoshi Hagiwara, MD, Department of Brain and Nerve Science, Anesthesiology, Oita University Faculty of Medicine, 1-1 Idaigaoka, Hasama-machi, Yufu 879-5593, Japan. E-mail: saku@med.oita-u.ac.jp

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp




Fig 1. Effects of lipopolysaccharide (LPS) treatment on the gross histopathology of cardiac tissue. At 12 h after LPS injection, cardiac tissue from the control (A: $\times 100$, C: $\times 400$; H&E) and LPS groups (B: $\times 100$, D: $\times 400$; H&E) was analyzed and no gross changes were found in either group.

Experimental Protocols

The rats were randomly assigned to 1 of 2 treatment groups. The 6 animals in the control group received an intravenous bolus injection of 0.9% NaCl solution alone. The 6 animals in the LPS group received an intravenous bolus injection of LPS dissolved in 0.9% NaCl solution (5 mg/kg)¹¹. All rats were killed under general 2% sevoflurane anesthesia 12 h following injection. Tissue specimens were immediately removed surgically and processed.

Histological Analysis

Heart tissue specimens were instilled with 10% formalin. The samples were embedded in paraffin and cut into 4- μ m sections for staining with hematoxylin and eosin.

Immunoblotting Analysis

The hearts were harvested from all animals. After the blood was washed out using saline perfusion, the heart was homogenized with a T-PER (Tissue Protein Extraction Reagent; Pierce, Rockford, IL, USA) in a polytron homogenator (IKA Labortechnik, Staufen, Germany). The homogenates were then centrifuged at 10,000 g for 5 min at 4°C. The concentration of protein in the collected supernatant was measured by absorbance at 562 nm using the BCA Protein Assay Reagent system (Pierce).

For gel electrophoresis, equal quantities of protein (100 μ g) were suspended in sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis (PAGE) buffer. Protein samples were boiled for 1 min and separated using a 10% SDS-polyacrylamide gel. Protein runs using SDS-PAGE were immediately electrotransferred to polyvinylidene difluoride membrane (PVDF) (Millipore, Bedford, MA, USA) at 60 V for 3 h in a wet transfer system containing 20 mmol/L Tris-HCl/0.2 mol/L glycine in 20% methanol as the transfer buffer. The membrane was blocked with 5% nonfat dry milk in Tris/Tween buffer [25 mmol/L Tris-HCl; 2% Tween 20 (Bio-Rad Lab, Hercules, CA, USA); 0.14 mol/L NaCl] overnight at 4°C.

Antibodies to HMGB1 (Shino-Test Corporation) or β -actin (Abcam, Cambridge, UK) were used as the primary

antibody. The immune serum was diluted 500-fold with 1% nonfat dry milk and incubated with gentle shaking for 1 h. The PVDF membrane was then rinsed 3 times with TBS/Tween buffer for 10 min. The secondary antibody was diluted 1,000-fold with 1% nonfat dry milk and incubated with the membrane for 1 h. The blot was then washed by rinsing with TBS/Tween 3 times for 10 min. The membrane was treated with enhanced chemiluminescence reagent (Amersham, Buckinghamshire, UK) and then exposed to X-ray film. After scanning the X-ray film, the band concentration was calculated by quantification of the integrated optical density using NIH Image J software package (NIH, Bethesda, MD, USA).

Measurement of HMGB1 Secretion

Serum samples were assayed using the ELISA sandwich method and 96-well plates that were precoated with monoclonal antibodies specific to rat HMGB1 (Shino-Test Corporation). The samples, negative controls, and diluted HMGB1 standard markers were added to each well. Detection of HMGB1 in the samples was performed according to the manufacturer's protocols. The A450 values were determined using an ELISA reader.

Isolated Heart Perfusion and Assessment of Cardiac Function

Cardiac function was determined by a modified isovolumetric Langendorff technique as described elsewhere¹²⁻¹⁴ and expressed as left ventricular developed pressure (LVDP), left ventricular end-diastolic pressure (LVEDP), left ventricular (LV)+dP/dt_{max}, and LV-dP/dt_{max}. At the termination of the experiments, beating hearts were rapidly excised into oxygenated Krebs-Henseleit solution containing 11 mmol/L glucose, 2.0 mmol/L CaCl₂, 4.3 mmol/L KCl, 25 mmol/L NaHCO₃, 118 mmol/L NaCl, 1.2 mmol/L MgSO₄, and 1.2 mmol/L KH₂PO₄. Normothermic retrograde perfusion was performed with this solution in an isovolumetric and nonrecirculating mode. The perfusion buffer was saturated with a gas mixture of 95% O₂-5% CO₂ at a pH of 7.4. Perfusion pressure was maintained at 75 mmHg.