

### Transcription activity of *Gas6* is regulated by adiponectin and *TNF $\alpha$* via AMPK

To investigate whether *Gas6* expression is transcriptionally regulated by adiponectin, *TNF $\alpha$* , and AMPK, a promoter study was undertaken. Reporter assay using the  $-1.9$ -kb *Gas6*-luciferase DNA construct revealed that adiponectin completely reversed the down-regulation of *Gas6* transcription activity by *TNF $\alpha$* . Furthermore, compound C abrogated the effect of adiponectin on *Gas6* transcription activity, indicating that adiponectin and *TNF $\alpha$*  regulate *Gas6* expression at the transcription level via AMPK activity (Fig. 6).

### Discussion

The present study showed that adiponectin has a protective effect against Pi-induced calcification and, furthermore, has an antagonistic effect on *TNF $\alpha$* -augmented calcification. Based on our previous finding that Pi-induced calcification is dependent on apoptotic cell death in HASMC, we examined the role of adiponectin and *TNF $\alpha$*  in Pi-induced apoptosis. As expected, we found that adiponectin had an inhibitory effect and *TNF $\alpha$*  had a stimulatory effect on Pi-induced apoptosis. This study also demonstrated the regulation of *Gas6* expression by *TNF $\alpha$*  and adiponectin, a suppressive effect and a promoting effect, respectively, at the transcriptional level. Akt, a critical downstream effector of *Gas6*, was activated by adiponectin, whereas *TNF $\alpha$*  had an opposite action on its phosphorylation. Given that adiponectin and *TNF $\alpha$*  did not affect *Gas6* expression and Akt phosphorylation in the absence of Pi (data not shown), the effects of adiponectin and *TNF $\alpha$*  on these molecules may depend on Pi-induced responses. These results suggest that *Gas6* is the target of adiponectin and *TNF $\alpha$*  in regulating Pi-induced apoptosis, accompanied by modulation of the Akt-dependent survival pathway.

As reported previously (6), Pi-induced VSMC calcification is associated with both phenotypic transition to osteoblastic cells via sodium-dependent phosphate cotransporter and apoptotic cell death. In our preliminary experiments, the expression of osteopontin, an osteoblastic marker, was not affected by *TNF $\alpha$*  and adiponectin (data not shown). Although this result suggests little influence of *TNF $\alpha$*  and adiponectin on osteoblastic differentiation of VSMC, extensive and systematic investigation including other markers of osteoblastic differentiation is needed to conclude this issue.

Multiple lines of clinical evidence show that adiponectin has protective actions on the cardiovascular system (26, 27). Circulating levels of adiponectin in humans are as high as 500–30,000  $\mu\text{g/ml}$  (28). Therefore, the concentration of adiponectin (300 ng/ml) used in this study are within physiological levels. Especially, consistent with our findings, adiponectin has been implicated in apoptosis of cardiovascular cells (19, 23, 29). Adiponectin inhibits apoptosis in cardiac myocytes and fibroblasts that are exposed to hypoxia-reoxygenation stress (19). In endothelial cells, adiponectin has been reported to inhibit serum starvation-induced apoptosis (23). *In vivo* experiments have also shown that adiponectin-deficient mice develop larger myocardial infarcts due to increased myocardial cell apoptosis and *TNF $\alpha$*  expression (17). Taking these observations together with

our results, the antiapoptotic actions of adiponectin contribute to the inhibition of VSMC calcification.

Most effects of adiponectin have been attributed to the activation of AMPK, which affects many aspects of cellular metabolism including glucose uptake (30, 31), glucose utilization (32), and fatty acid oxidation (33, 34). Recently, AMPK activation in VSMC has been suggested as a target to prevent or treat vascular disease (35, 36). AICAR-induced AMPK activation inhibited angiotensin II-stimulated VSMC proliferation, and administration of AICAR prevented neointimal formation in a rat balloon injury model (35). AMPK activation in VSMC elicited cell cycle arrest at the G1 phase and inhibited cell proliferation via p53 up-regulation (36). Furthermore, in the heart, the inhibitory effects of adiponectin on ischemic injury-induced apoptosis have been shown to be dependent on AMPK activation (19). The results of *in vitro* studies also revealed that AMPK signaling is essential for the antiapoptotic activities of adiponectin on endothelial cells (23). These observations are consistent with the finding of the present study that AMPK activated by adiponectin stimulated *Gas6* expression to restore the survival pathway, leading to the suppression of calcification.

In the present study, we further demonstrated that adiponectin significantly augmented the transcriptional activity of *Gas6* that was decreased by *TNF $\alpha$* . Indeed, suppression of AMPK by compound C clearly abrogated this beneficial effect of adiponectin. This result suggests that AMPK participates in the transcriptional regulation of *Gas6* by adiponectin and *TNF $\alpha$* . Several studies support that AMPK regulates the expression of particular genes at the transcriptional level (37–39). For example, AMPK activation by AICAR enhanced activation protein 1-mediated proopiomelanocortin promoter activities, which were completely abolished by compound C (37). AMPK has been shown to mediate the transcription signal that leads to the repression of phosphoenolpyruvate carboxykinase expression, a key enzyme of gluconeogenesis, through phosphorylation of





Fig. 6. Effect of adiponectin and *TNF $\alpha$*  on *Gas6* promoter activity. HASMC were transfected with the *Gas6* promoter-luciferase construct using lipofectamine 2000. Twenty-four hours after transfection, adiponectin (300 ng/ml), compound C (1  $\mu\text{M}$ ), and *TNF $\alpha$*  (20 ng/ml) were added. Cells were incubated for an additional 44 h. Luciferase activity was normalized to that of vehicle-treated cells. All values are presented as mean  $\pm$  SE (n = 4). \*,  $P < 0.05$  by Bonferroni test. Each experiment was performed in triplicate for each condition.



**FIG. 7.** Scheme of the effect of  $TNF\alpha$  and adiponectin on Pi-induced calcification. In HASMC, exogenous Pi is internalized by sodium-dependent phosphate cotransporter (NPC, such as Pit-1) and inhibits AMPK phosphorylation, followed by down-regulation of the Gas6-mediated survival pathway. This pathway stimulates apoptosis, leading to subsequent development of calcification.  $TNF\alpha$  directly suppresses AMPK activation by promoting PP2C activation via TNF receptor-1 (TNFR1). On the other hand, adiponectin activates LKB1-AMPK pathway via adiponectin receptors (AdipoR). AMPK activation modulated by  $TNF\alpha$  and adiponectin contributes to the regulation of Pi-induced calcification.

a transcription factor, AICAR-responsive element binding protein (38). It has also been observed that AICAR treatment is able to reduce nuclear factor- $\kappa$ B-regulated transcription, which is activated by  $TNF\alpha$  (39).

Consistent with our findings, it has been recently reported that  $TNF\alpha$  directly suppresses AMPK activation by promoting protein phosphatase 2C (PP2C) activity via TNF receptor-1 (40). PP2C has been proposed as one of modulators of the covalent regulation of AMPK (41). Increased PP2C levels account for the reduced AMPK activity and phosphorylation after  $TNF\alpha$  treatment (40). On the other hand, LKB1 is the well-known, principal upstream kinase of AMPK (42, 43) that is regulated by adiponectin (44). AMPK activation by adiponectin is considered to be mediated by the cell surface receptors adiponectin receptors 1 and 2 (45). Another adiponectin receptor, T-cadherin, has recently been identified (46). In preliminary experiments, we found that all of the three adiponectin receptors were endogenously expressed in HASMC, and Pi did not affect their expression (data not shown). Taking these observations together, we hypothesized the mechanism of regulation by adiponectin and  $TNF\alpha$  on Pi-induced vascular calcification (Fig. 7). However, further intensive investigations are required to elucidate the role of each player in VSMC calcification.

In summary, adiponectin inhibited VSMC calcification and antagonized the stimulatory effect of  $TNF\alpha$ . This action was caused by preventing apoptosis via AMPK activation, followed by restoration of the Gas6-mediated survival pathway. AMPK regulated Gas6 expression at the transcriptional level. AMPK activation regulated by adiponectin and  $TNF\alpha$  in vascular calcification might be a key to the management of cardiovascular disease.

We thank Yuki Ito for technical assistance.

Received July 25, 2007. Accepted December 26, 2007.

Address all correspondence and requests for reprints to: Masahiro Akishita, M.D., Ph.D., Department of Geriatric Medicine, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan. E-mail: akishita-ky@umin.ac.jp.

This study was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture of Japan (No. 18590801 and No. 19590854), Novartis Foundation for Gerontological Research, Kanazawa Medical Research Foundation, Ono Medical Research Foundation, and Takeda Research Foundation.

Disclosure Statement: The authors have nothing to disclose.

## References

- Wexler L, Brundage B, Crouse J, Detrano R, Fuster V, Maddahi J, Rumberger J, Stanford W, White R, Taubert K 1996 Coronary artery calcification: pathophysiology, epidemiology, imaging methods, and clinical implications. A statement for health professionals from the American Heart Association Writing Group. *Circulation* 94:1175–1192.
- Johnson RC, Leopold JA, Loscalzo J 2006 Vascular calcification: pathobiological mechanisms and clinical implications. *Circ Res* 99:1044–1059.
- van Poppel NM, Mattace-Raso FU, Vlietghart R, Grobbee DE, Asmar R, van der Kuip DA, Hofman A, de Feijter FJ, Oudkerk M, Witteman JC 2006 Aortic stiffness is associated with atherosclerosis of the coronary arteries in older adults: the Rotterdam Study. *J Hypertens* 24:2371–2376.
- Arad Y, Spadaro LA, Goodman K, Newstein D, Guerci AD 2000 Prediction of coronary events with electron beam computed tomography. *J Am Coll Cardiol* 36:1253–1260.
- Thompson GR, Partridge J 2004 Coronary calcification score: the coronary-risk impact factor. *Lancet* 363:557–559.
- Son BK, Kozaki K, Hijima K, Ito M, Kojima T, Ota H, Senda Y, Maemura K, Nakano T, Akishita M, Ouchi Y 2006 Statins protect human aortic smooth muscle cells from inorganic phosphate-induced calcification by restoring Gas6-Axl survival pathway. *Circ Res* 98:1024–1031.
- Nakamura Y, Shimada K, Fukuda D, Shimada Y, Ehara S, Hirose M, Kataoka T, Kamimori K, Shimodono S, Kobayashi Y, Yoshiyama M, Takeuchi K, Yoshikawa J 2004 Implications of plasma concentrations of adiponectin in patients with coronary artery disease. *Heart* 90:528–533.

8. Pischon T, Girman CJ, Hotamisligil GS, Rifai N, Hu FB, Rimm EB 2004 Plasma adiponectin levels and risk of myocardial infarction in men. *JAMA* 291:1730-1737
9. Schulze MB, Shai I, Rimm EB, Li T, Rifai N, Hu FB 2005 Adiponectin and future coronary heart disease events among men with type 2 diabetes. *Diabetes* 54:534-539
10. Kojima S, Funahashi T, Sakamoto T, Miyamoto S, Soejima H, Hokamaki J, Kajiwara I, Sugiyama S, Yoshimura M, Fujimoto K, Miyao Y, Suefuji H, Kitagawa A, Ouchi N, Kihara S, Matsuzawa Y, Ogawa H 2003 The variation of plasma concentrations of a novel, adipocyte derived protein, adiponectin, in patients with acute myocardial infarction. *Heart* 89:667
11. Ouchi N, Kihara S, Arita Y, Maeda K, Kuriyama H, Okamoto Y, Hotta K, Nishida M, Takahashi M, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y 1999 Novel modulator for endothelial adhesion molecules: adipocyte-derived plasma protein adiponectin. *Circulation* 100:2473-2476
12. Arita Y, Kihara S, Ouchi N, Maeda K, Kuriyama H, Okamoto Y, Kumada M, Hotta K, Nishida M, Takahashi M, Nakamura T, Shimomura I, Muraguchi M, Ohmoto Y, Funahashi T, Matsuzawa Y 2002 Adipocyte-derived plasma protein adiponectin acts as a platelet-derived growth factor- $\beta$ -binding protein and regulates growth factor-induced common postreceptor signal in vascular smooth muscle cell. *Circulation* 105:2893-2896
13. Maahs DM, Ogden LG, Kinney GL, Wadwa P, Snell-Bergeon JK, Dabelea D, Hokanson JE, Ehrlich J, Eckel RH, Rewers M 2005 Low plasma adiponectin levels predict progression of coronary artery calcification. *Circulation* 111:747-753
14. Ouchi N, Kihara S, Arita Y, Nishida M, Matsuyama A, Okamoto Y, Ishigami M, Kuriyama H, Kishida K, Nishizawa H, Hotta K, Muraguchi M, Ohmoto Y, Yamashita S, Funahashi T, Matsuzawa Y 2003 Adipocyte-derived plasma protein, adiponectin, suppresses lipid accumulation and class A scavenger receptor expression in human monocyte-derived macrophages. *Circulation* 107:1057-1063
15. Maeda N, Takahashi M, Funahashi T, Kihara S, Nishizawa H, Kishida K, Nagaretani H, Matsuda M, Komuro R, Ouchi N, Kuriyama H, Hotta K, Nakamura T, Shimomura I, Matsuzawa Y 2001 PP2A $\gamma$  increases expression and plasma concentrations of adiponectin, an adipose-derived protein. *Diabetes* 50:2094-2099
16. Kern PA, Di Gregorio GB, Lu T, Rassouli N, Ranganathan G 2003 Adiponectin expression from human adipose tissue: relation to obesity, insulin resistance, and tumor necrosis factor- $\alpha$  expression. *Diabetes* 52:1779-1785
17. Maeda N, Shimomura I, Kishida K, Nishizawa H, Matsuda M, Nagaretani H, Furuyama N, Kondo H, Takahashi M, Arita Y, Komuro R, Ouchi N, Kihara S, Tochino Y, Okamoto K, Horie M, Takeda S, Aoyama T, Funahashi T, Matsuzawa Y 2002 Diet-induced insulin resistance in mice lacking adiponectin/ACRP30. *Nat Med* 8:731-737
18. Ouchi N, Kihara S, Arita Y, Okamoto Y, Maeda K, Kuriyama H, Hotta K, Nishida M, Takahashi M, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Funahashi T, Matsuzawa Y 2000 Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF- $\kappa$ B signaling through a cAMP-dependent pathway. *Circulation* 102:1296-1301
19. Shibata R, Sato K, Pimentel DR, Takemura Y, Kihara S, Ohashi K, Funahashi T, Ouchi N, Walsh K 2005 Adiponectin protects against myocardial ischemia-reperfusion injury through AMPK- and COX-2-dependent mechanisms. *Nat Med* 11:1096-1103
20. Ouchi N, Kobayashi H, Kihara S, Kumada M, Sato K, Inoue T, Funahashi T, Walsh K 2004 Adiponectin stimulates angiogenesis by promoting cross-talk between AMP-activated protein kinase and Akt signaling in endothelial cells. *J Biol Chem* 279:1304-1309
21. Kudo N, Barr AJ, Barr RL, Desai S, Lopaschuk GD 1995 High rates of fatty acid oxidation during reperfusion of ischemic hearts are associated with a decrease in malonyl-CoA levels due to an increase in 5'-AMP-activated protein kinase inhibition of acetyl-CoA carboxylase. *J Biol Chem* 270:17513-17520
22. Ouchi N, Kihara S, Funahashi T, Matsuzawa Y, Walsh K 2003 Obesity, adiponectin and vascular inflammatory disease. *Curr Opin Lipidol* 14:561-566
23. Kobayashi H, Ouchi N, Kihara S, Walsh K, Kumada M, Abe Y, Funahashi T, Matsuzawa Y 2004 Selective suppression of endothelial cell apoptosis by the high molecular weight form of adiponectin. *Circ Res* 94:e27-e31
24. Son BK, Kozaki K, Iijima K, Eto M, Nakano T, Akishita M, Ouchi Y 2007 G $\alpha$ 6/ $\beta$ 1-P13K/Akt pathway plays a central role in the effect of statins on inorganic phosphate-induced calcification of vascular smooth muscle cells. *Eur J Pharmacol* 556:1-8
25. Corton JM, Gillespie JG, Hawley SA, Hardie DG 1995 5-Aminoimidazole-4-carboxamide ribonucleoside. A specific method for activating AMP-activated protein kinase in intact cells? *Eur J Biochem* 229:558-565
26. Guadillo O, González-Juanatey JR, Lago F 2007 The emerging role of adipokines as mediators of cardiovascular function: physiologic and clinical perspectives. *Trends Cardiovasc Med* 17:275-283
27. Inoue T, Kotooka N, Morooka T, Komoda H, Uchida T, Aso Y, Inukai T, Okuno T, Node K 2007 High molecular weight adiponectin as a predictor of long-term clinical outcome in patients with coronary artery disease. *Am J Cardiol* 100:569-574
28. Berg AH, Combs TP, Scherer PE 2002 ACRP 30/adiponectin: an adipokine regulating glucose and lipid metabolism. *Trends Endocrinol Metab* 13:84-89
29. Lin LY, Lin CY, Su TC, Liao CS 2004 Angiotensin II-induced apoptosis in human endothelial cells is inhibited by adiponectin through restoration of the association between endothelial nitric oxide synthase and heat shock protein 90. *FEBS Lett* 574:106-110
30. Russell 3rd RR, Li J, Coven DL, Pypaert M, Zechner C, Palmeri M, Giordano FJ, Mu J, Birnbaum MJ, Young LH 2004 AMP-activated protein kinase mediates ischemic glucose uptake and prevents postischemic cardiac dysfunction, apoptosis, and injury. *J Clin Invest* 114:495-503
31. Li J, Miller EJ, Ninomiya-Tsuji J, Russell RR 3rd, Young LH 2005 AMP-activated protein kinase activates p38 mitogen-activated protein kinase by increasing recruitment of p38 MAPK to TAB1 in the ischemic heart. *Circ Res* 97:872-879
32. Marsin AS, Bertrand L, Rider MH, Deprez J, Beauloye C, Vincent MF, Van den Bergh G, Carling D, Hue L 2000 Phosphorylation and activation of heart PFK-2 by AMPK has a role in the stimulation of glycolysis during ischaemia. *Curr Biol* 10:1247-1255
33. Kudo N, Gillespie JG, Kung L, Witters LA, Schulz R, Cianchan AS, Lopaschuk GD 1996 Characterization of 5'-AMP-activated protein kinase activity in the heart and its role in inhibiting acetyl-CoA carboxylase during reperfusion following ischemia. *Biochim Biophys Acta* 1301:67-75
34. Makinde AO, Gamble J, Lopaschuk GD 1997 Upregulation of 5'-AMP-activated protein kinase is responsible for the increase in myocardial fatty acid oxidation rates following birth in the newborn rabbit. *Circ Res* 80:482-489
35. Nagata D, Takeda R, Sata M, Satonaka H, Suzuki E, Nagano T, Hirata Y 2004 AMP-activated protein kinase inhibits angiotensin II-stimulated vascular smooth muscle proliferation. *Circulation* 110:444-451
36. Igata M, Motoshima H, Tsuruzoe K, Kojima K, Matsumura T, Kondo T, Taguchi T, Nakamaru K, Yano M, Kukidome D, Matsumoto K, Toyonaga T, Asano T, Nishikawa T, Arai T 2005 Adenosine monophosphate-activated protein kinase suppresses vascular smooth muscle cell proliferation through the inhibition of cell cycle progression. *Circ Res* 97:837-844
37. Iwasaki Y, Nishiyama M, Taguchi T, Kambayashi M, Asai M, Yoshida M, Nagawara T, Hashimoto K 2007 Activation of AMP-activated protein kinase stimulates proopiomelanocortin gene transcription in AT200 corticotroph cells. *Am J Physiol Endocrinol Metab* 292:E1899-E1905
38. Inoue E, Yamauchi J 2006 AMP-activated protein kinase regulates PEPCK gene expression by direct phosphorylation of a novel zinc finger transcription factor. *Biochem Biophys Res Commun* 351:793-799
39. Solaz-Fuster MC, Gimeno-Alcaniz JV, Casado M, Sanz P 2006 TRIP6 transcriptional co-activator is a novel substrate of AMP-activated protein kinase. *Cell Signal* 18:1702-1712
40. Steinberg GR, Michell BJ, van Denderen BJ, Watt MJ, Carey AL, Fam BC, Andrikopoulos S, Proietto J, Görgün CZ, Carling D, Hotamisligil GS, Febbraio MA, Kay TW, Kemp BE 2006 Tumor necrosis factor- $\alpha$ -induced skeletal muscle insulin resistance involves suppression of AMP-kinase signaling. *Cell Metab* 4:465-474
41. Davies SP, Helps NR, Cohen PT, Hardie DG 1995 5'-AMP inhibits dephosphorylation, as well as promoting phosphorylation, of the AMP-activated protein kinase. Studies using bacterially expressed human protein phosphatase-2C $\alpha$  and native bovine protein phosphatase-2AC. *FEBS Lett* 377:421-425
42. Hawley SA, Boudeau J, Reid JL, Mustard KJ, Udd L, Mäkelä TP, Alessi DR, Hardie DG 2003 Complexes between the LKB1 tumor suppressor, STRAD  $\alpha/\beta$  and MO25  $\alpha/\beta$  are upstream kinases in the AMP-activated protein kinase cascade. *J Biol* 2:28
43. Woods A, Johnston SR, Dickerson K, Leiper FC, Fryer LG, Neumann D, Schlatterer U, Wallimann T, Carlson M, Carling D 2003 LKB1 is the upstream kinase in the AMP-activated protein kinase cascade. *Curr Biol* 13:2004-2008
44. Imai K, Inukai K, Ikegami Y, Awata T, Katayama S 2006 LKB1, an upstream AMPK kinase, regulates glucose and lipid metabolism in cultured liver and muscle cells. *Biochem Biophys Res Commun* 351:595-601
45. Yamauchi T, Kamon J, Ito Y, Tsuchida A, Yokomizo T, Kita S, Sugiyama T, Miyagishi M, Hara K, Tsunoda M, Murakami K, Ohteki T, Uchida S, Takewaka S, Waki H, Tsuno NH, Shibata Y, Terachi Y, Froguel P, Tobe K, Koyasu S, Taira K, Kitamura T, Shimizu T, Nagai R, Kadowaki T 2003 Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. *Nature* 423:762-769
46. Hug C, Wang J, Ahmad NS, Bogan JS, Tsao TS, Lodish HF 2004 T-cadherin is a receptor for hexameric and high-molecular-weight forms of ACP30/adiponectin. *Proc Natl Acad Sci USA* 101:10308-10313

## AUTHOR QUERIES

### AUTHOR PLEASE ANSWER ALL QUERIES

1

A—Journal style does not allow abbreviations in the title. Please confirm or correct expansion of TNF, Gas6, and AMP.

B—Please confirm or correct postal codes added to the affiliations line.

C—Please confirm or correct expansion of TdT.

D—Please confirm or correct expansion of AP1.

E—Please confirm or correct expansion of NF- $\kappa$ B.

F—Please spell out LKB1.

---



## Original article

## Sirt1 modulates premature senescence-like phenotype in human endothelial cells

Hidetaka Ota<sup>a</sup>, Masahiro Akishita<sup>a,\*</sup>, Masato Eto<sup>a</sup>, Katsuya Iijima<sup>a</sup>,  
Masao Kaneki<sup>b</sup>, Yasuyoshi Ouchi<sup>a</sup><sup>a</sup> Department of Geriatric Medicine, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan<sup>b</sup> Department of Anesthesia and Critical Care, Massachusetts General Hospital, Harvard Medical School, 149 Thirteenth Street, Charlestown, MA 02129, USA

Received 6 July 2007; received in revised form 9 August 2007; accepted 10 August 2007

Available online 22 August 2007

## Abstract

Yeast Sir2 plays critical roles in gene silencing, stress resistance and longevity. Mammalian Sirt1 NAD<sup>+</sup>-dependent protein deacetylase, the closest homolog of Sir2, regulates cell cycle, cellular senescence, apoptosis and metabolism, by functional interactions with a number of biological molecules such as p53. To investigate a role of Sirt1 in endothelial dysfunction and premature senescence, we examined the effects of Sirt1 inhibition in human umbilical vein endothelial cells (HUVEC). Sirt1 inhibition by sirtinol, which is a 2-hydroxy-1-naphthaldehyde derivative, or siRNA for Sirt1-induced premature senescence-like phenotype, as judged by increased senescence-associated  $\beta$ -galactosidase (SA- $\beta$ -gal) activity, sustained growth arrest and enlarged and flattened cell morphology at 10 days after the treatment. Sixty-four percent of sirtinol (60  $\mu$ M/L)-treated HUVEC was SA- $\beta$ -gal-positive, whereas only 17% of vehicle-treated cells were positive. Sirt1 inhibition by sirtinol or Sirt1 siRNA increased PAI-1 expression and decreased both protein expression and activity of eNOS. Treatment with sirtinol or Sirt1 siRNA increased acetylation of p53, while p53 expression was unaltered. Impaired epidermal growth factor-induced activation of mitogen-activated protein kinases was associated with Sirt1 inhibition-induced senescence-like growth arrest. Conversely, overexpression of Sirt1 prevented hydrogen peroxide-induced SA- $\beta$ -gal activity, morphological changes and deranged expression of PAI-1 and eNOS. These results showed that Sirt1 inhibition increased p53 acetylation and induced premature senescence-like phenotype in parallel with increased PAI-1 and decreased eNOS expression. Our data suggest that Sirt1 may exert protective effects against endothelial dysfunction by preventing stress-induced premature senescence and deranged expression of PAI-1 and eNOS.

© 2007 Elsevier Inc. All rights reserved.

Keywords: Sirt1; Sirtinol; Premature senescence; Endothelial dysfunction; Oxidative stress

## 1. Introduction

Replicative senescence is the limited ability of mammalian cells to divide when cultured *in vitro* [1]. Cessation of cell division after extended propagation in weeks – or months – culture results from the attrition of telomeres and is termed replicative senescence. In contrast to replicative senescence, stressors such as oxidative stress elicit irreversible growth arrest within just a few days, which is referred to as stress-induced premature senescence (SIPS) [2,3]. Although cellular phenotypes of replicative senescence and SIPS are quite similar or indistinguishable, unlike replicative senescence, SIPS can be

induced independent of telomere attrition. In either type of cellular senescence, it is accompanied by a specific set of changes in cell function, morphology and gene expression [4,5]. Well-established biomarkers of cellular senescence include staining for senescence-associated  $\beta$ -galactosidase (SA- $\beta$ -gal) at pH 6.0 as opposed to endogenous lysosomal enzyme detected at pH 4.0 in normal cells.

Recently, premature senescence of endothelial cells has been proposed to be involved in endothelial dysfunction and atherogenesis [6]. Increased plasminogen activator inhibitor-1 (PAI-1) expression and reduced endothelial nitric oxide synthase (eNOS) activity, which play key roles in endothelial dysfunction and atherogenesis [7], are associated with premature senescence of vascular endothelial cells. Indeed, previous studies have shown that senescence of endothelial cells leads to endothelial

\* Corresponding author. Tel.: +81 3 5800 8832; fax: +81 3 5800 8831.

E-mail address: [akishita-tyk@umin.ac.jp](mailto:akishita-tyk@umin.ac.jp) (M. Akishita).

dysfunction and may contribute to the progression of atherosclerosis [8,9]. Human aortae with atherosclerotic lesion exhibited premature senescence-like phenotype, including SA- $\beta$ -gal activity and increased expression of PAI-1 [10], in association with decreased production of nitric oxide (NO) in endothelial cells [6]. In addition to the role as a prothrombotic factor, the induction of PAI-1 is part of cellular senescence program and has been used as an indicator of cellular senescence in various cell types, including endothelial cells [11,12]. Moreover, NO production [13] and eNOS expression [14,15] are markedly attenuated in senescent endothelial cells. However, the molecular mechanisms underlying premature senescence and endothelial dysfunction remain to be clarified.

Sirt2 (silent information regulator-2), an NAD<sup>+</sup>-dependent histone deacetylase, is highly conserved in organisms ranging from archaea to humans [16,17]. In yeast, Sir2 is responsible for silencing at repeated DNA sequences in mating-type loci, telomeres and rDNA and plays critical roles in DNA repair, stress resistance and longevity [18–20]. Mammalian Sirt1 NAD<sup>+</sup>-dependent protein deacetylase, the closest homolog of Sir2, regulates cell cycle, premature senescence, apoptosis and metabolism by interacting with a number of molecules, including p53, PML, Foxo, Ku70 and PPAR- $\gamma$  [21–26]. A previous study has shown that Sirt1 antagonizes p53-mediated premature senescence in mouse embryo fibroblasts [25]. Recently, we have demonstrated that Sirt1 inhibition induces premature senescence-like growth arrest in human cancer cells [28].

In the present study, to test the hypothesis that Sirt1 plays a role in endothelial premature senescence and dysfunction in vitro, we evaluated the effects of inhibition and overexpression of Sirt1. Here, we demonstrate in human vascular endothelial cells that inhibition of Sirt1-induced premature senescence-like phenotype in parallel with acetylation of p53, and that Sirt1 overexpression reverted premature senescence induced by hydrogen peroxide.

## 2. Materials and methods

### 2.1. Cell culture

Human umbilical vein endothelial cells (HUVEC) were purchased from CAMBREX (Walkersville, MD) and maintained in endothelial growth medium (EGM-2, EGM-2 singleQuots, CAMBREX). Population doubling levels (PDL) were calculated as described previously [29], and all experiments were performed at PDL of 8–9, unless otherwise stated.

### 2.2. Sirt1 inhibition and overexpression

Exponentially proliferating cells were washed three times with growth medium and exposed for 24 h to the indicated concentrations of sirtinol (Calbiochem, San Diego, CA), a cell permeable 2-hydroxy-1-naphthaldehyde derivative and a chemical inhibitor of NAD<sup>+</sup>-dependent protein deacetylases of sirtuins. Trichostatin A (TSA) was purchased from Sigma (St. Louis, MO). After exposure, the cells were washed three times with inhibitor-free medium and cultured for up to additional 9 days.

Proliferating cells were transfected with 200 pmol/L siRNA for Sirt1 (GAT GAA GTT GAC CTC CTC A [22] and TGA AGT GCC TCA GAT ATT A [28]), Sirt2 (sc-40988, Santa Cruz Biotechnology, Inc., CA), or control siRNA (Darmacom, Chicago, IL) using siMPORTER (Upstate, Charlottesville, VA). Sirt1 was overexpressed by transfection with pIRES-Sirt1, which was kindly provided by Dr. R.A. Weinberg [26,27], using jetPEI-HUVEC (Polyplustransfection, Illkirch, France) according to the manufacturer's instruction. Three days after the transfection, HUVEC were treated with 100  $\mu$ mol/L hydrogen peroxide (Sigma) for 1 h and then washed three times with medium and were cultured for up to 10 days. Inhibition of p53 was performed by transfection of 4  $\mu$ mol/L non-targeted oligonucleotide (5'-GGAGCCAGGGGGGAGGG-3') or p53 anti-sense oligonucleotide (5'-CCCTGCTCCCCCTGGCTCC-3').

### 2.3. Senescence-associated $\beta$ -galactosidase (SA- $\beta$ -gal)

#### Staining

At 10 days after the treatment, the proportion of SA- $\beta$ -gal-positive cells was determined as described by Dimri et al. [30].

### 2.4. BrdU incorporation assay

BrdU incorporation was analyzed using a commercially available kit (Roche, Indianapolis).

### 2.5. NOS activity assay

NOS activity of endothelial cells was determined using a NOS assay kit (Calbiochem) according to the manufacturer's instructions.

### 2.6. Immunoblotting

Cells were lysed on ice for 1 h in buffer (50 mmol/L Tris-HCl, pH 7.6, 150 mmol/L NaCl, 1%NP-40, 0.1% SDS, 1 mmol/L dithiothreitol, 1 mmol/L sodium vanadate, 1 mmol/L phenylmethylsulfonyl fluoride, 10  $\mu$ g/mL aprotinin, 10  $\mu$ g/mL leupeptin and 10 mmol/L sodium fluoride). Equal amounts of protein were separated by SDS-polyacrylamide gel electrophoresis and transferred to nitrocellulose membranes. After blocking, the filters were incubated with the following antibodies; anti-p53, anti-acetyl-histone H3 (Lys14), anti-phospho-p44/42 MAPK (Thr202/Tyr204), anti-p44/42 MAPK, anti-phospho-p38 MAPK (Thr180/Tyr182), anti-p38, anti-phospho-JNK (Thr183/Tyr185), anti-JNK (Cell Signaling, Danvers, MA), anti-PAI-1 (Molecular Innovations, Southfield, MI), anti-eNOS (BD Transduction Laboratories, San Jose, CA), anti-Sirt1 (rabbit monoclonal IgG, recognizes human and mouse Sirt1), anti-acetyl-p53 (Lys373/382 and Lys320), anti-acetyl-histone H4 (Lys16) (Upstate), anti-Sirt2 (Santa Cruz) and anti- $\beta$ -actin (Sigma). After washing and incubation with horseradish peroxidase-conjugated anti-rabbit or anti-mouse IgG (Amersham, Piscataway, NJ) for 1 h, the antigen-antibody complexes were visualized by an enhanced chemiluminescence system (Amersham).

## 2.7. Data analysis

Values are shown as mean  $\pm$  SEM in the text and figures. Differences between the groups were analyzed using one-way analysis of variance, followed by Bonferroni test. Probability values less than 0.05 were considered significant.

## 3. Results

### 3.1. Inhibition of Sirt1 induces premature senescence-like phenotype in HUVEC

To investigate whether Sirt1 modulates stress-induced premature senescence-like phenotype in HUVEC, we first examined the effect of Sirt1 inhibition. Sirt1 was inhibited by sirtinol, a specific chemical inhibitor of sirtuins, and Sirt1 siRNA. Sirtinol increased the acetylation of histone H3 (Lys14) and H4 (Lys16), endogenous substrates for Sirt1 (Fig. 1A), as expected. However, the protein expression of Sirt1 was not affected. Knockdown of Sirt1 with siRNA was confirmed by Western blotting (Fig. 1D). Treatment with sirtinol or Sirt1 siRNA induced sustained senescence-like growth arrest for up to 10 days (Fig. 1B). BrdU incorporation was also inhibited by sirtinol or Sirt1 siRNA at 10 days after treatment (Fig. 1C), indicating that the premature senescence-like phenotype and decreased cell number of HUVEC were accompanied by the attenuated DNA synthesis. We examined the effects of sirtinol or Sirt1 siRNA treatment on SA- $\beta$ -gal activity, a characteristic feature of senescence-related growth arrest. Sirtinol or Sirt1 siRNA significantly increased SA- $\beta$ -gal activity compared with

untreated cells or control siRNA at 10 days after treatment; 64% of sirtinol (60  $\mu$ mol/L)-treated HUVEC were SA- $\beta$ -gal-positive, whereas only 17% of vehicle-treated cells were positive (Figs. 2A and B). Sirtinol or Sirt1 siRNA increased SA- $\beta$ -gal activity in a time-dependent manner (data not shown). These cells also exhibited senescence-like morphological changes, that is, enlarged and flattened shapes (Fig. 2A). Next, because sirtinol can inhibit other NAD<sup>+</sup>-dependent protein deacetylases of sirtuin family, especially Sirt2 [32], we examined the effect of Sirt2 inhibition by Sirt2 siRNA on senescence-like phenotype at 10 days after treatment. Sirt2 siRNA successfully eliminated the protein expression of Sirt2 (Fig. 1D) but did not affect the rate of senescence-associated  $\beta$ -galactosidase-positive cells (Fig. 2B).

### 3.2. Senescent-like phenotype resulting from Sirt1 inhibition is mediated by p53

p53 plays a pivotal role in cellular senescence [11]. Previous studies have shown that Sirt1 inhibition promotes stress-induced apoptosis by enhancing p53 acetylation [21,26], while ectopically expressed Sirt1 blocks p53-mediated premature senescence in mouse embryo fibroblasts [25]. Therefore, we examined whether inhibition of Sirt1 affects p53 in HUVECs. Consistent with previous studies [21,25,26], Sirt1 inhibition by sirtinol or Sirt1 siRNA increased the acetylation of p53 after 3 days and the expression of p53 after 5 days (Figs. 3A and C). We also treated HUVEC with TSA, an inhibitor of classes I and II HDACs. Acetylation of p53 was increased at Lys373/382 and Lys320 in HUVECs treated with sirtinol but




Fig. 1. Effects of Sirt1 inhibition on Sirt1 expression, acetylation of histone H3, H4 and cell growth in HUVEC. (A) Western blot analysis for acetylated histone H3 at lysine 14 (K14) and histone H4 at lysine 16 (K16) at 24 h after treatment with sirtinol (30 and 60  $\mu$ mol/L) or nicotinamide (1 and 5 mmol/L), a physiological Sirt1 inhibitor. (B) The effects of sirtinol or Sirt1 siRNA on cell growth for 10 days. (C) BrdU incorporation analysis at 10 days after treatment with sirtinol or Sirt1 siRNA. (D) Western blot analysis for Sirt1 expression at 24 h after treatment with sirtinol or at 72 h after transfection of Sirt1 siRNA. Western blot analysis for Sirt2 expression at 3 days after transfection of Sirt2 siRNA.  $\beta$ -Actin served as a loading control.




Fig. 2. Effects of Sirt1 inhibition on senescence-associated  $\beta$ -galactosidase. (A) Photomicrographs of HUVEC stained for SA- $\beta$ -gal. Population doubling levels (PDLs) for the young and senescent cells are indicated, and sirtinol (60  $\mu$ mol/L), Sirt1 siRNA-treated cells were at PDL 8. (B) SA- $\beta$ -gal-positive cells at 10 days after sirtinol or Sirt1 or Sirt2 siRNA treatment. HUVEC treated with hydrogen peroxide ( $H_2O_2$ , 100  $\mu$ mol/L) for 1 h and the cells at PDL 8 served as positive controls for SA- $\beta$ -gal staining.  $n=3$  for each group. \* $p<0.05$  vs. control.

not with TSA, suggesting that endogenous Sirt1 plays an important role in deacetylation of p53 in HUVECs (Fig. 3D). We further examined whether the effects of Sirt1 inhibition were mediated by p53. We found that blockade of p53 by antisense oligonucleotides reversed the effects of Sirt1 inhibition on senescence associated  $\beta$ -galactosidase activity (Fig. 3E) and senescence-like morphological changes (data not shown) at 10 days after Sirt1 inhibition.

Next, we examined the activation status of mitogen-activated protein kinases (MAPKs) in response to epidermal growth factor (EGF). When untreated with sirtinol or Sirt1 siRNA, upon exposure to EGF, robust phosphorylation (activation) of extracellular-regulated protein kinase (ERK), c-Jun N-terminal kinase (JNK) and p38 MAPK was observed in HUVEC. By contrast, in sirtinol- or Sirt1 siRNA-treated cells, EGF-stimulated phosphorylation of ERK, JNK and p38 MAPK was markedly attenuated (Fig. 3F). The protein expression of ERK, JNK and p38 MAPK did not differ between sirtinol- or Sirt1 siRNA-treated and untreated cells.

### 3.3. Sirt1 inhibition promotes endothelial dysfunction

To investigate further the role of Sirt1 in premature senescence and endothelial function, we examined PAI-1 expression, eNOS expression and NOS activity in HUVEC. Sirt1 inhibition by sirtinol or Sirt1 siRNA increased PAI-1 expression by three-fold and decreased eNOS expression and NOS activity (Figs. 4A, B and D). These changes were similar to those observed in HUVEC that underwent hydrogen peroxide-induced premature senescence and also replicative senescence at 41 PDL (Figs. 4A, C and D). The protein expression of PAI-1 and eNOS were increased and decreased,

respectively, at 3 days after the addition of sirtinol or Sirt1 siRNA and further increased and reduced at 10 days after sirtinol or Sirt1 siRNA treatment (Fig. 5A). Likewise, the activity of NOS was reduced at 3 and 10 days after sirtinol or Sirt1 siRNA addition in a time-dependent manner (Fig. 5B). These results indicate that Sirt1 inhibition-induced premature senescence-like phenotype was associated with endothelial dysfunction.

To investigate the relationship between p53 and these protein expressions, we inhibited p53 by anti-sense oligonucleotide for p53. Anti-sense p53 reduced the expression of p53 and prevented induction of PAI-1 expression by Sirt1 inhibition, as compared with non-targeting oligonucleotide, although reduced eNOS expression was not reverted (Fig. 5C).

### 3.4. Overexpression of Sirt1 blocks oxidative stress-induced premature senescence

Next, we examined whether Sirt1 overexpression can protect HUVEC from oxidative stress-induced premature senescence. As shown previously [3,31], hydrogen peroxide induced premature senescence (Fig. 2B). Overexpression of Sirt1 significantly reduced the number of cells positive for SA- $\beta$ -gal and inhibited senescence-associated morphological changes (Figs. 6A and B). Moreover, overexpression of Sirt1 reversed the changes in eNOS and PAI-1 expression (Fig. 6C) and blocked hydrogen peroxide-induced growth arrest (Fig. 6D). By contrast, when untreated with hydrogen peroxide, overexpression of Sirt1 did not alter cell growth compared with empty vector. In HUVEC transfected with Sirt1, Sirt1 protein expression was increased up to 5 days after treatment with hydrogen peroxide, and then started declining and returned to






Fig. 3. Effects of Sirt1 inhibition on p53 and EGF-stimulated phosphorylation of MAPKs in HUVEC. (A, B) Acetylation of p53 (Ac-p53) at lysine 373/382 (K373/382) and total protein level of p53 were evaluated by Western blotting analysis at 24 h after treatment with sirtinol (A) and at 72 h after transfection with Sirt1 siRNA (B). Treatment with cisplatin (100 μmol/L) for 4 h served as a positive control. β-Actin served as a loading control. (C) Acetylation and total protein levels of p53 for 10 days after treatment with sirtinol or Sirt1 siRNA. (D) Acetylation of p53 at lysine 372/383 and lysine 320 (K372/383 and K320) was evaluated by Western blot analysis at 24 h after treatment with sirtinol (30 and 60 μmol/L) or Trichostatin A (TSA, 20 and 40 nmol/L). (E) Effect of transfection of non-targeted oligonucleotide (NT, 4.0 μmol/L, 5'-GGAGCCAGGGGGAGGG-3') or p53 anti-sense (p53, 4 μmol/L, 5'-CCCTGCTCCCCCTGGCTCC-3') on SA-β-gal activity in sirtinol- or Sirt1 siRNA-treated cells.  $n=3$  for each group. \* $p<0.05$  vs. NT control. (F) EGF-stimulated phosphorylation of MAPKs in sirtinol- or Sirt1 siRNA-treated cells. At 10 days after treatment with sirtinol or Sirt1 siRNA, following overnight serum starvation, the cells were exposed to EGF (50 ng/ml) for 20 min.

normal level at 10 days after hydrogen peroxide treatment (Fig. 6E).

#### 4. Discussion

We found that Sirt1 inhibition by a specific chemical inhibitor of sirtuins, sirtinol or gene knockdown by Sirt1 siRNA induced premature senescence-like phenotype, as judged by representative markers for premature senescence, SA-β-gal activity, enlarged and flattened cell morphology, sustained growth arrest, increased PAI-1 expression and suppressed activation of MAPKs in response to a growth factor, EGF. Another chemical inhibitor of sirtuins, splitomicin, also elicited

similar effects in HUVEC (data not shown). Sirt1 inhibition-induced premature senescence-like growth arrest was associated with acetylation of p53. These results are consistent with previous studies by others and us showing that: (1) overexpression of Sirt1 antagonizes premature senescence through deacetylation of p53 in mouse fibroblasts [25]; and (2) sirtinol induces premature senescence-like phenotype in human cancer cells [28]. Sirtinol is a cell-permeable 2-hydroxy-1-naphthaldehyde derivative that acts as a specific and direct inhibitor of all NAD<sup>+</sup>-dependent protein deacetylases of sirtuin family. Therefore, our results brought by sirtinol might involve the possible influence of other sirtuins, especially Sirt2 [32]. In this regard, we also examined the role of Sirt2 in premature senescence-like




Fig. 4. Effects of Sirt1 inhibition on PAI-1 and eNOS expression in HUVEC. (A, B) PAI-1 and eNOS expressions were evaluated by Western blotting at 10 days after treatment with sirtinol (A) or Sirt1 siRNA (B). Treatment with hydrogen peroxide ( $\text{H}_2\text{O}_2$ , 100  $\mu\text{mol/L}$ ) for 1 h served as a positive control for the phenotype of premature senescence. (C) The expression of PAI-1 and eNOS in young (PDL 8) and senescent (PDL 41) HUVECs. (D) The effects of sirtinol or Sirt1 siRNA treatment on NOS activity. HUVEC at PDL 8 and the cells treated with hydrogen peroxide served as positive controls.  $n=3$  for each group. \* $p<0.05$  vs. control.

phenotype in HUVEC. Little influence of gene knockdown by Sirt2 siRNA suggests that Sirt2 does not play a major role in our model of premature senescence, even though Sirt2 may be a target enzyme of sirtinol. However, there are 7 mammalian sirtuins and several might be involved in human endothelial senescence. Further studies are needed to address this point.

Moreover, overexpression of Sirt1 prevented premature senescence induced by hydrogen peroxide in HUVEC. This finding is in accord with a previous report that overexpression

of Sirt1 enhanced stress resistance against hydrogen peroxide through deacetylation of p53 in human mesangial cells [33]. Taken together, these findings suggest that Sirt1 may play an important role in stress resistance and the modulation of stress-induced premature senescence (SIPS), at least in part, through deacetylation of p53.

Sirtinol- or Sirt1 siRNA-induced premature senescence was accompanied by attenuated response to EGF in terms of activation of MAPKs signaling. A hallmark feature of senescent




Fig. 5. Time-dependent alterations in PAI-1 and eNOS expression by inhibition of Sirt1 in HUVEC. (A, B) PAI-1 and eNOS expression (A) and NOS activity (B) were evaluated after treatment with sirtinol or Sirt1 siRNA. (C) The effect of transfection of non-targeted oligonucleotide (NT oligo) and p53 antisense (4.0  $\mu\text{mol/L}$ ) on PAI-1 and eNOS expression in sirtinol-treated cells. \* $p<0.05$  vs. control.




Fig. 6. Effects of ectopically expressed Sirt1 on hydrogen peroxide-induced premature senescence in HUVEC. (A–C) The effects of Sirt1 overexpression on cell morphology (A), SA-β-gal staining (B) and the expression of PAI-1 and eNOS (C) in HUVEC treated with and without hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>, 100 μmol/L) for 1 h. The effects of Sirt1 overexpression on SA-β-gal-positive cells in hydrogen peroxide-treated HUVEC. \**p* < 0.05 vs. Mock. (D) The effects of Sirt1 overexpression on cell growth. *n* = 3 for each group. (E) Sirt1 protein expression in HUVEC transfected with empty vector and pIRES-Sirt1 during the 10-day culture.

cell is unresponsiveness to cell growth stimuli in induction of *c-fos* as well as cell proliferation. Our results are in agreement with previous studies that growth factor-stimulated activation of the ERK pathway is impaired in senescent human diploid fibroblasts [34].

Endothelial dysfunction has been implicated in the development of cardiovascular diseases, including hypertension and atherosclerosis [35]. Nevertheless, molecular mechanisms responsible for endothelial dysfunction remain to be determined. Recently, SIPS of endothelial cells has been proposed as a contributor to endothelial dysfunction [36,37]. We found that Sirt1 inhibition increased PAI-1 expression and decreased eNOS expression and NOS activity in HUVEC, and that overexpression of Sirt1 reverted hydrogen peroxide-induced alterations in these proteins. Collectively, these findings suggest that Sirt1 may protect from endothelial dysfunction by inhibiting premature senescence and thereby contribute to maintenance of vascular homeostasis. One can reasonably speculate therefore that Sirt1 may exert salutary actions against cardiovascular diseases (e.g., hypertension, atherosclerosis) by preventing SIPS-associated endothelial dysfunction. However, further studies will be required to clarify this point.

p53 is a major substrate for Sirt1, and that deacetylation of p53 regulates cell cycle, cellular senescence and stress resistance in various cell types, including cardiac myocytes [38]. Consistent with previous studies [21,25,26], Sirt1 inhibition by sirtinol or Sirt1 siRNA increased the acetylation of p53 after 3 days and the expression of p53 after 5 days. This result is consistent with the finding by van der Veer et al. [39] that nicotinamide phosphoribosyltransferase-induced and Sirt1-mediated p53 deacetylation was associated with the decreased expression of p53 in vascular smooth muscle cells. A recent

study has revealed that PAI-1 is a direct downstream target gene of p53 and a major player in the induction of cellular senescence [40]. In agreement with this, we found that inhibition of p53 reversed SA-β-gal activity and PAI-1 expression that were induced by Sirt1 inhibition. These results indicate that the acetylation of p53 may play a role in Sirt1 inhibition-induced premature senescence-like phenotype in HUVEC.

Mitosis-related telomere shortening is the critical event for replicative senescence. However, telomere attrition is dispensable for SIPS, which can be induced in a few days. A previous study has shown that changes in telomere length or in telomerase activity were not detected when SIPS was induced by oxidative stress in human endothelial cells [41]. Moreover, a recent study has demonstrated that telomere extension by overexpression of human telomerase does not affect SIPS [2], although it rescues replicative senescence [42]. These data clearly indicate that in contrast to replicative senescence, telomere erosion does not have an important role in the induction of SIPS.

Previous studies have shown senescence-like phenotype in endothelial cells in atherosclerotic lesion in humans and rodents [6,37], as discussed above. Oxidative stress and other stressors have been implicated in the pathogenesis of endothelial dysfunction, hypertension and atherosclerosis [35]. In contrast, there is no direct *in vivo* evidence indicating excessive replication of endothelial cells, which in turn causes replicative senescence, in cardiovascular diseases. In aggregate, it is reasonably conceivable that SIPS rather than genuine replicative senescence may be a major contributor to senescence-like phenotype of endothelial cells in cardiovascular diseases. Therefore, we focused on the role of Sirt1 in SIPS-like phenotype in endothelial cells in the present study.

In summary, our data showed that Sirt1 inhibition increased p53 acetylation and induced SIPS-like phenotype in HUVEC in parallel with increased PAI-1 and decreased eNOS expression, and that Sirt1 overexpression reversed the SIPS induced by oxidative stress. These results suggest that Sirt1 may exert protective effects against endothelial dysfunction by preventing premature senescence and deranged expression of PAI-1 and eNOS in endothelial cells.

#### 4.1. Perspectives

Prevention and reversal of endothelial dysfunction, a major pathogenic factor for cardiovascular diseases, has been an issue of intense investigation for a number of years. Recently, SIPS has been proposed as a contributor to endothelial dysfunction. Nonetheless, it remains largely unknown how stressors such as oxidative stress induces SIPS in endothelial cells. Our data suggest an important role for Sirt1 in the regulation of SIPS. Collectively, these findings highlight Sirt1 as a possible molecular target to prevent and/or treat cardiovascular diseases, including hypertension and atherosclerosis by protecting endothelial cells from SIPS-involved dysfunction. This possibility deserves further investigation.

#### Acknowledgments

We thank Dr. M. Takata (Department of Immunology and Molecular Genetics, Kawasaki Medical School, Okayama, Japan) and R.A. Weinberg for pIRES-Sirt1 plasmid. This work was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Culture and Sports of Japan (18590801).

#### References

- Hayflick L, Moorhead PS. The serial cultivation of human diploid cell strains. *Exp Cell Res* 1961;25:585–621.
- Gorbunova V, Seluanov A, Pereira-Smith OM. Expression of human telomerase (hTERT) does not prevent stress-induced senescence in normal human fibroblasts but protects the cells from stress-induced apoptosis and necrosis. *J Biol Chem* 2002;277:38540–9.
- Frippiat C, Chen QM, Zdanov S, Magalhaes JP, Remacle J, Toussaint O. Subcytotoxic H<sub>2</sub>O<sub>2</sub> stress triggers a release of transforming growth factor-beta 1, which induces biomarkers of cellular senescence of human diploid fibroblasts. *J Biol Chem* 2001;276:2531–7.
- Ben-Porath I, Weinberg RA. When cells get stressed. An integrative view of cellular senescence. *J Clin Invest* 2004;113:8–13.
- Shelton DN, Chang E, Whittier PS, Choi D, Punk WD. Microarray analysis of replicative senescence. *Curr Biol* 1999;9:939–45.
- Minamino T, Miyasuchi H, Yoshida T, Ishida Y, Yoshida H, Komuro I. Endothelial cell senescence in human atherosclerosis: role of telomere in endothelial dysfunction. *Circulation* 2002;105:1541–4.
- Foreman KE, Tang J. Molecular mechanisms of replicative senescence in endothelial cells. *Exp Gerontol* 2003;38:1251–7.
- Burnig KF. The endothelium of advanced arteriosclerotic plaques in humans. *Arterioscler Thromb* 1991;11:1678–89.
- Ross R. The pathogenesis of atherosclerosis—an update. *N Engl J Med* 1986;314:488–500.
- Cavallaro U, Castelli V, Del Monte U, Soria MR. Phenotypic alterations in senescent large-vessel and microvascular endothelial cells. *Mol Cell Biol Res Commun* 2000;4:117–21.
- Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. Oncogenic ras provokes premature cell senescence associated with accumulation of p53 and p16INK4a. *Cell* 1997;88:593–602.
- Comi P, Chiaramonte R, Maier JA. Senescence-dependent regulation of type 1 plasminogen activator inhibitor in human vascular endothelial cells. *Exp Cell Res* 1995;219:304–8.
- Albrecht EW, Stegeman CA, Heeringa P, Henning RH, van Goor H. Protective role of endothelial nitric oxide synthase. *J Pathol* 2003;199:8–17.
- Sato I, Morita I, Kaji K, Ikeda M, Nagao M, Murota S. Reduction of nitric oxide producing activity associated with in vitro aging in cultured human umbilical vein endothelial cell. *Biochem Biophys Res Commun* 1993;195:1070–6.
- Hoffmann J, Haendeler J, Aicher A, Rossig L, Vasa M, Zeiher AM, et al. Aging enhances the sensitivity of endothelial cells toward apoptotic stimuli: important role of nitric oxide. *Circ Res* 2001;89:709–15.
- Braunstein M, Rose AB, Holmes SG, Allis CD, Broach JR. Transcriptional silencing in yeast is associated with reduced nucleosome acetylation. *Genes Dev* 1993;7:592–604.
- Brachmann CB, Sherman JM, Devine SE, Cameron EE, Pillus L, Boeke JD. The SIR2 gene family, conserved from bacteria to humans, functions in silencing, cell cycle progression, and chromosome stability. *Genes Dev* 1995;9:2888–902.
- Rine J, Herakowitz I. Four genes responsible for a position effect on expression from HML and HMR in *Saccharomyces cerevisiae*. *Genetics* 1987;116:9–22.
- Aparicio OM, Billington BL, Gottschling DE. Modifiers of position effect are shared between telomeric and silent mating-type loci in *S. cerevisiae*. *Cell* 1991;66:1279–87.
- Smith JS, Boeke JD. An unusual form of transcriptional silencing in yeast ribosomal DNA. *Genes Dev* 1997;11:241–54.
- Luo J, Nikolaev AY, Imai S, Chen D, Su F, Shiloh A, et al. Negative control of p53 by Sir2alpha promotes cell survival under stress. *Cell* 2001;107:137–48.
- Picard F, Kurtev M, Chung N, Topark-Ngarm A, Senawong T, Machado De Oliveira R, et al. Sirt1 promotes fat mobilization in white adipocytes by repressing PPAR-gamma. *Nature* 2004;429:771–6.
- Cohen HY, Lavu S, Bitterman KJ, Hekking B, Imahiyero TA, Miller C, et al. Acetylation of the C terminus of Ku70 by CBP and PCAF controls Bax-mediated apoptosis. *Mol Cell* 2004;13:627–38.
- Brunet A, Sweeney LB, Sturgill JF, Chua KF, Greer PL, Lin Y, et al. Stress-dependent regulation of FOXO transcription factors by the SIRT1 deacetylase. *Science* 2004;303:2011–5.
- Langley E, Pearson M, Faretta M, Bauer UM, Frye RA, Minucci S, et al. Human SIR2 deacetylates p53 and antagonizes PML/p53-induced cellular senescence. *EMBO J* 2002;21:2383–96.
- Vaziri H, Dessain SK, Ng Eaton E, Imai SI, Frye RA, Pandita TK, et al. hSIR2(SIRT1) functions as an NAD-dependent p53 deacetylase. *Cell* 2001;107:149–59.
- Matsushita N, Takami Y, Kimura M, Tachiiri S, Ishiai M, Nakayama T, et al. Role of NAD-dependent deacetylases SIRT1 and SIRT2 in radiation and cisplatin-induced cell death in vertebrate cells. *Genes Cells* 2005;10:321–32.
- Ota H, Tokunaga E, Chang K, Hikasa M, Iijima K, Eto M, et al. Sirt1 inhibitor, Sirtinol, induces senescence-like growth arrest with attenuated Ras-MAPK signaling in human cancer cells. *Oncogene* 2006;25:176–85.
- Maciag T, Hoover GA, Stemerman MB, Weinstein R. Serial propagation of human endothelial cells in vitro. *J Cell Biol* 1981;91:420–6.
- Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, et al. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci U S A* 1995;92:9363–7.
- Chen QM, Liu J, Merrett JB. Apoptosis or senescence-like growth arrest: influence of cell-cycle position, p53, p21 and bax in H<sub>2</sub>O<sub>2</sub> response of normal human fibroblasts. *Biochem J* 2000;347:543–51.
- Mai A, Massa S, Lavu S, Pezzi R, Simeoni S, Ragno R, et al. Design, synthesis, and biological evaluation of sirtinol analogues as class III

- histone/protein deacetylase (Sirtuin) inhibitors. *J Med Chem* 2005;48: 7789–95.
- [33] Kurue S, Haneda M, Kanasaki K, Sugimoto T, Araki S, Isono M, et al. Silent information regulator 2 (SIRT1) attenuates oxidative stress-induced mesangial cell apoptosis via p53 deacetylation. *Free Radic Biol Med* 2006; 40:2175–82.
- [34] Torres C, Francis MK, Lorenzini A, Tresini M, Cristofalo VJ. Metabolic stabilization of MAP kinase phosphatase-2 in senescence of human fibroblasts. *Exp Cell Res* 2003;290:195–206.
- [35] Feletou M, Vanhoutte PM. Endothelial dysfunction: a multifaceted disorder (The Wiggers Award Lecture). *Am J Physiol: Heart Circ Physiol* 2006;291:H985–H1002.
- [36] Minamino T, Miyatachi H, Yoshida T, Tateno K, Komuro I. The role of vascular cell senescence in atherosclerosis. Antisenescence as a novel therapeutic strategy for vascular aging. *Curr Vasc Pharmacol* 2004;2: 141–8.
- [37] Brodsky SV, Gealekman O, Chen J, Zhang F, Togashi N, Crabtree M, et al. Prevention and reversal of premature endothelial cell senescence and vasculopathy in obesity-induced diabetes by ebselen. *Circ Res* 2004;94: 377–84.
- [38] Alcendor RR, Kirshenbaum LA, Imai S, Vatner SF, Sadoshima J. Silent information regulator 2alpha, a longevity factor and class III histone deacetylase, is an essential endogenous apoptosis inhibitor in cardiac myocytes. *Circ Res* 2004;95:971–80.
- [39] van der Veer E, Ho C, O'Neil C, Barbosa N, Scott R, Cregan SP, et al. Extension of human cell lifespan by nicotinamide phosphoribosyltransferase. *J Biol Chem* 2007;282:10841–5.
- [40] Kortlever RM, Higgins PJ, Bernards R. Plasminogen activator inhibitor-1 is a critical downstream target of p53 in the induction of replicative senescence. *Nat Cell Biol* 2006;8:877–84.
- [41] Chen J, Brodsky SV, Goligorsky DM, Hampel DJ, Li H, Gross SS, et al. Glycated collagen I induces premature senescence-like phenotypic changes in endothelial cells. *Circ Res* 2002;90:1290–8.
- [42] Bodnar AG, Ouellette M, Frolkis M, Holt SE, Chiu CP, Morin GB, et al. Extension of life-span by introduction of telomerase into normal human cells. *Science* 1998;279:349–52.

### EFFECTS OF AGE AND SEX ON PLASMA ADRENOMEDULLIN LEVELS IN PATIENTS WITH OBSTRUCTIVE SLEEP APNEA SYNDROME

*To the Editor:* It has been recognized that obstructive sleep apnea syndrome (OSAS) is one of the risk factors of cardiovascular disorders, including hypertension, ischemic heart disease, and cerebrovascular diseases.<sup>1-3</sup> Although the incidence of sleep apnea increases with age, the pathological roles of OSAS have not been fully established. OSAS-induced hypoxic stress and oxidative stress increase circulating inflammatory mediators, including adhesion molecules, inflammatory cytokines, and high-sensitivity C-reactive protein, leading to hypertension and cardiovascular events.<sup>4,5</sup> The stress and its related inflammatory molecules are implicated in the production of adrenomedullin (AM), which is a potent endothelial-derived vasodilator.<sup>6</sup> Circulating AM levels are higher in adults with untreated OSAS than in adults without OSAS.<sup>7</sup> Because plasma AM is closely correlated with pulse wave velocity and atherosclerosis progression in middle-aged and elderly patients, higher levels of AM may be a surrogate marker for hypertension and the progression of atherosclerosis.<sup>8</sup> Furthermore, treatment with nasal continuous positive airway pressure (nCPAP) mostly reversed the higher AM levels in subjects with OSAS. Thus, OSAS treatment may prevent atherosclerosis and cardiovascular events, although circulating AM levels and therapeutic response to OSAS have not been examined in elderly people. Furthermore, female sex hormones increase AM-induced vasodilation by increasing the expression of AM2 receptor components in rats.<sup>9</sup> Sex may affect circulating AM levels in humans.

Age and sex differences in AM levels in patients with OSAS were examined. Eighty middle-aged (aged 40-60)

and 80 elderly (aged 60) patients were compared with OSAS and 80 middle-aged and 80 elderly age- and body mass index (BMI)-matched subjects without OSAS. The patients had to fulfill the following criteria: absence of renal and renovascular hypertension, systolic blood pressure (BP) greater than 160 mmHg or diastolic BP greater than 95 mmHg, chronic renal and hepatic diseases, and diabetes mellitus. Patients who smoked or had systemic infections at the time of the study or within 4 weeks before the study were excluded. No patients were being treated with antihypertensive agents. These subjects were examined using polysomnography; subjects with an apnea-hypopnea index (AHI) less than five were controls, and those with an AHI of five or greater were determined to have OSAS. To assess OSAS-induced hypoxia quantitatively, the oxyhemoglobin desaturation index (ODI) was used in this study as previously described.<sup>4</sup> ODI was defined as  $ODI = \Sigma(90 - \text{oxygen saturation})/t$ , where  $t$  is time of desaturation (hours).<sup>4</sup>

Circulating AM levels and sleep study variables were compared. Then 3 months of nCPAP treatment was performed in the patients. Peripheral blood was obtained from the subjects at 7:30 a.m. to 8:00 a.m. before and after the 3 months treatment with nCPAP. The AM was measured using a specific radioimmunoassay.

There were no significant differences in BMI between the patients and controls in each age group, whereas AHI in subjects OSAS was markedly greater than in controls (Table 1). There were no significant differences in BP or metabolic indices. The AHI values in elderly ( $50.1 \pm 3.2$  events/h) and middle-aged ( $51.6 \pm 3.0$ ) subjects with OSAS were considerably greater than in age-matched controls ( $3.8 \pm 0.3$  and  $3.6 \pm 0.4$ , respectively). There were significant differences in baseline ODI between patients with OSAS and controls, suggesting that the patients with OSAS were exposed to a

**Table 1. Circulating Adrenomedullin (AM) Levels and Other Variables in Elderly and Middle-Aged Patients with Obstructive Sleep Apnea Syndrome (OSAS) and Controls**

Characteristic	Middle-Aged Subjects with OSAS (n = 80)	Elderly Subjects with OSAS (n = 80)	Middle-Aged Controls (n = 80)	Elderly Controls (n = 80)
	Mean $\pm$ Standard Error			
Age	46.8 $\pm$ 2.2	65.8 $\pm$ 2.2	45.1 $\pm$ 2.2	64.6 $\pm$ 2.1
Body mass index	33.4 $\pm$ 0.9	32.1 $\pm$ 0.9	32.8 $\pm$ 1.1	31.5 $\pm$ 1.1
Systolic blood pressure, mmHg	136.1 $\pm$ 3.7	139.1 $\pm$ 4.7	135.6 $\pm$ 4.1	137.1 $\pm$ 3.7
Diastolic blood pressure, mmHg	82.1 $\pm$ 3.2	80.1 $\pm$ 3.2	78.9 $\pm$ 3.8	78.1 $\pm$ 2.8
Total cholesterol, mg/dL	202.9 $\pm$ 7.9	199.1 $\pm$ 6.9	202.9 $\pm$ 7.9	198.3 $\pm$ 10.8
High-density lipoprotein cholesterol, mg/dL	43.0 $\pm$ 2.2	41.0 $\pm$ 2.1	43.6 $\pm$ 2.3	42.2 $\pm$ 2.1
Triglyceride, mg/dL	144.1 $\pm$ 10.7	137.1 $\pm$ 12.7	140.1 $\pm$ 10.7	132.0 $\pm$ 10.9
Fasting plasma glucose, mg/dL	98.4 $\pm$ 1.3	94.6 $\pm$ 1.3	97.6 $\pm$ 1.4	90.3 $\pm$ 1.1
Hemoglobin A1c, %	5.7 $\pm$ 0.1	5.8 $\pm$ 0.1	5.6 $\pm$ 0.1	5.27 $\pm$ 0.1
Total sleep time, min	368.1 $\pm$ 20.3*	348.1 $\pm$ 20.3*	440.3 $\pm$ 20.9	414.3 $\pm$ 20.9
Apnea-hypopnea index, events/h	51.6 $\pm$ 3.0*	50.1 $\pm$ 3.2*	3.8 $\pm$ 0.3	3.6 $\pm$ 0.4
Lowest oxygen saturation, %	67.2 $\pm$ 2.1*	68.9 $\pm$ 3.0*	95.8 $\pm$ 0.5	94.1 $\pm$ 0.5
Oxyhemoglobin desaturation index	2.45 $\pm$ 0.32*	2.41 $\pm$ 0.36*	0.02 $\pm$ 0.01	0.02 $\pm$ 0.01
Arousal index/h	42.4 $\pm$ 3.1*	40.2 $\pm$ 2.2*	8.3 $\pm$ 3.1	8.3 $\pm$ 3.1
AM level, pg/mL	49.1 $\pm$ 8.7	51.1 $\pm$ 4.1*	24.6 $\pm$ 1.9	25.7 $\pm$ 2.1

Note: There were 40 men and 40 women in each group.

\* $P < .001$  versus control group.

significantly greater degree of hypoxia than the control subjects. The magnitude of ODI was not different between middle-aged and elderly patients with OSAS.

Circulating AM levels in middle-aged and elderly patients with OSAS were significantly greater than in the age- and BMI-matched controls, although neither age nor sex affected them (Table 1). nCPAP treatment significantly decreased the higher levels of circulating AM in the patients irrespective of age and sex. After 3 months of treatment with nCPAP, AM levels in elderly patients ( $26.5 \pm 2.4$  pg/mL) were not different from those of middle-aged patients ( $24.7 \pm 2.1$  pg/mL).

These results indicated that plasma AM levels were higher in middle-aged and elderly patients with OSAS and could be decreased with nCPAP treatment, regardless of age and sex. The augmented increase in AM caused by severe nocturnal hypoxemia and oxidative stress due to OSA may overcome the age-dependent increase of AM levels in middle-aged and elderly patients with OSAS. Because AM is reported to induce cell surface expression of adhesion molecules, including E-selectin, vascular cell adhesion molecule-1, and intercellular adhesion molecule-1 (ICAM-1), on human endothelial cells, the higher level of AM is one of the mechanisms of higher levels of ICAM-1 in patients with OSAS.<sup>10</sup> The current study also indicates that treatment with nCPAP may be effective for the prevention of cardiovascular complications in elderly patients with OSAS.

Shinji Teramoto, MD  
Yasuhiro Yamaguchi, MD  
Hiroshi Yamamoto, MD  
Yoko Hanaoka, MD  
Masaki Ishii, MD  
Shinichiro Hibi, MD  
Department of Geriatric Medicine

Haruki Kume, MD  
Department of Urology

Masahiro Akishita, MD  
Yasuyoshi Ouchi, MD  
Department of Geriatric Medicine  
Graduate School of Medicine  
University of Tokyo  
Tokyo, Japan

#### ACKNOWLEDGMENTS

This work was supported in part by a research grant from Mitsui Life Social Welfare Foundation Japan fund and by research grant from Mitsukoshi Health and Welfare Foundation in Japan. There is no conflict of interest regarding the study.

**Conflict of Interest:** The editor in chief has determined that the authors have no conflict of interest regarding this study.

**Author Contributions:** study concept and design, Shinji Teramoto, MD, Yasuhiro Yamaguchi; acquisition of subjects and/or data analysis, Hiroshi Yamamoto, MD, Yoko Hanaoka, MD, Masaki Ishii, MD, Shinichiro Hibi, MD; interpretation of data, Haruki Kume, MD, Masahiro Akishita, MD; preparation of manuscript, Yasuyoshi Ouchi, MD.

**Sponsor's Role:** No sponsor in this study.

#### REFERENCES

- Chhajed PN, Tamm M, Strobel W. Sleep apnea and heart disease. *N Engl J Med* 2006;354:1086-1089.
- Teramoto S, Kume H, Matsuse T. Ambulatory blood pressure after sleep apnoea treatment. *Lancet* 2002;360:341-342.
- Teramoto S, Ohga E, Ouchi Y. Obstructive sleep apnoea. *Lancet* 1999;354:1213-1214.
- Ohga E, Nagase T, Tomita T et al. Increased levels of circulating ICAM-1, VCAM-1, and L-selectin in obstructive sleep apnoea syndrome. *J Appl Physiol* 1999;87:10-14.
- Teramoto S, Yamamoto H, Ouchi Y. Increased C-reactive protein and increased plasma interleukin-6 may synergistically affect the progression of coronary atherosclerosis in obstructive sleep apnoea syndrome. *Circulation* 2003;107:E40.
- Sugo S, Minamino N, Kangawa K et al. Endothelial cells actively synthesize and secrete adrenomedullin. *Biochem Biophys Res Commun* 1994;201:1160-1166.
- Schulz R, Flototto C, Jahn A et al. Circulating adrenomedullin in obstructive sleep apnoea. *J Sleep Res* 2006;15:89-95.
- Kita T, Kitamura K, Hashida S et al. Plasma adrenomedullin is closely correlated with pulse wave velocity in middle-aged and elderly patients. *Hypertens Res* 2003;26:887-893.
- Ross GR, Chauhan M, Gangula PR et al. Female sex steroids increase adrenomedullin-induced vasodilation by increasing the expression of adrenomedullin2 receptor components in rat mesenteric artery. *Endocrinology* 2006;147:389-396.
- Yoshii T, Iwai M, Li Z et al. Regression of atherosclerosis by amlodipine via anti-inflammatory and anti-oxidative stress actions. *Hypertens Res* 2006;29:457-466.

#### NATURAL THERAPIES—WHEN IGNORANCE IS NOT BLISS!!

*To the Editor:* A 72-year-old Caucasian man with history of hypertension, dyslipidemia, depression, gastroesophageal reflux disease, and mild obesity presented to the hospital with intractable singultus (hiccups) for approximately 9 days, associated with anorexia and epigastric pain. He had presented to his primary care provider the day before admission and was given prochlorperazine without relief. His daily medications were metoprolol, furosemide, fluvastatin, gemfibrozil, paroxetine, and omeprazole. A thorough history revealed that he had added 2 tablespoons daily of acetic acid (household vinegar) to his routine for the previous 2 weeks after reading an article in a health magazine that indicated that vinegar helps decrease food intake (by promoting satiety) and lowers cholesterol. Physical examination was unremarkable except for mild epigastric tenderness.

Computerized tomography of the abdomen with oral and intravenous contrast was suggestive of acute pancreatitis, and amylase and lipase levels were elevated, at 876 and 1,187 U/L, respectively. Other laboratory parameters were normal. Brain imaging and endoscopy were negative, and furosemide and paroxetine were discontinued on admission. The patient received nothing by mouth and was maintained with intravenous fluids for 2 days. He experienced complete resolution of the pancreatitis and the hiccups.

Vinegar has a long history in medicine, including use by the father of modern medicine, Hippocrates, as an agent to fight acute infections and chronic coughs. Some modern studies support antimicrobial effects with food preparation; others claim it is a nematocyst inhibitor and protective in stings by some jellyfish. Recently, vinegar has been identified as an antiglycemic in subjects with and without glucose intolerance. In addition, it has been shown to increase satiety in healthy patients.<sup>1</sup>

There have been various adverse effects of vinegar described in the literature. Some of these include corrosive

## Original Article

## Association of Plasma Dehydroepiandrosterone-Sulfate Levels with Endothelial Function in Postmenopausal Women with Coronary Risk Factors

Masahiro AKISHITA<sup>1)</sup>, Masayoshi HASHIMOTO<sup>2)</sup>, Yumiko OHIKE<sup>1)</sup>,  
Sumito OGAWA<sup>1)</sup>, Katsuya IJIMA<sup>1)</sup>, Masato ETO<sup>1)</sup>, and Yasuyoshi OUCHI<sup>1)</sup>

Age-related decline of plasma dehydroepiandrosterone-sulfate (DHEA-S) levels may be associated with the risk of cardiovascular disease in women. We investigated whether plasma DHEA-S levels are related to endothelial function in postmenopausal women with coronary risk factors. One hundred and fifteen postmenopausal women (mean age±SD: 57±5 years; range: 48–65 years) who underwent measurement of flow-mediated vasodilation (FMD) of the brachial artery using ultrasonography were enrolled. Plasma hormone levels were determined in the morning after a 14-h fast, and the relationship between hormone levels and FMD was analyzed. DHEA-S was significantly correlated with %FMD ( $r=0.392$ ,  $p<0.001$ ), while estradiol, total testosterone and cortisol were not. %FMD in the highest quartile of DHEA-S was 1.8-fold higher than that in the lowest quartile ( $5.3\pm 1.3$  vs.  $2.9\pm 2.0$  [means±SD],  $p<0.01$ ). Multiple regression analysis revealed that DHEA-S was related to %FMD independent of age, body mass index, hypertension, hyperlipidemia, diabetes mellitus and smoking ( $\beta=0.344$ ,  $p<0.01$ ), and was itself independent of age, body mass index, systolic blood pressure, total cholesterol, high-density lipoprotein (HDL) cholesterol, fasting plasma glucose and smoking ( $\beta=0.291$ ,  $p<0.05$ ). In conclusion, plasma DHEA-S levels were weakly but significantly related to endothelial function in postmenopausal women independent of other coronary risk factors, suggesting a protective effect of DHEA on the endothelium. (*Hypertens Res* 2008; 31: 69–74)

**Key Words:** endothelium, vasodilation, risk factor, man, nitric oxide

### Introduction

Plasma levels of dehydroepiandrosterone-sulfate (DHEA-S), the most abundant circulating steroid that is secreted from the adrenal cortex, decline with advancing age in men and women (1, 2). The age-related decrease in DHEA-S concentrations has often been associated with the pathological processes of aging, such as osteoporosis, depression and

dementia (3, 4). A number of studies have investigated the link between DHEA-S and cardiovascular disease, though with inconsistent results (5–7). Among them, Haffner *et al.* have shown that low DHEA-S levels predicted ischemic heart disease mortality in diabetic women (8). The association of low DHEA-S levels with carotid artery atherosclerosis (9), obesity (10) and decreased diurnal blood pressure variability (11) also suggests a vasoprotective role of DHEA(-S) in women. Furthermore, experimental studies showing endothe-

From the <sup>1)</sup>Department of Geriatric Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan; and <sup>2)</sup>Department of General Internal Medicine, Kobe University School of Medicine, Kobe, Japan.

This study was supported by Health and Labour Sciences Research Grants (H17-Chojo-046) from the Ministry of Health, Labour and Welfare of Japan, by the Gerontology Small Research Grant Program from the Division of Project Coordination, The University of Tokyo and by grants from the NOVARTIS Foundation for Gerontological Research and the Yamaguchi Endocrine Research Association.

Address for Reprints: Masahiro Akishita, M.D., Ph.D., Department of Geriatric Medicine, Graduate School of Medicine, the University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan. E-mail: akishita-cky@umin.ac.jp

Received January 16, 2007; Accepted in revised form August 3, 2007.



**Table 1. Characteristics of Study Subjects (N=115)**

Age, years	57±5	[48–65]
Body mass index, kg/m <sup>2</sup>	23.3±4.0	[17.6–35.0]
Risk factors		
Hypertension, n (%)	35 (30)	
Hyperlipidemia, n (%)	71 (62)	
Diabetes mellitus, n (%)	20 (17)	
Current smoker, n (%)	17 (15)	
Hemodynamic and vascular measurements		
Systolic blood pressure, mmHg	123±17	[93–170]
Diastolic blood pressure, mmHg	70±10	[52–100]
%FMD	4.8±2.4	[0.0–12.7]
%NTG	14.0±4.5	[4.1–22.5]
Carotid IMT, mm	0.92±0.22	[0.46–1.45]
Blood chemistry and hormones		
Total cholesterol, mmol/L	5.72±1.01	[3.73–8.96]
HDL cholesterol, mmol/L	1.66±0.42	[0.77–2.87]
Triglycerides, mmol/L	1.18±0.65	[0.36–3.49]
Fasting plasma glucose, mmol/L	5.42±1.22	[4.00–10.43]
Hemoglobin A1c, %	5.5±1.0	[4.3–9.6]
Estradiol, pmol/L	35±26	[18–160]
Testosterone, nmol/L	0.85±0.43	[0.21–2.01]
DHEA-S, μmol/L	2.28±1.07	[0.37–5.02]
Cortisol, nmol/L	316±121	[110–728]

Values except risk factors are expressed as mean±SD [range]. %FMD, percent flow-mediated dilation of brachial artery; %NTG, percent nitroglycerin-induced dilation of brachial artery; IMT, intima-media thickness of common carotid artery; HDL, high-density lipoprotein; DHEA-S, dehydroepiandrosterone-sulfate.

lium-dependent (12) and -independent (13) vasodilating effects of DHEA(-S) led us to hypothesize that postmenopausal women with low plasma DHEA-S levels would have impaired vasomotor function.

To test this hypothesis, we conducted a cross-sectional survey of 115 postmenopausal women by examining flow-mediated dilation (FMD) of the brachial artery and plasma sex hormones, and showed that low DHEA-S levels were associated with endothelial dysfunction.

## Methods

### Subjects

One hundred and fifteen postmenopausal women who underwent examination of vasomotor function of the brachial artery and intima-media thickness (IMT) of the carotid artery in our department were enrolled. The subjects were referred to our department to check their cardiovascular disease or risks. All of them were in chronic stable condition. A history was taken, and physical examination and laboratory tests were performed in all subjects. Subjects with a history of cardiovascu-

lar disease, including stroke, coronary heart disease, congestive heart failure and peripheral arterial disease, malignancy, overt endocrine disease or administration of steroid hormones, were excluded. The postmenopausal status of each subject was confirmed by the fact that at least 12 months had passed since her last menses and by the measurement of follicular stimulating hormone and estradiol. The characteristics of the study subjects are shown in Table 1.

Seventy-three percent of the subjects had one or more classical coronary risk factors, such as hypertension, hyperlipidemia, diabetes mellitus or current smoking. Hypertension, hyperlipidemia and diabetes mellitus were considered to be present based on the published diagnostic criteria (14–16) or if the subjects were taking any medications for these diseases. Eighty-six percent of the hypertensive subjects were treated: 75% with calcium antagonists, 19% with angiotensin-converting enzyme inhibitors, 10% with diuretics and 7% with β blockers. Eighty-six percent of the hyperlipidemic subjects were treated with statins, and 75% of the diabetic subjects were treated with oral hypoglycemic agents. None of the study subjects were taking nitrates. Each subject gave written informed consent before enrollment in this study. The study protocol was approved by the ethics committee of the Graduate School of Medicine, the University of Tokyo.

### Vascular Measurement

Vasomotor function of the brachial artery was evaluated using an ultrasound machine according to the method described previously (17). Briefly, endothelium-dependent flow-mediated vasodilation (%FMD) was measured as the maximal percent change of the vessel diameter after reactive hyperemia. The subjects were examined in the morning after a 14-h overnight fast with no medication. Subsequently, endothelium-independent nitroglycerin-induced vasodilation (%NTG) was measured as the maximal percent change of the vessel diameter after sublingual administration of nitroglycerin spray (0.3 mg; Toa Eiyo Co., Tokyo). Carotid IMT was evaluated using an ultrasound machine as described previously (17).

### Plasma Hormones

Blood sampling was performed in the morning of the vascular measurement after a 14-h overnight fast, and plasma was stored at -80°C until assay. Plasma estradiol, testosterone (total testosterone), DHEA-S and cortisol concentrations were determined using sensitive radioimmunoassays by a commercial laboratory (SRL Inc., Tokyo, Japan). The intra-assay coefficients of variation for these measurements were less than 5%.

Because plasma was deep-frozen for 3–7 years, we checked the change in titers using the stored samples, which had been measured at sampling 5–7 years before. The Pearson's correlation coefficients between the two measurements were 0.965

Table 2. Pearson's Correlation Coefficients between Age, Vascular Measurements and Plasma Hormones

	Age	%FMD	Carotid IMT
Estradiol	-0.108	0.041	-0.136
Testosterone	0.061	0.019	-0.088
DHEA-S	-0.198*	0.392 <sup>†</sup>	-0.187*
Cortisol	0.074	0.140	0.064

%FMD, percent flow-mediated dilation of brachial artery; IMT, intima-media thickness of common carotid artery. <sup>†</sup> $p < 0.001$ , \* $p < 0.05$ .

for estradiol ( $n=34$ ), 0.976 for testosterone ( $n=20$ ), 0.991 for DHEA-S ( $n=15$ ) and 0.937 for cortisol ( $n=16$ ), indicating that there was no significant change in plasma titers in our frozen samples.

#### Data Analysis

Pearson's simple correlation coefficients between age, vascular measurements and plasma hormones were determined. Standardized regression coefficients from multiple regression analysis of vascular measurements in relation to age, coronary risk factors and plasma hormones were determined. Differences between the groups were analyzed using one-factor ANOVA, followed by Newman-Keuls' test. A value of  $p < 0.05$  was considered statistically significant. Statistical analyses were performed using the SPSS ver. 11.0 software.

### Results

#### Changes in Plasma Hormones and Vascular Measurements According to Age and Coronary Risk Factors

Plasma levels of DHEA-S declined with age, while those of estradiol, testosterone and cortisol did not significantly change according to age (Table 2). %FMD decreased slightly but not significantly ( $r = -0.166$ ,  $p = 0.08$ ), while carotid IMT increased significantly ( $r = 0.337$ ,  $p < 0.01$ ) with advancing age.

The subjects with hypertension, hyperlipidemia or diabetes showed impaired %FMD compared to those without these diseases (hypertension,  $4.0 \pm 2.4$  vs.  $5.1 \pm 2.2$ ; hyperlipidemia,  $4.3 \pm 2.3$  vs.  $5.6 \pm 2.2$ ; diabetes mellitus,  $4.0 \pm 2.8$  vs.  $5.2 \pm 2.2$ ;  $p < 0.05$ ). %FMD in the patients taking antihypertensive agents, statins or hypoglycemic agents was comparable to or smaller than that in the patients without medical agents (hypertension,  $4.0 \pm 2.5$  vs.  $4.3 \pm 2.3$ , n.s.; hyperlipidemia,  $4.1 \pm 2.2$  vs.  $5.1 \pm 2.4$ , n.s.; diabetes,  $3.5 \pm 2.7$  vs.  $6.5 \pm 1.5$ ,  $p < 0.05$ ), suggesting that the favorable effects of medical treatment on endothelial function, if present, might have been lost in patients with a long history of coronary risk factors. In contrast, none of the plasma hormones were significantly

Table 3. Age-Adjusted Regression Coefficients between Vascular Measurements and Plasma Hormones

	%FMD	Carotid IMT
Estradiol	0.023	-0.098
Testosterone	0.029	-0.117
DHEA-S	0.366 <sup>†</sup>	-0.146
Cortisol	0.156	-0.062

%FMD, percent flow-mediated dilation of brachial artery; IMT, intima-media thickness of common carotid artery. Standardized regression coefficients by multiple regression analyses with %FMD or carotid IMT as a dependent variable and age and each of the hormones as independent variables are shown. <sup>†</sup> $p < 0.01$ .

changed according to coronary risk factors or medications (data not shown).

#### Relationship between Plasma Hormones and Vascular Measurements

First, simple correlation coefficients between plasma hormones and vascular measurements were determined. As shown in Table 2, %FMD was positively correlated with DHEA-S, whereas carotid IMT was negatively correlated with DHEA-S. There was no significant correlation between vascular measurements and other steroid hormones.

Next, age-adjusted regression coefficients were determined, because age was correlated with some of the hormones and vascular measurements, as mentioned above. The results showed that none of the hormones were significantly related to carotid IMT (Table 3). In contrast, DHEA-S was significantly related to %FMD, independent of age. Furthermore, the age-adjusted regression coefficients between DHEA-S and %FMD were 0.374 in the subjects with no coronary risk factor or medication ( $p < 0.05$ ,  $n = 31$ ) and 0.399 in those with coronary risk factors ( $p < 0.05$ ,  $n = 84$ ), most of whom were taking some medications.

Finally, multiple regression analyses were performed to exclude the influence of coronary risk factors on the relationship between hormones and %FMD. As shown in Table 4, DHEA-S was related to %FMD, independent of age, body mass index, hypertension, hyperlipidemia, diabetes mellitus and current smoking (Model 1), and were independent of age, body mass index, systolic blood pressure, total cholesterol, high-density lipoprotein (HDL) cholesterol, fasting plasma glucose and current smoking (Model 2). Furthermore, the relationship between DHEA-S and %FMD was significant after addition of carotid IMT to the model (Model 3), suggesting that the relationship was not attributable to the effect of DHEA(-S) on the development of subclinical atherosclerosis. Also, the statistical result was unchanged after addition of nitroglycerin-induced dilation of the brachial artery to the model (Model 4), indicating that DHEA-S is related to endothelial function independent of arterial compliance. Estradiol,

Table 4. Regression Coefficients between %FMD and Plasma Hormones Adjusted for Coronary Risk Factors

	Model 1	Model 2	Model 3	Model 4
Estradiol	0.065	0.042	0.025	0.094
Testosterone	0.036	0.052	0.028	0.148
DHEA-S	0.344 <sup>†</sup>	0.291 <sup>*</sup>	0.263 <sup>*</sup>	0.250 <sup>*</sup>
Cortisol	0.150	0.127	0.099	0.110

Standardized regression coefficients by multiple regression analyses with %FMD as a dependent variable and coronary risk factors (covariates used in each analysis are listed below) and each of the hormones as independent variables are shown. <sup>†</sup> $p < 0.01$ , <sup>\*</sup> $p < 0.05$ . Model 1: age, body mass index, hypertension, hyperlipidemia, diabetes mellitus, and current smoking. Model 2: age, body mass index, systolic blood pressure, total cholesterol, high-density lipoprotein cholesterol, fasting plasma glucose and current smoking. Model 3: Model 2 plus carotid intima-media thickness. Model 4: Model 2 plus percent nitroglycerin-induced dilation of brachial artery. %FMD, percent flow-mediated dilation of brachial artery.

testosterone and cortisol were not significantly related to %FMD in similar multivariate analyses (Table 4). As shown in Fig. 1, %FMD showed a stepwise increment according to quartiles of DHEA-S, and %FMD in the highest quartile of free testosterone was 1.8-fold higher than that in the lowest quartile ( $5.3 \pm 1.3$  vs.  $2.9 \pm 2.0$ ,  $p < 0.01$ ).

All the variables except hormones were mandatorily incorporated into the multiple regression models shown in Table 4, because they are classical coronary risk factors. In forward and backward stepwise models including age, body mass index, hypertension, hyperlipidemia, diabetes mellitus, smoking and the four steroid hormones, DHEA-S, hypertension and hyperlipidemia ( $\beta = 0.390$ ,  $-0.224$  and  $-0.295$ , respectively) were selected as significant variables. Next, in forward and backward stepwise models including age, body mass index, systolic blood pressure, total cholesterol, HDL cholesterol, fasting plasma glucose, smoking and the four steroid hormones, only DHEA-S and glucose were selected as significant variables ( $\beta = 0.330$  and  $-0.325$ , respectively). Taken together, these results show that DHEA-S was consistently related to %FMD in the multivariate analyses.

### Discussion

In this cross-sectional study, plasma DHEA-S levels were positively correlated with %FMD, a surrogate marker of clinical atherosclerosis that reflects endothelial function (18, 19). Adjustment for potential confounders such as age, coronary risk factors and nitroglycerin-induced dilation of the brachial artery had little influence on the results. These results suggest that endogenous DHEA(-S) plays a vasoprotective role in postmenopausal women.

The majority of DHEA exists as the sulfated form (DHEA-S) in circulation, and its concentration is reported to be 100-




Fig. 1. Percent flow-mediated dilation of the brachial artery (%FMD) according to quartiles of serum dehydroepiandrosterone-sulfate (DHEA-S). Values are expressed as the means  $\pm$  SEM. <sup>\*</sup> $p < 0.05$ , <sup>\*\*</sup> $p < 0.01$  vs. Q1.

to 500-fold higher than that of testosterone and 1,000- to 10,000-fold higher than that of estradiol (4), suggesting an important physiologic role of DHEA(-S). Plasma DHEA-S levels do not drop during the menopause transition, but decline with aging (1, 2, 20). A recent study has shown a positive correlation between DHEA-S levels and the duration of reactive hyperemia, another less specific marker of endothelial function compared to FMD, in young women ( $26 \pm 6$  years old,  $n = 27$ ) with polycystic ovary syndrome (21). In postmenopausal women, however, a correlation between DHEA-S levels and endothelial vasomotor function has not yet been reported. On the other hand, two studies have reported the effects of DHEA supplementation on FMD in postmenopausal women (12, 22). Williams *et al.* (12) showed a significant increase in FMD after the 12-week administration of DHEA (50 mg/day) in healthy postmenopausal women, while Silvestri *et al.* (22) observed no effect using the same dose of DHEA for 4 weeks in postmenopausal women with increased cardiovascular risk. In middle-aged men, it has been reported that low-dose DHEA supplementation (25 mg/day) for 12 weeks improved FMD (23). Experimental studies have also demonstrated the effects of DHEA on endothelial proliferation (12) and endothelial nitric oxide synthase (26), further supporting the notion that DHEA(-S) has a protective effect on the endothelium.

At this moment, intrinsic receptors for DHEA(-S) have not been identified. Thus, DHEA(-S) may exert its activity after conversion to androgens or estrogens via androgen or estrogen receptors, although in previous reports neither an androgen receptor antagonist nor an estrogen receptor antagonist blocked some of the *in vitro* effects of DHEA on endothelial cells (12, 23). Testosterone supplementation in men (23, 24) and estrogen replacement in postmenopausal women (27, 28) improves endothelial vasomotor function. Activation of endothelial nitric oxide synthase and stimulation of nitric oxide production via androgen and estrogen receptors (29) might be attributable to the effect of DHEA(-S) on FMD. For

this reason, we added testosterone and estradiol into the analysis. Plasma testosterone and estradiol levels, however, were correlated neither with %FMD nor with DHEA-S (data not shown), although tissue conversion of DHEA(-S) into androgens or estrogens might play a role. Further studies are needed to clarify the molecular mechanism underlying the association between DHEA-S and FMD.

The incidence of cardiovascular disease is lower in premenopausal women than in men of the same age, and increases after menopause (30), indicating the cardioprotective effect of endogenous estrogen in premenopausal women. Accordingly, estrogen replacement therapy had been expected to prevent cardiovascular disease in postmenopausal women, with disappointing results in randomized controlled trials (31). Alternatively, there has been a resurgence of interest in DHEA as an anti-aging hormone, and DHEA has been widely used in this context in the USA, despite the dearth of information on its physiologic and pharmacologic effects (4). Although the present study indicated a cardioprotective effect of endogenous DHEA(-S), we do not suggest that DHEA be prescribed for postmenopausal women with coronary risk factors until the efficacy and safety of DHEA supplementation has been established. At present, life-style modification such as exercise is preferable, because exercise can increase both DHEA levels (32) and nitric oxide production (33) in elderly women.

The results of this study do not imply that DHEA(-S) has favorable effects on endothelial function in men. In fact, the plasma DHEA-S level was not related to FMD in middle-aged men after adjustments for age and coronary risk factors (our unpublished observation). Consequently, the meaning of DHEA-S in association with endothelial function may be different between men and women. Gender differences in other steroid hormones and steroid hormone receptor expression in arteries (34, 35) might play a mechanistic role.

This study had some limitations. First, since this was a cross-sectional study, the causal relationship between DHEA-S and vasomotor function could not be determined. Endothelial dysfunction might be associated with a reduction in blood flow of the adrenal glands, leading to decreased hormone production. Longitudinal studies following the subjects might add some information in this regard. Secondly, a population bias was possible. Three-quarters of the study subjects had one or more of the coronary risk factors, and most of them were taking medications. Consequently, the results might have been different if subjects homogeneous in terms of health status and medications had been studied, although the association of DHEA-S with %FMD was consistent in the subjects with or without coronary risk factors and medications, as mentioned above. The subjects with a history of cardiovascular disease were excluded from the study, because they might have low plasma DHEA-S levels as a result of advanced atherosclerosis and reduced blood flow of the adrenal glands, although the inclusion of those subjects did not fundamentally alter the statistical results (data not shown).

In summary, low plasma DHEA-S levels were associated with endothelial dysfunction in postmenopausal women independent of other risk factors, suggesting a protective effect of DHEA(-S) on the endothelium. This finding provides a mechanistic insight into the role of endogenous DHEA in the development of cardiovascular disease in postmenopausal women.

### Acknowledgements

We thank Ms. Yuki Ito for her excellent technical assistance.

### References

1. Bjornerem A, Straume B, Midtby M, et al: Endogenous sex hormones in relation to age, sex, lifestyle factors, and chronic diseases in a general population: the Tromso Study. *J Clin Endocrinol Metab* 2004; **89**: 6039-6047.
2. Tannenbaum C, Barrett-Connor E, Laughlin GA, Platt RW: A longitudinal study of dehydroepiandrosterone sulphate (DHEAS) change in older men and women: the Rancho Bernardo Study. *Eur J Endocrinol* 2004; **151**: 717-725.
3. Lamberts SW, van den Beld AW, van der Lely AJ: The endocrinology of aging. *Science* 1997; **278**: 419-424.
4. Hinson JP, Raven FW: DHEA deficiency syndrome: a new term for old age? *J Endocrinol* 1999; **163**: 1-5.
5. Wu FC, von Eckardstein A: Androgens and coronary artery disease. *Endocr Rev* 2003; **24**: 183-217.
6. Liu PY, Death AK, Handelsman DJ: Androgens and cardiovascular disease. *Endocr Rev* 2003; **24**: 313-340.
7. Tchermof A, Labrie F: Dehydroepiandrosterone, obesity and cardiovascular disease risk: a review of human studies. *Eur J Endocrinol* 2004; **151**: 1-14.
8. Haffner SM, Most SE, Klein BE, Klein R: Sex hormones and DHEA-SO4 in relation to ischemic heart disease mortality in diabetic subjects. The Wisconsin Epidemiologic Study of Diabetic Retinopathy. *Diabetes Care* 1996; **19**: 1045-1050.
9. Bernini GP, Sgro' M, Moretti A, et al: Endogenous androgens and carotid intimal-medial thickness in women. *J Clin Endocrinol Metab* 1999; **84**: 2008-2012.
10. Santoro N, Torrens J, Crawford S, et al: Correlates of circulating androgens in mid-life women: the study of women's health across the nation. *J Clin Endocrinol Metab* 2005; **90**: 4836-4845.
11. Barna I, Feher T, de Chatel R: Relationship between blood pressure variability and serum dehydroepiandrosterone sulfate levels. *Am J Hypertens* 1998; **11**: 532-538.
12. Williams MR, Dawood T, Ling S, et al: Dehydroepiandrosterone increases endothelial cell proliferation *in vitro* and improves endothelial function *in vivo* by mechanisms independent of androgen and estrogen receptors. *J Clin Endocrinol Metab* 2004; **89**: 4708-4715.
13. Hutchison SJ, Browne AE, Ko E, et al: Dehydroepiandrosterone sulfate induces acute vasodilation of porcine coronary arteries *in vitro* and *in vivo*. *J Cardiovasc Pharmacol* 2005; **46**: 325-332.
14. Japanese Society of Hypertension: Japanese Society of Hypertension Guidelines for the Management of Hyperten-