

TABLE 2. SERUM 25(OH)D LEVEL AND CHARACTERISTICS FOR MALES AND FEMALES

Characteristics	Male			Female		
	Lower ( $\leq 25.0$ ng/ml) (n = 249)	Higher ( $\geq 26.0$ ng/ml) (n = 701)	p	Lower ( $\leq 21.0$ ng/ml) (n = 576)	Higher ( $\geq 22.0$ ng/ml) (n = 1431)	p
Fall experience over the previous year (yes, n, %)	27 (10.8)	76 (10.8)	0.938*	129 (22.4)	243 (17.0)	0.020*
Average number of falls (times, mean $\pm$ SD)	2.1 $\pm$ 2.4	1.7 $\pm$ 1.1	0.422 <sup>†</sup>	1.6 $\pm$ 1.2	1.4 $\pm$ 0.8	0.021 <sup>b)</sup>
Hand grip strength (kg, mean $\pm$ SD)	30.5 $\pm$ 6.7	31.7 $\pm$ 6.5	0.020 <sup>b)</sup>	17.9 $\pm$ 4.6	19.2 $\pm$ 4.6	0.002 <sup>b)</sup>
Stork standing time with eye open (s, mean $\pm$ SD)	34.6 $\pm$ 22.5	38.2 $\pm$ 22.5	0.046 <sup>b)</sup>	31.7 $\pm$ 23.5	37.7 $\pm$ 23.0	<0.001 <sup>†</sup>
Normal walking speed (m/s, mean $\pm$ SD)	1.19 $\pm$ 0.26	1.25 $\pm$ 0.26	0.061 <sup>†</sup>	1.12 $\pm$ 0.28	1.21 $\pm$ 0.27	<0.001 <sup>†</sup>
Serum albumin (g/dl, mean $\pm$ SD)	4.34 $\pm$ 0.24	4.35 $\pm$ 0.26	0.616 <sup>†</sup>	4.28 $\pm$ 0.23	4.33 $\pm$ 0.21	<0.001 <sup>†</sup>
	Insufficiency ( $< 20.0$ ng/ml) (n = 46)	Normal ( $\geq 20.0$ ng/ml) (n = 904)	p	Insufficiency ( $< 20.0$ ng/ml) (n = 356)	Normal ( $\geq 20.0$ ng/ml) (n = 1651)	p
Fall experience over the previous year (yes, n, %)	3 (6.5)	100 (11.1)	0.454*	92 (25.8)	280 (17.0)	0.001*
Average number of falls (times, mean $\pm$ SD)	2.7 $\pm$ 0.6	1.8 $\pm$ 1.5	0.338 <sup>†</sup>	1.7 $\pm$ 1.3	1.4 $\pm$ 1.5	0.006 <sup>†</sup>
Hand grip strength (kg, mean $\pm$ SD)	28.5 $\pm$ 6.4	31.5 $\pm$ 6.5	0.003 <sup>†</sup>	18.1 $\pm$ 4.7	19.0 $\pm$ 4.6	0.420 <sup>b)</sup>
Stork standing time with eye open (s, mean $\pm$ SD)	31.4 $\pm$ 22.9	37.5 $\pm$ 22.5	0.124 <sup>†</sup>	29.8 $\pm$ 22.9	37.2 $\pm$ 23.2	<0.001 <sup>†</sup>
Normal walking speed (m/s, mean $\pm$ SD)	1.16 $\pm$ 0.79	1.24 $\pm$ 0.26	0.138 <sup>†</sup>	1.11 $\pm$ 0.29	1.20 $\pm$ 0.27	<0.001 <sup>†</sup>
Serum albumin (g/dl, mean $\pm$ SD)	4.27 $\pm$ 0.26	4.35 $\pm$ 0.22	0.027 <sup>†</sup>	4.27 $\pm$ 0.23	4.32 $\pm$ 0.21	<0.002 <sup>†</sup>

\* The Mantel-Haenszel  $\chi^2$  test adjusted for age.<sup>†</sup> ANCOVA adjusted for age.

TABLE 3. ASSOCIATION OF SERUM ALBUMIN AND 25(OH)D LEVELS WITH PHYSICAL PERFORMANCE FOR MALES AND FEMALES

	Handgrip strength			Stork standing time			Normal walking speed		
	$\beta$	SE	p	$\beta$	SE	p	$\beta$	SE	p
Men									
Albumin	0.096	0.852	0.001	0.002	3.644	0.947	0.025	0.053	0.576
25(OH)D	0.067	0.037	0.020	0.075	0.152	0.030	0.111	0.002	0.012
Women									
Albumin	0.109	0.459	<0.001	0.037	2.502	0.106	0.045	0.030	0.051
25(OH)D	0.062	0.020	0.003	0.109	0.105	<0.001	0.143	0.001	<0.001

Values are adjusted for age. p values are derived from multiple regression analysis.

 $\beta$ , standardized regression coefficient.

carried out to confirm the correlation among the three physical performance tests. The results showed high and significant intercorrelation for these three variables; correlation coefficients were from 0.23 (handgrip strength and stork standing time) to 0.35 (stork standing time and normal walking speed) in men and from 0.31 (handgrip strength and stork standing time) to 0.47 (stork standing time and normal walking speed) in women, and all were significant at  $p < 0.001$ . Therefore, we adopted only "normal walking speed" to represent the physical performance tests as well as the independent variable for the final multiple logistic regression model.

Table 4 shows the associations of fall experience over the previous year with normal walking speed, serum albumin, and 25(OH)D levels by multiple logistic regression models with age adjustment. Normal walking speed (unit = 0.1 m/s) showed a significant protective effect against falls in

TABLE 4. MULTIPLE LOGISTIC REGRESSION MODEL OF FACTORS ASSOCIATED WITH FALL EXPERIENCE OVER THE PREVIOUS YEAR

Risk factor	Male			Female		
	OR	95% CI	p	OR	95% CI	p
Age (yr)	1.02	0.95-1.10	NS	1.02	0.99-1.06	NS
Normal walking speed (0.1 m/s)	0.87	0.77-0.97	0.015	0.92	0.88-0.97	0.001
Albumin (g/dl)	1.69	0.45-6.33	NS	1.60	0.88-2.90	NS
25(OH)D (ng/ml)	1.00	0.95-1.06	NS	0.97	0.94-0.99	0.010

Dependent variable was "fall experience over the previous year" (yes = 1, no = 0).

The unit of normal walking speed was transferred from meters per second to 0.1 m/s in this final multiple logistic regression model.

NS, not significant.

both men (OR = 0.87, 95% CI = 0.77-0.97) and women (OR = 0.92, 95% CI = 0.88-0.97). Serum 25(OH)D level (unit = 1 ng/ml) also had a significant and independent protective effect for falls found only in women (OR = 0.97, 95% CI = 0.94-0.99,  $p = 0.01$ ).

### DISCUSSION

Maintenance of physical performance in old age is an important factor not only for a healthy and independent life in the community but also a way to prevent falls that can lead to a marked decline in activities of daily living (ADLs). A national survey in Japan has shown that the annual frequency of falls is >20% in those >65 yr of age and that ~10% of these falls result in fractures.<sup>(27)</sup>

This study showed that the proportion of people who reported falls in the previous year increased with age and that falls were more common in women than in men.<sup>(20)</sup> These findings are consistent with the results of other studies of falls among community-dwelling elderly.<sup>(28-32)</sup> Aoyagi et al.<sup>(33)</sup> reported that the proportion of falls in the previous year after age standardization for Japanese was about one half of that of whites. Furthermore, the incidence of hip fracture among Japanese elderly was found to be much lower than that reported for whites in North America and Europe.<sup>(34)</sup> This difference is probably partly the result of the lower fall rate among Japanese, suggesting that both ethnicity (genetics) and lifestyle (environmental) factors may be involved.<sup>(14)</sup> Recently, some studies have shown that lower serum vitamin D level is a risk factor for falls and fall-associated physical performance among the elderly.<sup>(9,10,35)</sup> At present, however, there are few studies on the association between serum 25(OH)D level and falls in Japanese community elderly, whose frequency of falls is less than that observed in Europe and the United States.<sup>(6-8,13)</sup>

In this study, we found that there were significant sex differences of 25(OH)D level on average and in a pattern of decline along with aging; namely, women had significant lower serum 25(OH)D levels at any age group and showed remarkable declines with aging. One of the reasons for this sex difference may be general inactivity and lower intake of vitamin D from daily food among Japanese elderly women compared with men. One Japanese national survey showed that, compared with 39.1% of men, 32.6% of women engage in physical activity for at least 30 min two or more times a week.<sup>(36)</sup> Our previous study also reported that women had significantly lower rates of regular sports activity than men (13.8% versus 21.5%).<sup>(37)</sup> Furthermore, we recently reported that one of the significant predictors for cessation of regular activity was "female sex" as well as "smoking" and "slow walking speed" from a population-based, 2-yr follow-up study.<sup>(38)</sup> The national survey also showed that women took smaller amounts of vitamin D than men ( $8.6 \pm 9.0$  versus  $8.9 \pm 10.0$   $\mu\text{g/d}$  on average). In particular, for elderly respondents  $\geq 70$  yr of age, the average intake of vitamin D in women ( $8.6 \pm 9.0$   $\mu\text{g/d}$ ) was much less than that in men ( $10.3 \pm 10.1$   $\mu\text{g/d}$ ).<sup>(36)</sup> These factors of physical inactivity and lower intake of daily vitamin D in elderly women may have caused their observed higher frequency of 25(OH)D insufficiency compared with men. It is

known that the main source of vitamin D in humans is considered to be through the skin, where vitamin D is produced during exposure to UVB sunlight.<sup>(39,40)</sup> In our study, to avoid any seasonal variation of serum 25(OH)D, data collection was carried out only during autumn (October/November), which meant that serum 25(OH)D levels would be almost stable and at an average throughout the year among a normal Japanese population.<sup>(41)</sup>

The range of serum 25(OH)D levels was 8-42 ng/ml in men and 9-38 ng/ml in women. Although it is still uncertain what an optimal 25(OH)D level is, it has been suggested that the range of 25(OH)D levels should be 32-100 ng/ml, with a lower limit somewhere between 15 and 36 ng/ml.<sup>(41-43)</sup> In this study, we defined a 25(OH)D level of <20 ng/ml as insufficiency. From this, we judged that the prevalence of 25(OH)D insufficiency was significantly predominant in women (17.7%) compared with men (4.8%;  $p < 0.001$ ). Studies concerning the prevalence of 25(OH)D insufficiency in various populations have been challenged because of the lack of standardization of assays and different cut-off points.<sup>(44)</sup> A comparison of serum 25(OH)D level between hip fracture patients and nonhip fracture controls in Japan showed that average serum 25(OH)D concentrations were significantly different: 17.8 ng/ml in hip fracture patients and 25.8 ng/ml in nonhip fracture controls.<sup>(45)</sup> Furthermore, 62% of the hip fracture patients ( $N = 50$ ) had 25(OH)D insufficiency, defined as having a serum 25(OH)D concentration <20 ng/ml. In this context, the elderly whose 25(OH)D levels were <20 ng/ml can be considered to be in insufficiency and at high risk of a hip fracture, which indeed has increased sharply during last two decades in Japan.<sup>(46)</sup>

In comparisons of serum 25(OH)D levels and fall-associated variables between the group of participants in the lowest quartile and the groups of the three other higher quartiles of serum 25(OH)D level, all variables but normal walking speed in men were significantly lower in the lower 25(OH)D group than in the higher group. Our findings are consistent with the results from a Swedish population-based study of 986 community-living elderly women<sup>(47)</sup> that showed that the lower 25(OH)D group was significantly correlated with inferior gait speed, inferior balance test, and lower knee extension/flexion strength results, all of which are fall-associated variables.

The rate of fall experience over the previous year by serum 25(OH)D level was significantly different only in women [i.e., 22.4% in the lower 25(OH)D group and 17.0% in the higher group ( $p = 0.02$ ) and also 25.8% in the insufficiency group and 17.0% in the normal group ( $p = 0.001$ ), respectively]. The average numbers of falls were also significantly different between the lower and higher groups ( $p = 0.021$ ) and between the insufficient and normal groups ( $p = 0.006$ ), respectively. This finding that lower serum 25(OH)D level or 25(OH)D insufficiency status is associated with falls among elderly women is consistent with many previous studies.<sup>(3,4,11)</sup> However, there is a controversy about which type of vitamin D [i.e., 25(OH)D or 1,25(OH)<sub>2</sub>D<sub>3</sub>] is associated with fall risk. Faulkner et al.,<sup>(48)</sup> who examined the relationship of vitamin D supplementation and the serum concentration of vitamin D metabolites

with falls in older white, community-dwelling women ( $n = 389$ ) in the United States, reported that only the higher serum  $1,25(\text{OH})_2\text{D}_3$  concentration was associated with a lower fall risk but that  $25(\text{OH})\text{D}$  concentration was not associated with falls. In our study, as one of the study limitations, serum  $1,25(\text{OH})_2\text{D}_3$  was not assessed in the participants undergoing the mass health examination.

Further analysis on the association of serum albumin and  $25(\text{OH})\text{D}$  levels with the fall-associated variables in this study showed that only serum  $25(\text{OH})\text{D}$  level had a significant association with all three fall-associated variables in both sexes. On the other hand, serum albumin had a significant association only with handgrip strength in both sexes. The mechanism connecting serum albumin and muscle mass or power is not clear.<sup>(49,50)</sup> However, serum albumin concentration may be a marker of the protein status of an individual, with lower values indicating a diminished protein reserve and stimulated catabolic processes leading to muscle break down and also muscle strength decline.<sup>(50)</sup> Thus, low serum albumin, even within a normal range, is independently associated with weaker muscle strength and future decline in older men and women.<sup>(51,52)</sup>

Serum  $25(\text{OH})\text{D}$  concentration is an important determinant of muscle mass and sarcopenia. In an observational study of community-dwelling elderly in the Netherlands, incident sarcopenia, defined as a minimum of 40% decline in muscle strength and 3% decline in muscle mass, was found to be twice as likely among  $25(\text{OH})\text{D}$ -deficient elderly [ $25(\text{OH})\text{D}$  level < 10 ng/ml] than among elderly with a  $25(\text{OH})\text{D}$  level of >20 ng/ml.<sup>(7)</sup> With respect to the role of vitamin D in muscle strength, the majority of the actions of vitamin D are mediated through  $1,25(\text{OH})_2\text{D}_3$  binding to nuclear vitamin D receptor (VDR) that can directly modulate the transcription of the gene possessing a functional binding site for VDR in its regulatory region.<sup>(8,53)</sup> Therefore, muscle strength seems to be influenced by the VDR genotype in the muscle cell. With the use of specific restriction endonucleases, several VDR polymorphisms have been determined. In nonobese, older women, a 23% difference in quadriceps strength and a 7% difference in grip strength between two homozygote types of a restriction site have been found.<sup>(54)</sup> The action of vitamin D is affected by allelic variance of the VDR. A genomic study on VDR polymorphism has shown that Japanese women had much lower frequency of homozygote BB (1.4%) than white women (16.7%).<sup>(55)</sup> In this context, if the VDR polymorphism affects not only the BMD but also muscle strength of elderly women, Japanese women in general may have an advantage with respect to their lower frequency of falls and associated hip fractures.

As for the association of serum  $25(\text{OH})\text{D}$  level and fall experience over the previous year by multiple logistic regression models, even after the adjustment of other fall-associated variables, serum  $25(\text{OH})\text{D}$  level was independently associated with falls in women as well as normal walking speed in this study. A considerable number of population-based studies have also been conducted on walking ability in relation to the occurrence of falls in the elderly.<sup>(56-60)</sup> Increased body sway, uneven distances, and uneven timing during walking were identified as risk factors

for falls.<sup>(61,62)</sup> The authors have previously reported that both fall experience over the previous year and a decline in walking speed were very strong predictors of the occurrence of frequent falls in a 5-yr follow-up cohort study among Japanese community-living elderly.<sup>(18)</sup> Muscle function or strength as the single most important component for walking ability has been consistently identified as a risk factor for hip fractures as a consequence of falls in the elderly. Subclinical  $25(\text{OH})\text{D}$  insufficiency is also considered to be an important risk factor for hip fractures in elderly people in both white<sup>(63,64)</sup> and Japanese<sup>(45)</sup> populations. We have suggested that elderly women with lower  $25(\text{OH})\text{D}$  levels and with a significant decline in their fall-associated variables tend to decline in walking capability and be more vulnerable to falls even in elderly populations of Japanese women, whose fall rate has been reported to be low.

A relevant issue regarding the role of  $25(\text{OH})\text{D}$  in physical performance is that vitamin D supplementation has been reported to be significantly effective in maintaining or improving physical performance and preventing falls among the elderly. In a recent randomized and multiple-dose study, Broe et al.<sup>(65)</sup> reported that a high dose of vitamin D (800 IU/d) reduced the risk of falls dramatically by 72% lower (adjusted-incidence rate ratio) than participants taking a placebo over the same 5-mo period in a nursing home. However, the question of which type of vitamin D supplementation (i.e., either cholecalciferol or calciferol) is more effective at reducing falls among community-living elderly is still controversial. The association of vitamin D supplementation with better physical performance related to the risk of falls and/or reduced falls is unclear at present. For example, a meta-analysis of five randomized controlled trials provided some evidence that vitamin D supplementation might reduce falls,<sup>(10)</sup> whereas the results of a second meta-analysis of four randomized controlled trials of vitamin D found no such evidence.<sup>(66)</sup>

Our findings suggesting that there are significant relationships between serum  $25(\text{OH})\text{D}$  and fall-associated physical performance and with falls themselves could provide guidance about how to prevent falls and fractures, particularly hip fractures resulting from falls among community-living elderly. Two such countermeasures are to improve muscle strength, especially in the lower extremities, and to enhance balance ability.<sup>(67,68)</sup> For example, from a randomized controlled exercise intervention trial for Japanese community-dwelling elderly, we have proven that a moderate exercise intervention program in addition to a home-based program significantly improved fall-associated variables and, consequently, decreased the incidence of falls for 1.5 yr after the intervention.<sup>(67)</sup>

Another countermeasure would be to maintain a high level of serum  $25(\text{OH})\text{D}$  by adequate intake of foods containing vitamin D, supplementation, and exposure to sunlight. Incidentally, fish consumption seems to play an important role in maintaining adequate vitamin D nutrition among elderly Japanese.<sup>(69)</sup> In general, these countermeasures seem to be consistent with the traditional Japanese lifestyle. Our earlier study using risk factor analysis of hip fractures in elderly Japanese showed that such a traditional

Japanese lifestyle, including living on Japanese tatami mats and eating fish daily, was strongly associated with a decrease in the risk of hip fractures.<sup>(14)</sup> Living on Japanese tatami mats, including futon-style bedding, seems have great benefits for preventing falls and hip fractures by continuously strengthening the muscles of the hip girdle and lower extremities by sitting, squatting, and frequent standing over the course of many years. Furthermore, consumption of dark-meat fish, which is rich in vitamin D, also seems to be beneficial for maintaining adequate 25(OH)D levels in the elderly, especially in the winter season.<sup>(69)</sup>

Before detailing our final conclusions, some limitations of our study must be considered. (1) The subjects analyzed were not selected randomly from the study population; as well, they were relatively healthy elderly persons who were able to travel from their homes to the health checkup venue. As a result, elderly persons with lower physical functional capacity were excluded. (2) Plasma 1,25(OH)<sub>2</sub>D<sub>3</sub>, albumin-corrected calcium, and PTH, which would provide information on the extent of any primary vitamin D deficiency,<sup>(8,48,70)</sup> and creatine clearance that may affect the metabolism of vitamin D through the kidney,<sup>(71)</sup> were not assessed in this study. It is well known that increased secretion of PTH is associated with decreased serum 25(OH)D levels, which may commonly occur in the elderly. However, according to a survey on the nutritional status of vitamin D among Japanese community-dwelling elderly, Nakamura et al.<sup>(72)</sup> reported that only 1.8% of the subjects had elevated intact PTH levels. (3) We did not analyze the genotype of the VDR that could influence muscle strength and, likely, the fall rate as well. (4) This study was cross-sectional and therefore did not provide cause/effect relationships, although we showed a significant correlation between physical performance and serum 25(OH)D levels in Japanese community-dwelling elderly. Therefore, a longitudinal follow-up study and controlled clinical trials would seem necessary to confirm the role of serum 25(OH)D in falls and its association with the physical performance of the elderly.

In conclusion, our findings showed that a lower serum 25(OH)D level was significantly associated with fall experience over the previous year and with fall-associated variables in Japanese women whose fall rate has been reported to be about one half that of white women. This indicates that serum 25(OH)D level has a common and positive relationship with the occurrence of falls in elderly women, and probably beyond any genetic background represented by VDR phenotype differences and anthropometric and nutritional differences.

Muscle weakness or sarcopenia, frailty, and falls, all of which can be frequent among the elderly and therefore are often categorized as geriatric syndrome, have a major impact on the elderly in terms of both morbidity and mortality. The connections between these items related to geriatric syndrome and 25(OH)D has been well established by many population-based epidemiological studies including this one. Such geriatric syndromes could be prevented by both exercise interventions and adequate levels of serum 25(OH)D; these would help maintain good physical performance and functional capacity for a high quality of life among community-dwelling elderly.

## ACKNOWLEDGMENTS

This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education and Culture of Japan (16390187) and a Research Grant from the Research Society for Metabolic Bone Diseases in Japan.

## REFERENCES

- Riggs BL 2003 Role of the vitamin-D-endocrine system in the pathophysiology of postmenopausal osteoporosis. *J Cell Biochem* 88:209-215.
- Lips P 2001 Vitamin D deficiency and secondary hyperparathyroidism in the elderly: Consequences for bone loss and fractures and therapeutic implications. *Endocr Rev* 22:477-501.
- Bischoff HA, Stahelin HB, Urscheler N, Ehrensam R, Vonthein R, Perrig-Chiello P, Tyndall A, Theiler R 1999 Muscle strength in the elderly: Its relation to vitamin D metabolites. *Arch Phys Med Rehabil* 80:54-58.
- Mowe M, Haug E, Bohmer T 1999 Low serum, calcidiol concentration in older adults with reduced muscular function. *J Am Geriatr Soc* 47:220-226.
- Verhaar HJ, Samson M, Lansens PA, de Vreede PL, Macnen JW, Duursma SA 2001 Muscle strength, functional mobility and vitamin D in older women. *Aging Clin Exp Res* 12:455-460.
- Janssen H, Samson MM, Verhaar HJ 2002 Vitamin D deficiency, muscle function, and falls in elderly people. *Am J Clin Nutr* 75:611-615.
- Visser M, Deeg D, Lips P 2003 Low vitamin D and high parathyroid hormone levels as determinants of loss of muscle strength and muscle mass (sarcopenia): The Longitudinal Ageing Study Amsterdam. *J Clin Endocrinol Metab* 88:5766-5772.
- Campbell PMF, Allain TJ 2006 Muscle strength and vitamin D in older people. *Gerontology* 52:335-338.
- Dhesi JK, Bearne LM, Moniz C, Hurley MV, Jackson SH, Swift CG, Allain TJ 2002 Neuromuscular and psychomotor function in elderly subjects who fall and the relationship with vitamin D status. *J Bone Miner Res* 17:891-897.
- Bischoff HA, Stahelin HB, Dick W, Akos R, Knecht M, Salis C, Nebiker M, Theiler R, Pfeifer M, Begerow B, Lew RA, Conzelmann M 2003 Effects of vitamin D and calcium supplementation on falls: A randomized controlled trial. *J Bone Miner Res* 18:343-351.
- Stein MS, Wark JD, Scherer SC, Walton SL, Chick P, Di Carantonio M, Zajac JD, Flicker L 1999 Falls relate to vitamin D and parathyroid hormone in an Australian nursing home and hostel. *J Am Geriatr Soc* 47:1195-1201.
- Kwon J, Suzuki T, Yoshida H, Kim H, Yoshida Y, Iwasa H 2007 Concomitant lower serum albumin and vitamin D levels are associated with decreased objective physical performance among Japanese community-dwelling elderly. *Gerontology* 53:322-328.
- Lipsitz LA, Nakajima I, Gagnon M, Hirayama T, Connelly CM, Izumo H, Hirayama T 1994 Muscle strength and fall rates among residents of Japanese and American nursing homes: An international cross-cultural study. *J Am Geriatr Soc* 42:953-959.
- Suzuki T, Yoshida H, Hashimoto T, Yoshimura N, Fujiwara S, Fukunaga M, Nakamura T, Yoh K, Inoue T, Hosoi T, Orimo H 1997 Case-control study of risk factor for hip fracture in Japanese elderly by Mediterranean Osteoporosis Study (MEDOS) questionnaire. *Bone* 21:461-467.
- Suzuki T, Iwasa H, Yoshida H, Kim H, Watanabe S, Yukawa H 2003 Comprehensive health examination ('Otashakenshin') for the prevention of geriatric syndromes and a bedridden state in the community elderly. 1. Differences in characteristics between participants and non-participants (in Japanese with English summary). *Jpn J Public Health* 50:39-48.


16. Iwasa H, Suzuki T, Yoshida H, Kim H, Yoshida Y 2003 Cognitive function as the factor determining higher level competence in community-dwelling elderly (In Japanese with English summary). *Jpn J Public Health* 50:950-958.
17. Kwon J, Suzuki T, Kim H, Yoshida H, Yoshida Y, Furuna T 2005 Health status and physical fitness undernourished community-dwelling elderly people. (In Japanese with English summary). *Jpn J Phys Fitness Sports Med* 54:99-106.
18. Suzuki T, Sugiura M, Furuna T, Nishizawa S, Yoshida H, Ishizaki T, Kim H, Yukawa H 1999 Association of physical performance and falls among the community elderly in Japan in a 5 year follow-up study (in Japanese with English summary). *Jpn J Geriatr* 36:472-478.
19. Obuchi S, Shibata H, Suzuki T, Yasumura S 1994 Relationship between walking ability and risk of falls in community-dwelling elderly in Japan. *J Phys Ther Sci* 6:39-44.
20. Yasumura S, Haga H, Suzuki T, Shibata H 1994 Rate of falls: The correlates among elderly people living in an urban community in Japan. *Age Ageing* 23:323-327.
21. Nagasaki H, Itoh H, Furuna T 1995 The structure underlying physical performance measures for older adults in the community. *Aging Clin Exp Res* 7:451-458.
22. Thomas MK, Demay MB 2000 Vitamin D deficiency and disorders of vitamin D metabolism. *Endocrinol Metab Clin North Am* 29:611-627.
23. Lips P, Chapuy MC, Dawson-Hughes B, Pols HA, Holick MF 1999 An international comparison of serum 25-hydroxyvitamin D measurements. *Osteoporos Int* 9:394-397.
24. Nakamura K, Nashimoto M, Yamamoto M 2000 Summer/winter differences in the serum 25-hydroxyvitamin D3 and parathyroid hormone levels of Japanese women. *Int J Biometeorol* 44:186-189.
25. Malabanan A, Veronikis IE, Holick MF 1998 Redefining vitamin D insufficiency. *Lancet* 351:805-806.
26. Need AG, Horowitz M, Morris HA, Nordin BC 2000 Vitamin D status: Effects on parathyroid hormone and 1,25-dihydroxyvitamin D in postmenopausal women. *Am J Clin Nutr* 71:1577-1581.
27. Shibata H (ed.) 1997 An Overall Survey on Falls and Fractures of Community Elderly. Report on the Studies by National Science Research Grants. No. 07307007. Ministry of Education, Culture, Sports, Science and Technology, Tokyo, Japan.
28. Blake AJ, Morgan K, Bendall MJ, Dallosso H, Ebrahim SB, Aric TH, Fentem PH, Bassey EJ 1988 Falls by elderly people at home: Prevalence and associated factors. *Age Ageing* 17:365-372.
29. Wickham C, Cooper C, Margetts BM, Barker DJ 1989 Muscle strength, activity, housing and the risk of falls in elderly people. *Age Ageing* 18:47-51.
30. Prudham D, Evans JG 1981 Factors associated with falls in the elderly: A community study. *Age Ageing* 10:141-146.
31. Winner SJ, Morgan CA, Evans JG 1989 Perimenopausal risk of falling and incidence of distal forearm fracture. *BMJ* 298:1486-1488.
32. O'Loughlin JL, Robitaille Y, Boivin JF, Suissa S 1993 Incidence of and risk factors for falls and injurious falls among the community-dwelling elderly. *Am J Epidemiol* 137:342-354.
33. Aoyagi K, Ross P, Davis JW, Wasnich RD, Hayashi T, Takemoto T 1998 Falls among community-dwelling elderly in Japan. *J Bone Miner Res* 13:1468-1474.
34. Schwartz AV, Kelsey JL, Maggi S, Tuttleman M, Ho SC, Jonsson PV, Poor G, Sisson JA, Xu L, Matkin CC, Nelson LM, Heyse SP 1999 International variation in the incidence of hip fractures: Cross-national project on osteoporosis for the WHO program for research on aging. *Osteoporos Int* 9:242-253.
35. Pfeifer M, Begerow B, Minne HW, Abrams C, Nachtigall D, Hansen C 2000 Effects of a short-term vitamin D and calcium supplementation on body sway and secondary hyperparathyroidism in elderly women. *J Bone Miner Res* 15:1113-1118.
36. Ministry of Health Labour and Welfare 2006 The National Health and Nutrition Survey, Japan, 2005. Tokyo, Ministry Health, Labour and Welfare, Tokyo, Japan
37. Kim H, Suzuki T, Iwasa H, Yoshida H 2003 A prospective study of the effects of regular sports practice on mortality among the elderly in a rural community in Japan: An 8-year follow-up study. *Geriatr Gerontol Int* 3:S15-S20.
38. Shimada H, Lord S, Yoshida H, Kim H, Suzuki T 2007 Predictors of cessation of regular leisure-time physical activity in community-dwelling elderly people. *Gerontology* 53:293-297.
39. Vieth R 2000 Problems with direct 25-hydroxyvitamin D assays. And the target amount of vitamin D nutrition desirable for patients with osteoporosis. *Osteoporos Int* 11:635-636.
40. Holick MF 2004 Sunlight and vitamin D for bone health status and ramifications of vitamin deficiency. *Am J Clin Nutr* 80(Suppl):1706S-1709S.
41. Ono Y, Suzuki A, Kotake M, Zang X, Nishiwaki K, Ishikawa Y, Imamura S, Nagata M, Takamoto S, Itoh M 2005 Seasonal changes of serum 25-hydroxyvitamin D and intact parathyroid hormone level in a normal Japanese population. *J Bone Miner Metab* 23:147-151.
42. Hollis BW 2005 Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: Implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr* 135:317-322.
43. Vieth R 1999 Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr* 69:842-856.
44. Binkly N, Krueger D, Cowgill CS 2004 Assay variation confounds the diagnosis of hypovitamin D: A call for standardization. *J Clin Endocrinol Metab* 89:3152-3157.
45. Sakuma M, Endo N, Oinuma T, Hayami T, Endo E, Yazawa T, Watanabe K, Watanabe S 2006 Vitamin D and intact PTH status in patients with hip fracture. *Osteoporos Int* 17:1608-1614.
46. Orimo H, Hashimoto T, Sakata K, Yoshimura N, Hosoi T, Suzuki T 2000 Trends in the incidence of hip fracture in Japan, 1987-1997: The third nationwide survey. *J Bone Miner Metab* 18:126-131.
47. Gerdhem P, Ringsberg KAM, Obrant KJ, Akesson K 2005 Association between 25-hydroxyvitamin D level, physical activity, muscle strength and fractures in the prospective population-based OPRA study of elderly women. *Osteoporos Int* 16:1425-1431.
48. Faulkner KA, Cauley JA, Zmuda JM, Landsittel DP, Newman AB, Studenski SA, Redfern MS, Ensrud KE, Fink HA, Lane NE, Nevitt MC 2006 Higher 1,25-dihydroxyvitamin D<sub>3</sub> concentrations associated with lower fall rates in older community-dwelling women. *Osteoporos Int* 17:1318-1328.
49. Baumgartner RN, Koehler KM, Romero L, Garry PJ 1996 Serum albumin is associated with skeletal muscle in elderly men and women. *Am J Clin Nutr* 64:552-558.
50. Visser M, Kritchevsky S, Newman A, Goodpaster B, Tyllavsky F, Nevitt M, Harris T 2005 Lower serum albumin concentration and change in muscle mass: The health, aging and body composition study. *Am J Clin Nutr* 82:531-537.
51. Kwon J, Suzuki T, Kim H, Yoshida H, Kumagai S, Yoshida Y, Furuna T, Sugiura M 2005 Health status and physical fitness of undernourished community-dwelling elderly people (in Japanese with English summary). *Jpn J Phys Fitness Sports Med* 54:99-106.
52. Schalk BW, Deeg DJ, Penninx BW, Bouter L, Visser M 2005 Serum albumin and muscle strength: A longitudinal study in older men and women. *J Am Geriatr Soc* 53:1331-1338.
53. Bischoff HA, Borchers M, Gudat F, Deurmueller U, Theiler R, Staehelin HB, Dick W 2001 In situ detection of the 1,25-dihydroxyvitamin D<sub>3</sub> receptor in human skeletal muscle tissue. *Histochem J* 33:19-24.
54. Janssen H, Samson M, Verhaar H 2002 Vitamin D deficiency, muscle function, and falls in elderly people. *Am J Clin Nutr* 75:611-615.
55. Tokita A, Matsumoto H, Morrison NA, Tawa T, Miura Y, Irimoto M, Yamamori S, Watanabe T, Kuwabara Y, Yabuta K, Eisman JA 1997 Vitamin D receptor alleles, bone mineral density and turnover in premenopausal Japanese women. *J Bone Miner Res* 12:494-495.
56. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace

- RB 1995 Lower extremity function in persons over the age of 70 year as a predictor of subsequent disability. *N Engl J Med* 332:556-561.
57. Suzuki T, Yoshida H, Kim H, Yukawa H, Furuta T, Shinkai S, Ishizaki T, Watanabe S, Shibata H 2003 Walking speed as a good predictor for maintenance of I-ADL among the rural community elderly in Japan: A 5-year follow-up study from TMIG-LISA. *Geriatr Gerontol Int* 3:S6-S14.
58. Buchner DM, Cress ME, Esselman PC, Margherita AT, de Lateur BJ, Cambell AJ, Wagner EH 1996 Factors associated with changes in gait speed in older adults. *J Gerontol A Biol Sci Med Sci* 51:M297-M302.
59. Gibbs J, Hughes S, Dunlop D, Singer R, Chang RW 1996 Predictors of change in walking velocity in older adults. *J Am Geriatr Soc* 44:126-132.
60. Kwon J, Suzuki T, Yoshida H, Kim H, Yoshida Y, Iwasa H, Furuta T 2007 Association between change in bone mineral density and decline in usual walking speed in elderly community dwelling Japanese women during 2 year of follow-up. *J Am Geriatr Soc* 55:240-244.
61. Tinetti ME, Speechly M, Ginter SF 1988 Risk factors for falls among elderly persons living in the community. *N Engl J Med* 319:1701-1707.
62. Topper AK, Maki BE, Hokkiday PJ 1993 Are activity-based assessments of balance and gait in the elderly predictive of risk of falling and/or type of fall? *J Am Geriatr Soc* 41:479-487.
63. LeBoff MS, Kohlmeier L, Hurwitz S, Franklin J, Wright J, Glowacki J 1999 Occult vitamin D deficiency in postmenopausal US women with acute hip fracture. *JAMA* 281:1505-1511.
64. Nuti R, Martini G, Valenti R, Gambera D, Gennari L, Salvadori S, Avanzati A 2004 Vitamin D status and bone turnover in women with acute hip fracture. *Clin Orthop* 422:208-213.
65. Broe KE, Chen TC, Weinberg J, Bischoff-Ferrari HA, Hollick MF, Kiel DP 2007 A higher dose of vitamin D reduces the risk of falls in nursing home residents: A randomized, multiple-dose study. *J Am Geriatr Soc* 55:234-239.
66. Latham NK, Anderson CS, Reid IR 2003 Effects of Vitamin D supplementation on strength, physical performance, and falls in older persons: A systematic review. *J Am Geriatr Soc* 51:1219-1226.
67. Suzuki T, Kim H, Yoshida H, Ishizaki T 2004 Randomized controlled trial of exercise intervention for the prevention of falls in community-dwelling elderly Japanese women. *J Bone Miner Metab* 22:602-611.
68. Rubenstein LZ, Josephson KR, Trueblood PR, Loys S, Harker JO, Pietruszka FM, Robbins AS 2000 Effects of group exercise program on strength, morbidity, and falls among fall-prone elderly men. *J Gerontol A Biol Sci Med Sci* 55:M317-M321.
69. Nakamura K 2006 Vitamin D insufficiency in Japanese populations: From the viewpoint of the prevention of osteoporosis. *J Bone Miner Metab* 24:1-6.
70. Nakamura K, Nishiwaki T, Ueno K, Yamamoto M 2005 Serum 25-hydroxyvitamin D and parathyroid hormone levels of non-institutionalized elderly Japanese requiring care. *J Bone Miner Metab* 23:488-494.
71. Dukas LC, Schacht E, Mazor Z, Stahelin HB 2005 A new significant and independent risk factor for falls in elderly men and women: A low creatinine clearance of less than 65 ml/min. *Osteoporos Int* 16:332-338.
72. Nakamura K, Nashimoto M, Hori Y, Yamamoto M 2000 Serum parathyroid hormone in healthy female Japanese in relation to serum 25-hydroxy vitamin D. *Int J Vitamin Nutr Res* 70:287-292.

Address reprint requests to:  
Takao Suzuki, MD, PhD  
Tokyo Metropolitan Institute of Gerontology  
35-2 Sakaecho, Itabashi-ku  
Tokyo 173-0015, Japan  
E-mail: suzuta@tmig.or.jp

Received in original form August 15, 2007; revised form March 14, 2008; accepted March 21, 2008.

# 生活習慣病 ガイドブック



編集 兵庫県医師会生活習慣病対策プロジェクト会議

# VIII 脳卒中

## —脳血管障害の診断と危険因子について—

櫻井 孝

### 1 脳卒中の概説

1. 致死的な脳血管障害は減少しているが、後遺症を残す患者が増加している。機能予後の改善のため、超急性期治療の可能性を考える。
2. 超急性期に診断に必要なすべての所見をとることは不可能である。治療方針を決定するための必要最小限の診断を行い、一刻も早く必要な画像検査を行う。
3. 急性発症する神経精神症候には脳卒中以外の疾患の可能性も考える。
4. 脳卒中の局在診断のみならず、閉塞血管、発症機序を明らかにし、病型分類に応じた治療、再発予防を行う。

### 2 脳卒中超急性期の診断と全身管理

1. 問診のポイント：「今回、何が起きたか」を明確にする。家族の立ち会いを得て、発症前のADLと比較して今回の発症後にできなくなったことを明らかにする。「急性発症か、緩徐進行性か、階段状悪化、何をしていたときか?」、発症時刻の特定、随伴症状（頭痛、頸部痛、発熱）など。高血圧、糖尿病、高脂血症、心疾患、心房細動、内服薬などの既往。
2. 診察のポイント：一般身体所見、意識・精神状態（Japan Coma Scale）、注意・見当識、精神状態、認知機能。神経学的診察では、脳神経系、運動系（Barre 徴候、Mingazzini 徴候）、感覚系（顔面・体幹・四肢の最低各一点で確認）、反射（深部腱反射、病的反射）、協調運動（指鼻・膝踵試験、つぎ足歩行）。
3. 脳梗塞全般としては、発症直後には絶対安静が必要（15度程度の頭部挙上は可能）。
4. 脳卒中発症直後の高血圧治療は、高血圧性脳症、クモ膜下出血が強く疑われる場合以外は病型診断が確定してから行う。
5. 脳梗塞超急性期の組織プラスミノゲンアクチベーター（t-PA）の静注療法：治療開始時刻は発症後3時間以内。アテローム血栓性脳梗塞、ラクナ梗塞、心原性脳塞栓症はいずれも対象となる。

## Japan Coma Scale (JCS)

### III. 刺激しても覚醒しない状態(3桁の点数で表現)

(deep coma, coma, semicoma)

- 300. 痛み刺激に全く反応しない
- 200. 痛み刺激で少し手足を動かしたり顔をしかめる
- 100. 痛み刺激に対し、払いのけるような動作をする

### II. 刺激すると覚醒する状態(2桁の点数で表現)

(stupor, lethargy, hypersomnia, somnolence, drowsiness)

- 30. 痛み刺激を加えつつ呼びかけを繰り返すと辛うじて開眼する
- 20. 大きな声または体を揺さぶることにより開眼する
- 10. 普通の呼びかけで容易に開眼する

### I. 刺激しなくても覚醒している状態(1桁の点数で表現)

(delirium, confusion, senselessness)

- 3. 自分の名前、生年月日が言えない
- 2. 見当識障害がある
- 1. 意識清明とは言えない

注 R : Restlessness(不穏)、I : Incontinence(失禁)、A : Apallic state または Akinetic mutism  
たとえば30Rまたは30不穏とか、20Iまたは20失禁として表す。

## 3 脳血管障害(脳卒中)の分類

National Institute of Neurological Disorders and Stroke の分類による(注1)

- A. 無症候性脳血管障害
- B. 局所障害型の脳血管障害
  - 1. 一過性脳虚血発作(注2)
  - 2. 脳卒中
    - a. くも膜下出血
    - b. 脳出血
    - c. 脳梗塞
      - (1) アテローム血栓性
      - (2) 心原性血栓性
      - (3) ラクナ梗塞
      - (4) その他
- C. 脳血管性痴呆
- D. 高血圧性脳症

注1 : 脳の血管の破綻により生じる疾患を脳血管障害という。脳血管障害の中で突然発症をするものだけを脳卒中として区別する。

注2 : 脳虚血によると考えられる局所脳神経症状が一過性に出現し、24時間以内に完全に消失するもの。

## 4 脳卒中慢性期：危険因子の管理と予防

### 1. 脳梗塞

- 1) 高血圧は脳梗塞の最大の危険因子であり、降圧療法が推奨される（グレードA）。脳卒中の予防には血圧は低ければ低いほどよい（The lower, the better）。降圧目標は140/90mmHg未満（高齢者）、130/85mmHg未満（若年・中年）。
- 2) 糖尿病は脳梗塞発症のリスク（2～3倍）。血糖の管理だけでは脳卒中を予防しにくく、合併した危険因子すべてに対処する（血圧は130/80mmHg未満）。
- 3) 脳卒中予防効果が証明された脂質異常治療薬はスタチンのみ。
- 4) 喫煙は脳梗塞発症のリスク（2～4倍）。禁煙により脳卒中発症リスクは禁煙後2年以内に急速に減少し、5年以内に非喫煙者と同じレベルになる。
- 5) 適量を超える飲酒は脳梗塞のリスク。日本酒1日3合（エタノール60～70g）以上の飲酒で、全脳卒中の発症増加（1.9倍）。少量の飲酒は脳梗塞のリスクを下げるという。
- 6) 弁膜症を伴わない心房細動（NVAf）を伴う脳梗塞の予防にワルファリンが有効（international normalized ratio（INR）2.0～3.0でコントロール）。しかし日本人、特に70歳以上では、やや低用量のワルファリン（INR 1.6～2.6）が勧められる。INR >2.6にて出血性合併症は急増する。
- 7) 無症候性脳梗塞は、脳梗塞の危険因子。無症候性脳梗塞から発症する脳卒中の約2割が脳出血であり、抗血小板薬の投与は高血圧の管理を厳重に行った上で行う。
- 8) 非心原性脳梗塞（アテローム血栓性脳梗塞、ラクナ梗塞）の再発予防上、最も有効な抗血小板療法はアスピリン（75～150mg/日）。ラクナ梗塞の再発予防に対してエビデンスを有する抗血小板薬は現時点ではシロスタゾールのみ。抗血小板薬（アスピリン、チクロピジン、クロピドグレル）による頭蓋内出血の危険性は0.2～0.3%/年、ワルファリンでは0.3～1.1%/年。
- 9) 脳循環代謝改善薬：イブジラスト、酒石酸イフェンプロジルは脳梗塞後のめまいに、ニセルゴリンは脳梗塞後の認知障害に有効。
- 10) 脳卒中の18～62%にうつ状態を合併する。うつ状態は、認知機能低下、ADL回復の遅延、死亡（3倍）のリスクとなる。選択的セロトニン再取り込み阻害薬（SSRI）を含む抗うつ薬の投与が推奨される。
- 11) 中等度以上の症候性頸動脈狭窄病変（狭窄率50%以上、特に70%以上）では、内科的治療（抗血小板療法を含む）＋頸動脈内膜剥離術（CEA）が有効。

### 2. 脳出血

- 1) 脳出血の発症の危険因子：年齢、性別、高血圧、過量飲酒、血清総コレステロール低値。拡張期血圧 >95mmHgでは、40～59歳で9.0倍、60歳以上で3.4倍増加。多量飲酒は脳出血のリスク（1日60g以上で2.18倍）。
- 2) 定期的な運動は、脳出血の予防に有効である可能性がある。

- 3) 1日3種類の果物や野菜を取ることに脳出血の危険性が51%減少するという。
- 4) 脳出血後の脳卒中再発では、約半数が脳出血、残り半数が脳梗塞である。
- 5) 高血圧性脳出血では血圧のコントロール不良例で再発が多い。脳出血の再発予知因子としては、脳葉出血が重要であり、ApoEの遺伝子多型が関連する。
- 6) 維持透析患者では、脳出血を発症する危険性が高い(5~10倍)。

### 3. クモ膜下出血

- 1) 女性に多い(男女比1:2)。
- 2) 予後不良例が約40%存在する。発症時の意識障害の程度が予後と相関する。
- 3) クモ膜下出血の危険因子: 脳動脈瘤や脳動静脈奇形の存在の他、喫煙習慣、高血圧保有、一週間に150g以上の飲酒。
- 4) 一般医療機関から脳神経外科専門施設への搬送は、速やかに行われるべきである。移送中の再出血の予防のため、血圧管理、鎮痛鎮静、医師の同乗が望ましい。

### 参考資料

- 1) 日本脳卒中学会: 脳卒中治療ガイドライン2004 (<http://www.jsts.gr.jp/jss08.html>)
- 2) 日本高血圧学会: 高血圧治療ガイドライン2004

# 生活習慣病 ガイドブック



編集 兵庫県医師会生活習慣病対策プロジェクト会議

# IX 認知症

## —認知症の診断と予防—

櫻井 孝

### 1 認知症の概説

1. 認知症とは、いったん獲得した知的機能が持続的に低下し、複数の認知障害があるために社会生活に支障をきたすようになった状態である。
2. 認知症の頻度は65歳以上の高齢者で約6～7%であり、加齢とともに増加する。
3. 近年、認知症に対する社会の見方も一変しており、全診療科の積極的な関わりが求められる。

### 2 認知症を来す疾患

1. アルツハイマー病と血管性認知症が最多（両疾患を合わせると80%程度）。
2. 脳腫瘍・慢性硬膜下血腫・正常圧水頭症などの脳外科的治療の対象となる疾患、甲状腺機能低下症などの内分泌疾患、重症の肝臓病や腎臓病、ビタミン欠乏症などの「治療により脳機能が改善する認知症」を見逃さない（表1）。

表1 日常診療でよく見られる認知症の原因疾患

1. 脳血管性の疾患  
脳血管障害（脳出血、脳梗塞など）による認知症
2. 退行変性疾患  
アルツハイマー病、パーキンソン病、びまん性レビー小体病など
3. 内分泌・代謝性中毒性疾患  
甲状腺機能低下症、下垂体機能低下症、ビタミンB<sub>12</sub>欠乏症、ビタミンB<sub>1</sub>欠乏症、ペラグラ、ミトコンドリア脳筋症、肝性脳症、透析脳症、低酸素症、低血糖症、アルコール脳症、薬物中毒など
4. 感染性疾患  
脳炎・髄膜炎、脳膿瘍、進行麻痺、クロイツフェルト・ヤコブ病など
5. 腫瘍性疾患  
脳腫瘍、髄膜浸潤など
6. 外傷性疾患  
慢性硬膜下血腫、頭部外傷後遺症など
7. その他  
正常圧水頭症、神経ペーチェット、サルコイドーシス、シェーグレン症候群など

3. 脳血管障害、脳症や薬剤の副作用（表2）によるせん妄状態など治療可能な状態も少なくない。
4. うつ病や他の統合失調症は認知症と混乱されることがある（仮性認知症）。

**表2 高齢者のせん妄の原因**

1. 薬剤性：抗パーキンソン病薬、抗コリン薬、抗潰瘍薬（H<sub>2</sub>ブロッカー）、抗ヒスタミン薬、解熱鎮痛消炎剤、抗精神薬、抗不安・鎮静・睡眠薬、抗うつ薬、抗ウイルス薬（アシクロビル）、抗腫瘍薬（植物アルカロイド、L-asparaginase）、循環器用薬：（抗不整脈薬、β遮断薬、ジギタリス、硝酸薬）、気管支拡張剤（キサンチン）、咳止め（リン酸コデイン）、副腎皮質ステロイド、インターフェロン、甲状腺薬、抗結核薬（エタンブトール、イソニアジド、サイクロセリン）、アルコールなど
2. 感染症：肺炎、尿路感染症、敗血症など
3. 代謝異常：脱水、電解質異常、低血糖、尿毒症、肝性脳症、甲状腺機能亢進症、クッシング症候群、ビタミン欠乏症
4. 循環器疾患：心不全、心筋梗塞、不整脈
5. 中枢神経疾患：脳血管障害、認知症、慢性硬膜下血腫、脳髄膜炎、脳腫瘍
6. 呼吸不全、CO<sub>2</sub>ナルコーシス
7. SLEなどの自己免疫疾患
8. 悪性腫瘍
9. 手術・全身麻酔

### 3 認知症の診断

#### 日常診療上のヒント

認知症の診断において、詳細な問診を行うことが診断に到達する早道である。またすべての一般内科、神経内科的診察を行うことは困難であり、主要症候をまずスクリーニングする。

1. 問診のポイント：1）もともとの認知能（職歴、学歴、生活歴を参考）、2）発症時期、3）発症後の経過（進行の速さ、階段状悪化）、4）生活上の支障（ADLの低下）、5）記憶、見当識、注意、言語、遂行機能障害、失認、失行を示すエピソード、6）神経脱落症状、7）他の身体症候、8）体重変化など。他の身体・神経疾患、頭部外傷、生活習慣病などの既往、内服薬、飲酒、喫煙などの情報を得る。
2. 診察のポイント：脳神経（眼球運動、眼振、対光反射、構音・嚥下障害）、パーキンソン症候（安静時振戦、筋トーンス）、筋力（Barre test）、深部腱反射、病的反射、歩行、協調運動、構成機能（狐、リング、鳩の手まね）など。
3. 高次脳機能のスクリーニングには、改訂長谷川式簡易知能評価スケール（HDS-R）（表3）、ミニメンタルテスト（MMSE）が有用である。
4. 採血検査で治療可能な認知症をチェックする（WBC、RBC、Hb、Hct、Plt、

APTT、PT、Fib、総蛋白、アルブミン、AST、ALT、LDH、 $\gamma$ -GTP、ALP、BUN、Cr、CK、Na、K、Ca、P、T-Chol、TG、HDL-C、血糖、葉酸、ビタミンB<sub>1</sub>、ビタミンB<sub>12</sub>、fT<sub>4</sub>、TSH、アンモニア、RPR、尿検査一般)。

5. 脳画像検査(頭部CT/MRI)で、慢性硬膜下血腫、脳腫瘍などの疾患を除外する。血管病変の程度を評価する。
6. 高齢者のアルツハイマー病では、遅延再生(短期記憶)、時間見当識の障害が早期から現れ、認知障害は緩徐に進行する。
7. 脳血管性認知症の診断には、①認知症の確認、②血管病変に伴う神経所見と脳画像で示される脳血管障害の存在、③脳血管障害との時間的關係(3ヵ月以内の認知症発症、または認知機能の急速な低下、動揺性・段階的な悪化)を示すことが要件となる。

### 日常診療上のヒント

皮質下白質病変(ラクナー梗塞を含む)は認知障害の原因となる。しかし脳画像検査でみられる白質病変の程度のみから、血管性認知症の診断はできない。

表3 改訂長谷川式簡易知能評価スケール

氏名	年齢 (男・女)	年 月 日生	質問内容	配点
1			お年はいくつですか？(2年までの誤差は正解)	0, 1
2			今日は何年の何日ですか？何曜日ですか？ (年、月、日、曜日が正解でそれぞれ1点ずつ)	0, 1 0, 1 0, 1 0, 1
3			私たちがいまいる所はどこですか？ (自発的にできれば2点、5秒おいて家ですか？病院ですか？施設ですか？のなかから正しい選択をすれば1点)	0, 1, 2
4			これから言う3つの言葉を言ってみて下さい。あとでまた聞きますのでよく覚えておいて下さい。 (以下の系列のいずれか1つで、採用した系列に○印をつけておく) 1: a) 桜 b) 猫 c) 電車 2: a) 梅 b) 犬 c) 自動車	0, 1 0, 1 0, 1
5			100から7を順番に引いて下さい。 (100-7は？、それからまた7をひくと？と質問する。最初の答えが不正解の場合、打ち切る。それぞれ1点。)	0, 1 0, 1
6			私がこれから言う数字を逆から言って下さい。 (6-8-2, 3-5-2-9を逆に言ってもらう。3桁逆唱に失敗したら、打ち切る)	0, 1 0, 1
7			先ほど覚えてもらった言葉をもう一度言ってみて下さい。(自発的に回答があれば各2点、もし回答が無い場合以下のヒントを与え正解であれば1点) a) 植物 b) 動物 c) 乗り物	a: 0, 1, 2 b: 0, 1, 2 c: 0, 1, 2
8			これから5つの品物を見せます。それを隠しますのでなにがあったか言って下さい。(時計、鍵、タバコ、硬貨など必ず相互に無関係なもの)	0, 1, 2 3, 4, 5
9			知っている野菜の名前をできるだけ多く言って下さい。(答えた野菜の名前を右欄に記入する。途中で詰まったり、約10秒間待っても答えがない場合はそこで打ち切る) 0~5=0点、6=1点、7=2点、8=3点、9=4点、10=5点	0, 1, 2 3, 4, 5
満点30点 20点以下 認知症の疑い				合計点数


## 4 認知症の治療

1. 高齢者および認知症への処方若年者よりも少量から始める。また服薬法をできるだけ簡単にする。
2. アルツハイマー病では、ドネペジルが第1選択薬。著効は数10%であるが、多くの症例で会話力や協調性、社会的活動の改善が期待できる。
3. 焦燥、妄想、攻撃性に対しては抗精神病薬（ハロペリドール、オランザピン、リスペリドンなど）が用いられるが、生命予後を低下させるとの報告があり、慎重な投与が必要。
4. 非薬物療法（回想法、音楽・絵画など）は、患者のストレス減少、残在能力の活用などに有用とされる。またデイ・ケアなどでのグループ活動への参加はADL低下の抑制に効果がある。介護保険を積極的に利用し、精神的・身体的な機能維持を計る。
5. 成年後見制度は、医療や財政に関する意志決定の代行制度など、本人の権利を守る意義がある。本人や家族へ助言することが必要。

## 5 認知症の予防

1. 脳血管障害は血管性認知症のリスクである。
2. アルツハイマー病の危険因子として、介入できない危険因子（年齢、遺伝子異常：アポE $\epsilon$ 4保持）の他に、環境因子が重要である。即ち、動脈硬化、脳卒中、糖尿病、高血圧・低血圧、喫煙、飲酒、高コレステロール、心臓病、高脂肪食、血

図1 アルツハイマー病と血管性認知症の境界



- 栓症、高ホモシスチン、高フィブリノーゲン、頭部外傷、抑うつ、閉経、低教育レベルなどで、これらは脳血管障害のリスクとほぼ共通する。アルツハイマー病による神経変性と血管障害は、相加的・相乗的に脳の障害を増悪させる（図1）。
3. 壮年期よりの生活習慣病をきめ細かく管理することで、高齢者になった際の認知症の発症を抑制し得る可能性がある。

# 認知症治療の最前線

ニコチン治療を中心として

監修：広島大学名誉教授／洛和会京都治験・臨床研究支援センター所長 中村 重信

メヂカルビュー社

## 合併症のある認知症患者への対応

### 1) 糖尿病

神戸大学大学院医学系研究科内科学講座老年内科学分野/教授\*  
芳野 弘・櫻井 孝・横野 浩一\*

#### はじめに

認知症では、さまざまな療養上の問題が生じる。認知症の前期～中期では記憶、見当識障害などに伴う薬剤管理の困難、判断能力の低下が問題となり、後期となると広範な認知機能の低下に伴い日常生活動作が制限され、家族の介護負担は多大なものとなる。一方、糖尿病は全身に多彩な合併症をきたす疾患であり、認知症性疾患に糖尿病が合併すると、実に全人的な問題点を考慮せねばならない。

認知症と糖尿病の合併を考えると、多くが70歳以降に発症する認知症性疾患から遅れて発症する高齢者糖尿病と、成年期に発症して経過の長い糖尿病が高齢化して、その後認知症性疾患が合併したものに分ける必要がある。高齢発症の糖尿病では、血管合併症の進行が緩徐であることが多く、その療養では過度の高血糖、低血糖、あるいは感染症に対する対策が中心となる。しかし後者においては、すでに細小血管合併症や動脈硬化性病変が存在し、日常生活動作(ADL)が低下していることが多く、認知症の合併により糖尿病自体の療養が困難となることがある。そこで本稿では糖尿病の経過中に認知症性疾患を合併した高齢者の療養を中心に、その問題点と対策についてまとめたい。

#### 糖尿病の認知障害と認知症

高齢糖尿病患者では、非糖尿病高齢者に比べて認知機能が低下していることが知られている。注意力、前頭葉機能、記憶の低下などさまざまな認知機能の低下が生じるが、その程度は比較的軽度であり、多くは糖尿病の療養の支障となるものではない。しかし経口血糖降下薬の内服管理を忘れる、インスリン自己注射に失敗が多いなどのエピソードの原因となることもある。

糖尿病と認知症の合併については、これまでに多くの疫学研究が報告されている。いずれの研究でも血管性認知症の合併は有意に高値である。最近の前向きコホート研究の結果では、アルツハイマー病(AD)の合併も高いことが報告されている。1999年のロッテルダム研究では、特にインスリン使用者で認知症の発症が高いことが報告され、多くの関心を集めている<sup>1)</sup>。すなわち、糖尿病は認知症の危険因子であるとする立場から、糖尿病の認知障害から認知症に至るプロセスについて、多くの研究が世界中で進行している。

#### 糖尿病に伴う認知障害、認知症の発症機序と予防

高齢者糖尿病患者の脳には図1に示したさまざまな要因が脳機能を修飾しているものと考えられる<sup>2)3)</sup>。①低血糖、②高血糖に伴う代謝障害、③脳血管障害、④インスリン、IGF-1などの成長因子の作用、⑤βアミロイドカスケードなどが想定される。低血糖は重要で、大脳皮質、大脳基底核、海馬などの神経細胞は低酸素、低グルコースに対して脆弱である。成人では低血糖に陥ると、動悸、冷汗などの交感神経症状により低血糖を自覚するが、低血糖を繰り返す患者や高齢者では、交感神経症状が欠如し、いきなり意識障害、認知障害として気づかれることがある(無自覚低血糖)。認知症を合併した高齢者糖尿病患者では低血糖の訴えも曖昧であることが多く、低血糖が遅延すると脳機能は急速に低下することがある。

高血糖が持続すると脳内でもポリオール代謝経路、酸化ストレス、グリケーションなどの糖代謝異常が進行する。これらの代謝異常はADの発症機序の1つとしても指摘されており、高血糖の持続は脳機能を低下させることが想定される。また高齢者糖尿病では脳の大血管、小血管に多

様な病変をきたし、神経組織は低酸素に陥りやすい。糖尿病では脳出血の頻度はむしろ低下しているが、脳血栓症、ラクナ梗塞は増加する。その予防には血圧管理が重要であるが、脳血管の狭窄性病変を十分に把握したうえでの適正な血圧コントロールが求められる。特に後期高齢者では拡張期血圧の低下は認知障害のリスクであるとの報告もあり、過度の降圧療法には注意を要する。

近年、ADと血管性認知症の危険因子がほぼ共通していることが明らかとなった。ADの原因となるβアミロイドカスケードは低酸素、高血糖、高コレステロール、また高血圧と関連する。すなわち、糖尿病のみならず、合併する代謝異常、血圧、食生活の改善は、認知症の発症を抑制することが期待される。現在多くの医師が糖尿病性腎症の進行を抑制するためにACE阻害薬を処方するのと同じく、高齢者糖尿病の脳機能を守るためにいかなる治療を行うべきか、一定の指針が提言されることが期待されている。


## 糖尿病性認知症患者への対応

高齢者糖尿病で認知症を合併した患者の療養では、個々の疾患の治療を行うのみではなく、身体的、精神的、社会的な問題が重要である。これらの問題点をもれなく評価するための手段として、高齢者総合機能評価は有用である。高齢者総合機能評価を行い、問題点を整理してその対策を包括的に考える。患者の療養におけるキーパーソンを特定し、主治医を中心とし、精神科医、看護師、薬剤師、栄養士、理学療法士などのコメディカルからなるチーム医療を実践することが重要である。糖尿病性認知症患者の医療の質を向上させることは、患者のQOLを改善し、家族の介護負担を軽減させるのみならず、医療経済の効率改善にも強く寄与する。

### 文献

- 1) Ott A, Stolk RP, van Harskamp F, et al : Diabetes mellitus and the risk of dementia ; the Rotterdam Study. *Neurology* 53 : 1937-1942, 1999
- 2) 櫻井 孝 : 高齢者糖尿病と認知機能障害. *Diabetes Frontiers* 13 (3) : 337-340, 2002
- 3) 横野浩一, 櫻井 孝 : 高齢者糖尿病の治療と研究. *日老医誌* 41 : 369-371, 2004

図1 高齢者糖尿病とアルツハイマー病



# 老年医学 update 2008-09

編集 日本老年医学会雑誌編集委員会

特集

高齢者総合的機能評価の臨床応用  
高齢者糖尿病の新たな展開