

Fig. 5. Effects of PD98059 and SB203580 on either the ADP-induced PDGF-AB or 5-HT secretion in human platelets. Various dose of (A and C) PD98059, a specific inhibitor of MEK1/2 or (B and D) SB203580, a specific inhibitor of p38 MAPK, were added to PRP and incubated at 37 °C for 5 min without stirring. PRP was followed in an aggregometer at 37 °C for 5 min with a stirring speed of 800 rpm; PRP was preincubated for 1 min with stirring, then platelet aggregation was monitored for 4 min after the addition of 3 μM of ADP (●) or vehicle (○). Aggregation was terminated by the addition of an ice-cold EDTA (10 mM) solution. The mixture was centrifuged at 10,000g at 4 °C for 2 min and the supernatants were then subjected to ELISA for (A and B) PDGF-AB and (C and D) 5-HT. Representative results from at least three independent experiments are shown. Bars indicate standard deviations of triplicate assays. The asterisk (*) indicates a significant decrease ($p < 0.05$) with respect to the value of ADP (3 μM) alone.

granule populations has not yet been elucidated. These previous observations led to the speculation that there are differential mechanisms of granule secretion among these three granule populations. In the present study, both the serum levels of PDGF-AB and 5-HT, secreted from α -granules and dense granules respectively, decreased in parallel with the decreased levels of HSP27 phosphorylation. These results strongly suggest that HSP27 phosphorylation may therefore play an important role in the granule secretion of both dense granules and α -granules. Dense granule contains ADP and release one which is recognized to be released through a positive feedback mechanism. However, since we can not strictly distinguish exogenous ADP from endogenous one, we do not exclude the possibility that the latter one might affect platelet aggregation. In conclusion, these results indicate that the ADP-induced phosphorylation of HSP27 via p38 MAPK and/or p44/p42 MAPK is sufficient for platelet granule secretion but not for the platelet aggregation in humans. In consideration with the fact that the specific antagonist for 5-HT has already been clinically applied for amelioration of ischemic symptoms of arterial thrombotic dis-

eases [33], we might be able to establish the new clinical target of arterial thrombotic diseases by the investigation of the relationship between HSP27 and platelet granule secretion.

References

- [1] F. Rendu, B. Brohard-Bohn, *Platelets* 12 (2001) 261–273.
- [2] B. Hechler, C. Leon, C. Vial, P. Vigne, C. Frelin, J.P. Cazenave, C. Gachet, *Blood* 92 (1998) 152–159.
- [3] J.L. Daniel, C. Dangelmaier, J. Jin, Y.B. Kim, S.P. Kunapuli, *Thromb. Haemost.* 82 (1999) 1322–1326.
- [4] A.J. Schafer, *Am. J. Med.* 101 (1996) 199–209.
- [5] E.J. Topol, *N. Engl. J. Med.* 347 (2002) 1359–1360.
- [6] R.W. Colman, *FASEB J.* 4 (1990) 1425–1435.
- [7] M.Z. Kowalska, M.Z. Ratajczak, M. Majka, J. Jin, S.P. Kunapuli, L. Brass, M. Poncz, *Blood* 96 (2000) 50–57.
- [8] C. Dangelmaier, J. Jin, J.L. Daniel, J.R. Smith, S.P. Kunapuli, *Eur. J. Biochem.* 267 (2000) 2283–2289.
- [9] K. Falke, D. Lange, P. Presek, *Thromb. Haemost.* 92 (2004) 114–123.
- [10] J.P. Hendrick, F.U. Hartl, *Annu. Rev. Biochem.* 62 (1993) 349–384.
- [11] I.J. Benjamin, D.R. McMillan, *Circ. Res.* 83 (1998) 117–132.
- [12] Y. Inaguma, S. Goto, H. Shinohara, K. Hasegawa, K. Ohshima, K. Kato, *J. Biochem. (Tokyo)* 114 (1993) 378–384.

- [14] M. Gaestel, W. Schroder, R. Benndorf, C. Lippmann, R. Buchner, F. Huch, V.A. Fischmann, H. Bielek, *J. Biol. Chem.* 266 (1991) 14721–14724.
- [15] J. Landry, H. Lambert, M. Zhou, J.N. Lavie, F. Hokey, L.A. Weber, C.W. Anderson, *J. Biol. Chem.* 267 (1992) 794–801.
- [16] W.J. Welch, *J. Biol. Chem.* 260 (1985) 3058–3062.
- [16] K. Kato, K. Hasegawa, S. Goto, Y. Inaguma, *J. Biol. Chem.* 269 (1994) 11274–11278.
- [17] T. Kato, E.A. M. Ehrnsperger, X. Peville, A. Kotlyarov, G. Tutich, C. Dacasse, C. Paul, M. Waske, A.P. Arrigo, J. Buchner, M. Gaestel, *J. Biol. Chem.* 274 (1999) 18947–18950.
- [18] Y. Zhu, S. O'Neill, J. Nakayama, I. Lassi, M.E. Mendicino, *Blood* 84 (1994) 1115–1121.
- [19] J.M. Kyriakis, *J. Virol.* 71 (1995) 24313–24316.
- [20] J. Gaay, H. Lambert, G. Gungas-Breton, J.N. Lavie, J. Hokey, J. Landry, *J. Cell Sci.* 110 (1997) 157–168.
- [21] K. Kato, H. Ito, K. Hasegawa, Y. Inaguma, O. Kozawa, T. Asano, *J. Neurochem.* 66 (1996) 946–950.
- [22] U.S. Lammli, *Naturwissenschaften* 227 (1970) 680–685.
- [23] D.R. Alessi, A. Cuenda, P. Cohen, D.L. Dolley, A.R. Sartori, *J. Biol. Chem.* 270 (1995) 27484–27491.
- [24] A. Cuenda, J. Rouse, Y.S. Doza, R. Meier, P. Cohen, T.F. Gallagher, P.R. Young, J.C. Lee, *FEBS Lett.* 364 (1995) 229–233.
- [25] S.P. Kenapala, *Scientific World J.* 2 (2002) 424–431.
- [26] A. McNicol, E.C.G. Jackson, *Biochem. Pharmacol.* 65 (2003) 1243–1250.
- [27] B.G. Lovell, H. Biesemann, K.B. Pollitt, *J. Biol. Chem.* 263 (1988) 90–94.
- [28] J. Miron, K. Vancompernelle, J. Vandekerckhove, M. Wolchek, B. Geiger, *J. Cell Biol.* 114 (1991) 275–291.
- [29] J. Miron, M. Wolchek, B. Geiger, *Int. J. Biochem.* 178 (1988) 543–551.
- [30] J.N. Lavie, E. Hokey, L.A. Weber, J. Landry, *J. Biol. Chem.* 268 (1993) 24210–24214.
- [31] F. Renfu, P. Marche, J. Maclouf, A. Gaard, S. Levy-Toledano, *Biochem. Biophys. Res. Commun.* 116 (1983) 513–519.
- [32] F. Renfu, P. Marche, J. Viret, J. Maclouf, M. Lebrat, D. Tenza, J. Caen, S. Levy-Toledano, *Biochimie* 69 (1987) 305–313.
- [33] K. Satoh, Y. Yatomi, Y. Ozaki, J. Hirumi, *Haemostasis* 4 (2006) 479–481.



Involvement of Rho-kinase in prostaglandin $F_{2\alpha}$ -stimulated interleukin-6 synthesis via p38 mitogen-activated protein kinase in osteoblasts

Chiho Minamitani^{a,b}, Takano Otsuka^a, Shinji Takai^b, Rie Matsushima-Nishiwaki^b, Seiji Adachi^b, Yoshiteru Hanai^{b,c}, Jun Mizutani^a, Haruhiko Tokuda^{b,c}, Osamu Kozawa^{b,*}

^a Department of Orthopedic Surgery, Nagoya City University Graduate School of Medical Sciences, Nagoya 467-8601, Japan

^b Department of Pharmacology, Gifu University Graduate School of Medicine, 1-1 Yanagido, Gifu 501-1194, Japan

^c Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu, Aichi 474-8511, Japan

ARTICLE INFO

Article history:

Received 10 September 2007

Received in revised form 24 April 2008

Accepted 19 May 2008

Keywords:

PGF_{2α}
Rho-kinase
MAP kinase
IL-6
Osteoblast

ABSTRACT

We have previously reported that prostaglandin $F_{2\alpha}$ (PGF_{2α}) stimulates interleukin-6 (IL-6), a potent bone resorptive agent, through p44/p42 mitogen-activated protein (MAP) kinase in osteoblast-like MC3T3-E1 cells. In the present study, we investigated whether Rho-kinase is implicated in the PGF_{2α}-stimulated IL-6 synthesis in MC3T3-E1 cells. PGF_{2α} time-dependently induced the phosphorylation of myosin phosphatase targeting subunit (MYPT-1), a Rho-kinase substrate. Y27632, a specific Rho-kinase inhibitor, significantly reduced the PGF_{2α}-stimulated IL-6 synthesis as well as the MYPT-1 phosphorylation. Fasudil, another inhibitor of Rho-kinase, suppressed the PGF_{2α}-stimulated IL-6 synthesis. Y27632 and fasudil failed to affect the PGF_{2α}-induced phosphorylation of p44/p42 MAP kinase. SB203580 and BIRB0796, potent inhibitors of p38 MAP kinase, suppressed the IL-6 synthesis induced by PGF_{2α}. While SP600125, an inhibitor of stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK), failed to reduce the synthesis. Y27632 as well as fasudil attenuated the PGF_{2α}-induced phosphorylation of p38 MAP kinase. These results strongly suggest that Rho-kinase regulates PGF_{2α}-stimulated IL-6 synthesis via p38 MAP kinase activation in osteoblasts.

© 2008 Published by Elsevier Ireland Ltd.

1. Introduction

It is well recognized that interleukin-6 (IL-6) is a multifunctional cytokine that has crucial effects on a wide range of functions such as promoting B cell differentiation, T cell activation and inducing acute phase proteins (Akira et al., 1993; Heymann and Rousselle, 2000; Kwan Tat et al., 2004). The bone metabolism is regulated mainly by two functional cells, osteoblasts and osteoclasts, responsible for bone formation and bone resorption, respectively (Nijweide et al., 1986). As for bone metabolism, IL-6 has been shown to stimulate bone resorption and promote osteoclast formation (Ishimi et al., 1990; Roodman, 1992; Heymann and Rousselle, 2000; Kwan Tat et al., 2004). It has been reported that potent bone resorptive agents such as tumor necrosis factor- α and IL-1 stimulate IL-6 synthesis in osteoblasts (Ishimi et al., 1990; Littlewood et al., 1991; Helle et al., 1998). Currently, evidence is accumulating that IL-6 secreted from osteoblasts plays a pivotal role as a downstream effector of bone resorptive agents in bone metabolism.

It is well known that prostaglandins (PGs) act as autocrine/paracrine modulators of osteoblasts (Nijweide et al., 1986; Pilbeam

et al., 1996). Among them, PGF_{2α} is recognized to be a potent bone resorptive agent in bone metabolism (Pilbeam et al., 1996). It has been reported that PGF_{2α} stimulates the proliferation of osteoblasts and inhibits the differentiation (Pilbeam et al., 1996). In our previous studies (Miwa et al., 1990; Kozawa et al., 1994), we showed that PGF_{2α} induces the activation of protein kinase C via phosphoinositide-hydrolyzing phospholipase C and phosphatidylcholine-hydrolyzing phospholipase D in osteoblast-like MC3T3-E1 cells. Regarding with IL-6 synthesis in osteoblasts, we have reported that PGF_{2α} stimulates IL-6 synthesis via p44/p42 MAP kinase in MC3T3-E1 cells (Kozawa et al., 1997; Tokuda et al., 1999). However, the exact mechanism behind PGF_{2α}-stimulated IL-6 synthesis in osteoblasts remains to be clarified.

Recent studies suggest that Rho and the down-stream effector, Rho-associated kinase (Rho-kinase) play crucial roles in a variety of cellular functions such as cell motility and smooth muscle contraction (Fukata et al., 2001; Riento and Ridley, 2003; Shimokawa and Rashid, 2007). As for osteoblasts, it has been demonstrated that Rho-kinase and p38 MAP kinase are involved in the endothelin-1-induced expression of PG endoperoxide G/H synthase mRNA in osteoblasts (Windischhofer et al., 2002). In addition, it has been shown that the Rho/Rho-kinase pathway stimulates osteoblast proliferation whereas it inhibits osteoblast differentiation (Harney et

* Corresponding author. Tel.: +81 58 230 6214; fax: +81 58 230 6215.
E-mail address: okozawa@gifu-u.ac.jp (O. Kozawa).

al., 2004). However, the exact role of Rho-kinase in osteoblasts has not yet been fully clarified.

In the present study, we further investigated the exact mechanism behind $\text{PGF}_{2\alpha}$ -stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells. We here show that Rho-kinase regulates $\text{PGF}_{2\alpha}$ -stimulated IL-6 synthesis through p38 MAP kinase activation in these cells.

2. Materials and methods

2.1. Materials

$\text{PGF}_{2\alpha}$ and mouse IL-6 enzyme immunoassay (ELISA) kit were purchased from R&D Systems, Inc. (Minneapolis, MN). Y27632, SB203580 and SP600125 were obtained from Calbiochem–Novabiochem, Co. (La Jolla, CA). Hydroxyfasudil (fasudil) was purchased from Sigma–Aldrich, Co. (St. Louis, MO). BIRB0796 was obtained from Dr. Philip Cohen (University of Dundee, UK). Phospho-specific MYPT-1 antibodies were purchased from Millipore, Co. (Billerica, MA). MYPT-1 antibodies were obtained from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies and p38 MAP kinase antibodies were purchased from Cell-Signaling Technology, Inc. (Beverly, MA). ECL Western blotting detection system was purchased from GE Healthcare UK, Ltd. (Buckinghamshire, England). Other materials and chemicals were obtained from commercial sources. Y27632, SB203580, SP600125 and BIRB0796 were dissolved in dimethylsulfoxide. The maximum concentration of dimethylsulfoxide was 0.1%, which did not affect the assay for IL-6 or Western blot analysis.

2.2. Cell culture

Cloned osteoblast-like MC3T3-E1 cells derived from newborn mouse calvaria (Sudo et al., 1983) were maintained as previously described (Kozawa et al., 1992). Briefly, the cells were cultured in α -minimum essential medium (α -MEM) containing 10% fetal calf serum (FCS) at 37 °C in a humidified atmosphere of 5% $\text{CO}_2/95\%$ air. The cells were seeded into 35-mm or 90-mm diameter dishes in α -MEM containing 10% FCS. After 5 days, the medium was exchanged for α -MEM containing 0.3% FCS. The cells were used for experiments after 48 h.

2.3. IL-6 assay

The cultured cells were stimulated by 10 μM $\text{PGF}_{2\alpha}$ in 1 ml of α -MEM containing 0.3% FCS for the indicated periods. When indicated, the cells were pretreated with various doses of Y27632, fasudil, SB203580, BIRB0796 or SP600125 for 60 min. The conditioned medium was collected at the end of the incubation, and the IL-6 concentration was measured by ELISA kit.

2.4. Western blot analysis

Western blot analysis was performed as follows. The cultured cells were pretreated with various doses of Y27632 or fasudil for 60 min, and then stimulated by $\text{PGF}_{2\alpha}$ in α -MEM containing 0.3% FCS for the indicated periods. The cells were washed twice with phosphate-buffered saline and then lysed, homogenized and sonicated in a lysis buffer containing 62.5 mM Tris/HCl; pH 6.8, 2% sodium dodecyl sulfate (SDS), 50 mM dithiothreitol and 10% glycerol. The cytosolic fraction was collected as a supernatant after centrifugation at 125,000 $\times g$ for 10 min at 4 °C. Twenty micrograms of cytosolic protein was charged, and SDS-polyacrylamide gel electrophoresis (PAGE) was performed according to Laemmli (1970) in 10% polyacrylamide gel. The fractionated proteins were then transferred onto a PVDF Membrane (Bio-Rad, Hercules, CA). The protein-transferred membrane was blocked with 5% fat-free dry milk in Tris-buffered saline–Tween 20 (TBS-T; 20 mM Tris/HCl, pH 7.6, 137 mM NaCl, 0.1% Tween-20) for 2 h before incubation with the primary antibodies. The rabbit polyclonal phospho-specific MYPT-1 antibodies, MYPT-1 antibodies, phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies or p38 MAP kinase antibodies were used as primary antibodies. Peroxidase-labeled antibodies raised in goat against rabbit IgG (KPL Inc., Geithersburg, MD) were used as second antibodies. The first and second antibodies were diluted at 1:1000 with 5% fat-free dry milk in TBS-T. Peroxidase activity on the membrane was visualized on X-ray film by means of the ECL Western blotting detection system.

2.5. Determination

The absorbance of enzyme immunoassay samples was measured at 450 nm with EL 340 Bio Kinetic Reader (Bio-Tek Instruments, Inc., Winooski, VT). The densitometric analysis was performed using Molecular Analyst/Macintosh (Bio-Rad Laboratories). Protein concentrations were determined by using a protein assay reagent kit (Thermo Fisher Scientific K.K., Rockford, IL) with BSA as a reference protein.




Fig. 1. Effects of $\text{PGF}_{2\alpha}$ on the phosphorylation of MYPT-1 in MC3T3-E1 cells. The cultured cells were stimulated by 10 μM $\text{PGF}_{2\alpha}$ for the indicated periods. Twenty micrograms of cytosolic protein were subjected to SDS-PAGE with subsequent Western blot analysis with antibodies against phospho-specific MYPT-1 or MYPT-1. The histogram shows quantitative representations of the levels of $\text{PGF}_{2\alpha}$ -induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. Values represent the amount of phospho-MYPT-1 divided by those of total MYPT-1. * $p < 0.05$, compared to the value of control (lane 1).

2.6. Statistical analysis

The data were analyzed by ANOVA followed by the Bonferroni method for multiple comparisons between pairs, and a $p < 0.05$ was considered significant. All data are presented as the mean \pm S.E.M. of triplicate determinations. Each experiment was repeated three times with similar results.

3. Results

3.1. Effects of $\text{PGF}_{2\alpha}$ on the phosphorylation of MYPT-1 in MC3T3-E1 cells

It is generally recognized that MYPT-1, which is a component of myosin phosphatase, is a down-stream substrate of Rho-kinase (Fukata et al., 2001; Ito et al., 2004). In order to clarify whether $\text{PGF}_{2\alpha}$ activates Rho-kinase in osteoblast-like MC3T3-E1 cells, we examined the effect of $\text{PGF}_{2\alpha}$ on the phosphorylation of MYPT-1. $\text{PGF}_{2\alpha}$ markedly induced the phosphorylation of MYPT-1 in a time-dependent manner (Fig. 1). The effect of $\text{PGF}_{2\alpha}$ on the phosphorylation of MYPT-1 reached its peak within 2 min and decreased thereafter (Fig. 1).

Y27632, an inhibitor of Rho-kinase (Shimokawa and Rashid, 2007), suppressed the $\text{PGF}_{2\alpha}$ -induced phosphorylation levels of MYPT-1 (Fig. 2A). In addition, fasudil, another inhibitor of Rho-kinase (Shimokawa and Rashid, 2007), reduced the $\text{PGF}_{2\alpha}$ -induced levels of MYPT-1 phosphorylation (Fig. 2B).

3.2. Effects of Y27632 or fasudil on the $\text{PGF}_{2\alpha}$ -stimulated IL-6 synthesis in MC3T3-E1 cells

We previously showed that $\text{PGF}_{2\alpha}$ stimulates IL-6 synthesis in osteoblast-like MC3T3-E1 cells (Kozawa et al., 1997). In order to investigate whether Rho-kinase is involved in the $\text{PGF}_{2\alpha}$ -induced




Fig. 2. Effects of Y27632 or fasudil on the PGF_{2α}-induced phosphorylation of MYPT-1 in MC3T3-E1 cells. The cultured cells were pretreated with 10 μM Y27632 (A), 10 μM fasudil (B) or vehicle for 60 min, and then stimulated by 10 μM PGF_{2α} or vehicle for 2 min. Twenty micrograms of cytosolic protein were subjected to SDS-PAGE with subsequent Western blot analysis with antibodies against phospho-specific MYPT-1 or MYPT-1. The histogram shows quantitative representations of the levels of PGF_{2α}-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean ± S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. Values represent the amount of phospho-MYPT-1 divided by those of total MYPT-1. **p* < 0.05, compared to the control (lanes 1). ***p* < 0.05, compared to the value of PGF_{2α} alone (lanes 2).

synthesis of IL-6 in MC3T3-E1 cells, we next examined the effect of Y27632 on the synthesis of IL-6 induced by PGF_{2α}. Y27632, which by itself had little effect on the IL-6 levels, significantly suppressed the PGF_{2α}-induced synthesis of IL-6 (Fig. 3). The inhibitory effect of Y27632 was dose-dependent in the range between 0.1 and 10 μM. Y27632 (10 μM) caused approximately 80% inhibition in the PGF_{2α}-effect.

Fasudil as well as Y27632 reduced the PGF_{2α}-stimulated IL-6 synthesis in these cells (Fig. 4). The effect of fasudil on the IL-6 synthesis was dose-dependent in the range between 0.1 and 10 μM. Fasudil (10 μM) caused about 90% inhibition in the PGF_{2α}-effect.




Fig. 3. Effect of Y27632 on the PGF_{2α}-induced IL-6 synthesis in MC3T3-E1 cells. The cultured cells were pretreated with various doses of Y27632 for 60 min, and then stimulated by 10 μM PGF_{2α} or vehicle for 48 h. Each value represents the mean ± S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. **p* < 0.05, compared to the value of PGF_{2α} alone.

3.3. Effects of Y27632 or fasudil on the PGF_{2α}-induced phosphorylation of p44/p42 MAP kinase in MC3T3-E1 cells

We have previously reported that PGF_{2α} stimulates IL-6 synthesis via p44/p42 MAP kinase in osteoblast-like MC3T3-E1 cells




Fig. 4. Effect of fasudil on the PGF_{2α}-induced IL-6 synthesis in MC3T3-E1 cells. The cultured cells were pretreated with various doses of fasudil for 60 min, and then stimulated by 10 μM PGF_{2α} or vehicle for 48 h. Each value represents the mean ± S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. **p* < 0.05, compared to the value of PGF_{2α} alone.




Fig. 5. Effect of Y27632 on the $\text{PGF}_{2\alpha}$ -induced phosphorylation of p44/p42 MAP kinase in MC3T3-E1 cells. The cultured cells were pretreated with various doses of Y27632 for 60 min, and then stimulated by $10 \mu\text{M}$ $\text{PGF}_{2\alpha}$ or vehicle for 20 min. Twenty micrograms of cytosolic protein were subjected to SDS-PAGE with subsequent Western blot analysis with antibodies against phospho-specific p44/p42 MAP kinase or p44/p42 MAP kinase. The histogram shows quantitative representations of the levels of $\text{PGF}_{2\alpha}$ -induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. Values represent the amount of phospho-p44/p42 MAP kinase divided by those of total p44/p42 MAP kinase.

(Tokuda et al., 1999). In order to investigate whether Rho-kinase-effect on the $\text{PGF}_{2\alpha}$ -stimulated IL-6 synthesis is dependent upon the activation of p44/p42 MAP kinase in MC3T3-E1 cells, we next examined the effect of Y27632 on the phosphorylation of p44/p42 MAP kinase by $\text{PGF}_{2\alpha}$. However, Y27632 did not affect the $\text{PGF}_{2\alpha}$ -induced phosphorylation of p44/p42 MAP kinase (Fig. 5). In addition, fasudil had little effect on the phosphorylation levels of p44/p42 MAP kinase (data not shown).

3.4. Effects of SB203580, BIRB0796 or SP600125 on the $\text{PGF}_{2\alpha}$ -stimulated IL-6 synthesis in MC3T3-E1 cells

It is currently known that three MAP kinases, p44/p42 MAP kinase, p38 MAP kinase and stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) are known as central elements used by mammalian cells to transduce the various messages of a variety of agonists (Widmann et al., 1999). To investigate the involvement of other MAP kinases in the $\text{PGF}_{2\alpha}$ -stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells, we examined the effect of SB203580, a specific p38 MAP kinase inhibitor (Cuenda et al., 1995), on the IL-6 synthesis. We have previously shown that $\text{PGF}_{2\alpha}$ stimulates the activation of p38 MAP kinase and SAPK/JNK in MC3T3-E1 cells (Tokuda et al., 2007). SB203580 significantly suppressed the $\text{PGF}_{2\alpha}$ -stimulated synthesis of IL-6 (Table 1). The inhibitory effect of SB203580 was dose-dependent in the range between 1 and $10 \mu\text{M}$. BIRB0796, a more potent inhibitor of p38 α and p38 β MAP kinases than SB203580 (Bain et al., 2007), significantly suppressed the $\text{PGF}_{2\alpha}$ -stimulated synthesis of IL-6 (Fig. 6). The inhibitory effect of BIRB0796 was dose-dependent in the range between 0.1 and $1 \mu\text{M}$. On the contrary, SP600125, a specific inhibitor of SAPK/JNK (Bennett et al., 2001), did not suppress the IL-6 synthesis stimulated by $\text{PGF}_{2\alpha}$ (data not shown).

Table 1

Effect of SB203580 on the $\text{PGF}_{2\alpha}$ -stimulated IL-6 synthesis in MC3T3-E1 cells

SB203580 (μM)	$\text{PGF}_{2\alpha}$	IL-6 (pg/ml)
0	-	<7.8
0	+	697 \pm 62*
1	-	<7.8
1	+	377 \pm 3**
3	-	<7.8
3	+	213 \pm 11**
10	-	<7.8
10	+	154 \pm 5**

The cultured cells were pretreated with various doses of SB203580 for 60 min, and then stimulated by $10 \mu\text{M}$ $\text{PGF}_{2\alpha}$ or vehicle for 48 h. Each value represents the mean \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. * $p < 0.05$, compared to the control. ** $p < 0.05$, compared to the value of $\text{PGF}_{2\alpha}$ alone.

3.5. Effects of Y27632 or fasudil on the $\text{PGF}_{2\alpha}$ -induced phosphorylation of p38 MAP kinase in MC3T3-E1 cells

In addition, we examined effect of Y27632 on the $\text{PGF}_{2\alpha}$ -induced phosphorylation of p38 MAP kinase in MC3T3-E1 cells. Y27632 markedly suppressed the $\text{PGF}_{2\alpha}$ -induced phosphorylation of p38 MAP kinase (Fig. 7A). The Y27632-effect on the phosphorylation levels was dose-dependent in the range between 1 and $10 \mu\text{M}$. Y27632 ($3 \mu\text{M}$) caused about 80% inhibition in the $\text{PGF}_{2\alpha}$ -effect.

Fasudil reduced the $\text{PGF}_{2\alpha}$ -induced levels of phosphorylated-p38 MAP kinase (Fig. 7B). The inhibitory effect of fasudil was dose-dependent in the range between 1 and $10 \mu\text{M}$. Fasudil ($10 \mu\text{M}$) caused approximately 90% inhibition in the $\text{PGF}_{2\alpha}$ -effect.

4. Discussion

In the present study, we showed that $\text{PGF}_{2\alpha}$ time-dependently induced the phosphorylation of MYPT-1 in osteoblast-like




Fig. 6. Effect of BIRB0796 on the $\text{PGF}_{2\alpha}$ -induced IL-6 synthesis in MC3T3-E1 cells. The cultured cells were pretreated with various doses of BIRB0796 for 60 min, and then stimulated by $10 \mu\text{M}$ $\text{PGF}_{2\alpha}$ or vehicle for 48 h. Each value represents the mean \pm S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. * $p < 0.05$, compared to the value of $\text{PGF}_{2\alpha}$ alone.




Fig. 7. Effects of Y27632 or fasudil on the PGF_{2α}-induced phosphorylation of p38 MAP kinase in MC3T3-E1 cells. The cultured cells were pretreated with various doses of Y27632 (A) or fasudil (B) for 60 min, and then stimulated by 10 μM PGF_{2α} or vehicle for 10 min. Twenty micrograms of cytosolic protein were subjected to SDS-PAGE with subsequent Western blot analysis with antibodies against phospho-specific p38 MAP kinase or p38 MAP kinase. The histogram shows quantitative representations of the levels of PGF_{2α}-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean ± S.E.M. of triplicate determinations. Similar results were obtained with two additional and different cell preparations. Values represent the amount of phospho-p38 MAP kinase divided by those of total p38 MAP kinase. **p* < 0.05, compared to the control (lanes 1). ***p* < 0.05, compared to the value of PGF_{2α} alone (lanes 2).

MC3T3-E1 cells, using phospho-specific MYPT-1 (Thr850) antibodies. MYPT, which is a myosin-binding subunit of myosin phosphatase and regulates the interaction of actin and myosin, is well known to be a downstream target of Rho-kinase (Fukata et al., 2001; Ito et al., 2004). Additionally, we next demonstrated that Y27632 and fasudil, inhibitors of Rho-kinase (Shimokawa and Rashid, 2007), truly suppressed the PGF_{2α}-induced phosphorylation of MYPT-1. Y27632 also reduced the basal MYPT-1 phosphorylation, whereas fasudil did not suppress it. It has been reported that Y27632 also inhibits other kinases like PKCδ with a

similar potency to that for Rho kinase 2 (Davies et al., 2000; Eto et al., 2001). The discrepancy between the effects of Y27632 and fasudil on basal phosphorylation of MYPT-1 might be due to the unselectivity of Y27632. Therefore, our results indicate that PGF_{2α} elicits the activation of Rho-kinase in osteoblast-like MC3T3-E1 cells.

We next investigated the involvement of Rho-kinase in the PGF_{2α}-stimulated IL-6 synthesis or not in osteoblast-like MC3T3-E1 cells. Y27632 (Shimokawa and Rashid, 2007) significantly reduced the PGF_{2α}-stimulated synthesis of IL-6. This finding suggests that the PGF_{2α}-activated Rho-kinase functions as a positive regulator in the IL-6 synthesis in these cells. In addition, we showed that the IL-6 synthesis stimulated by PGF_{2α} was inhibited by fasudil (Shimokawa and Rashid, 2007) as well as Y27632. Therefore, taking our results into account, it is most likely that PGF_{2α} activates Rho-kinase, resulting in up-regulation of IL-6 synthesis in osteoblast-like MC3T3-E1 cells.

With regard to IL-6 synthesis in osteoblasts, we have previously reported that the activation of p44/p42 MAP kinase is involved in the PGF_{2α}-stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells (Tokuda et al., 1999). Additionally, we investigated the relationship between Rho-kinase and p44/p42 MAP kinase in the PGF_{2α}-stimulated IL-6 synthesis in these cells. However, Y27632 or fasudil had little effect on the PGF_{2α}-induced phosphorylation levels of p44/p42 MAP kinase. Based on our findings, it seems unlikely that Rho-kinase affects the PGF_{2α}-stimulated IL-6 synthesis through the modulation of p44/p42 MAP kinase in osteoblast-like MC3T3-E1 cells. It is generally recognized that SAPK/JNK and p38 MAP kinase in addition to p44/p42 MAP kinase are known as central elements used by mammalian cells to transduce the various messages (Widmann et al., 1999). We have shown that PGF_{2α} stimulates the activation of SAPK/JNK and p38 MAP kinase in MC3T3-E1 cells (Tokuda et al., 2007). In the present study, the IL-6 synthesis induced by PGF_{2α} was suppressed by SB203580, a specific inhibitor of p38 MAP kinase (Cuenda et al., 1995) but not SP600125, a specific inhibitor of SAPK/JNK (Bennett et al., 2001). It has been reported that SB203580 inhibits other than p38 MAP kinase protein kinases with similar or even greater (receptor-interacting protein 2, RIP2) potency (Godl et al., 2003). As RIP2-mediated NF-κB activation reportedly leads to increase IL-6 (Sarkar et al., 2006), it is possible that SB203580-induced suppression of IL-6 synthesis is due to RIP2 blocking by SB203580. Thus, we further examined the effect of BIRB0796 (Bain et al., 2007) on the PGF_{2α}-stimulated IL-6 synthesis in MC3T3-E1 cells, and found that BIRB0796 significantly suppressed the PGF_{2α}-stimulated IL-6 synthesis. These results strongly suggest that p38 MAP kinase as well as p44/p42 MAP kinase plays a part in the PGF_{2α}-stimulated IL-6 synthesis. Next, we tried to elucidate the relationship between Rho-kinase and p38 MAP kinase in the PGF_{2α}-stimulated IL-6 synthesis in these cells. The PGF_{2α}-induced phosphorylation levels of p38 MAP kinase were markedly suppressed by Y27632 and fasudil. Therefore, it is probable that Rho-kinase regulates the PGF_{2α}-stimulated IL-6 synthesis via p38 MAP kinase. In the present study, the maximum effect on the phosphorylation of MYPT-1, a well-known downstream target of Rho-kinase (Fukata et al., 2001), was observed within 2 min after the PGF_{2α} stimulation. In our previous study (Tokuda et al., 2007), we have shown that the phosphorylation of p38 MAP kinase reach the peak at 10 min after the stimulation of PGF_{2α} in MC3T3-E1 cells. The time course of the PGF_{2α}-induced phosphorylation of MYPT-1 seems to be faster than that of p38 MAP kinase. Therefore, it seems reasonable that PGF_{2α}-induced activation of p38 MAP kinase subsequently occurs after the Rho-kinase activation. Taking our findings into account as a whole, our results strongly suggest that Rho-kinase acts at a point upstream from p38 MAP kinase in the PGF_{2α}-stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells.

It is currently recognized that Rho-kinase plays an important role in a variety of cellular functions, especially vascular smooth muscle contraction (Fukata et al., 2001; Riento and Ridley, 2003; Shimokawa and Rashid, 2007). In bone metabolism, it has been reported that the activation of Rho-kinase suppresses the differentiation of osteoblasts and induces their proliferation (Harmey et al., 2004). Our present results suggest that Rho-kinase in osteoblasts acts as positive regulator in the synthesis of IL-6, one of the central modulators of bone metabolism. It is well known that IL-6 produced by osteoblasts is a potent bone resorptive agent and induces osteoclast formation (Ishimi et al., 1990; Kwan Tat et al., 2004). In addition, it is recognized that PGF_{2α}, an inducer of IL-6, acts as a bone resorptive agent in bone metabolism (Nijweide et al., 1986; Pilbeam et al., 1996). During the fracture healing, once mineralized cartilage resorption is completed and woven bone formation has occurred, a subsequent phase of bone resorption takes place in which IL-6 increases in their expression (Louis and Einhorn, 2006). Therefore, our present findings lead us to speculate that PGF_{2α}-activated Rho-kinase in osteoblasts functions as a positive regulator of bone resorption via the fine-tuning of the local cytokine network, such as induction of IL-6. The Rho-kinase pathway in osteoblasts might be a new candidate as a molecular therapeutic target of fracture healing, bone formation and bone metabolic disease such as osteoporosis. The exact mechanism of Rho-kinase activation in osteoblasts is still not precisely known. Further investigations are required to elucidate the exact role of Rho-kinase in bone metabolism.

Taken together, our results strongly suggest that Rho-kinase regulates the PGF_{2α}-stimulated IL-6 synthesis via p38 MAP kinase in osteoblasts.

Acknowledgements

We are very grateful to Yoko Kawamura for her skillful technical assistance. This investigation was supported in part by Grant-in-Aid for Scientific Research (16590873 and 16591482) for the Ministry of Education, Science, Sports and Culture of Japan, the Research Grants for Longevity Sciences (17A-3), Research Grant on Proteomics and Research Grant on Longevity Sciences from the Ministry of Health, Labour and Welfare of Japan.

References

Akita, S., Taga, T., Kishimoto, T., 1993. Interleukin-6 in biology and medicine. *Adv. Immunol.* 54, 1–78.

Bain, J., Plater, L., Matt, E., Shpiro, N., Hastie, J., McLaughlan, H., Klevernic, I., Arthur, S., Alessi, D., Cohen, P., 2007. The selectivity of protein kinase inhibitors: a further update. *Biochem. J.* 408, 297–315.

Bennett, B.L., Sasaki, D.T., Murray, B.W., O'Leary, E.C., Sakata, S.T., Xu, W., Leisten, J.C., Motiwala, A., Pierce, S., Satoh, Y., Bhagwat, S.S., Manning, A.M., Anderson, D.W., 2001. SP600125, an anthrapyrazolone inhibitor of Jun N-terminal Kinase. *Proc. Natl. Acad. Sci. U.S.A.* 98, 13681–13686.

Cuenda, A., Rouse, J., Doza, Y.N., Meier, R., Cohen, P., Gallagher, T.F., Young, P.R., Lee, J.C., 1995. SB203580 is a specific inhibitor of a MAP kinase homologue which is stimulated by cellular stresses and interleukin-1. *FEBS Lett.* 364, 229–233.

Davies, S., Reddy, H., Caivano, M., Cohen, P., 2000. Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem. J.* 351, 95–105.

Eto, M., Kitagawa, T., Yazawa, M., Mukai, H., Ono, Y., Brautigan, David, L., 2001. Histamine-induced vasoconstriction involves phosphorylation of a specific inhibitor protein for myosin phosphatase by protein kinase C α and δ isoforms. *J. Biol. Chem.* 276, 29072–29078.

Fukata, Y., Amano, M., Kaibuchi, K., 2001. Rho-Rho-kinase pathway in smooth muscle contraction and cytoskeletal reorganization of non-muscle cells. *Trends Pharmacol. Sci.* 22, 32–39.

Godl, K., Wissing, J., Kurtenbach, A., Habenberger, P., Blencke, S., Gutbrod, H., Salasid, K., Stein-Gerlach, M., Missio, A., Cotton, M., Daub, H., 2003. An efficient proteomics method to identify the cellular targets of protein kinase inhibitors. *Proc. Natl. Acad. Sci. U.S.A.* 100, 15434–15439.

Harmey, D., Stenbeck, G., Nobes, C.D., Lax, A.J., Grigoriadis, A.E., 2004. Regulation of osteoblast differentiation by Pasteurellamultocida toxin (PMT): a role for Rho GTPase in bone formation. *J. Bone Miner. Res.* 19, 661–670.

Heile, M., Brakenhoff, J.P.J., DeGroot, E.R., Aarden, L.A., 1998. Interleukin 6 is involved in interleukin 1-induced activities. *Eur. J. Immunol.* 18, 957–959.

Heymann, D., Rousselle, A.V., 2000. gp130 Cytokine family and bone cells. *Cytokine* 12, 1455–1468.

Ishimi, Y., Miyaura, C., Jin, C.H., Akatsu, T., Abe, F., Nakamura, Y., Yamaguchi, Y., Yoshiki, S., Matsuda, T., Hirano, T., Kishimoto, T., Suda, T., 1990. IL-6 is produced by osteoblasts and induces bone resorption. *J. Immunol.* 145, 3297–3303.

Ito, M., Nakano, T., Erdodi, F., Hartshorne, D.J., 2004. Myosin phosphatase: structure, regulation and function. *Mol. Cell. Biochem.* 259, 197–209.

Kozawa, O., Tokuda, H., Miwa, M., Kotoyori, J., Oiso, Y., 1992. Cross-talk regulation between cyclic AMP production and phosphoinositide hydrolysis induced by prostaglandin E₂ in osteoblast-like cells. *Exp. Cell Res.* 198, 130–134.

Kozawa, O., Suzuki, A., Kotoyori, J., Tokuda, H., Watanabe, Y., Ito, Y., Oiso, Y., 1994. Prostaglandin F_{2α} activates phospholipase D independently from activation of protein kinase C in osteoblast-like cells. *J. Cell. Biochem.* 55, 373–379.

Kozawa, O., Suzuki, A., Tokuda, H., Uematsu, T., 1997. Prostaglandin F_{2α} stimulates interleukin-6 via activation of PKC in osteoblast-like cells. *Am. J. Physiol.* 272, E208–E211.

Kwan Tat, S., Padrines, M., Theoleyre, S., Heymann, D., Fortun, Y., 2004. IL-6, RANKL, TNF- α /IL-1: interrelations in bone resorption pathophysiology. *Cytokine Growth Factor Rev.* 15, 49–60.

Laemmli, U.K., 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.

Littlewood, A.J., Russil, J., Harvey, G.R., Hughes, D.E., Russel, R.G.G., Gowen, M., 1991. The modulation of the expression of IL-6 and its receptor in human osteoblasts in vitro. *Endocrinology* 129, 1513–1520.

Louis, C., Einhorn, A.T., 2006. In: Kleerekoper, M., Lane, N. (Eds.), *Fracture Healing: The Biology of Bone Repair and Regeneration in Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*, sixth ed. The American Society for Bone and Mineral Research, USA, pp. 42–48.

Miwa, M., Tokuda, H., Tsushita, K., Kotoyori, J., Takahashi, Y., Ozaki, N., Kozawa, O., Oiso, Y., 1990. Involvement of pertussis-toxin-sensitive GTP-binding protein in prostaglandin F_{2α}-induced phosphoinositidylhydrolysis in osteoblast-like cells. *Biochem. Biophys. Res. Commun.* 171, 1229–1235.

Nijweide, P.J., Burger, E.H., Feyen, J.H.M., 1986. Cells of bone: proliferation, differentiation, and hormonal regulation. *Physiol. Rev.* 66, 855–886.

Pilbeam, C.C., Harrison, J.R., Raisz, L.G., 1996. *Prostaglandins and bone metabolism*. In: Raisz, L.G., Rodan, G.A. (Eds.), *Principles of Bone Biology*. Academic Press, San Diego, USA, pp. 715–728.

Riento, K., Ridley, A.J., 2003. Rocks: multifunctional kinases in cell behaviour. *Nat. Rev. Mol. Cell Biol.* 4, 446–456.

Roodman, G.D., 1992. Interleukin-6: an osteotropic factor? *J. Bone Miner. Res.* 7, 475–478.

Sarkar, A., Duncan, M., Hart, J., Hertlein, E., Guttridge, D.C., Wewers, M.D., 2006. ASC directs NF- κ B activation by regulating receptor interacting protein-2 (RIP2) caspase-1 interactions. *J. Immunol.* 176, 4979–4986.

Shimokawa, H., Rashid, M., 2007. Development of Rho-kinase inhibitors for cardiovascular medicine. *Trends Pharmacol. Sci.* 28, 296–302.

Sudo, H., Kodama, H., Amagai, Y., Yamamoto, S., Kasai, S., 1983. In vitro differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria. *J. Cell Biol.* 96, 191–198.

Tokuda, H., Kozawa, O., Harada, A., Uematsu, T., 1999. p42/p44 mitogen-activated protein kinase activation is involved in prostaglandin F_{2α}-induced interleukin-6 synthesis in osteoblasts. *Cell. Signal.* 11, 325–330.

Tokuda, H., Takai, S., Matsushima-Nishiwaki, R., Akamatsu, S., Hanai, Y., Hosoi, T., Harada, A., Ohta, T., Kozawa, O., 2007. (-)-Epigallocatechingallate enhances prostaglandin F_{2α}-induced VEGF synthesis via up-regulating SAPK/JNK activation in osteoblasts. *J. Cell Biochem.* 100, 1146–1153.

Widmann, C., Gibson, S., Jarpe, M.B., Johnson, G.L., 1999. Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human. *Physiol. Rev.* 79, 143–180.

Windischhofer, W., Zach, D., Fauler, G., Rasporinig, G., Kofeler, H., Leis, H.J., 2002. Involvement of Rho and p38 MAPK in endothelin-1-induced expression of PGHS-2 mRNA in osteoblast-like cells. *J. Bone Miner. Res.* 17, 1774–1784.

Raloxifene-Induced Acceleration of Platelet Aggregation

Chiho Minamitani^{1,3}, Shinji Takai², Rie Matsushima-Nishiwaki², Yoshiteru Hanai^{1,2},
Takanobu Otuka³, Osamu Kozawa² and Haruhiko Tokuda^{1,2}

Abstract

A 59-year-old postmenopausal woman diagnosed to have primary osteoporosis began to take 60 mg daily of oral raloxifene. The platelet aggregation induced by 1 μ M adenosine diphosphate (ADP) and the α 2-antiplasmin activity were accelerated significantly after 8 weeks from the beginning of raloxifene-treatment, and gradually deteriorated up to 24 weeks. ADP markedly caused the phosphorylation of Akt in the platelets obtained at 24 weeks. Although there were no subjective complaints at 24 weeks, the medication was stopped with her consent to avoid any adverse effects due to thrombus formation. The platelet hyper-aggregability and Akt phosphorylation induced by ADP disappeared at 4 weeks after the cessation of medication. These results strongly suggest that raloxifene caused the acceleration of platelet aggregation and subclinical thrombus formation through the Akt signal pathway in this case.

Key words: raloxifene, osteoporosis, Akt, platelet aggregation

(Inter Med 47: 1523-1528, 2008)

(DOI: 10.2169/internalmedicine.47.0732)

Introduction

Raloxifene, a selective estrogen receptor modulator (SERM), has been well recognized to be an efficient therapeutic agent for postmenopausal osteoporosis (1). Raloxifene reportedly increases the vertebra and femoral neck bone mineral density, and reduces the risk of vertebral fracture (1). In addition, raloxifene has been reported to significantly reduce the incidence of breast cancer (1). Raloxifene is thought to be a preferable medicine for postmenopausal women. However, the use of this drug as well as estrogen is associated with an increased risk of developing venous thromboembolus (2). Platelet aggregation and the coagulation system are important for the formation of the thrombus. Although significant attention has been paid to the coagulation system, the influence of raloxifene on platelet aggregation is not known.

Platelet aggregation is usually measured using either the optical density (OD) method (3) or the impedance method (4), both of which are indispensable for the clinical evaluation of platelet function. However, these methods provide

little information about subtle but crucial changes in the number of platelet microaggregates in response to small changes in the aggregating stimuli or the proaggregatory status. The light-scattering (LS) method, which has primarily been used for experimental research, provides a tool with a greater sensitivity for detecting microaggregates of platelets than the conventional light transmittance method (4). A particle-counting method that employs LS has recently been developed (5), thus allowing identification of particle size in terms of light intensity and minimizing the interference by neighboring platelets, which may attenuate the high intensity light scattered by larger particles (6). Recently, an important role of platelet microaggregation in the circulation and a significant alteration of platelet microaggregation in pathogenesis were reported by the creation of a laser LS system for microaggregates (7-9). Therefore, an LS system was employed for the detection of microaggregates of platelets in our clinical study (10).

Adenosine diphosphate (ADP) is considered to be a weak physiological agonist for platelet aggregation by itself in comparison, for example, to thrombin or collagen (11). However, ADP is a necessary cofactor for normal activation

¹Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, Obu, ²Department of Pharmacology, Gifu University Graduate School of Medicine, Gifu and ³Department of Orthopedic Surgery, Nagoya City University Graduate School of Medical Sciences, Nagoya

Received for publication November 6, 2007; Accepted for publication May 22, 2008

Correspondence to Dr. Haruhiko Tokuda, tokuda@ncgg.go.jp

of platelets by other agonists. Even if it is weak agonist for platelet aggregation, low-concentration ADP amplifies the effects (12). A recent investigation revealed that purinoreceptor P2Y₁₂ is a principal receptor for mediating ADP-induced platelet aggregation and the thromboembolism (13, 14). Phosphatidylinositol 3-kinase (PI3K) effectors, such as the serine/threonine kinase Akt, play important roles in platelet aggregation as an intracellular signal system downstream of the P2Y₁₂ receptor (14). The current report describes the acceleration of low-dose ADP-induced platelet aggregation with the increasing Akt activation in a patient who was treated with a SERM, raloxifene.

Case Report

A 59-year-old postmenopausal woman, who had suffered from intermittent lumbago for several years, was referred to the outpatient service for osteoporosis. X-ray examination showed a vertebral fracture on T-12. Bone marrow depression (BMD) determined by dual-energy X-ray absorptiometry (QDR-2000) was: lumbar spine 0.803 g/cm³ (T-score, -2.6); total femur 0.598 g/cm³ (T-score, -2.8; Table 1). The blood analysis appeared to be in normal range except the elevation of alkaline phosphatase (ALP) and bone type ALP (Table 2). According to the diagnostic criteria for primary osteoporosis in Japan (15), she was diagnosed to have primary osteoporosis with a high turnover bone metabolism, and then was administered 60 mg daily of oral raloxifene. She agreed to undergo an assessment of her platelet function before and after the treatment, and signed an informed consent agreement as approved by the local institutional review board after receiving a detailed explanation of the protocol. Platelet-rich plasma (PRP) was chronologically obtained from freshly drawn venous blood samples. Platelet aggregation was assessed by PA-200 aggregometer (Kowa, Tokyo, Japan), which can determine the size of platelet aggregates based upon particle counting using light scatter (13). The platelets were preincubated for 2 minutes, then platelet aggregation was monitored for 5 minutes after the addition of various doses of ADP (0.3-30 μM). The percentage of transmittance of the isolated platelets was recorded as 0%, and that of the appropriate platelet-free plasma (blank) was recorded as 100%. ADP concentration that caused 50% of transmittance of the isolated platelets was considered as ED 50. Just after evaluating platelet aggregation, the remaining PRP was sufficiently washed with phosphate buffered saline, separated by sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis (PAGE), and a Western blot analysis was performed using antibodies against phospho-specific Akt (Thr308) to detect the activated form and with Akt antibodies for determination of the total amount, as previously described (16).

Figures 1, 2 show that the ED₅₀ of ADP for the platelet aggregation gradually decreased after treatment with raloxifene. After 8 weeks after the initiation of treatment, less than 1 μM ADP caused the significant acceleration of

Table 1. Clinical Feature

Age	59 years
Sex	female
Height	152.9 cm
Weight	56.4 kg
Body mass index	24.12 kg/m ²
Lumber spine L2-L4 BMD	0.803 g/cm ³
YAM	72%

Footnote: YAM: young adult mean

the platelet aggregation, and it deteriorated up to 24 weeks. Considering these results, although she did not show any symptoms of thrombosis, the administration of raloxifene was terminated to avoid major clinical problems. Four weeks after the cessation of raloxifene, the level of the platelet aggregation was markedly improved to the level before the raloxifene-treatment. According to the analysis of the size of platelet aggregates, a decrease of small aggregates (9-25 μm) and the increase of large aggregates (50-70 μm) were observed after 2 weeks, and an increase of medium aggregates (25-50 μm) was subsequently observed after 4 weeks (Fig. 2), suggesting that the pathological change of platelet function occurred within at least 2 weeks of the raloxifene treatment. In addition, ADP (1 μM) markedly caused the phosphorylation of Akt in the platelets obtained at 24 weeks, but the ADP-induced Akt phosphorylation disappeared in the platelets obtained at 4 weeks after the cessation of raloxifene (Fig. 3).

It is well known that a lack of α₂-antiplasmin (α₂AP), a major component of fibrinolytic system, causes an excess of plasmin, and that plasmin plays an important role in the formation of platelet aggregates. Consequently, the reduction of α₂AP could be a risk factor for the activation of platelets resulting in thrombus formation (9). The α₂AP activity was examined in this case to determine the effect of raloxifene treatment on the fibrinolytic system. The α₂AP activity gradually increased from the beginning of raloxifene-treatment, and it was sustained up to 4 weeks after the cessation of the medication (Fig. 4).

Discussion

This report presents a case of primary osteoporosis showing platelet hyper-aggregability to ADP, closely related to the administration of the standard dose of raloxifene. This is probably the first report clearly showing raloxifene-induced unusual platelet aggregation, which was detectable using the LS method. Interestingly, the hyper-sensitivity was correlated with the enhancement of Akt phosphorylation induced by ADP in platelets. The platelets play an important role in normal homeostasis and abnormal activation of platelets thus leading to thrombosis. During vascular injury or conditions of high shear, exposure of the collagen-rich subendothelium

Table 2. Laboratory Data

Parameters		normal range
Peripheral Blood		
WBC	4600/ μ L	3500-9200/ μ L
RBC	431 x 10 ⁴ / μ L	388-488 x 10 ⁴ / μ L
Hb	13.3 g/dL	11.3-15.5 g/dL
Hct	39.9%	34.4-45.6%
Plt	16.4 x 10 ⁴ / μ L	15.5-36.5 x 10 ⁴ / μ L
Blood Chemistry		
Total protein	7.7 g/dL	6.3-8.1 g/dL
Albumin	4.7 g/dL	3.7-4.9 g/dL
AST	24 IU/L	9-38 IU/L
ALT	33 IU/L	4-36 IU/L
ALP	361 IU/L	60-201 IU/L
Urinary nitrogen	13 mg/dL	9-21 mg/dL
Creatinine	0.5 mg/dL	0.4-0.9 mg/dL
Serum calcium	9.5 mg/dL	8.4-9.7 mg/dL
Serum iP	3.5 mg/dL	2.5-4.5 mg/dL
Endocrinology		
free T ₃	2.79 pg/mL	2.84-4.45 pg/mL
free T ₄	1.19 pg/mL	0.82-1.72 pg/mL
TSH	0.73 mIU/mL	0.49-3.83 mIU/mL
Cortisol	14 ng/mL	7-25 ng/mL
High sensitivity PTH	212 pg/mL	90-270 pg/mL
Bone Metabolic Markers		
BAP	46.4 IU/L	7.9-29.0 IU/L
Serum NTx	13.3 nmolBCE/L	10.7-24.0 nmolBCE/L

Footnote: RBC: Red blood cell, WBC: White blood cell, Hb: Hemoglobin, Hct: Hematocrit, Plt: Platelet, AST: L-asparatate aminotransferase, ALT: L-alanine aminotransferase, ALP: Alkaline phosphatase, free T₃: free tri-iodothyronine, free T₄: free thyroxin, TSH: thyroid-stimulating hormone, PTH: parathyroid hormone, BAP: bone type alkaline phosphatase, Serum NTx: Serum cross-linked N-telopeptides of I type collagen.

activates the platelets and results in the formation of a stable thrombus due to the combined action of ADP secreted from platelet-dense granules and generated thrombin (14, 17). The present results suggest that raloxifene could induce hyper-reactivity of platelet aggregation by increased sensitivity to ADP stimulation and by Akt activation. The hyper-sensitivity of platelets shown here might lead us to pay attention to the risk of arterial thrombosis, in addition to the increased hazard ratio of venous thromboembolism, under standard raloxifene treatment for osteoporosis.

Prior studies have shown that PI3K is a necessary component which plays a pivotal role in the signaling of purinore-

ceptor P2Y₁₂ in platelets (14, 18). A potential downstream effector for PI3K is Akt, which contributes to fibrinogen receptor activation and platelet aggregation. In addition, PI3K/Akt signaling is also reportedly involved in the release of ADP-containing granules (18). The platelet-aggregated form consists of two types of platelet membrane receptors, glycoprotein (GP) IIb/IIIa and GPIb/V/IX. GPIIb/IIIa not only binds fibrinogen and von Willebrand factor (vWF) to mediate platelet aggregation and adhesion, but it also serves as a signaling receptor. ADP induces the signaling, which activates the receptor function of GPIIb/IIIa for soluble fibrinogen, and ADP-induced platelet aggregation promotes the




Figure 1. The change in ED50 for ADP-induced platelet aggregation. PRP samples were obtained during the treatment and 4 weeks after the cessation of raloxifene administration. PRP was chronologically obtained from freshly drawn venous blood samples. The ED50 for ADP was determined.

binding of fibrinogen to GPIIb/IIIa (19). The P2Y12 receptor is involved in the constitution of stable macroaggregates (irreversible change) through full activation of the GPIIb/IIIa (13, 20). In the present case, raloxifene probably increased the platelet sensitivity to ADP and induced the enhancement of Akt activity in platelets, thus suggesting that the up-regulation in signaling events of P2Y12 receptor occurred under raloxifene administration. Blockade of P2Y12 receptor signaling might be a possible therapeutic strategy to avoid the adverse effects of this useful agent. The accumulation of more such cases and further investigations will thus be required to clarify the above mechanism of action in detail.

On the other hand, an increase of α 2AP activity was also observed in this case. Plasmin plays a crucial role in the formation of platelet aggregation. It is generally recognized that increase of α 2AP, which is produced by the liver, causes the reduction of plasmin, resulting in the inhibition of fibrinolytic system. Thus, it seems that an inhibition of the fibrinolytic system occurred in parallel with the increase of platelet aggregation under the treatment with raloxifene. As plasmin is known to play an important role in the formation of platelet aggregation, it is possible that the increase of α 2AP activity resulting in the reduction of plasmin in this case is a patho-physiological defensive mechanism against the raloxifene-induced platelet hyper-aggregability. In addition, the inhibition of fibrinolytic system might reduce the capacity of thrombolysis, resulting in the thrombus formation. In the present case, raloxifene treatment was continued for 6 months and the increased activity of α 2AP sustained after the cessation of raloxifene. It seems that the 4 weeks cessation of raloxifene after 6 months administration is not sufficient for the recovery of fibrinolytic system. The moni-




Figure 2. Platelet aggregability after the addition of ADP measured with the laser-light scattering method. Blue line indicates small aggregates; green line, medium aggregates; red line, large aggregates; and black line, optical density (%T). PRP samples were obtained before (A), after 2 weeks (B) and after 4 weeks (C) of raloxifene administration, and the ADP (1 μ M)-induced platelet aggregation was determined using a PA-200 aggregometer.

toring of α 2AP, in addition to the platelet microaggregates by a LS system, is probably a useful tool for the detection of the subclinical thrombus formation in the clinical use of raloxifene.

In conclusion, a standard dosage of raloxifene, one type




Figure 3. The change in ADP-induced platelet aggregation and the phosphorylation of Akt in the platelets under the treatment of raloxifene. Under the treatment of raloxifene and 4 weeks after the cessation, PRP sample were obtained at the indicated time. The platelet aggregation (%) induced by ADP (1 μM) was determined using a PA-200 aggregometer, and then with the same samples, the phosphorylation of Akt induced by ADP (1 μM) was examined at 24 weeks and 4 weeks after the cessation of the treatment. The platelet extracts were separated by SDS-PAGE and a Western blot analysis was performed with antibodies against either phospho-specific Akt or Akt.




Figure 4. The change in the α2-antiplasmin (α2AP) activity. Under the treatment of raloxifene and 4 weeks after the cessation, PRP samples were obtained at the indicated week. PRP was chronologically obtained from freshly drawn venous blood samples.

of SERM, was found to induce platelet hyper-aggregability during the conventional therapeutic management for post-menopausal osteoporosis. The clinicians using SERM must be aware of the acceleration of platelet aggregation in addition to that of coagulation which may occur during the ad-

ministration of this regimen.

Acknowledgement

This study was supported in part by Lilly Fellowship and Research Grant Program for Bone & Mineral in Japan.

References

1. Ettinger B, Black DM, Mitlak BH, et al. Reduction of vertebral fracture risk in postmenopausal women with osteoporosis treated with raloxifene: results from a 3-year randomized clinical trial. Multiple Outcomes of Raloxifene Evaluation (MORE) investigators. *JAMA* **282**: 637-645, 1999.
2. Peverill RE. Hormone therapy and venous thromboembolism. *Best Pract Res Clin Endocrinol Metab* **17**: 149-164, 2003.
3. Born GV, Hume M. Effects of the numbers and sizes of platelet aggregates on the optical density of plasma. *Nature* **215**: 1027-1029, 1967.
4. Cardinal DC, Flower RJ. The study of platelet aggregation in whole blood. *Br J Pharmacol* **66**: 94-95, 1979.
5. Ozaki Y, Satoh K, Yatomi Y, et al. Detection of platelet aggregates with a particle counting method using light scattering. *Anal Biochem* **218**: 284-294, 1994.
6. Tohgi H, Takahashi H, Watanabe K, Kuki H, Shirasawa Y. Development of large platelet aggregates from small aggregates as determined by laser-light scattering: effects of aggregant concentration and antiplatelet medication. *Thromb Haemost* **75**: 838-843, 1996.
7. Haque SF, Matubayashi H, Izumi S, et al. Sex difference in platelet aggregation detected by new aggregometry using light scattering. *Endocr J* **48**: 33-41, 2001.
8. Sakamoto T, Ogawa H, Kawano H, et al. Rapid change of platelet aggregability in acute hyperglycemia. Detection by a novel laser-light scattering method. *Thromb Haemost* **83**: 475-479, 2000.
9. Takei M, Matsuno H, Okada K, et al. Lack of alpha 2-antiplasmin enhances ADP induced platelet micro-aggregation through the presence of excess active plasmin in mice. *J Thromb Thrombolysis* **14**: 205-211, 2002.
10. Matsuno H, Tokuda H, Ishisaki A, et al. P2Y₁₂ receptors play a significant role in the development of platelet microaggregation in patients with diabetes. *J Clin Endocrinol Metab* **90**: 920-927, 2005.
11. Colman RW. Aggregin: a platelet ADP receptor that mediates activation. *FASEB J* **4**: 1425-1435, 1990.
12. Kowalska MA, Ratajczak MZ, Majka M, et al. Stromal cell-derived factor-1 and macrophage-derived chemokine: 2 factor-1 and macrophage-derived chemokine: 2 chemokines that activate platelets. *Blood* **96**: 50-57, 2000.
13. Fabre JE, Nguyen M, Latour A, et al. Decreased platelet aggregation, increased bleeding time and resistance to thromboembolism in P2Y₁-deficient mice. *Nat Med* **5**: 1199-1202, 1999.
14. Dorsam RT, Kunapuli SP. Central role of the P2Y₁₂ receptor in platelet aggregation. *J Clin Invest* **113**: 340-345, 2004.
15. Orimo H, Hayashi Y, Fukunaga M, et al. Osteoporosis diagnostic criteria review committee: Japanese Society for Bone and Mineral Research Diagnostic criteria for primary osteoporosis. Year 2000 version. *J Bone Miner Metab* **19**: 331-337, 2001.
16. Nakajima K, Hirade K, Kozawa O, et al. Akt regulates thrombin-induced HSP27 phosphorylation in aortic smooth muscle cells: Function at a point downstream from p38 MAP kinase. *Life Sci* **77**: 96-107, 2005.
17. Shanker H, Murugappan S, Kim S, et al. Role of G protein-gated inwardly rectifying potassium channels in P2Y₁₂ receptor-mediated platelet functional responses. *Blood* **104**: 1335-1343, 2004.
18. Woulfe D, Jiang H, Morgans A, et al. Defects in secretion, aggregation, and thrombus formation in platelets from mice lacking Akt2. *J Clin Invest* **113**: 441-450, 2004.
19. Naik UP, Naik MU. Association of CIB with GPIIb/IIIa during outside-in signaling is required for platelet spreading on fibrinogen. *Blood* **102**: 1335-1362, 2003.
20. Eckly A, Gendreau JL, Hechler B, et al. Differential involvement of the P2Y₁ and P2Y₇ receptors in the morphological changes of platelet aggregation. *Thromb Haemost* **85**: 694-701, 2001.



Contents lists available at ScienceDirect

Prostaglandins, Leukotrienes and Essential Fatty Acids

journal homepage: www.elsevier.com/locate/plefa

Function of Rho-kinase in prostaglandin D₂-induced interleukin-6 synthesis in osteoblasts

Haruhiko Tokuda^{a,b,*}, Shinji Takai^b, Rie Matsushima-Nishiwaki^b, Yoshiteru Hanai^{a,b}, Seiji Adachi^b, Chiho Minamitani^c, Jun Mizutani^c, Takanobu Otsuka^c, Osamu Kozawa^b

^a Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, 36-3 Gengo, Obu, Aichi 474-8511, Japan

^b Department of Pharmacology, Gifu University, Graduate School of Medicine, Gifu 501-1194, Japan

^c Department of Orthopedic Surgery, Nagoya City University Graduate School of Medical Sciences, Nagoya 467-8601, Japan

ARTICLE INFO

Article history:

Received 24 January 2008

Received in revised form

27 June 2008

Accepted 10 July 2008

ABSTRACT

We have previously reported that prostaglandin D₂ (PGD₂) stimulates interleukin-6 (IL-6), a potent bone resorptive agent, in osteoblast-like MC3T3-E1 cells. In the present study, we investigated whether Rho-kinase is implicated in the PGD₂-stimulated IL-6 synthesis in MC3T3-E1 cells. PGD₂ time-dependently induced the phosphorylation of myosin phosphatase targeting subunit (MYPT-1), a Rho-kinase substrate. Y27632, a specific Rho-kinase inhibitor, significantly reduced the PGD₂-stimulated IL-6 synthesis as well as the MYPT-1 phosphorylation. Fasudil, another inhibitor of Rho-kinase, suppressed the PGD₂-stimulated IL-6 synthesis. The PGD₂-stimulated IL-6 synthesis was reduced by PD98059, a MEK inhibitor, and SB203580, an inhibitor of p38 mitogen-activated protein (MAP) kinase, but not SP600125, an inhibitor of stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK). However, Y27632 and fasudil failed to affect the PGD₂-induced phosphorylation of p44/p42 MAP kinase. On the other hand, Y27632 as well as fasudil markedly attenuated the PGD₂-induced phosphorylation of p38 MAP kinase. In addition, PGD₂ additively induced IL-6 synthesis in combination with endothelin-1 which induces IL-6 synthesis through p38 MAP kinase regulated by Rho-kinase. These results strongly suggest that Rho-kinase regulates PGD₂-stimulated IL-6 synthesis via p38 MAP kinase activation in osteoblasts.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

It is well recognized that interleukin-6 (IL-6) is a multi-functional cytokine that has crucial effects on a wide range of functions such as promoting B-cell differentiation, T-cell activation and inducing acute phase proteins [1–3]. Bone homeostasis is maintained by the coordinated actions of osteoblast-mediated formation and osteoclast-mediated bone removal [4]. In the process of bone remodeling, IL-6 has been shown to stimulate bone resorption and promote osteoclast formation [2,3,5,6]. It has been reported that potent bone resorptive agents such as tumor-necrosis factor- α and interleukin-1 stimulate IL-6 synthesis in osteoblasts [5,7,8]. Currently, evidence is accumulating that IL-6 secreted from osteoblasts plays a pivotal role as a local downstream effector of bone resorptive agents in bone remodeling.

It is well known that prostaglandins act as local factors, autacoids, of osteoblasts [4,9]. Among them, prostaglandin D₂ (PGD₂) is recognized to be implicated in the control of osteoblast function and bone anabolism [9]. It has been reported that PGD₂ stimulates collagen synthesis during calcification of osteoblasts [10]. In addition, PGD₂ produced by osteoblasts reportedly modulates expression of osteoprotegerin and RANKL in osteoblasts [11]. In our previous study [12], we reported that PGD₂ stimulates IL-6 synthesis via Ca²⁺ mobilization in osteoblast-like MC3T3-E1 cells. However, the exact mechanism behind PGD₂-stimulated IL-6 synthesis in osteoblasts remains to be clarified.

Accumulating evidence suggests that Rho and the downstream effector, Rho-associated kinase (Rho-kinase) play crucial roles in a variety of cellular functions such as cell motility and smooth muscle contraction [13–15]. As for osteoblasts, it has been demonstrated that Rho and p38 mitogen-activated protein (MAP) kinase are involved in the endothelin-1 (ET-1)-induced expression of prostaglandin endoperoxide G/H synthase mRNA in osteoblasts [16]. In addition, it has been shown that the Rho/Rho-kinase pathway stimulates osteoblast proliferation whereas it inhibits osteoblast differentiation [17]. We have recently reported that Rho-kinase regulates ET-1-stimulated IL-6 synthesis through p38 MAP kinase

* Corresponding author at: Department of Clinical Laboratory, National Hospital for Geriatric Medicine, National Center for Geriatrics and Gerontology, 36-3 Gengo, Obu, Aichi 474-8511, Japan. Tel.: +81 562 46 2311; fax: +81 562 46 8396.
E-mail address: tokuda@ncgg.go.jp (H. Tokuda).

in osteoblast-like MC3T3-E1 cells [18]. However, the exact role of Rho-kinase in osteoblasts has not yet been fully clarified.

In the present study, we further investigated the exact mechanism behind PGD₂-stimulated IL-6 synthesis in osteoblast-like MC3T3-E1 cells. We here show that Rho-kinase regulates PGD₂-stimulated IL-6 synthesis through p38 MAP kinase activation in these cells.

2. Materials and methods

2.1. Materials

PGD₂, ET-1 and mouse IL-6 enzyme immunoassay (ELISA) kit were purchased from R&D Systems, Inc. (Minneapolis, MN). Y27632, an inhibitor of Rho-kinase [15], PD98059, an inhibitor of MEK [19], SB203580, an inhibitor of p38 MAP kinase [20], and SP600125, an inhibitor of stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) [21], were obtained from Calbiochem-Novabiochem Co. (La Jolla, CA). Hydroxyfasudil (fasudil), another inhibitor of Rho-kinase [15], was purchased from Sigma (St. Louis, MO). Phospho-specific myosin phosphatase targeting subunit (MYPT-1) antibodies were purchased from Upstate (Lake Placid, NY). MYPT-1 antibodies were obtained from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA). Phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies and p38 MAP kinase antibodies were purchased from Cell Signaling, Inc. (Beverly, MA). ECL Western blotting detection system was purchased from Amersham Biosciences (Piscataway, NJ). Other materials and chemicals were obtained from commercial sources. Y27632, PD98059, SB203580 and SP600125 were dissolved in dimethyl sulfoxide. The maximum concentration of dimethyl sulfoxide was 0.1%, which did not affect the assay for IL-6 or Western blot analysis.

2.2. Cell culture

Cloned osteoblast-like MC3T3-E1 cells derived from newborn mouse calvaria [22] were maintained as previously described [23]. Briefly, the cells were cultured in α -minimum essential medium (α -MEM) containing 10% fetal calf serum (FCS) at 37 °C in a humidified atmosphere of 5% CO₂/95% air. The cells were seeded into 35-mm or 90-mm diameter dishes in α -MEM containing 10% FCS. After 5 days, the medium was exchanged for α -MEM containing 0.3% FCS. The cells were used for experiments after 48 h.

2.3. IL-6 assay

The cultured cells were stimulated by 10 μ M PGD₂, 0.1 μ M ET-1, their combination or vehicle in 1 ml of α -MEM containing 0.3% FCS for the indicated periods. When indicated, the cells were pretreated with various doses of Y27632, fasudil, PD98059, SB203580 or SP600125 for 60 min. The conditioned medium was collected at the end of the incubation, and the IL-6 concentration was measured by ELISA kit.

2.4. Western blot analysis

The cultured cells were pretreated with various doses of Y27632 or fasudil for 60 min, and then stimulated by PGD₂ in α -MEM containing 0.3% FCS for the indicated periods. The cells were washed twice with phosphate-buffered saline and then lysed, homogenized and sonicated in a lysis buffer containing 62.5 mM Tris/HCl, pH 6.8, 2% sodium dodecyl sulfate (SDS), 50 mM

dithiothreitol and 10% glycerol. The cytosolic fraction was collected as a supernatant after centrifugation at 125,000g for 10 min at 4 °C. SDS-polyacrylamide gel electrophoresis (PAGE) was performed by Laemmli [24] in 10% polyacrylamide gel. Western blotting analysis was performed as described previously [25] by using phospho-specific MYPT-1 antibodies, MYPT-1 antibodies, phospho-specific p44/p42 MAP kinase antibodies, p44/p42 MAP kinase antibodies, phospho-specific p38 MAP kinase antibodies or p38 MAP kinase antibodies with peroxidase-labeled antibodies raised in goat against rabbit IgG being used as second antibodies. Peroxidase activity on the PVDF sheet was visualized on X-ray film by means of the ECL Western blotting detection system.

2.5. Determination

The absorbance of enzyme immunoassay samples was measured at 450 nm with EL 340 Bio Kinetic Reader (Bio-Tek Instruments, Inc., Winooski, VT). The densitometric analysis of the band on film was performed using Molecular Analyst/Macintosh (Bio-Rad Laboratories, Hercules, CA).

2.6. Statistical analysis

The data were analyzed by ANOVA followed by the Bonferroni method for multiple comparisons between pairs, and a $p < 0.05$ was considered significant. All data are presented as the mean \pm S.E.M. of triplicate independent experiments. Each experiment was repeated three times with similar results.

3. Results

3.1. Effect of PGD₂ on the phosphorylation of MYPT-1 in MC3T3-E1 cells

It is well accepted that MYPT-1, which is a component of myosin phosphatase, is a downstream substrate of Rho-kinase [13,26]. In order to clarify whether PGD₂ activates Rho-kinase in osteoblast-like MC3T3-E1 cells, we examined the effect of PGD₂ on the phosphorylation of MYPT-1. PGD₂ markedly induced the phosphorylation of MYPT-1 in a time-dependent manner (Fig. 1A). The effect of PGD₂ on the phosphorylation of MYPT-1 reached its peak within 2 min and decreased thereafter (Fig. 1A).

Y27632, a specific inhibitor of Rho-kinase [15], suppressed the PGD₂-induced phosphorylation levels of MYPT-1 (Fig. 1B). In addition, fasudil, another inhibitor of Rho-kinase [15], reduced the PGD₂-induced levels of phosphorylated MYPT-1 (data not shown).

3.2. Effects of Y27632 or fasudil on the PGD₂-stimulated IL-6 synthesis in MC3T3-E1 cells

We previously showed that PGD₂ stimulates IL-6 synthesis in osteoblast-like MC3T3-E1 cells [12]. In order to investigate whether Rho-kinase is implicated in the PGD₂-induced synthesis of IL-6 in MC3T3-E1 cells, we next examined the effect of Y27632 on the synthesis of IL-6 induced by PGD₂. Y27632, which by itself had little effect on the IL-6 levels, significantly suppressed the PGD₂-induced synthesis of IL-6 (Fig. 2A). The inhibitory effect of Y27632 was dose dependent in the range between 0.1 and 10 μ M. Y27632 (10 μ M) caused approximately 80% inhibition in the PGD₂-effect.

In addition, fasudil as well as Y27632, which by itself had little effect on IL-6 synthesis, reduced the PGD₂-stimulated IL-6 synthesis in these cells (Fig. 2B). The effect of fasudil on the IL-6 synthesis was dose dependent in the range between 0.1 and

10 μ M. Fasudil (10 μ M) caused about 80% suppression in the PGD₂-effect.

3.3. Effect of a combination of PGD₂ and ET-1 on IL-6 synthesis in MC3T3-E1 cells

We have recently reported that Rho-kinase regulates ET-1-stimulated IL-6 synthesis through p38 MAP kinase in osteoblast-

like MC3T3-E1 cells [18]. Therefore, we further examined the effect of a combination of PGD₂ and ET-1 on IL-6 synthesis in these cells. PGD₂ and ET-1 additionally induced IL-6 synthesis (Table 1).

3.4. Effects of PD98059, SB203580 or SP600125 on the PGD₂-stimulated IL-6 synthesis in MC3T3-E1 cells

It is currently known that three MAP kinases, p44/p42 MAP kinase, p38 MAP kinase and SAPK/JNK among the MAP kinase superfamily are known as central elements used by mammalian cells to transduce the various messages of extracellular stimuli [27]. We have previously reported that PGD₂ stimulates the activation of p44/p42 MAP kinase, p38 MAP kinase and SAPK/JNK in osteoblast-like MC3T3-E1 cells [28,29]. To investigate the involvement of three MAP kinases in the PGD₂-stimulated IL-6 synthesis in these cells, we examined the effect of PD98059, a potent MEK inhibitor [19], SB203580, a specific inhibitor of p38 MAP kinase [20], or SP600125, a specific inhibitor of SAPK/JNK [21], on the IL-6 synthesis. PD98059 and SB203580 significantly suppressed the PGD₂-stimulated synthesis of IL-6 (Table 2). On the contrary, SP600125, a specific inhibitor of SAPK/JNK [21], failed to suppress the IL-6 synthesis stimulated by PGD₂ (data not shown).

3.5. Effects of Y27632 or fasudil on the PGD₂-induced phosphorylation of p44/p42 MAP kinase in MC3T3-E1 cells

In order to elucidate whether Rho-kinase effect on the PGD₂-stimulated IL-6 synthesis is due to the activation of p44/p42 MAP kinase or not in MC3T3-E1 cells, we next examined the effect of Y27632 on the phosphorylation of p44/p42 MAP kinase by PGD₂. However, Y27632 hardly affected the PGD₂-induced phosphorylation of p44/p42 MAP kinase (Fig. 3A). Additionally, the phosphorylation of p44/p42 MAP kinase by PGD₂ was not reduced by fasudil (Fig. 3B).

3.6. Effects of Y27632 or fasudil on the PGD₂-induced phosphorylation of p38 MAP kinase in MC3T3-E1 cells

Furthermore, we examined the effect of Y27632 on the PGD₂-induced phosphorylation of p38 MAP kinase in MC3T3-E1 cells. Y27632 markedly suppressed the PGD₂-induced levels of p38 MAP kinase phosphorylation (Fig. 4A). The inhibitory effect of Y27632 on the phosphorylated levels was dose dependent in the range between 1 and 10 μ M. Y27632 (10 μ M) caused more than 90% inhibition in the PGD₂-effect (Fig. 4A).

Fasudil suppressed the PGD₂-induced levels of phosphorylated-p38 MAP kinase (Fig. 4B). The suppressive effect of fasudil as well as Y27632 was dose dependent in the range between 1 and 10 μ M. Fasudil (10 μ M) reduced the PGD₂-induced levels almost to the control levels (Fig. 4B).




Fig. 1. Effect of Y27632 on the PGD₂-induced phosphorylation of MYPT-1 in MC3T3-E1 cells. (A) The cultured cells were stimulated by 10 μ M PGD₂ for the indicated periods. (B) The cultured cells were pretreated with 10 μ M Y27632 or vehicle for 60 min, and then stimulated by 10 μ M PGD₂ or vehicle for 2 min. The extracts of cells were subjected to SDS-PAGE with subsequent Western blotting analysis with antibodies against phospho-specific MYPT-1 or MYPT-1. The histogram shows quantitative representations of the levels of PGD₂-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean \pm S.E.M. of triplicate independent experiments. Similar results were obtained with two additional and different cell preparations. * p < 0.05, compared with the control. # p < 0.05, compared with the value of PGD₂ alone.




Fig. 2. Effects of Y27632 or fasudil on the PGD₂-induced IL-6 synthesis in MC3T3-E1 cells. The cultured cells were pretreated with various doses of Y27632 (A) or fasudil (B) for 60 min, and then stimulated by 10 μM PGD₂ (●) or vehicle (○) for 24 h. Each value represents the mean ± S.E.M. of triplicate independent experiments. Similar results were obtained with two additional and different cell preparations. **p* < 0.05, compared to the value of PGD₂ alone.

Table 1
Effect of a combination of PGD₂ and ET-1 on IL-6 synthesis in MC3T3-E1 cells

ET-1	PGD ₂	IL-6 (pg/ml)
–	–	<7.8
–	+	427 ± 25*
+	–	296 ± 18*
+	+	991 ± 41**

The cultured cells were stimulated by 0.1 μM ET-1, 10 μM PGD₂, their combination or vehicle for 24 h. Each value represents the mean ± S.E.M. of triplicate independent experiments. Similar results were obtained with two additional and different cell preparations.

* *p* < 0.05, compared to the control.

** *p* < 0.05, compared to the value of PGD₂ alone.

Table 2
Effects of PD98059 or SB203580 on the PGD₂-stimulated IL-6 synthesis in MC3T3-E1 cells

Inhibitors	PGD ₂	IL-6 (pg/ml)
–	–	<7.8
–	+	447 ± 29*
PD98059	–	<7.8
PD98059	+	282 ± 16**
SB203580	–	<7.8
SB203580	+	78 ± 5**

The cultured cells were pretreated with 30 μM PD98059 or 30 μM SB203580 for 60 min, and then stimulated by 10 μM PGD₂ or vehicle for 24 h. Each value represents the mean ± S.E.M. of triplicate independent experiments. Similar results were obtained with two additional and different cell preparations.

* *p* < 0.05, compared to the control.

** *p* < 0.05, compared to the value of PGD₂ alone.

4. Discussion

In the present study, we demonstrated that PGD₂ time-dependently induced the phosphorylation of MYPT-1 in osteoblast-like MC3T3-E1 cells, using antibodies against phospho-MYPT-1 (Thr850). MYPT, which is a myosin-binding subunit of myosin phosphatase and regulates the interaction of actin and

myosin, is well known to be a downstream target of Rho-kinase [13,26]. In addition, we next found that Y27632 and fasudil, inhibitors of Rho-kinase [15], truly attenuated the PGD₂-induced phosphorylation of MYPT-1. Based on our results, it is most likely that PGD₂ induces the activation of Rho-kinase in osteoblast-like MC3T3-E1 cells.

We tried to clarify whether Rho-kinase functions in the PGD₂-stimulated IL-6 synthesis or not in osteoblast-like MC3T3-E1 cells. The PGD₂-stimulated synthesis of IL-6 was significantly suppressed by Y27632 [15] in a dose-dependent manner. This finding suggests that Rho-kinase activation is involved in the PGD₂-stimulated IL-6 synthesis in MC3T3-E1 cells. In addition, we demonstrated that fasudil [15] as well as Y27632 inhibited the IL-6 synthesis stimulated by PGD₂. Therefore, taking our findings into account, it is most likely that the PGD₂-activated Rho-kinase functions as a positive regulator in the IL-6 synthesis in osteoblast-like MC3T3-E1 cells.

Regarding about intracellular signaling system of PGD₂ in osteoblasts, we have previously reported that three MAP kinases, p44/p42 MAP kinase, p38 MAP kinase and SAPK/JNK among the MAP kinase superfamily are activated by PGD₂ in osteoblast-like MC3T3-E1 cells [28,29]. Therefore, we investigated the roles of three MAP kinases in the PGD₂-stimulated IL-6 synthesis in MC3T3-E1 cells. However, SP600125 [21] failed to reduce the PGD₂-stimulated IL-6 synthesis. Based on our finding, it seems unlikely that PGD₂ stimulates IL-6 synthesis through the activation of SAPK/JNK in osteoblast-like MC3T3-E1 cells. On the contrary, in the present study, we showed that the IL-6 synthesis induced by PGD₂ was suppressed by PD98059 [19] or SB203580 [20]. Thus, our results suggest that p44/p42 MAP kinase and p38 MAP kinase play a part in the PGD₂-stimulated IL-6 synthesis in these cells.

Furthermore, we tried to elucidate the relationship between Rho-kinase and these MAP kinase, p44/p42 MAP kinase and p38 MAP kinase in the PGD₂-stimulated IL-6 synthesis in MC3T3-E1 cells. However, Y27632 or fasudil did not affect the PGD₂-induced phosphorylation of p44/p42 MAP kinase. On the other hand, we demonstrated that the PGD₂-induced phosphorylation levels of p38 MAP kinase were markedly reduced by Y27632 and fasudil. Therefore, our findings suggest that Rho-kinase regulates the




Fig. 3. Effects of Y27632 or fasudil on the PGD₂-induced phosphorylation of p44/p42 MAP kinase in MC3T3-E1 cells. The cultured cells were pretreated with various doses of Y27632 (A) or fasudil (B) for 60 min, and then stimulated by 10 μM PGD₂ or vehicle for 20 min. The histogram shows quantitative representations of the levels of PGD₂-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean \pm S.E.M. of triplicate independent experiments. Similar results were obtained with two additional and different cell preparations.

PGD₂-stimulated IL-6 synthesis via p38 MAP kinase but not p44/p42 MAP kinase. With regard to the time-dependent effect of PGD₂, we showed that the maximum effect of PGD₂ on the




Fig. 4. Effects of Y27632 or fasudil on the PGD₂-induced phosphorylation of p38 MAP kinase in MC3T3-E1 cells. The cultured cells were pretreated with various doses of Y27632 (A) or fasudil (B) for 60 min, and then stimulated by 10 μM PGD₂ or vehicle for 10 min. The histogram shows quantitative representations of the levels of PGD₂-induced phosphorylation obtained from laser densitometric analysis of three independent experiments. Each value represents the mean \pm S.E.M. of triplicate independent experiments. Similar results were obtained with two additional and different cell preparations. * $p < 0.05$, compared with the control. # $p < 0.05$, compared with the value of PGD₂ alone.

phosphorylation of MYPT-1, a well-known downstream target of Rho-kinase [13], was observed within 2 min after the stimulation. In our previous study [29], we have reported that the

phosphorylation of p38 MAP kinase reaches the peak at 20 min after the stimulation of PGD₂ in these cells. The time course of the PGD₂-induced phosphorylation of MYPT-1 appears to be faster than that of p38 MAP kinase. Thus, it is reasonable that PGD₂-induced activation of p38 MAP kinase subsequently occurs via the activation of Rho-kinase. In addition, we showed that PGD₂ additively induced IL-6 synthesis in combination with ET-1 which induces IL-6 synthesis through p38 MAP kinase regulated by Rho-kinase [18]. It seems that Rho-kinase-mediated signaling induced from different pathways could enhance the downstream signaling in osteoblasts, resulting in the amplification of IL-6 synthesis. Based on our results as a whole, it is most likely that PGD₂-activated Rho-kinase functions at a point upstream from p38 MAP kinase in the IL-6 synthesis in osteoblast-like MC3T3-E1 cells.

It is well known that Rho-kinase plays an important role in a variety of cellular functions, especially vascular smooth muscle contraction [13–15]. In bone metabolism, Rho-kinase reportedly plays a suppressive role in the differentiation of osteoblasts while it induces their proliferation [17]. Bone formation is recognized to be performed by differentiated mature osteoblasts, suggesting that Rho-kinase activation regulates bone formation toward the suppression. Our present results indicate that PGD₂ induces Rho-kinase activation in osteoblasts, resulting in synthesizing IL-6, a potent modulator of bone metabolism. IL-6 produced by osteoblasts is generally recognized to be a bone resorptive agent and induces osteoclast differentiation [3,5]. Taking our present findings into account as a whole, it is probable that PGD₂ plays a regulator of bone metabolism toward up-regulating bone resorption and down-regulating bone formation through the activation of Rho-kinase. It has recently been reported that human osteoblasts produce PGD₂, which acts on the DP receptor to decrease osteoprotegerin production and on the CRTH2 receptor to decrease RANKL expression [11], suggesting that PGD₂ might elicit divergent effects on osteoclastogenesis. Further investigations are necessary to elucidate the exact role of Rho-kinase in osteoblasts.

In conclusion, our results strongly suggest that Rho-kinase functions as a positive regulator in the PGD₂-stimulated IL-6 synthesis in osteoblasts.

Acknowledgments

We are very grateful to Yoko Kawamura for her skillful technical assistance. This investigation was supported in part by Grant-in-Aid for Scientific Research (16590873 and 16591482) for the Ministry of Education, Science, Sports and Culture of Japan, the Research Grants for Longevity Sciences (15A-1 and 15C-2), Research Grant on Proteomics and Research Grant on Longevity Sciences from the Ministry of Health, Labour and Welfare of Japan.

References

[1] S. Akira, T. Taga, T. Kishimoto, Interleukin-6 in biology and medicine, *Adv. Immunol.* 54 (1993) 1–78.
 [2] D. Heymann, A.V. Rousselle, gp130 Cytokine family and bone cells, *Cytokine* 12 (2000) 1455–1468.
 [3] S. Kwan Tat, M. Padrine, S. Theoleyre, D. Heymann, Y. Fortin, IL-6 is produced by osteoblasts and induces bone resorption, *Cytokine Growth Factor Rev.* 15 (2004) 49–60.

[4] E.F. Morgan, G.L. Barnes, T.A. Einhorn, The bone organ system: form and function, in: R. Marcus, D. Feldman, D. Nelson, C.J. Rosen (Eds.), *Osteoporosis*, third ed., 2008, pp. 3–25.
 [5] Y. Ishimi, C. Miyaura, C.H. Jin, T. Akatsu, F. Abe, Y. Nakamura, Y. Yamaguchi, S. Yoshiki, T. Matsuda, T. Hirano, T. Kishimoto, T. Suda, IL-6 is produced by osteoblasts and induces bone resorption, *J. Immunol.* 145 (1990) 3297–3303.
 [6] G.D. Roodman, Interleukin-6: an osteotropic factor?, *J. Bone Miner. Res.* 7 (1992) 475–478.
 [7] A.J. Littlewood, J. Russell, G.R. Harvey, D.E. Hughes, R.G.G. Russel, M. Gowen, The modulation of the expression of IL-6 and its receptor in human osteoblasts in vitro, *Endocrinology* 129 (1991) 1513–1520.
 [8] M. Helle, J.P. Brakenhoff, E.R. DeGroot, L.A. Aarden, Interleukin 6 is involved in interleukin 1-induced activities, *Eur. J. Immunol.* 18 (1988) 957–959.
 [9] H. Hiki, T. Takato, T. Shimizu, S. Ishii, The roles of prostanooids, leukotrienes and platelet-activating factor in bone metabolism and disease, *Prog. Lipid Res.* 47 (2008) 107–126.
 [10] Y. Tasaki, R. Takamori, Y. Koshihara, Prostaglandin D₂ metabolite stimulates collagen synthesis by human osteoblasts during calcification, *Prostaglandins* 41 (1991) 303–313.
 [11] M.A. Gallant, R. Samadifard, J.A. Hackett, J. Antoniou, J.L. Parent, A.J. de Brum-Fernandes, Production of prostaglandin D₂ by human osteoblasts and modulation of osteoprotegerin, RANKL and cellular migration by DP and CRTH2 receptors, *J. Bone Miner. Res.* 20 (2005) 672–681.
 [12] H. Tokuda, O. Kozawa, A. Harada, T. Uematsu, Prostaglandin D₂ induces interleukin-6 synthesis via Ca²⁺ mobilization in osteoblasts: regulation of protein kinase C, *Prostaglandins Leukot. Essent. Fatty Acids* 61 (1999) 189–194.
 [13] Y. Fukata, M. Amano, K. Kaibuchi, Rho-Rho-kinase pathway in smooth muscle contraction and cytoskeletal reorganization of non-muscle cells, *Trends Pharmacol. Sci.* 22 (2001) 32–39.
 [14] K. Riento, A.J. Ridley, Rho GTPases: multifunctional kinases in cell behaviour, *Nat. Rev. Mol. Cell Biol.* 4 (2003) 446–456.
 [15] H. Shimokawa, M. Rashid, Development of Rho-kinase inhibitors for cardiovascular medicine, *Trends Pharmacol. Sci.* 28 (2007) 296–302.
 [16] W. Windischhofer, D. Zach, C. Fauler, G. Rasputnik, H. Kofeler, H.J. Leis, Involvement of Rho and p38 MAPK in endothelin-1-induced expression of PGHS-2 mRNA in osteoblast-like cells, *J. Bone Miner. Res.* 17 (2002) 1774–1784.
 [17] D. Harmey, G. Stenbeck, C.D. Nobes, A.J. Lax, A.E. Grigoriadis, Regulation of osteoblast differentiation by *Pasteurella multocida* toxin (PMT): a role for Rho GTPase in bone formation, *J. Bone Miner. Res.* 19 (2004) 661–670.
 [18] H. Tokuda, Y. Hanai, R. Matsushima-Nishiwaki, J. Yamauchi, T. Doi, A. Harada, S. Takai, O. Kozawa, Rho-kinase regulates endothelin-1-stimulated IL-6 synthesis via p38 MAP kinase in osteoblasts, *Biochem. Biophys. Res. Commun.* 362 (2007) 799–804.
 [19] D.R. Alessi, A. Cuenda, P. Cohen, D.T. Dudley, A.R. Saitli, PD98059 is a specific inhibitor of the activation of mitogen-activated protein kinase in vitro and in vivo, *J. Biol. Chem.* 270 (1995) 27489–27494.
 [20] A. Cuenda, J. Rouse, Y.N. Doza, R. Meier, P. Cohen, T.F. Gallagher, P.R. Young, J.C. Lee, SB203580 is a specific inhibitor of a MAP kinase homologue which is stimulated by cellular stresses and interleukin-1, *FEBS Lett.* 364 (1995) 229–233.
 [21] B.L. Bennett, D.T. Sasaki, B.W. Murray, E.C. O'Leary, S.T. Sakata, W. Xu, J.C. Leisten, A. Motiwala, S. Pierce, Y. Satoh, S.S. Bhagwat, A.M. Manning, D.W. Anderson, SP600125, an anthracycline inhibitor of Jun N-terminal Kinase, *Proc. Natl. Acad. Sci. USA* 98 (2001) 13681–13686.
 [22] H. Sudo, H. Kodama, Y. Amagai, S. Yamamoto, S. Kasai, In vitro differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria, *J. Cell Biol.* 96 (1993) 191–198.
 [23] O. Kozawa, H. Tokuda, M. Miwa, J. Kotyynri, Y. Oiso, Cross-talk regulation between cyclic AMP production and phosphoinositide hydrolysis induced by prostaglandin E₂ in osteoblast-like cells, *Exp. Cell Res.* 198 (1992) 130–134.
 [24] U.K. Laemmli, Cleavage of structural proteins during the assembly of the head of bacteriophage T4, *Nature* 227 (1970) 680–685.
 [25] K. Kato, H. Ito, K. Hasegawa, Y. Inaguma, O. Kozawa, T. Asano, Modulation of the stress-induced synthesis of hsp27 and alpha B-crystallin by cyclic AMP in C6 rat glioma cells, *J. Neurochem.* 66 (1996) 946–950.
 [26] M. Ito, T. Nakano, F. Erdodi, D.J. Hartshorne, Myosin phosphatase: structure, regulation and function, *Mol. Cell. Biochem.* 259 (2004) 197–209.
 [27] C. Widmann, S. Gibson, M.B. Jarpe, G.L. Johnson, Mitogen-activated protein kinase: conservation of a three-kinase module from yeast to human, *Physiol. Rev.* 79 (1999) 143–180.
 [28] O. Kozawa, T. Otsuka, D. Hatakeyama, M. Niwa, H. Matsuno, H. Ito, K. Kato, N. Matsui, T. Uematsu, Mechanism of prostaglandin D₂-stimulated heat shock protein 27 induction in osteoblasts, *Cell. Signal.* 13 (2001) 535–541.
 [29] M. Yoshida, M. Niwa, A. Ishisaki, K. Hirade, H. Ito, K. Shimizu, K. Kato, O. Kozawa, Methotrexate enhances prostaglandin D₂-stimulated heat shock protein 27 induction in osteoblasts, *Prostaglandins Leukot. Essent. Fatty Acids* 71 (2004) 351–362.