

- Harding AE, Thomas PK, Baraitser M, Bradbury PG, Morgan-Hughes JA, Ponsford JR. X-linked recessive bulbospinal neuronopathy: a report of ten cases. *J Neurol Neurosurg Psychiatry* 1982; 45: 1012–9.
- Ikeuchi T, Koide R, Tanaka H, Onodera O, Igarashi S, Takahashi H, et al. Dentatorubral-pallidolusian atrophy: clinical features are closely related to unstable expansions of trinucleotide (CAG) repeat. *Ann Neurol* 1995; 37: 769–75.
- Johansson J, Forsgren L, Sandgren O, Brice A, Holmgren G, Holmberg M. Expanded CAG repeats in Swedish spinocerebellar ataxia type 7 (SCA7) patients: effect of CAG repeat length on the clinical manifestation. *Hum Mol Genet* 1998; 7: 171–6.
- Kachi T, Sobue G, Sobue I. Central motor and sensory conduction in X-linked recessive bulbospinal neuronopathy. *J Neurol Neurosurg Psychiatry* 1992; 55: 394–7.
- Katsuno M, Adachi H, Kume A, Li M, Nakagomi Y, Niwa H, et al. Testosterone reduction prevents phenotypic expression in a transgenic mouse model of spinal and bulbar muscular atrophy. *Neuron* 2002; 35: 843–54.
- Katsuno M, Adachi H, Doyu M, Minamiyama M, Sang C, Kobayashi Y, et al. Leuprorelin rescues polyglutamine-dependent phenotypes in a transgenic mouse model of spinal and bulbar muscular atrophy. *Nat Med* 2003; 9: 768–73.
- Katsuno M, Sang C, Adachi H, Minamiyama M, Waza M, Tanaka F, et al. Pharmacological induction of heat-shock proteins alleviates polyglutamine-mediated motor neuron disease. *Proc Natl Acad Sci USA* 2005; 102: 16801–6.
- Katsuno M, Adachi H, Minamiyama M, Waza M, Tokui K, Banno H, et al. Pathogenesis, animal models and therapeutics in spinal and bulbar muscular atrophy (SBMA) [Review]. *Exp Neurol* 2006; 200: 8–18.
- Kennedy WR, Alter M, Sung JH. Progressive proximal spinal and bulbar muscular atrophy of late onset. A sex-linked recessive trait. *Neurology* 1968; 18: 671–680.
- Kimura J. Principles and variations of nerve conduction studies. Electrodiagnosis in diseases of nerve and muscle: principles and practice. 3rd edn. New York: Oxford University Press; 2001a. p. 91–129.
- Kimura J. Assessment of individual nerves. In: Kimura J, editor. Electrodiagnosis in diseases of nerve and muscle: principles and practice. 3rd edn. New York: Oxford University Press; 2001b. p. 130–77.
- Kimura J. The F wave and the A wave. In: Kimura J, editor. Electrodiagnosis in diseases of nerve and muscle: principles and practice. 3rd edn. New York: Oxford University Press; 2001c. p. 439–65.
- Kimura J. The F wave and the A wave. In: Kimura J, editor. Electrodiagnosis in diseases of nerve and muscle: principles and practice. 3rd edn. New York: Oxford University Press; 2001d. p. 307–38.
- Koike H, Mori K, Misu K, Hattori N, Ito H, Hirayama M, et al. Painful alcoholic polyneuropathy with predominant small-fiber loss and normal thiamine status. *Neurology* 2001; 56: 1727–32.
- Koike H, Iijima M, Sugiura M, Mori K, Hattori N, Ito H, et al. Alcoholic neuropathy is clinicopathologically distinct from thiamine-deficiency neuropathy. *Ann Neurol* 2003; 54: 19–29.
- La Spada AR, Wilson EM, Lubahn DB, Harding AE, Fischbeck KH. Androgen receptor gene mutations in X-linked spinal and bulbar muscular atrophy. *Nature* 1991; 352: 77–9.
- Li M, Sobue G, Doyu M, Mukai E, Hashizume Y, Mitsuma T. Primary sensory neurons in X-linked recessive bulbospinal neuropathy: histopathology and androgen receptor gene expression. *Muscle Nerve* 1995; 18: 301–8.
- Li M, Miwa S, Kobayashi Y, Merry DE, Yamamoto M, Tanaka F, et al. Nuclear inclusions of the androgen receptor protein in spinal and bulbar muscular atrophy. *Ann Neurol* 1998; 44: 249–54.
- Mahant N, McCusker EA, Byth K, Graham S; Huntington Study Group. Huntington's disease: clinical correlates of disability and progression. *Neurology* 2003; 61: 1085–92.
- Minamiyama M, Katsuno M, Adachi H, Waza M, Sang C, Kobayashi Y, et al. Sodium butyrate ameliorates phenotypic expression in a transgenic mouse model of spinal and bulbar muscular atrophy. *Hum Mol Genet* 2004; 13: 1183–92.
- Mori K, Iijima M, Koike H, Hattori N, Tanaka F, Watanabe H, et al. The wide spectrum of clinical manifestations in Sjogren's syndrome-associated neuropathy. *Brain* 2005; 128: 2518–34.
- Norris FH, Calanchini PR, Fallat RJ, Panchari S, Jewett B. The administration of guanidine in amyotrophic lateral sclerosis. *Neurology* 1974; 24: 721–8.
- Olney RK, Aminoff MJ, So YT. Clinical and electrodiagnostic features of X-linked recessive bulbospinal neuronopathy. *Neurology* 1991; 41: 823–8.
- Perutz MF, Pope BJ, Owen D, Wanker EE, Scherzinger E. Aggregation of proteins with expanded glutamine and alanine repeats of the glutamine-rich and asparagine-rich domains of Sup35 and of the amyloid beta-peptide of amyloid plaques. *Proc Natl Acad Sci USA* 2002; 99: 5596–600.
- Polo A, Teatini F, D'Anna S, Manganotti P, Salvati A, Dallapiccola B, et al. Sensory involvement in X-linked spino-bulbar muscular atrophy (Kennedy's syndrome): an electrophysiological study. *J Neurol* 1996; 243: 388–92.
- Sakahira H, Breuer P, Hayer-Hartl MK, Hartl FU. Molecular chaperones as modulators of polyglutamine protein aggregation and toxicity. [Review]. *Proc Natl Acad Sci USA* 2002; 99 (Suppl 4): 16412–8.
- Sobue G, Hashizume Y, Mukai E, Hirayama M, Mitsuma T, Takahashi A. X-linked recessive bulbospinal neuronopathy. A clinicopathological study. *Brain* 1989; 112: 209–32.
- Sobue G, Doyu M, Kachi T, Yasuda T, Mukai E, Kumagai T, et al. Subclinical phenotypic expressions in heterozygous females of X-linked recessive bulbospinal neuronopathy. *J Neurol Sci* 1993; 117: 74–8.
- Sone J, Hishikawa N, Koike H, Hattori N, Hirayama M, Nagamatsu M, et al. Neuronal intranuclear hyaline inclusion disease showing motor-sensory and autonomic neuropathy. *Neurology* 2005; 65: 1538–43.
- Sperfeld AD, Karitzky J, Brummer D, Schreiber H, Haussler J, Ludolph AC, et al. X-linked bulbospinal neuronopathy: Kennedy disease. *Arch Neurol* 2002; 59: 1921–6.
- Takeyama K, Ito S, Yamamoto A, Tanimoto H, Furutani T, Kanuka H, et al. Androgen-dependent neurodegeneration by polyglutamine-expanded human androgen receptor in *Drosophila*. *Neuron* 2002; 35: 855–64.
- Tanaka F, Doyu M, Ito Y, Matsumoto M, Mitsuma T, Abe K, et al. Founder effect in spinal and bulbar muscular atrophy (SBMA). *Hum Mol Genet* 1996; 5: 1253–7.
- Tanaka F, Reeves MF, Ito Y, Matsumoto M, Li M, Miwa S, et al. Tissue-specific somatic mosaicism in spinal and bulbar muscular atrophy is dependent on CAG-repeat length and androgen receptor-gene expression level. *Am J Hum Genet* 1999; 65: 966–73.
- The ALS CNTF Treatment Study (ACTS) Phase I-II Study Group. The amyotrophic lateral sclerosis functional rating scale: Assessment of activities of daily living in patients with ALS. *Arch Neurol* 1996; 53: 141–7.
- Waza M, Adachi H, Katsuno M, Minamiyama M, Sang C, Tanaka F, et al. 17-AAG, an Hsp90 inhibitor, ameliorates polyglutamine-mediated motor neuron degeneration. *Nat Med* 2005; 11: 1088–95.
- Zoghbi HY, Orr HT. Glutamine repeats and neurodegeneration [Review]. *Annu Rev Neurosci* 2000; 23: 217–47.