


Fig. 3



Effects of the genetic marker of *UGT1A1* *6 or *28' on the area under concentration curve (AUC) ratios of SN-38G/SN-38 (a) and SN-38/irinotecan (b), and SN-38 by irinotecan dosage (c) in 176 Japanese cancer patients after irinotecan treatment.

Table 5 Association of *UGT1A1**6 and *28 with irinotecan toxicities

Diplotype (+ = *6 or *28)	Number of patients	Diarrhea (grade 3)	Neutropenia (grade 3 or 4)
Irinotecan monotherapy			
-/-	21	3 (14.3%) ^a	3 (14.3%)
+/-	29	2 (6.90%)	7 (24.1%)
+/+	5	1 (20.0%)	4 (80.0%)
		<i>P</i> -value ^b	0.0117
			0.0124
With cisplatin			
-/-	35	1 (2.9%)	20 (57.1%)
+/-	20	2 (10.0%)	14 (70.0%)
+/+	7	1 (14.3%)	7 (100%)
		<i>P</i> -value ^b	0.0315
			0.0883

^aPercentage of the patient number in each diplotype is indicated in parentheses.

^bChi-squared test for trend.

^cFisher's exact test, (-/- and +/-) vs. +/+.

of diarrhea in the irinotecan/capecitabine regimen, in which diarrhea was a major toxicity [20]. A highly frequent allele *IA9**22 with an insertion of T into the nine T repeats in the promoter region (-126_-118T₉ > T₁₀) was shown to have an enhanced promoter activity in an in-vitro reporter assay [21], whereas *IA9* protein expression levels did not change in the clinical samples [22]. Rare variations, *IA9**5 [766G > A(D256N)] and *UGT1A10**3 [605C > T(T2021)], were shown to cause reduced activity *in vitro*, but their clinical importance is still unknown [23,24]. Moreover, close linkages among *IA9*, *IA7*, and *IA1*

Table 6 Multiple regression analysis of the nadir of absolute neutrophil counts in the patients with irinotecan monotherapy

Variable	Coefficient	F-value	<i>P</i> -value	R ²	Intercept	N
Serum ALP ^a	-349.9	12.2	0.0010	0.3942	643	53
Neutrophil count before irinotecan treatment	0.2466	13.5	0.0006			
*6 or *28	-369.1	6.40	0.0146			

^aGrade 1 or greater scores of serum ALP before irinotecan treatment.

polymorphisms were found in Caucasians and Asians in an ethnic-specific manner [20,25-28].

Our study also revealed close linkages between *IA9**22 and *IA7**1, *IA7**3 and *IA1**6 or *28 [28]. This fact makes it difficult to draw firm conclusions about the effects of *IA7**3 and *IA9**22 themselves. It is, however, reasonable to conclude that the degree of neutropenia depends on the activity of *UGT1A1*, because *UGT1A1* is a major *UGT1A* enzyme in the liver and plays a primary role for regulating plasma concentrations of SN-38.

Taken together, for practical application to individualized irinotecan therapy, genotyping of *UGT1A1**6 and *28 would be beneficial and necessary in Japanese cancer patients to avoid severe adverse reactions. The frequency

of homozygotes for *6 or *28 (namely, *6/*6, *6/*28, and *28/*28) is approximately 10%, which is comparable to the frequency of *28 homozygotes in Caucasian populations. In our study, it may be difficult to establish definite guidelines for dose reductions of irinotecan for patients homozygous for *6 or *28. Considering, however, 2.4-fold steep relationship between the dose of irinotecan and the AUC of SN-38 for patients homozygous for *6 or *28 compared with patients without *6 or *28 (Fig. 3c), the dose for patients homozygous for *6 or *28 should be reduced to a half of the dosage recommended for other patients. Prospective studies are necessary to confirm the validity of the recommendation for dose reduction in Japanese cancer patients homozygous for *6 or *28.

Acknowledgements

We thank Ms Chie Sudo for her secretarial assistance. This study was supported in part by the Program for the Promotion of Fundamental Studies in Health Sciences and by the Program for the Promotion of Studies in Health Sciences of the Ministry of Health, Labor and Welfare of Japan. Analytical standards of irinotecan and its metabolites were kindly supplied by Yakult Honsha Co. Ltd. (Tokyo, Japan).

References

- Garcia-Carbonero R, Supko JG. Current perspectives on the clinical experience, pharmacology, and continued development of the camptothecins. *Clin Cancer Res* 2002; 8:641-661.
- Slatter JG, Su P, Sams JP, Schaaf LJ, Wieners LC. Bioactivation of the anticancer agent CPT-11 to SN-38 by human hepatic microsomal carboxylesterases and the in vitro assessment of potential drug interactions. *Drug Metab Dispos* 1997; 25:1157-1164.
- Iyer L, King CD, Whittington PF, Green MD, Roy SK, Tephly TR, et al. Genetic predisposition to the metabolism of irinotecan (CPT-11). Role of uridine diphosphate glucuronosyltransferase isoform 1A1 in the glucuronidation of its active metabolite (SN-38) in human liver microsomes. *J Clin Invest* 1998; 101:847-854.
- De Forni M, Bugat R, Chabot GG, Culine S, Extra JM, Gouyette A, et al. Phase I and pharmacokinetic study of the camptothecin derivative irinotecan, administered on a weekly schedule in cancer patients. *Cancer Res* 1994; 54:4347-4354.
- Gupta E, Lestingi TM, Mick R, Ramirez J, Vokes EE, Ratain MJ. Metabolic fate of irinotecan in humans: correlation of glucuronidation with diarrhea. *Cancer Res* 1994; 54:3723-3725.
- Hanioka N, Ozawa S, Jinno H, Ando M, Saito Y, Sawada J. Human liver UDP-glucuronosyltransferase isoforms involved in the glucuronidation of 7-ethyl-10-hydroxycamptothecin. *Xenobiotica* 2001; 31:687-699.
- Fisher MB, VandenBranden M, Findlay K, Burchell B, Thummel KE, Hall SD, et al. Tissue distribution and interindividual variation in human UDP-glucuronosyltransferase activity: relationship between UGT1A1 promoter genotype and variability in a liver bank. *Pharmacogenetics* 2000; 10:727-739.
- Ando Y, Saka H, Ando M, Sawa T, Muro K, Ueoka H, et al. Polymorphisms of UDP-glucuronosyltransferase gene and irinotecan toxicity: a pharmacogenetic analysis. *Cancer Res* 2000; 60:6921-6926.
- Iyer L, Das S, Janisch L, Wen M, Ramirez J, Karrison T, et al. UGT1A1*28 polymorphism as a determinant of irinotecan disposition and toxicity. *Pharmacogenetics* J 2002; 2:43-47.
- Innocenti F, Undevia SD, Iyer L, Chen PX, Das S, Kocherginsky M, et al. Genetic variants in the UDP-glucuronosyltransferase 1A1 gene predict the risk of severe neutropenia of irinotecan. *J Clin Oncol* 2004; 22:1382-1388.
- Marcuello E, Altes A, Menoyo A, del Rio E, Gomez-Pardo M, Baiget M. UGT1A1 gene variations and irinotecan treatment in patients with metastatic colorectal cancer. *Br J Cancer* 2004; 91:678-682.
- Rouits E, Boisdron-Celle M, Dumont A, Guerin O, Morel A, Gamelin E. Relevance of different UGT1A1 polymorphisms in irinotecan-induced toxicity: a molecular and clinical study of 75 patients. *Clin Cancer Res* 2004; 10:5151-5159.
- Kaniwa N, Kurose K, Jinno H, Tanaka-Kagawa T, Saito Y, Saeki M, et al. Racial variability in haplotype frequencies of UGT1A1 and glucuronidation activity of a novel single nucleotide polymorphism 686C>T (P229L) found in an African-American. *Drug Metab Dispos* 2005; 33:458-465.
- Sai K, Saeki M, Saito Y, Ozawa S, Katori N, Jinno H, et al. UGT1A1 haplotypes associated with reduced glucuronidation and increased serum bilirubin in irinotecan-administered Japanese patients with cancer. *Clin Pharmacol Ther* 2004; 75:501-515.
- Ciotti M, Basu N, Brangi M, Owens IS. Glucuronidation of 7-ethyl-10-hydroxycamptothecin (SN-38) by the human UDP-glucuronosyltransferases encoded at the UGT1 locus. *Biochem Biophys Res Commun* 1999; 260:199-202.
- Gagne JF, Montminy V, Belanger P, Journault K, Gaucher G, Guillemette C. Common human UGT1A polymorphisms and the altered metabolism of irinotecan active metabolite 7-ethyl-10-hydroxycamptothecin (SN-38). *Mol Pharmacol* 2002; 62:608-617.
- Oguri T, Takahashi T, Miyazaki M, Isobe T, Kohno N, Mackenzie PL. UGT1A10 is responsible for SN-38 glucuronidation and its expression in human lung cancers. *Anticancer Res* 2004; 24:2893-2896.
- Basu NK, Ciotti M, Hwang MS, Kole L, Mitra PS, Cho JW, et al. Differential and special properties of the major human UGT1-encoded gastrointestinal UDP-glucuronosyltransferases enhance potential to control chemical uptake. *J Biol Chem* 2004; 279:1429-1441.
- Ando M, Ando Y, Sekido Y, Ando M, Shimokata K, Hasegawa Y. Genetic polymorphisms of the UDP-glucuronosyltransferase 1A7 gene and irinotecan toxicity in Japanese cancer patients. *Jpn J Cancer Res* 2002; 93:591-597.
- Carlini LE, Meropol NJ, Bever J, Andria ML, Hill T, Gold P, et al. UGT1A7 and UGT1A9 polymorphisms predict response and toxicity in colorectal cancer patients treated with capecitabine/irinotecan. *Clin Cancer Res* 2005; 11:1226-1236.
- Yamanaka H, Nakajima M, Katoh M, Hara Y, Tachibana O, Yamashita J, et al. A novel polymorphism in the promoter region of human UGT1A9 gene (UGT1A9*22) and its effects on the transcriptional activity. *Pharmacogenetics* 2004; 14:329-332.
- Girard H, Court MH, Bernard O, Fortier LS, Villeneuve L, Hao Q, et al. Identification of common polymorphisms in the promoter of the UGT1A9 gene: evidence that UGT1A9 protein and activity levels are strongly genetically controlled in the liver. *Pharmacogenetics* 2004; 14:501-515.
- Jinno H, Saeki M, Saito Y, Tanaka-Kagawa T, Hanioka N, Sai K, et al. Functional characterization of human UDP-glucuronosyltransferase 1A9 variant, D256N, found in Japanese cancer patients. *J Pharmacol Exp Ther* 2003; 306:688-693.
- Jinno H, Saeki M, Tanaka-Kagawa T, Hanioka N, Saito Y, Ozawa S, et al. Functional characterization of wild-type and variant (T202I and M59I) human UDP-glucuronosyltransferase 1A10. *Drug Metab Dispos* 2003; 31:528-532.
- Kohle C, Mohrle B, Munzel PA, Schwab M, Wernet D, Badary OA, et al. Frequent co-occurrence of the TATA box mutation associated with Gilbert's syndrome (UGT1A1*28) with other polymorphisms of the UDP-glucuronosyltransferase-1 locus (UGT1A6*2 and UGT1A7*3) in Caucasians and Egyptians. *Biochem Pharmacol* 2003; 65: 1521-1527.
- Huang MJ, Yang SS, Lin MS, Huang CS. Polymorphisms of uridine-diphosphoglucuronosyltransferase 1A7 gene in Taiwan Chinese. *World J Gastroenterol* 2005; 11:797-802.
- Innocenti F, Liu W, Chen P, Dedai AA, Das S, Ratain MJ. Haplotypes of variants in the UDP-glucuronosyltransferase 1A9 and 1A1 genes. *Pharmacogenet Genomics* 2005; 15:295-301.
- Saeki M, Saito Y, Jinno H, Sai K, Ozawa S, Kurose K, et al. Haplotype structures of the UGT1A gene complex in a Japanese population. *Pharmacogenomics J* 2006; 6:63-75.
- Sai K, Kaniwa N, Ozawa S, Sawada J. An analytical method for irinotecan (CPT-11) and its metabolites using a high-performance liquid chromatography: parallel detection with fluorescence and mass spectrometry. *Biomed Chromatogr* 2002; 16:209-218.
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Rpy Stat Soc B* 1995; 57:289-300.
- Han JY, Lim HS, Shin ES, Yoo YK, Park YH, Lee JE, et al. Comprehensive analysis of UGT1A polymorphisms predictive for pharmacokinetics and treatment outcome in patients with non-small-cell lung cancer treated with irinotecan and cisplatin. *J Clin Oncol* 2006; 24:2237-2244.
- Kitagawa C, Ando M, Ando Y, Sekido Y, Wakai K, Imaizumi K, et al. Genetic polymorphism in the phenobarbital-responsive enhancer module of the UDP-glucuronosyltransferase 1A1 gene and irinotecan toxicity. *Pharmacogenet Genomics* 2005; 15:35-41.



Randomised phase III trial of carboplatin plus etoposide vs split doses of cisplatin plus etoposide in elderly or poor-risk patients with extensive disease small-cell lung cancer: JCOG 9702

H Okamoto^{*,1}, K Watanabe¹, H Kunikane¹, A Yokoyama², S Kudoh³, T Asakawa⁴, T Shibata⁴, H Kunitoh⁵, T Tamura⁵ and N Saijo⁶, on Behalf of the Japan Clinical Oncology Group (JCOG)-Lung Cancer Study Group

¹Department of Respiratory Medicine, Yokohama Municipal Citizen's Hospital, 56 Okazawa-cho, Hodogaya-ku, Yokohama, Kanagawa 240-8555, Japan; ²Niigata Cancer Center Hospital, Niigata-city, Japan; ³Osaka City University Medical School, Osaka-city, Japan; ⁴National Cancer Center, Tokyo, Japan; ⁵National Cancer Center Hospital, Tokyo, Japan; ⁶National Cancer Center East Hospital, Kashiwa, Japan

We compared the efficacy and the safety of a carboplatin plus etoposide regimen (CE) vs split doses of cisplatin plus etoposide (SPE) in elderly or poor-risk patients with extensive disease small-cell lung cancer (ED-SCLC). Eligibility criteria included: untreated ED-SCLC; age ≥ 70 and performance status 0–2, or age < 70 and PS 3. The CE arm received carboplatin area under the curve of five intravenously (IV) on day 1 and etoposide 80 mg m⁻² IV on days 1–3. The SPE arm received cisplatin 25 mg m⁻² IV on days 1–3 and etoposide 80 mg m⁻² IV on days 1–3. Both regimens were given with granulocyte colony-stimulating factor support in a 21–28 day cycle for four courses. A total of 220 patients were randomised. Median age was 74 years and 74% had a PS of 0 or 1. Major grade 3–4 toxicities were (%CE/%SPE): leucopenia 54/51, neutropenia 95/90, thrombocytopenia 56/16, infection 7/6. There was no significant difference (CE/SPE) in the response rate (73/73%) and overall survival (median 10.6/9.9 mo; $P = 0.54$). Palliation scores were very similar between the arms. Although the SPE regimen is still considered to be the standard treatment in elderly or poor-risk patients with ED-SCLC, the CE regimen can be an alternative for this population considering the risk–benefit balance.

British Journal of Cancer (2007) 97, 162–169. doi:10.1038/sj.bjc.6603810 www.bjancer.com

Published online 19 June 2007

© 2007 Cancer Research UK

Keywords: small-cell lung cancer; carboplatin; cisplatin; etoposide; elderly; poor-risk

Approximately half of patients with small-cell lung cancer (SCLC) are older than 70 years, and the proportion of elderly SCLC patients is continuously increasing in Japan (Morita, 2002). However, since many investigators have arbitrarily excluded elderly patients from clinical trials, no standard chemotherapeutic regimen has been established for elderly patients with SCLC. The Japan Clinical Oncology Group (JCOG) has reported that carboplatin plus etoposide (CE) is an active and less toxic regimen in elderly patients with SCLC (Okamoto *et al*, 1999). However, other clinical trials have indicated that the combination chemotherapy of reduced (Souhami *et al*, 1997) or split doses of cisplatin plus etoposide (SPE) (Murray *et al*, 1998; Westeel *et al*, 1998) can be safely and effectively administered in elderly or poor-risk patients with SCLC. Therefore, we conducted a phase III trial comparing CE with SPE in elderly or poor-risk patients with SCLC. Although elderly is not the same as poor-risk, many clinical trials for the elderly have included both types of patients. Therefore, we

decided to include both elderly and poor-risk patients with SCLC at the time of proposal for this phase III trial.

PATIENTS AND METHODS

Patient selection

Eligibility criteria included patients with histologically or cytologically confirmed SCLC who were ≥ 70 years of age and had an Eastern Cooperative Oncology Group performance status (PS) of 0–2, or who were < 70 years in age and had a PS of 3. Additional criteria consisted of extensive disease (ED), chemotherapy-naïve, evaluable or measurable disease, expected survival ≥ 2 months, adequate organ functions (leucocyte count ≥ 4000 mm⁻³, platelet count $\geq 100\,000$ mm⁻³, haemoglobin level ≥ 9.0 g dl⁻¹, AST/ALT $\leq 2 \times$ upper limit of normal range, total bilirubin ≤ 1.5 mg dl⁻¹, creatinine ≤ 1.5 mg dl⁻¹, 24-h creatinine clearance (Cr) ≥ 50 ml min⁻¹, and PaO₂ ≥ 60 mmHg), no symptomatic pericardial or pleural effusion requiring drainage, no active concomitant malignancy, no senile dementia, and written informed consent. Exclusion criteria included brain metastases requiring radiotherapy, superior vena cava (SVC) syndrome requiring radiotherapy, serious medical or psychiatric illness, or pregnancy or lactation. Staging procedures included chest X-ray, computed tomography (CT) scan of the chest, CT scan or magnetic resonance

*Correspondence: Dr H Okamoto;

E-mail address: scyooka@alles.or.jp

Presented in part at the Forty-first Annual Meeting of the American Society of Clinical Oncology, Orlando, FL, May 13–17, 2005.

Received 18 October 2006; revised 25 April 2007; accepted 26 April 2007; published online 19 June 2007

imaging (MRI) of the brain, CT scan or ultrasound of the abdomen, isotope bone scanning, and bone marrow aspiration or biopsy.

Treatment protocol

Patients were randomised to either the CE arm or the SPE arm. The CE regimen consisted of carboplatin area under the curve (AUC) of five intravenously (IV) on day 1 and etoposide 80 mg m^{-2} IV on days 1, 2, and 3. The SPE regimen consisted of cisplatin 25 mg m^{-2} IV on days 1, 2, and 3 and etoposide 80 mg m^{-2} IV on days 1, 2, and 3. Cycles were repeated every 3–4 weeks for up to four courses. In our previous phase II study using the CE regimen for elderly patients with SCLC, carboplatin AUC of 5 on day 1 and etoposide 100 mg m^{-2} on days 1, 2, and 3 were administered every 4 weeks (Okamoto *et al*, 1999). However, because grade 3 or 4 neutropenia occurred in 91% of the patients, in the current phase III trial we decided to reduce the etoposide dosage to 80 mg m^{-2} on days 1, 2, and 3, and repeat the cycle every 3–4 weeks instead of every 4 weeks. Twenty-four-hour Ccr was substituted for glomerular filtration rate (GFR) in Calvert's formula. Antiemetic prophylaxis with 5-HT₃ antagonists plus dexamethasone was used at the treating physician's discretion. According to the Japanese approved guideline, prophylactic use of recombinant human granulocyte colony-stimulating factor (G-CSF) was recommended for daily administration after day 4 until the leucocyte (neutrophil) count exceeded $10\,000 (5000) \text{ mm}^{-3}$. If the leucocyte (neutrophil) count decreased to less than $3000 (1500) \text{ mm}^{-3}$, then G-CSF was restarted. However, the actual use of G-CSF was left at the discretion of the treating physician. Subsequent courses of chemotherapy were initiated when leucocyte count $\geq 3000 \text{ mm}^{-3}$; platelet count $\geq 75\,000 \text{ mm}^{-3}$; $\text{Cr} \leq 1.5 \text{ mg dl}^{-1}$; $\text{AST/ALT} \leq 2.5 \times$ upper limit of normal range; and either $\text{PS} \leq 2$ and age ≥ 70 years, or $\text{PS} \leq 3$ and age < 70 years were satisfied both after day 21 and two or more days after the discontinuation of G-CSF. If the above criteria were not satisfied by the first day of the next course, treatment was withheld until full recovery. If more than 6 weeks passed from day 1 of the last course, the patient was removed from protocol treatment. Dose modifications were made based only on grade 4 haematologic toxicities. If grade 4 leucopenia or neutropenia lasting 4 days or more was present, or grade 4 thrombocytopenia occurred, the doses for the next course were carboplatin AUC of 4 on day 1, cisplatin 20 mg m^{-2} for 3 days, and etoposide 60 mg m^{-2} for 3 days. If the same haematologic toxicity was observed after dose reduction, the patient was removed from protocol treatment. If grade 3 or 4 non-haematologic toxicities, except for nausea/vomiting and hyponatraemia, occurred, the patient was removed from protocol treatment even if the toxicities improved thereafter.

Responders after four courses were not allowed to receive further chemotherapy until progressive disease (PD) developed. Although post-protocol treatment was left at the discretion of the physician, crossover treatment was prohibited.

Evaluation

Tumour responses were evaluated according to World Health Organization criteria (World Health Organization, 1979). Toxicities were evaluated according to JCOG Toxicity Criteria (Tobinai *et al*, 1993), which are similar to the National Cancer Institute-Common Toxicity Criteria (NCI-CTC ver 1) for the grading of toxicities.

Palliation score

Study-specific eight-item palliation scores were completed by patients before treatment and 3 weeks after the third course of chemotherapy. The attending physicians were not allowed to complete the scores. The items consisted of cough, pain, anorexia, shortness of breath, well-being, nausea, diarrhoea or constipation, and sleep. The items were scored as not at all present (0), a little

(1), moderate (2), and very much (3). The sum of the total score for all eight items was compared between the baseline and post-treatment assessments. If the post-treatment score was below the baseline score, the palliation score for that patient was judged as having shown improvement.

Study design and statistics

This trial was designed as a multicentre, prospective, randomised phase III trial. The study protocol was approved by the Clinical Trial Review Committee of JCOG and the institutional review board of each participating institution before the initiation of the study. The primary endpoint was overall survival (OS). In this study, the experimental arm was the CE arm and the control was the SPE arm. The MST of our previous phase II trial for elderly patients with extensive disease small-cell lung cancer (ED-SCLC) using the CE regimen was 10.1 months. The MST of the SPE regimen for a similar population was not available at the time of the study proposal. Although Westeel and co-workers in 1998 and Murray and co-workers in 1998 reported an excellent MST of SPE plus concurrent chest radiotherapy for elderly or frail patients with limited disease (LD)-SCLC, an MST of the SPE regimen for elderly or frail patients with ED-SCLC was not available at that time. The only data available on the CAV/PE regimen for elderly or poor-risk patients with SCLC using reduced cisplatin (60 mg m^{-2} IV on day 1) were reported by Souhami and co-workers in 1997 and the MST of that study was 5.9 months. Therefore, for statistical calculations in the current phase III trial, we used the MST value of the Souhami trial for the control arm instead of the MST of the SPE regimen. In addition, an individualised AUC-based dosing strategy of carboplatin was expected to have greater efficacy and less toxicity compared with the SPE regimen at that time. This trial was designed as a superiority trial and the planned sample size was 110 patients in each arm for 80% power to detect a 0.67 hazard ratio for CE to SPE in OS at an alpha of 0.025 (one sided) (Schoenfeld and Richter, 1982). Patients were randomised to receive either CE or SPE with a minimisation method for balancing centre, PS (0–1 vs 2–3) and age (≥ 70 years vs < 70 years).

Survival distributions were compared by unstratified log-rank test. Proportion of improvement in palliation score was evaluated by Fisher's exact test. The change in each symptom score by treatment arm was evaluated by the Wilcoxon rank-sum test. The relationship between the interval of each chemotherapy course and the two regimens was evaluated by the Wilcoxon rank-sum test. Multivariate analysis was performed using Cox's proportional hazards model to evaluate the importance of seven clinically selected variables (treatment arm, PS, age, sex, lactate dehydrogenase level, alkaline phosphatase level, and leucocyte count) as prognostic factors. All *P*-values in this report are two sided, excluding *P*-values for OS and progression-free survival (PFS).

The interim analysis was performed after half of the planned number of patients had been enrolled in March 2002, with adjustment for multiplicity by the alpha-spending function (DeMets and Lan, 1994) with an O'Brien-Fleming type boundary. Because the interim analysis did not meet the prespecified stopping criteria, the study was continued and the planned accrual of 220 patients was randomised in this trial.

RESULTS

Patient characteristics

Between August 1998 and February 2004, a total of 220 patients were registered from 24 institutions. Baseline characteristics were well balanced between the arms. Median age was 74 years, 92% were 70 years or older, 88% were male, and 74% had a PS of 0 or 1 (Table 1). One patient in the CE arm was found to have LD after the completion of protocol chemotherapy due to protocol violation, and this patient was considered ineligible (Figure 1).

Delivery of treatment

Reasons for termination of treatment are listed in Figure 1, and there were no major differences between the arms. Of the patients, 63% in the CE arm and 67% in the SPE arm completed four courses, and 11% in the CE arm and 8% in the SPE arm did not complete treatment because of toxicity or complications. Treatment-related death (TRD) occurred in four patients; three patients in the CE arm and one in the SPE arm. All TRDs of patients who were ≥ 70 years old with a good pretreatment PS (all PS 1) were associated with neutropenic infection, which occurred after the first course of chemotherapy. Although the median interval of chemotherapy was slightly prolonged in the CE arm than in the SPE arm, total delivered courses were similar between the arms (Table 2). One patient in the SPE arm never received chemotherapy due to the occurrence of delirium after registration. Dose reduction was more frequently observed in the CE arm than in the SPE arm: 29% vs 10%, $P < 0.01$. Course delay, G-CSF delivery and total courses with G-CSF delivery were similar between the arms.

Toxicity and palliation score

Toxicities are listed in Table 3. Grade 3 or 4 leucopenia and neutropenia occurred in 54 and 95% of the CE arm vs 51 and 90% of the SPE arm, respectively. Grade 3 or 4 thrombocytopenia occurred more frequently in the CE arm than in the SPE arm: 56 vs 16%, $P < 0.01$. Gastrointestinal toxicities including nausea or

vomiting and diarrhoea were mild in both arms. There were few grade 3 or 4 toxicities and no remarkable differences between the arms. Other non-haematologic toxicities were similarly distributed between the arms. Grade 3–4 hyponatraemia, mainly caused by syndrome of inappropriate antidiuretic hormone (SIADH) secretion, occurred in 14–16% of the patients. More importantly, thrombocytopenia occurred more frequently in the CE arm, but none of the patients in either arm showed grade 3 or 4 bleeding. Only one patient in the CE arm showed grade 2 bleeding. Because no grading of febrile neutropenia was listed in JCOG toxicity criteria, the rate of the toxicity was not investigated in this study.

Baseline and post-treatment palliation scores were evaluated in 220/220 (100%) and 208/220 (95%) patients, respectively. We handled missing values by imputing the worst score. Improvement was achieved in 69 (63%) patients in the CE arm vs 61 (56%) patients in the SPE arm, although the difference was not statistically significant ($P = 0.34$). Similarly, there were no statistical differences in the change of each symptom score between the arms (Table 4).

Objective tumour response, PFS and OS

The objective response rate of 73% was quite similar between the arms. Five CRs and 75 PRs were observed in each arm (Table 5). Progression-free survival curves and OS curves are shown in Figure 2A and B. Ninety-seven percent of the patients had progressed or died at the time of final analysis. Progression-free survival was quite similar between the arms ($P = 0.20$, one sided).

Table 1 Patient characteristics

	CE (n = 110)	SPE (n = 110)	P-value
Age (years)			
Median (range)	74 (56–86)	73.5 (55–85)	0.34
≥ 70 years old (%)	102 (93)	100 (91)	0.81
Sex (male/female)	95/15	98/12	0.68
ECOG PS, 0–1/2/3	81/21/8	81/19/10	0.80
$\geq 5\%$ weight loss	26	38	0.18
LN metastasis			
Contralateral mediastinum	71	59	0.13
Supraclavicular	89	79	0.15
Distant metastasis			
Liver	30	30	1.0
Lung	31	30	1.0
Brain	18	18	1.0
Bone	25	17	0.23
Adrenal	13	7	0.24
Bone marrow	12	12	1.0

CE, carboplatin plus etoposide; ECOG, Eastern Cooperative Oncology Group; LN, lymph node; PS, performance status; SPE, split doses of cisplatin plus etoposide.

Table 2 Compliance and drug delivery

	CE (n = 110)	SPE (n = 109*)	P-value
Median interval of each chemotherapy (days) (range)			
1–2	27 (14–35)	23 (20–37)	0.02 ^b
2–3	25 (21–56)	22 (20–35)	0.07 ^b
3–4	27 (21–36)	24 (21–38)	0.05 ^b
Total delivered courses/projected courses	353/440 (80%)	360/436 (83%)	
Dose reduction	32 (29%)	11 (10%)	<0.01 ^c
Course delay	45 (41%)	40 (37%)	0.58 ^c
G-CSF delivery	81 (74%)	84 (77%)	0.64 ^c
No. of courses with G-CSF delivery/number of total courses	183/354 (52%)	203/362 (56%)	

CE, carboplatin plus etoposide; G-CSF, granulocyte colony-stimulating factor; SPE, split doses of cisplatin plus etoposide. *One patient never received chemotherapy due to delirium after registration. ^bWilcoxon rank-sum test. ^cFisher's exact test.




Figure 1 Flow diagram of randomised phase III trial of CE vs SPE in elderly or poor-risk patients with extensive disease SCLC.

Table 3 Toxicities (JCOG Toxicity Criteria, Worst Grade of Any Course)

Toxicity	CE					SPE					P-value
						Grade					
	1	2	3	4	3+4 (%)	1	2	3	4	3+4 (%)	
Haematologic											
Leucopenia	5	45	46	13	(54)	8	43	49	7	(51)	0.79
Neutropenia	0	5	46	58	(95)	4	7	41	57	(90)	0.22
Anaemia	9	58	32	—	(29)	20	45	27	—	(25)	0.54
Thrombocytopenia	20	18	29	32	(56)	16	15	12	5	(16)	<0.01
Non-haematologic											
Nausea/vomiting	40	24	2	—	(2)	46	28	3	—	(3)	0.68
Diarrhoea	8	9	1	0	(1)	11	3	1	0	(1)	1.0
Bilirubin	—	31	0	0	(0)	—	16	1	0	(1)	0.50
AST	47	9	3	0	(3)	30	8	6	0	(6)	0.33
ALT	40	9	2	0	(2)	38	8	4	0	(4)	0.45
Creatinine	10	2	0	0	(0)	27	3	1	0	(1)	0.50
Hyponatremia	38	11	7	11	(16)	46	20	6	9	(14)	0.58
PaO ₂	39	21	7	1	(10)	44	23	2	1	(4)	0.22
Fever	15	15	0	0	(0)	21	16	0	0	(0)	—
Infection	12	15	5	3	(7)	16	7	5	1	(6)	0.78
Bleeding	8	1	0	0	(0)	4	0	0	0	(0)	—
Neurologic-sensory	2	1	0	—	(0)	3	2	0	—	(0)	—
Alopecia	67	22	—	—	—	66	15	—	—	—	—

CE, carboplatin plus etoposide; JCOG, Japan Clinical Oncology Group; PaO₂, partial pressure of oxygen; SPE, split doses of cisplatin plus etoposide.

Table 4 Palliation score

Symptom	CE		SPE		P ^a
	Change from baseline		Change from baseline		
	Mean (s.d.)	Median (range)	Mean (s.d.)	Median (range)	
Cough	-0.38 (1.16)	0 (-3 to 3)	-0.54 (1.06)	0 (-3 to 3)	0.51
Pain	-0.19 (1.00)	0 (-3 to 3)	-0.19 (0.96)	0 (-3 to 3)	0.96
Anorexia	-0.07 (1.16)	0 (-3 to 3)	0.08 (1.22)	0 (-3 to 3)	0.37
Shortness of breath	-0.05 (1.02)	0 (-2 to 3)	-0.31 (0.95)	0 (-3 to 3)	0.12
Well-being	-0.15 (1.13)	0 (-3 to 3)	-0.02 (1.14)	0 (-3 to 3)	0.48
Nausea	0.16 (0.84)	0 (-2 to 3)	0.26 (0.80)	0 (-1 to 3)	0.21
Diarrhoea or constipation	0.05 (1.07)	0 (-3 to 3)	0.04 (0.99)	0 (-3 to 3)	0.69
Sleep	-0.15 (1.08)	0 (-3 to 3)	-0.04 (0.89)	0 (-3 to 2)	0.10
Total	-0.80 (6.04)	-2 (-12 to 22)	-0.71 (5.35)	-1 (-15 to 21)	0.32

CE, carboplatin plus etoposide; s.d., standard deviation; SPE, split doses of cisplatin plus etoposide. ^aWilcoxon rank-sum test.

The MST was 5.2 months in the CE arm vs 4.7 months in the SPE arm. OS was very similar between the arms ($P=0.54$, one sided). The MST and 1-year survival rate was 10.6 months and 41% in the CE arm vs 9.9 months and 35% in the SPE arm.

Second-line chemotherapy

According to an *ad-hoc* survey (not pre-specified in the protocol), 130 (59%) patients (68 (62%) patients in the CE arm and 62 (56%) in the SPE arm) received second-line chemotherapy after relapse and the regimens were almost equally distributed between the arms. The same regimen as the initial chemotherapy, platinum-based combinations, and irinotecan regimens with or without other agents were administered in 17 (15%), 48 (44%), and 40 (36%) patients in the CE arm vs 10 (9%), 44 (40%), and 40 (36%) in

Table 5 Therapeutic response (WHO)

	CE	SPE	Total
CR	5	5	10
PR	75	75	150
NC	17	11	28
PD	11	16	27
NE	2	3	5
Total	110	110	220
Response rate	73%	73%	
95% CI	63-81%	63-81%	

CE, carboplatin plus etoposide; CI, confidence interval; CR, complete response; NE, no change; NC, not evaluable; PD, progressive disease; PR, partial response; SPE, split doses of cisplatin plus etoposide; WHO, World Health Organization.




Figure 2 (A) PFS curves ($n=220$). (B) OS curves ($n=220$). (C) Survival curves of the patients ≥ 70 years of age with a PS of 0-2 ($n=202$).

the SPE arm. Other chemotherapy regimens included topotecan monotherapy, amrubicin monotherapy, or other regimens.

Subset analysis and multivariate analysis

Subset analysis was performed according to PS and age (Table 6). There were no differences in OS between the arms in any subset; thus, an interaction between treatment and PS is unlikely. The survival curves of the patients ≥ 70 years of age with a PS of 0-2 are shown in Figure 2C, and the survival curves were very

Table 6 Subset analysis – overall survival

Subgroup	Number of patients (%)	MST (months)	
		CE	SPE
PS 0-1	162 (74)	10.9	10.1
PS 2-3	58 (26)	8.3	8.1
<70 years and PS 3	18 (8)	7.1	6.9
≥ 70 years and PS 0-2	202 (92)	10.8	10.0

CE, carboplatin plus etoposide; MST, median survival time; PS, performance status; SPE, split doses of cisplatin plus etoposide.

Table 7 Multivariate analysis with baseline prognostic factors

Variables	P-value	Hazard ratio	95% CI
Treatment arm (CE vs. SPE)	0.99	0.99	0.75-1.33
Alkaline phosphatase level (normal vs abnormal)	0.97	0.99	0.68-1.46
Lactate dehydrogenase level ($\geq \times 1.5$ vs $< \times 1.5$)	<0.001	1.69	1.23-2.26
Leucocyte count ($\geq 10000/\text{mm}^3$ vs $< 10000/\text{mm}^3$)	0.06	1.82	0.99-3.36
Age (≥ 75 years vs < 75 years)	0.77	1.05	0.78-1.41
PS (2-3 vs 0-1)	0.41	1.15	0.82-1.61
Sex (female vs male)	0.13	0.70	0.45-1.11

CE = carboplatin plus etoposide; SPE = split doses of cisplatin plus etoposide; PS = performance status; CI = confidence interval.

similar with that of original overall populations. Even in the multivariate analysis with seven selected baseline variables, there was no difference in OS between the arms. High lactate dehydrogenase level was most strongly associated with poor prognosis (Table 7).

DISCUSSION

Until recently, there was no standard chemotherapeutic regimen for elderly SCLC patients. Two phase III (Medical Research Council Lung Cancer Working Party, 1996; Souhami *et al*, 1997) and two randomised phase II trials (Pfeiffer *et al*, 1997; Arzizzone *et al*, 2005) have shown that suboptimal chemotherapies, such as oral etoposide monotherapy or attenuated doses of combination chemotherapy, may lead to reduced survival in elderly or poor-risk SCLC patients when compared with standard doses of combination chemotherapies. The CE regimen, which has acceptable toxicities and reproducible efficacy, has been used in elderly or poor-risk patients with SCLC worldwide, although there have been substantial differences in toxicities and efficacy between the reported phase II trials. Four trials demonstrated both favourable toxicities and efficacy (Carney, 1995; Evans *et al*, 1995; Matsui *et al*, 1998; Okamoto *et al*, 1999) and three showed somewhat disappointing results because of suboptimal doses of oral etoposide (Larive *et al*, 2002), greater inclusion of patients with poor prognostic factors (Samantas *et al*, 1999), and deterioration of comorbidities as a result of chemotherapy (Quoix *et al*, 2001). No phase III trial evaluating the role of the CE regimen in this population has been reported until now.

This is the first phase III trial comparing carboplatin-based CE and cisplatin-based SPE regimens in elderly or poor-risk patients with ED-SCLC. In addition, this is also the largest randomised trial specifically designed for elderly or poor-risk SCLC patients. Although there was no significant difference in the palliation scores, response rate, and OS between the arms, the efficacy of

both regimens was promising, as this study included only elderly or poor-risk patients with SCLC. Most toxicities were tolerable and the treatment compliance was also favourable in both arms. Approximately two-thirds of the patients received all four cycles of treatment. The CE arm in the current trial had more pronounced thrombocytopenia, which was considered manageable because none of the patients in the CE arm showed grade 3 or 4 bleeding, and the CE arm had a slightly prolonged course interval and a slightly greater incidence of dose reduction. However, in our opinion, these toxicities are less meaningful in clinical practice. More importantly, the CE regimen does not require hydration and can be given in an outpatient setting. Based on the results of this study, many JCOG members prefer the CE regimen to the SPE regimen and consider it to be more suitable for the control arm of future phase III trials.

The MST of each regimen (10.6 months for CE vs 9.9 months for SPE) was promising considering that this study included only elderly or frail patients with ED-SCLC. However, some retrospective studies have shown that fit elderly patients who have adequate organ functions, a good PS, and no comorbidity are able to tolerate intensive chemotherapy well and show a similar therapeutic response and survival rate as younger patients (Siu *et al*, 1996; Yuen *et al*, 2000). In fact, in this trial the MST of fit elderly patients ≥ 70 years of age with a PS of 0–1 was 10.9 months for the CE arm and 10.1 months for the SPE arm. In contrast, the MST of patients with a PS of 3 was only approximately 7 months. Furthermore, the group of fit elderly patients comprised 74% of the patients in this study. Therefore, the favourable survival rates in our trial may be attributable to patient selection. In other words, one limitation of this study is that the results of this trial cannot be extrapolated to frail elderly with a poor PS and/or comorbid illness because of the likelihood of greater inclusion of fit elderly patients in this trial.

Although the total dose in both the CE and SPE arms was slightly lower than the standard regimen, 92% of the patients showed grade 3 or 4 neutropenia, and dose reduction and course delay occurred frequently. However, the MST of both regimens was comparable with that of non-elderly or non-selected patients with ED-SCLC in historical reports (Noda *et al*, 2002; Niell *et al*, 2005). These findings suggest that both regimens are not suboptimal, but are near-full and effective doses for elderly or poor-risk patients with ED-SCLC. The CE arm in the current trial had a slightly prolonged course interval and a slightly greater incidence of dose reduction when compared to the SPE regimen. However, 95% of the patients showed grade 3 or 4 neutropenia and 56% showed grade 3 or 4 thrombocytopenia. Therefore, we believe that the dose escalation of the CE regimen may be difficult in this trial.

It remains unclear whether the elderly are able to tolerate a single modest dose of cisplatin ($60\text{--}80\text{ mg m}^{-2}$ IV) on day 1. We feel that a fit elderly person who passes strict eligibility criteria can receive a modest dose of cisplatin IV on day 1. However, the more common situation is of elderly patients who have comorbidity and a poor PS, and cannot tolerate a standard single dose of cisplatin. Westeel *et al* (1998) and Murray *et al* (1998) reported that split doses of cisplatin were safely and effectively administered in elderly or frail patients with LD-SCLC. The SPE regimen appeared to be an appropriate treatment for elderly patients with SCLC who cannot tolerate a standard single dose of cisplatin. However, it remains unclear whether fit elderly patients in our trial can tolerate a standard single dose of cisplatin, and if so, it also remains unclear whether fit elderly patients who receive a standard single dose of cisplatin are able to achieve a more improved survival than those who receive SPE. Unfortunately, no randomised study comparing a single standard dose of cisplatin with SPE has been reported in fit elderly patients with SCLC.

There are some problems with the design in this study. The hypothesis was that carboplatin would improve survival, and

the design of the trial was a superiority design with survival as the primary end point. However, this hypothesis was based on two possible misconceptions. First, carboplatin could be better dosed and might be more efficacious than cisplatin in SCLC. Unfortunately, this hypothesis could not be sustained on the basis of the available literatures. A number of clinical trials have indicated that carboplatin-based combination chemotherapy has a similar or slightly reduced efficacy compared with cisplatin-based combination chemotherapy against various tumours (Go and Adjei, 1999; Hotta *et al*, 2004). Therefore, our trial should have been designed as a non-inferiority trial. However, if this trial were planned as a non-inferiority trial, a total sample size would be about 500 to 1000 patients, with equal expected survival and a non-inferiority margin for hazard ratio ranging from 1.2 to 1.3. Second, the cisplatin dose in the control arm was an attenuated dose. Souhami *et al* (1997) used reduced dose of cisplatin (60 mg m^{-2} IV on day 1) and Murray *et al* (1998) used a single course of a split cisplatin dose in their studies. These regimens were completely different from the control arm in the present study. A standard dose of cisplatin given in 3 days is the best way of giving standard cisplatin (30 mg m^{-2} IV on days 1–3) with etoposide (130 mg m^{-2} IV on days 1–3), according to the North Central Cancer Treatment Group (Maksmiuk *et al*, 1994). Had standard SPE been used for the control arm, better survival might have been achieved with increased toxicities. Another problem with the design was the inclusion of patients with a PS of 3, even if they were less than 70 years old. This made the target population heterogeneous. The number of such patients actually recruited was quite small, so emphasising the inappropriateness of their inclusion. A further limitation of this study may be a long accrual period of five-and-a-half years. Because our oncologists might have been afraid of the risk of TRD or increased toxicities in frail elderly with a poor PS and/or comorbid illness, more fit elderly patients were selectively registered and consequently the accrual rate was very slow.

In our trial, although both regimens were well-tolerated and efficacy was promising, over 90% of the patients in both arms showed grade 3 or 4 neutropenia, which may be justified and acceptable for a clinical trial involving elderly or poor risk patients with ED-SCLC, because only 6% of the patients showed grade 3 or 4 infection and TRD occurred in only four (1.8%) patients. Because all TRD occurred after the first course of chemotherapy, careful monitoring and management is necessary, particularly in the first course, if CE or SPE are administered to elderly or frail patients. Several retrospective analyses (Findlay *et al*, 1991; Radford *et al*, 1992) and a prospective study (Timmer-Bonte *et al*, 2005) have shown that standard-dose chemotherapy without G-CSF support causes more risk of early death and sepsis in the older population. Moreover, the American Society of Clinical Oncology (ASCO) guideline recommends the use of prophylactic G-CSF in patients at higher risk for chemotherapy-induced infection, such as those having a poor PS, older age, or comorbid illness (Smith *et al*, 2006). In this trial, the prophylactic use of G-CSF was recommended, but the actual use was left to the discretion of the treating physician because the use of G-CSF leads to increased drug cost. Although G-CSF was administered in only 54% of the total courses, we believe that the prophylactic use of G-CSF with CE regimen should be recommended in a new trial or clinical practice.

In conclusion, although the SPE regimen is still considered to be the standard treatment for elderly or poor-risk patients with ED-SCLC, the CE regimen can be an alternative for this population considering the risk-benefit balance. Based on the results of our trial, a phase III trial of the CE regimen vs amrubicin monotherapy, supported by a pharmaceutical company, is now ongoing in elderly patients with ED-SCLC in Japan, and a comparative trial of the CE regimen vs carboplatin plus irinotecan regimen (Okamoto *et al*, 2006) is being discussed for a future trial in our group.

ACKNOWLEDGEMENTS

We are indebted to Ms Mieko Imai and Ms Tomoko Yamabe for data management, and to Dr Haruhiko Fukuda for direction of the JCOG

REFERENCES

- Ardizzoni A, Favaretto A, Boni L, Baldini E, Castiglioni F, Antonelli P, Pari F, Tibaldi C, Altieri AM, Barbera S, Cacciani G, Raimondi M, Tixi L, Stefani M, Monfardini S, Antilli A, Rosso R, Paccagnella A (2005) Platinum-etoposide chemotherapy in elderly patients with small-cell lung cancer: results of a randomized multicenter phase II study assessing attenuated-dose or full-dose with lenograstim prophylaxis-A Forza Operativa Nazionale Italiana Carcinoma Polmonare and Gruppo Studio Tumori Polmonari Veneto (FONICAP-GSTPV) study. *J Clin Oncol* 23: 569-575
- Carney DN (1995) Carboplatin/etoposide combination chemotherapy in the treatment of poor prognosis patients with small cell lung cancer. *Lung Cancer* 12(Suppl 3): S77-S83
- DeMets DL, Lan KK (1994) Interim analysis: the alpha spending function approach. *Stat Med* 13: 1341-1352
- Evans WK, Radwi A, Tomiak E, Logan DM, Martins H, Stewart DJ, Goss G, Maroun JA, Dahrouge S (1995) Oral etoposide and carboplatin: effective therapy for elderly patients with small cell lung cancer. *Am J Clin Oncol* 18: 149-155
- Findlay MP, Griffin AM, Raghavan D, McDonald KE, Coates AS, Duval PJ, Gianoutsos P (1991) Retrospective review of chemotherapy for small cell lung cancer in the elderly: dose the end justify the means? *Eur J Cancer* 27: 1597-1601
- Go RS, Adjei AA (1999) Review of the comparative pharmacology and clinical activity of cisplatin and carboplatin. *J Clin Oncol* 17: 409-422
- Hotta K, Matsuo K, Ueoka H, Kiura K, Tabata M, Tanimoto M (2004) Meta-analysis of randomized clinical trials comparing cisplatin to carboplatin in patients with advanced non-small-cell lung cancer. *J Clin Oncol* 22: 3838-3852
- Larive S, Bombaron P, Riou R, Fournel P, Perol M, Lena H, Dussopt C, Phillip-Joet F, Touraine F, Lecaer H, Souquet PJ (2002) Carboplatin-etoposide combination in small cell lung cancer patients older than 70 years: a phase II trial. *Lung Cancer* 35: 1-7
- Maksmiuk AW, Jett JR, Earle JD, Su JQ, Diegert FA, Mailliard JA, Kardinal CG, Krook JE, Veeder MH, Wiesenfeld M, Tschetter LK, Levitt R (1994) Sequencing and schedule effects of cisplatin plus etoposide in small-cell lung cancer: results of a North Central Cancer Treatment Group randomized clinical trial. *J Clin Oncol* 12: 70-76
- Matsui K, Masuda N, Fukuoka M, Yana T, Hirashima T, Komiya T, Kobayashi M, Kawahara M, Atagi S, Ogawara M, Negoro S, Kudoh S, Furuse K (1998) Phase II trial of carboplatin plus oral etoposide for elderly patients with small-cell lung cancer. *Br J Cancer* 77: 1961-1965
- Medical Research Council Lung Cancer Working Party (1996) Comparison of oral etoposide and standard intravenous multidrug chemotherapy for small-cell lung cancer: a stopped multicentre randomised trial. *Lancet* 348: 563-566
- Morita T (2002) A statistical study of lung cancer in the annual of pathological autopsy cases in Japan, from 1958 to 1997, with reference to time trends of lung cancer in the world. *Jpn J cancer Res* 93: 15-23
- Murray N, Grafton C, Shah A, Gelmon K, Kostashuk E, Brown E, Coppin C, Goldman A, Page R (1998) Abbreviated treatment for elderly, infirm, or noncompliant patients with limited-stage small-cell lung cancer. *J Clin Oncol* 16: 3323-3328
- Niell HB, Herndon II JE, Miller AA, Watson DM, Sandler AB, Kelly K, Marks RS, Perry MC, Ansari RH, Otterson G, Ellerton J, Vokes EE, Green MR (2005) Randomized phase III intergroup trial of etoposide and cisplatin with or without paclitaxel and granulocyte colony-stimulating factor in patients with extensive-stage small-cell lung cancer: Cancer and Leukemia Group B Trial 9732. *J Clin Oncol* 23: 3752-3759
- Noda K, Nishiwaki Y, Kawahara M, Negoro S, Sugiura T, Yokoyama A, Fukuoka M, Mori K, Watanabe K, Tamura T, Yamamoto S, Saijo N (2002) Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med* 346: 85-91
- Okamoto H, Naoki K, Narita Y, Hida N, Kunikane H, Watanabe K (2006) A combination chemotherapy of carboplatin and irinotecan with granulocyte colony-stimulating factor (G-CSF) support in elderly patients with small cell lung cancer. *Lung Cancer* 53: 197-203
- Okamoto H, Watanabe K, Nishiwaki Y, Mori K, Kurita Y, Hayashi I, Masutani M, Nakata K, Tsuchiya S, Isobe H, Saijo N (1999) Phase II study of area under the plasma-concentration-vs-time curve-based carboplatin plus standard-dose intravenous etoposide in elderly patients with small-cell lung cancer. *J Clin Oncol* 17: 3540-3545
- Pfeiffer P, Rytter C, Madsen EL, Moeholt K, Hansen O, Bentzen S, Palshof T, Rose C (1997) Re: five-day oral etoposide treatment for advanced small-cell lung cancer: randomized comparison with intravenous chemotherapy. *J Natl Cancer Inst* 89: 1892-1893
- Quoix E, Breton JL, Daniel C, Jacoulet P, Debieve D, Paillet N, Kessler R, Moreau L, Coetmeur D, Lemarie E, Milleron B (2001) Etoposide phosphate with carboplatin in the treatment of elderly patients with small-cell lung cancer: a phase II study. *Ann Oncol* 12: 957-962
- Radford JA, Ryder WD, Dodwell D, Anderson H, Thatcher N (1992) Predicting septic complications of chemotherapy: An analysis of 382 patients treated for small cell lung cancer without dose reduction after major sepsis. *Eur J Cancer* 29A: 81-86
- Samantas E, Skarlos DV, Pectasides D, Nicolaidis P, Kalofonos H, Mylonakis N, Vardoulakis Th, Kosmidis P, Pavlidis N, Fountzilas G (1999) Combination chemotherapy with low doses of weekly carboplatin and oral etoposide in poor risk small cell lung cancer. *Lung Cancer* 23: 159-168
- Schoenfeld DA, Richter JR (1982) Nomograms for calculating the number of patients needed for a clinical trial with survival as an endpoint. *Biometrics* 38: 163-170
- Siu LL, Shepherd FA, Murray N, Feld R, Pater J, Zee B (1996) Influence of age on the treatment of limited-stage small-cell lung cancer. *J Clin Oncol* 14: 821-828
- Smith TJ, Khatcheressian J, Lyman GH, Ozer H, Armitage JO, Balducci L, Bennett CL, Cantor SB, Crawford J, Cross SJ, Demetri G, Desch CE, Pizzo PA, Schiffer CA, Schwartzberg L, Somerfield MR, Somlo G, Wade JC, Wade JL, Winn RJ, Wozniak AJ, Wolff AC (2006) 2006 update of recommendations for the use of white blood cell growth factors: an evidence-based clinical practice guideline. *J Clin Oncol* 24: 3187-3205
- Souhami RL, Spiro SG, Rudd RM, Ruiz de Elvira MC, James LE, Gower NH, Lamont A, Harper PG (1997) Five-day oral etoposide treatment for advanced small-cell lung cancer: randomized comparison with intravenous chemotherapy. *J Natl Cancer Inst* 89: 577-580
- Timmer-Bonte JN, de Boo TM, Smit HJ, Biesma B, Wilschut FA, Cheragwandi SA, Termeer A, Hensing CA, Akkermans J, Adang EM, Bootsma GP, Tjan-Heijnen VC (2005) Prevention of chemotherapy-induced febrile neutropenia by prophylactic antibiotics plus or minus granulocyte colony-stimulating factor in small-cell lung cancer: a dutch randomized phase III study. *J Clin Oncol* 23: 7974-7984
- Tobinai K, Kohno A, Shimada Y, Watanabe T, Tamura T, Takeyama K, Narabayashi M, Fukutomi T, Kondo H, Shimoyama M, Suemasu K (1993) Toxicity grading criteria of the Japan Clinical Oncology Group. *Jpn J Clin Oncol* 23: 250-257
- Westeel V, Murray N, Gelmon K, Shah A, Sheehan F, McKenzie M, Wong F, Morris J, Grafton C, Tsang V, Goddard K, Murphy K, Parsons C, Amy R, Page R (1998) New combination of the old drugs for elderly patients with small-cell lung cancer: a phase II study of the PAVE regimen. *J Clin Oncol* 16: 1940-1947
- World Health Organization (1979) *WHO Handbook for Reporting Results of Cancer Treatment* WHO offset publication No. 48 Geneva: World Health Organization
- Yuen AR, Zou G, Turrisi AT, Sause W, Komaki R, Wagner H, Aisner SC, Livingston RB, Blum R, Johnson DH (2000) Similar outcome of elderly patients in Intergroup Trial 0096: cisplatin, etoposide, and thoracic radiotherapy administered once or twice daily in limited stage small cell lung carcinoma. *Cancer* 89: 1953-1960

data center and oversight of management of the study. This work was supported in part by Grants-in-Aid for Cancer Research and for the Second-Term Comprehensive 10-Year Strategy for Cancer Control from the Ministry of Health, Labour, and Welfare (Tokyo).

Appendix

This study was coordinated by the Japan Clinical Oncology Group (N Saijo, Chairperson) and was performed with the cooperation of the following institutions and investigators: Tochigi Cancer Center Hospital, Tochigi (K Mori, M Noda, T Kondo, and Y Kamiyama); National Nishi-Gunma Hospital, Gunma (S Tsuchiya, Y Koike, K Satoh, A Tohi, and K Kaira); Gunma Cancer Center Hospital, Gunma (K Minato); Saitama Cancer Center Hospital, Saitama (H Sakai, K Kobayashi, and R Kuroki); National Cancer Center, Central Hospital, Tokyo (T Tamura, Y Ohe, H Kunitoh, I Sekine, H Nokihara, and H Murakami); National Cancer Center Hospital East, Chiba (R Kakinuma, K Kubota, H Ohmatsu, K Gotoh, and S Niho); National International Medical Center, Tokyo (Y Takeda, S Izumi, A Kawana, M Kamimura, and M Iikura); Toranomon Hospital, Tokyo (K Kishi, and M Kawabata); Kanagawa Cancer Center Hospital, Kanagawa (K Yamada, I Nomura, F Oshita, and M Ikehara); Yokohama Municipal Citizen's Hospital, Kanagawa (K Watanabe, H Kunikane, H Okamoto, A Nagatomo, and H Aono); Niigata Cancer Center Hospital, Niigata (A Yokoyama, H Tsukada, M Makino, T Shinbo, S Kinebuchi, J Tanaka, M Tango, and

T Ohara); Gifu City Hospital, Gifu (T Sawa, M Miwa, T Ishiguro, M Sawada, and T Yoshida); Aichi Cancer Center Central Hospital, Aichi (K Yoshida, and T Hida); Aichi Cancer Center Aichi Hospital, Aichi (H Saitoh, and M Okuno); Osaka City University Medical School, Osaka (S Kudoh, S Kyoh, H Kamoi, N Yoshimura, T Kodama, K Ohtani, S Shiraishi, S Nomura, S Enomoto, H Matsuura, and R Wake); Kinki University Medical School, Osaka (T Nogami, N Yamamoto, S Sakai, K Kodama, K Akiyama, J Tsurutani, K Tamura); Osaka Prefectural Adult Disease Center, Osaka (S Nakamura, F Imamura, M Yoshimura, S Yamamoto, K Ueno, H Ohmiya, H Matsuoka, and H Uda); Osaka Prefectural Respiratory and Allergy Medical Center, Osaka (M Furukawa, T Yamadori, T Takimoto, and T Hirashima); National Kinki Central Thoracic Disease Center, Osaka (S Minami, N Naka, T Kawaguchi, and H Ishikawa); National Toneyama Hospital, Osaka (Y Okano); Osaka City General Medical Center, Osaka (N Takifuji, and M Miyazaki); Kobe City Central Hospital, Kobe (T Nishimura, Y Okazaki, D Kinose, H Fujii, S Takakura, and M Hayashi); Sasebo City General Hospital, Nagasaki (J Araki); Kumamoto Regional Medical Center, Kumamoto (H Senba, T Seto, and S Fujii).



Detection of unsuspected distant metastases and/or regional nodes by FDG-PET in LD-SCLC scan in apparent limited-disease small-cell lung cancer

Seiji Niho^{a,*}, Hirofumi Fujii^b, Koji Murakami^{b,c}, Seisuke Nagase^a, Kiyotaka Yoh^a, Koichi Goto^a, Hironobu Ohmatsu^a, Kaoru Kubota^a, Ryuzo Sekiguchi^d, Shigeru Nawano^d, Nagahiro Saijo^a, Yutaka Nishiwaki^a

^a Division of Thoracic Oncology, National Cancer Center Hospital East, Kashiwanoha 6-5-1, Kashiwa, Chiba 277-8577, Japan

^b Functional Imaging Division, National Cancer Center Research Center for Innovative Oncology, Chiba, Japan

^c PET Center, Dokkyo Medical University, Tochigi, Japan

^d Department of Radiology, National Cancer Center Hospital East, Chiba, Japan

Received 12 January 2007; received in revised form 30 March 2007; accepted 6 April 2007

KEYWORDS

Small-cell lung cancer;
Limited-disease;
FDG-PET;
CT;
Staging;
Occult distant metastasis

Summary We retrospectively investigated the clinical usefulness of fluorodeoxyglucose positron emission tomography (FDG-PET) for evaluation of patients with limited-disease small-cell lung cancer (LD-SCLC) diagnosed by conventional staging procedures. Sixty-three patients received whole body FDG-PET scans after routine initial staging procedures. The findings of FDG-PET scans suggesting extensive-stage disease were confirmed by other imaging tests or by the patient's clinical course. FDG-PET scan findings indicated distant metastases in 6 of 63 patients. Metastatic disease was confirmed in five of these six patients (8%, 95% confidence interval: 3–18%). FDG-PET scan also detected regional lymph node metastases even in nine patients (14%) in whom computed tomography images had been negative, including contralateral lymph node metastasis in three patients. FDG-PET scan detected additional lesions in patients diagnosed as having LD-SCLC by conventional staging procedures. The therapeutic strategies were changed in 8% of patients based on the results of FDG-PET. FDG-PET scan is recommended as an initial staging tool for patients with this disease.

© 2007 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Small-cell lung cancer (SCLC) accounts for 15–20% of all lung cancers. SCLC shows more aggressive biological behaviour than non-small cell lung cancer (NSCLC). A clinical two-stage system proposed by the Veterans Administration Lung

* Corresponding author. Tel.: +81 4 7133 1111;
fax: +81 4 7131 4724.
E-mail address: siniho@east.ncc.go.jp (S. Niho).

Study Group (VALSG) distinguishes limited-disease (LD) and extensive-disease (ED) in SCLC [1]. LD is defined as limited to one hemithorax, including mediastinal, contralateral hilar and ipsilateral supraclavicular lymph nodes, while ED represents tumour spread beyond these regions. Approximately two-thirds of patients with SCLC are diagnosed as having ED at the initial staging. The current standard care for LD-SCLC is a combination of chemotherapy and chest irradiation. With current treatment, patients with LD have a median survival of 23–27 months [2,3], compared to 10–12 months for those with ED [4]. Therefore, accurate pretreatment staging is important for patients with SCLC in order to determine the appropriate therapy.

Conventional staging procedures for lung cancer consist of computed tomography (CT) of the chest and upper abdomen, bone scan, and CT scan or magnetic resonance imaging (MRI) of the brain. Recently, fluorodeoxyglucose positron emission tomography (FDG-PET) was introduced as a staging tool for NSCLC. According to the guidelines of the American Society of Clinical Oncology, PET scan is recommended for survey occult locoregional lesions and distant metastases in patients with NSCLC [5]. Two separate prospective studies demonstrated that FDG-PET detected unsuspected distant metastases in 24% of patients with apparent stage III NSCLC [6,7]. Another study showed that FDG-PET changed or influenced management decisions in 67% of patients with NSCLC. PET plays an important role in staging of NSCLC [8]. However, previous PET studies of SCLC involved only a relatively small number of patients [9–17]. In a prospective study, FDG-PET was performed for 24 patients diagnosed as having LD-SCLC by conventional staging procedures [9]. Based on FDG-PET findings, two of these 24 patients were upstaged to ED. Bone metastases were found in one patient, and contralateral supraclavicular lymph node metastasis in another. Larger studies are required to confirm the role of FDG-PET in the staging of LD-SCLC. In this study, we retrospectively investigated the usefulness of FDG-PET to detect distant metastases or unsuspected regional nodal metastases in patients with LD-SCLC diagnosed by conventional staging procedures.

2. Patients and methods

2.1. Patients

Seventy patients were newly diagnosed as having LD-SCLC by conventional staging procedures at the National Cancer Center Hospital East between July 2003 and December 2006. Conventional staging procedures included history and physical examination, chest radiography, CT scan of the chest, CT scan or ultrasound (US) of the abdomen, bone scan, and CT scan or MRI of the brain. CT scan and MR images were enhanced with contrast media. LD is defined in this study as disease limited to one hemithorax, including mediastinal, contralateral hilar and supraclavicular lymph nodes, ipsilateral pleural effusion, and pericardial effusion, while ED represents tumour spread beyond these manifestations [18]. This study included 63 patients who received whole body FDG-PET scan after the routine initial staging procedures. Fifty-seven were male and the remaining 6 were

female. Median age was 64 years, range 48–80 years. Forty-two patients received FDG-PET before commencement of chemotherapy. The remaining 21 patients received FDG-PET 1 to 11 days (median: 4 days) after commencement of chemotherapy. Forty-four and 19 patients received CT scan and US of the abdomen, respectively.

2.2. FDG-PET scan

FDG-PET scans were performed before March 2005 (patients No. 1–25), and FDG-PET/CT scans were performed after April 2005 (patients No. 26–63). Three hundred MBq of F-18 FDG were intravenously injected after at least 6 h of fasting. Acquisition was initiated 60 min after the injection. FDG-PET imaging was performed using a GE Advance Scanner (General Electric Medical System, Milwaukee, WI), whose axial field of view was 15.2 cm and spatial resolution 4.9 mm of full-width-half-maximum. Scans were performed using two-dimensional acquisition mode from the thigh to the skull base with seven bed positions. Each bed position was composed of 1 min of transmission scanning and 5 min of emission scanning.

FDG-PET/CT imaging was performed using a GE Discovery LS Scanner (General Electric Medical System, Milwaukee, WI) or a GE Discovery ST Scanner (the same manufacturer). The PET component of the GE Discovery LS Scanner was the same as that of the GE Advance Scanner. For the PET component of the GE Discovery ST Scanner, the axial field of view was 15.7 cm and the spatial resolution was 6.2 mm of full-width-half-maximum. PET scans were performed with both scanners using 2-dimensional acquisition mode from the thigh to the skull base with 7 bed positions. Each bed position was composed of 4 min of emission scanning. The CT component of both PET/CT scanners was a 16-row multi-detector CT scanner and CT images were acquired with a tube voltage of 140 kV, and the tube current was automatically set using the auto-exposure control function so that the number of standard deviations of noise was limited to 10. Attenuation correction of PET images was performed using the data from CT images.

Image reconstruction was performed using an ordered subsets expectation maximization (OSEM) algorithm with subset and iteration values of 14 and 2, respectively.

2.3. Image interpretation

All PET and CT images were interpreted by experienced radiologists and physicians. The 4.25 mm-thick images of axial, coronal and sagittal planes on hard copy films were reviewed. Uptake stronger than mediastinal blood pool activity was diagnosed as malignancy by the visual estimation. Symmetrical activities observed in both hilar regions were considered to be benign reactive changes. Any discrepancies between the radiologist and physician were resolved by discussion. The findings detected by FDG-PET were confirmed by other image tests or observation of the clinical course. FDG-PET was conducted after conventional staging procedures. CT, US and bone scans were interpreted without the FDG-PET findings. However, FDG-PET scan was interpreted in comparison with CT findings, while PET/CT findings were interpreted independently.

Table 1 Discrepancy between FDG-PET and conventional staging procedures (distant metastases)

Patient no.	Age (years)	Gender	CT N	PET N	PET M	Interval between conventional staging procedures and FDG-PET (days)	Comments
2	61	Male	2	2	1	20	Multiple bone metastases (PET)
6	68	Male	2	2	1	7	Lymph node metastasis around the cardia (PET)
47	61	Male	3	3	1	28	Multiple bone metastases (PET)
55	68	Male	2	2	1	20 (CT) and 14 (bone scan)	Liver, axillary lymph node, and iliac bone metastases (PET)
59	52	Male	3	3	1	13	Adrenal, cervical and mandibular lymph node metastases (PET)
63	59	Male	3	3	1	18 (CT) and 11 (bone scan)	Multiple bone and liver metastases (PET)

FDG, fluorodeoxyglucose; PET, positron emission tomography; CT, computed tomography; N, node; M, metastasis.

* Diagnosis of lymph node metastasis was not confirmed by other imaging modalities or observation of the clinical course.

3. Results

3.1. Detection of distant metastasis

FDG-PET showed results different from those of conventional staging procedures in 17 of 63 patients. PET scan demonstrated findings suggesting distant metastases in 6 of 63 patients (Table 1). The median interval between conventional staging procedures and FDG-PET was 16 days (range: 7–28). Abnormal uptake was observed around the cardia in one of these six patients (No. 6). A repeat FDG-PET study demonstrated a longer uptake stripe indicating radiation-induced oesophagitis and the diagnosis could not be established, as there was a remaining possibility of physiological uptake in the oesophagus. The diagnosis of metastatic disease was confirmed in the remaining five patients (8%, 95% confidence interval (CI): 3–18%). Among these five patients, four had bone metastases, two had liver metastases, one had adrenal metastasis, and two had lymph node metastases in the cervical or axillary region. The therapeutic strategy for these five patients was changed and they received only chemotherapy without thoracic radiotherapy. One patient (No. 47) had shown negative findings on bone scintigraphy four weeks before the FDG-PET study, but PET scan demonstrated increased FDG uptake in bones throughout the body. MRI of the spine confirmed the diagnosis of multiple bone metastases (Fig. 1). A repeat bone scan after three months detected obvious multiple bone metastases in No. 2 patient. Two hepatic lesions, as well as the primary tumour, mediastinal and hilar lymph nodes, had all increased in size after two cycles of chemotherapy in patient No. 55. A hepatic lesion, as well as the primary tumour, had decreased in size after two cycles of chemotherapy in patient No. 63. These hepatic lesions were compatible with liver metastases. Abnormal uptake by the right adrenal gland disappeared on repeat PET/CT after four cycles of chemotherapy in patient No. 59. Abnormal uptake in primary and mediastinal lesions was extremely decreased in

this patient. The right adrenal gland lesion was compatible with metastasis.

FDG-PET detected liver metastasis in one of 44 patients staged by CT scan of the abdomen (No. 55), and liver or adrenal metastasis in two of 19 patients staged by US (Nos. 59 and 63). Liver and adrenal metastases not detected by US were small, such that the CT part of PET/CT could not detect them as metastases. Ratios of upstaging by FDG-PET between initial CT scan and US of the abdomen were not statistically significant (1/44 versus 2/19, $P=0.214$).

3.2. Detection of regional lymph node metastases

FDG-PET scans detected regional lymph node metastases that had been negative on CT scans in nine patients (14%) (Table 2). The median interval between CT of the chest and FDG-PET was 19 days (range: 7–34). FDG-PET scans newly detected ipsilateral supraclavicular lymph node metastasis in four patients, contralateral lymph node metastasis in three, and mediastinal lymph node metastasis in two. These nine patients all underwent curative chemoradiotherapy, and abnormal FDG uptake in mediastinal and/or supraclavicular lymph nodes disappeared or decreased on repeat PET scans after chemoradiotherapy. These lymph nodes were considered positive for metastasis.

CT scan detected swollen mediastinal lymph nodes without abnormal FDG uptake in two patients. One patient had a past history of pulmonary tuberculosis complicated by pulmonary fibrosis. The swollen pretracheal lymph node was considered negative for metastasis because the node size remained unchanged after four cycles of chemotherapy although the primary tumour shrank. This case showed false positive findings on CT whereas FDG-PET correctly diagnosed the extent of disease (No. 43). The other patient had atelectasis of the right middle lobe due to the primary tumour. Superior mediastinal and subcarinal lymph nodes were considered to be metastatic on CT, but abnormal FDG uptake was absent. After three cycles of chemotherapy the




Fig. 1 A 61-year-old man with small-cell lung cancer. Bone scintigraphy was negative for osseous metastasis (a). However, PET scan demonstrated increased FDG uptake in bones throughout the body (b). MRI of the spine confirmed multiple bone metastases (c).

mediastinal lesion showed no change although the primary tumour had decreased in size and atelectasis of the right middle lobe was improved. The mediastinal lymph nodes were considered negative for metastasis (No. 61).

4. Discussion

SCLC tends to disseminate early in the disease course and displays a more aggressive clinical behaviour than NSCLC. Local treatment modalities alone such as radiotherapy or surgery are not effective in prolonging survival beyond a few weeks. Systemic chemotherapy is the mainstay of treatment for patients in all stages of SCLC. A combination of chemotherapy and thoracic irradiation can promote long-term survival for patients diagnosed as having limited disease and recent clinical trials of chemoradiotherapy for LD-SCLC obtained 5-year survival rates of 24–26% [2,3]. However, thoracic irradiation might cause severe radiation pneumonitis, resulting in respiratory failure and/or treatment-related death. Furthermore, thoracic irradiation might also cause oesophagitis which worsens patient quality of life. Accurate clinical staging is important to determine the indications for chemoradiotherapy in SCLC. Our study demonstrated that FDG-PET scan detected unsuspected distant metastases in 8% of patients with LD-SCLC based on conventional staging procedures and that the detection of these new lesions changed their therapeutic strategies. Furthermore, FDG-PET scan detected regional lymph node

metastases which had not been visualized on CT scan in 14% of patients. The radiation field could be appropriately set to cover the positive nodes based on the PET study results. Our results reconfirmed those of a previous preliminary study with a smaller number of patients [9].

Is the rate of the detection of unsuspected distant metastases (8%) clinically significant? Previous studies demonstrated that FDG-PET scan detected unsuspected distant metastases in 24% of patients with stage III NSCLC [6,7]. Compared to this result, the impact of FDG-PET on the staging of SCLC seems to be weaker. SCLC tends to have more obvious distant metastases than NSCLC, because of the aggressive biological behaviour of SCLC. Therefore, FDG-PET might detect unsuspected distant metastases at a relatively low rate. The most common region for unsuspected PET-detected metastasis in NSCLC was the abdomen, with 53% of patients having adrenal, liver, and other lesions [6]. In our study, FDG-PET detected bone metastases in four of five patients who were upstaged from LD to ED. These lesions might reflect metastasis to the bone marrow, although no pathological evidence was obtained, because neither bone marrow biopsy nor aspiration cytology was routinely conducted for the initial clinical staging.

Our retrospective analyses have several limitations. We did not confirm histologically regional lymph node or distant metastases detected by FDG-PET or CT. These lesions were not routinely biopsied and most metastatic lesions were chemosensitive and radiosensitive. Our confirmation was inevitably based on observation of the clinical course.

Table 2 Disagreement between FDG-PET and conventional staging procedures (regional lymph node metastases)

Patient no.	Age (years)	Gender	CT N	PET N	PET M	Interval between CT scan of the chest and FDG-PET (days)	Comments
1	63	Male	3	3	0	8	Contralateral supraclavicular lymph node metastasis (PET)
5	64	Female	1	2	0	34	Subcarinal lymph node metastasis (PET)
16	71	Male	3	3	0	7	Contralateral supraclavicular lymph node metastasis (PET)
20	69	Male	3	3	0	20	Ipsilateral supraclavicular lymph node metastasis (PET)
25	60	Male	3	3	0	27	Ipsilateral supraclavicular lymph node metastasis (PET)
30	66	Male	2	2	0	7	Pretracheal lymph node metastasis (PET)
33	72	Male	3	3	0	13	Ipsilateral supraclavicular lymph node metastasis (PET)
41	49	Female	3	3	0	19	Contralateral supraclavicular lymph node metastasis (PET)
43	73	Male	2	0	0	34	False-positive pretracheal lymph node metastasis (CT)
56	48	Female	3	3	0	11	Ipsilateral supraclavicular lymph node metastasis (PET)
61	74	Male	2	0	0	27	False-positive superior mediastinal and subcarinal lymph nodes (CT).

FDG, fluorodeoxyglucose; PET, positron emission tomography; CT, computed tomography; N, node; M, metastasis.

We employed no special strategies to reduce the bias of PET readers. PET readers might have reported in such a way as to reduce or increase the impact of PET. One-third of patients received FDG-PET after commencement of chemotherapy. However, the median interval between commencement of chemotherapy and FDG-PET was 4 days (range: 1–11 days). We considered the chemotherapy to have had no effects on the findings of FDG-PET in such a short time after the initiation of chemotherapy.

FDG-PET is expected to have the potentially to both up- and downstage patients with SCLC as well as NSCLC. A previous study demonstrated that FDG-PET correctly downstaged ED to LD in three of 120 patients with SCLC [10]. These three patients had adrenal swelling on CT scan, but these lesions were negative on FDG-PET. On the other hand, FDG-PET correctly upstaged LD to ED in 10 of 120 patients with SCLC. It seems that SCLC seldom has a solitary distant metastasis because of its aggressive clinical behaviour. Most ED-SCLC has multiple, not solitary, or obvious distant metastasis. Furthermore, the health insurance system does not allow patients who obviously have metastatic lung cancer to receive FDG-PET in Japan. Therefore, we did not include

patients with ED-SCLC in our analysis. Needless to say, FDG-PET is considered to be useful in patients with possible, but not evident, distant metastasis on other imaging tests, such as a solitary adrenal swelling.

According to the VALSG system, LD-SCLC is defined as a tumour confined to one hemithorax and regional lymph nodes [1]. Contralateral hilar or contralateral supraclavicular nodal involvement was classified as ED. According to the International Association for the Study of Lung Cancer (IASLC) consensus report, the classification of LD-SCLC includes bilateral hilar and/or supraclavicular nodal involvement, and ipsilateral pleural effusion [18]. A previous retrospective study demonstrated that the IASLC staging criteria for SCLC patients had a higher prognostic impact than VALSG criteria [19]. Therefore, we adopted the IASLC staging criteria for SCLC in our study.

In conclusion, FDG-PET scans detected unsuspected distant metastases in five of 63 patients with LD-SCLC (95% CI: 3–18%) and these findings resulted in a change of therapeutic strategies in these five patients. FDG-PET scans also detected contralateral supraclavicular lymph node metastases that had been negative on CT scans in three other

patients. These additional findings facilitated setting appropriate irradiation fields. FDG-PET scan is recommended as an initial staging tool in patients with apparent LD-SCLC.

Conflict of interest

The authors certify that there are no potential conflicts of interest.

Acknowledgments

We thank Takashi Sato, R.T., Hideaki Kitamura, R.T., Kazumasa Inoue, R.T. and Akira Hirayama, R.T. and Yoshiaki Kojima, Ph.D. for technical assistance. This work was supported in part by a Grant from the Ministry of Health, Labour, and Welfare for the third term Comprehensive Strategy for Cancer Control and a Grant-in-Aid for Cancer Research from the Ministry of Health, Labour, and Welfare, Japan.

References

- [1] Zelen M. Keynote address on biostatistics and data retrieval. *Cancer Chemother Rep* 1973;3(4):31-42.
- [2] Takada M, Fukuoka M, Kawahara M, Sugiura T, Yokoyama A, Yokota S, et al. Phase III study of concurrent versus sequential thoracic radiotherapy in combination with cisplatin and etoposide for limited-stage small-cell lung cancer: results of the Japan Clinical Oncology Group Study 9104. *J Clin Oncol* 2002;20:3054-60.
- [3] Turrisi 3rd AT, Kim K, Blum R, Sause WT, Livingston RB, Komaki R, et al. Twice-daily compared with once-daily thoracic radiotherapy in limited small-cell lung cancer treated concurrently with cisplatin and etoposide. *N Engl J Med* 1999;340:265-71.
- [4] Noda K, Nishiwaki Y, Kawahara M, Negoro S, Sugiura T, Yokoyama A, et al. Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N Engl J Med* 2002;346:85-91.
- [5] Pfister DG, Johnson DH, Azzoli CG, Sause W, Smith TJ, Baker Jr S, et al. American Society of Clinical Oncology treatment of unresectable non-small-cell lung cancer guideline: update 2003. *J Clin Oncol* 2004;22:330-53.
- [6] MacManus MP, Hicks RJ, Matthews JP, Hogg A, McKenzie AF, Wirth A, et al. High rate of detection of unsuspected distant metastases by pet in apparent stage III non-small-cell lung cancer: implications for radical radiation therapy. *Int J Radiat Oncol Biol Phys* 2001;50:287-93.
- [7] Eschmann SM, Friedel G, Paulsen F, Reimold M, Hehr T, Scheiderbauer J, et al. Impact of staging with ^{18}F -FDG-PET on outcome of patients with stage III non-small cell lung cancer: PET identifies potential survivors. *Eur J Nucl Med Mol Imaging* 2007;34:54-9.
- [8] Kalff V, Hicks RJ, MacManus MP, Binns DS, McKenzie AF, Ware RE, et al. Clinical impact of ^{18}F fluorodeoxyglucose positron emission tomography in patients with non-small-cell lung cancer: a prospective study. *J Clin Oncol* 2001;19:111-8.
- [9] Bradley JD, Dehdashti F, Mintun MA, Govindan R, Trinkaus K, Siegel BA. Positron emission tomography in limited-stage small-cell lung cancer: a prospective study. *J Clin Oncol* 2004;22:3248-54.
- [10] Brink I, Schumacher T, Mix M, Ruhland S, Stoelben E, Digel W, et al. Impact of ^{18}F FDG-PET on the primary staging of small-cell lung cancer. *Eur J Nucl Med Mol Imaging* 2004;31:1614-20.
- [11] Blum R, MacManus MP, Rischin D, Michael M, Ball D, Hicks RJ. Impact of positron emission tomography on the management of patients with small-cell lung cancer: preliminary experience. *Am J Clin Oncol* 2004;27:164-71.
- [12] Chin Jr R, McCain TW, Miller AA, Dunagan DP, Acostamadiedo J, Douglas Case L, et al. Whole body FDG-PET for the evaluation and staging of small cell lung cancer: a preliminary study. *Lung Cancer* 2002;37:1-6.
- [13] Hauber HP, Bohuslavizki KH, Lund CH, Fritscher-Ravens A, Meyer A, Pforte A. Positron emission tomography in the staging of small-cell lung cancer: a preliminary study. *Chest* 2001;119:950-4.
- [14] Kamel EM, Zwahlen D, Wyss MT, Stumpe KD, von Schulthess GK, Steinert HC. Whole-body ^{18}F -FDG PET improves the management of patients with small cell lung cancer. *J Nucl Med* 2003;44:1911-7.
- [15] Pandit N, Gonen M, Krug L, Larson SM. Prognostic value of ^{18}F FDG-PET imaging in small cell lung cancer. *Eur J Nucl Med Mol Imag* 2003;30:78-84.
- [16] Schumacher T, Brink I, Mix M, Reinhardt M, Herget G, Digel W, et al. FDG-PET imaging for the staging and follow-up of small cell lung cancer. *Eur J Nucl Med* 2001;28:483-8.
- [17] Shen YY, Shiau YC, Wang JJ, Ho ST, Kao CH. Whole-body ^{18}F -2-deoxyglucose positron emission tomography in primary staging small cell lung cancer. *Anticancer Res* 2002;22:1257-64.
- [18] Stahel RA, Ginsberg R, Havemann K, Hirsch FR, Ihde DC, Jassem J, et al. Staging and prognostic factors in small cell lung cancer: a consensus report. *Lung Cancer* 1989;5:119-26.
- [19] Micke P, Faldum A, Metz T, Beeh KM, Bittinger F, Hengstler JG, et al. Staging small cell lung cancer: veterans administration lung study group versus international association for the study of lung cancer—what limits limited disease? *Lung Cancer* 2002;37:271-6.

Review

Problems with Registration-Directed Clinical Trials for Lung Cancer in Japan

IKUO SEKINE,¹ HIROSHI NOKIHARA,¹ NOBORU YAMAMOTO,¹ HIDEO KUNITOH,¹
YUICHIRO OHE,¹ NAGAIHIRO SAJO² and TOMOHIDE TAMURA¹

¹Division of Internal Medicine and Thoracic Oncology, National Cancer Center Hospital,
Tokyo, Japan

²Division of Internal Medicine, National Cancer Center Hospital East, Kashiwa, Japan

SEKINE, I., NOKIHARA, H., YAMAMOTO, N., KUNITOH, H., OHE, Y., SAJO, N. and TAMURA, T. *Problems with Registration-Directed Clinical Trials for Lung Cancer in Japan*. Tohoku J. Exp. Med., 2007, 213 (1), 17-23 — New anticancer agents against lung cancer are needed because efficacy of chemotherapy is limited. The long time required, low quality, and considerable costs of registration-directed clinical trials in Japan ("Chicken") have been pointed out. The quality of 24 phase I and 41 phase II trials of an anticancer drug for lung cancer were analyzed according to the approval year of the drug. The human resources and infrastructure to support oncology clinical practice and clinical trials were compared between Japan and the USA. A maximum tolerated dose was not defined in any of seven phase I trials before 1989, and was determined in two of six trials between 1989 and 1996 and in seven of 10 trials thereafter. Before 1989, 29 (20%) of 142 patients registered in two trials were ineligible, and the number of ineligible patients was not reported in the five trials. Sample size calculations were not performed in any of seven phase II trials before 1989 and were performed in only four of 10 trials between 1989 and 1996 and in all 23 trials conducted thereafter. The shortage of human resources, including medical oncologists, oncology nurse practitioners and clinical research coordinators, is serious and acute. The infrastructure to support clinical trials also remains insufficient in Japan. In conclusion, registration-directed clinical trials of anticancer agents have advanced significantly during last three decades but remain unsatisfactory. The development of infrastructure and human resources is an urgent task to ensure high-quality clinical trials without unnecessary delays. ——— clinical trials; medical oncologists; nurse practitioners; lung cancer; anticancer agents

© 2007 Tohoku University Medical Press

Lung cancer is one of the most common malignancies and the leading cause of cancer-related deaths in many countries. In the year 2000, the annual number of deaths from lung cancer was estimated to be 1.1 million worldwide,

and global lung cancer incidence is increasing at a rate of 0.5% per year (Schottenfeld and Searle 2005). About 80% of patients with lung cancer have already developed distant metastases or pleural effusion, either by the time of the initial

Received June 13, 2007; revision accepted for publication July 11, 2007.

Correspondence: Ikuo Sekine, Division of Internal Medicine and Thoracic Oncology, National Cancer Center Hospital, Tsukiji 5-1-1, Chuo-ku, Tokyo 104-0045, Japan.
e-mail: isekine@nce.go.jp

diagnosis or by the time recurrence is detected after surgery for local disease. These patients can be treated with systemic chemotherapy, but the efficacy of currently available anticancer agents is limited to the extent that patients with advanced disease rarely live long. Therefore, new chemotherapeutic agents continue to be developed against lung cancer (Sekine and Saijo 2000).

The Japanese Pharmaceutical Affairs Law (PAL) was enacted in 1948, and was first amended in 1960 to provide for regulations to ensure the maintenance of the quality, efficacy, and safety of drugs and medical devices, and to promote research and development of these medical and pharmaceutical products. Good Clinical Practice (GCP) was enforced by the Bureau Notification of the Ministry of Health and Welfare of Japan ("Kyokuchou-Tsuuchi") in 1989 (the former GCP). In 1996, the PAL and its related laws were amended to strengthen GCP (the new GCP), Good Laboratory Practice, Good Post-Marketing Surveillance Practice, and standard compliance

reviews, conforming to the International Conference on Harmonization of Technical Requirements for Registration of Pharmaceuticals for Human Use. In contrast to the laws prevailing in the US and EU, marketing approval for anticancer agents in Japan has been granted based on reports of the anti-tumor effects of the new agents in phase II trials (Fujiwara and Kobayashi 2002).

Under this Japanese drug approval system regulated by the PAL, 23 anticancer drugs have been approved for use against lung cancer during the last five decades (Fig. 1). Of these, 9 drugs are original to Japan, some of which are routinely used all over the world. Several problems, however, have been pointed out in registration-directed clinical trials in Japan ("Chiken"), including the long time required, low quality, and considerable cost (The Ministry of Health, Labour and Welfare of Japan 2002; The Ministry of Education, Science and Culture and the Ministry of Health, Labour and Welfare 2003). As a result, Japanese cancer patients must wait for a long time




Fig. 1. Anticancer drugs approved for lung cancer in Japan. Bold: original to Japan. Dotted line: case series studies, solid thick line: investigational new drug phase I-II trials for approval, and dotted thin line: post-marketing sponsored phase III trials. Vertical dotted lines indicate the year when the former and new GCP were issued.

until they receive new anticancer drugs which have been approved long before in other countries (The Ministry of Health, Labour and Welfare of Japan 2005). We discuss the aspects and issues of registration-directed trials in Japan by reviewing such trials for the 23 anticancer drugs.

Review of registration-directed clinical trials for the 23 anticancer drugs

A total of 65 phase I and II trials of an anticancer drug for approval were reviewed in terms of definition of eligibility criteria, maximum tolerated dose (MTD), sample size, response criteria, and extramural review for tumor responses. The MTD is the dose associated with serious but reversible toxicities in a sizeable proportion of patients and the one that offers the best chance for a favorable therapeutic ratio (Piantadosi 1997). The number of patients accrued in a trial, percentage of ineligible patients, number of participant hospitals in a trial, and the study period defined as the months between the first and last patient accrual were also analyzed. They were obtained from a published paper for 53 trials, from a meeting abstract and in-company resource for one trial, and from in-company resource alone for the remaining 11 trials. The clinical developmental period of an anticancer drug was defined as years between the start month of the first phase I trial and the month of the approval for lung cancer.

These parameters are compared according to the approval year of the drug. We categorized three periods of approval: 1) before 1989, 2) between 1989 and 1996, and 3) between 1997 and 2004, because the former GCP was enforced in 1989, and the new GCP in 1997 (Fujiwara et al. 2002).

Of the 23 anticancer drugs, six drugs whose clinical development started before 1974 were approved on the basis of the clinical experience of the use of the drug without clinical trials (Fig. 1). A total of 24 phase I trials were identified (Table 1). The MTD was not defined in the protocol of any trials before 1989, but was defined in 33% of trials between 1989 and 1996, and in 70% of trials after 1996. Instead of the MTD, maximum acceptable dose, defined as the dose associated with grade 2 or severer toxicity in two thirds or more patients, was used in a trial after 1996. About twice more patients were registered in a trial before 1989 than thereafter, but 20% of the registered patients before 1989 were ineligible. The study period of a phase I trial got longer as the number of participant hospitals decreased, from 7 months and 11 hospitals before 1989 to 13 months and 4 hospitals after 1996, respectively.

In this review, 41 phase II trials for approval were analyzed (Table 2). Calculation of the sample size was not made in any trials before 1989, was seen in 40% of trials between 1989 and 1996, and in all trials thereafter. Response criteria were

TABLE 1. Investigational new drug phase I trials for approval.

	Before 1989	1989-1996	1997 or thereafter
Total number of trials	7	6	11
Defined, number (%) of trials			
Eligibility criteria	4 (57)	6 (100)	11 (100)
Maximum tolerated dose*	0 (0)	2 (33)	7 (70) [†]
Results of trials, median (range)			
Number of patients**	61 (32-170)	24 (18-54)	29 (9-43)
% of ineligible patients	20 (20-21) [†]	8 (0-33)	6 (0-22)
Number of hospitals	11 (1-21)	9 (1-18)	4 (1-17)
Study period in months	7 (5-30)	10 (5-11)	13 (8-24)

*Statistically significant difference obtained ($p = 0.014$ by the chi-square test); **Statistically significant difference obtained ($p < 0.01$ by the Kruskal Wallis test); [†]Data were available in 2 trials only; [‡]Data were available in 10 trials only.

TABLE 2. Investigational new drug phase II trials for approval.

	Before 1989	1989-1996	1997 or thereafter
Total number of trials	7	11	23
Defined, number (%) of trials			
Eligibility criteria	4 (57)	11 (100)	23 (100)
Sample size calculation*	0 (0)	4 (40) [†]	23 (100)
Response criteria	6 (86)	11 (100)	23 (100)
Extramural review	3 (43)	9 (82)	23 (100)
Results of trials, median (range)			
Number of patients	71 (10-127)	68 (18-153)	61 (11-102)
% of ineligible patients	18 (0-29) [†]	3 (0-22)	3 (0-12)
Number of hospitals	27 (3-103)	17 (1-30)	20 (5-46)
Study period in months	18 (12-36)	12 (6-34)	26 (4-48) [‡]

*Statistically significant difference obtained ($p < 0.01$ by the chi-square test); [†]Data were available in 5 trials only; [‡]Data were available in 10 trials only; [§]Data were available in 22 trials only.

defined in almost all studies, but an extramural review was conducted only after 1989. The median number of registered patients in a trial was constant through the three periods, but the percentage of ineligible patients was high in trials conducted before 1989. The number of patients in a trial, and the number of hospitals in a trial were similar regardless of the year. The median study period in recent trials was 26 months.

The clinical development period was evaluated in the 23 drugs. Cisplatin was approved for germ cell tumors in 1983 and additionally approved for non-small cell lung cancer (NSCLC) in 1986. S-1 was firstly approved for gastric cancer in 1999, and additionally approved for NSCLC in 2004. The other drugs were approved for lung cancer for the first time. The median (range) clinical development period was 5.2 (3.2-14.5) years before 1989, 6.0 (4.8-9.1) years between 1989 and 1996, and 9.0 (3.9-15.4) years in 1997 or thereafter.

Development and recent problems of phase I and phase II trials in Japan

The concept of the "clinical trial" was not widely followed in Japan until 1974, when a phase I trial of nimustine hydrochloride (ACNU) was launched as one part of the United States-Japan Cooperation Cancer Research Program on

the basis of the agreement between the National Cancer Institute and Japan Society for the Promotion of Science (Sugano 1982; Niitani 1999). Phase I trials before 1989 required the accrual of many patients, because 1) the maximum tolerated dose was not defined, 2) many patients were treated at unnecessary dose levels because the modified Fibonacci dose escalation schedule was not applied, and 3) the percentage of ineligible patients was high. Some of these issues were improved in 1997 or thereafter, but the maximum tolerated dose is still not defined in as many as 40% of trials. Recently, oncology phase I trials came to be conducted among fewer hospitals than before, as more participants were recruited in each hospital. This facilitated communication among phase I investigators, which is important to complete phase I trials safely.

Phase II trials play the central role in anti-cancer agent approval in Japan, because the approval can be granted based on the response rate in these trials. The quality of protocols for phase II trials suggested by eligibility criteria, sample size calculation, response criteria, and extramural review has been improved significantly. The study period of phase II trials, however, was and is still too long, as long as 4 years in recent trials. To increase participant hospitals, however, is not necessarily a desirable solution,