

lower (dorsomedial) nerve tissue. The space between the pelvic plexus and the paracolpium was developed anteroposteriorly by using Kelly forceps or Metzenbaum scissors. From this approach, the pelvic plexus was put away laterally. Then the uterine branch of the pelvic plexus was cut, which enabled dissection of the paracolpium without involving the pelvic plexus.

For the five patients of group B, the nerve-sparing procedure could not be successfully performed due to bleeding from the paracolpium and/or the surgeon's inexperience, although we had an intension to preserve the nerves. The space between the pelvic plexus and the paracolpium could not be developed, and therefore, the pelvic plexus could not be put aside. The vaginal canal was cut without selective dissection of the uterine branches of the pelvic plexus for the cases of group B.

#### Postoperative treatment

Postoperative whole-pelvic external radiation therapy (50 Gy) was employed when there was lymph node metastasis or histologically confirmed parametrial invasion. When the tumor had invaded the lymphatic or vascular channels, we treated the patient with cisplatin-based chemotherapy for 3-6 cycles, unless the patient refused this treatment. The follow-up period ranged from 12 to 48 months (median: 29 months).

#### Urodynamic study

Urodynamic studies were carried out before the operation and at 1 month and, 3, 6, and 12 months after the operation. The urodynamic studies were carried out using a Urolab spectrum™ (Life-Tech Inc., Stafford, TX) by the following procedure. A 7F double-lumen catheter for cystometry (DLC-7D; Life-Tech Inc.) was placed in the urethra after collecting residual urine, and a balloon catheter for measuring rectal internal pressure (RPC-9D; Life-Tech Inc.) was inserted into the rectum. While the patient was sitting on a toilet seat, saline solution was continuously infused through the catheter into the urethra at a rate of 30 mL/min. Maximum cystometric capacity was defined as the capacity at the time of the strongest desire to void, the time when 700 mL of saline solution had been infused in the patients having no desire to void, or the time of persistent urinary incontinence if urinary incontinence occurred before the desire to void. Vesical volume, vesical pressure, Pabd, and urinary flow volume were monitored simultaneously. Detrusor contraction pressure was automatically calculated by subtracting Pabd from vesical pressure. Urodynamic evaluators were bladder compliance (Cves) at

the moment of strong desire to void, maximum flow rate (MFR), detrusor contraction pressure at maximum flow (Pdet Qmax), abdominal pressure at maximum flow (Pabd Qmax), and residual urine volume. The problem in analysis of the voiding phase is that there is a lack of data for patients in whom voiding is impossible due to psychologic stress. Since no data for Pabd Qmax and Pdet Qmax could be obtained from some of the present patients in the early postoperative period due to voiding incompetence, these pressures were analyzed preoperatively and at 6 and 12 months postoperatively. Maximum urinary flow rate was taken as 0.0 mL/sec for patients who were not able to void in tests carried out at 1 month and at 3 months after the operation. Bladder sensation was categorized into four levels: (1) normal: a patient felt a desire to void when 200 mL or more of saline solution was infused; (2) increased: a patient felt a strong desire to void when less than 200 mL of saline solution was infused; (3) reduced: a patient merely felt an obscure tension of the bladder when 400 mL or more of saline solution was infused; and (4) absent: a patient did not feel a tension of the bladder when 400 mL or more of saline solution was infused or persistent urinary incontinence occurred before the desire to void. All the terms and symbols used for parameters in this paper were taken from the 1988 report by the International Continence Society.

#### Statistical analysis

Categorical variables were analyzed using the  $\chi^2$  test or Fisher exact test. Median values of continuous variables were compared by the Mann-Whitney *U* test. Whether our procedure had an impact on each parameter or not was analyzed by repeated measure one-way analysis of variance (ANOVA), and significance in differences between time points was examined using Fisher's protected least significant difference test when significant differences were indicated by the results of repeated measure one-way ANOVA. The comparison between group A and group B was analyzed by repeated measure two-way ANOVA. A result was deemed significant when  $P < 0.05$ .

#### Results

##### Comparison of the results of urodynamic study between patients with and without successful autonomic nerve preservation

Results of urodynamic studies performed 110 times in group A are shown in Table 2. Data were obtained

**Table 2.** Urodynamic values of 22 patients in whom autonomic nerves had been preserved

	Number of cases	Before surgery (mean $\pm$ SD)	After surgery (months)			
			1	3	6	12
<b>Storage phase</b>						
Cves at SDV (mL/cmH <sub>2</sub> O)	22	64.0 $\pm$ 63.4	12.9 $\pm$ 14.2*	31.4 $\pm$ 39.1*	41.3 $\pm$ 45.9*	67.8 $\pm$ 71.8
<b>Voiding phase</b>						
MFR (mL/sec)	21	19.6 $\pm$ 8.5	8.9 $\pm$ 7.3*	10.2 $\pm$ 7.2*	13.8 $\pm$ 8.7*	16.5 $\pm$ 10.1
Pabd Qmax (cmH <sub>2</sub> O)	20	5.5 $\pm$ 24.2			19.4 $\pm$ 24.4*	13.9 $\pm$ 22.5*
Pde Qmax (cmH <sub>2</sub> O)	20	38.1 $\pm$ 17.1			26.5 $\pm$ 17.4*	26.8 $\pm$ 14.7*
Residual urine	22	6.8 $\pm$ 10.6	126.9 $\pm$ 125.2*	90.5 $\pm$ 115.2*	34.6 $\pm$ 58.9*	16.5 $\pm$ 34.4

SDV, strong desire to void.

\* $P < 0.05$  compared to preoperative value.

from all the patients in the storage phase, but data for three patients (one patient who could not void in the preoperative test, one patient for whom a problem occurred in testing before the operation, and one patient for whom a problem occurred in testing at 12 months after the operation) were not included in the analysis in the voiding phase. Data for two patients, one who could not void in the test carried out before the operation and one who could not void in the test carried out at 12 months after the operation, were not included in the analysis at the time of MFR. Results of urodynamic studies performed 25 times in group B are shown in Table 3.

There was no significant difference in Cves of group A between before the operation and at 12 months after the operation, but Cves of group B at 12 months after the operation was significantly lower than that before the operation. Repeated measure two-way ANOVA revealed that the adverse effect on Cves in group B was significantly more than that in group A ( $P < 0.05$ , Fig. 1).

There was no significant difference in MFR of group A between before the operation and at 12 months after the operation, but MFR of group B at 12 months after the operation was significantly lower than that before the operation. There was no significant difference in MFR between groups A and B (Fig. 2).

The Pabd Qmax of either group A or group B at 12 months after the operation was significantly higher than that before the operation. The adverse effect on Pabd in group B was significantly more than that in group A ( $P < 0.05$ , Fig. 3).

The Pdet Qmax of either group A or group B at 12 months after the operation was significantly lower than that before the operation. The adverse effect on Pdet Qmax in group B was significantly more than that in group A ( $P < 0.01$ , Fig. 4).

The residual urine volume of group A was almost the same as the preoperative volume at 12 months after the operation. The adverse effect on residual urine volume in group B was significantly more than that in group A ( $P < 0.05$ , Fig. 5).

#### The effect of postoperative treatment on the results of postoperative urodynamic study

The radiation therapy was given to 3 of the 22 patients in group A postoperatively. Among the urodynamic variables, Cves seemed to be adversely affected by radiation therapy. Cves in 19 nonirradiated patients decreased to the nadir value at 1 month after operation and increased steadily thereafter during the follow-up period after operation and returned to preoperative value at 12 months after the operation. On

**Table 3.** Urodynamic values of five patients in whom autonomic nerves had not been preserved

	Number of cases	Before surgery (mean $\pm$ SD)	After surgery (months)			
			1	3	6	12
<b>Storage phase</b>						
Cves at SDV (mL/cmH <sub>2</sub> O)	5	163.8 $\pm$ 167.9	9.8 $\pm$ 7.6*	32.2 $\pm$ 19.7*	32.3 $\pm$ 28.4*	59.3 $\pm$ 36.9*
<b>Voiding phase</b>						
MFR (mL/sec)	5	17.0 $\pm$ 3.3	3.1 $\pm$ 3.6*	6.2 $\pm$ 7.2*	7.7 $\pm$ 6.8*	10.8 $\pm$ 5.0*
Pabd Qmax (cmH <sub>2</sub> O)	5	1.0 $\pm$ 18.1			21.8 $\pm$ 24.2	40.2 $\pm$ 27.5*
Pde Qmax (cmH <sub>2</sub> O)	5	44.0 $\pm$ 15.7			28.6 $\pm$ 20.0	8.6 $\pm$ 12.2*
Residual urine	5	3.6 $\pm$ 4.0	251.2 $\pm$ 227.0*	274.2 $\pm$ 245.6*	145.2 $\pm$ 136.5	88.2 $\pm$ 87.8

SDV, strong desire to void.

\* $P < 0.05$  compared to preoperative value.




Figure 1. Cves at strong desire to void. a) Preoperative and postoperative measurement in group A patients. b) Comparison of changes in measurement between groups A and B.

contrary, Cves in three irradiated patients further decreased after 1 month of operation and reached nadir value at 6 months after the operation. The Cves remained at decreased level even 12 months after the operation. The postoperative radiation was suggested to impair the recovery of Cves, although the difference between the two patients groups did not reach a statistically significant level ( $P = 0.087$ ).

## Discussion

The incidence of urinary tract injuries in radical hysterectomy has been decreasing in recent years due to improvements in surgical techniques. Much interest has recently been shown in autonomic nerve-sparing surgical techniques<sup>(2-6)</sup>.

There have been many studies on urodynamics after radical hysterectomy, but a consensus regarding urodynamics has not been reached. One reason for this is that surgical techniques used for radical hysterectomy have not been clearly described in many reports. Piver *et al.* proposed five classes for the extent of radical hysterectomy<sup>(7)</sup>, and this classification is often used now for discriminating types of operation. The surgical technique of Wertheim-Meigs is considered to cor-

respond to type III<sup>(8)</sup>. In radical hysterectomy, pelvic plexus is in close proximity to the paracolpium at the depth at which the vagina should be dissected. If separation of the pelvic plexus from the paracolpium is insufficient, the pelvic plexus will be injured at the time of amputation of vagina. Possover *et al.* recently reported that preservation of the pelvic splanchnic nerves and pelvic plexus, with the middle rectal artery serving as a landmark for identification, is important for preserving bladder function<sup>(6)</sup>. If the cardinal ligament below the middle rectal artery is dissected, the pelvic splanchnic nerves will be injured. Moreover, if the uterosacral ligaments are excised at their sacral attachments, the hypogastric nerves and the pelvic plexus may be also excised. Therefore, it is not clear in a simple statement that a type III operation was performed, whether hypogastric nerves, the pelvic splanchnic nerves, and the pelvic plexus were preserved or not. Thus, accurate evaluation of bladder dysfunction after radical hysterectomy is not possible without detailed information on the surgical procedure used.

Another important point in discussing urodynamics after radical hysterectomy is comparison of preoperative and postoperative test results. It has been reported





Figure 2. MFR. a) Preoperative and postoperative measurement in group A patients. b) Comparison of changes in measurement between groups A and B.

Figure 3. Pabd Qmax. a) Preoperative and postoperative measurement in group A patients. b) Comparison of changes in measurement between groups A and B.



that about 80% of patients with uterine cervical cancer already have some degree of bladder dysfunction before undergoing surgery<sup>(9)</sup>. It would not be possible to determine whether poor Cves is due to the operation or due to a dysfunction that already existed before the operation, if preoperative tests were not carried out. There has been criticism of studies in which urodynamic analysis was only carried out after the operation<sup>(10)</sup>. Other important points for urodynamic studies of urinary function after radical hysterectomy are to conduct tests on the subjects at the same time for comparison of results among the subjects and to carry out not only short-term but also long-term assessments. Voiding dysfunction symptoms after radical hysterectomy have been thought to disappear within 6–12 months after the operation<sup>(11,12)</sup>. However, urodynamic studies have revealed that there does exist underlying bladder dysfunction after 12 months of operation<sup>(13)</sup>.


In this study, the results showed that our technique does not cause a deterioration in Cves and MFR and an increase in residual urine volume at 12 months after the operation, although it does cause some amount of increase in Pabd Qmax and reduction in Pdet Qmax. Actually, there was no patient who complained of urinary difficulty at 12 months after the operation, which was confirmed by a urodynamic

study and a questionnaire. That is to say, radical hysterectomy combined with autonomic nerve preservation did not cause significant deterioration in QoL of our patients. There have been very few reports in which the surgical technique used and the timing of tests are clearly described, preoperative and postoperative values for all the patients are compared, and results of both short-term and long-term tests are presented. Such description and presentation of results appear in a report by Scotti *et al.*, but, unfortunately, type II operations were performed on half of their patients<sup>(14)</sup>. In our experience, bladder dysfunction is rarely a problem in a type II operation.

We employ our technique for autonomic nerve preservation to the uninvaded side for patients with stage IIB uterine cervical cancer. It is reported that the normal urinary function could be maintained when at least one side of the sympathetic nerve was preserved in experimental animals<sup>(15)</sup>. This data suggest that the normal urinary function can be maintained by applying the operation with autonomic nerve preservation to the noninvaded side in patients with stage IIB cervical cancer who had parametrial invasion only at one side.

Piver's type III (Wertheim–Meigs) operation is the treatment of choice for FIGO stages IB–IIA cervical cancer in Western countries. The standard treatment

Figure 4. Pde Qmax. a) Preoperative and postoperative measurement in group A patients. b) Comparison of changes in measurement between groups A and B.






Figure 5. Residual urine volume. a) Preoperative and postoperative measurement in group A patients. b) Comparison of changes in measurement between groups A and B.

for patients with stages IB–IIB cervical cancer in Japan is radical hysterectomy originally described by Okabayashi<sup>(16)</sup>. This surgery is more radical than Wertheim hysterectomy. According to the recent statistics regarding treatment of cervical cancer announced from the Japan Society of Obstetrics and Gynecology<sup>(17)</sup>, only 4% of patients with stage IIB cervical cancer received primary radiotherapy. Concurrent chemoradiation is not a standard treatment in Japan at this moment. The radiation therapy will cause permanent loss of ovarian function and sexual disturbance due to fibrous stenosis of vaginal canal, and these are quite detrimental for sexually active women. We have performed procedures for preserving ovarian function and preventing shortening of the vagina<sup>(8–21)</sup> in an attempt to maintain QoL of patients with cervical cancer treated with radical hysterectomy. Regarding chemoradiation, however, a large-scale clinical trial needs to be conducted in Japan too.

In conclusion, our aim seems to have been achieved because the urinary function, both subjectively and objectively, at 12 months after operation is almost normal.

### Acknowledgment

The authors wish to thank Dr Kenichi Oguchi for his contribution to introducing urodynamic analysis in the care of patients undergoing radical hysterectomy, without whom our attempts would not have been started and completed.

### References

- Ralph G, Tamussino K, Lichtenegger W. Urological complications following radical abdominal hysterectomy for cervical cancer. *Arch Gynecol Obstet* 1988;243:215–20.
- Yabuki Y, Asamoto A, Hoshihara T, Nishimoto H, Satou N. A new proposal for radical hysterectomy. *Gynecol Oncol* 1996;62:370–8.

- Kuwabara Y, Suzuki M, Hashimoto M, Furugen Y, Yoshida K, Mitsuhashi N. New method to prevent bladder dysfunction after radical hysterectomy for uterine cervical cancer. *J Obstet Gynaecol Res* 2000;26:1–8.
- Trimbos JB, Maas CP, DeRuiter MC, Peters AAW, Kenter GG. A nerve-sparing radical hysterectomy: guidelines and feasibility in Western patients. *Int J Gynecol Cancer* 2001;11:180–6.
- Hockel M, Konerding MA, Heussel CP. Liposuction-assisted nerve-sparing extended radical hysterectomy: oncologic rationale, surgical anatomy, and feasibility study. *Am J Obstet Gynecol* 1998;178:971–6.
- Posslover M, Stober S, Plaut K, Schneider A. Identification and preservation of the motoric innervation of the bladder in radical hysterectomy type III. *Gynecol Oncol* 2000;79:154–7.
- Piver MS, Rutledge F, Smith JP. Five classes of extended hysterectomy for women with cervical cancer. *Obstet Gynecol* 1974;44:265–72.
- Wertheim E. The extended abdominal operation for carcinoma uteri (based on 500 cases). *Am J Obstet Gynecol* 1912;66:169–232.
- Lin HH, Yu HJ, Sheu BC, Huang SC. Importance of urodynamic study before radical hysterectomy for cervical cancer. *Gynecol Oncol* 2001;81:270–2.
- Leveckis J, Boucher NR, Parys BT, Reed MW, Shorthouse AJ, Anderson JB. Bladder and erectile dysfunction before and after rectal surgery for cancer. *Br J Urol* 1995;76:752–6.
- Glahn BE. The neurogenic factor in vesical dysfunction following radical hysterectomy for carcinoma of the cervix. *Scand J Urol Nephrol* 1970;4:107–16.
- Green TH, Meigs JV, Ulfelder H et al. Urologic complications of radical Wertheim hysterectomy: incidence, etiology, management and preservation. *Obstet Gynecol* 1962;20:293–312.
- Sekido N, Kawai K, Akaza H. Lower urinary dysfunction as persistent complication of radical hysterectomy. *Int J Urol* 1997;4:259–64.
- Scotti RJ, Bergman A, Bhatia NN, Ostergard DR. Urodynamic changes in urethrovaginal function after radical hysterectomy. *Obstet Gynecol* 1986;68:111–20.
- Liang JT, Chien CT, Chang KJ et al. Neurophysiological basis of sympathetic nerve-preserving surgery for lower rectal cancer—a canine model. *Hepato-gastroenterology* 1998;45:2206–14.
- Okabayashi H. Radical abdominal hysterectomy for cancer of the cervix uteri. *Surg Gynecol Obstet* 1921;33:335–41.
- Kanazawa K. Annual Report on the Treatment for Cervical Cancer in 2000. Report of Gynecologic Cancer Committee. *Acta Obstet Gynecol Jpn* 2004;56:15–8. In Japanese.
- Tabata M, Ichinoe K, Sakuragi N, Shiina Y, Yamaguchi T, Mabuchi Y. Incidence of ovarian metastasis in patients with cancer of the uterine cervix. *Gynecol Oncol* 1987;28:255–61.
- Yamamoto R, Okamoto K, Todo Y et al. A study of risk factors for ovarian metastases in stage Ib–IIIb cervical carcinoma and analysis of ovarian function after a transposition. *Gynecol Oncol* 2001;82:312–6.
- Yamamoto R, Okamoto K, Ebina Y, Shirato H, Sakuragi N, Fujimoto S. Prevention of vaginal shortening following radical hysterectomy. *Br J Obstet Gynaecol* 2000;107:841–5.
- Okamoto K, Sakuragi N, Fujimoto S. Ovarian function and survival of patients with cervical carcinoma treated with radical hysterectomy and ovarian transposition. In: Leung PCK, Adashi EY, eds. *The ovary*. San Diego: Elsevier Academic Press, 2004:559–67.

Accepted for publication July 12, 2004

EDITORIAL

## Is paclitaxel/carboplatin really a useful regimen for ovarian cancer compared with platinum/doxorubicin/cyclophosphamide?

Article on page 221

### Historical control study of paclitaxel-carboplatin (TJ) versus conventional platinum-based chemotherapy (CAP) for epithelial ovarian cancer

R. Numazaki, E. Miyagi, R. Onose, T. Nakazawa, K. Sugiura, K. Asukai, H. Nakayama, A. Miyamatsu, N. Okamoto, F. Hirahara

In this issue of the *International Journal of Clinical Oncology*, Numazaki et al.<sup>1</sup> report the efficacy and toxicity of paclitaxel/carboplatin compared with platinum/doxorubicin/cyclophosphamide. In general, the purpose of this study and the direction of the discussion are well considered and documented, and we can know, in general, the efficacy and toxicity of practical treatments with paclitaxel/carboplatin in Japanese women with ovarian cancer. Unfortunately, three major problems in this study need to be clarified.

1. This is not a protocol study, but, rather, a comparison of practical treatments. On this point, the article seems confused.
2. As for the anti-tumor effects of paclitaxel/carboplatin reported, they neither fit the response evaluation criteria in solid tumors (RECIST) or the criteria set by the Japan Society of Clinical Oncology (JSCO), but are evaluated by the authors' specific criteria. Although all the patients with disease stages III/IV were evaluated for effects, in terms of lesions on images, it cannot be considered as common sense that all patients with III/IV disease after standard surgery will have measurable lesions, as was described in this article.
3. Because there were differences in the use of granulocyte-colony-stimulating factor (G-CSF) and a 5-HT<sub>3</sub> antagonist over time, a toxicity comparison of the groups cannot be done. In fact, there were no treatment discontinuations in the group with paclitaxel-carboplatin (TJ) which showed severe blood toxicity (grade 3/4), whereas there were patients who discontinued the platinum-based chemotherapy (CAP).

I feel that this article could be more useful if a study of the effects of TJ treatment and toxicity, and the results in subjects with stage III/IV cancers were to be addressed and compared with the results of past (CAP) therapy, which has been described in the Discussion section.

### References

1. Numazaki R, Miyagi E, Onose R, et al. (2006) Historical control study of paclitaxel-carboplatin (TJ) versus conventional platinum-based chemotherapy (CAP) for epithelial ovarian cancer. *Int J Clin Oncol* 11:221-228
2. McGuire WP, Hoskins WJ, Brady MF, et al. (1996) Cyclophosphamide and cisplatin compared with paclitaxel and cisplatin in patients with stage III and stage IV ovarian cancer (GOG 111). *N Engl J Med* 334:1-6
3. Piccart MJ, Bertelsen K, James K, et al. (2000) Randomized intergroup trial of cisplatin-paclitaxel versus cisplatin-cyclophosphamide in women with advanced epithelial ovarian cancer: 3-year results (OV-10). *J Natl Cancer Inst* 92:699-708
4. The International Collaborative Ovarian Neoplasm (ICON) Group (2002) Paclitaxel plus carboplatin versus standard chemotherapy with either single-agent carboplatin or cyclophosphamide, doxorubicin, and cisplatin in women with ovarian cancer: the ICON3 randomised trial. *Lancet* 360:505-515
5. Ozols RF, Bundy BN, Greer BE, et al. (2003) Phase III trial carboplatin and paclitaxel compared with cisplatin and paclitaxel in patients with optimally resected stage III ovarian cancer: a Gynecologic Oncology Group study (GOG 158). *J Clin Oncol* 21:3194-3200
6. Du Bois A, Luck HJ, Meier W, et al. (2003) A randomized clinical trial of cisplatin/paclitaxel versus carboplatin/paclitaxel as first-line treatment of ovarian cancer (AGO). *J Natl Cancer Inst* 95:1320-1329
7. Vasey PA, Jayson GC, Gordon A, et al. (2004) Phase III randomized trial of docetaxel-carboplatin versus paclitaxel-carboplatin as first-line chemotherapy for ovarian carcinoma (SCOTROC). *J Natl Cancer Inst* 96:1682-1691
8. Yamamoto R, Minobe S, Kaneuchi M, et al. (2002) A phase I/II study of carboplatin and paclitaxel in patients with epithelial ovarian cancer. *Jpn J Clin Oncol* 32:128-134
9. Takei Y, Suzuki M, Ohwada Y, et al. (2003) A feasibility study of paclitaxel and carboplatin therapy in Japanese patients with epithelial ovarian cancer. *Oncol Rep* 10:951-955

Toru Sugiyama, M.D.  
Iwate Medical University School of Medicine,  
19-1 Uchimarui, Morioka, Iwate 020-8505, Japan  
Tel. +81-19-651-5111; Fax +81-19-622-1900  
e-mail: sugiyama@iwate-med.ac.jp

## Treatment results of adjuvant chemotherapy after radical hysterectomy for intermediate- and high-risk stage IB–IIA cervical cancer

Nobuhiro Takeshima\*, Kenji Umayahara, Kiyoshi Fujiwara, Yasuo Hirai,  
Ken Takizawa, Katsuhiko Hasumi

Department of Gynecology, Cancer Institute Hospital, 3-10-6, Ariake, Koto-Ku, Tokyo 135-8550, Japan

Received 28 January 2006

Available online 14 June 2006

### Abstract

**Objective.** To determine the effectiveness of chemotherapy alone as postoperative adjuvant therapy for intermediate- and high-risk cervical cancer.

**Methods.** The study group comprised of 65 consecutive patients with stage IB or IIA squamous cell or adenosquamous cervical cancer who were initially treated with radical hysterectomy and pelvic lymphadenectomy between 1993 and 2002. Tumors were of intermediate-risk (stromal invasion > 50%,  $n = 30$ ) or high-risk (positive surgical margin, parametrial invasion, and/or lymph node involvement,  $n = 35$ ). In all cases, chemotherapy was administered adjuvantly; three courses of bleomycin, vincristine, mitomycin, and cisplatin for intermediate-risk cases and five courses for high-risk cases. Disease-free survival and complications of the combined therapy were investigated.

**Results.** Estimated 5-year disease-free survival was 93.3% for the 30 patients with intermediate-risk tumors (100% for those with squamous cell carcinoma and 71.4% for those with adenosquamous carcinoma) and 85.7% for the 35 patients with high-risk tumors (89.3% for those with squamous cell carcinoma and 71.4% for those with adenosquamous carcinoma). The incidence of locoregional recurrence was 3.3% in the intermediate-risk group and 8.6% in the high-risk group. Side effects of chemotherapy and complications of the combined therapy were within acceptable limits. No patient had severe bleomycin-related pulmonary toxicity. Only 1.5% of patients developed small bowel obstruction, which was cured by conservative therapy.

**Conclusions.** The treatment results suggest the potential role of adjuvant chemotherapy alone for patients with cervical cancer.

© 2006 Elsevier Inc. All rights reserved.

**Keywords:** Cervical cancer; Chemotherapy; Adjuvant therapy

### Introduction

Radiotherapy (RT) has been used as postoperative adjuvant therapy to reduce recurrence in patients with cervical cancer. For patients with intermediate-risk tumors, i.e., cervical cancer showing deep stromal invasion and/or capillary space involvement but no lymph node metastasis, the effectiveness of RT has been widely accepted, based on the cure rate [1] and the results of a randomized study reported by Sedlis et al. [2]. For patients with node-positive cervical cancer, however, reports on the outcome of postoperative RT have implied that adjuvant RT alone may change the pattern of recurrence from intrapelvic to

extrapelvic but offer little survival benefit [3–6]. In 2000, Peters et al. [7] advocated a new strategy for treating high-risk cervical cancer. They showed a significant survival advantage with the use of concurrent chemoradiation (CCR) rather than RT alone in patients with high-risk cervical cancer.

The debate regarding the role of adjuvant therapy for cervical cancer patients has generally focused on RT or CCR, but several studies have suggested the utility of chemotherapy (CT) alone as postoperative adjuvant therapy for cervical cancer. Lahousen et al. [8] prospectively compared adjuvant CT with RT in patients with high-risk cervical cancer and concluded that neither of the regimens improved survival or the rate of recurrence. In that study, however, the 5-year survival rate was about 80% among 19 patients with nodal disease who received CT. Iwasaka et al. [9] reported an 83% 5-year survival rate among

\* Corresponding author. Fax: +81 3 3570 0343.

E-mail address: [nobuhiro.takeshima@jicr.or.jp](mailto:nobuhiro.takeshima@jicr.or.jp) (N. Takeshima).

patients with high-risk cervical cancer treated adjuvantly with CT. They also reported an advantage of CT in the treatment of local failure. These findings suggest that the value of CT as adjuvant therapy for cervical cancer is worth reexamining.

At our institute, CT alone has been used as postoperative adjuvant therapy for cervical cancer since 1993. Herein, we report the treatment results among patients with stage IB–IIA squamous cell or adenosquamous cervical cancer, all of whom were treated initially by radical hysterectomy, had intermediate- or high-risk tumors, and received CT postoperatively.

### Patients and methods

Between 1993 and 2002, 233 patients with stage IB–IIA squamous cell or adenosquamous cervical cancer underwent radical hysterectomy and pelvic lymphadenectomy at the Cancer Institute Hospital (Tokyo, Japan). None of the patients had received any treatment prior to surgery. After pathologic assessment, patients who were identified as having intermediate- or high-risk tumors were scheduled to receive postoperative adjuvant CT without RT.

The criteria for intermediate-risk tumors were the presence of more than 50% cervical stromal invasion and the absence of high-risk factors. The presence or absence of lymph-vascular space involvement and tumor size does not alter these criteria. The criterion for high-risk tumors was the presence of at least one of the three following factors: histologically confirmed positive lymph nodes, parametrial invasion, and a positive surgical margin.

Of the 233 patients, 35 were identified as having intermediate-risk tumors, but five of these patients were excluded from the study. None of the five patients received CT; three refused it and two had medical problems. One of the five patients received RT. Forty patients were identified as having high-risk tumors, but five were excluded from the study. One patient refused CT and ultimately received no adjuvant therapy. The remaining four patients received CT, but one patient also received RT, two received a different regimen of CT, and the last patient died of primary lung cancer 6 months after treatment. Thus, 30 patients with intermediate-risk tumors and 35 patients with high-risk tumors were the study subjects. Patient characteristics are listed in Table 1.

Table 1  
Patient characteristics

	Intermediate-risk <i>n</i> = 30	High-risk <i>n</i> = 35
Median age in years (range)	49 (28–70)	43 (25–68)
Clinical stage		
Stage IB1	16	20
Stage IB2	12	12
Stage IIA	2	3
Cell type		
Squamous	23	28
Adenosquamous	7	7
Stromal invasion		
Inner half	0	10
Outer half	30	25
Lymph-vascular space involvement		
No	13	7
Yes	17	28
Positive pelvic nodes		
No	30	4
Yes	0	31
Parametrial involvement		
No	30	27
Yes	0	8
Positive margins		
No	30	30
Yes	0	5

Table 2

Relapse rates and sites of failure

Recurrence	Intermediate-risk ( <i>n</i> = 30)	High-risk ( <i>n</i> = 35)
Locoregional recurrence	1 (3.3%)	3 (8.6%)
Distant recurrence	1 (3.3%)	2 (5.7%)
Both	0	0
Total	2 (6.7%)	5 (14.3%)

Throughout the study period, only CT was administered as postoperative adjuvant therapy, with a single regimen (modified BOMP) being used. The chemotherapy regimen has been described elsewhere [10]. In brief, it consisted of bleomycin (5 mg in 500 mL saline infused intravenously [IV] continuously for 7 consecutive days), vincristine (0.7 mg/m<sup>2</sup> as an IV bolus on day 7), mitomycin (7 mg/m<sup>2</sup> as an IV bolus on day 7), and cisplatin (10 mg/m<sup>2</sup> dissolved in 500 mL saline and infused over 4 h on days 1 through 7). This regimen was scheduled to be repeated every 4 weeks for three cycles in patients with intermediate-risk tumors and five cycles in those with high-risk tumors. The number of CT cycles was based on our prior experience with this CT regimen. For objective targets, the median number of cycles per patient was four [10].

The disease-free interval, sites of relapse, and the morbidity after treatment were investigated in the study patients. Median follow-up time for surviving patients was 75 (range, 39–132) months in the intermediate-risk group and 74 (range, 38–126) months in the high-risk group. Disease-free survival was calculated by the Kaplan–Meier method and was compared by the log-rank test.

### Results

#### Treatment outcome

Two (6.7%) of the 30 patients with intermediate-risk disease suffered recurrence (Table 2). One of these two patients suffered recurrence in the pelvic cavity 18 months after treatment, and the other suffered multifocal recurrence in the liver and lung 11 months after treatment. These two cases were of the adenosquamous cell type. None of the 23 patients with intermediate-risk squamous cell carcinoma suffered recurrence. The estimated 5-year disease-free survival rate in this group was 93.3% (100% for patients with squamous cell carcinoma and 71.4% for those with adenosquamous carcinoma) (Fig. 1).




Fig. 1. Disease-free survival of patients with intermediate- and high-risk cervical cancer. Estimated 5-year disease-free survival rates were 93.3% for the 30 patients with intermediate-risk disease and 85.7% for the 35 patients with high-risk disease. No significant difference was noted between the two groups ( $P = 0.3122$ , log-rank test).



Five (14.3%) of the 35 patients with high-risk disease suffered recurrence. Three suffered locoregional recurrence, and the remaining two patients suffered distant recurrence (Table 2). Of the 28 patients with high-risk squamous cell carcinoma, three (10.7%) suffered recurrence, whereas two of the seven patients with high-risk adenosquamous carcinoma (28.6%) suffered recurrence. The estimated 5-year disease-free survival rate in the high-risk group was 85.7% (89.3% for patients with squamous cell carcinoma and 71.4% for those with adenosquamous carcinoma). Survival did not differ significantly between the intermediate- and high-risk groups ( $P = 0.3122$ ) (Fig. 1).

Four patients with locoregional recurrence were treated chiefly with RT, but none survived. The three patients with distant recurrence were treated by CT other than the BOMP regimen, but all died from the disease.

#### Chemotherapy tolerance and toxicity

Of the intermediate-risk patients, seven (23.3%) were unable to complete three cycles of CT; one completed one cycle and six completed two cycles. Of the high-risk patients, 10 (28.6%) were unable to complete five cycles of CT; two completed two cycles, three completed three cycles, and five completed four cycles. In most cases, the patients refused to continue chemotherapy due to CT-related nausea and vomiting.

Side effects of adjuvant CT were within acceptable limits. No patient had severe bleomycin-related pulmonary toxicity or cisplatin-related renal dysfunction. Bleomycin-related dermatitis was observed in one patient. Grade 3 or 4 myelosuppression was observed in 16.7% (5/30) of the intermediate-risk group and 31.4% (11/35) of the high-risk group, respectively.

#### Complications of the combined therapy

Complications requiring surgical intervention occurred in 2 (3.1%) of the 65 patients, none of which was related to tumor recurrence (Table 3). In one of the two patients, herniation of the abdominal wall was surgically repaired 7 months after the initial surgery. In the other patient, hydronephrosis requiring catheterization between the bladder and kidney was found 10 days after surgery. No patient developed urinary fistula. One patient (1.5%) suffered small bowel obstruction, but this was cured by conservative therapy. Three patients (4.6%) suffered locoregional infection (pelvic lymphocyst,  $n = 2$ ; leg cellulitis,  $n = 1$ ), all of whom were cured without surgical intervention.

Table 3  
Complications of combined therapy ( $n = 65$ )

Complication	No. of cases (requiring surgery)
Gastrointestinal	
Small bowel obstruction	1 (0)
Genitourinary	
Hydronephrosis	3 (1)
Infectious	
Pelvic lymphocyst	2 (0)
Leg cellulitis	1 (0)
Other	
Herniation of the abdominal wall	1 (1)

#### Discussion

RT has long been used as postoperative adjuvant therapy for stages IB and IIA cervical cancer. In a prospective study of postsurgical patients treated for intermediate-risk cervical cancer, Sedlis et al. [2] showed that adjuvant pelvic RT reduced recurrence from 28% to 15% at 2 years after treatment. Because node-positive cases were excluded from that study, the effectiveness of RT alone for high-risk cervical cancer was unclear, but results suggest that RT has a role in adjuvant therapy. For postsurgical patients with high-risk disease, however, it has been reported that RT alone may decrease the local failure rate, but provide no survival benefit (3–6). Although the necessity of systemic therapy for patients with high-risk disease has been suggested, several studies prospectively examined the effectiveness of CT followed by RT in patients with high-risk cervical cancer in an attempt to reduce both local and distant recurrence, but most of these studies failed to show any survival advantage [11,12]. This problem was resolved by Peters et al. [7]. They examined two treatment modalities, RT and CCR, in 243 patients with high-risk cervical cancer (207 node-positive patients) and obtained 4-year disease-free survival rates of 63% and 80%, respectively. Thus, RT for intermediate-risk postsurgical patients and CCR for high-risk postsurgical patients are at this time considered the optimum treatments. CCR may also be applicable for intermediate-risk tumors.

Despite this worldwide trend, we have consistently treated patients who have undergone surgery for cervical cancer with CT alone for several reasons. First, distant metastasis is a major problem when RT alone is used to treat high-risk cervical cancer, and CT is considered the most powerful means of eradicating subclinical distant metastases. If RT is added to CT, the CT dosage is limited. Second, our CT regimen yielded a 76% response rate in patients with recurrent cervical cancer and an acceptable incidence of side effects [10]; this response rate is much better than that of other recent studies [13]. Third, when CT alone is used for adjuvant therapy, RT can be reserved for local recurrence, simplifying treatment of local recurrence should it occur. Finally, this treatment strategy may provide a better postoperative quality of life by precluding radiation-related morbidity, such as small bowel obstruction or leg lymphedema. Conversely, the biggest concern is that the possibility for local recurrence may increase in the absence of pelvic RT.

Data from the current study indicate that suppression of distant metastasis by means of postoperative CT is feasible in both intermediate- and high-risk patients. One of thirty patients with intermediate-risk disease and 2 of 35 patients with high-risk suffered distant metastasis. More importantly, few patients suffered local recurrence. Despite the absence of pelvic RT, only 3.3% (1/30) of intermediate-risk patients and 8.6% (3/35) of high-risk patients developed locoregional recurrence. Iwasaka et al. [9] performed adjuvant CT (three courses of cisplatin, vincristine, mitomycin, and peplomycin) for high-risk cervical cancer and compared their patients with those from another hospital where adjuvant RT was performed during the same period. The results of CT (an 83% 5-year

survival rate) were similar to those of RT (an 81.7% 5-year survival rate). In that study, however, intra- and extrapelvic recurrences accounted for 85% and 23%, respectively, of all recurrences in the CT group. A relatively high incidence of local recurrence in patients with high-risk cervical cancer (14.3%, 4/28) treated with CT alone (six courses of carboplatin and bleomycin) was also reported by Lahousen et al. [8]. Both studies included stage IIB cancers, and we assume that the majority of cases of local recurrence were cases of stage IIB disease. Data from the current study, which included stage IB and IIA cancers, suggest that intrapelvic recurrence is not a major obstacle when CT alone is used as adjuvant therapy.

With respect to treatment of local recurrence, Iwasaka et al. [9] reported that 45% of patients with local recurrence in the CT group survived. Because there were only four cases with local recurrence in the current series, we were unable to confirm this finding. In our study, all patients with local recurrence were treated by full dose of RT, but none survived. However, in the Lahousen et al. study [8], four of the eight patients with pelvic recurrence in their CT group survived. Although further details were not given, it is probable that the surviving patients with local failure were treated successfully by RT.

It must be emphasized that the current study excluded patients with adenocarcinoma; these patients are treated with a different CT regimen at our institute. However, the treatment outcome of these patients is considerably worse than that of patients with squamous cell carcinoma (data not shown). Peters et al. [7] reported no survival difference between patients with high-risk adenocarcinoma and those with high-risk squamous cell carcinoma when CCR was used. These findings, along with the fact that more than half of the recurrences in the current study stemmed from adenosquamous carcinoma, suggest that the survival advantage of CT indicated in the current study may apply only to squamous cell carcinoma.

Another issue that must be discussed is complications of adjuvant therapy following surgery. Radical hysterectomy followed by RT is reported to be associated with a relatively high incidence of morbidity [14–16]. Barter et al. [15] reported that 30% of patients treated had serious complications; small bowel obstruction in particular has been one of the major problems of that treatment combination. Data from the current study well reflect the characteristics of our combined therapy. A very low incidence of small bowel obstruction or genitourinary complication was observed. Conversely, a relatively high incidence of locoregional infection was noted. These findings are probably associated with the absence of pelvic RT and the use of postoperative CT. As for leg lymphedema, Soisson et al. [3] reported that the incidence of lymphedema requiring medical therapy was significantly increased from 5.2% in patients treated with surgery alone to 22% in those receiving surgery plus adjuvant pelvic RT. Because there was no control group in our study, we were unable to show an advantage of CT with respect to leg lymphedema, but we do have the impression

that CT has little if any adverse effect pertaining to leg lymphedema. Further studies are needed to clarify this point.

In summary, our results suggest the potential role of adjuvant CT alone for patients with cervical cancer. Because of the absence of whole pelvic radiation, this treatment is likely to be advantageous in terms of treatment-related complications. We suggest that CT alone is worth reconsidering as an alternative postoperative therapy for patients with cervical cancer.

## Acknowledgment

This work was supported in part by grants from the Smoking Research Foundation (Japan).

## References

- [1] Chung CK, Nahhas WA, Stryker JA, Curry SL, Abt AB, Mortel R. Analysis of factors contributing to treatment failure in stages IB and IIA carcinoma of the cervix. *Am J Obstet Gynecol* 1980;138:550–6.
- [2] Sedlis A, Bundy BN, Rotman MZ, Lentz SS, Mudderspach LI, Zaino RJ, et al. A randomized trial of pelvic radiation therapy versus no further therapy in selected patients with stage IB carcinoma of the cervix after radical hysterectomy and pelvic lymphadenectomy. *Gynecol Oncol* 1999;73:177–83.
- [3] Soisson AP, Soper JT, Clarke-Pearson DL, Berchuk A, Montana G, Creasman WT. Adjuvant radiotherapy following radical hysterectomy for patients with stage IB and IIA cervical cancer. *Gynecol Oncol* 1990;37:390–5.
- [4] Thomas GM, Dembo AJ. Is there a role for adjuvant pelvic radiotherapy after radical hysterectomy in early stage cervical cancer. *Int J Gynecol Cancer* 1991;1:1–8.
- [5] Kinney WK, Alvarez RD, Reid GC, Schray MF, Soong SJ, Morley GW, et al. Value of adjuvant whole-pelvic irradiation after Wertheim hysterectomy for early-stage squamous carcinoma of the cervix with pelvic nodal metastasis: matched-control study. *Gynecol Oncol* 1989;34:258–62.
- [6] Uno T, Ito H, Itami J, Yasuda S, Isobe K, Hara R, et al. Postoperative radiation therapy for stage IB–IIB carcinoma of the cervix with poor prognostic factors. *Anticancer Res* 2000;20:2235–9.
- [7] Peters III WA, Liu PY, Barrett II RJ, Stock RJ, Monk BJ, Berek JS, et al. Concurrent chemotherapy and pelvic radiation therapy compared with pelvic radiation therapy alone as adjuvant therapy after radical surgery in high-risk early-stage cancer of the cervix. *J Clin Oncol* 2000;18:1606–13.
- [8] Lahousen M, Haas J, Pickel H, Hackel A, Kurz C, Ogris H, et al. Chemotherapy versus radiotherapy versus observation for high-risk cervical carcinoma after radical hysterectomy: a randomized, prospective, multicenter trial. *Gynecol Oncol* 1999;73:196–201.
- [9] Iwasaka T, Kamura T, Yokoyama M, Matsuo N, Nakano H, Sugimori H. Adjuvant chemotherapy after radical hysterectomy for cervical carcinoma: comparison with effects of radiotherapy. *Obstet Gynecol* 1988;91:977–81.
- [10] Shimizu Y, Akiyama F, Umezawa S, Ishiya T, Utusgi K, Hasumi K. Combination of consecutive low-dose cisplatin with bleomycin, vincristine, and mitomycin for recurrent cervical carcinoma. *J Clin Oncol* 1998;16:1869–78.
- [11] Tattersall MH, Ramirez C, Coppleson M. A randomized trial of adjuvant chemotherapy after radical hysterectomy in stage Ib–IIa cervical cancer patients with pelvic lymph node metastasis. *Gynecol Oncol* 1992;46:176–81.
- [12] Curtin JP, Hoskins WJ, Venkatraman ES, Almadrones L, Podraz KC, Long H, et al. Adjuvant chemotherapy versus chemotherapy plus pelvic irradiation for high-risk cervical cancer patients after radical hysterectomy and pelvic lymphadenectomy (RH-PLND): randomized phase III trial. *Gynecol Oncol* 1996;61:3–10.
- [13] Long III HJ, Bundy BN, Grendys EC, Benda JA, McMeekin DS, Sorosky J, et al. Randomized phase III trial of cisplatin with or without topotecan in carcinoma of the uterine cervix. *J Clin Oncol* 2005;23:4626–33.

- [14] Dische S. Radiotherapy of cervical cancer. *Clin Obstet Gynaecol* 1985;12:203–27.
- [15] Barter JF, Soong SJ, Shingleton HM, Hatch KD, Orr Jr JW. Complications of combined radical hysterectomy–postoperative radiation therapy in women with early stage cervical cancer. *Gynecol Oncol* 1989;32:292–6.
- [16] Monk BJ, Cha DS, Walker JL, Burger RA, Ramsinghani NS, Manetta A, et al. Extent of disease as an indication for pelvic radiation following radical hysterectomy and bilateral pelvic lymph node dissection in the treatment of stage IB and IIA cervical carcinoma. *Gynecol Oncol* 1994;54:4–9.

EDITORIAL

## Concurrent chemoradiation for cervical cancer: what should we do next?

Article on page 309

**Feasibility of concurrent cisplatin use during primary and adjuvant chemoradiation therapy: a phase I study in Japanese patients with cancer of the uterine cervix**

Y. Watanabe, H. Nakai, M. Shimaoka, T. Tobiume, I. Tsuji, H. Hoshiai

Received: March 29, 2006

Several randomized trials have demonstrated the therapeutic advantage of concurrent chemoradiotherapy (CCRT) over radiation alone in the treatment of locoregionally advanced uterine cervical cancer.<sup>1–4</sup> Consequently, CCRT is now considered a standard treatment component for patients with uterine cervical cancer. In these trials, cisplatin was used alone or in combination with 5-fluorouracil (5-FU). Rose et al.<sup>2</sup> demonstrated similar survival results and less toxicity with weekly cisplatin compared with moderate-dose cisplatin and 5-FU in a Gynecologic Oncology Group (GOG) study (GOG-120). Based on those results, weekly cisplatin (40 mg/m<sup>2</sup>) is now a standard regimen in the United States. However, it should be noted that the average patient age in these studies is younger than that seen in clinical practice in Japan. The Japanese Patterns of Care Study (JPCS) demonstrated that the median age of Japanese cervical cancer patients treated with definitive radiotherapy was 70 years.<sup>5</sup> In contrast, 80%–90% of patients entered in the GOG studies were below 60 years of age.<sup>2–4</sup> Therefore, an original prospective clinical study of CCRT for Japanese patients has been encouraged to assess the feasibility and toxicity of weekly cisplatin at 40 mg/m<sup>2</sup>.

In this issue of the *International Journal of Clinical Oncology*, Watanabe et al.<sup>6</sup> report the results of a phase I study of CCRT with single-agent cisplatin in Japanese patients with cervical cancer. They designed two dose-escalation programs for weekly and monthly schedules. Dose-limiting toxicities (DLTs) were investigated separately for both definitive and postoperative settings. Their DLT criteria were designed somewhat similarly to those of recent GOG phase I studies. Treatment delay (i.e., radiotherapy and chemotherapy) of greater than 7 days due to toxicity was defined as a DLT. The investigators observed that the two patients in the definitive CCRT arm and two of the three in the postoperative CCRT arm experienced DLT at the dose level of 40 mg/m<sup>2</sup> (maximum tolerated dose; MTD). Therefore, they recommended a cisplatin dose of 30 mg/m<sup>2</sup> for the weekly schedule. Similarly, they concluded that 75 mg/m<sup>2</sup> should be the recommended dose for the monthly schedule. From these results, they concluded that weekly cisplatin at a dose of 40 mg/m<sup>2</sup> was not feasible for Japanese patients.

To my knowledge, the GOG has not officially performed a phase I/II study of CCRT with weekly cisplatin at a dose of 40 mg/m<sup>2</sup>. Therefore, there are no feasibility data for this protocol evaluated by standard DLT criteria, as Watanabe et al.<sup>6</sup> stated in their report. Keys et al.,<sup>3</sup> in GOG-123, showed that 90% of patients were able to receive four or more courses of cisplatin. Similarly, Rose et al.<sup>2</sup> reported that, in GOG-120, four or more courses could be delivered in over 90% of patients and that nearly half of all patients received all the planned courses (six courses) of cisplatin. However, these studies<sup>2,3</sup> did not describe the details or the number of patients who required treatment modification and/or treatment delay, especially in regard to cisplatin administration. As mentioned above, Watanabe et al.<sup>6</sup> have now stated explicitly that more than a 7-day treatment delay constitutes a DLT. Can we not conclude that 8–14 days of delay in cisplatin administration is clinically acceptable in CCRT? Watanabe et al.<sup>6</sup> reported that all patients with DLT experienced granulocytopenia on the day of the planned fourth cycle of cisplatin. If further waiting had been allowed for these patients, the fourth cycle might have been safely given, and this phase I study could have reached a different conclusion. Ohno et al.<sup>7</sup> reported another Japanese phase I study, from the National Institute of Radiological Sciences in Japan. They concluded that weekly cisplatin at 40 mg/m<sup>2</sup> was feasible for Japanese patients. In the Ohno study, a longer treatment delay was allowed; namely, 2 weeks for radiotherapy and 3 weeks for chemotherapy. It should be noted that these authors reported that four of five patients who developed DLT were able to receive the full course of radiotherapy without interruption. In their report, Watanabe et al.<sup>6</sup> also showed that most patients were able to receive the full dose of radiation despite developing DLT.

Regrettably, in their current article, Watanabe et al. did not provide a detailed toxicity profile. Table 1 shows the toxicities that have been reported for CCRT using weekly cisplatin at 40 mg/m<sup>2</sup>. The article by Ikushima et al.<sup>8</sup> reports retrospective toxicity data for Japanese patients with cervical cancer who were given concurrent weekly cisplatin and radiotherapy. These authors decided to determine cisplatin

**Table 1.** Acute toxicity (grade 3 or more) of CCRT using weekly cisplatin at 40 mg/m<sup>2</sup>

Author (study)	Year	No. of patients	Toxicity (grade 3/4) %			
			Hematological	Leukopenia	Thrombocytopenia	Gastrointestinal
Rose <sup>2</sup> (GOG-120)	1999	176	—	21/2	2/0	8/4
Keys <sup>3</sup> (GOG-123)	1999	183	18/3	—	—	9/5
Pearcey <sup>10</sup>	2002	127	6/0	—	—	11/5
Serkies <sup>9</sup>	2004	112	—	4/2	0	4/2*
Ohno <sup>7</sup>	2005	6	—	83/0	0	16/0
Ikushima <sup>8</sup>	2006	11	91/9	—	—	9/0

\*Nausea/vomiting

dose according to patient age, with 40 mg/m<sup>2</sup> the dose for younger patients (under 65 years) and 30 mg/m<sup>2</sup> for older patients (65 years or over). They observed that all 11 patients treated with 40 mg/m<sup>2</sup> cisplatin developed grade 3 or greater hematological toxicity. In contrast to these Japanese series, severe hematologic toxicities have been reported to be less frequent in studies conducted in North America and Europe.<sup>2,3,9,10</sup> In Poland, Serkies et al.<sup>9</sup> reported data, from their routine clinical practice, on patient compliance and an acute toxicity profile for CCRT including weekly 40 mg/m<sup>2</sup> cisplatin. Seventy-four percent of their patients with cervical cancer received at least four cycles of cisplatin. They showed that nearly 40% of patients who did not receive the five planned cycles had reasons other than toxicity, such as delayed administration of the first cycle of chemotherapy. They reported that only 5% of patients experienced grade 3 or 4 leukopenia.

On the basis of these findings, what should we do next in CCRT for cervical cancer? I have concluded that a new phase I study seeking an optimal weekly cisplatin dose would have little value. It is important to note that a survival advantage has been demonstrated with cisplatin at the very dose of 40 mg/m<sup>2</sup>, but not at a compromised lower dose. On the other hand, it is true that Japanese patients have experienced severe hematological toxicity more frequently than patients in North America and Europe. However, life-threatening toxicity has been uncommon, even in Japanese patients; most importantly, radiotherapy has rarely been interrupted by toxicities. Thus, I believe we should conduct a proper phase II study of CCRT with weekly cisplatin at 40 mg/m<sup>2</sup> for Japanese patients, to determine its toxicity profile and survival outcome. In designing the trial, a treatment modification rule for cisplatin administration should be carefully planned. For this purpose, a detailed description of supportive treatment, such as the use of granulocyte-colony-stimulating factor (G-CSF) should be included in the protocol.

In addition to the scheduling of chemotherapy, the radiotherapy method is another important problem to be investigated in a future trial of CCRT for cervical cancer. There are several differences between the United States and Japan in the radiotherapy methods used. The most notable differences are in the dose rates of intracavitary brachytherapy (ICBT) and the total dose of radiotherapy. Although a National Cancer Institute of Canada

(NCIC) study<sup>10</sup> employed not only low-dose-rate (LDR) but also high-dose-rate (HDR) and medium-dose-rate (MDR) ICBT, all randomized studies from the United States have only utilized LDR.<sup>1-4</sup> In contrast, almost all patients in Japan have been treated with HDR-ICBT.<sup>5</sup> Therefore, it should be investigated whether CCRT with HDR-ICBT leads to the same favorable outcome and acceptable toxicity as those demonstrated in earlier randomized studies utilizing LDR-ICBT. In addition, a large variance between the United States and Japan in the total radiotherapy dose has been identified.<sup>11</sup> This is another reason to conduct a prospective phase II study using weekly cisplatin with HDR-ICBT in Japan. Trials of several new chemotherapeutic regimens for CCRT are now underway. Baseline CCRT data with a "standard chemotherapeutic regimen" obtained with Japanese patients will be essential to make proper comparisons with investigational CCRT regimens in future.

## References

- Morris M, Eifel PJ, Lu J, et al. (1999) Pelvic radiation with concurrent chemotherapy compared with pelvic and para-aortic radiation for high-risk cervical cancer. *N Engl J Med* 340:1137-1143
- Rose PG, Bundy BN, Watkins EB, et al. (1999) Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer. *N Engl J Med* 340:1144-1153
- Keys HM, Bundy BN, Stehman FB, et al. (1999) Cisplatin, radiation, and adjuvant hysterectomy compared with radiation and adjuvant hysterectomy for bulky stage IB cervical carcinoma. *N Engl J Med* 340:1154-1161
- Whitney CW, Sause W, Bundy BN, et al. (1999) Randomized comparison of fluorouracil plus cisplatin versus hydroxyurea as an adjunct to radiation therapy in stage IIB-IVA carcinoma of the cervix with negative para-aortic lymph nodes: a Gynecologic Oncology Group and Southwest Oncology Group study. *J Clin Oncol* 17:1339-1348
- Toita T, Nakamura K, Uno T, et al. (2005) Radiotherapy for uterine cervical cancer: results of the 1995-1997 Patterns of Care Process survey in Japan. *Jpn J Clin Oncol* 35:139-148
- Watanabe Y, Nakai H, Shimaoka M, et al. (2006) Feasibility of concurrent cisplatin use during primary and adjuvant chemoradiotherapy: a phase I study in Japanese patients with cancer of the uterine cervix. *Int J Clin Oncol* 11:309-313
- Ohno T, Kato S, Tsujii H (2005) Phase I study of weekly cisplatin plus radiotherapy for locally advanced cervical cancer (in Japanese). *Jpn J Gynecol Oncol* 23:564-571

8. Ikushima H, Osaki K, Furutani S, et al. (2006) Chemoradiation therapy for cervical cancer: toxicity of concurrent weekly cisplatin. *Radiat Med* 24:115-121.
9. Serkies K, Jassem J (2004) Concurrent weekly cisplatin and radiotherapy in routine management of cervical cancer: a report on patient compliance and acute toxicity. *Int J Radiat Oncol Biol Phys* 60:814-821.
10. Pearcey R, Brundage M, Drouin P, et al. (2002) Phase III trial comparing radical radiotherapy with and without cisplatin chemotherapy in patients with advanced squamous cell cancer of the cervix. *J Clin Oncol* 20:966-972.
11. Toita T, Kakinohana Y, Ogawa K, et al. (2003) Combination external beam radiotherapy and high-dose-rate intracavitary brachytherapy for uterine cervical cancer: analysis of dose and fractionation schedule. *Int J Radiat Oncol Biol Phys* 56:1344-1353.

Takafumi Toita, M.D.

Department of Radiology, Graduate School of Medical Science, University of the Ryukyus, 207 Uehara, Nishihara-cho, Okinawa 903-0215, Japan  
Tel. +81-98-895-1162; Fax +81-98-895-1420  
e-mail: b983255@med.u-ryukyu.ac.jp



## Frequency and characteristics of isolated para-aortic lymph node recurrence in patients with uterine cervical carcinoma in Japan: A multi-institutional study

Yuzuru Niibe <sup>a,\*</sup>, Tomoko Kazumoto <sup>b</sup>, Takafumi Toita <sup>c</sup>, Hideya Yamazaki <sup>d</sup>, Keiko Higuchi <sup>e</sup>,  
Noriko Ii <sup>f</sup>, Kazunori Suzuki <sup>g</sup>, Takashi Uno <sup>h</sup>, Sunao Tokumaru <sup>i</sup>, Makoto Takayama <sup>j</sup>,  
Kenji Sekiguchi <sup>k</sup>, Yasuo Matsumoto <sup>l</sup>, Koichi Michimoto <sup>m</sup>,  
Masahiko Oguchi <sup>m</sup>, Kazushige Hayakawa <sup>a</sup>

<sup>a</sup> Department of Radiology, Kitasato University School of Medicine, 1-15-1, Kitasato, Sagami-hara, Kanagawa 228-8555, Japan

<sup>b</sup> Department of Radiology, Saitama Cancer Center, Japan

<sup>c</sup> Department of Radiology, Ryuky University School of Medicine, Japan

<sup>d</sup> Department of Radiation Oncology, Osaka Medical Center for Cancer and Cardiovascular Diseases, Japan

<sup>e</sup> Department of Radiology, Gunma Cancer Center, Japan

<sup>f</sup> Department of Radiology, Mie University School of Medicine, Japan

<sup>g</sup> Department of Radiology, Hamamatsu University School of Medicine, Japan

<sup>h</sup> Department of Radiation Oncology, Chiba University Graduate School of Medicine, Japan

<sup>i</sup> Department of Radiology, Saga University School of Medicine, Japan

<sup>j</sup> Department of Radiology, Kyorin University School of Medicine, Japan

<sup>k</sup> Department of Radiology, St. Luke's Hospital, Japan

<sup>l</sup> Department of Radiology, Niigata Cancer Center, Japan

<sup>m</sup> Department of Radiation Oncology, Cancer Institute Hospital, the Japanese Foundation for Cancer Research, Japan

Received 2 October 2005

Available online 4 May 2006

### Abstract

**Objective.** In most cases of uterine cervical carcinoma recurrence, the first site of distant metastasis or recurrence is reported to be the para-aortic region. Some reports have demonstrated that, in cases of isolated para-aortic lymph node recurrence treated by radiation therapy, patients survived for a long period, which suggests that isolated para-aortic lymph node recurrence in uterine cervical carcinoma is a regional disease rather than systemic disease. Determining the predictive characteristics of isolated para-aortic lymph node recurrence in patients at the time of the initial treatment for primary uterine cervical carcinoma is important, so we conducted the current multi-institutional study.

**Patients and methods.** Patients ( $n=3137$ ) with uterine cervical carcinoma of stages Ia to IVa were treated in twelve Japanese hospitals between 1994 and 2003. The current study investigated the frequency and characteristics of patients with isolated para-aortic lymph node recurrence as well as the characteristics of clinical stage, histopathology, serum squamous cell carcinoma antigen level, the treatment method at the initial treatment, the duration between the initial treatment and the recurrence, and the serum squamous cell carcinoma antigen level at the recurrence.

**Results.** Of the 3137 patients with uterine cervical carcinoma in stages Ia–IVa, 67 (2.1%) experienced recurrence in isolated para-aortic lymph nodes. Stratified by clinical stage, none of the 613 patients with stage Ia experienced recurrence in isolated para-aortic lymph nodes. However, recurrence was experienced by 14 (1.4%) of the 966 patients with stage Ib, 7 (3.5%) of the 199 patients with stage IIa, 14 (2.3%) of the 613 patients with stage IIb, 1 (2.1%) of the 48 patients with stage IIIa, 26 (4.6%) of the 538 patients with stage IIIb, and 5 (5%) of the 100 patients with stage IVa. The mean duration time between the initial treatment and isolated para-aortic recurrence was 20 months (range, 2–49 months). The correlations between duration time and the clinico-pathological factors (clinical stage, histopathology, serum squamous cell carcinoma antigen level, and treatment method) at the initial treatment were investigated. No statistically significant factors have been revealed in the current study.

\* Corresponding author. Fax: +81 42 778 9436.

E-mail address: [joe-n@hkg.odn.ne.jp](mailto:joe-n@hkg.odn.ne.jp) (Y. Niibe).

**Conclusions.** The frequency of isolated para-aortic lymph node recurrence was 2.1% and increased with increasing clinical stage at the initial treatment (stage IVa: 5%) in the current study.

© 2006 Elsevier Inc. All rights reserved.

**Keywords:** Isolated para-aortic lymph node recurrence; Uterine cervical carcinoma; Radiation therapy

## Introduction

Recently, oligo-metastasis/oligo-recurrence has been one of the most important concerns of oncology, especially in radiation oncology [1]. Advances of diagnostic imaging and biochemical diagnosis for various carcinomas have enabled us to detect isolated metastasis or recurrence, although some decades ago, metastasis and recurrence meant systemic disease in almost all cases. However, no strategy has been established for treating oligo-metastasis/oligo-recurrence in any kind of carcinoma. Most oncologists select systemic chemotherapy for these patients as a community standard. Nonetheless, oligo-metastasis/oligo-recurrence, especially isolated metastasis or recurrence, is considered to not always mean systemic disease. Singh et al. reported that patients with isolated para-aortic lymph node recurrence in uterine cervical carcinoma achieved 100% of 5-year survival when the recurrent site was treated with chemoradiotherapy [2]. Niibe et al. reported that the survival rates of patients with metastatic brain tumors with controlled primary lesions and no other distant metastasis were 88.9% after 1 year and 51.9% after 3 years respectively [3]. These findings suggested that some oligo-metastasis/oligo-recurrence patients could survive for as long a period as the patients with primary carcinoma. Thus, these patients must be treated curatively.

Uterine cervical carcinoma was reported to spread more by the lymphatic route than by the hematogenous route [4]. In most cases, the first site of distant metastasis or recurrence is the para-aortic region. As mentioned above, Singh et al. reported the long-term survival of patients with isolated para-aortic lymph node recurrence, which meant that isolated para-aortic lymph node recurrence in uterine cervical carcinoma is a regional disease rather than a systemic disease. Determining the characteristics of isolated para-aortic lymph node recurrence in patients at the time of the initial treatment for primary uterine cervical carcinoma is important.

Thus, we conducted the current multi-institutional study to reveal the frequency and characteristics of isolated para-aortic lymph node recurrence in uterine cervical carcinoma.

## Patients and methods

Patients ( $n=3137$ ) with uterine cervical carcinoma of stages Ia to IVa were treated in twelve Japanese university hospitals, cancer centers, and major general hospitals between 1994 and 2003. The current study investigated the frequency and characteristics of isolated para-aortic lymph node recurrence as well as the clinical stage, histopathology, serum squamous cell carcinoma (SCC) antigen level, initial treatment method, duration between the initial treatment and the recurrence, and serum SCC antigen level at the time of recurrence (Table 1).

Data were collected on data sheets from these twelve hospitals. Data sheets included patient age, serum SCC antigen level, treatment method, and the date at the initial treatment and patient age, serum SCC antigen level, and the date at the time of detection of isolated para-aortic recurrence. A data center was established at the Department of Radiology, Kitasato University Hospital.

## Results

Of the 3137 patients with uterine cervical carcinoma in stages I–IVa, 67 (2.1%) experienced recurrence in isolated para-aortic lymph nodes. Stratified by clinical stage, none of the 613 patients with stage Ia experienced recurrence in isolated para-aortic lymph nodes. However, recurrence was experienced by 14 (1.4%) of the 966 patients with stage Ib, 7 (3.5%) of the 199 patients with stage IIa, 14 (2.3%) of the 613 patients with stage IIb, 1 (2.1%) of the 48 patients with stage IIIa, 26 (4.6%) of the 538 patients with stage IIIb, and 5 (5%) of the 100 patients with stage IVa. These results suggested that patients with more locally advanced stages (IIIb and IVa) were more likely to experience recurrence in isolated para-aortic lymph nodes than patients with early locally invasive stages (I–II).

Other patients characteristics are summarized in Table 2. The mean age was 55.7 years (range, 25–86 years). The mean duration time between the initial treatment and isolated para-aortic recurrence was 20 months (range, 2–49 months) (Fig. 1). As for the initial treatment, 32 patients underwent external radiation therapy combined with intracavity radiation therapy alone; 20 patients underwent surgery combined with external radiation therapy; 12 patients underwent concurrent chemoradiation therapy (radiation therapy: external radiation therapy combined with intracavity radiation therapy); and 3 patients underwent surgery only. As for histopathology, 56 patients were found to have squamous cell carcinoma; 5 patients had adenocarcinoma; 5 patients had adenosquamous cell carcinoma; and 1 patient had a malignancy that was unclassified. The mean serum SCC antigen level at the start of the initial treatment was 17.3 ng/dl (range, 0.5–100 ng/dl), and the mean serum SCC antigen level at the time of isolated para-aortic lymph node recurrence was 9.5 ng/dl (range, 0–120 ng/dl). These results indicate that the serum SCC antigen level at the time of isolated para-aortic lymph node recurrence tended to be lower than that at the initial treatment. As for symptoms of the isolated para-aortic lymph node recurrence, 20 patients had symptoms with recurrence (Table 3). Lumbago was the most frequent symptom,

Table 1  
The frequency of isolated para-aortic lymph node recurrence

Clinical stage	Frequency of isolated para-aortic lymph node recurrence
Ia	0/613 (0%)
Ib	14/966 (1.4%)
IIa	7/199 (3.5%)
IIb	14/613 (2.3%)
IIIa	1/48 (2.1%)
IIIb	26/538 (4.6%)
IVa	5/100 (5%)
Ia–IVa	67/3137 (2.1%)



Table 2  
Patients characteristics of isolated para-aortic lymph node recurrence

Mean age	55.7 years (range; 25–86 years)
Histopathology	
Squamous cell carcinoma	56
Adenocarcinoma	5
Adenosquamous cell carcinoma	5
Unclassified	1
Initial treatment	
Radiation therapy alone	32
Chemoradiation therapy	12
Surgery followed by radiation therapy	20
Surgery alone	3
Mean serum SCC level	
Initial treatment	17.3 ng/dl (range; 0.5–100 ng/dl)
Recurrence	9.5 ng/dl (range; 0–120 ng/dl)
Mean DT*	20 months (range; 2–49 months)

\* Mean DT: the mean duration time between the initial treatment and isolated para-aortic recurrence.

seen in 14 patients. Three patients experienced edema of the lower extremities, and three patients experienced pain in the lower extremities. The correlations between duration time and the clinico-pathological factors (clinical stage, histopathology, serum SCC antigen level, and treatment method) at the initial treatment were investigated. No statistically significant factors have been revealed in the current study.

The correlation between serum SCC antigen level at the initial treatment and that at the time of isolated para-aortic lymph node recurrence was statistically significant ( $r = 0.492$ ,  $P = 0.01$ ) (Fig. 2).

The correlation between higher serum SCC antigen level ( $>10$  ng/dl) at the time of isolated para-aortic lymph node recurrence and coexisting symptoms at the time of recurrence was statistically significant ( $P = 0.05$ ).

## Discussion

Some patients with uterine cervical carcinoma and isolated para-aortic lymph node recurrence were reported to survive for a long period and were considered to be cured [2,5–7]. Singh

Table 3  
Symptom at the isolated para-aortic recurrence

Symptom	Number of patients
Lumbago	14
Edema of lower extremities	3
Pain of lower extremities	3

et al. reported that 100% of patients with uterine cervical carcinoma and isolated para-aortic lymph node recurrence treated with concurrent chemoradiotherapy achieved 5-year survival, although patients treated with only chemotherapy died within 1.5 years [2]. Niibe et al. reported that, in cases of advanced uterine cervical carcinoma with isolated para-aortic lymph node recurrence or metastasis treated with radiation therapy, 38% of patients achieved 5-year survival and the authors pointed out that c-erb B-2/HER2 expression in tumor tissues had prognostic significance, suggesting that anti-c-erb B-2/HER2 therapy, molecule-targeting therapy, such as with trastuzumab, might have an influence on survival [5]. These findings indicated that isolated para-aortic lymph node recurrence in uterine cervical carcinoma was not considered to be a systemic disease but to be a loco-regional disease. The detection of isolated para-aortic lymph node recurrence is important, so we investigated the frequency and characteristics of isolated para-aortic lymph node recurrence in patients with uterine cervical carcinoma.

The current multi-institutional study revealed that 2.1% of patients with uterine cervical carcinoma treated with curative therapy (including radiation therapy, chemoradiation therapy, surgery, and combined therapy) experienced recurrence in isolated para-aortic lymph nodes. This is the clinical demonstration in the largest population ( $n = 3137$ ). Others report on this theme in a large population as follows. Chou et al. reported in 2001 that 26 out of 867 patients (3%) who received pelvic radiotherapy after the diagnosis of primary cervical carcinoma were found to have isolated para-aortic lymph node recurrence in Taiwan [7]. Hong et al. reported in 2004 that 46 out of 1292 patients (3.6%) with uterine cervical carcinoma who underwent curative intended radiation therapy were found to have para-aortic lymph node recurrence in Taiwan [8]. Tsai et al. reported




Fig. 1. The cumulative mean duration time between the initial treatment and isolated para-aortic lymph node recurrence was demonstrated. The mean duration was 20 months (range, 2–49 months).




Fig. 2. The correlation between serum SCC antigen level at the start of the initial treatment and serum SCC antigen level at the isolated para-aortic lymph node recurrence. The positive correlation was recognized ( $r = 0.492$ ,  $P = 0.01$ ).

in 2005 that 14 out of 816 patients (1.7%) with uterine cervical carcinoma who received curative intended radiation therapy were found to have para-aortic lymph node recurrence in USA [2]. Furthermore, patients with more locally advanced stages at the initial treatment had a higher frequency of isolated para-aortic lymph node recurrence (stage Ib: 1.4% versus stage IVa: 5%). These results suggested the hypothesis that even locally advanced patients spread with lymphatic route not with hematogenous systemic route that existed not in a small number. Ikeda et al. reported that only 8 of 1961 patients with uterine cervical carcinoma experienced recurrence in the brain and that there were no patients with uterine cervical carcinoma that experienced isolated brain recurrence [9]. These results support the above-mentioned hypothesis.

Regarding the characteristics of isolated para-aortic lymph node recurrence in patients with uterine cervical carcinoma, about only one-third of these had symptoms. The others had no symptoms. Even no coexisting serum-SCC antigen elevation was not rare. Thus, routine examination by not only pelvic, but abdominal computed tomography or magnetic resonance imaging of uterine cervical carcinoma patients treated with curative therapy was considered to be required. Furthermore, coexisting symptoms in patients with isolated para-aortic lymph node recurrence were reported to be correlated with a much worse prognosis [2,10]. The mean duration time between isolated para-aortic recurrence and the initial treatment was 20 months (range, 2–49 months) in the current study. However, there were no correlations between duration time and various clinico-pathological factors (clinical stage, histopathology, serum SCC antigen level, and treatment method). These results suggested that long-term routine follow-up was required to detect isolated para-aortic lymph node recurrence. On the other hand, the correlation between the serum SCC antigen level at the time of initial treatment and that at the time of isolated para-aortic lymph node recurrence was statistically significant ( $r = 0.492$ ,  $P = 0.01$ ) (Fig. 2), and the correlation between higher serum SCC antigen level ( $>10$  ng/dl) at the time of isolated para-aortic lymph node recurrence and coexisting symptoms at the time of recurrence was also statistically significant ( $P = 0.05$ ). These suggested that routine follow-up with serum SCC antigen testing is important when the serum SCC antigen level is elevated at the time of initial treatment.

In conclusion, the frequency of isolated para-aortic lymph node recurrence was 2.1% and increased with increasing

clinical stage at the initial treatment (stage IVa: 5%) in the current study. Two-thirds of patients with isolated para-aortic lymph node recurrence had no symptoms. Routine follow-up with serum SCC antigen testing and abdominal computed tomography or magnetic resonance imaging is important except stage Ia.

#### Acknowledgments

This study was supported in part by grants from Ministry of Health, Labour and Welfare, Japan and from Kitasato University Research Grant for Young Researchers, Japan.

#### References

- [1] Hellman S, Weichselbaum RR. Oligometastases. *J Clin Oncol* 1995;13: 8–10.
- [2] Singh AK, Grigsby PW, Rader JS, Powell A. Cervix carcinoma, concurrent chemoradiotherapy, and salvage node recurrence. *Int J Radiat Oncol Biol Phys* 2005;61:450–5.
- [3] Niibe Y, Karasawa K, Nakamura O, Shinoura N, Okamoto K, Yamada R, et al. Survival benefit of stereotactic radiosurgery for metastatic brain tumors in patients with controlled primary lesions and no other distant metastases. *Anticancer Res* 2003;23:4157–60.
- [4] Robert JK, Henry JN, Edward W. Tumors of the cervix. In: Bethesda MD, editor. Atlas of tumor pathology. Washington, D.C.: Armed Forces Institute of Pathology; 1992. p. 37–139.
- [5] Niibe Y, Nakano T, Ohno T, Suzuki Y, Oka K, Tsujii H. Prognostic significance of c-erb B-2/HER2 expression in advanced uterine cervical carcinoma with para-aortic lymph node metastasis treated with radiation therapy. *Int J Gynecol Cancer* 2003;13:849–55.
- [6] Kim JS, Kim JS, Kim SY, Kim KH, Cho MJ. Hyperfractionated radiotherapy with concurrent chemotherapy for para-aortic lymph node recurrence in carcinoma of the cervix. *Int J Radiat Oncol Biol Phys* 2003; 55:1247–53.
- [7] Chou HH, Wang CC, Lai CH, Hong JH, Ng KK, Chang TC, et al. Isolated para-aortic lymph node recurrence after definitive irradiation for cervical carcinoma. *Int J Radiat Oncol Biol Phys* 2001;51:442–8.
- [8] Hong JH, Tsai CS, Lai CH, Chang TC, Wang CC, Chou HH, et al. Recurrence squamous cell carcinoma of cervix after definitive radiotherapy. *Int J Radiat Oncol Biol Phys* 2004;60:249–57.
- [9] Ikeda S, Yamada T, Katsumata N, Hida K, Tanemura K, Tsunematu R, et al. Cerebral metastasis in patients with uterine cervical cancer. *Jpn J Clin Oncol* 1998;28:27–9.
- [10] Grigsby PW, Vest ML, Perez CA. Recurrent carcinoma of the cervix exclusively in the para-aortic nodes following radiation therapy. *Int J Radiat Oncol Biol Phys* 1994;28:451–5.

# Peptide YY producing strumal carcinoid of the ovary as the cause of severe constipation with contralateral epithelial ovarian cancer

Kouichiro Kawano, Kimio Ushijima, Takefumi Fujimoto, Kan Komai and Toshiharu Kamura

*Department of Obstetrics and Gynecology, Kurume University, School of Medicine, Kurume, Fukuoka, Japan*

## Abstract

Primary ovarian carcinoid tumors are rare. It has been reported that constipation was a presenting symptom in some patients with ovarian carcinoid. A case of strumal carcinoid of the ovary with contralateral clear cell adenocarcinoma of the ovary discovered with a complaint of constipation is described. Constipation was dramatically improved by resectioning the tumor. The tumor cells were positive for peptide YY (PYY) in the carcinoid component, but not in any other components. The present case could provide evidence of the correlation between constipation and PYY that has been reported elsewhere. Interestingly, the constipation caused by PYY also helped in discovering epithelial ovarian cancer.

**Key words:** carcinoid tumor, constipation, endometriosis, ovarian cancer, peptide YY.

## Introduction

Primary ovarian carcinoid tumors comprise less than 0.1% of ovarian cancers.<sup>1</sup> The histologic patterns have been divided into four groups: insular, trabecular, strumal, and mucinous with or without associated mature teratoma. The insular type is the most common type of ovarian carcinoid, usually with carcinoid syndrome. Strumal carcinoids are the second most common type, and this type of carcinoid is usually a trabecular type carcinoid rather than an insular type. A number of cases of ovarian carcinoid with the chief complaint of severe constipation correlating with the production of peptide YY (PYY) have been reported.<sup>2-4</sup> A patient with ovarian carcinoid with synchronous contralateral epithelial ovarian cancer who suffered from severe constipation, which was relieved by removal of the tumor, is described.

## Case Report


A 47-year-old nulliparous woman visited the gastroenterology clinic because of persistent severe constipation. The patient had defecation once in 10 days despite taking 2 g of magnesium oxide every day. She was referred to the authors' University Hospital because an ovarian tumor had been suspected by ultrasonography. There was no clinical evidence of carcinoid syndrome (flushing, diarrhea, or cardiac disease) or abnormal thyroid function.

On physical examination, a cystic, immobile mass that was larger than fist-size and had a smooth surface was palpated in the lower abdomen. The uterus was enlarged to over goose-egg size with poor mobility. Computed tomography revealed a 17 × 9-cm sized right ovarian tumor and a 12 × 8-cm sized left ovarian tumor. The right ovarian tumor showed a cystic mass

Received: June 1 2006.

Accepted: September 22 2006.


Reprint request to: Dr Kouichiro Kawano, Department of Obstetrics and Gynecology, Kurume University, School of Medicine, 67 Asahi-machi, Kurume, Fukuoka 830-0011, Japan. Email: [kkawano@mbp.ocn.ne.jp](mailto:kkawano@mbp.ocn.ne.jp)



**Figure 1** Pelvic computed tomography. (a) Right ovarian tumor: cystic tumor containing solid part (b) Left ovarian tumor: multicystic tumor containing solid part

with a papillary part suggestive of a malignant tumor (Fig. 1a). The left ovarian tumor showed a multi-cystic mass containing a solid part (Fig. 1b). Serum testing for tumor markers showed a carbohydrate antigen (CA)-19-9 level of 251.2 IU/mL, a CA-125 level of 283 IU/mL, and a carcinoembryonic antigen (CEA) level of 2.3 ng/mL. Dripped infusion of pyelography revealed a deviation of the running of the right ureter due to the pelvic mass. Mucosal invasion of the colon was negative, but barium enemas revealed deviation of the sigmoid colon due to the pelvic mass. Malignant ovarian tumor was suspected from preoperative examination.

At laparotomy, an enlarged uterus and bilateral ovarian tumors were revealed. The rectum was involved by the adhesion in the uterus and bilateral ovarian tumors. Only a small amount of ascites was observed and was collected for cytological examina-



**Figure 2** Macroscopic finding of excised uterus with bilateral tubes and ovaries.

tion. No metastatic or implanted lesions were observed in the abdominal cavity. The excised right ovarian tumor consisted of a mainly cystic part with some solid part, and a brown serous fluid was observed (Fig. 2). The left ovary was divided by the solid part and the cystic part. The solid part of the left ovarian tumor was firm, tan to yellow in color, and its cut surface was homogeneous. The frozen section gave rise to a diagnosis of clear cell adenocarcinoma of the right ovary and a sex cord tumor (e.g. granulosa cell tumor) of the left ovary. A total abdominal hysterectomy, bilateral salpingo-oophorectomy, pelvic lymphadenectomy, para-aortic lymph node sampling, omentectomy and appendectomy were performed. Pelvic and para-aortic lymph node metastases were not observed. Cytological examination of the cells in the ascites was negative. The excised uterus had an enlarged myometrium at the posterior wall without obvious tumor.

Histopathologically, papillary proliferation was observed inside the right ovarian tumor. The papillae were lined by cells that had a clear cytoplasm or showed a hobnail pattern (Fig. 3), which is the typical appearance of clear cell adenocarcinoma. The left ovarian tumor exhibited various histologic patterns. The solid part of the left ovarian tumor consisted of two carcinoid and thyroid components (Fig. 4a). The carcinoid was composed of uniform, medium-sized, cuboidal to columnar cells arranged in a trabecular pattern. The cells had round uniform nuclei that contained finely dispersed chromatin and inconspicuous nucleoli. The cytoplasm appeared granular and eosinophilic to amphophilic. The thyroid component demonstrated what appeared to be normal thyroid tissue. The size of the colloid was somewhat small where the