

Table 1. Distributions of TC cases diagnosed during 1993–1999 by histological classification and their relative 5-year survival

Classification	Testicular neoplasia (TC) (ICD-O-M: 9060–9102, 8000–8004, etc.)	Germ cell origin (TGCC)			Non-germ cell origin and NOS		
		Total (ICD-O-M: 9060–9102)	Seminoma (ICD-O-M: 9060–9064)	Non-seminoma (ICD-O-M: 9065–9102)	Total	Other specified cancers (ICD-O-M: 8140, etc.)	NOS (ICD-O-M: 8000–8004)
Number of cases	352	326	214	112	26	4 ¹	22
Relative survival rate at 5 years after diagnosis (%) (95% confidence interval)	88.9 (88.0–90.8)	91.0 (89.2–92.8)	95.6 (93.9–97.3)	82.2 (78.2–86.2)	62.5 (52.5–62.8)	33.4 (6.2–60.6)	66.9 (56.1–77.7)

¹All four cases were histopathologic ally identified to be adenocarcinoma. TC, testicular cancer; TGCC, testicular germ-cell cancer; NOS, non-germ cell origin or not otherwise specified; ICD-O-M, International Classification of Disease for Oncology in Morphology.

Table 2. Relative survival¹ of testicular germ-cell cancer (TGCC) and testicular cancer (TC) cases categorized according to hospital procedure volume

	Group A	Group B	Group C	Hospital unknown	Total number
Number of cases procured by hospital throughout the period	≥ 25	24 ≥, ≥ 8	7 ≥		
Number of patients: X ⁽ⁿ⁾	111 (116)	97 (101)	108 (117)	10 (15)	326 (352)
Number of deaths within 5-year period after diagnosis: Y ⁽ⁿ⁾	2 (4)	4 (5)	21 (25)	2 (5)	29 (38)
Proportion of deaths within 5 years after diagnosis: Y/X% ⁽ⁿ⁾	1.8 (3.4)	4.1 (5.0)	19.4 (20.5)	20.0 (27.8)	8.9 (10.8)
Relative survival rate at 5 years after diagnosis (%) ⁽ⁿ⁾	98.8 (96.9)	95.3 (94.3)	79.7 (79.4)	80.7 (68.6)	91.0 (88.9)
95% confidence interval ⁽ⁿ⁾	97.3–100.3 (96.0–98.8)	92.7–98.0 (91.5–97.1)	75.4–84.0 (75.3–83.5)	67.9–93.5 (57.4–79.8)	88.2–92.8 (88.0–90.8)

¹TC (including other specified and NOS, see Table 1).

Table 3. Crude and adjusted hazard ratios (HR) of testicular germ-cell cancer (TGCC) cases categorized according to hospital procedure volume

Hospital procedure volume	Crude HR ¹ (95% CI)	Adjusted HR ² (95% CI)
A	0.083 (0.019–0.352)	0.111 (0.025–0.495)
B	0.203 (0.070–0.593)	0.360 (0.120–1.084)
C	1	1

¹AIC = 277.7 (crude model), AIC = 217.7 (adjusted model). AIC, Akaike Information Criterion.

²Adjusted with clinical stage (distant/localized and regional), histology (non-seminoma/seminoma) and age at diagnosis.

for how smaller number of hospitals that the total TC patients) at quartile points using Armitage's test for trends in proportions and Gini's coefficient.⁽¹¹⁾

Statistical analysis. All statistical analysis was done using StatMate III (ATMS, Tokyo, Japan) and *P*-value for statistical significance was judged as less than 0.05.

Results

Table 1 shows distributions of TC cases diagnosed during 1993–1999 by histological classification and their relative 5-year survival. Most of TC cases were TGCC. Non-germ cell neoplasm and NOS cases accounted for less than 8%. Approximately two-thirds of TGCC cases were those of seminoma. The relative 5-year survival of TGCC cases was a little higher than that of the whole TC cases. The relative survival of seminoma cases was significantly higher than those of non-seminoma, non-germ cell origin or NOS.

Relative 5-year survival of TGCC cases was also calculated by hospital procedure volume (Table 2). Survival in group A and group B seemed significantly higher than that in group C (see 95% CI of TGCC in Table 2). Even when non-germ cell cancer and NOS cases were added to TGCC cases (TC cases), the difference was still remarkable.

Crude and adjusted hazard ratios (HR) of TGCC cases by hospital procedure volume were calculated using the Cox propor-

tional hazard model (Table 3). Hazard ratios for the cases treated in hospital group A and B were much lower than that for those in group C. Stage and age-adjusted HR for those treated in group A were also much lower than that for those in group C. Although the age-adjusted HR for those treated in group B was not significantly lower than that in group C, there seemed to be some association between hospital procedure volume and prognosis.

Hospital referral situation of the TC patients was evaluated in Osaka through three consecutive 4-year periods during 1990–2001 (Fig. 1, Table 4). There seemed to be no impressive change in referral centralization according to Gini's coefficient, or significant trends in the cumulative relative frequency of hospitals corresponding to the 2nd and the 3rd quartile of respective 4-year-period cases. Half of the cases were treated in hospitals with an experience volume of less than 3 persons per year. Furthermore, one-quarter of cases were treated in hospitals with a procedure volume of no more than 1 person per year during the period 1990–2001. In the zone between the 2nd and the 3rd quartiles, relative survival rate seemed to decrease remarkably (see hazard ratio of B/C in Table 3).

Discussion

Our previous study on the OCR database with respect to the prognosis of TC cases during 1975–1992 suggested that survival rates in smaller hospital were lower than those reported in SEER.⁽⁵⁾ The 5-year relative survival of TC in Osaka during 1993–1999 (91.0%) was remarkably higher than that reported during 1975–1992 (75.2%), which can be attributed to further diffusion of standard chemotherapy with cisplatin and subsequent combination of salvage chemotherapy methods, stereotactic radiotherapy and highly sensitive diagnostic modalities for detecting distant metastasis and so on. Nevertheless, 5-year relative survival of TC in Osaka during 1993–1999 remained lower than in all European countries (93.8%, 1995–1999)⁽⁷⁾ and SEER (95.7%, 1993–1995)⁽¹²⁾ registries. Hospital procedure volume was reported to potentially be a better index of different clinical outcome for some cancer sites.⁽¹³⁾ In accordance with that report, our findings with respect to the relative survival difference by hospital procedure volume

Table 4. Patients' referral situation through three consecutive 4-year periods, as to hospital procedure volume of testicular cancer (TC) cases in Osaka during 1990–2001

	1990–1993	1994–1997	1998–2001	P-value ⁴
Number of patients for respective period	200	182	180	
Number of hospitals for respective period	45	40	42	
Cumulative relative hospital frequency at 2nd quartile of the periodical patients' (%)	14.5	10.8	13.8	0.860
Patients number treated at the corresponding hospitals ¹	2.8	2.5	2.5	
Cumulative relative hospital frequency at 3rd quartile of the periodical patients' (%)	24.6	22.1	24.6	0.827
Patients number treated at the corresponding hospitals	1.0	1.0	1.0	
Gini's coefficient ²	0.58	0.60	0.57	

¹Linearly interpolated.

²Average number of patients treated per year at the hospitals corresponding to 2nd and 3rd quartile point respectively, see Figure 1.

³Armitage's χ^2 test for trend (two-sided).

⁴Proportion of space: S (surrounded by linear OB and respective curve)/ ΔOAB , see Figure 1.




Fig. 1. Relative cumulative frequency curve between testicular cancer cases and number of hospitals in order to evaluate the referral centralization among three consecutive 4-year periods: 1990–2001.

and the significantly lower adjusted hazard ratio of 'A/C' strongly suggest that the poor prognosis in hospital group C, where at most one or two patients per year were treated, may be attributed to immature experience.

Findings in our study are subject to at least the following five limitations.

Were there sufficient death cases to evaluate the survival? Incidence of TGCC is relatively rare and the cases have become ever more curable, consequently it is difficult to obtain many death cases even in a population as large as Osaka. In our study, statistically significant results were obtained for relative 5-year survival 'A, B versus C' and adjusted hazard ratio 'A/C'. However, the study would need to include more subjects to obtain sufficient statistical power to examine the ratio 'B/C'.

Do the OCR data reflect the current prognoses? As the current OCR database scheme requires 6–7 years to compile and report complete information of 5-year prognosis, the latest patients for whom relative survival rate can be obtained were diagnosed by 1999.

References

- 1 Einhorn LH, Donohue J. *cis*-Diamminedichloroplatinum, vinblastine and bleomycin combination chemotherapy in disseminated testicular cancer. *Ann Intern Med* 1977; 87: 293–8.
- 2 Motzer RJ, Scheinfeld J, Mazumdar M *et al*. Paclitaxel, ifosfamide, and cisplatin second-Line therapy for patients with relapsed testicular germ cell cancer. *J Clin Oncol* 2000; 18: 2413–8.

This scheme might be insufficient for evaluating more recent advancement in cancer treatments. The 'Period-analysis' method,⁽¹⁴⁾ which can reflect the recent advancements in diagnosis and treatment, will be needed for cancer registry system in Japan.

Are the OCR data sufficiently detailed? Some data items obtained from the current OCR system might contain insufficient detail, which complicated our interpretation. For example, 'chemotherapy' just meant initial treatment and did not include second-line chemotherapy, whether adjuvant, salvage, palliative or anything else.

Are there any other possible selection biases distorting the survival difference by hospital procedure volume? First, in the process of registration, the possible different character of 'death certificated only' (DCO) cases might distort the findings toward unpredictable direction. Actually, Cancer Incidence in Five Continents VIII⁽¹⁵⁾ showed 2% DCO for TC in Osaka. Second, if the completeness of the registry was different among the three groups, it also might affect some unpredictable bias, possibly toward overestimation. Nevertheless, we believe that it would not have so much magnitude as to affect our conclusion greatly. Third, there existed some NOS cases of histology whose survival (66.9%) was lower than that of TGCC (91.0%). We cannot tell what effect these cases might have on our findings of survival difference by hospital procedure volume. The proportion was less than 7% of total cases, therefore the distortion might be small.

Is the patients' hospital referral situation adequately evaluated? Trends in hospital referrals can be usefully quantified by Gini's coefficient, despite the fact that the coefficient is generally cited for socioeconomic analysis of disparity trends.⁽¹¹⁾ Though the quantification may not be warranted if the numbers of hospitals differ largely in the respective period, those considered here were almost the same. The coefficient may be worthwhile for the evaluation of overall referral situation as well as that in details at the quartile points.

In conclusion, the prognosis of TC in Osaka has remarkably improved, but it still remains lower than those in the EU and/or USA. TC cases in Osaka should be treated collectively in more experienced hospitals like those in group A, especially for cases with advanced disease.

- 3 Mencil PJ, Motzer RJ, Mazumdar M, Vlamis V, Bajorin DF, Bosl GJ. Advanced seminoma: treatment results, survival, and prognostic factors in 142 patients. *J Clin Oncol* 1994; 12: 120–6 (1994 deleted).
- 4 Hain SF, O's Doherty MJ. Fluorodeoxyglucose positron emission tomography in the evaluation of germ cell tumours at relapse. *Br J Cancer* 2000; 83: 863–9.
- 5 Ries LAG, Eisner MP, Kosary CL *et al*. *SEER Cancer Statistics Review, 1973–98*. Bethesda, MD: National Cancer Institute, 2001.

- 6 Oshima A, Kitagawa T, Ajiki W, Tsukuma H, Takenaka S, Iura A. Survival of testicular cancer patients in Osaka Japan. *Jpn J Clin Oncol* 2001; 31: 438-43.
- 7 Berrino F, Angelis RD, Sant M *et al*. Survival for eight major cancers and all cancers combined for European adults diagnosed in 1995-99: results of the Eurocare-4 study. *Oncol Lancet Com* 2007; 8: 773-83.
- 8 Esteve J, Benhamou E, Raymond L. *Statistical Methods in Cancer Research*, Vol. IV *Descriptive Epidemiology*. Lyon: IARC, 1994; 231-45.
- 9 Editors in Japan National Cancer Center. Survival probability table. [Cited 20 March 2008.] Available from URL: <http://ganjoho.ncc.go.jp/professional/statistics/cohort01.html>
- 10 Akaike H, Amari T, Kitagawa G, Shimohira H. *Akaike's Information Criterion (AIC)* [in Japanese]. Tokyo: Kyoritsu-syuppan, 2007.
- 11 Uwafuji I, Morimoto E, Tsunekane M. *Statistics for Investigation and Analysis* [in Japanese]. Tokyo: Maruzen, 2006.
- 12 Editors in USA. National Cancer Institute. *Surveillance Epidemiology and End Results Cancer Statistics Review*. [Cited 20 March 2008.] Available from URL: http://seer.cancer.gov/csr/1975_2004/
- 13 Ioka A, Tsukuma H, Ajiki W, Oshima A. Hospital procedure volume and survival of cancer patients in Osaka, Japan: a population-based study with latest cases. *Jpn J Clin Oncol* 2007; 37: 544-53.
- 14 Brenner H, Hakulinen T. Period versus cohort modeling of up-to-date cancer survival. *Int J Cancer* 2007; 122: 898-904.
- 15 Parkin DM, Whelan SL, Ferlay J, Teppo L, Thomas DB. *Cancer Incidence in Five Continents*, Vol. III. Lyon: IARC. 2003; 264-5.

Cancer survival in five continents: a worldwide population-based study (CONCORD)



Michel P Coleman, Manuela Quaresma, Franco Berrino, Jean-Michel Lutz, Roberta De Angelis, Riccardo Capocaccia, Paolo Baili, Bernard Rachet, Gemma Gatta, Timo Hakulinen, Andrea Micheli, Milena Sant, Hannah K Weir, J Mark Elwood, Hideaki Tsukuma, Sergio Koifman, Gulnar Azevedo e Silva, Silvia Francisci, Mariano Santaquilani, Arduino Verdecchia, Hans H Storm, John L Young, and the CONCORD Working Group*

Summary

Background Cancer survival varies widely between countries. The CONCORD study provides survival estimates for 1.9 million adults (aged 15–99 years) diagnosed with a first, primary, invasive cancer of the breast (women), colon, rectum, or prostate during 1990–94 and followed up to 1999, by use of individual tumour records from 101 population-based cancer registries in 31 countries on five continents. This is, to our knowledge, the first worldwide analysis of cancer survival, with standard quality-control procedures and identical analytic methods for all datasets.

Methods To compensate for wide international differences in general population (background) mortality by age, sex, country, region, calendar period, and (in the USA) ethnic origin, we estimated relative survival, the ratio of survival noted in the patients with cancer, and the survival that would have been expected had they been subject only to the background mortality rates. 2800 life tables were constructed. Survival estimates were also adjusted for differences in the age structure of populations of patients with cancer.

Findings Global variation in cancer survival was very wide. 5-year relative survival for breast, colorectal, and prostate cancer was generally higher in North America, Australia, Japan, and northern, western, and southern Europe, and lower in Algeria, Brazil, and eastern Europe. CONCORD has provided the first opportunity to estimate cancer survival in 11 states in USA covered by the National Program of Cancer Registries (NPCR), and the study covers 42% of the US population, four-fold more than previously available. Cancer survival in black men and women was systematically and substantially lower than in white men and women in all 16 states and six metropolitan areas included. Relative survival for all ethnicities combined was 2–4% lower in states covered by NPCR than in areas covered by the Surveillance Epidemiology and End Results (SEER) Program. Age-standardised relative survival by use of the appropriate race-specific and state-specific life tables was up to 2% lower for breast cancer and up to 5% lower for prostate cancer than with the census-derived national life tables used by the SEER Program. These differences in population coverage and analytical method have both contributed to the survival deficit noted between Europe and the USA, from which only SEER data have been available until now.

Interpretation Until now, direct comparisons of cancer survival between high-income and low-income countries have not generally been available. The information provided here might therefore be a useful stimulus for change. The findings should eventually facilitate joint assessment of international trends in incidence, survival, and mortality as indicators of cancer control.

Funding Centers for Disease Control and Prevention (Atlanta, GA, USA), Department of Health (London, UK), Cancer Research UK (London, UK).

Introduction

International comparisons of population-based cancer survival have been rare,^{1,2} but large and unexplained differences in survival have been reported for many cancers from individual studies and cancer registries in Europe and North America.³ For example, 5-year relative survival for women diagnosed with breast cancer during 1985–89 was 73% in Europe (weighted mean for 17 countries)⁴ and 84% in the USA.⁵ The CONCORD study provides a systematic comparison of survival between Europe and North America,^{6–8} extended to countries in all other continents.

The first international comparison of cancer survival, published in 1964,⁹ was a study of patients diagnosed with one of 15 common cancers in Denmark, England, Finland, France, Norway, Sweden, and the USA, mainly during

1945–54. It was the first study in which relative survival techniques, first described in the 1950s,^{10–12} were used to correct the survival estimates for differences in background mortality between participant countries. The findings are mainly of historical interest, but survival in the USA (represented by Connecticut) was generally higher than in the European countries.

Cancer survival is known to vary between the regions of the USA covered by the US National Cancer Institute's (NCI) Surveillance, Epidemiology and End Results (SEER) Program,¹³ but the range of survival in Europe is much wider. Furthermore, survival from breast cancer during 1985–94 was higher in each of the nine SEER areas than in any of the 22 countries participating in the European study of cancer survival (EUROCORE).¹⁴ The differences were

Published Online
July 17, 2008
DOI:10.1016/S1470-2045(08)70179-7

*Members of the CONCORD Working Group are listed in the webappendix

Cancer Research UK Cancer Survival Group, Non-Communicable Disease Epidemiology Unit, London School of Hygiene and Tropical Medicine, London, UK (Prof M P Coleman FPPH, M Quaresma MSc, B Rachet MD); Department of Preventive and Predictive Medicine (F Berrino MD, G Gatta MD, M Sant MD), and Descriptive Epidemiology and Health Planning Unit (P Baili PhD, A Micheli PhD), Fondazione IRCCS Istituto Nazionale Tumori, Milan, Italy; Geneva Cancer Registry, Geneva, Switzerland (J-M Lutz MD); National Centre for Epidemiology, Surveillance and Health Promotion, Department of Cancer Epidemiology, Istituto Superiore di Sanità, Rome, Italy (R De Angelis BSc, R Capocaccia PhD, S Francisci PhD, M Santaquilani PhD, A Verdecchia PhD); Finnish Cancer Registry, Helsinki, Finland (Prof T Hakulinen PhD); Division of Cancer Prevention and Control, Centers for Disease Control and Prevention, Atlanta, GA, USA (H K Weir PhD); British Columbia Cancer Agency, Vancouver, BC, Canada (Prof J M Elwood MD); Osaka Cancer Registry, Department of Cancer Control and Statistics, Osaka Medical Centre for Cancer and Cardiovascular Diseases, Osaka, Japan (H Tsukuma MD); Department of Epidemiology, National School of Public Health, Oswaldo Cruz Foundation, Ministry of Health, Rio de Janeiro, Brazil (S Koifman PhD); Institute of Social Medicine, University of Rio de Janeiro,

Rio de Janeiro, Brazil
(G Azevedo e Silva PhD);
Department of Cancer
Prevention and
Documentation, Danish Cancer
Society, Copenhagen, Denmark
(HH Storm MD); Metropolitan
Atlanta SEER Registry, Georgia
Center for Cancer Statistics,
Department of Epidemiology,
Rollins School of Public Health
at Emory University, Atlanta,
GA, USA (Prof J L Young PhD)

Correspondence to:
Prof Michel P Coleman,
Cancer Research UK Cancer
Survival Group, Non-
Communicable Disease
Epidemiology Unit, London
School of Hygiene and Tropical
Medicine, London WC1E 7HT, UK
michel.coleman@lshtm.ac.uk

See Online for webfigure 1

often more marked in elderly patients;⁸ for several cancers, 5-year survival for patients diagnosed aged 75 years or older during the 1990s was nearly 20% higher in the USA than in Europe.²¹

The CONCORD study began in 1999 as an extension of the EUROCARE-3 study, then just starting. EUROCARE has published systematic comparisons of survival for most adult and childhood cancers in Europe since 1995.²⁴ The first EUROCARE study involved patients diagnosed in 1978–84 in 12 countries;²⁵ EUROCARE-2 covered patients diagnosed during 1985–89 in 17 countries,²⁶ and EUROCARE-3 involved 22 countries, with patients diagnosed in 1990–94 and followed up to 1999.^{27,28} More recently, EUROCARE-4 has included patients diagnosed in 23 countries during all or part of 1995–2002 and followed up to 2003.^{29,30}

CONCORD was originally designed to assess the survival of adults (aged 15–99 years) diagnosed with cancer of the breast (women), colorectum, or prostate during 1990–94 in Europe and the USA, using population-based data and standardised quality control, and with identical analysis for all datasets, adjusted for differences in general population (background) mortality by country, region, race, and calendar period, and also for differences in the age structure of patient populations. CONCORD also enables comparison of cancer survival between five states and four metropolitan areas in the USA covered by the SEER Program (SEER-9) and 11 states covered by the Centers for Disease Control and Prevention's (CDC) National Program of Cancer Registries (NPCR). It also provides a wider comparison of cancer survival between black and white patients in the USA than has previously been possible.

CONCORD includes data from one or more countries on all five continents. To our knowledge, it is the first attempt at a global comparison of cancer survival.

Methods

Cancer registries

In 1999, we identified at international cancer meetings in Atlanta (USA) and Lisbon (Portugal), and from published studies, population-based cancer registries that had published survival data and were operational during 1990–99. Registries that had met the quality criteria for inclusion in *Cancer Incidence in Five Continents* (volume VII, 1988–92)³¹ were eligible. We obtained data from 19 other registries. Most had met comparable criteria, such as those in the EUROCARE-3 study (patients diagnosed during 1990–94 with follow-up to 1999).²⁸ North American registries were eligible if they had met the standards required for Cancer Incidence in North America, 1991–95,³² and could provide complete follow-up to the end of 1999. In total, we identified 112 registries, but 11 were withdrawn or excluded: no response (one); withdrawal for legal reasons (one); incomplete registration before 1995 (four); follow-up activity stopped before 1999 (two); data not supplied by the September, 2005 deadline (three).

A pilot study of 50 registries in 2000 obtained a 100% response. All registries were able to provide data for

patients diagnosed during all or part of the period 1990–94, and had access to various data sources to obtain follow-up information for all patients for at least 5 years or to the end of 1999. After further recruitment, a detailed questionnaire was obtained for 100 of the 101 registries finally included in the analyses, covering data definitions and methods of operation, including data collection, coding of tumour site, morphology, behaviour, and stage at diagnosis, tracing of registered patients to ascertain their vital status, and linkage between data on the incident tumour and data on subsequent death or loss to follow-up. The procedures and definitions used, the stated quality and completeness of data on the registration of incident cancers, and of the follow-up of those patients over the next 5 years, were deemed adequate to attempt cancer-survival analysis, subject to central quality control of the data. The pilot study confirmed the feasibility of the CONCORD protocol³³ and the active support of cancer registries for wider international comparisons of cancer survival. The questionnaire and detailed findings are available online.³⁴

Data sources

Anonymised individual tumour records were obtained from population-based cancer registries in all five continents, as defined on UN guidelines:³⁵ Africa, America (Central and South, including the Caribbean), America (North), Asia, Europe, and Oceania (table 1 and webfigure 1). We retained Hawaii (USA) with North America rather than Oceania.

Africa was represented by a single cancer registry, for the wilaya (département, or state) of Sétif (Algeria).

Central and South America, including the Caribbean, were represented by the national cancer registry of Cuba and two regional registries in Brazil: the Goiânia (Goiás state) registry is one of 20 registries in state capitals, whereas the Campinas (São Paulo state) registry is the only one in Brazil that is not in a state capital.

Data from North America include five of the seven largest provinces in Canada (British Columbia, Manitoba, Nova Scotia, Ontario, and Saskatchewan). Data for the USA came from 22 registries covering 16 states (California, Colorado, Connecticut, Florida, Hawaii, Idaho, Iowa, Louisiana, Michigan, Nebraska, New Jersey, New Mexico, New York State, Rhode Island, Utah, and Wyoming) and six metropolitan areas (Atlanta, GA, Los Angeles, CA, San Francisco, CA, Detroit, MI, New York City, NY, and Seattle, WA).

Population-based cancer registries in the USA receive support from either or both of the two federal cancer-surveillance programmes, the NCI's SEER Program and the CDC's NPCR.³⁶ As of 1990, the SEER Program included nine population-based cancer registries covering some 10% of the US population (SEER-9): the states of Connecticut, Hawaii, Iowa, New Mexico, and Utah, and the metropolitan areas of Atlanta, GA, Detroit, MI, San Francisco, CA, and Seattle, WA. The Los Angeles cancer registry became a SEER registry in 1992, but we opted to retain it with the NPCR data, so that the SEER grouping

we used was identical with that for which SEER data had been published in the past (SEER-9). The NPCR at the CDC began more recently, and this is the first cancer-

survival analysis for 11 states: California, Colorado, Florida, Idaho, Louisiana, Michigan, Nebraska, New Jersey, New York, Rhode Island, and Wyoming.

	Population covered by registry	% of national population	Breast		Colon		Rectum		Colorectum			Prostate	Total
			Women	Men	Women	Men	Women	Men	Women	Men	Women		
Africa													
Algeria (Sétif)	1104 561	4.2	180	10	14	30	30	40	44	36	300		
America (Central and South)													
Brazilian registries	1795 387	1.2	806	130	194	50	69	180	263	474	1723		
Campinas	870 380	0.6	175	61	82	—	—	—	—	149	467		
Goiania	925 007	0.6	631	69	112	50	69	119	181	325	1256		
Cuba	10 754 868	100.0	6461	1083	1516	674	734	1757	2250	4341	14 809		
South American registries	12 550 255	—	7267	1213	1710	724	803	1937	2513	4815	16 532		
America (North)													
Canadian registries	16 474 543	58.1	44 620	13 989	13 819	6272	4220	20 261	18 039	45 999	128 919		
British Columbia	3 131 700	11.0	9141	2223	2178	625	412	2848	2590	11 496	26 075		
Manitoba	1 109 998	3.9	2932	954	957	556	343	1510	1300	3761	9503		
Nova Scotia	918 000	3.2	2316	771	829	—	—	—	—	2243	6159		
Ontario	10 298 801	36.3	27 389	9214	9069	4613	3154	13 827	12 223	25 310	78 749		
Saskatchewan	1 016 044	3.6	2842	827	786	478	311	1305	1097	3189	8433		
US registries	108 775 729	42.4	324 551	89 673	96 186	40 149	32 774	129 822	128 960	356 881	940 214		
Atlanta,† GA	2 315 961	0.9	5747	1215	1473	474	496	1689	1969	6406	15 811		
California	30 974 659	12.1	85 143	21 384	22 351	9999	8172	31 383	30 523	95 707	242 756		
Los Angeles, CA	9 055 424	—	22 587	5741	6136	2659	2233	8400	8369	25 789	65 145		
San Francisco, CA	3 805 588	—	12 321	3165	3375	1463	1194	4628	4569	12 733	34 251		
Colorado	3 495 939	1.4	9117	2084	2183	944	751	3028	2934	11 433	26 512		
Connecticut	3 300 712	1.3	11 335	3112	3299	1458	1128	4570	4427	11 357	31 689		
Florida	13 650 553	5.3	46 065	14 845	15 007	6007	4790	20 852	19 797	64 256	150 970		
Hawaii	1 158 613	0.5	2857	986	808	508	279	1494	1087	3482	8920		
Idaho	1 071 685	0.4	2689	676	681	331	239	1007	920	3899	8515		
Iowa	2 818 401	1.1	9133	2776	3522	1267	989	4043	4521	10 743	28 440		
Louisiana	4 293 003	1.7	11 204	3302	3780	1374	1186	4676	4966	13 059	33 905		
Michigan	9 479 065	3.7	31 183	8821	9323	3791	3162	12 612	12 485	23 705	79 985		
Detroit, MI	3 969 304	—	12 247	3223	3534	1499	1213	4722	4747	17 162	38 878		
Nebraska	1 611 687	0.6	5242	1625	1801	776	544	2401	2345	6828	16 816		
New Jersey	7 880 508	3.1	27 125	8110	8670	3694	3091	11 804	11 761	29 877	80 567		
New Mexico	1 595 442	0.6	3796	901	892	436	323	1337	1215	5393	11 741		
New York State	18 246 653	7.1	55 404	15 191	17 426	6936	5889	22 127	23 315	47 096	147 942		
New York City	7 322 564	—	21 644	5821	7048	2335	2253	8156	9301	16 770	55 871		
Rhode Island	1 012 581	0.4	3466	1113	1280	477	440	1590	1720	3449	10 225		
Seattle,† WA	3 567 217	1.4	10 451	2415	2577	1168	893	3583	3470	12 818	30 322		
Utah	1 836 799	0.7	3506	866	805	393	293	1259	1098	5779	11 642		
Wyoming	466 251	0.2	1088	251	298	116	109	367	407	1594	3456		
North American registries	125 250 272	44.0	369 171	103 662	110 005	46 421	36 994	150 083	146 999	402 880	1 069 133		
Asia													
Japanese registries	10 819 997	8.7	7179	5469	4588	3510	2248	8979	6836	1691	24 685		
Fukuji	827 000	0.7	840	738	709	477	310	1215	1019	325	3399		
Osaka	8 734 516	7.0	5112	3337	2593	2075	1283	5412	3876	920	15 320		
Yamagata	1 258 481	1.0	1227	1394	1286	958	655	2352	1941	446	5966		

(Continues on next page)

	Population covered by registry	% of national population	Breast		Colon		Rectum		Colorectum		Prostate	Total
			Women	Men	Women	Men	Women	Men	Women			
(Continued from previous page)												
Europe												
Austria (Tirol)	624939	8.0	1559	416	483	261	237	677	720	1432	4388	
Czech Republic (West Bohemia)	861000	8.3	1543	672	601	681	416	1353	1017	693	4606	
Denmark	5145160	100.0	14686	3954	4822	3308	2495	7262	7317	6503	35768	
Estonia	1562468	100.0	2205	598	845	479	553	1077	1398	1143	5823	
Finland	5070000	100.0	12214	1907	2639	1687	1561	3594	4200	7544	27552	
French registries	3098526	5.6	6359	1675	1544	1164	876	2839	2420	2909	14527	
Bas-Rhin	954710	1.8	2591	848	730	522	379	1370	1109	1626	6696	
Calvados	618353	1.1	1640	440	448	345	309	785	757	1283	4465	
Côte d'Or	507147	0.9	791	387	366	297	188	684	554	..	2029	
Isère	1018316	1.8	1337	1337	
Germany (Saarland)	1067027	1.3	2957	1035	1237	712	656	1747	1893	1610	8207	
Iceland	254960	100.0	504	125	128	37	47	162	175	493	1334	
Ireland	3609000	100.0	1513	587	534	382	224	969	758	1062	4302	
Italian registries	8944772	15.3	26403	8713	8672	4743	3887	13456	12559	10671	63089	
Ferrara	355479	0.6	1321	488	486	200	158	688	644	438	3091	
Genoa	695981	1.3	2571	892	894	442	380	1334	1274	1122	6301	
Latina	468865	0.8	657	199	182	135	84	334	266	197	1454	
Macerata	281537	0.5	629	296	283	168	119	464	402	435	1930	
Modena	602570	0.5	1887	641	654	361	275	1002	929	810	4628	
Parma	391237	0.7	1318	480	410	256	204	736	614	456	3124	
Ragusa	140537	0.5	513	159	171	123	82	282	253	227	1275	
Romagna	604488	0.8	1347	498	549	226	226	724	775	740	3586	
Sassari	469570	0.8	591	143	128	126	62	269	190	198	1248	
Turin	996443	1.8	3009	868	904	500	457	1368	1361	1030	6768	
Tuscany	1167687	2.1	3807	1420	1446	854	702	2274	2148	1797	10026	
Varese	793378	1.4	2400	691	710	410	344	1101	1054	803	5358	
Veneto	1977000	3.5	6353	1938	1855	942	794	2880	2649	2418	14300	
Malta	365000	100.0	359	76	73	53	31	129	104	111	703	
Netherlands registries	5158472	34.3	15862	2418	2791	1471	1271	3889	4062	5353	29166	
Amsterdam	2620000	17.4	7509	1764	2117	1020	946	2784	3063	4171	17527	
Netherlands (North)	1602661	10.6	5999	5999	
Netherlands (South)	935811	6.3	2354	654	674	451	325	1105	999	1182	5640	
Norway	4245180	100.0	9193	3590	4136	2536	2048	6126	6184	9841	31344	
Polish registries	2373190	6.1	4220	1080	1152	827	773	1907	1925	1159	9211	
Cracow	747985	1.9	1205	240	243	203	168	443	411	253	2312	
Warsaw	1625205	4.2	3015	840	909	624	605	1464	1514	906	6899	
Portugal (South)	1145000	11.4	1219	364	355	327	236	691	591	344	2845	
Slovakia	5297774	100.0	6079	2572	2126	2646	1815	5218	3941	2821	18059	
Slovenia	2072000	100.0	3327	914	898	1025	851	1939	1749	160	8175	
Spanish registries	5566140	14.4	9744	3439	2934	2502	1613	5941	4547	4273	24505	
Basque Country	2097000	5.4	3816	1321	1027	1057	589	2378	1616	1721	9531	
Granada	787898	2.0	879	299	255	219	152	518	407	..	1804	
Mallorca	582655	1.5	1143	447	394	296	213	743	607	617	3110	
Murcia	1036966	2.8	1485	505	512	397	330	902	842	643	3872	
Navarra	520300	1.3	1229	404	304	249	167	653	471	688	3041	
Tarragona	541321	1.4	1192	463	442	284	162	747	604	604	3147	

(Continues on next page)

Survival estimates reported from the SEER Program have until now been the only population-based cancer survival data from the USA.^{31,37} We wanted to compare survival between the areas covered by registries in the NPCR and the SEER Program during 1990-94. We received separate datasets from Detroit, MI, San Francisco, CA (SEER registries), and Los Angeles, CA (NPCR), and these were included in the respective totals for SEER and NPCR.

However, the data from these metropolitan areas could not be separately identified in the state-wide datasets we received from California and Michigan, therefore, the non-metropolitan data for those states could not be included with the other NPCR data. Data from all nine SEER registries were available.³⁸

Survival in the SEER-9 areas was therefore compared with survival in nine states and one metropolitan area covered by

	Population covered by registry	% of national population	Breast		Colon		Rectum		Colorectum		Prostate	Total
			Women	Men	Women	Men	Women	Men	Women			
(Continued from previous page)												
Sweden	8 826 939	100.0	24 170	6 112	6 685	4 401	3 578	10 513	10 263	24 041	68 987	
Swiss registries	1 758 249	25.8	4 847	--	--	--	--	--	--	--	4 847	
Basel	429 104	6.3	1 365	--	--	--	--	--	--	--	1 365	
Geneva	381 492	5.6	1 275	--	--	--	--	--	--	--	1 275	
Graubünden-Glarus	210 485	3.1	544	--	--	--	--	--	--	--	544	
St Gallen-Appenzel	483 801	7.1	1 027	--	--	--	--	--	--	--	1 027	
Valais	253 367	3.7	636	--	--	--	--	--	--	--	636	
UK	58 984 046	--	154 867	41 499	45 729	30 600	22 556	72 099	68 285	78 608	373 859	
England (national)	49 310 000	100.0	129 703	33 983	37 334	25 618	18 780	59 601	56 114	66 181	311 599	
East Anglia	2 089 000	4.2	6 330	1 820	2 060	1 245	954	3 065	3 014	3 897	16 306	
Mersey	2 412 000	4.9	6 561	1 932	2 080	1 425	1 069	3 357	3 149	3 242	16 309	
Oxford	2 582 000	5.2	7 458	1 737	1 934	1 193	929	2 930	2 863	3 612	16 863	
South Thames	6 756 000	13.7	17 002	3 880	4 689	2 824	2 328	6 704	7 017	8 232	38 955	
South West	3 320 000	6.7	19 203	5 630	6 215	3 869	2 917	9 499	9 132	11 766	49 600	
Trent	4 745 000	9.6	13 360	3 523	3 793	3 045	2 087	6 568	5 880	6 774	32 582	
West Midlands	5 278 000	10.7	13 561	4 397	4 482	3 272	2 066	7 669	6 548	7 315	35 093	
Yorkshire	3 698 000	7.5	9 473	2 599	2 910	2 121	1 574	4 720	4 484	5 165	23 842	
English registries	30 880 000	62.5	92 948	25 518	28 163	18 994	13 924	44 512	42 087	50 003	229 550	
Northern Ireland	1 648 960	100.0	1 527	562	576	328	224	890	800	888	4 105	
Scotland	5 100 086	100.0	14 254	4 441	5 089	2 671	2 124	7 112	7 213	6 855	35 434	
Wales	2 925 000	100.0	9 383	2 513	2 730	1 983	1 428	4 496	4 158	4 684	22 721	
European registries	126 029 842	--	303 830	81 746	88 384	59 842	45 724	141 588	134 108	161 771	741 297	
Oceania												
Australia	18 071 422	100.0	41 090	15 200	15 098	9 911	6 904	25 111	22 002	42 890	131 093	
Australian Capital Territory	304 371	1.7	548	180	160	99	78	279	238	414	1 479	
New South Wales	6 133 913	33.9	14 382	5 358	5 066	3 478	2 354	8 836	7 420	15 507	46 145	
Northern Territory	178 062	1.0	165	46	41	41	20	87	61	78	391	
Queensland	3 252 245	18.0	7 052	2 783	2 743	1 619	998	4 402	3 741	7 468	22 663	
Southern Australia	1 473 966	8.2	3 688	1 323	1 335	937	734	2 260	2 069	4 228	12 245	
Tasmania	472 971	2.6	1 081	474	453	242	171	716	624	1 321	3 742	
Victoria	4 521 392	25.0	10 583	3 865	4 103	2 683	1 978	6 548	6 081	9 826	33 038	
Western Australia	1 734 502	9.6	3 591	1 171	1 157	812	571	1 983	1 768	4 048	11 390	
CONCORD												
CONCORD total	293 826 349	--	728 717	207 300	219 799	120 438	92 703	327 738	312 502	614 083	1 983 040	

*Some registries provided data for shorter periods: ie, 4 years: Campinas, Macerata, Granada (1991-94); 3 years: Isère (1990-92); Portugal (1991-93); Serrif, Sassari (1992-94); 2 years: Malta, Northern Ireland (1993-94); 1 year: Ireland (1994). No state-wide data available for this city. Where a registry did not provide data for a given cancer, cell entries for numbers of patients and survival estimates are left blank. National percentages are derived from the raw data and can differ from the sum of regional percentages because of rounding. Row totals avoid double counting of colon and rectal tumours; also shown in the table as colon and rectum combined.

Table 1: Population coverage and number of adults (aged 15-99 years) diagnosed with cancer of the breast, colon, rectum, or prostate during 1990-94* and included in the analyses: continent, country, and region

NPCR: Colorado, Florida, Idaho, Los Angeles, CA, Louisiana, Nebraska, New Jersey, New York, Rhode Island, and Wyoming. For this comparison, data from the non-metropolitan areas of California and Michigan were excluded to ensure that the two sets of data were mutually exclusive.

In Asia, Japan was represented by three of the prefectural (state) registries: Fukui, Osaka, and Yamagata.

In Europe, the 53 cancer registries that contributed data to EURO-CARE-3²⁸ on cancers of the breast, colon, rectum, or prostate all participated in the CONCORD study. Six other registries also provided data: two national registries (Northern Ireland and Ireland) and four regional registries from the Netherlands (North) and Switzerland (Graubunden-Glarus, St Gallen-Appenzell, Valais). As in the EURO-CARE study, the UK is considered as its four constituent countries (England, Scotland, Wales, Northern Ireland), each of which has a national registry. In England, both the national cancer registry and eight of the regional cancer registries submitted datasets.

Oceania was represented by the national cancer registry of Australia, with data from each of the eight population-based state or territorial registries.

Quality control

Procedures used in the EURO-CARE-3 study were applied to all datasets. Tumour records were supplied with the anatomical site coded to the ninth revision of the International Classification of Diseases (ICD-9²⁹) for four index tumours: cancers of the breast (women) (ICD-9 174.0–174.9), colon (153.0–153.9), rectum (including the anus, 154.0–154.9), and prostate (185). Tumour morphology and behaviour were coded to the first or second revision of ICD-Oncology (ICD-O,³⁰ ICD-O-2³¹). Only invasive malignant tumours (behaviour code 3) were included. Patients with an index tumour had sometimes been registered with another malignancy, either before or after the index tumour. Data on those other cancers in index patients were also submitted. Only the first, primary, invasive, malignant tumour diagnosed in each patient was retained for analysis. Patients registered with a malignant neoplasm before the index tumour were excluded, although non-melanoma skin cancer was not counted as a previous tumour for this purpose. Bilateral breast cancers and multiple colon cancers were included as a single tumour if synchronous; otherwise, only the earliest tumour was considered. The duration of survival was taken from the date of diagnosis of the index tumour until death from any cause, or until the patient was censored from the analysis as alive, either at loss to follow-up or after Dec 31, 1999, whichever came first; any subsequent tumour occurring in the same patient during that period was ignored.

Standard quality-control routines, based on those developed by the International Agency for Research on Cancer,³² were applied to each tumour record. Records with invalid codes, impossible sequences of dates, or improbable combinations of tumour site and morphology were returned to the registry for checking. Usually, the registry provided a

correction or an explanation. Corrected tumour records were checked again: those which still had missing, invalid or inconsistent values for sex, site, morphology, or dates were flagged as major errors and excluded from analysis. Records for which an unlikely combination of age, site and morphology had nonetheless been confirmed as correct were flagged as minor errors, and included in the analyses. Details of the approach have been published elsewhere.³³ Detailed quality-control findings are available online.³⁴

Follow-up

All registries used more than one mechanism of follow-up to ascertain the vital status (alive, dead, emigrated, lost to follow-up) and the date of the last vital status for each registered patient. The mechanisms varied between countries, usually linkage between the registry's database and a variety of other data sources, especially the national index of deaths. Secure linkage of a tumour record and a record of death, based on a set of identifiers such as name, sex, date of birth, and personal identity number, enabled the registry to update the tumour record accordingly. Direct contact with the patient or their family to establish vital status was unusual, although home visits by registry staff were done in Algeria. Enquiries to the patient's primary care physician or hospital consultant were frequently used. A wide variety of administrative databases was also used, such as social insurance, health insurance, motor vehicle records, drivers' licences, hospital discharge records, national primary-care databases, electoral registers (those eligible to vote), and voter registration records (those who voted in the last election). The presence of a person's record in such administrative databases on a given date is taken as evidence that the person was alive on that date. This is subject to administrative error (failure to remove in timely fashion the record of a person known to be dead) and fraud (by someone seeking to retain access to benefits received by the deceased), but in most instances the risks are small. If coverage of the databases was known to be high, and especially if a person was present in more than one such database, the risk of error decreased further.

In the USA, a match to an administrative database might show that an event occurred during a certain quarter of a year (eg, an insurance claim paid, a licence renewed), but the exact date might not be known; the date of last vital status was then set to the first day of the quarter, ie, Jan 1, April 1, July 1, Sept 1. This approach can give rise to irregular distributions of the day of last known vital status, but it is a conservative approach to establishing when patients were last known to be alive, because patients are censored from survival analysis on the latest of any such dates in the record.

The proportion of patients not known to be dead and for whom the registry could not be certain that the date of last vital status was at least 5 years after diagnosis was less than 1% overall. The proportion was often zero (follow-up for at least 5 years was established for every patient not known to be dead), the highest proportion was 4%, and only in a

	Breast	Colon	Rectum		Colorectum		Prostate	
	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	RS (%) [95% CI]
Africa								
Algeria (Sétif)	38.8 (31.4-46.2) R	11.4 (0.7-40.9) R	30.6 (9.5-56.1) R	25.9 (11.4-43.7) R	18.2 (6.6-34.6) R	22.5 (10.6-37.7) R	22.6 (11.2-36.7) R	21.4 (8.7-38.9) R
America (Central and South)								
Brazilian registries	58.4 (52.7-64.6)	33.1 (24.2-45.3)	32.7 (26.1-40.8)	49.3 (34.8-69.8)	38.4 (27.3-53.9)	47.3 (37.5-59.6)	43.5 (35.7-53.1)	49.3 (43.6-55.8)
Campinas	36.6 (27.8-48.3)	23.8 (13.1-36.8) R	21.4 (12.6-31.9) R	34.4 (25.2-47.0)
Goiânia	65.4 (58.3-73.2)	48.1 (36.7-63.1)	44.8 (35.2-56.9)	49.3 (34.8-69.8)	38.4 (27.3-53.9)	47.3 (37.5-59.6)	43.5 (35.7-53.1)	55.7 (49.0-63.3)
Cuba	84.0 (82.9-85.2)	59.3 (55.8-63.1)	61.4 (58.3-64.5)	59.2 (55.1-63.7)	62.8 (58.6-67.4)	59.5 (56.8-62.5)	62.0 (59.5-64.6)	69.7 (67.1-72.3)
America (North)								
North American registries	83.7 (83.5-83.9)	59.5 (59.1-59.9)	59.9 (59.5-60.3)	56.4 (55.8-56.9)	59.7 (59.1-60.3)	58.6 (58.3-58.9)	60.0 (59.7-60.3)	91.1 (90.9-91.3)
Canadian registries								
British Columbia	82.5 (81.9-83.0)	56.1 (55.1-57.2)	58.7 (57.7-59.7)	53.1 (51.5-54.6)	58.7 (57.0-60.4)	55.3 (54.4-56.2)	58.9 (58.0-59.8)	85.1 (84.4-85.7)
Manitoba	85.4 (84.2-86.5)	57.0 (54.5-59.6)	59.2 (56.8-61.7)	64.6 (59.9-69.7)	62.8 (57.5-68.6)	58.7 (56.4-61.0)	59.9 (57.7-62.2)	89.3 (88.1-90.5)
Nova Scotia	82.9 (80.9-85.0)	57.4 (53.4-61.6)	59.8 (56.1-63.8)	54.6 (49.6-60.1)	58.1 (52.3-64.6)	56.4 (53.3-59.7)	59.5 (56.4-62.8)	87.5 (85.5-89.6)
Ontario	79.3 (77.0-81.8)	54.3 (50.0-58.9)	58.2 (54.3-62.4)	84.7 (81.8-87.6)
Saskatchewan	81.6 (80.9-82.3)	56.0 (54.8-57.3)	58.5 (57.3-59.7)	51.1 (49.3-52.9)	57.8 (55.8-59.8)	54.5 (53.5-55.6)	58.6 (57.5-59.6)	83.4 (82.5-84.3)
US registries	82.8 (80.8-84.8)	55.4 (51.3-59.7)	58.0 (53.9-62.4)	54.8 (49.6-60.6)	61.1 (55.1-67.7)	55.2 (52.0-58.6)	59.1 (55.6-62.7)	77.5 (74.4-80.8)
Atlanta,† GA	83.9 (83.7-84.1)	60.1 (59.6-60.5)	60.1 (59.7-60.5)	56.9 (56.3-57.5)	59.8 (59.2-60.4)	59.1 (58.8-59.5)	60.2 (59.8-60.5)	91.9 (91.7-92.1)
Los Angeles, CA	85.7 (84.0-87.4)	63.9 (60.2-67.7)	60.7 (57.8-63.7)	56.5 (50.9-62.7)	64.3 (59.4-69.7)	62.3 (59.3-65.6)	62.0 (59.4-64.7)	93.4 (91.8-94.9)
San Francisco, CA	84.6 (84.3-85.0)	60.4 (59.5-61.2)	59.5 (58.7-60.3)	57.2 (56.0-58.5)	60.1 (58.8-61.4)	59.4 (58.7-60.1)	59.9 (59.2-60.5)	90.4 (90.0-90.8)
Colorado	83.4 (82.6-84.2)	61.2 (59.6-62.9)	58.4 (56.9-60.0)	55.7 (53.3-58.1)	58.5 (56.1-61.0)	59.5 (58.1-60.8)	58.5 (57.2-59.8)	90.7 (89.9-91.5)
Connecticut	86.2 (85.2-87.2)	59.2 (57.1-61.4)	59.9 (57.9-62.0)	56.5 (53.4-59.8)	60.3 (57.1-63.7)	58.4 (56.6-60.2)	60.2 (58.4-62.0)	89.5 (88.4-90.6)
Florida	87.0 (85.8-88.2)	61.6 (59.0-64.4)	62.0 (59.5-64.6)	55.6 (51.7-59.8)	59.8 (55.9-64.0)	59.7 (57.5-62.0)	61.7 (59.6-63.8)	92.8 (91.6-93.9)
Hawaii	85.7 (84.7-86.7)	62.3 (60.1-64.7)	63.4 (61.3-65.6)	61.3 (58.1-64.6)	62.4 (59.1-65.8)	62.0 (60.2-63.9)	63.4 (61.6-65.2)	91.7 (90.5-93.0)
Idaho	84.0 (83.5-84.5)	60.2 (59.2-61.3)	61.0 (60.0-62.0)	57.1 (55.5-58.7)	61.0 (59.4-62.6)	59.4 (58.5-60.2)	61.2 (60.3-62.1)	89.0 (88.4-89.5)
Iowa	89.3 (87.3-91.4)	67.9 (64.2-71.8)	66.5 (62.6-70.6)	59.3 (54.2-64.8)	61.0 (54.7-68.0)	65.0 (61.9-68.1)	65.5 (62.2-69.0)	90.9 (88.7-93.2)
Louisiana	86.3 (84.2-88.5)	61.4 (56.9-66.3)	63.4 (59.1-68.0)	66.9 (60.8-73.6)	60.0 (53.3-67.6)	63.6 (59.9-67.6)	62.8 (59.2-66.7)	91.7 (89.8-93.7)
Michigan	86.6 (85.8-87.7)	60.8 (58.4-63.3)	64.8 (62.7-67.0)	59.0 (55.6-62.6)	63.8 (60.2-67.6)	60.3 (58.3-62.3)	64.7 (62.9-66.6)	92.6 (91.4-93.8)
Minnesota	81.0 (79.9-82.2)	59.8 (57.5-62.1)	58.8 (56.8-60.7)	57.3 (53.9-60.9)	58.7 (55.5-62.1)	59.1 (57.3-61.1)	58.9 (57.2-60.6)	88.4 (87.2-89.6)
New Jersey	82.3 (81.7-83.0)	58.7 (57.4-60.1)	59.3 (58.0-60.5)	55.2 (53.2-57.2)	59.2 (57.2-61.3)	57.8 (56.7-58.9)	59.4 (58.4-60.5)	100.0 (99.8-100.0)
New Mexico	83.0 (82.0-84.1)	60.5 (58.3-62.8)	58.0 (56.0-60.1)	55.8 (52.6-59.1)	57.5 (54.2-60.9)	59.1 (57.3-61.0)	57.9 (56.2-59.6)	93.4 (92.4-94.4)
New York State	85.4 (84.0-86.9)	60.4 (57.3-63.7)	64.2 (61.4-67.2)	58.3 (56.0-63.0)	60.6 (56.0-65.7)	59.8 (57.3-62.5)	63.6 (61.1-66.1)	92.8 (91.3-94.4)
North Carolina	83.3 (82.6-84.0)	61.3 (59.9-62.7)	61.1 (59.8-62.5)	56.1 (54.0-58.2)	58.4 (56.3-60.5)	59.6 (58.4-60.8)	60.5 (59.4-61.6)	90.8 (90.1-91.6)
Ohio	84.6 (82.7-86.4)	62.0 (58.1-66.2)	61.6 (57.8-65.7)	52.6 (47.2-58.7)	59.1 (53.0-65.8)	59.0 (55.7-62.4)	61.0 (57.8-64.4)	92.4 (90.7-94.1)
Pennsylvania	81.0 (80.5-81.5)	56.6 (55.6-57.7)	56.4 (55.5-57.4)	54.9 (53.4-56.4)	56.7 (55.2-58.2)	56.1 (55.3-57.0)	56.6 (55.8-57.4)	85.6 (85.0-86.2)
Rhode Island	77.4 (76.6-78.2)	54.2 (52.6-55.9)	53.6 (52.1-55.1)	50.6 (48.2-53.2)	52.4 (50.0-54.9)	53.2 (51.8-54.5)	53.3 (52.1-54.6)	81.6 (80.5-82.7)
Texas	84.6 (82.8-86.4)	64.7 (60.9-68.7)	63.5 (60.0-67.2)	60.1 (54.5-66.3)	59.9 (54.5-65.8)	63.3 (60.2-66.7)	62.8 (59.8-65.8)	90.8 (88.4-93.2)
Virginia	88.6 (87.5-89.7)	63.7 (61.3-66.2)	64.1 (61.9-66.5)	60.7 (57.2-64.4)	65.4 (61.9-69.2)	63.0 (60.9-65.1)	64.8 (62.9-66.8)	95.0 (94.0-96.0)
Washington	85.8 (84.0-87.7)	60.8 (56.8-65.1)	58.6 (54.5-63.0)	59.9 (54.2-66.2)	61.3 (55.0-68.2)	61.1 (57.8-64.6)	59.6 (56.2-63.3)	93.7 (92.2-95.2)
Wisconsin	84.3 (80.9-87.8)	59.5 (52.5-67.4)	58.5 (52.2-65.6)	46.5 (37.3-57.9)	52.3 (42.7-64.0)	56.0 (50.1-62.5)	57.8 (52.4-63.7)	92.2 (89.3-95.3)
Asia								
Japanese registries	81.6 (79.7-83.5)	63.0 (61.3-64.8)	57.1 (55.5-58.8)	58.2 (55.9-60.5)	57.6 (55.2-60.1)	61.1 (59.7-62.5)	57.3 (55.9-58.6)	50.4 (46.3-54.9)
Fukui	83.1 (78.3-88.2)	68.5 (64.2-73.0)	62.8 (58.8-67.0)	59.6 (54.1-65.7)	61.6 (56.0-67.8)	65.3 (61.8-68.9)	62.4 (59.1-65.9)	54.1(46.6-61.6)R
Osaka	79.4 (77.1-81.9)	59.6 (57.3-62.0)	52.5 (50.4-54.7)	54.4 (51.3-57.7)	55.2 (51.9-58.7)	57.6 (55.7-59.5)	53.3 (51.5-55.2)	51.1 (46.1-56.6)
Yamagata	87.3 (83.4-91.4)	67.5 (64.3-70.8)	63.7 (60.7-66.8)	63.7 (59.8-67.9)	61.8 (57.6-66.3)	66.0 (63.5-68.5)	63.0 (60.5-65.5)	49.4(43.2-55.6)R
Europe								
European registries	73.1 (72.9-73.4)	46.8 (46.3-47.2)	48.4 (48.0-48.8)	43.2 (42.7-43.7)	47.4 (46.9-48.0)	45.3 (45.0-45.6)	48.1 (47.7-48.4)	57.1 (56.7-57.6)
Austria (Tirol)	74.9 (71.9-78.1)	57.0 (51.5-63.0)	59.3 (54.3-64.7)	45.8 (39.1-53.8)	45.2 (37.6-52.8) R	52.7 (48.2-57.6)	55.1 (50.8-59.7)	86.1 (82.9-89.4)
Czech Republic (West Bohemia)	62.9 (58.9-67.1)	37.7 (33.0-43.0)	37.6 (33.3-42.5)	29.3 (25.2-34.1)	39.1 (33.8-45.2)	33.8 (30.5-37.6)	38.3 (34.9-42.0)	50.7 (44.4-58.0)
Denmark	73.6 (72.5-74.7)	44.7 (42.7-46.7)	48.6 (46.8-50.4)	43.4 (41.2-45.6)	45.9 (43.6-48.3)	44.2 (42.7-45.7)	47.7 (46.3-49.2)	38.4 (36.3-40.6)
Estonia	61.3 (57.9-64.8)	38.5 (33.7-43.1)	39.1 (35.3-43.2)	33.6 (28.4-39.7)	30.2 (26.0-35.1)	36.4 (32.8-40.4)	35.5 (32.6-38.6)	56.5 (52.3-60.9)

(Continues on next page)

	Breast	Colon	Rectum	Colorectum	Prostate			
	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	RS (%) (95% CI)
(Continued from previous page)								
Finland	80.2 (79.0-81.4)	54.6 (51.6-57.8)	54.7 (52.5-57.1)	49.8 (46.8-53.0)	52.6 (49.7-55.6)	52.5 (50.4-54.7)	54.0 (52.2-55.8)	62.9 (60.6-65.2)
French registries	79.8 (78.2-81.4)	57.4 (54.4-60.7)	60.1 (57.2-63.2)	52.8 (49.3-56.7)	63.9 (60.1-67.8)	55.6 (53.3-58.1)	61.5 (59.2-64.0)	73.7 (70.5-77.1)
Bas-Rhin	82.2 (79.7-84.7)	57.8 (53.5-62.5)	62.7 (58.8-66.9)	57.9 (52.6-63.7)	61.7 (56.0-67.9)	57.8 (54.4-61.4)	63.0 (59.6-66.6)	73.8 (69.4-78.4)
Calvados	75.6 (72.5-78.8)	62.0 (56.0-68.5)	61.3 (56.0-67.1)	52.2 (45.6-59.8)	67.9 (62.0-74.5)	57.6 (53.1-62.5)	64.2 (60.1-68.5)	73.1 (68.4-78.2)
Côte d'Or	78.1 (74.1-82.3)	50.6 (44.6-57.5)	52.6 (46.7-59.4)	45.3 (38.8-53.0)	61.3 (53.3-70.5)	48.7 (44.1-53.7)	55.3 (50.5-60.6)	--
Isère	81.9 (78.6-85.2)	--	--	--	--	--	--	--
Germany (Saarland)	75.5 (73.3-77.8)	52.0 (48.2-56.0)	56.2 (52.9-59.7)	47.8 (43.0-53.1)	52.5 (48.1-57.3)	50.1 (47.2-53.2)	55.0 (52.3-57.9)	76.4 (72.7-80.4)
Iceland	79.0 (73.5-85.0)	48.1 (39.0-59.3)	54.9 (45.2-66.6)	52.1 (31.9-71.4) R	48.4 (31.7-64.6) R	49.5 (41.0-59.9)	54.0 (45.9-63.6)	69.7 (62.2-78.1)
Ireland	69.6 (66.1-73.3)	49.1 (44.0-54.8)	48.5 (43.7-53.8)	41.1 (35.0-48.2)	52.5 (44.6-60.3) R	46.0 (42.0-50.4)	50.0 (45.9-54.5)	62.8 (58.0-68.0)
Italian registries	79.5 (78.8-80.3)	52.4 (51.1-53.8)	53.8 (52.6-55.0)	47.4 (45.7-49.2)	50.4 (48.6-52.3)	50.7 (49.7-51.8)	52.7 (51.7-53.8)	65.4 (63.7-67.2)
Ferrara	78.5 (75.6-82.2)	48.5 (43.2-54.5)	54.9 (49.8-60.5)	44.6 (37.1-53.6)	48.0 (40.5-57.0)	47.3 (42.8-52.2)	53.6 (49.2-58.4)	69.8 (63.2-76.0) R
Genoa	80.6 (78.3-83.0)	49.9 (45.9-54.2)	51.2 (47.5-55.3)	40.5 (35.2-46.6)	45.4 (40.0-51.5)	46.8 (43.5-50.3)	49.5 (46.3-52.9)	66.2 (61.0-71.9)
Latina	81.8 (76.4-87.5)	52.7 (45.3-61.3)	57.4 (49.9-65.9)	46.3 (36.3-56.2) R	45.1 (34.7-58.5)	51.2 (45.0-58.2)	53.3 (47.1-60.3)	61.0 (53.9-69.1)
Macerata	77.5 (73.0-82.4)	48.9 (42.8-55.9)	57.9 (51.7-65.0)	42.0 (34.1-51.8)	52.1 (41.2-62.6) R	46.7 (41.6-52.3)	56.8 (51.4-62.7)	69.7 (63.1-76.0) R
Modena	83.1 (80.4-85.8)	55.0 (50.5-59.9)	52.0 (47.7-56.5)	48.4 (42.5-55.1)	45.3 (39.0-52.5)	52.8 (49.2-56.7)	49.8 (46.2-53.7)	68.7 (61.7-76.6)
Parma	81.2 (78.1-84.4)	50.7 (45.6-56.4)	53.7 (48.3-59.7)	47.4 (39.9-54.9) R	41.6 (34.7-49.7)	49.8 (45.6-54.5)	49.3 (44.9-54.2)	56.1 (48.0-65.6)
Ragusa	68.9 (63.2-75.1)	39.5 (37.0-48.8)	44.0 (36.8-52.6)	50.3 (40.8-61.9)	37.8 (26.0-50.3) R	44.9 (38.7-52.1)	41.9 (35.9-48.9)	49.9 (41.0-58.9) R
Romagna	87.4 (84.4-90.4)	51.4 (46.2-57.1)	58.7 (54.0-63.8)	51.0 (42.9-59.0) R	57.9 (50.8-65.9)	50.9 (46.6-55.5)	58.4 (54.4-62.7)	73.3 (67.9-79.2)
Sassari	76.4 (71.3-81.9)	39.9 (31.2-51.0)	41.5 (32.0-51.0) R	44.5 (34.2-54.8) R	42.8 (31.5-58.0)	42.3 (35.9-50.1)	43.5 (36.5-51.8)	52.2 (42.8-61.5) R
Turin	79.4 (77.1-81.7)	50.1 (46.1-54.5)	51.4 (47.8-55.4)	43.7 (39.0-49.0)	54.0 (48.8-59.6)	47.8 (44.7-51.2)	52.4 (49.3-55.6)	63.2 (58.1-68.8)
Tuscany	80.8 (78.9-82.7)	55.6 (52.5-58.9)	54.4 (51.4-57.5)	50.8 (46.9-55.0)	48.7 (44.6-53.2)	53.8 (51.4-56.4)	52.5 (50.1-55.1)	66.4 (62.4-70.7)
Varese	77.6 (75.2-80.0)	55.3 (51.0-59.9)	55.1 (51.1-59.5)	52.4 (46.5-59.0)	53.4 (47.8-59.6)	54.5 (51.1-58.2)	54.5 (51.1-58.1)	72.2 (66.7-78.2)
Veneto	77.5 (76.2-79.1)	53.7 (50.9-56.7)	54.6 (52.0-57.3)	48.4 (44.6-52.5)	55.7 (51.7-60.0)	52.0 (49.8-54.4)	55.0 (52.8-57.2)	61.8 (58.5-65.3)
Malta	73.5 (66.7-81.1)	38.0 (25.9-50.7) R	58.0 (46.5-72.4)	34.7 (20.8-49.9) R	52.5 (31.9-71.4) R	35.7 (27.0-47.1)	55.5 (46.1-66.8)	44.3 (32.3-56.9) R
Netherlands registries	77.6 (76.6-78.6)	52.7 (50.1-55.4)	55.4 (53.2-57.7)	55.0 (51.6-58.6)	54.5 (51.3-57.9)	53.6 (51.5-55.7)	55.1 (53.3-57.0)	69.5 (67.2-71.9)
Amsterdam	78.0 (76.5-79.4)	52.1 (49.1-55.2)	54.1 (51.6-56.7)	51.5 (47.6-55.7)	56.4 (52.7-60.3)	51.9 (49.5-54.3)	54.8 (52.7-57.0)	68.1 (65.4-70.8)
Netherlands (North)	77.8 (76.2-79.4)	--	--	--	--	--	--	--
Netherlands (South)	75.7 (72.9-78.5)	54.2 (49.2-59.8)	59.4 (54.9-64.2)	62.1 (56.6-68.1)	49.2 (43.1-56.1)	58.0 (54.2-62.2)	56.1 (52.5-60.0)	74.9 (70.3-79.8)
Norway	76.3 (75.1-77.6)	50.8 (48.7-53.0)	54.4 (52.5-56.3)	51.3 (48.9-53.9)	56.9 (54.3-59.6)	51.1 (49.5-52.8)	55.3 (53.8-56.9)	63.0 (60.9-65.1)
Polish registries	62.9 (60.6-65.3)	28.5 (25.3-32.1)	30.9 (28.0-34.2)	28.4 (24.7-32.7)	30.2 (26.7-34.1)	28.6 (26.1-31.3)	30.6 (28.3-33.0)	37.1 (33.0-41.6)
Cracow	54.7 (50.6-59.1)	24.6 (18.8-32.1)	23.4 (17.9-30.7)	25.0 (18.9-33.3)	22.9 (16.8-31.1)	25.7 (21.5-30.8)	22.5 (18.3-27.6)	21.3 (15.2-29.9)
Warsaw	66.1 (63.4-68.9)	29.7 (26.1-33.9)	33.6 (30.3-37.4)	29.2 (24.9-34.2)	32.6 (28.6-37.3)	29.6 (26.8-32.7)	33.0 (30.3-35.8)	41.4 (36.5-46.8)
Portugal (South)	72.2 (68.2-76.5)	48.6 (42.6-55.4)	44.8 (39.1-51.3)	42.3 (35.5-50.4)	44.5 (37.8-52.4)	46.5 (41.8-51.8)	44.7 (40.2-49.7)	47.7 (40.7-54.8) R
Slovakia	57.9 (55.9-59.9)	40.1 (37.7-42.7)	44.1 (41.7-46.7)	27.6 (25.5-29.8)	32.3 (29.9-34.8)	34.0 (32.3-35.8)	38.7 (37.0-40.5)	45.7 (42.7-49.0)
Slovenia	66.3 (63.8-68.9)	37.3 (33.5-41.5)	39.8 (36.3-43.6)	34.0 (30.5-38.0)	35.6 (32.1-39.5)	35.7 (33.1-38.5)	37.7 (35.3-40.4)	43.7 (39.4-48.4)
Spanish registries	77.7 (76.4-79.0)	54.2 (52.2-56.3)	56.3 (54.2-58.4)	50.0 (47.7-52.4)	51.8 (49.1-54.6)	52.5 (51.0-54.1)	54.7 (53.1-56.4)	60.5 (57.6-63.6)
Basque Country	79.5 (77.6-81.5)	59.0 (55.8-62.3)	58.3 (55.0-61.8)	53.3 (49.6-57.3)	52.2 (47.8-56.9)	56.5 (54.1-59.0)	56.2 (53.5-58.9)	63.0 (58.8-67.4)
Granada	71.8 (67.0-77.0)	50.6 (44.3-57.8)	50.9 (44.5-58.2)	45.7 (38.1-54.8)	51.1 (43.0-60.8)	48.2 (43.3-53.7)	51.1 (46.0-56.8)	--
Mallorca	80.1 (77.2-83.2)	51.4 (46.4-57.1)	57.4 (52.2-63.0)	48.9 (42.5-56.2)	51.7 (44.5-59.9)	50.9 (46.9-55.3)	56.1 (51.8-60.7)	68.2 (60.7-76.6)
Murcia	72.8 (69.1-76.8)	49.7 (44.4-55.7)	54.8 (50.2-59.9)	49.2 (43.4-55.8)	47.8 (42.0-54.4)	49.7 (45.5-54.3)	52.3 (48.7-56.3)	52.0 (45.4-59.4)
Navarra	78.3 (74.9-81.8)	50.6 (45.1-56.8)	53.3 (46.8-60.8)	42.7 (36.4-50.1)	58.1 (49.1-66.5) R	47.7 (43.4-52.4)	55.6 (50.4-61.3)	54.6 (47.2-63.0)
Tarragona	76.4 (73.0-80.0)	49.2 (43.9-55.1)	52.8 (47.8-58.3)	50.1 (43.2-58.3)	49.8 (40.9-58.4) R	49.6 (45.4-54.3)	53.7 (47.4-56.4)	54.6 (46.3-64.3)
Sweden	82.0 (81.2-82.7)	52.5 (50.9-54.2)	54.8 (53.3-56.3)	53.0 (51.2-55.0)	58.2 (56.3-60.2)	52.8 (51.6-54.1)	56.2 (55.0-57.4)	66.0 (64.7-67.3)
Swiss registries	76.0 (74.3-77.7)	--	--	--	--	--	--	--
Basel	78.2 (75.1-81.4)	--	--	--	--	--	--	--
Geneva	79.1 (76.0-82.4)	--	--	--	--	--	--	--

(Continues on next page)

very few registries was it greater than 1% (available online¹⁴). Such patients are described as censored from the analysis.

Statistical analysis

We estimated relative survival up to 5 years after diagnosis from the individual tumour data, using the Hakulinen approach¹⁵ embedded in the US National Cancer Institute's publicly accessible SEER*Stat software.¹⁶ SEER*Stat is the standard tool used for cancer-survival estimation by the SEER Program cancer registries, and we used it to ensure that survival estimates for US registries would be seen as comparable with those already published by the SEER Program. Survival estimates were also derived by race for the USA (black and white).

Relative survival is the ratio of the survival noted in the patients with cancer and the survival that would have been

expected had they been subject only to the mortality rates of the general population (background mortality). It is a measure of the excess mortality in patients with cancer over and above the background mortality, and can be interpreted as survival from the cancer after correction for other causes of death. This approach is crucial for international comparisons of cancer survival, because the background risks of death from all causes in adults often differ very widely. Background mortality was taken from life tables developed specially for the CONCORD study, specific for sex, calendar year, region, and race.¹⁶

The probability of survival in successive years after diagnosis was estimated in survivors to the start of each year. We report the cumulative relative survival at 5 years. Survival was not estimated if fewer than five patients with a given cancer were available for analysis in any category defined by age, sex, and race. Relative survival was adjusted

	Breast	Colon	Rectum	Colorectum	Prostate			
	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	RS (%) (95% CI)
(Continued from previous page)								
Graubünden-Glarus	71.7 (66.8-77.0)	--	--	--	--	--	--	--
St Gallen-Appenzell	71.7 (68.1-75.5)	--	--	--	--	--	--	--
Valais	75.3 (70.4-80.6)	--	--	--	--	--	--	--
UK	69.7 (69.4-70.1)	43.5 (42.9-44.1)	44.4 (43.8-45.0)	40.6 (39.9-41.3)	45.3 (44.5-46.1)	42.3 (41.8-42.8)	44.7 (44.3-45.2)	51.1 (50.4-51.8)
England (national)	69.8 (69.5-70.2)	43.4 (42.8-44.1)	44.3 (43.7-45.0)	40.4 (39.6-41.2)	45.4 (44.6-46.3)	42.2 (41.7-42.7)	44.7 (44.2-45.3)	50.9 (50.1-51.7)
East Anglia	70.8 (69.2-72.4)	43.6 (40.8-46.7)	42.9 (40.2-45.8)	46.0 (42.4-49.8)	49.8 (46.1-53.9)	44.6 (42.4-47.0)	45.2 (43.0-47.6)	51.9 (48.4-55.7)
Mersey	69.4 (67.8-71.1)	43.8 (41.0-46.9)	43.6 (41.0-46.4)	41.2 (38.1-44.5)	44.5 (41.0-48.2)	43.0 (40.9-45.1)	44.0 (41.8-46.2)	52.6 (49.3-56.1)
Oxford	71.1 (69.6-72.6)	44.8 (42.1-47.8)	45.0 (42.4-47.8)	43.1 (39.8-46.6)	45.6 (41.8-49.7)	44.3 (42.1-46.6)	45.3 (43.2-47.6)	50.4 (47.4-53.6)
South Thames	73.9 (73.0-74.9)	45.5 (43.6-47.6)	48.3 (46.5-50.2)	45.3 (43.0-47.8)	51.1 (48.6-53.6)	45.5 (44.0-47.1)	49.3 (47.9-50.8)	56.1 (54.0-58.2)
South West	73.4 (72.5-74.2)	51.5 (49.8-53.1)	51.6 (50.1-53.2)	48.6 (46.7-50.6)	52.0 (49.8-54.2)	50.3 (49.0-51.5)	51.8 (50.5-53.1)	55.8 (53.9-57.9)
Trent	68.2 (67.2-69.3)	40.3 (38.3-42.5)	42.2 (40.2-44.2)	39.3 (37.1-41.6)	43.8 (41.3-46.5)	39.8 (38.3-41.4)	42.9 (41.3-44.5)	47.0 (44.8-49.4)
West Midlands	75.4 (74.2-76.5)	48.0 (46.2-49.9)	48.4 (46.6-50.2)	44.4 (42.2-46.7)	46.9 (44.3-49.6)	46.6 (45.2-48.1)	48.0 (46.5-49.5)	55.4 (53.2-57.7)
Yorkshire	71.4 (70.1-72.8)	45.5 (43.1-48.1)	45.4 (43.1-47.8)	43.8 (41.1-46.7)	49.8 (46.8-53.0)	44.7 (42.9-46.6)	47.0 (45.1-48.9)	53.3 (50.5-56.4)
Northern Ireland	72.0 (68.9-75.3)	47.3 (42.1-53.0)	49.0 (44.3-54.3)	48.2 (41.6-55.8)	43.8 (37.0-51.9)	47.8 (43.7-52.3)	47.8 (43.8-52.2)	54.0 (48.7-59.9)
Scotland	70.6 (69.5-71.8)	45.9 (44.0-47.9)	47.8 (46.1-49.6)	42.3 (39.9-44.9)	46.9 (44.4-49.6)	44.6 (43.1-46.2)	47.7 (46.2-49.2)	54.2 (52.0-56.5)
Wales	67.1 (65.8-68.4)	39.9 (37.5-42.6)	38.0 (35.7-40.4)	39.5 (36.8-42.3)	41.9 (38.8-45.2)	39.8 (38.0-41.8)	39.3 (37.5-41.3)	47.9 (44.9-51.1)
Oceania								
Australia (national)	80.7 (80.1-81.3)	57.8 (56.8-58.8)	57.7 (56.7-58.6)	54.8 (53.6-56.1)	59.2 (57.8-60.6)	56.7 (55.9-57.5)	58.2 (57.4-58.9)	77.4 (76.6-78.2)
Australian Capital Territory	80.4 (74.3-87.0)	62.0 (53.8-71.5)	59.1 (51.2-68.2)	57.2 (45.5-68.1) R	61.3 (49.8-75.5)	56.5 (49.1-65.1)	59.8 (53.0-67.5)	78.7 (72.5-85.5)
New South Wales	80.4 (79.4-81.5)	60.8 (59.1-62.6)	58.2 (56.6-59.9)	56.9 (54.7-59.1)	59.6 (57.3-61.9)	59.3 (57.9-60.7)	58.7 (57.4-60.0)	78.3 (77.0-79.6)
Northern Territory	71.9 (58.7-88.0)	53.5 (36.3-69.4) R	51.7 (34.2-67.5) R	46.3 (28.9-63.4) R	66.5 (39.6-86.0) R	52.1 (38.6-70.5)	53.2 (39.9-70.9)	63.7 (49.0-77.0) R
Queensland	80.5 (79.0-82.0)	59.8 (57.5-62.3)	60.6 (58.6-62.8)	53.7 (50.7-56.9)	61.2 (57.7-64.8)	57.7 (55.8-59.6)	60.7 (58.9-62.5)	75.7 (73.9-77.6)
South Australia	80.0 (78.0-82.0)	56.3 (53.0-59.8)	58.6 (55.5-61.8)	55.2 (51.3-59.4)	59.2 (55.1-63.6)	55.8 (53.3-58.4)	58.6 (56.1-61.2)	77.1 (74.3-80.1)
Tasmania	77.1 (73.4-81.1)	52.4 (46.8-58.6)	50.0 (44.9-55.6)	44.9 (37.5-53.6)	55.0 (46.8-64.6)	50.2 (45.7-55.1)	51.8 (47.4-56.6)	70.2 (65.8-74.8)
Victoria	81.5 (80.4-82.7)	54.7 (52.7-56.7)	56.1 (54.3-57.9)	54.9 (52.5-57.4)	59.0 (56.5-61.6)	54.8 (53.3-56.4)	57.2 (55.7-58.6)	76.8 (75.2-78.4)
Western Australia	81.4 (79.3-83.5)	53.2 (49.7-56.9)	54.5 (51.4-57.8)	50.9 (46.8-55.3)	54.8 (50.3-59.7)	52.5 (49.8-55.3)	54.8 (52.1-57.5)	80.0 (77.7-82.3)

RS=relative survival. R=raw (not age-standardised) survival estimate; too few cases in one or more age groups. *International Cancer Survival Standard (see text). †No state-wide data available for this city. ‡Survival truncated if greater than 1.0 (100%). 95% CIs were calculated by use of a logarithmic transformation (see text).

Table 2: 5-year relative survival (%), age-standardised to ICSS weights* with 95% CIs for adults (aged 15-99 years) diagnosed with cancer of the breast (women), colon, rectum, or prostate during 1990-94 and followed up to Dec 31, 1999: continent, country, and region

for heterogeneity in the withdrawal of patients from follow-up and consequent changes in the age-sex-race distribution of patients with cancer in successive calendar years, by use of the exact method.⁴⁴

Expected survival was derived from complete life tables that contained the probabilities of death or the central death rates for the general population of the registry's territory, by single year of age, sex and (where possible) race, and single calendar year between 1990 and 1999. Many registries provided complete life tables. For some registries, complete life tables were constructed from raw data obtained from published sources on the numbers of deaths by age, sex, and race in the relevant year(s) or period, and the corresponding populations. For the remaining registries, abridged (5-year or 10-year age groups) life tables from published sources were smoothed to produce complete life tables. In some registries, life tables were interpolated, as required, to provide life tables by single calendar year throughout the decade 1990–99. Details are provided in an accompanying paper.⁴⁵

Cancer survival is known to vary with race,^{46–48} and we assessed racial differences in survival where possible. Individual tumour records were coded by race only in the data from the USA (black, white, other). Race-specific estimates of relative survival were produced with separate life tables for each race, constructed from the raw data on populations and the number of deaths.⁴⁶

In the USA, race-specific mortality in the general population also varies between states.⁴⁹ We developed separate sets of complete life tables for each state and metropolitan area and for each sex. This approach was designed to enable the closest possible adjustment of relative survival estimates in the USA for geographic variation in background mortality in both blacks and whites, by age, sex, and calendar period. Race-specific life tables for both blacks and whites were developed for 11 of the 16 states and all six metropolitan areas. Where race-specific life tables were available, they were used in the estimation of relative survival for patients of that race. For other patients, the all-races life table for that population was used. For five less populous states (Hawaii, Idaho, New Mexico, Utah, and Wyoming: 6% of the 109 million population covered by participating registries; webtable), only the life tables for whites were sufficiently robust, and relative survival estimates for blacks are not separately presented.

Relative survival measures the extent to which patients with cancer have a higher death rate than the general population of the country or region in which they live.⁵⁰ Occasionally, despite use of the most appropriate life table, this excess death rate can be negative in a given time interval since diagnosis, implying that the death rate of cancer survivors during that interval is actually lower than that of the general population. This situation can arise from random variation in the death rate when the number of deaths in the interval is small,⁵¹ either because the

interval is very short, or because survival is poor and most patients have already died before the start of the interval, or because survival is high and there are very few deaths. In such situations, we present by default the estimate derived by use of the SEER*Stat option to constrain the excess mortality rate to zero, which imposes a plateau in the relative survival curve. The unconstrained estimate was also obtained for comparison.

Even though relative survival is already adjusted for age-specific differences in background mortality, robust international comparison of relative survival requires age-standardisation,²³ because the age distribution of patients with cancer varies between countries, and because relative survival also varies widely by age, at least in Europe.²⁷ Conventional age-specific weights used to standardise incidence or mortality rates (eg, the national population or the hypothetical world standard population²⁸) are unsuitable because patients with cancer have a very different age profile from that of the general population.

A cancer-survival comparison of such wide scope has not been done before and the choice of weights for age-standardisation was not straightforward. International standard cancer-patient populations have been proposed, with different sets of weights in 5-year or 10-year age bands for each of 20 common cancers, derived from their worldwide distribution.²⁹ The weights used for the EURO-CARE-3 study were derived from the age distribution of all patients included in that study for each cancer, and were thus cancer-specific.⁴¹ The disadvantage of these standards is either that a unique set of weights is required for each cancer (cancer-specific), or else that the standards are arbitrary (study-specific), vitiating comparison between studies.

We chose the recently developed International Cancer Survival Standard (ICSS) weights.³⁰ These comprise just three sets of age weights, derived from discriminant analysis to find the smallest number of sets of standard age weights that enable adequate standardisation of survival. Each standard is applicable to a range of different cancers, and provides age-standardised survival estimates that are not too different from the unstandardised estimates. The first ICSS standard applies to cancers for which incidence rises rapidly with age, and we used this in all analyses. For cancers of the breast, colon, and rectum, we used five age groups: 15–44, 45–54, 55–64, 65–74, and 75–99 years. For prostate cancer, which occurs mainly in older men, we used four age groups: 15–54, 55–64, 65–74, and 75–99 years. Where data were too sparse for standardisation, the raw (unstandardised) survival estimate is presented, flagged with "R".

The same age weights were used for men and women, and for each race, enabling direct comparison of age-standardised relative survival between patient groups defined by sex and race. Because identical weights were used for breast, colon, and rectal cancer, the age-standardised estimates of survival for these cancers can also be directly compared. This would not be possible if cancer-specific weights were used.

See Online for webtable





Figure 1: 5-year relative survival (%), age-standardised to the ICSS weights* with 95% CIs for adults (aged 15–99 years) diagnosed with cancer of the breast (women), colorectum, or prostate during 1990–94 and followed up to Dec 31, 1999; country. Vertical bar on the right of each graphic shows the contribution (%) of each continent to the total number of cases analysed (contributions under 1% are not labelled). Red vertical line represents mean survival for the 22 European countries that participated in EUROCARE-3, age-standardised to ICSS weights. Switzerland only provided data for breast cancer. *Age-standardised to ICSS weights, except for Sétif, Algeria (all cancers), Malta (prostate), and Portugal (prostate), which were unstandardised values (see text). †Problems with data quality might have led to over-estimation (see text).

Figure 2: 5-year relative survival (%), using state-specific and race-specific life tables and age-standardised to the ICSS weights* for adults (aged 15–99 years) diagnosed with cancer of the breast (women), colon, rectum, and rectum combined, or prostate during 1990–94 and followed up to Dec 31, 1999: 16 US States and six metropolitan areas. Vertical lines represent mean survival for SEER (red) and NPCR (green) registries, age-standardised to ICSS weights (see text). *Age-standardised to ICSS weights (see text). †Problems with data quality might have led to over-estimation (see text).



For countries represented by more than one regional cancer registry, we provide a survival estimate derived from the pooled data for all contributing registries, age-standardised in the same way. This is an overall estimate of survival in the combined territories providing data from

that country, not a weighted mean of the various regional estimates. The combined estimate should not be considered as necessarily representative of survival in the country as a whole, except where the regional registries cover the entire country.

	Breast	Colon	Rectum		Colorectum		Prostate	
	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI) See Online for webpanel	RS (%) (95% CI)
Atlanta, GA (all races), S	85.7 (84.0-87.4)	64.7 (60.5-68.0)	60.9 (58.0-63.9)	56.6 (51.0-62.8)	64.5 (59.5-69.8)	62.5 (59.4-65.8)	62.2 (59.6-64.9)	94.0 (92.4-95.6)
Black	71.1 (67.1-75.4)	59.5 (52.3-68.5)	52.6 (47.2-58.6)	45.5 (35.3-58.6)	52.1 (42.5-63.7)	56.8 (50.3-64.2)	52.9 (48.1-58.2)	86.5 (83.3-89.8)
White	89.6 (87.8-91.5)	65.4 (61.3-69.8)	63.9 (60.5-67.6)	59.4 (52.9-66.7)	67.9 (62.3-74.1)	64.1 (60.6-67.8)	65.4 (62.4-68.4)	96.1 (94.4-97.9)
California (all races), N	84.9 (84.5-85.3)	60.8 (59.9-61.6)	59.8 (59.0-60.6)	57.5 (56.3-58.8)	60.3 (59.0-61.6)	59.8 (59.1-60.5)	60.1 (59.4-60.8)	91.1 (90.6-91.5)
Black	73.4 (71.4-75.6)	54.8 (51.4-58.4)	51.1 (48.3-54.2)	50.3 (44.9-56.4)	50.9 (45.9-56.4)	53.6 (50.7-56.7)	51.2 (48.7-53.8)	84.5 (82.9-86.1)
White	85.3 (84.9-85.7)	60.7 (59.8-61.6)	60.1 (59.2-61.0)	57.4 (56.1-58.7)	60.4 (59.0-61.8)	59.7 (58.9-60.4)	60.3 (59.6-61.0)	90.8 (90.3-91.2)
Los Angeles, CA (all races), N	83.8 (83.0-84.6)	61.9 (60.2-63.6)	58.8 (57.3-60.3)	56.2 (53.8-58.6)	58.8 (56.4-61.3)	60.0 (58.7-61.4)	58.8 (57.6-60.2)	91.7 (90.9-92.6)
Black	72.5 (69.6-75.6)	58.9 (54.5-63.8)	52.1 (48.2-56.3)	49.8 (42.3-58.7)	50.1 (43.4-57.9)	57.0 (53.2-61.2)	51.7 (48.2-55.3)	84.8 (82.5-87.3)
White	84.7 (83.9-85.5)	61.5 (59.7-63.4)	59.4 (57.6-61.2)	55.4 (52.8-58.2)	58.5 (55.7-61.4)	59.6 (58.0-61.1)	59.2 (57.7-60.7)	92.3 (91.4-93.2)
San Francisco, CA (all races), S	86.6 (85.6-87.6)	59.8 (57.6-62.0)	60.3 (58.2-62.5)	57.0 (53.8-60.3)	60.6 (57.4-64.0)	58.9 (57.2-60.6)	60.5 (58.8-62.3)	90.5 (89.4-91.6)
Black	77.2 (73.2-81.4)	47.4 (41.0-54.7)	50.0 (44.4-56.4)	54.7 (43.7-68.5)	52.3 (41.5-66.0)	49.7 (44.2-56.0)	50.6 (45.6-56.2)	83.7 (80.4-87.1)
White	87.5 (86.5-88.6)	60.3 (57.9-62.9)	61.1 (58.7-63.6)	56.8 (53.2-60.5)	61.5 (57.9-65.4)	59.3 (57.3-61.4)	61.4 (59.4-63.5)	90.2 (88.9-91.4)
Colorado (all races), N	87.0 (85.8-88.2)	61.7 (59.2-64.5)	62.0 (59.6-64.6)	55.6 (51.7-59.9)	59.8 (55.9-64.0)	59.8 (57.6-62.1)	61.7 (59.6-63.9)	92.9 (91.8-94.1)
Black	81.6 (74.1-89.9)	45.0 (34.3-58.8)	48.0 (36.9-62.5)	76.8 (44.8-97.7) R	39.6 (16.2-64.9) R	49.7 (39.3-63.0)	46.7 (36.2-60.2)	80.7 (74.6-87.4)
White	87.0 (85.8-88.2)	62.1 (59.4-65.0)	62.3 (59.8-65.0)	54.9 (50.9-59.2)	60.6 (56.6-64.9)	59.8 (57.6-62.2)	62.1 (60.0-64.4)	92.8 (91.6-94.0)
Connecticut (all races), S	85.7 (84.7-86.8)	62.4 (60.2-64.7)	63.5 (61.4-65.7)	61.3 (58.1-64.6)	62.4 (59.1-65.9)	62.1 (60.3-64.0)	63.4 (61.6-65.2)	91.9 (90.7-93.2)
Black	75.2 (69.3-81.6)	51.1 (41.9-62.3)	52.7 (44.8-61.9)	63.5 (47.5-85.0)	73.3 (56.8-86.2) R	54.4 (46.0-64.3)	56.5 (49.4-64.6)	82.3 (77.6-87.2)
White	86.3 (85.3-87.3)	62.9 (60.6-65.3)	64.1 (61.9-66.4)	61.3 (58.1-64.7)	61.9 (58.5-65.5)	62.4 (60.5-64.3)	63.7 (61.9-65.6)	92.3 (91.0-93.6)
Florida (all races), N	84.0 (83.5-84.5)	60.2 (59.2-61.2)	61.0 (60.0-62.1)	57.0 (55.5-58.6)	61.0 (59.4-62.7)	59.4 (58.5-60.2)	61.2 (60.4-62.1)	89.2 (88.7-89.8)
Black	72.7 (70.1-75.3)	54.4 (50.0-59.1)	54.3 (50.9-57.9)	44.8 (37.7-53.1)	54.5 (48.4-61.3)	51.6 (47.8-55.6)	54.8 (51.9-58.0)	84.7 (82.7-86.7)
White	84.7 (84.2-85.2)	60.5 (59.4-61.6)	61.6 (60.5-62.7)	57.6 (56.2-59.5)	61.3 (59.6-63.1)	59.8 (59.0-60.8)	61.7 (60.8-62.6)	89.7 (89.1-90.3)
Hawaii (all races), S	90.2 (88.1-92.3)	68.4 (64.7-72.3)	67.2 (63.3-71.3)	59.6 (54.5-65.2)	61.5 (55.1-68.6)	65.4 (62.4-68.6)	66.2 (62.8-69.7)	91.8 (89.6-94.1)
White	90.2 (86.5-94.1)	67.9 (61.2-75.2)	61.6 (54.1-70.1)	54.0 (44.3-65.8)	66.0 (50.8-79.0) R	64.6 (58.6-71.1)	62.9 (56.2-70.3)	92.4 (89.0-96.0)
Idaho (all races), N	86.3 (84.2-88.5)	61.4 (56.9-66.3)	63.4 (59.1-68.0)	66.9 (60.8-73.6)	60.0 (53.3-67.6)	63.6 (59.9-67.6)	62.8 (59.1-66.7)	91.7 (89.8-93.7)
White	86.3 (84.2-88.5)	61.4 (56.8-66.4)	63.4 (59.1-68.1)	66.7 (60.5-73.4)	59.9 (53.1-67.5)	63.6 (59.8-67.5)	62.8 (59.1-66.8)	91.5 (89.5-93.5)
Iowa (all races), S	86.6 (85.5-87.7)	60.8 (58.4-63.3)	64.8 (62.7-67.0)	59.0 (55.6-62.6)	63.8 (60.2-67.6)	60.3 (58.3-62.3)	64.7 (62.9-66.6)	92.7 (91.5-93.9)
Black	60.1 (46.6-77.5)	66.8 (39.0-89.6) R	75.2 (51.7-94.1) R	56.5 (17.3-91.4) R	40.7 (12.5-71.8) R	66.9 (43.7-86.2) R	65.9 (46.5-82.8) R	85.8 (72.3-97.6) R
White	86.8 (85.7-87.8)	60.8 (58.4-63.3)	64.6 (62.5-66.8)	58.7 (55.3-62.4)	63.8 (60.2-67.7)	60.2 (58.2-62.2)	64.6 (62.7-66.5)	92.6 (91.4-93.8)
Louisiana (all races), N	81.0 (79.8-82.1)	59.9 (57.6-62.2)	58.8 (56.9-60.8)	57.2 (53.8-60.9)	58.7 (55.5-62.1)	59.2 (57.3-61.1)	58.9 (57.2-60.6)	88.6 (87.4-89.9)
Black	69.9 (67.2-72.7)	54.2 (49.6-59.3)	53.1 (49.6-56.9)	48.0 (40.8-56.4)	48.2 (41.9-55.4)	53.1 (49.2-57.7)	52.4 (49.2-55.8)	80.6 (78.1-83.2)
White	84.0 (82.8-85.3)	61.6 (59.2-64.3)	60.6 (58.4-63.0)	58.4 (54.6-62.4)	61.4 (57.8-65.3)	60.7 (58.6-62.9)	61.1 (59.2-63.1)	91.0 (89.6-92.4)
Michigan (all races)†, N	82.3 (81.6-82.9)	58.8 (57.5-60.2)	59.3 (58.1-60.6)	55.2 (53.2-57.2)	59.2 (57.2-61.3)	57.8 (56.7-59.0)	59.5 (58.4-60.6)	100 (99.8-100)
Black†	69.6 (67.2-72.1)	47.9 (44.2-51.9)	51.8 (48.5-55.4)	46.1 (39.1-51.9)	45.1 (39.3-51.8)	47.1 (43.9-50.6)	50.5 (47.6-53.6)	100 (99.3-100)
White†	83.3 (82.6-84.0)	59.7 (58.3-61.2)	60.2 (58.9-61.6)	55.9 (53.8-58.1)	60.2 (58.1-62.4)	58.7 (57.5-59.9)	60.4 (59.3-61.6)	100 (99.8-100)
Detroit, MI (all races), S	83.0 (81.9-84.0)	60.6 (58.4-62.9)	58.2 (56.2-60.3)	55.7 (52.5-59.0)	57.4 (54.2-60.9)	59.2 (57.3-61.0)	58.0 (56.3-59.8)	93.8 (92.8-94.8)
Black	71.7 (68.9-74.6)	50.6 (45.9-55.8)	51.3 (47.6-55.4)	48.4 (40.9-57.2)	44.5 (37.4-53.0)	49.8 (45.7-54.2)	50.5 (47.1-54.3)	88.7 (86.4-91.1)
White	85.4 (84.3-86.5)	62.7 (60.2-65.3)	60.7 (58.3-63.2)	57.4 (53.9-61.0)	59.6 (55.9-63.4)	61.1 (59.1-63.2)	60.3 (58.3-62.4)	95.3 (94.2-96.4)
Nebraska (all races), N	85.4 (84.0-86.8)	60.4 (57.3-63.7)	64.3 (61.4-67.2)	58.3 (53.9-63.0)	60.6 (55.9-65.7)	58.8 (57.3-62.5)	63.6 (61.1-66.1)	92.9 (91.3-94.4)
Black	83.1 (72.7-94.9)	69.6 (46.5-88.2) R	48.2 (29.9-66.4) R	60.0 (24.9-90.5) R	77.4 (22.6-100) R	66.9 (47.5-83.5) R	52.6 (34.9-69.7) R	78.7 (68.4-90.6)
White	85.4 (83.9-86.8)	59.9 (56.7-63.2)	64.9 (62.1-67.9)	57.8 (53.4-62.6)	60.5 (55.7-65.7)	58.3 (56.7-62.0)	64.0 (61.5-66.6)	93.1 (91.6-94.7)
New Jersey (all races), N	82.4 (82.7-84.1)	61.5 (60.1-62.9)	61.2 (59.9-62.6)	56.1 (54.1-58.3)	58.4 (56.4-60.6)	59.7 (58.6-60.9)	60.6 (59.5-61.7)	91.2 (90.4-91.9)
Black	73.1 (70.2-76.1)	51.6 (46.4-57.4)	51.5 (47.7-55.6)	46.4 (38.3-56.2)	45.1 (38.9-53.0)	50.3 (45.8-55.2)	50.3 (46.9-53.9)	81.0 (78.5-83.5)
White	83.8 (83.1-84.6)	61.4 (60.0-62.9)	61.8 (60.4-63.2)	56.0 (53.9-58.3)	58.9 (56.7-61.1)	59.6 (58.4-60.9)	61.1 (59.9-62.3)	90.8 (90.0-91.7)

(Continues on next page)

The proportion of survivors is constrained in the range zero to one (or 0 to 100%), but confidence intervals (CIs) for relative survival derived in the usual way, from the Normal approximation, can produce implausible values (<0 or >1). SEER*Stat provided the standard error of the cumulative relative survival based on the Greenwood formula,⁴¹ but did not provide CIs. We used these standard errors to estimate CIs on the logarithmic scale (webpanel).

For the USA, we constructed funnel plots of relative survival for each cancer and sex, to obtain further insight into the variability of survival by race and state, and to avoid spurious ranking of the survival estimates.⁴² The plots show how much a particular survival estimate deviates from the pooled US value, given the precision of

each estimate. The precision depends on the number of deaths included in the analysis, which depends in turn on the size of the population and the frequency and lethality of the cancer in that population. 5-year relative survival estimates for each population, age-standardised and adjusted for race-specific and state-specific background mortality, were plotted against the precision of the estimates, taken as the inverse square of their standard errors (webpanel). The horizontal line in each plot, the target, was estimated as the pooled 5-year relative survival for all participating US populations, age-standardised to the same weights. Raw survival estimates were not plotted. The 99.8% control limits superimposed on each plot represent about three standard deviations from the pooled US survival at each level of precision.

	Breast	Colon	Rectum	Colorectum	Prostate			
	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	Men RS (%) (95% CI)	Women RS (%) (95% CI)	RS (%) (95% CI)		
(Continued from previous page)								
New Mexico (all races), S	84.6 (82.8-86.4)	62.1 (58.1-66.3)	61.7 (57.9-65.7)	52.6 (47.2-58.7)	59.1 (53.0-65.8)	59.0 (55.8-62.4)	61.0 (57.8-64.4)	92.4 (90.7-94.2)
White	84.7 (82.8-86.6)	61.7 (57.7-66.0)	61.4 (57.5-65.6)	52.7 (47.1-58.9)	59.0 (52.8-65.9)	58.9 (55.6-62.4)	60.9 (57.6-64.4)	92.7 (91.0-94.4)
New York State (all races), N	83.0 (80.5-85.5)	56.8 (55.8-57.8)	56.5 (55.6-57.5)	55.0 (53.5-56.5)	56.7 (55.2-58.3)	56.3 (55.4-57.1)	56.7 (55.9-57.5)	85.9 (85.3-86.5)
Black	67.2 (65.4-69.1)	45.9 (42.8-49.2)	46.0 (43.5-48.5)	42.8 (37.6-48.7)	46.8 (42.2-51.9)	45.1 (42.4-47.9)	46.2 (44.0-48.5)	75.9 (74.2-77.7)
White	82.1 (81.5-82.6)	57.3 (56.2-58.4)	57.2 (56.2-58.2)	55.8 (54.2-57.4)	57.4 (55.8-59.1)	56.9 (56.0-57.8)	57.4 (56.5-58.2)	86.5 (85.8-87.2)
New York City, NY (all races), N	77.6 (76.8-78.4)	54.5 (52.9-56.2)	53.8 (52.3-55.3)	50.9 (48.4-53.5)	52.6 (50.2-55.1)	53.5 (52.1-54.9)	53.5 (52.3-54.8)	82.3 (81.2-83.4)
Black	65.8 (63.7-67.9)	45.2 (41.7-49.1)	45.0 (42.2-48.0)	44.4 (38.3-51.4)	46.5 (41.3-52.3)	45.0 (41.9-48.3)	45.3 (42.8-48.0)	74.0 (71.9-76.1)
White	79.6 (78.7-80.6)	55.6 (53.7-57.6)	54.9 (53.1-56.7)	51.9 (49.1-54.9)	53.3 (50.5-56.3)	54.5 (52.9-56.2)	54.5 (53.0-56.1)	83.3 (81.8-84.7)
Rhode Island (all races), N	84.6 (82.7-86.4)	64.7 (60.9-68.7)	63.4 (60.0-67.1)	60.1 (54.5-66.3)	59.9 (54.5-65.7)	63.4 (60.2-66.7)	62.7 (59.8-65.8)	90.9 (88.5-93.3)
Black±	82.9 (65.8-100)	58.6 (28.5-85.9) R	45.0 (26.5-71.8) R	NA	79.6 (28.8-100) R	65.5 (35.6-90.7) R	57.5 (31.4-78.5) R	75.5 (59.5-89.0) R
White	84.9 (83.1-86.8)	64.9 (61.1-69.0)	63.7 (60.2-67.4)	60.2 (54.5-66.4)	59.3 (53.9-65.3)	63.6 (60.4-66.9)	62.8 (59.8-65.9)	91.4 (89.0-93.9)
Seattle, WA (all races), S	88.7 (87.6-89.8)	63.9 (61.5-66.4)	64.2 (61.9-66.6)	60.8 (57.4-64.5)	65.5 (61.9-69.3)	63.2 (61.1-65.3)	64.9 (62.9-66.9)	95.3 (94.3-96.4)
Black	64.7 (55.5-75.3)	54.9 (42.5-71.0)	63.9 (45.4-80.2) R	46.9 (26.6-67.1) R	48.7 (20.1-75.1) R	51.9 (41.0-65.6)	54.9 (42.1-71.8)	89.6 (84.0-95.4)
White	89.3 (88.2-90.4)	64.4 (61.9-67.0)	64.1 (61.7-66.5)	61.7 (58.1-65.6)	65.7 (61.9-69.6)	63.8 (61.7-66.0)	64.8 (62.8-66.9)	95.4 (94.3-96.4)
Utah (all races), S	85.8 (84.0-87.7)	60.8 (56.8-65.1)	58.6 (54.5-63.0)	59.9 (54.2-66.2)	61.3 (55.0-68.2)	61.1 (57.8-64.6)	59.6 (56.2-63.3)	93.7 (92.1-95.2)
White	85.9 (84.0-87.9)	60.7 (56.6-65.1)	58.7 (54.6-63.2)	59.7 (53.8-66.2)	62.6 (56.3-69.8)	61.0 (57.6-64.6)	60.2 (56.6-63.9)	93.5 (91.9-95.1)
Wyoming (all races), N	84.2 (80.8-87.7)	59.5 (52.5-67.4)	58.5 (52.1-65.6)	46.4 (37.2-57.9)	52.2 (42.7-63.9)	55.9 (50.1-62.4)	57.7 (52.2-63.8)	92.2 (89.2-95.2)
White	84.3 (80.9-87.8)	60.5 (53.5-68.4)	58.1 (51.7-65.3)	46.2 (37.0-57.9)	52.5 (42.7-64.4)	56.5 (50.6-63.0)	57.5 (52.0-63.7)	92.1 (89.2-95.1)
NPCR (all races)	83.1 (82.8-83.4)	59.8 (59.3-60.4)	59.6 (59.1-60.1)	56.3 (55.5-57.1)	58.8 (58.0-59.7)	58.8 (58.3-59.2)	59.6 (59.1-60.0)	89.5 (89.2-89.8)
Black	70.7 (69.6-71.8)	52.1 (50.2-54.1)	50.5 (49.0-52.0)	46.9 (43.7-50.2)	49.2 (46.4-51.9)	50.7 (49.1-52.5)	50.3 (49.0-51.6)	81.1 (80.2-82.1)
White	84.0 (83.7-84.3)	60.1 (59.6-60.7)	60.4 (59.8-60.9)	56.7 (55.8-57.5)	59.4 (58.5-60.3)	59.1 (58.6-59.6)	60.2 (59.8-60.7)	90.0 (89.7-90.3)
SEER (all races)	86.1 (85.6-86.5)	61.9 (61.0-62.8)	62.1 (61.2-62.9)	58.5 (57.1-59.8)	61.8 (60.4-63.2)	60.9 (60.2-61.7)	62.2 (61.5-62.9)	93.1 (92.7-93.5)
Black	72.6 (70.8-74.5)	52.1 (48.9-55.5)	52.8 (50.2-55.5)	51.1 (45.8-56.9)	50.0 (45.1-55.6)	51.9 (49.2-54.9)	52.5 (50.2-55.0)	87.2 (85.7-88.7)
White	87.0 (86.6-87.5)	62.3 (61.3-63.3)	62.8 (61.9-63.8)	58.9 (57.5-60.4)	62.7 (61.2-64.2)	61.3 (60.5-62.2)	63.0 (62.2-63.8)	93.5 (93.0-93.9)
US registries (all races)	84.0 (83.8-84.2)	60.2 (59.8-60.6)	60.2 (59.8-60.6)	57.0 (56.4-57.6)	59.3 (59.2-60.5)	59.3 (58.9-59.6)	60.3 (60.0-60.6)	93.3 (92.1-92.5)
Black	70.9 (70.0-71.8)	51.5 (50.0-53.1)	51.0 (49.8-52.3)	47.4 (44.7-50.1)	49.4 (47.3-51.7)	50.5 (49.1-51.8)	50.8 (49.7-51.9)	85.8 (85.0-86.6)
White	84.7 (84.5-84.9)	60.5 (60.0-60.9)	60.8 (60.4-61.2)	57.3 (56.6-57.9)	60.4 (59.7-61.1)	59.6 (59.2-59.9)	60.8 (60.5-61.2)	92.4 (92.2-92.7)

RS=relative survival. S=Surveillance, Epidemiology and End Results (SEER) registry. N=National Program of Cancer Registries (NPCR) registry. See text for attribution of registries to NPCR and SEER. R=raw (not age-standardised) survival estimate; too few cases in one or more age groups. *International Cancer Survival Standard (see text). †Survival truncated if greater than 1.0 (100%). ‡Survival estimates based on fewer than five patients are not shown (NA=not applicable). Black populations are not shown separately for Hawaii, Idaho, New Mexico, Utah, or Wyoming, because it was not possible to estimate relative survival for blacks in these states with race-specific life tables (see text).

Table 3: 5-year relative survival (%) by use of state-specific and race-specific life tables and age-standardised to ICSS weights* with 95% CIs† for adults (aged 15-99 years) diagnosed with cancer of the breast (women), colon, rectum, or prostate during 1990-94 and followed up to Dec 31, 1999, by race: US populations

Differences between survival estimates are presented as the absolute value, eg, 15% is given as 5% (not 50%) higher than 10%.

We analysed individual data for almost 2 million adults who were diagnosed with a first, primary, malignant, invasive neoplasm of the breast (women), colon, rectum, or prostate during the period 1990–94 and who had been followed up to ascertain their vital status for at least 5 years after diagnosis until the end of 1999 or later. Data were contributed by 101 population-based cancer registries covering a combined population of almost 300 million persons living in 31 countries (table 1 and webfigure 1). Canada and the USA contributed 1.07 million patients (54% of the total) from a population base of 125 million. The 24 European countries contributed 740 000 patients (37%) from a population base of 126 million, indicating lower mean incidence of cancer than in North America.

The smallest dataset came from Sétif (Algeria), covering a population of 1.1 million, some 4% of the national population. The registry could only provide data for the period 1992–94, the population is young, and cancer risks are currently low on the global scale.⁴³ The dataset was therefore small, a total of 300 patients. This decreases the statistical precision of survival estimates, but no patient was detected solely at death certification or autopsy, and the vital status of every patient was ascertained at a home visit by registry staff, something no other registry could deliver. Some of the datasets for black patients in US states were of similar size (webtable). California provided the largest single dataset of 240 000 patients diagnosed during 1990–94 in a population of 31 million (12% of the US population), with a very high cancer risk on the global scale (table 1).

For 16 of the 31 countries, the data covered 100% of the national population (table 1). The proportion of the national population covered by the data for the other 15 countries ranged from less than 10% (Algeria, Brazil, Japan, Austria, Czech Republic, France, Germany, Poland) to 10–29% (Italy, Portugal, Spain, Switzerland) and 30% or more (Canada, USA, the Netherlands).

Most registries provided data on patients diagnosed during the entire 5-year period 1990–94, but ten registries provided data for shorter periods (table 1).

Data for all four index cancers were provided by 89 of the 101 registries. Two specialised registries in Côte-d'Or (France) only collect data on cancers of the breast or colorectum, respectively, whereas ten general registries that collect data for all cancers only contributed data for selected cancers: breast (Isère, France; northern Netherlands; all five Swiss registries); breast, colon, and prostate (Campinas, Brazil; Nova Scotia, Canada), or breast, colon, and rectum (Granada, Spain; table 1).

Ethical approval for the CONCORD study³¹ was obtained from the Istituto Superiore di Sanità, Rome, Italy (CE-ISS 02/03, May 20, 2002) and from the Institutional Review Board of the CDC, Atlanta, GA, USA (IRB #3551, July 24, 2002). SEER data were obtained from the public-use dataset.³⁸ For other registries, anonymised data were trans-

mitted to the CONCORD Data Analysis Centre at the Istituto Superiore di Sanità by use of special courier delivery of encrypted and password-protected CD-ROMs with separate email transmission of the password, or pre-planned deposition of password-protected files on a specially created File Transfer Protocol (FTP) site from which the data were immediately removed in Rome. Each tumour record included a serial number for the purposes of quality control with the originating cancer registry.

Role of the funding source

The pilot study (January, 2000 to March, 2000) was funded by the UK Department of Health (£75 000). The CDC funded data collection and the costs of linkage to the National Death Index for the phase I study in participating registries in the National Program of Cancer Registries (US\$3 million). The Cancer Survival Group (including BR, MQ) in the London School of Hygiene and Tropical Medicine, London, UK, has been funded by Cancer Research UK (grant C1336/A5735) since April, 2005. Funding applications were open and competitive. None of the funding sources had any role in design, data collection, analysis, interpretation of the data, or writing of this article. MPC, MQ, RdA, RC, SF, MSantaquilani, and AV had access to the raw data. The corresponding author had full access to all of the data and the final responsibility to submit for publication.

Results

The background risk of death in the general population varied widely between the participating countries and regions. The mean life expectancy at birth during the decade 1990–99 ranged from 63.7 to 77.6 years in men and from 70.9 to 83.7 years in women.⁴⁶ In the USA, the range of life expectancies in white and black populations did not overlap at all in the states and metropolitan areas for which life tables could be constructed for both groups. The ranges for men were 64.0–70.1 years in blacks and 71.1–75.9 years in whites, whereas the ranges in women were 73.3–76.5 years in blacks and 78.8–80.9 years in whites.

The cumulative risk of death from all causes over the age range 15–59 years in the general population of the participating countries and regions ranged widely, from 9% to 34% in men and from 5% to 17% in women. Over the age range 60–84 years, the cumulative risk of death ranged from 60% to 86% in men and from 40% to 75% in women.⁴⁶

Of 785 255 records of breast cancer submitted for analysis, 45 020 (6%) related to women registered with a previous primary cancer, and were excluded (available online⁴⁶). Of the 740 235 eligible first primary invasive breast cancers, 9215 records were excluded as death-certificate-only (DCO) registrations (1%), 239 as autopsy-detected tumours (<1%) and 2064 with major errors (<1%), leaving 728 717 patients for analysis (98% of those eligible), of whom 370 000 (51%) were resident in North America and 304 000 (42%) in Europe (table 1). Almost all (97%) of the

tumours included in the analyses were microscopically verified, less than 1% of patients were censored from the analysis within 5 years of diagnosis, and 2.3% died within 1 month of diagnosis.

Relative survival at 5 years, age-standardised to the International Cancer Survival Standard weights, ranged from 80% or over in North America, Sweden, Japan, Finland, and Australia to less than 60% in Brazil and Slovakia, and below 40% in Algeria (table 2 and figure 1). Survival in the 24 European countries that contributed to CONCORD was mostly in the range 70–79%.

The survival estimate of 38.8% (95% CI 31.4–46.2) for Sétif (Algeria) is based on 180 patients, and it is not age-standardised because there were too few patients for analysis in some age groups, but age-standardisation for breast cancer in other datasets rarely altered the raw estimate by more than 5% in either direction (data not shown), and survival from breast cancer was undoubtedly much lower in Algeria than in all the other countries.

The pooled estimate of 5-year survival for the two Brazilian registries was 58.4%, but the estimate for Goiânia (65.4%) is more reliable than the very low figure for Campinas (36.6%), where high proportions of patients were excluded as DCO or with major errors (available online¹⁶). The proportion of metastatic tumours was higher in Campinas, however. The 5-year survival estimate for Cuba was 84.0%, but this was likely to be an over-estimate: some 28% of records were excluded because they were registered solely from a death certificate.

The pooled estimate of 5-year survival for Canada was 82.5%, with a narrow range from 79.3% in Nova Scotia to 85.4% in British Columbia (table 2 and figure 1). In the USA, 5-year relative survival for all races combined ranged from 78–81% in New York City, New York State, and Louisiana to 89–90% in Hawaii and Seattle, WA (table 2), but most of the estimates were within a fairly narrow range, from 82% to 87% (figure 2). Survival in metropolitan areas covered by SEER registries was similar to that in the respective states: Detroit, MI 83.0% and Michigan State 82.3%; San Francisco, CA 86.2% and California State 84.6%. 5-year survival was 77.4% for residents of New York City, NY (with 40% of the state population), slightly lower than for New York State as a whole, 81.0%.

Survival was lower for blacks than for whites in all 17 populations in the USA for which this could be assessed with race-specific life tables (webfigure 2). The age-adjusted pooled estimate of 5-year survival was 84.7% for whites (range 80–90%) and 70.9% for blacks (table 3). The range in survival was wider for blacks (60–83%), but the values at both extremes of the range were based on relatively few patients and have wider CIs. Within a given US population, the absolute difference in age-adjusted relative survival between blacks and whites ranged from 2% (Rhode Island, Nebraska) to 25–27% (Iowa and Seattle, WA; table 3 and webfigure 2). Even in areas where blacks comprise 25% or more of the population, survival for black women was 8–14% below the lowest estimate for white women (79.6%)

in any of the participating areas: Atlanta, GA (71.1%), Detroit, MI (71.7%), New York City, NY (65.8%), and Louisiana (69.9%). The pooled estimate of 5-year survival for the USA was 84.0%, with 86.1% in areas covered by SEER and 83.1% in areas covered by NPCR (table 3).

Survival in black women was always lower than the mean survival for all US populations included, and often more than three standard deviations below it (below the 99.8% control limits), after controlling for the precision of the estimates. Survival in white women is generally within or above the upper control limits, especially in territories covered by the SEER Program. A notable exception is for white women in New York State, including New York City, where the survival estimates are precise, but well below the lower control limits (webfigure 3).

The pooled estimate of 5-year survival for breast cancer in Japan was 81.6%. Survival in Osaka (79.4%) was lower than in Fukui (83.1%) and Yamagata (87.3%; table 2 and figure 1).

5-year relative survival for breast cancer in Europe, age-standardised to the ICSS weights, ranged from 57.9% in Slovakia to 82.0% in Sweden (table 2 and figure 1), whereas the pooled estimate derived from the data of 58 registries in the 24 participating European countries was 73.1%. Survival estimates for most of these countries have been reported.¹⁷ The CONCORD study includes additional data from four countries: 5-year survival was 69.6% in Ireland and 72.0% in Northern Ireland, similar to the UK mean value of 69.7% (table 2). In Switzerland, 5-year survival in the cantons of St Gallen-Appenzell, Graubünden-Glarus, and Valais was 72–75%, about 4–7% lower than in Geneva or Basel. 5-year survival was 77.8% in northern Netherlands, similar to that in Amsterdam and southern Netherlands (76–78%).

The national estimate of 5-year survival for breast cancer in Australia was 80.7%. Survival was virtually identical in the six largest states (96% of the national population), in the range 80–82%, but notably lower in the two smallest regions: 71.9% in Northern Territory (1.0% of the population) and 77.1% in Tasmania (2.6%).

Of 488741 colon cancer records submitted for analysis, 45862 records (9%) were excluded for a previous cancer, leaving 442879 first, primary, invasive colon cancers eligible for analysis (available online¹⁶). A further 13102 (3%) were excluded as DCO registrations, 1534 (<1%) as autopsy-detected tumours, and 1144 (<1%) as major errors, leaving 427099 patients for inclusion in the analyses (96% of those eligible). Of these, 214000 (50%) were resident in North America, 170000 (40%) in Europe, and 30300 (7%) in Australia. Cancers of the colon comprised 67% of all colorectal tumours (table 1). Microscopic verification was high (94%), and less than 1% of patients were censored from the analysis within 5 years of diagnosis. Almost 11% of patients died within the first month after diagnosis.

Relative survival at 5 years, age-standardised to the ICSS weights, ranged from about 60% in North America, Japan, Australia, and some western European countries down to

40% or less in Algeria, Brazil, Czech Republic, Estonia, Poland, Slovenia, and Wales (table 2 and figure 3).

The survival estimates of 11.4% (95% CI 0.7–40.9) for men and 30.6% (9.5–56.1) for women in Sétif (Algeria) were based on fewer than 20 patients, and are not age-standardised, but survival was clearly lower in Algeria than in all the other countries.

The estimate of 5-year relative survival for Goiânia (48.1% in men, 44.8% in women) was more plausible for Brazil than the low estimates for Campinas, where 26% of patients had to be excluded with errors.³⁴ 5-year survival in Cuba was about 60% in both sexes, although more than half the patients were excluded from analysis as DCOs.³⁴

In Canada, the pooled estimate of 5-year survival was 56.1% for men and 58.7% for women. Variation between provinces was small, from 54–57% in men and 58–60% in women (table 2 and figure 3). In the USA, 5-year relative survival for all races combined was 60.1% in both sexes, with a range from 53.6% for women in New York City to 67.9% for men in Hawaii (table 2 and figure 2). Again, most of the estimates were within a narrow range, 59–64%.

5-year survival for colon cancer among blacks in the USA was lower than among whites. In 34 paired observations of this difference in survival (men and women in 17 populations), only three exceptions were noted, in men and women in Iowa and men in Nebraska. The estimates for blacks in those three areas were based on fewer than 50 patients, have wide confidence intervals and were not age-standardised (table 3 and webfigure 4). The pooled estimate of age-adjusted 5-year relative survival for the USA was 61% for white men and women, and 51–52% for black men and women. Within a given population, the absolute difference between blacks and whites ranged from 2.6% in men and 7.3% in women in Los Angeles, CA to 14.3–17.2% in Colorado. The geographical range in black-white differences in survival is affected by small populations to some extent, but even in areas where blacks comprise 25% or more of the population (Atlanta, GA, Detroit, MI, New York City, NY, Louisiana), 5-year survival from colon cancer in blacks was 6–12% lower than for whites in the same population (table 3). The pooled estimate of 5-year survival in areas covered by NPCR was 59.8% for men and 59.6% for women, and 61.9% for men and 62.1% for women in SEER areas.

Age-standardised survival in whites ranged from 54.9% to 67.9% (table 3 and webfigure 4). The range of age-standardised survival for blacks was 45.0% to 59.9%. Survival in blacks was generally lower than the mean value for all included US populations and often more than three standard deviations below the mean, after controlling for precision of the survival estimates. Survival in whites was generally within the control limits. The main exception was for white women in New York State, including New York City, NY, where survival estimates were precise, but more than three standard deviations below the lower control limits around the pooled US estimate (webfigure 3).

In Japan, the pooled survival estimate was 63.0% in men and 57.1% in women, although survival was about 10% lower in Osaka prefecture than in Fukui or Yamagata (table 2 and figure 3).

In Europe, 5-year relative survival for colon cancer in men ranged from 28.5% in Poland to 54–57% in Spain, Finland, Austria, and France. In women, the lowest estimate was also for Poland (30.9%), while survival was in the range 55–60% in nine countries (table 2 and figure 3). The pooled estimates for the 51 contributing registries in 23 European countries were 46.8% in men and 48.4% in women. Data on colon cancer were not available for the five Swiss registries, Isère (France), or northern Netherlands. Survival estimates for most of these countries have been published elsewhere.²⁷ This study included additional data from two countries. 5-year survival in Ireland was 49.1% in men and 48.5% in women. The estimates for Northern Ireland were 47.3% in men and 49.0% in women, slightly higher than the pooled estimate for the UK, 43.5% in men and 44.4% in women (table 2).

The national estimate of 5-year survival for colon cancer in Australia was 57.8% in men and 57.7% in women. Survival ranged from 50–62% in the eight states and territories: it was highest in New South Wales, the Australian Capital Territory, and Queensland, and lowest in Tasmania, Northern Territory, and Western Australia (9.6% of the population; table 1).

Of 233 176 rectal-cancer records submitted for analysis, 15 731 records (7%) were excluded for a previous cancer, leaving 217 445 first, primary, invasive rectal cancers eligible for analysis (available online³⁵). A further 3213 (1%) were excluded as DCO registrations, 517 (<1%) as autopsy-detected tumours and 574 (<1%) as major errors, leaving 213 141 patients for inclusion in the analyses (98% of those eligible). Of these, 83 000 (39%) were resident in North America, 106 000 (50%) in Europe, and 16 800 (8%) in Australia. Cancers of the rectum comprised 33% of all colorectal tumours (table 1). Microscopic verification was high (96%). Less than 1% of patients were censored from the analysis within 5 years of diagnosis. Almost 8% died within the first month after diagnosis.

5-year relative survival from rectal cancer, age-standardised to the ICSS weights, ranged from about 60% to around 20% in both sexes, with Japan, Canada, the USA, France, the Netherlands, Sweden, and Australia at the upper end of the range, and Algeria, Estonia, Poland, and Slovakia at the lower end (table 2 and figure 3).

The 5-year survival estimates of 25.9% (95% CI 11.4–43.7) for men and 18.2% (6.6–34.6) for women in Sétif (Algeria) were each based on 30 patients, and were not age-standardised because data were too sparse in some age groups.

5-year relative survival in Goiânia, Brazil, was 49.3% in men and 38.4% in women. No data were available for Campinas. 5-year survival in Cuba was 59.2% in men and 62.8% in women, based on analysis of about 700 patients

See Online for webfigure 4