

Current Status of Emergency Care for ST-Elevation Myocardial Infarction in an Urban Setting in Japan

Naoki Nakayama, MD¹; Kazuo Kimura, MD¹; Tsutomu Endo, MD²; Kazuki Fukui, MD³; Hideo Himeno, MD⁴; Yuji Iwasawa, MD⁵; Yasuyuki Mochida, MD⁶; Yukiko Morita, MD⁷; Makoto Shimizu, MD⁸; Tomoaki Shimizu, MD⁹; Takahiro Takei, MD¹⁰; Keiko Yoshida, MD⁸; Atsushi Wada, MD¹¹; Satoshi Umemura, MD¹²

Background The door-to-balloon time (DTB) is an important predictor of the outcome for patients with ST-elevation myocardial infarction (STEMI). In Japan, percutaneous coronary intervention (PCI) can be performed at many hospitals, and the predominant strategy for reperfusion therapy is primary PCI. However, it remains unclear how rapidly reperfusion is achieved at these hospitals.

Methods and Results The study group comprised 369 patients with STEMI who presented within 12 h of symptom onset to a tertiary emergency center (TEC) or at 11 community hospitals (CHs) in 2006 and underwent emergency coronary angiography. Median DTB was shorter in the TEC (63 vs 104 min, $P < 0.001$), and the rate of DTB within 90 min was higher in the TEC (96% vs 39%, $P < 0.001$). Lateral myocardial infarction, presentation during off-hours, and non-cardiologist as the first-contact physician were significantly associated with a prolonged DTB in CHs. There was a trend toward lower 30-day mortality from all causes in the TEC (2.0% vs 4.8%, $P = 0.08$). Multiple logistic regression analysis demonstrated that prolonged DTB (> 90 min) was an independent predictor of 30-day mortality (odds ratio 12.6; 95% confidence interval 1.85–86.2, $P = 0.01$).

Conclusions Establishment of emergency cardiac care systems with the goal of DTB within 90 min is required in PCI-capable hospitals to improve clinical outcomes.

Key Words: Acute myocardial infarction (AMI); Door-to-balloon time; Mortality; Reperfusion

The time from hospital admission to reperfusion is an important predictor of the outcome for patients with ST-elevation myocardial infarction (STEMI)^{1–3} Guidelines for the management of STEMI issued by the American College of Cardiology/American Heart Association and the European Society of Cardiology recommend primary percutaneous coronary intervention (PCI) for reperfusion therapy when it can be performed in a timely manner (ie, door-to-balloon time (DTB) or first medical contact-to-balloon time within 90 min) by experienced operators at high-volume centers.^{4,5} However, emergency

primary PCI is not always feasible, and treatment is frequently delayed. In Japan, PCI can be performed at many hospitals, and the predominant strategy for reperfusion therapy is primary PCI^{6–9} but how rapidly reperfusion is achieved in Japanese hospitals that routinely perform PCI remains unclear. Drugs, devices, and techniques for primary PCI have been actively debated, whereas shortening the time to reperfusion has only recently received attention. This study evaluated DTB and analyzed its components in patients with STEMI who were admitted to emergency hospitals in an urban setting in Japan and examined differences in DTB according to the type of hospital.

Methods

Study Population

We retrospectively studied 369 patients (283 men, 86 women; mean age, 65 years) with STEMI who were admitted to 12 emergency hospitals in an urban setting in Japan and underwent emergency coronary angiography within 12 h after symptom onset. Patients with out-of-hospital cardiac arrest were excluded. All subjects had to have typical chest pain lasting for 20 min or more and ST-segment elevation in at least 2 adjacent leads (> 0.1 mV in leads II, III, aVF, aVL, V4–6 and > 0.2 mV in leads V1–3) on the admission electrocardiogram (ECG). The diagnosis of acute myocardial infarction (AMI) was confirmed by an increase in the creatine kinase level to more than twice the upper limit of normal. Another 20 patients with STEMI within 12 h after symptom onset were hospitalized during the study period, but did not undergo urgent cardiac catheterization because of advanced age or comorbidity. No patient

(Received June 4, 2008; revised manuscript received September 29, 2008; accepted October 15, 2008; released online January 15, 2009)

¹Division of Cardiology, Yokohama City University Medical Center,

²Department of Cardiology, Saiseikai Yokohama Nambu Hospital, ³De-

partment of Cardiology, Kanagawa Prefectural Cardiovascular and

Respiratory Center, Yokohama, ⁴Department of Cardiology, Fujisawa

Municipal Hospital, Fujisawa, ⁵Department of Cardiology, Yokosuka

Municipal Hospital, Yokosuka, ⁶Department of Cardiology, Omori Red

Cross Hospital, Tokyo, ⁷Department of Cardiology, National Hospital

Organization Sagami National Hospital, Sagami, ⁸Depart-

ment of Cardiology, International Goodwill Hospital, Yokohama,

⁹Department of Cardiology, Prefectural Ashigara-kami Hospital,

Kanagawa, ¹⁰Department of Cardiology, Fujisawa Shounandai Hospi-

tal, Fujisawa, ¹¹Department of Cardiovascular Medicine, Chigasaki

Municipal Hospital, Chigasaki and ¹²Department of Medical Science

and Cardiorespiratory Medicine, Yokohama City University Graduate School

of Medicine, Yokohama, Japan

Mailing address: Kazuo Kimura, MD, Division of Cardiology,

Yokohama City University Medical Center, 4-57 Urafune-cho,

Minami-ku, Yokohama 232-0024, Japan. E-mail: c-kimura@urahp.

yokohama-cu.ac.jp

All rights are reserved to the Japanese Circulation Society. For permis-

sions, please e-mail: cj@j-circ.or.jp

Table 1 Capabilities of the 2 Types of Hospitals

	TEC (n=1)	CHs (n=11)
No. beds	720	239-601*
No. cardiologists	14	4-8*
Annual no. total PCI \geq 200	+	3/11
Interventional capability for 24 h/7 days	+	10/11
Cardiologists on duty for 24 h/7 days	+	3/11
Cardiologists always initially evaluate patients with chest discomfort in the emergency department	+	1/11
On-site cardiac surgical back-up	+	3/11
Pre-hospital 12-lead ECG transmission system	+	-

*Range.

TEC, tertiary emergency center; CHs, community hospitals; PCI, percutaneous coronary intervention; ECG, electrocardiogram.

Table 2 Baseline Characteristics of the Patients

	TEC (n=98)	CHs (n=271)	P value
Age (years)	62.3 \pm 10.8	66.6 \pm 11.7	0.002
Gender, M	80 (82%)	203 (75%)	0.18
Medical history			
Hypertension	57 (58%)	148 (55%)	0.54
Dyslipidemia	44 (45%)	116 (43%)	0.66
Diabetes mellitus	36 (37%)	102 (38%)	0.93
Previous myocardial infarction	8 (8%)	20 (7%)	0.81
Previous bypass surgery	2 (2%)	4 (1%)	0.51
Infarct location			
Anterior	38 (39%)	130 (48%)	0.12
Inferior	48 (49%)	105 (39%)	0.08
Lateral	7 (7%)	21 (8%)	0.85
Time and mode of presentation			
Presentation during regular hours	49 (50%)	134 (49%)	0.93
Presentation by self-transport	4 (4%)	69 (25%)	<0.001
Inter-hospital transfer	30 (31%)	57 (21%)	0.06
Cardiologist as 'first-contact' physician	94 (96%)	117 (45%)	<0.001
Killip class \geq 2	10 (10%)	34 (13%)	0.54
Fibrinolytic therapy with facilitated PCI	42 (43%)	93 (34%)	0.13

Abbreviations see in Table 1.

received thrombolytic therapy without coronary angiography. All clinical data were collected from medical records by a single researcher. The 12 participating institutions consisted of 1 tertiary emergency center (TEC) and 11 community hospitals (CHs), all of which could perform PCI. The study period was from January through December 2006.

Data Analysis and Definition

We compared the institutional and patients' baseline characteristics between the TEC and CHs. Time factors such as DTB and its components were also compared. We then analyzed variables associated with treatment delays and clinical outcomes.

Door time was defined as the arrival time at the emergency department where emergency cardiac catheterization was performed (not the arrival time at referral hospitals for transferred patients). Balloon time was defined as the time of first balloon inflation, or the time of stent implantation if the patient underwent primary stenting. PCI was considered successful if a Thrombolysis In Myocardial Infarction (TIMI) flow grade of 2-3 was obtained in the infarct-related artery in the presence of <50% residual stenosis with balloon angioplasty or <20% residual stenosis with stent implantation. If initial angiography showed TIMI grade 3 flow with >75% stenosis, additional PCI was left to the discretion of the attending physician.

Presentation time was divided into "regular hours"

(defined as weekdays from 08.30-19.59 h) and "off-hours" (weekdays from 20.00-08.29 h and weekends).

Statistical Analysis

Statistical analysis was performed with SPSS 11.0 for Windows (SPSS, Inc, Chicago, IL, USA). The Kolmogorov-Sminov test was used to analyze the normality of the distribution of the study variables. Values are expressed as mean \pm SD or median with interquartile ranges. Univariate comparisons of continuous variables between groups were done with unpaired t-tests for normally distributed variables and Mann-Whitney tests for skewed variables. Categorical variables were compared with chi-square tests or Fisher's exact test. Multivariable logistic-regression analysis was performed with the variables that were significantly related to outcome variables in the univariate analyses to determine predictors of DTB within 90 min and of 30-day mortality. Differences were considered significant at $P < 0.05$.

Results

Hospital Characteristics

Table 1 shows the characteristics of the hospitals. The TEC was a high-volume center with many strategies to reduce the DTB. An attending cardiologist is always available to examine patients who present with chest discomfort at the emergency department, as compared with only 1 of the 11

Table 3 Angiographic Findings and Interventional Procedure Rates

	TEC (n=98)	CHs (n=271)	P value
Initial TIMI flow grade 0	46 (47%)	124 (46%)	0.84
POBA and/or stent implantation	80 (82%)	231 (85%)	0.40
Thrombectomy	13 (13%)	83 (31%)	0.001
Distal protection	9 (9%)	3 (1%)	<0.001
Pre intravascular ultrasound	74 (76%)	47 (17%)	<0.001
Temporary pacing	1 (1%)	22 (8%)	0.001
Final TIMI flow grade 3	86 (88%)	242 (89%)	0.68
PCI success	77 (96%)	226 (98%)	0.34

TIMI, Thrombolysis In Myocardial Infarction; POBA, plain old balloon angioplasty. Other abbreviations see in Table 1.

Table 4 Time Factors Related to Reperfusion

	TEC	CHs	P value
Symptom onset-admission (min)	83 (51, 142)	100 (56, 180)	0.08
Admission-ECG (min)	4 (3, 5)	4 (2, 7)	0.67
ECG-catheterization laboratory (min)	20 (16, 25)	45 (35, 69)	<0.001
Catheterization laboratory-CAG (min)	13 (11, 16)	18 (15, 23)	<0.001
CAG-balloon (min)	24 (20, 31)	26 (19, 38)	0.28
Door-balloon (min)	63 (55, 70)	101 (83, 134)	<0.001
DTB within 90 min	96%	39%	<0.001

Values are median with interquartile ranges.

CAG, first coronary angiography; DTB, door-to-balloon time. Other abbreviations see in Table 1.




Fig 1. Median door-to-balloon time (DTB) during regular hours and off-hours in each hospital type. TEC, tertiary emergency center; CHs, community hospitals.

CHs. The TEC has a pre-hospital 12-lead ECG transmission system from the ambulance to the emergency department, and ECG transmission was performed in approximately 50% of the TEC patients.

Patients' Characteristics

Within the overall study group, 98 (27%) patients were admitted to the TEC and 271 (73%) to CHs. There was no difference in sex, infarct location, or the rate of Killip class ≥ 2 between the 2 types of institution. The patients admitted to TEC were younger than those admitted to CHs. Fibrinolytic therapy with low-dose tissue plasminogen activator with facilitated PCI was performed in 42 patients (43%) in the TEC and in 93 (34%) patients in the CHs. Overall, 15 patients who received fibrinolytic agents did not undergo PCI

because of successful reperfusion as confirmed by coronary angiography, without balloon angioplasty. These patients were excluded from the analysis regarding the time from CAG to first balloon inflation. First-contact physicians were more frequently cardiologists in the TEC. The proportion of patients transferred from other hospitals was higher in the TEC, whereas the rate of self-transport to the hospital was significantly lower (Table 2).

Angiographic Findings and Interventional Procedure Rate

The frequency of PCI-related procedures such as thrombectomy, use of distal protection devices, and pre-intravascular ultrasound differed between the 2 types of institution. Balloon angioplasty (POBA), stent implantation, or both were performed in 82% of the patients in the TEC and in

Table 5 Variables Associated With Prolonged DTB (>90 min) in Patients Admitted to CHs

	Univariate analysis	P value	Multivariate analysis	95%CI	P value
Annual PCI procedures <200 in admitting hospital*	1.09	0.76	—	—	—
Cardiogenic shock	0.36	0.045	—	—	0.16
Lateral wall infarction	3.45	0.043	3.85	1.01–14.8	0.049
Presentation during off-hours	3.39	<0.001	2.94	1.63–5.31	<0.001
Presentation by self-transport	2.3	0.014	—	—	0.10
Inter-hospital transfer	0.41	0.004	—	—	0.37
Non-cardiologist as 'first-contact physician'	4.13	<0.001	3.63	2.02–6.52	<0.001

ORs of prolonged DTB are shown. Variables reaching $P < 0.05$ on univariate analysis were entered into the multivariate analysis.

*Although annual PCI volume is included, it was not associated with prolonged DTB and was not entered into the multivariate analysis.

CI, confidence interval; OR, odds ratio. Other abbreviations see in Tables 1, 4.

Table 6 Multivariate Predictors of All-Cause 30-Day Mortality

	OR	95%CI	P value
Age >75 years	11.4	2.05–63.3	0.005
DTB >90 min	12.6	1.85–86.2	0.01
Killip class ≥ 2	34.3	5.56–211.2	<0.001
Final TIMI flow grade ≤ 2	25.5	2.75–236.7	0.004
Anterior myocardial infarction	11.8	1.34–103.3	0.03

Variables reaching $P < 0.05$ on univariate analysis were entered into the multivariate analysis.

Abbreviations see in Tables 3, 4.

85% of those in CHs. PCI was successful in 95% of the patients treated in the TEC and in 98% of those treated in CHs. The proportion of patients with a final TIMI flow grade of 3 did not differ between the TEC and CHs (Table 3).

Time Factors Related to Reperfusion and Clinical Outcome

The time from symptom onset to admission was slightly but not significantly longer in the CHs. The time from admission to ECG recording and the time from first coronary angiography to first balloon inflation were similar in the 2 types of hospital. However, the time from ECG recording to the cardiac catheterization laboratory was significantly longer in the CHs, resulting in a longer DTB. The TEC had a median DTB of 63 min and achieved a guideline-recommended DTB within 90 min in 96% of patients, whereas the CHs had a median DTB of 101 min and only 39% of patients achieved a DTB within 90 min (Table 4). Presentation during off-hours significantly prolonged the DTB for 27 min as compared with presentation in regular hours in the CHs, whereas the time of presentation did not influence the DTB in the TEC (Fig 1). Moreover, treatment was markedly delayed (DTB >150 min) in 18% of the patients treated at CHs, and this increased to 30% when a non-cardiologist was the first-contact physician. There was a trend toward lower 30-day mortality from all causes in the TEC in the overall study group (2.0% vs 4.8%, $P = 0.08$).

Variables Influencing DTB in CHs

Because a guideline-recommended DTB was achieved in most of the patients treated in the TEC, the variables that influence the DTB were analyzed among the patients who were admitted to a CH. Hospitals that performed more than 200 PCI per year did not have a shorter DTB than hospitals that performed less procedures. After multivariable adjustment, lateral MI, presentation during off-hours, and non-cardiologist as first-contact physician were significantly

associated with a prolonged DTB (Table 5). Inter-hospital transportation occurred in 21% of patients admitted to a CH, but was not a significant factor for DTB >90 min on multivariate analysis. After excluding referred patients and analyzing data only from patients with direct transportation, the results for time factors were similar to those obtained for the study group as a whole.

Analysis of Predictors of 30-Day Mortality

Table 6 shows the predictors of 30-day mortality from all causes in patients who underwent emergency PCI. Multiple logistic regression analysis showed that age >75 years, anterior MI, Killip class ≥ 2 , final TIMI flow grade ≤ 2 , and DTB >90 min were independent predictors of mortality within 30 days.

Discussion

Our study showed that CHs achieved the guideline-recommended DTB in less than 40% of patients with STEMI, and that prolonged DTB was an independent predictor of 30-day mortality. Delayed treatment was associated with patient presentation during off-hours and non-cardiologists as the first-contact physician. In contrast, the TEC, which specialized in cardiac emergency care, performed rapid reperfusion therapy in most patients.

Prompt PCI significantly reduces mortality and morbidity in patients with STEMI, and prolonged DTB has been associated with poor outcomes.^{1–5,10} However, most previous studies were performed in Western countries at medical centers with experienced PCI providers. We investigated whether a similar situation exists in Japanese hospitals. In contrast to Western countries, many hospitals in Japan can perform PCI and consequently, 75–94% of patients with STEMI undergo primary PCI for reperfusion therapy^{6–8} including those treated at low-volume centers.^{11,12} Our findings showed that PCI volume at the CHs was unrelated to

either DTB or 30-day mortality in contrast to results obtained by Western studies, where a prolonged DTB was associated with higher 30-day mortality. This outcome suggests that the establishment of emergency cardiac care systems is a more important factor in the reduction of DTB than the treatment of patients with STEMI at high-volume centers, consistent with the recommendations of "Development of System of Care for ST-Elevation Myocardial Infarction," issued by the American Heart Association.¹³ A recent study has shown that hospital policies that emphasize having a well-equipped catheterization laboratory staffed by attending cardiologists and other dedicated care providers contributes to a reduced DTB.¹⁴ In another study, the establishment of protocols requiring that the cardiac catheterization laboratory is activated by emergency department physicians and that patients with cardiac problems are immediately transferred and treated at catheterization laboratory significantly reduced DTB by 38 min overall,¹⁵ contributing to a decrease in mean infarct size.

Our study found that non-cardiologists as first-contact physician and off-hour presentation were associated with prolonged DTB in Japan. In the CHs, more than half of the patients with STEMI were initially treated by a non-cardiologist, who quickly obtained a 12-lead ECG for patients who had symptoms suggesting STEMI, but the time from ECG to the catheterization laboratory remained prolonged. The time from first coronary angiography to first balloon inflation, the PCI success rate, and the proportion of patients with a final TMI flow grade of 3 did not differ between the CHs and TEC, suggesting the cardiologists were unaware of treatment delays that occurred before they were consulted. Ideally, all patients with symptoms suggesting STEMI should be examined by a cardiologist immediately after admission; however, most CHs do not have a 24-h on-site attending cardiologist 7 days per week because of limited human and financial resources. Non-cardiologists should therefore be aware of the clinical significance of the DTB and immediately transfer patients suspected to have STEMI to the cardiac catheterization laboratory.

In Vienna, Austria, the establishment of a cooperative network and implementation of the guidelines for the treatment of STEMI led to earlier reperfusion therapy and significantly improved clinical outcomes.¹⁶ Pre-hospital ECG transmission enables on-call cardiologists to be notified before the patient arrives in the emergency department and reduces PCI-related delays, with earlier activation of the catheterization laboratory.¹⁷⁻¹⁹ We believe that pre-hospital ECG transmission and analysis contributed substantially to the higher rate of a DTB within 90 min at the TEC in our study.

In conclusion, our results suggest that the design and implementation of appropriate guidelines and the improvement of hospital performance through the organization and establishment of emergency cardiac care systems will consistently ensure rapid reperfusion in patients with STEMI. Optimal policies tailored to meet the needs of specific communities are likely to reduce the DTB and lead to improved outcomes.

Study Limitations

This study included a small number of patients and was performed in a single region of Japan. Overall, 37% of patients underwent facilitated PCI with a half-dose of tissue plasminogen activator; negative results after facilitated PCI have been reported,^{20,21} but several studies, including the

Comparison of Angioplasty and Prehospital Thrombolysis In acute Myocardial infarction (CAPTIM) trial,²² the Vienna registry,¹⁶ and the National Registry of Myocardial Infarction (NRFMI),¹⁸ have demonstrated the effectiveness of fibrinolytic therapy early after the onset of MI. In addition, the former 2 trials also confirmed the efficacy of immediate PCI after fibrinolytic therapy.^{16,22,23} We therefore administered a half-dose of fibrinolytic therapy followed by PCI in younger patients who presented early after onset.

Although the DTB at the TEC was within guideline recommendations in nearly all patients with STEMI, a direct comparison with the results of previous Western studies is precluded by differences in the definition of the DTB for transferred patients. Indeed, inter-hospital transfer has been found to be a strong independent predictor of a delayed time from first hospital arrival to first balloon inflation,²⁴ but the present study focused on hospital performance to identify in-hospital problems that need to be solved. Furthermore, we must recognize that emergency care systems in Japan might differ considerably from those in Western countries.

Conclusion

A DTB longer than 90 min was associated with a higher 30-day mortality in Japanese patients with STEMI. Guideline-recommended DTB was achieved in most of the patients in the TEC, as compared with less than 40% in CHs. Although the DTB was frequently longer in CHs, there was no correlation between PCI volume and DTB. Establishment of emergency systems for STEMI care with the goal of a DTB within 90 min is required in PCI-capable hospitals to improve clinical outcomes.

References

- Berger PB, Ellis SG, Holmes DR Jr, Granger CB, Criger DA, Betriu A, et al. Relationship between delay in performing direct coronary angioplasty and early clinical outcome in patients with acute myocardial infarction: Results from the Global Use of Strategies to Open Occluded Arteries in Acute Coronary Syndromes (GUSTO-IIb) trial. *Circulation* 1999; 100: 14-20.
- Cannon CP, Gibson CM, Lambrew CT, Shoultz DA, Levy D, French WJ, et al. Relationship of symptom-onset-to-balloon time and door-to-balloon time with mortality in patients undergoing angioplasty for acute myocardial infarction. *JAMA* 2000; 283: 2941-2947.
- Shavelle DM, Rasouli ML, Frederick P, Gibson CM, French WJ, National Registry of Myocardial Infarction Investigators. Outcome in patients transferred for percutaneous coronary intervention (a national registry of myocardial infarction 2/3/4 analysis). *Am J Cardiol* 2005; 96: 1227-1232.
- Antman EM, Anbe DT, Armstrong PW, Bates ER, Green LA, Hand M, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: A report of the American College of Cardiology/American Heart Association task force on practice guidelines (Committee to revise the 1999 guidelines for management of patients with acute myocardial infarction). *J Am Coll Cardiol* 2004; 44: E1-E211.
- Van de Werf F, Ardissino D, Betriu A, Cokkino DV, Falk E, Fox KA, et al. Management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force on the Management of Acute Myocardial Infarction of the European Society of Cardiology. *Eur Heart J* 2003; 24: 28-66.
- Sakurai K, Watanabe J, Iwabuchi K, Koseki Y, Kon-no Y, Fukuchi M, et al. Comparison of the efficacy of reperfusion therapies for early mortality from acute myocardial infarction in Japan: Registry of Miyagi Study Group for AMI (MsAMI). *Circ J* 2003; 67: 209-214.
- Ue S, Chino M, Ishiki T. Rates of primary percutaneous coronary intervention worldwide. *Circ J* 2005; 69: 95-100.
- Kasanuki H, Honda T, Haze K, Sumiyoshi T, Horie T, Yagi M, et al. A large-scale prospective cohort study on the current status of therapeutic modalities for acute myocardial infarction in Japan: Rationale and initial results of the HJAMI Registry. *Am Heart J* 2005; 150:

- 411–418.
9. Matsui K, Kojima S, Sakamoto T, Ishihara M, Kimura K, Miyazaki S, The Japanese Acute Coronary Syndrome Study (JACSS) Investigators. Weekend onset of acute myocardial infarction does not have a negative impact on outcome in Japan. *Circ J* 2007; 71: 1841–1844.
 10. Keeley EC, Boura JA, Grimes CL. Primary angioplasty vs intravenous thrombolytic therapy for acute myocardial infarction: A quantitative review of 23 randomised trials. *Lancet* 2003; 361: 13–20.
 11. Nishigaki K, Yamazaki T, Fukunishi M, Tanihata S, Fujiwara H, for the Japanese Coronary Intervention Study (JCIS) Group. Assessment of acute myocardial infarction in Japan by the Japanese Coronary Intervention Study (JCIS) Group. *Circ J* 2004; 68: 515–519.
 12. Shiraishi J, Kohno Y, Sawada T, Nishizawa S, Arihara M, Hadase M, The AMI-Kyoto Multi-Center Risk Study Group. In-hospital outcomes of primary percutaneous coronary interventions performed at hospitals with and without on-site coronary artery bypass graft surgery. *Circ J* 2007; 71: 1208–1212.
 13. Jacobs AK, Antman EM, Faxon DP, Gregory T, Solis P. Development of systems of care for ST-elevation myocardial infarction patients: Executive summary. *Circulation* 2007; 116: 217–230.
 14. Bradley EH, Herrin J, Wang Y, Barton BA, Webster TR, Mattera JA, et al. Strategies for reducing the door-to-balloon time in acute myocardial infarction. *N Engl J Med* 2006; 355: 2308–2320.
 15. Khot UN, Johnson ML, Ramsey C, Khot MB, Todd R, Shaikh SR, et al. Emergency department physician activation of the catheterization laboratory and immediate transfer to an immediately available catheterization laboratory reduce door-to-balloon time in ST-elevation myocardial infarction. *Circulation* 2007; 116: 67–76.
 16. Kalla K, Christ G, Karnik R, Malzer R, Norman G, Prachar H, et al. Implementation of guidelines improves the standard of care: The Viennese registry on reperfusion strategies in ST-elevation myocardial infarction (Vienna STEMI registry). *Circulation* 2006; 113: 2398–2405.
 17. Bradley EH, Roumanis SA, Radford MJ, Webster TR, McNamara RL, Mattera JA, et al. Achieving door-to-balloon times that meet quality guidelines: How do successful hospitals do it? *J Am Coll Cardiol* 2005; 46: 1236–1241.
 18. Curtis JP, Portnay EL, Wang Y, McNamara RL, Herrin J, Bradley EH, et al. The pre-hospital electrocardiogram and time to reperfusion in patients with acute myocardial infarction, 2000–2002: Findings from the National Registry of Myocardial Infarction-4. *J Am Coll Cardiol* 2006; 47: 1544–1552.
 19. Afolabi BA, Novaro GM, Pinski SL, Fromkin KR, Bush HS. Use of the prehospital ECG improves door-to-balloon times in ST segment elevation myocardial infarction irrespective of time of day or day of week. *Emerg Med J* 2007; 24: 588–591.
 20. Assessment of the Safety and Efficacy of a New Treatment Strategy with Percutaneous Coronary Intervention (ASSENT-4 PCI) investigators. Primary vs tenecteplase-facilitated percutaneous coronary intervention in patients with ST-segment elevation acute myocardial infarction (ASSENT-4 PCI): Randomised trial. *Lancet* 2006; 367: 569–578.
 21. Ellis SG, Tendera M, de Belder MA, van Boven AJ, Widimsky P, Janssens L, et al. Facilitated PCI in patients with ST-elevation myocardial infarction. *N Engl J Med* 2008; 358: 2205–2217.
 22. Steg PG, Bonney E, Chabaud S, Lapostolle F, Dubien PY, Cristofini P, Prehospital Thrombolysis in acute Myocardial infarction (CAPTIM) Investigators. Impact of time to treatment on mortality after prehospital fibrinolysis or primary angioplasty: Data from the CAPTIM randomized clinical trial. *Circulation* 2003; 108: 2851–2856.
 23. Pinto DS, Kirtane AJ, Nallamothu BK, Murphy SA, Cohen DJ, Laham RJ, et al. Hospital delays in reperfusion for ST-elevation myocardial infarction: Implications when selecting a reperfusion strategy. *Circulation* 2006; 114: 2019–2025.
 24. Angeja BG, Gibson CM, Chin R, Frederick PD, Every NR, Ross AM, et al. Predictors of door-to-balloon delay in primary angioplasty. *Am J Cardiol* 2002; 89: 1156–1161.

8. 縦隔炎と感染予防

石橋和幸 山本文雄*

I. 基本的な考え方

近年、心臓血管外科領域においては手術手技の改良や人工材料の発達により、その手術成績は飛躍的に向上してきている。しかしながら現在においても、術後に遭遇する感染症は患者の予後を左右するもっとも重篤な合併症の一つである。心臓血管外科手術は本来無菌手術であるが、体外循環や多数の血管内留置カテーテル、さまざまな人工材料の体内留置などにより、術中・術後に体内に病原微生物が移入されやすく、また感染への抵抗性も低下して宿主と病原菌の力関係が逆転し、術後に感染症を発症するリスクは高い。心臓血管外科医にとって、この感染ならびに感染準備状態をいかにコントロールするかは、手術成績を左右する重大な問題となる。

本稿においては、心臓血管外科周術期における感染予防対策、術後に問題となる感染症対策および縦隔炎の診断と治療方針について述べる。

II. 心臓血管外科手術後の感染症予防対策

術後の感染には、患者自身が有する常在菌による感染（内因性感染）と、ICUや病室に存在する院内環境汚染菌による感染（外因性感染）がある。これらの術後感染を防止するためには、すべての体内への細菌の侵入をいかに防止するかにある。その対策として、これまでに Centers for Disease Control and Prevention (CDC) をはじめとするさまざまな「感染症対策ガイドライン」¹⁾

が示されているが、院内環境や感染対策状況は施設によってさまざまであり、通常の標準的感染予防対策（standard precaution）に加えて、各施設、各診療科に応じた「感染対策マニュアル」を作成し対応していくことが必要である（表1）。そこで、心臓血管外科手術後の感染症予防対策について、その特徴的対策としてわれわれの施設が行っている方法を中心に、術前・術中・術後について述べる。

① 予防的抗菌薬投与

心臓血管外科手術は無菌手術であるが、皮膚切開を必ず伴うため皮膚に常在する表皮ブドウ球菌や黄色ブドウ球菌による内因性感染を対象とし抗菌薬を投与する。一般的に第一世代セフェム系薬（cefazolin）やペニシリン系薬（ampicillinとsulbactamの合剤）などが用いられる。定期的なメチシリン耐性黄色ブドウ球菌（MRSA）prevalence rate 調査を行うことを原則とし、特にMRSA 保菌例や院内環境でMRSAが多いと判断される状況のときは、発症阻止のため vancomycin を使用する²⁾（健康保険非適用）。手術部位の細菌汚染が始まるのは皮膚切開時であり、そのときに抗菌薬が血中および組織中の殺菌的濃度に達していることが必要であり、皮膚切開の30分前に投与する。Cefazolinは半減期が約2時間であり、組織中の抗菌薬濃度を閉創まで十分に殺菌的な水準に維持するため、手術開始3~4時間後には追加投与が必要となる。半減期の長い vancomycin においても手術開始6時間後には追加投

キーワード：心臓血管外科手術、予防的抗菌薬、縦隔炎、リスクファクター

* K. Ishibashi (講師), F. Yamamoto (教授) : 秋田大学心臓血管外科。

表 1. 心臓血管外科手術後の感染症予防対策

1. 環境対策
standard precaution の徹底
感染対策マニュアルの作成
2. 術前対策
術前禁煙, 術前シャワー浴
手術直前にクリッパーによる除毛
術前監視培養(mupirocin 塗布, povidone-iodine 希釈液によるうがい)
3. 術中対策
maximal barrier precaution の徹底
手術時間の短縮
3時間ごとの手袋の交換
4. 術後対策
気管チューブ・血管カテーテル・尿道カテーテルの早期抜去
早期離床, 早期経口摂取開始, 厳格な血糖コントロール
適切な創部管理

与が必要である。投与期間については、これまでにさまざまな臨床研究がなされてきたが、米国胸部外科学会のガイドラインでは、耐性菌の出現も考慮し、心臓血管外科手術後 48 時間までの投与が推奨されている³⁾。

② 術前予防対策

術後の感染症を予防するために、入院時に菌の有無、喫煙、血糖管理状況など術前のリスクファクターについて評価する。MRSA の鼻腔内保菌が手術部位感染 (surgical site infection: SSI) のリスクファクターであることは多数報告されており⁴⁾、鼻腔・咽頭の監視培養を行う。監視培養で MRSA が検出された場合には、緊急例以外は mupirocin 塗布による除菌や povidone-iodine 希釈液によるうがいを行う。当施設では週 1 回の検査で連続 2 回の陰性を確認してから手術に臨んでいる。原則的に、手術前日には液体石鹸を用いシャワー浴を行う。術前の剃毛は原則的に行わず、必要な場合のみ手術当日、手術室にてハサミやクリッパーを用いて行う。

③ 術中予防対策

十分な手洗いの後でも 3 時間経過すると手指に表皮常在細菌が認められることが報告されており、手術時間の短縮を心がけ、約 3 時間ごとに速乾性手指消毒剤を使用し手袋の交換や、再度手洗いをするなど配慮が必要である。閉胸前に縦隔、心嚢内を生理食塩水で十分に洗浄し、落下細菌や血腫などを十分に除去する。術後感染症の原因菌の多くは術中・術後に体内に移入するので、外科医のみならず麻酔医、看護師、臨床工学技師などの無菌操作もきわめて重要である。手術室へ

の入室人数が増加すると浮遊粉塵が多くなり、病原微生物が混入しやすくなるため、手術室への入室を制限することも重要である。血管内用具留置時、静脈内投与薬剤の準備、および投与に際しては、無菌法の原則を順守する。特に中心静脈カテーテル挿入の際には、maximal barrier precaution に沿って帽子、マスクを着用し、手洗いの後、滅菌ガウンと滅菌手袋を着用して行う。すなわち術中においては、①手術室の環境対策、②血管確保時の無菌対策、③手術操作に関する感染対策が重要である。

④ 術後予防対策

感染起因菌の侵入部位は、創部のみならず血管ライン刺入部、ドレーン挿入部、気管チューブ、尿道カテーテルなどさまざまであり、それらのラインを早期に抜去することが重要である。また栄養管理を考慮し、早期離床、早期経口摂取開始を行う。一期的に閉鎖した切開創は、72 時間以内に接合面が接着する。そのため原則的に術後 3 日間、術中の滅菌ドレッシング剤で保護し、それ以降は開放とする。ただし創部より滲出液などがあった場合には、その症例ごとに対応する。包交の前後では必ず手洗いをを行い、手袋を着用し包交を行う。高血糖時には白血球の機能(遊走、接着、貪食、殺菌能)が障害されると考えられており、周術期において感染を生じやすい状態となるため、特に術後 48 時間以内は血糖を厳格にコントロール(血糖値を 200 mg/dl 以下)する⁵⁾。

すなわち術後は、①体内への挿入ライン(カテーテルやドレーン)および創部の管理、②術後の栄養、血糖管理が重要である。



図1. 術後感染症に対する抗菌薬投与方針

⑤ 術後感染症対策

予防的抗菌薬を終了後、発熱や炎症所見が遷延、再上昇した場合には、盲目的に広域スペクトラムの抗菌薬を用いるのではなく、感染部位、起原菌の同定に努め、感受性、感染部位に応じた抗菌薬を選択することが重要である(図1)。起原菌検索のため、病歴の間診と同時に必ず身体診察を行い、局所の圧痛、腫脹、痰、咽頭痛、歯痛、排尿痛、頭痛、下痢、関節痛など丹念に聴取しながら診察する。臨床検査では末梢血液検査、尿検査などのほかに、喀痰、尿、血液の細菌培養検査は必須である。それ以外に、局所に排膿があった場合には同部位の培養、または必要に応じて便培養を採取する。起原菌の特定がもっとも基本であるが、培養検査では菌の同定や感受性試験の結果までに時間がかかるため、Gram染色による鏡検が抗菌薬投与の手がかりとなる。鏡検によってGram陽性球菌による重症感染症が疑われた場合には、初期治療から vancomycin を投与する。起原菌と感受性が判明した時点において、より抗菌力が強く、毒性が低く、抗菌スペクトラムの狭い薬剤に変更する。抗菌薬を投与する場合には、pharmacokinetics (PK)/pharmacodynamics (PD) を考慮し、投与回数、投与量を十分に検討することが重要である。心臓血管外科手術後の感染症には、① SSI と② 術野外感染 (remote infection) があり、その双方への留意が必要である。

表2. 縦隔炎発症のリスクファクター

宿主要因	手術要因	術後要因
高齢 肥満 MRSA 保菌 糖尿病 慢性閉塞性肺障害 喫煙 術前心不全 腎不全 再手術 緊急手術	手術時間延長 術後出血 両側内胸動脈使用 輸血	長期 ICU 滞在 (長期挿管) 気管切開 術後肺炎

Ⅲ. 術後感染性心内膜炎

心臓血管外科手術後に感染性心内膜炎を合併する危険性は2%以下⁶⁾と報告されているが、一度罹患した場合には敗血症を伴い、死亡率が約20%と重篤な合併症である。特に人工弁置換術後の心内膜炎では、予後は不良である。原因菌としては連鎖球菌、ブドウ球菌、腸球菌が多く、菌種に応じた抗菌薬の投与が重要である。原因菌が同定できない場合においても、ガイドラインに従い抗菌薬を投与する⁷⁾。診断は、心臓超音波検査で vegetation を認め、血液培養で菌血症を証明する。内科治療に抵抗性で抗菌薬の投与でも炎症反応が改善しない場合、vegetation が増大傾向にあるときや心不全がコントロールできない場合には外科治療が必要となる。

Ⅳ. 縦 隔 炎

① リスクファクターと診断

縦隔炎は、現在においても心臓血管外科手術後の0.5~3.3%に発症し、その死亡率も5.4~23%と高く⁸⁻¹²⁾、術後の SSI の中でもっとも重篤な感染症の一つである。宿主側の要因としては高齢、肥満、MRSA 保菌、糖尿病、慢性閉塞性肺障害、喫煙、腎不全、再手術、緊急手術などであり、手術要因としては手術時間、術後出血、両側内胸動脈使用、輸血であり、術後要因としては長期 ICU 滞在(長期挿管)、気管切開などがあげられている(表2)。

手術後に発熱や WBC 数や CRP の上昇に加え、胸骨の痛みや発赤および動揺、創部やドレーンから膿汁が認められた場合には縦隔炎を疑う。診断

には局所の細菌培養検査のほか、胸部CTが有用である。胸部CTで縦隔内に air bubble や ring enhance される液体の貯留、胸骨離開や骨髄炎像が認められた場合には縦隔炎と診断する。臨床症状が乏しくCTなどで診断が困難な場合には、FDG-PETによって胸骨や胸骨裏面のブドウ糖の異常消費を調べることも、診断の補助手段として有用である。起病菌は表皮ブドウ球菌と黄色ブドウ球菌が約40~70%と多く、その半数がMRSAと報告されている^{6,8-12)}。

② 縦隔炎の治療

治療には、抗菌薬に加えて外科的処置が必要になる。胸骨の破壊が軽度の場合や膿瘍が限局している場合には、局所のデブリドマン、生理食塩水による洗浄およびSBバック（住友ベークライト社、東京）などによる持続陰圧吸引で治療する。消毒薬は創傷治癒に働く細胞を傷害し創の治癒を遷延させるため、局所の消毒薬での洗浄や抗菌薬の塗布は行わない。炎症が強く胸骨が大きく破壊されている場合には、これまでに抗菌薬や消毒薬による洗浄の後、二次的な胸骨閉鎖や抗菌薬による閉鎖式縦隔洗浄、大網や大胸筋充填などが行われてきたが、最近では胸骨を開放しスポンジとSBバックなどを用いた陰圧閉鎖式持続吸引ドレナージ（vacuum assisted closure : VAC）が行われるようになり、治癒率をより向上させている。またVAC療法と併用し、死腔に大網や大胸筋などを充填することや、胸骨閉鎖、有茎筋皮弁なども有効であると報告されている¹³⁻¹⁵⁾。胸骨の破壊や炎症の程度は症例によってさまざまであるため、十分なデブリドマンの後、①胸骨閉鎖、②VAC、③大網充填などを組み合わせて対応していくことが肝要である。

V. 結 論

心臓血管外科手術後の感染を予防するためには、以下のことを日々実践していくことが肝要である。

1) ガイドラインに沿って各施設、各診療科ごとの「感染対策マニュアル」を作成し、感染予防に努める。

2) 予防的抗菌薬はcefazolinまたはampicillinとsulbactamの合剤を用い、手術開始前に投与し術後48時間で終了とする。

3) 術後感染症が疑われた場合には、必ず各種細菌培養検査を行ってから抗菌薬を投与する。

4) 縦隔炎が疑われた場合にはただちにCTを撮影し、外科的治療を開始する。症状に応じてvacuum-assisted systemを用いた持続陰圧吸引療法を行う。

文 献

- 1) Mangram AJ, Horan TC, Pearson ML et al : Guideline for prevention of surgical site infection, 1999 : Centers for Disease Control and Prevention (CDC) Hospital Infection Control Practices Advisory Committee. *Am J Infect Control* 27 : 97-132, 1999
- 2) Engelman R, Shahian D, Shemin R et al : The Society of Thoracic Surgeons practice guideline series : antibiotic prophylaxis in cardiac surgery ; part II. antibiotic choice. *Ann Thorac Surg* 83 : 1569-1576, 2007
- 3) Edwards FH, Engelman RM, Houck P et al : The Society of Thoracic Surgeons practice guideline series : antibiotic prophylaxis in cardiac surgery ; part I. duration. *Ann Thorac Surg* 81 : 397-404, 2006
- 4) Shrestha NK, Banbury MK, Weber M et al : Safety of targeted perioperative mupirocin treatment for preventing infections after cardiac surgery. *Ann Thorac Surg* 81 : 2183-2188, 2006
- 5) Zerr KJ, Furnary AP, Grunkemeier GL et al : Glucose control lowers the risk of wound infection in diabetics after open heart operations. *Ann Thorac Surg* 63 : 356-361, 1997
- 6) Saadatian-Elahi M, Teyssou R, Vanhems P : *Staphylococcus aureus*, the major pathogen in orthopaedic and cardiac surgical site infections : a literature review. *Int J Surg* 22 : 405-409, 2007
- 7) 宮武邦夫, 北村惣一郎, 中澤 誠ほか : 感染性心内膜炎の予防と治療に関するガイドライン. *Circ J* 67 [Suppl] : 1039-1082, 2003
- 8) Tang GH, Maganti M, Weisel RD et al : Prevention and management of deep sternal wound infection. *Semin Thorac Cardiovasc Surg* 16 : 62-69, 2004
- 9) Abboud CS, Wey SB, Balter VT : Risk factors for mediastinitis after cardiac surgery. *Ann Thorac Surg* 77 : 676-683, 2004
- 10) Crabtree TD, Codd JE, Fraser VJ et al : Multivariate analysis of risk factors for deep and

superficial sternal infection after coronary artery bypass grafting at a tertiary care medical center. *Semin Thorac Cardiovasc Surg* 16 : 53-56, 2004

- 11) Fowler Jr VG, O'Brien SM, Muhlbauer LH et al : Clinical predictors of major infections after cardiac surgery. *Circulation* 112 [Suppl] : I358-I365, 2005
- 12) Ridderstolpe L, Gill H, Granfeldt H et al : Superficial and deep sternal wound complications : incidence, risk factors and mortality. *Eur J Cardiothorac Surg* 20 : 1168-1175, 2001
- 13) Immer FF, Durrer M, Mühlemann KS et al : Deep sternal wound infection after cardiac

surgery : modality of treatment and outcome. *Ann Thorac Surg* 80 : 957-961, 2005

- 14) Fleck TM, Fleck M, Moidl R et al : The vacuum-assisted closure system for the treatment of deep sternal wound infections after cardiac surgery. *Ann Thorac Surg* 74 : 1596-1600, 2002
- 15) Fuchs U, Zittermann A, Stuetzgen B et al : Clinical outcome of patients with deep sternal wound infection managed by vacuum-assisted closure compared to conventional therapy with open packing : a retrospective analysis. *Ann Thorac Surg* 79 : 526-531, 2005

SUMMARY

Management of Mediastinitis and Preventions of Perioperative Nosocomial Infection after Cardiovascular Surgery

Kazuyuki Ishibashi et al., Department of Cardiovascular Surgery, Akita University, Akita, Japan

This article reviews risk factors and treatment of perioperative nosocomial infections. The primary prophylactic antibiotic is recommended to be a cefazolin or penicillin with sulbactam, because the most frequent organism cultured in cardiac surgical site infection (SSI) is *Staphylococcus*. Antibiotic prophylaxis of 48 hours' duration after cardiovascular surgery is clinically effective in minimizing infectious complication. In patients considered at high risk for a staphylococcal infection, vancomycin may be recommended. In the treatment of postoperative infections, Gram-stain-based antibiotic selection is necessary and the initial empirical therapy to ensure adequate coverage of potentially infective organisms should be accompanied by de-escalation until microbiological data become available. Mediastinitis, which is one of the important infectious complications after cardiovascular surgery, requires surgical drainage and debridement, as well as antimicrobial therapy. Vacuum-assisted closure (VAC) is an effective therapy compared with the conventional technique of open packing. Continuous clinician's efforts and prolonged infection control programs are very important for prevention of perioperative nosocomial infection after cardiac surgery.

KEY WORDS

cardiovascular surgery/antibiotic prophylaxis/mediastinitis/risk factor

胸部外科

2007年5月号 (Vol.60 No.5)

定価2,625円(税込)

南江堂 TEL.03-3811-7239 (営業)

700号記念

特集〈胸部外科10年の進歩と将来展望〉

- 特集「胸部外科10年の進歩と将来展望」によせて
.....幕内晴朗/近藤丘
- 癌の外科治療における分子生物学
- 心臓血管外科における再生医療と分子生物学
- 人工心臓
- 同種移植と異種移植

- 胸部外科の診療システムの課題と展望
—日本胸部外科学会からのメッセージ
- 胸部外科の魅力ある卒前・卒後教育
- 胸部外科と専門医制度—心臓血管外科
- 胸部外科医と専門医制度—呼吸器外科
- 胸部外科における国際交流

未破裂の心外型 Valsalva 洞動脈瘤

相田 弘秋 加賀谷 聡 山本文雄 山浦 玄武*

はじめに

Valsalva 洞動脈瘤はまれな疾患であり、その未破裂例はさらにまれである。未破裂例は無症状で偶然発見されるために、その治療方針は議論のあるところである。今回、未破裂の心外型 Valsalva 洞動脈瘤を経験したので報告する。

I. 症 例

症 例 49 歳，女。

主 訴：労作時息切れ。

家族歴：特記すべきことはない。

既往歴：36 歳時に子宮腺筋症，慢性閉塞性肺疾患。

現病歴：2005 年 1 月ごろから労作時息切れを自覚し，当院内科を受診した。慢性閉塞性肺疾患として外来治療していたが，2007 年 9 月にたまたま施行した心臓超音波検査で Valsalva 洞動脈瘤を指摘され，手術目的にて当科へ紹介となった。

現 症：身長 160 cm，体重 41.2 kg，体温 36.8℃，血圧 136/76 mmHg，脈拍 60 回/分・整であった。心雑音はなかった。漏斗胸と両下肢静脈瘤を認めた。

血液・生化学所見：血液一般，生化学，血清検査はすべて正常で，炎症反応も陰性であった。




図 1. 術前左室造影像
無冠状動脈洞の瘤化を認める。

胸部 X 線所見：心胸郭比 45.7%，肺気腫様変化を認めた。

呼吸機能所見：1 秒率 48.2%，%肺活量 92%であった。

心電図所見：洞調律，不完全右脚ブロックを認めた。

心臓超音波所見：無冠状動脈洞に 3.5×4.2 cm の動脈瘤を認めた。左室駆出率は 77%と良好で、

キーワード：Valsalva 洞動脈瘤，心外型動脈瘤，パッチ閉鎖

* H. Aida (部長)，S. Kagaya (科長)：厚生連平鹿総合病院心臓血管外科 (〒013-8610 横手市前郷字八ツ口 3-1)
F. Yamamoto (教授)，G. Yamaura：秋田大学心臓血管外科。



図2. 術中所見

無冠状動脈洞に20×15 mmの瘤口があり、その右側に瘤を認める。



図3. 術後CT

無冠状動脈洞動脈瘤の血栓化(矢印)を認める。

大動脈弁閉鎖不全を認めなかった。

心カテーテル所見: 無冠状動脈洞から生じた巨大な Valsalva 洞動脈瘤を認めた(図1)。

以上から無症状であるものの、巨大瘤破裂の危険性を考慮して手術適応とした。

手術所見: 胸骨正中切開にて開胸した。Valsalva 洞動脈瘤は心内腔と接しない心外型であった。上行大動脈送血, 上下大静脈脱血にて人工心肺を確立した。上行大動脈を遮断し, 右上肺静脈ベントを行いながら心筋保護液を順行性および逆行性に注入して心停止を得た。上行大動脈を横切すると無冠状動脈洞に20×15 mmの瘤口を認めた(図2)。ウマ心膜パッチ(XAG-400: Edwards Lifesciences社, Irvine)を用い, 5-0プロリン糸の連続縫合で瘤口を閉鎖した。人工心肺からの離脱は容易であった。

術後経過: 術後は順調に経過し, CTで瘤の血栓閉塞を確認した(図3)。

II. 考 察

Valsalva 洞動脈瘤は井上¹⁾の分類が一般的に用いられ, 心外型と心内型に分類される。心外型は通常1個の Valsalva 洞から発生した囊状動脈瘤であり, 心外に発育してしばしば巨大化するが, 右心房を外から圧迫することはあっても右心房内に突出することはない。一方, 心内型は瘤口を有する瘤が心腔内に突出するが, 通常みられる破裂性の瘤では小指頭ないし母指頭大で, wind-sock, balloon type と表現されるように吹き流し

状あるいは指サック状を呈し, その先端が穿孔することが特徴とされる¹⁾。

Valsalva 洞動脈瘤の原因は先天性と後天性に分けられる。先天性としては胎生期の円錐部隆起および心内膜床の融合不全や Marfan 症候群などの中膜変性によるものが考えられ, 後天性としては外傷, 大動脈炎症候群, 梅毒, 結核などによるものが考えられる^{2,3)}。心外型の原因は大動脈壁と大動脈弁輪間の中膜の連続性が欠如していることであり⁴⁾。心内型はこれに中隔筋性組織の欠損が加わるとする説⁵⁾があり, 合理的な説明であると考えられる。

本稿では特に心外型について述べることにする。本邦ではこれまでに文献上20例の心外型 Valsalva 洞動脈瘤が報告されている(表1)。平均年齢51.5(24~69)歳, 男女比は10:10と同等であった。主訴は合併する心疾患によるものが多く, 14例(70%)に大動脈弁閉鎖不全症, 8例(40%)に冠状動脈病変, このほかに右室流出路狭窄2例, 上行大動脈瘤2例を認めた。瘤の発生部位は左冠状動脈洞5例, 右冠状動脈洞7例, 無冠状動脈洞6例とほぼ同等の比率であったが, すべての冠状動脈洞に認めたものが1例, 右~無冠状動脈洞に発生したものが1例あった。

表1. 本邦における未破裂の心外型 Valsalva 洞動脈瘤の報告例

症例名	直径(mm)	瘤の部位	合併心疾患	手術
1988)	60	L	AR, LMT 圧迫	パッチ閉鎖, AVR
1987)	100	N	上行瘤	パッチ閉鎖 & 上行置換術
1989)	100	R	AR, RCA 閉塞	パッチ閉鎖
1989)	80	L	AR, LMT 閉塞	パッチ閉鎖, AVR, CABG
加藤 (1993)	30	R	AR, 右室流出路狭窄, 狭心症	パッチ閉鎖, AVR, CABG
Okita (1995)	70	R	RCA 圧迫	パッチ閉鎖, RCA 再移植
浜田 (1995)	70	N	AR	パッチ閉鎖
水島 ⁷⁾ (1996)	140	R	なし	パッチ閉鎖, 瘤縮小
本多 (1997)	60	R-N	AR	パッチ閉鎖, AVR
Yasuda ⁸⁾ (2000)	70	L	AR, RCA 拡張症	大動脈基部置換術
	50	N	AR, 上行瘤	パッチ閉鎖, 上行置換術
下山 (2001)	80	R	AR, RCA 閉塞	パッチ閉鎖, AVR, CABG
Kusuyama ⁹⁾ (2002)	15	R	AR	パッチ閉鎖
佐藤 (2002)	35	N	なし	パッチ閉鎖, 瘤切除
横山 (2003)	50	L	RCA 瘤	パッチ閉鎖, CABG
内川 ⁶⁾ (2003)	35	N	AR	パッチ閉鎖, AVR→出血: 上行置換術
近江 (2004)	66	L	AR	大動脈基部置換術
石神 (2004)	70	R	AR, 右室流出路狭窄	パッチ閉鎖, AVR, CABG
Akashi (2005)	60, 42, 20	N, L, R	AR	大動脈基部置換術
木村 ¹⁰⁾ (2006)	52	N	なし	大動脈基部置換術

L: 左冠状動脈洞, R: 右冠状動脈洞, N: 無冠状動脈洞, AR: 大動脈弁閉鎖不全, RCA: 右冠状動脈, LMT: 左冠状動脈主幹部, AVR: 大動脈弁置換, CABG: 冠動脈バイパス術

瘤径は15~140 mm とさまざまであった。手術術式は16例(80%)でパッチ閉鎖が行われており、これに合併心疾患に対するAVRやCABGが追加されていた。パッチ使用例の中には閉鎖部位からの出血のため上行置換へ変更した症例もあった⁶⁾。パッチ閉鎖が困難な4例に対しては大動脈基部置換術が行われた。

病理組織学的に動脈瘤壁が特異的所見を示したのは5例あったが(Ehlers-Danlos症候群, 好酸球増多症候群, Marfan症候群, Behçet病, 大動脈炎症候群)⁷⁻¹⁰⁾。動脈瘤様変化の所見が8例と多く、ほかに粘液腫様変性が2例、記載なしが5例であった。

手術術式はパッチ閉鎖術が標準と思われるが、合併心疾患やValsalva洞動脈瘤の性状, 上行大動脈の性状を考慮して決定すべきであろう。手術適応については議論のあるところである。本症はまれな疾患であり、動脈硬化性動脈瘤のような瘤径に基づいた手術適応基準はない。本症の原因が大動脈壁と大動脈弁輪の中膜の連続性の欠如であることから、瘤破裂のリスクは高いと考えられる。本症の報告例中、Behçet病に伴ったValsalva洞動脈瘤では瘤径15 mmで手術が施行されて

いた⁹⁾。合併疾患を考慮しながら、発見後早期の手術が望ましいと考えられた。

本例は大動脈壁や瘤壁を採取していないため病理組織学的検討ができず、Valsalva洞動脈瘤の原因確定にはいたらなかった。

おわりに

未破裂の心外型 Valsalva 洞動脈瘤の1例に対して瘤口パッチ閉鎖術を行い、良好な結果を得た。

文 献

- 1) 井上 正: 非破裂性心内型バルサルバ洞動脈瘤. 心臓 29: 415-417, 1997
- 2) 榎原 任, 今野草二: 先天性 Valsalva 洞動脈瘤. 胸部外科 21: 26-30, 1968
- 3) Kirklin JW, Barratt-Boyes BE: Congenital sinus of Valsalva aneurysm and aortico-left ventricular tunnel. Cardiac Surgery, 3rd Ed, Churchill Livingstone, New York, p911-927, 2003
- 4) Edwards JE, Burchell HB: Specimen exhibiting the essential lesion in aneurysm of the aortic sinus. Proc Staff Meet Mayo Clinic 31:

- 407-421, 1956
- 5) 正津 晃, 竹内成之, 井上 正ほか: 特異的な心内異常を伴う心室中隔欠損症—Valsalva 洞動脈瘤その他. 胸部外科 25: 465-475, 1972
 - 6) 内川 伸, 伊藤良明, 林 和秀ほか: 非破裂心外型 Valsalva 洞動脈瘤の経験. 胸部外科 56: 879-882, 2003
 - 7) 水島恒和, 高橋俊樹, 島崎靖之ほか: 心外型非破裂性巨大 Valsalva 洞動脈瘤—1 手術例及び文献的考察. 日胸外会誌 44: 202-207, 1996
 - 8) Yasuda F, Shimono T, Adachi K et al: Surgical repair of extracardiac unruptured acquired Valsalva aneurysms. Ann Thorac Surg 70: 1696-1698, 2000
 - 9) Kusuyama T, Nakamura Y, Yamagishi H et al: Unruptured aneurysm of the sinus of Valsalva with Behçet's disease. Circ J 66: 107-108, 2002
 - 10) 木村知恵理, 小宮達彦, 田村暢成ほか: 大動脈炎による未破裂 Valsalva 洞動脈瘤に対し大動脈基部再建術を施行した 1 例. 日心臓血管外会誌 35: 271-274, 2006

SUMMARY

Surgical Repair of Extracardiac Unruptured Aneurysm of the Sinus of Valsalva

Hiroaki Aida et al., Department of Cardiovascular Surgery, Hiraka General Hospital, Yokote, Japan

An isolated extracardiac unruptured aneurysm of the non-coronary sinus of Valsalva was detected incidentally in a 49-year-old woman with chronic obstructive pulmonary disease.

The orifice of the aneurysm was closed with XAG-400 patch. The aortic valve was normal. 20 cases of successful operation have been reported in Japan with extracardiac unruptured aneurysm. In the review of the literature, patch closure of the orifice of the aneurysm was the standard procedure in most cases. However, aortic root replacement was required in some cases with concomitant cardiac and/or aortic disease.

KEY WORDS

aneurysm of sinus of Valsalva/extracardiac aneurysm/patch closure

*

*

*

胸部外科

2006年4月号 (Vol.59 No.4)

定価2,520円 (税込)

nhp 南江堂

TEL.03-3611-7239 (夜間)

特集〈狭小大動脈弁輪に対する弁置換術〉

■特集「狭小大動脈弁輪に対する弁置換術」
によせて……………幕内晴朗

1 弁輪拡大術

Patient-prosthesis mismatchの回避/狭小大動脈弁輪に対する人工弁置換術/高齢者に対する大動脈弁輪拡大術

2 術式の工夫

高齢者狭小大動脈弁輪に対する弁置換術/狭小弁輪を伴う大動脈弁狭窄に対する弁輪上拡大の有用性/高齢者狭小大動脈弁輪に対するapicoaortic conduitの成績と問題点/石灰化狭小大動脈弁に対する弁置換術

3 小口径人工弁

狭小弁輪に対するconventional大動脈弁置換術の遠隔成績/高齢者に対する19mm人工弁

を用いた大動脈弁置換術

4 心機能

大動脈弁置換術後のpatient-prosthesis mismatch/左室拡張機能障害からみた大動脈弁手術

5 高齢者

高齢者の弁置換術において有効弁口面積指数0.85超は必要か/高齢者高度石灰化狭小大動脈弁輪に対する弁置換術