

Table 3. Age- and Sex-Adjusted Mean Values of Heart Rate–Corrected QT Interval Duration According to Brachial-Ankle Pulse Wave Velocity Quartiles and Relevant Factors

	Quartiles of brachial-ankle pulse wave velocity				<i>p</i> for trend	<i>p</i> for homogeneity
	Q1	Q2	Q3	Q4		
Hypertension						
No (<i>n</i> =1,618)	402.4±0.9	405.8±0.9	408.8±1.3	410.8±1.9	<0.0001	0.43
Yes (<i>n</i> =1,048)	402.8±3.9	408.8±2.0	409.7±1.3	412.8±1.2	0.01	
Dyslipidemia						
No (<i>n</i> =1,202)	403.1±1.4	405.9±1.3	410.0±1.4	408.6±1.8	0.03	0.14
Yes (<i>n</i> =1,464)	401.3±1.4	407.0±1.2	408.8±1.1	414.5±1.2	<0.0001	
Diabetes						
No (<i>n</i> =2,243)	402.4±1.0	406.4±0.9	408.7±0.9	411.9±1.2	<0.0001	0.39
Yes (<i>n</i> =423)	404.1±3.8	407.6±2.7	411.7±2.2	412.9±1.9	0.06	
Obesity						
No (<i>n</i> =1,938)	402.7±1.1	406.4±1.0	408.7±1.0	411.2±1.2	<0.0001	0.19
Yes (<i>n</i> =728)	401.7±2.0	406.9±1.7	410.5±1.6	414.8±1.8	<0.0001	
ECG abnormalities						
No (<i>n</i> =2,196)	402.2±1.0	405.8±0.9	409.1±0.9	411.8±1.2	<0.0001	0.43
Yes (<i>n</i> =470)	402.3±3.7	411.1±2.9	410.6±2.3	413.7±2.1	0.04	
Alcohol intake						
No (<i>n</i> =1,504)	404.3±1.4	408.1±1.2	410.6±1.2	413.6±1.3	<0.0001	0.34
Yes (<i>n</i> =1,162)	399.8±1.4	404.4±1.3	407.6±1.3	410.9±1.6	<0.0001	
Habitual smoking						
No (<i>n</i> =2,068)	403.9±1.2	407.2±1.0	409.7±1.0	413.7±1.1	<0.0001	0.21
Yes (<i>n</i> =598)	397.2±1.6	404.1±1.7	408.1±1.8	407.5±2.3	0.0003	

Values are age- and sex-adjusted means±SEM.

Table 4. Age- and Sex-Adjusted Mean Values of Heart Rate–Corrected QT Interval Duration According to Brachial-Ankle Pulse Wave Velocity Quartiles and the Number of Relevant Factors

Number of relevant factors	Quartiles of brachial-ankle pulse wave velocity				<i>p</i> for trend
	Q1	Q2	Q3	Q4	
0–1 (<i>n</i> =903)	406.0±1.6	404.0±1.4	408.1±1.4	411.6±1.7	0.001
2–3 (<i>n</i> =1,313)	402.5±1.4	404.8±1.3	409.4±1.3	410.4±1.5	0.003
4–7 (<i>n</i> =450)	407.4±2.3	407.7±2.2	413.9±2.2	413.5±2.4	0.02

Values are age- and sex-adjusted means±SEM. Relevant factors: hypertension, dyslipidemia, diabetes, obesity, ECG abnormalities, alcohol intake, and habitual smoking.

(*p*<0.0001 for trend in both sexes). In the following analyses, male and female subjects were combined because the relationships of baPWV to QTc were comparable between men and women.

Table 3 shows the age- and sex-adjusted mean values of QTc according to quartiles of the baPWV levels for subgroups of participants defined on the basis of the presence or absence of hypertension, dyslipidemia, diabetes, obesity, ECG abnormalities, alcohol intake, or smoking habits. There were comparable relationships between baPWV and QTc for participants who were and were not hypertensive. Likewise, there were no interactions in the relationships of baPWV with QTc between subgroups defined by every other relevant factor (all *p* values for interaction >0.05). There were also com-

parable relationships of baPWV with QTc between participants who were and were not taking antihypertensive agents or lipid-lowering agents (*p* for interaction >0.5). We also estimated the age- and sex-adjusted mean values of QTc according to quartiles of the baPWV levels by the number of relevant factors (Table 4). There was a significantly positive relationship between baPWV and QTc in each of the groups defined by a number of cardiovascular risk factors of 0–1, 2–3, and 4–7.

Figure 2 shows the multivariate-adjusted mean values of QTc according to quartiles of the baPWV levels. The multivariate-adjusted mean values of QTc significantly increased with rising baPWV levels, even after controlling for age, sex, hypertension, ECG abnormalities, dyslipidemia, diabetes,




Fig. 2. Multivariate-adjusted mean values of heart rate-corrected QT interval duration according to quartiles of brachial-ankle pulse wave velocity levels. The centers of the boxes are placed at the estimates of mean values. Other conventions are the same as in Fig. 1. Mean values and *p* values are adjusted for age, sex, hypertension, ECG abnormalities, dyslipidemia, diabetes mellitus, obesity, serum calcium and potassium, alcohol intake, and smoking habits. **p* < 0.01 vs. the lowest quartile group. *p* < 0.0001 for trend.

obesity, serum calcium, serum potassium, alcohol intake, and smoking habits (*p* < 0.0001 for trend).

Discussion

To our knowledge, this is the first study to address the associations between baPWV and QTc in a general population without preexisting cardiovascular disease. In the present analysis, the mean values of QTc increased with rising baPWV levels for both men and women. These associations remained strong and continuous, even after controlling for traditional cardiovascular risk factors, suggesting an independent relationship between subclinical arterial disease (atherosclerosis) and QT interval prolongation.

In the present study, there were strong and continuous relationships between QTc and baPWV, which has been shown to be a functional marker for subclinical atherosclerotic disease in central and peripheral arteries (16, 21). Ours is the largest study to have investigated the association between subclinical arterial disease and QT interval prolongation, but there have been a few other cross-sectional studies addressing this question using other structural markers of subclinical arterial disease (10–12). The Insulin Resistance Atherosclerosis Study (IRAS) investigated the association between carotid intima media thickness and QTc in 912 nondiabetic subjects without coronary artery disease and found a close association between carotid atherosclerosis and QT interval prolongation (10). The Salzburg Atherosclerosis Prevention Program in Subjects at High Individual Risk also showed a positive correlation

between carotid intima media thickness and QT interval duration in 1,199 clinically healthy subjects (11). These observational data support our hypothesis that subclinical arterial disease is associated with QT interval prolongation.

It is well known that the QT interval is affected by heart rate (18, 22). In order to control for the confounding effects of heart rate, we used QTc, which was estimated by Bazett's formula, and found significant associations between baPWV and QTc. When the Friedrich formula was used for estimation of QTc instead of Bazett's formula, similar associations were observed. We also investigated the association between baPWV and crude QT interval duration and found significantly positive relationships even after adjustment for heart rate, ECG abnormalities, and other cardiovascular risk factors (data not shown). These results suggest that baPWV is significantly associated with QT interval duration and this association is independent of the effects of heart rate.

The mechanism underlying the association between subclinical arterial disease and the acquired form of QT interval prolongation has not been clearly defined. Subclinical arterial disease and subsequent arterial stiffness may increase ventricular load and, as a consequence, may promote myocardial and electrophysiological remodeling, resulting in QT interval prolongation (23, 24). Another possible mechanism is that microvascular atherosclerosis in the coronary artery, which is strongly related to systemic arterial disease, may lead to sub-endocardial ischemia and thus extend QT interval duration (25).

One limitation of our study is that we have no information on subjects with congenital long QT syndrome. However, the prevalence of the congenital long QT syndrome has been reported to be less than 0.1% (26). Furthermore, in our subjects the relationship between baPWV and QTc was strong and continuous, even after excluding participants with QT intervals of 440 ms or more (*p* < 0.0001 for trend). Thus, the influence of congenital long QT syndrome would seem to have been negligible. Another limitation is that information on repeated measurements of baPWV and QTc is limited. This fact made it difficult for us to conduct longitudinal analysis.

In conclusion, we found close associations between baPWV and QTc for men and women without histories of cardiovascular disease. These associations were independent of hypertension, ECG abnormalities, dyslipidemia, diabetes, obesity, alcohol intake, and smoking habits. Thus, subclinical arterial disease appears to contribute to the pathogenesis of QT interval prolongation. Future longitudinal studies are necessary to clarify the causal relationship between subclinical arterial disease and QT interval prolongation.

Acknowledgements

The authors thank the residents of Hisayama for their participation in the survey and the staff of the Division of Health and Welfare of Hisayama for their cooperation in this study.

References

- Lepeschkin E, Surawicz B: The measurement of the Q-T interval of the electrocardiogram. *Circulation* 1952; **6**: 378–388.
- Al-Khatib SM, LaPointe NM, Kramer JM, Califf RM: What clinicians should know about the QT interval. *JAMA* 2003; **289**: 2120–2127.
- Ewing DJ, Boland O, Neilson JM, Cho CG, Clarke BF: Autonomic neuropathy, QT interval lengthening, and unexpected deaths in male diabetic patients. *Diabetologia* 1991; **34**: 182–185.
- Tomassoni G, Pisano E, Gardner L, Krucoff MW, Natale A: QT prolongation and dispersion in myocardial ischemia and infarction. *J Electrocardiol* 1998; **30** (Suppl): 187–190.
- Dekker JM, Schouten EG, Klootwijk P, Pool J, Kromhout D: Association between QT interval and coronary heart disease in middle-aged and elderly men. The Zutphen Study. *Circulation* 1994; **90**: 779–785.
- Dekker JM, Crow RS, Hannan PJ, Schouten EG, Folsom AR: Heart rate–corrected QT interval prolongation predicts risk of coronary heart disease in black and white middle-aged men and women: the ARIC study. *J Am Coll Cardiol* 2004; **43**: 565–571.
- de Bruyne MC, Hoes AW, Kors JA, Hofman A, van Bemmel JH, Grobbee DE: Prolonged QT interval predicts cardiac and all-cause mortality in the elderly. The Rotterdam Study. *Eur Heart J* 1999; **20**: 278–284.
- Robbins J, Nelson JC, Rautaharju PM, Gottdiener JS: The association between the length of the QT interval and mortality in the Cardiovascular Health Study. *Am J Med* 2003; **115**: 689–694.
- Okin PM, Devereux RB, Howard BV, Fabsitz RR, Lee ET, Welty TK: Assessment of QT interval and QT dispersion for prediction of all-cause and cardiovascular mortality in American Indians: the Strong Heart Study. *Circulation* 2000; **101**: 61–66.
- Strohmer B, Pichler M, Iglseider B, Paulweber B: Relationship of QT interval duration with carotid intima media thickness in a clinically healthy population undergoing cardiovascular risk screening. *J Intern Med* 2005; **257**: 238–246.
- Festa A, D'Agostino R Jr, Rautaharju P, et al: QT interval a marker of subclinical atherosclerosis in nondiabetic subjects? The Insulin Resistance Atherosclerosis Study (IRAS). *Stroke* 1999; **30**: 1566–1571.
- Takebayashi K, Aso Y, Matsutomo R, Wakabayashi S, Inukai T: Association between the corrected QT intervals and combined intimal-medial thickness of the carotid artery in patients with type 2 diabetes. *Metabolism* 2004; **53**: 1152–1157.
- Nichols WW: Clinical measurement of arterial stiffness obtained from noninvasive pressure waveforms. *Am J Hypertens* 2005; **18**: 3S–10S.
- Sawabe M, Takahashi R, Matsushita S, et al: Aortic pulse wave velocity and the degree of atherosclerosis in the elderly: a pathological study based on 304 autopsy cases. *Atherosclerosis* 2005; **179**: 345–351.
- Sugawara J, Hayashi K, Yokoi T, et al: Brachial-ankle pulse wave velocity: an index of central arterial stiffness? *J Hum Hypertens* 2005; **19**: 401–406.
- Yamashina A, Tomiyama H, Takeda K, et al: Validity, reproducibility, and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. *Hypertens Res* 2002; **25**: 359–364.
- Arima H, Tanizaki Y, Kiyohara Y, et al: Validity of the JNC VI recommendations for the management of hypertension in a general population of Japanese elderly: the Hisayama study. *Arch Intern Med* 2003; **163**: 361–366.
- Bazett HC: An analysis of time relations of the electrocardiogram. *Heart* 1920; **7**: 353–370.
- Kobayashi K, Akishita M, Yu W, Hashimoto M, Ohni M, Toba K: Interrelationship between non-invasive measurements of atherosclerosis: flow-mediated dilation of brachial artery, carotid intima-media thickness and pulse wave velocity. *Atherosclerosis* 2004; **173**: 13–18.
- The Expert Committee on the Diagnosis and Classification: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 1997; **20**: 1183–1197.
- Tomiyama H, Yamashina A, Arai T, et al: Influences of age and gender on results of noninvasive brachial-ankle pulse wave velocity measurement—a survey of 12517 subjects. *Atherosclerosis* 2003; **166**: 303–309.
- Oikarinen L, Nieminen MS, Viitasalo M, et al: QRS duration and QT interval predict mortality in hypertensive patients with left ventricular hypertrophy: the Losartan Intervention for Endpoint Reduction in Hypertension Study. *Hypertension* 2004; **43**: 1029–1034.
- Westerhof N, O'Rourke MF: Haemodynamic basis for the development of left ventricular failure in systolic hypertension and for its logical therapy. *J Hypertens* 1995; **13**: 943–952.
- Watabe D, Hashimoto J, Hatanaka R, et al: Electrocardiographic left ventricular hypertrophy and arterial stiffness: the Ohasama study. *Am J Hypertens* 2006; **19**: 1199–1205.
- Watanabe H, Ohtsuka S, Kakihana M, Sugishita Y: Coronary circulation in dogs with an experimental decrease in aortic compliance. *J Am Coll Cardiol* 1993; **21**: 1497–1506.
- Skinner J: Guidelines for the Diagnosis and Management of Familial Long QT Syndrome. *Heart Lung Circ* 2007; **16**: 22–24.

Secular Trends in the Incidence of and Risk Factors for Ischemic Stroke and Its Subtypes in Japanese Population

Michiaki Kubo, MD, PhD*; Jun Hata, MD, PhD*; Yasufumi Doi, MD, PhD;
Yumihiro Tanizaki, MD, PhD; Mitsuo Iida, MD, PhD; Yutaka Kiyohara, MD, PhD

Background—The study of long-term trends in the incidence of and risk factors for ischemic stroke subtypes could offer insights into primary and secondary prevention.

Methods and Results—We established 3 cohorts of residents ≥ 40 years of age in 1961, 1974, and 1988 in the Japanese community of Hisayama. Morphological examinations by autopsy or brain imaging were performed on most of the ischemic stroke cases developed in these cohorts. When 13-year follow-up data were compared, the age-adjusted incidence of ischemic stroke and lacunar infarction declined significantly from the first to the third cohort for both sexes, whereas the incidences of atherothrombotic and cardioembolic infarction did not change during this period. Hypertension was a powerful risk factor for the development of ischemic stroke, and improvement of hypertension control would have largely influenced this declining trend: The age- and sex-adjusted hazard ratio of hypertension decreased from 3.25 (95% CI 2.17 to 4.86) in the first cohort to 1.83 (1.29 to 2.58) in the third cohort. A rapid increase in the prevalence of metabolic disorders may have offset the impact of improvements in hypertension control and resulted in a slowdown of the decline in the incidence of ischemic stroke in the cohorts in the present study; however, hypertension still makes a large contribution to the development of ischemic stroke.

Conclusions—These findings suggest that in the Japanese population, the incidence of ischemic stroke has declined significantly over the past 40 years, probably owing to better management of hypertension. There is a need for greater primary prevention efforts in the treatment of hypertension and metabolic disorders. (*Circulation*. 2008;118:2672-2678.)

Key Words: cerebral infarction ■ morbidity ■ risk factors ■ hypertension ■ trend

Stroke continues to be a major public health concern worldwide. In Japan, it is the third leading cause of death and a major neurological cause of long-term disability.¹ The increase in the elderly population that accompanies the improvement in life expectancy is expected to further increase stroke prevalence. On the other hand, there have been major advances in the identification and management of stroke risk factors and the treatment of acute stroke. The study of temporal trends in stroke incidence provides insights into the effect of these factors. Several epidemiological studies have reported that the declining or stable incidence of stroke is likely attributable to better treatment of risk factors over time.²⁻⁸ On the basis of their 50 years of follow-up data, the authors of the Framingham Study recently showed that the age-adjusted incidence of stroke decreased significantly in men and women owing to the improved control of hypertension and smoking.² In Japan, the incidence of stroke declined by 60% from 1964 to 1983 in a rural population.⁷ We also found in a Japanese urban area that the incidence of ischemic stroke declined markedly between the 1960s and

1970s as a result of hypertension control, but this declining trend was slowed in the late 1980s and 1990s, probably because of an increase in metabolic disorders.⁸

Clinical Perspective p 2678

Because the pathogenesis, prognosis, and treatment differ among ischemic stroke subtypes,^{9,10} the evaluation of temporal trends in the incidence of and risk factors for ischemic stroke subtypes may contribute to more effective primary and secondary prevention of ischemic stroke. However, morphological features of the brain were not readily available before the widespread use of computed tomography and magnetic resonance imaging, and the definition of ischemic stroke subtypes was not determined until the early 1990s.¹¹⁻¹³ Therefore, there is little information on the effect of the changes in cardiovascular risk factors on secular trends in the incidence of ischemic stroke and its subtypes.

The Hisayama Study is a population-based study that has established several cohorts at times that correspond to periods of remarkable lifestyle changes in Japan.^{8,14-16} One of the

From the Department of Environmental Medicine (M.K., J.H., Y.T., Y.K.) and Department of Medicine and Clinical Science (Y.D., M.I.), Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan, and Laboratory for Genotyping Development (M.K.), Center for Genomic Medicine, The Institute of Physical and Chemical Research (RIKEN), Yokohama, Kanagawa, Japan.

*The first 2 authors contributed equally to this work.

Correspondence to Michiaki Kubo, MD, Laboratory for Genotyping Development, Center for Genomic Medicine, RIKEN, Suehiro-cho 1-7-22, Tsurumi, Yokohama, Kanagawa, 230-0045, Japan. E-mail mkubo@src.riken.jp

© 2008 American Heart Association, Inc.

Circulation is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.107.743211

characteristics of this study is that most of the deceased study subjects underwent autopsy examination from the beginning of the study, and thus, the morphological features of the brains examined by autopsy or brain imaging are available for most of the stroke cases in each cohort.^{8,14} Furthermore, study-team physicians performed physical and neurological examinations on the subjects who developed stroke and collected detailed clinical information throughout the study period. These characteristics of the study design enabled us to examine secular trends in the incidence of and risk factors for ischemic stroke subtypes. We previously reported the steadily declining incidence of lacunar infarction (LAI) using 12-year follow-up data of the first 3 cohorts.¹⁷ In this article, we extend the follow-up period of these cohorts to 13 years and compare the impact of cardiovascular risk factors on the incidence of ischemic stroke subtypes.

Methods

Study Population

The Hisayama Study, an epidemiological study of cerebrovascular-cardiovascular diseases, was established in 1961 in Hisayama Town, a suburban community adjacent to Fukuoka City, a metropolitan area on Kyushu Island in southern Japan. The population of the town was ~8000 in 2007, and full community surveys of the residents have been repeated since 1961. The study design and characteristics of the subject population have been described in detail elsewhere.¹⁴⁻¹⁶ Briefly, we established 4 study cohorts from Hisayama residents ≥ 40 years of age in 1961, 1974, 1988, and 2002 after screening examinations. In 1961, a total of 1658 subjects in that age group consented to participate in the screening examination (participation rate 90.1%). After the exclusion of subjects with a history of stroke or myocardial infarction and subjects who died or moved out of town during the examination, 1618 subjects were enrolled as the first cohort. Similarly, after excluding subjects with a history of stroke or myocardial infarction, we established a second cohort consisting of 2038 subjects from 2135 participants (participation rate 81.2%) in 1974, a third cohort of 2637 subjects from 2742 participants (participation rate 80.9%) in 1988, and a fourth cohort of 3123 subjects from 3328 participants (participation rate 77.6%) in 2002. The health status of these cohort populations was followed up every year by repeated health examinations or by mail or telephone for any subjects who did not undergo a regular examination or who moved out of town. Only 2 subjects in the first cohort, 2 in the second cohort, and 1 in the third cohort were lost to follow-up. The development of cardiovascular diseases in the study populations was also checked by a daily monitoring system organized by the study team, local physicians, and members of the local health and welfare office. When the subjects died, autopsy examinations were performed at the Department of Pathology, Kyushu University.

Measurement of Cardiovascular Risk Factors

Details of the measurement of cardiovascular risk factors in each cohort were published previously.^{8,14-16} In brief, blood pressures were measured 3 times with subjects in a recumbent position in 1961 and in a sitting position in 1974, 1988, and 2002, and hypertension was defined as a mean systolic blood pressure ≥ 140 mm Hg, a mean diastolic blood pressure ≥ 90 mm Hg, or current use of antihypertensive agents. Glucose intolerance was defined by an oral glucose tolerance test in subjects with glycosuria in 1961, by fasting and postprandial glucose concentrations in 1974, and by a 75-g oral glucose tolerance test in 1988 and 2002, in addition to medical history of diabetes. Serum cholesterol levels were measured by the Zak-Henly method with the modification by Yoshikawa in 1961, by the Zurkowski method in 1974, and by the enzymatic method in 1988 and 2002. Hypercholesterolemia was defined as total cholesterol ≥ 5.7 mmol/L (220 mg/dL). Body height and weight were

measured with subjects in light clothing without shoes, and obesity was defined as body mass index ≥ 25.0 kg/m². Information on antihypertensive treatment, alcohol intake, and smoking habits was obtained with the use of a standardized questionnaire and was categorized as current habitual use or not. Current drinking was also categorized as light (1 to 33 g/d) or heavy (≥ 34 g/d) drinking according to daily ethanol intake.

Definition of Ischemic Stroke Subtypes

Stroke was defined as a sudden onset of nonconvulsive and focal neurological deficit that persisted for >24 hours and was classified as ischemic stroke, cerebral hemorrhage, subarachnoid hemorrhage, or undetermined type.⁸ The diagnoses of ischemic stroke subtypes were made on the basis of the Classification of Cerebrovascular Disease III proposed by the National Institute of Neurological Disorders and Stroke,¹¹ as well as on the basis of the diagnostic criteria of the Trial of Org 10172 in Acute Stroke Treatment (TOAST) study¹² and Cerebral Embolism Task Force.¹³ We classified ischemic stroke subtypes into 4 categories: LAI, atherothrombotic infarction (ATI), cardioembolic infarction (CEI), and undetermined subtype. Details of the diagnostic criteria of ischemic stroke subtypes have been published previously.¹⁰ Briefly, LAI was diagnosed as the presence of a relevant brain stem or subcortical hemispheric lesion with a diameter of <1.5 cm demonstrated on brain imaging or autopsy and no evidence of cerebral cortical or cerebellar impairment. ATI was diagnosed when the subject had significant stenosis ($>50\%$) or occlusion of a major cerebral artery with infarct size ≥ 1.5 cm on brain imaging or autopsy. The diagnosis of CEI was made on the basis of primary and secondary clinical features suggestive of CEI as reported by the Cerebral Embolism Task Force.¹³ The category of undetermined stroke included all ischemic stroke cases for which the subtype could not be determined because of insufficient clinical or morphological information. We considered morphological findings significant and used clinical features as reference information.

During the 13-year follow-up period, first-ever ischemic stroke developed in 134 subjects (83 cases of LAI, 28 of ATI, 17 of CEI, and 6 of undetermined subtype) in the first cohort, in 142 subjects in the second cohort (76 cases of LAI, 29 of ATI, 34 of CEI, and 3 of undetermined subtype), and in 154 subjects in the third cohort (74 cases of LAI, 42 of ATI, 38 of CEI, and 0 of undetermined subtype). Among these, morphological examinations by autopsy or brain imaging were performed on 90.3% (autopsy rate 90.3%) in the first cohort, 97.2% (autopsy rate 87.5%) in the second cohort, and 100.0% (autopsy rate 72.4%) in the third cohort.

Statistical Analysis

The prevalences of possible risk factors were adjusted for age by the direct method and were examined for trends across cohorts by the Cochran-Mantel-Haenszel χ^2 test with 10-year age groupings. Age-adjusted mean values of risk factors were calculated by the covariance method, and their trends were tested by the linear regression model. The incidences of first-ever ischemic stroke and its subtypes were calculated by the person-year method with adjustment for age by the direct method. The world standard population was used as a standard population. The age-adjusted incidences among the first 3 cohorts were compared with the use of the Cox proportional hazards model. Age and sex-adjusted hazard ratios (HRs) and 95% CIs of cardiovascular risk factors for the development of ischemic stroke and its subtypes were estimated by the Cox proportional hazards model in each cohort, and the population attributable risk fraction of each risk factor was calculated.

Ethical Considerations

The study protocol was approved by the Human Ethics Review Committee of the Graduate School of Medical Sciences, Kyushu University.

The authors had full access to the data and take full responsibility for its integrity. All authors have read and agree to the manuscript as written.

Table 1. Trends in Age-Adjusted Prevalence of Cardiovascular Risk Factors Among 4 Examinations of the Hisayama Study by Sex

Variables	Men				<i>P</i> for Trend	Women				<i>P</i> for Trend
	1961 (n=705)	1974 (n=855)	1988 (n=1110)	2002 (n=1315)		1961 (n=913)	1974 (n=1183)	1988 (n=1527)	2002 (n=1808)	
Age, y	55±11	56±11	57±12	60±12	<0.001	57±12	58±12	59±12	62±13	<0.001
Hypertension, %	38.4	43.1	44.1	42.0	0.25	35.9	40.1	35.1	31.3	<0.001
Antihypertensive agents, %	2.0	8.4	13.2	18.2	<0.001	2.1	7.4	13.4	16.6	<0.001
Systolic BP, mm Hg*	162±18	157±18	151±18	148±18	<0.001	163±19	161±19	154±19	149±19	<0.001
Diastolic BP, mm Hg*	91±11	90±11	87±11	89±11	0.011	88±11	87±11	83±11	86±11	<0.001
Glucose intolerance, %	11.6	14.1	39.3	54.5	<0.001	4.8	7.9	30.0	35.5	<0.001
Obesity, %	7.0	11.6	24.1	29.3	<0.001	12.9	21.5	23.8	24.0	<0.001
Body mass index, kg/m ²	21.3±2.8	21.7±2.8	22.8±2.8	23.5±2.8	<0.001	21.7±3.4	22.5±3.3	22.9±3.3	22.9±3.4	<0.001
Hypercholesterolemia, %	2.8	12.2	26.9	25.8	<0.001	6.6	19.9	41.6	41.6	<0.001
Total cholesterol, mmol/L	3.9±0.9	4.7±0.9	5.1±0.9	5.1±0.9	<0.001	4.2±1.0	5.0±1.0	5.5±1.0	5.4±1.0	<0.001
Atrial fibrillation, %	0.7	1.6	1.6	1.1	0.84	0.5	0.4	0.9	0.6	0.55
Current smoking, %	75.0	73.3	50.4	46.9	<0.001	16.6	10.2	6.9	8.5	<0.001
Current drinking, %	69.6	63.8	61.5	71.7	0.043	8.3	5.7	9.5	29.1	<0.001
Light drinking, %	43.4	31.9	29.5	37.7	...	8.2	5.5	8.0	27.1	...
Heavy drinking, %	26.3	31.9	32.0	34.0	...	0.1	0.2	1.5	2.0	...

BP indicates blood pressure. Hypertension was defined as systolic BP \geq 140 mm Hg or diastolic BP \geq 90 mm Hg or current use of antihypertensive agents. Hypercholesterolemia was defined as total cholesterol level \geq 5.7 mmol/L (220 mg/dL). Obesity was defined as body mass index \geq 25.0 kg/m². Current drinking was divided into light (1 to 33 g) and heavy (\geq 34 g) drinking according to daily ethanol intake.

*Mean systolic and diastolic BPs among hypertensive subjects in each examination.

Results

Trends in Cardiovascular Risk Factors

We compared the age-adjusted prevalence of cardiovascular risk factors at baseline examination among the 4 cohorts by sex (Table 1). During the 40-year period from 1961 to 2002, the populations grew 5 years older in both sexes. The age-adjusted prevalence of hypertension was stable at \approx 40% in men (*P* for trend=0.25) and decreased significantly in women (*P* for trend <0.001), whereas the proportion of individuals using antihypertensive agents increased consistently with time in both men and women. As a result, age-adjusted mean blood pressures among hypertensive men and women decreased significantly throughout the study period. In contrast, the age-adjusted prevalence of glucose intolerance and obesity increased greatly over the study period for both sexes. More than half of men and one third of women had glucose intolerance in 2002. The age-adjusted prevalence of hypercholesterolemia increased 10-fold in men and 6-fold in women from 1961 to 1988 but was unchanged in 2002. The age-adjusted prevalence of current smoking for men was 4-fold higher than that for women in 1961, and it decreased significantly with time for both sexes. The prevalence of current drinking increased significantly for both sexes in 2002.

Trends in Incidence of Ischemic Stroke Subtypes

We then compared the age-adjusted incidence of ischemic stroke using the results of a 13-year follow-up in the first 3 cohorts (1st, 2nd, and 3rd cohort). The age-adjusted incidence of ischemic stroke declined significantly for both sexes throughout the cohorts: It significantly declined by 56% for men and by 40% for women from the first to the third cohort

(*P* for trend <0.001 for either sex; Table 2). In regard to ischemic stroke subtypes, the age-adjusted incidence of LAI for men declined significantly by 54% from the first to the second cohort, and it continued to decline by 39% from the second to the third cohort (*P* for trend <0.001). The age-adjusted incidence of LAI for women also declined by 25% from the first to the second cohort, and it continued to decline by 17% from the second to the third cohort (*P* for trend=0.003). The age-adjusted incidence of ATI and CEI did not change significantly among the cohorts for either sex.

Trends in Proportion of Ischemic Stroke Subtype

The proportions of ischemic stroke subtypes among the cohorts are shown by sex in the Figure. For men, the proportion of subjects with LAI decreased steadily from the first to the third cohort, whereas the proportions with ATI and CEI increased. For women, the proportion of the subjects with CEI increased slightly from the first to the third cohort, but the proportions of those with the other subtypes were constant among the cohorts.

Trends in the Effect of Cardiovascular Risk Factors on Ischemic Stroke

Because both cardiovascular risk factors and the incidence of ischemic stroke changed dramatically, we compared the impact of cardiovascular risk factors on the development of ischemic stroke among the first 3 cohorts (Table 3). In the first cohort, hypertension was a powerful risk factor for ischemic stroke (age- and sex-adjusted HR 3.25, 95% CI 2.17 to 4.86) and largely contributed to its occurrence (population attributable risk fraction 51%). The impact of hypertension gradually declined during the study period; however, hyper-

Table 2. Age-Adjusted Incidence Rate (per 1000 Person-Years) of Ischemic Stroke and Its Subtypes Among 3 Cohorts of the Hisayama Study by Sex, With a 13-Year Follow-Up in Each Cohort

	Men			P for Trend	Women			P for Trend
	1st Cohort (7456 PY)	2nd Cohort (9655 PY)	3rd Cohort (12 333 PY)		1st Cohort (10 294 PY)	2nd Cohort (13 762 PY)	3rd Cohort (17 953 PY)	
Ischemic stroke								
No. of events	72	70	70		62	72	84	
Incidence rate	8.73	5.44	3.85	<0.001	4.28	3.06	2.57	<0.001
LAI								
No. of events	48	34	30		35	42	44	
Incidence rate	5.68	2.59	1.59	<0.001	2.41	1.81	1.50	0.003
ATI								
No. of events	14	14	22		14	15	20	
Incidence rate	1.88	1.03	1.23	0.27	0.96	0.61	0.54	0.084
CEI								
No. of events	9	21	18		8	13	20	
Incidence rate	1.08	1.74	1.03	0.43	0.58	0.56	0.53	0.86
Undetermined subtype								
No. of events	1	1	0		5	2	0	
Incidence rate	0.09	0.09	0.00	0.20	0.33	0.08	0.00	0.004

PY indicates person-years.

tension was still a significant risk factor and made the largest contribution to the development of ischemic stroke even in the third cohort (HR 1.83, 95% CI 1.29 to 2.58, population attributable risk fraction 30%). Glucose intolerance was also a significant risk factor for ischemic stroke in the first cohort. The effect of glucose intolerance on the occurrence of ischemic stroke was reduced and was not significant in the second cohort, but it appeared to be a significant risk factor in the third cohort. The population attributable risk fraction for glucose intolerance decreased from 13% in the first cohort to 4% in the second cohort and then increased to 13% in the third cohort. Obesity appeared to be a significant risk factor for ischemic stroke in every cohort, and its population attributable risk fraction was increased gradually from 6% in the first cohort to 9% in the third cohort. Hypercholesterol-

emia, smoking habits, and alcohol intake were not significant risk factors for ischemic stroke in any of the cohorts. In the multivariate analysis that included all risk factors, hypertension was a significant risk factor for ischemic stroke, and its HR decreased from 2.92 (95% CI 1.93 to 4.41) in the first cohort to 1.71 (95% CI 1.20 to 2.45) in the third cohort. Glucose intolerance was an independent risk factor for ischemic stroke in the first cohort (HR 1.91, 95% CI 1.23 to 2.95) but was not significant in the third cohort (HR 1.28, 95% CI 0.93 to 1.78). Obesity was not a significant risk factor in any of the cohorts after adjustment for other risk factors. We tried to investigate the effect of cardiovascular risk factors on ischemic stroke subtypes, but we could not find reliable evidence of an effect of these risk factors on the development of each subtype, probably because of the small number of events.




Figure. Proportion of ischemic stroke subtypes among the 3 cohorts of the Hisayama Study by sex.

Table 3. Age- and Sex-Adjusted HRs of Cardiovascular Risk Factors for Ischemic Stroke Among 3 Cohorts of the Hisayama Study

	1st Cohort			2nd Cohort			3rd Cohort		
	HR (95% CI)	<i>P</i>	PAF	HR (95% CI)	<i>P</i>	PAF	HR (95% CI)	<i>P</i>	PAF
Hypertension	3.25 (2.17–4.86)	<0.001	0.51	3.09 (2.05–4.65)	<0.001	0.53	1.83 (1.29–2.58)	<0.001	0.30
Glucose intolerance	2.45 (1.60–3.74)	<0.001	0.13	1.38 (0.87–2.17)	0.17	0.04	1.41 (1.02–1.94)	0.036	0.13
Obesity	1.83 (1.12–3.00)	0.017	0.06	1.63 (1.04–2.57)	0.034	0.07	1.54 (1.07–2.21)	0.021	0.09
Hypercholesterolemia	1.07 (0.50–2.29)	0.87	0.00	1.42 (0.95–2.12)	0.085	0.07	0.96 (0.68–1.35)	0.80	–0.02
Current smoker	1.27 (0.85–1.90)	0.24	0.10	0.83 (0.55–1.24)	0.36	–0.08	1.33 (0.89–1.98)	0.16	0.07
Current drinker	0.99 (0.65–1.51)	0.94	–0.01	1.45 (0.96–2.19)	0.081	0.12	1.09 (0.72–1.64)	0.70	0.02

PAF indicates the population attributable risk fraction.

Discussion

By comparing the incidence of ischemic stroke subtypes among 3 cohorts established at different times in a Japanese community, we demonstrated that the incidence of LAI declined significantly from the first to the third cohort for both sexes, whereas the incidence of ATI and CEI remained stable. During the study period, blood pressure levels among hypertensive subjects decreased significantly with time as a result of the popularization of antihypertensive medication. The prevalence of smoking habits declined steadily for both sexes. Contrary to these declining trends, the prevalence of metabolic disorders, namely, obesity, glucose intolerance, and hypercholesterolemia, increased steeply with time. These changes in cardiovascular risk factors might affect the incidence of ischemic stroke and its subtypes.

Hypertension is the most powerful risk factor for ischemic stroke.⁹ In the first cohort, hypertension contributed to approximately half of the occurrence of ischemic stroke. During the study period, the age-adjusted prevalence of hypertension declined in women, and the proportion of all participants receiving hypertensive treatment increased steeply in both sexes. This improvement of hypertension control resulted in a decrease in age-adjusted mean systolic blood pressure level of 14 mm Hg among hypertensive subjects in both sexes. Because of this improved control of hypertension, the impact of the disease on the development of ischemic stroke was seen to weaken in the third cohort. The Framingham Study also showed a decline in the annual incidence of nonembolic stroke during a follow-up period of 50 years or more.² During this period, the mean systolic blood pressure level, prevalence of hypertension, and proportion of all participants receiving treatment for hypertension improved significantly. These reductions in the incidence of ischemic stroke and improvements in treatment for hypertension were similar to the findings of the present study. Our previous study showed that the impact of hypertension was similar for all ischemic stroke subtypes.¹⁰ These results suggest that better management of hypertension might have made the biggest contribution to the declining trend in the incidence of ischemic stroke, especially of LAI; however, hypertension was still a significant risk factor in the third cohort and had a large attributable risk fraction for ischemic stroke. Because half of the hypertensive subjects did not undergo treatment for hypertension in the third cohort, there is a need for greater primary prevention efforts to improve the treatment of hypertension.

In subjects in the present study, the age-adjusted prevalence of metabolic disorders, such as obesity, hypercholesterolemia, and glucose intolerance, increased greatly during the past 40 years, probably owing to the westernization of the Japanese lifestyle. When we examined the impact of these metabolic disorders on the development of ischemic stroke, glucose intolerance was a significant risk factor in the first and the third cohort, and the impact of obesity was constant throughout the study period. Both glucose intolerance and body mass index have been shown to be significant risk factors for ischemic stroke and LAI.^{10,18} Moreover, obesity is closely related to other cardiovascular risk factors and jointly increases the risk of ischemic stroke.¹⁹ Our previous study also showed that the accumulation of metabolic disorders (that is, metabolic syndrome) was a significant risk factor for the development of ischemic stroke in our third cohort.²⁰ We speculate that the improved management of hypertension and the worsening of metabolic disorders cancelled each other out and resulted in the slowdown of the declining trend of the incidence of LAI and the sustained incidence of ATI.

Smoking is a widely accepted risk factor for ischemic stroke in Western populations, but this relationship is controversial for Japanese.^{10,21,22} In the present study cohorts, the declining prevalence of smoking habits closely mirrored the declining trend in the incidence of ischemic stroke; however, smoking habits had little impact on the incidence of ischemic stroke in the present study cohorts. One possible explanation is that the association between smoking and the risk of ischemic stroke is only evident in populations with moderate to high levels of serum cholesterol.²³ A recent review of cardiovascular mortality trends in Japan²³ showed that the increase in serum cholesterol appeared mainly in young to middle-aged people. In contrast, elderly people, a high-risk group for ischemic stroke, continued to maintain a lower cholesterol level. However, the prevalence of smoking habits is still high in Japanese men, and therefore, the adverse influence of smoking might appear in the current generation of younger men, with a higher cholesterol level to be seen in the future.

LAI is the most common subtype of ischemic stroke in the Japanese population, unlike in Western populations.¹ Among subjects in the present study, because of the decreased incidence of LAI and the sustained incidences of ATI and CEI, the proportion of ischemic stroke subtypes has become closer to that of Western populations in men (Figure).

However, the pattern of ischemic stroke subtypes differed from that of Western populations, with subjects in the present study showing a high proportion of LAI even in recent years (43% for men and 52% for women in the third cohort). A recent hospital-based registration study in an urban area²⁴ and a study of 16 992 patients with acute ischemic stroke from rural areas in Japan²⁵ also showed a higher prevalence of LAI than of other subtypes. One possible explanation for this is the racial difference in the genetic susceptibility of LAI. We recently found 2 susceptibility genes for ischemic stroke, *PRKCH* and *AGTRL1*, in a genome-wide association study.^{26,27} A single-nucleotide polymorphism in the *PRKCH* gene increased the risk of LAI, but this single-nucleotide polymorphism is specific to Asian populations.²⁷

The present study has several limitations. First, the number of events of subtypes other than LAI was relatively small, and therefore, the power to assess trends in the incidence of and risk factors for ischemic stroke subtypes was weak. Second, there were a large number of subjects overlapping among the cohorts. Indeed, 916 of the subjects in the first cohort also accounted for 45% of the population of the second cohort. In addition, a total of 1229 subjects in the second cohort also participated in the third cohort (47% of the third cohort). However, we treated the overlapping subjects as in any life table analysis, establishing every cohort after excluding subjects with prior stroke or myocardial infarction at baseline. Therefore, these overlapping populations were not considered to distort the incidence trends in the present study. Third, the measurement of blood glucose and the criteria for glucose intolerance were different among the cohorts, which suggests an underestimation of the prevalence of glucose intolerance in the former cohorts. Nevertheless, the rapid changes in other risk factors in the present study are in accordance with the results of the National Nutritional Survey and other surveys of Japan.²³ Finally, the methods of case ascertainment and the diagnostic sensitivity of imaging techniques changed dramatically during the study period. The proportion of case subjects with of incident ischemic stroke who received diagnostic imaging tests increased over time. Echocardiography and carotid scanning were rarely performed in the former cohorts (3.0% and 0% in the first cohort, 29.6% and 4.2% in the second cohort, and 61.7% and 27.3% in the third cohort, respectively). Therefore, it is possible that the trends in the incidence of ATI and CEI were less accurate than the trends for LAI. Nonetheless, we believe that the findings of the present study reflect the actual secular trends in the incidence of ischemic stroke subtypes and their risk factors in the Japanese population, because we performed comprehensive surveillance, including autopsy examinations, in most of the cases.

Conclusions

By comparing the incidence of and risk factors for ischemic stroke subtypes among 3 cohorts established at different times in a Japanese community, we demonstrated that the incidence of LAI declined significantly from the 1960s to the late 1990s, but LAI remained the most frequent subtype of ischemic stroke in the Japanese. The improvement in hypertension control might have had a major influence on this

declining trend. However, hypertension still has a large impact on ischemic stroke, and the increasing prevalence of metabolic disorders might emerge as an additional risk in future cohorts. The present study indicates the need for continued primary prevention efforts, particularly with respect to hypertension and metabolic disorders.

Sources of Funding

This study was supported in part by a grant-in-aid for scientific research A (No. 18209024) from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

Disclosure

None.

References

- Hachinski V. Stroke in Japanese. *Stroke*. 2006;37:1143.
- Carandang R, Seshadri S, Beiser A, Kelly-Hayes M, Kase CS, Kannel WB, Wolf PA. Trends in incidence, lifetime risk, severity, and 30-day mortality of stroke over the past 50 years. *JAMA*. 2006;296:2939–2946.
- Mayo NE, Nadeau L, Daskalopoulou SS, Cote R. The evolution of stroke in Quebec: a 5-year perspective. *Neurology*. 2007;68:1122–1127.
- Kleindorfer D, Broderick J, Khoury J, Flaherty M, Woo D, Alwell K, Moomaw CJ, Schneider A, Miller R, Shukla R, Kissela B. The unchanging incidence and case-fatality of stroke in the 1990s: a population-based study. *Stroke*. 2006;37:2473–2478.
- Kagan A, Popper J, Reed DM, MacLean CJ, Grove JS. Trends in stroke incidence and mortality in Hawaiian Japanese men. *Stroke*. 1994;25:1170–1175.
- Sytkowski PA, D'Agostino RB, Belanger A, Kannel WB. Sex and time trends in cardiovascular disease incidence and mortality: the Framingham Heart Study, 1950–1989. *Am J Epidemiol*. 1996;143:338–350.
- Shimamoto T, Komachi Y, Inada H, Doi M, Iso H, Sato S, Kitamura A, Iida M, Konishi M, Nakanishi N, Terao A, Naito Y, Kojima S. Trends for coronary heart disease and stroke and their risk factors in Japan. *Circulation*. 1989;79:503–515.
- Kubo M, Kiyohara Y, Kato I, Tanizaki Y, Arima H, Tanaka K, Nakamura H, Okubo K, Iida M. Trends in the incidence, mortality, and survival rate of cardiovascular disease in a Japanese community: the Hisayama Study. *Stroke*. 2003;34:2349–2354.
- Sacco RL. Risk factors, outcomes, and stroke subtypes for ischemic stroke. *Neurology*. 1997;49(suppl 4):S39–S44.
- Tanizaki Y, Kiyohara Y, Kato I, Iwamoto H, Nakayama K, Shinohara N, Arima H, Tanaka K, Ibayashi S, Fujishima M. Incidence and risk factors for subtypes of cerebral infarction in a general population: the Hisayama Study. *Stroke*. 2000;31:2616–2622.
- Special report from the National Institute of Neurological Disorders and Stroke: classification of cerebrovascular disease III. *Stroke*. 1990;21:637–676.
- Adams HP Jr, Bendixen BH, Kappelle LJ, Biller J, Love BB, Gordon DL, Marsh EE III; TOAST Investigators. Classification of subtype of acute ischemic stroke: definitions for use in a multicenter clinical trial. *Stroke*. 1993;24:35–41.
- Cerebral Embolism Task Force. Cardiogenic brain embolism. *Arch Neurol*. 1986;43:71–84.
- Katsuki S. Epidemiological and clinicopathological study on cerebrovascular disease in Japan. *Prog Brain Res*. 1966;21:64–89.
- Omae T, Ueda K, Kikumura T, Shikata T, Fujii I, Yanai T, Hasuo Y. Cardiovascular deaths among hypertensive subjects of middle to old age: a long-term follow-up study in a Japanese community. In: Onesti G, Kim KE, eds. *Hypertension in the Young and Old*. New York, NY: Grunc & Stratton; 1981:285–297.
- Fujishima M, Kiyohara Y, Kato I, Ohmura T, Iwamoto H, Nakayama K, Ohmori S, Yoshitake T. Diabetes and cardiovascular disease in a prospective population survey in Japan: the Hisayama Study. *Diabetes*. 1996;45(suppl 3):S14–S16.
- Kubo M, Kiyohara Y, Ninomiya T, Tanizaki Y, Yonemoto K, Doi Y, Hata J, Oishi Y, Shikata K, Iida M. Decreasing incidence of lacunar vs other types of cerebral infarction in a Japanese population. *Neurology*. 2006;66:1539–1544.
- Oki I, Nakamura Y, Okamura T, Okayama A, Hayakawa T, Kita Y, Ueshima H; NIPPON DATA80 Research Group. Body mass index and risk

- of stroke mortality among a random sample of Japanese adults: 19-year follow-up of NIPPON DATA80. *Cerebrovasc Dis*. 2006;22:409–415.
19. Miyamatsu N, Kadowaki T, Okamura T, Hayakawa T, Kita Y, Okayama A, Nakamura Y, Oki I, Ueshima H. Different effects of blood pressure on mortality from stroke subtypes depending on BMI levels: a 19-year cohort study in the Japanese general population - NIPPON DATA80. *J Hum Hypertens*. 2005;19:285–291.
 20. Ninomiya T, Kubo M, Doi Y, Yonemoto K, Tanizaki Y, Rahman M, Arima H, Tsuruya K, Iida M, Kiyohara Y. Impact of metabolic syndrome on the development of cardiovascular disease in a general Japanese population: the Hisayama Study. *Stroke*. 2007;38:2063–2069.
 21. Kiyohara Y, Ueda K, Fujishima M. Smoking and cardiovascular disease in general population in Japan. *J Hypertens*. 1990;8:S9–S15.
 22. Ueshima H, Choudhury SR, Okayama A, Hayakawa T, Kita Y, Kadowaki T, Okamura T, Minowa M, Iimura O; NIPPON DATA80 Research Group. Cigarette smoking as a risk factor for stroke death in Japan: NIPPON DATA80. *Stroke*. 2004;35:1836–1841.
 23. Ueshima H. Explanation for the Japanese paradox: prevention of increase in coronary heart disease and reduction in stroke. *J Atheroscler Thromb*. 2007;14:278–286.
 24. Kitamura A, Nakagawa Y, Sato M, Iso H, Sato S, Imano H, Kiyama M, Okada T, Okada H, Iida M, Shimamoto T. Proportions of stroke subtypes among men and women ≥ 40 years of age in an urban Japanese city in 1992, 1997, and 2002. *Stroke*. 2006;37:1374–1378.
 25. Kimura K, Kazui S, Minematsu K, Yamaguchi T; for the Japan Multi-center Stroke Investigators' Collaboration (J-MUSIC). Analysis of 16992 patients with acute ischemic stroke and transient ischemic attack in a hospital-based prospective registration study. *Cerebrovasc Dis*. 2004;18:47–56.
 26. Hata J, Matsuda K, Ninomiya T, Yonemoto K, Matsushita T, Ohnishi Y, Saito S, Kitazono T, Ibayashi S, Iida M, Kiyohara Y, Nakamura Y, Kubo M. Functional SNP in an Sp1-binding site of *AGTRL1* gene is associated with susceptibility to brain infarction. *Hum Mol Genet*. 2007;16:630–639.
 27. Kubo M, Hata J, Ninomiya T, Matsuda K, Yonemoto K, Nakano T, Matsushita T, Yamazaki K, Ohnishi Y, Saito S, Kitazono T, Ibayashi S, Sueishi K, Iida M, Nakamura Y, Kiyohara Y. A nonsynonymous SNP in *PRKCH* (protein kinase C η) increases the risk of cerebral infarction. *Nat Genet*. 2007;39:212–217.

CLINICAL PERSPECTIVE

Stroke continues to be a major public health concern worldwide. Several epidemiological studies have reported that the declining or stable incidence of stroke is most often attributed to better treatment of risk factors over time. Here, by comparing the incidence of and risk factors for ischemic stroke subtypes among 3 cohorts established at different times in a Japanese community, we demonstrate that the age-adjusted incidence of ischemic stroke and of lacunar infarction declined significantly from the 1960s to the late 1990s, but lacunar infarction remains the most frequent subtype of ischemic stroke in the Japanese. Hypertension was a powerful risk factor for the development of ischemic stroke, and improvement of hypertension control would have largely influenced this declining trend: The age- and sex-adjusted hazard ratio of hypertension decreased from 3.25 (95% CI 2.17 to 4.86) in the first cohort to 1.83 (1.29 to 2.58) in the third cohort. However, hypertension still has a large impact on ischemic stroke, and the increase in metabolic disorders might emerge as an additional risk in the third cohort. The present study indicates the need for continued primary prevention efforts, particularly with respect to hypertension and metabolic disorders.

Impact of Kidney Disease and Blood Pressure on the Development of Cardiovascular Disease

An Overview From the Japan Arteriosclerosis Longitudinal Study

Toshiharu Ninomiya, MD, PhD; Yutaka Kiyohara, MD, PhD; Yosuke Tokuda, MS;
Yasufumi Doi, MD, PhD; Hisatomi Arima, MD, PhD; Akiko Harada, PhD;
Yasuo Ohashi, PhD; Hirotsugu Ueshima, MD, PhD;
for the Japan Arteriosclerosis Longitudinal Study Group

Background—Kidney disease is associated with an increased risk of cardiovascular disease (CVD); however, there have been few well-designed prospective studies of this issue in Asian populations. Recent epidemiological studies have suggested that a lower blood pressure level may be associated with an increased risk of CVD in individuals with kidney dysfunction.

Methods and Results—Using data from 10 community-based cohort studies in Japan, we conducted follow-up on a total of 30 657 individuals 40 to 89 years of age without preexisting CVD or kidney failure and examined the relationship between reduced glomerular filtration rate (GFR) and the risk of CVD. During an average 7.4-year follow-up, 727 individuals experienced CVD. The age- and sex-adjusted incidence of CVD increased significantly in subjects with GFR of 60 to 89 mL · min⁻¹ · 1.73 m⁻² (4.3 per 1000 person-years, *P*=0.002) and in those with a GFR <60 mL · min⁻¹ · 1.73 m⁻² (6.5, *P*<0.001) compared with those with a GFR ≥90 mL · min⁻¹ · 1.73 m⁻² (2.9). Even after adjustment for potential confounding factors, subjects with a GFR <60 mL · min⁻¹ · 1.73 m⁻² had a 57% (95% CI 14% to 115%) greater risk of CVD than those with a GFR ≥90 mL · min⁻¹ · 1.73 m⁻². The multivariate-adjusted hazard ratios of CVD increased in a log-linear manner with elevations in blood pressure levels, regardless of GFR levels (all *P* for trend <0.01).

Conclusions—Our findings suggest that a reduced GFR is a significant risk factor for CVD in the general Japanese population. Additionally, a log-linear association of blood pressure level with CVD risk was observed, without evidence of a J-curve association, regardless of GFR levels. (*Circulation*. 2008;118:2694-2701.)

Key Words: cardiovascular diseases ■ blood pressure ■ kidney ■ meta-analysis

Kidney disease is increasingly being recognized as a leading public health issue. Chronic kidney disease, most commonly defined by a reduction in glomerular filtration rate (GFR) or the presence of proteinuria, affects 10% to 15% of the adult population in Western countries^{1,2} and is associated with an increased risk of cardiovascular disease (CVD)³⁻⁵; however, there have been few well-designed large prospective studies in general Asian communities to date.⁶⁻⁸

Clinical Perspective p 2701

Blood pressure is an important determinant of the risk of CVD in the general population,^{9,10} in which it has been well established that treatment for high blood pressure prevents CVD.¹¹ Blood pressure is commonly elevated in individuals

with a reduced GFR,^{4,5} which suggests that lowering blood pressure may offer significant benefits in this population. Recent prospective cohort studies, however, have reported that the risk of stroke or death for individuals with a reduced GFR is greater among those with systolic blood pressure levels below 120 mm Hg than among those with higher levels.^{12,13} These data have raised concerns that lowering blood pressure may provide less benefit than previously believed, or may even be hazardous, in individuals with kidney dysfunction.

In the present report, we discuss the results of a pooling analysis from the Japan Arteriosclerosis Longitudinal Study—Existing Cohorts Combine (JALS-ECC), which is an overview of individual participant data from 21 community-based

Continuing medical education (CME) credit is available for this article. Go to <http://cme.ahajournals.org> to take the quiz.

From the Department of Environmental Medicine (T.N., Y.K., H.A.) and Department of Medicine and Clinical Science (Y.D.), Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan; Department of Biostatistics/Epidemiology and Preventive Health Sciences (Y.T., A.H., Y.O.), School of Health Sciences and Nursing, University of Tokyo, Tokyo, Japan; and Department of Health Science (H.U.), Shiga University of Medical Science, Otsu, Japan.

The online-only Data Supplement is available with this article at <http://circ.ahajournals.org/cgi/content/full/118/25/2694/DC1>.

Correspondence to Yutaka Kiyohara, MD, PhD, Department of Environmental Medicine, Graduate School of Medical Sciences, Kyushu University, Maidashi 3-1-1, Higashi-ku, Fukuoka, 812-8582 Japan. E-mail kiyohara@envmed.med.kyushu-u.ac.jp

© 2008 American Heart Association, Inc.

Circulation is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.108.792903

Downloaded from circ.ahajournals.org at KYUSHU UNIVERSITY on February 3, 2009

longitudinal observational studies in Japan.¹⁴ Our aims were to assess the impact of a reduction in GFR on the development of CVD in the general population and to examine the association of blood pressure with the risk of CVD in individuals with a reduced GFR.

Methods

Study Population

The rationale, study design, and methods of the JALS-ECC have been described elsewhere.¹⁴ In brief, cohort studies were eligible for inclusion in this project if they satisfied the following criteria: (1) Japanese population; (2) prospective cohort study; (3) at least 3000 person-years of follow-up; (4) date of birth, sex, height, weight, blood pressure, and serum total cholesterol recorded at baseline; and (5) date of death or the age at death recorded during a follow-up. Quality control of the collected cohort data were performed at the JALS Coordinating Center. The individual records of 66 691 participants in 21 cohort studies were included in the present project, with 82.7% of the participants from 17 community-based cohorts and 17.3% from 4 work-site-based cohorts. Permission to submit each collection of cohort data to the JALS Coordinating Center was obtained from the relevant institutional review boards for ethical issues.

Of the 21 cohort studies, 11 cohorts were excluded from the present analysis for the following reasons: 4 were work-site-based cohorts, 3 did not include creatinine data, 3 lacked many values for relevant variables, and 1 included no event data for either stroke or myocardial infarction. From the remaining 10 cohorts, we excluded participants less than 40 years of age and those 90 years of age or older, those with unavailable examination data at baseline or unavailable event data, those with a history of CVD, and those with an estimated GFR $<15 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$. For the analysis regarding CVD and stroke, the Niigata cohort was excluded, because information on stroke events was unavailable. A final total of 23 033 participants were enrolled in the CVD analysis, 23 084 in the stroke analysis, 30 657 in the myocardial infarction analysis, and 31 374 in the all-cause death analysis. The average follow-up period was 7.4 years.

Risk Factors

The JALS Coordinating Center requested individual participant data from the collaborating investigators. Serum creatinine was measured by Jaffe's method in 8 cohorts, by the enzymatic method in 1 cohort, and by both methods in 1 cohort. Serum creatinine values measured by the enzymatic method were corrected for Japanese subjects by the addition of $18.3 \mu\text{mol/L}$.¹⁵ GFR was estimated with the 4-variable Modification of Diet in Renal Disease study equation.³ In accordance with the National Kidney Foundation Kidney Disease Outcomes Quality Initiative guidelines,¹⁶ GFR levels were classified in the following ranges: ≥ 90 , 60 to 89, and $<60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$. Blood pressure was measured by a standard sphygmomanometer in all cohorts. Mean values were used in several cohorts that measured 2 or more blood pressure values. Blood pressure levels at baseline were classified into 4 categories (normal, prehypertension, stage 1 hypertension, and stage 2 hypertension) according to the criteria of the seventh report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure.¹⁷ Diabetes was defined as a fasting blood glucose level of $\geq 7.0 \text{ mmol/L}$, a casual blood glucose level of $\geq 11.1 \text{ mmol/L}$, current use of insulin or oral medication for diabetes, and/or a history of diabetes. Serum total cholesterol was determined enzymatically. Information on smoking habit was obtained through a standard questionnaire and classified as current habitual use or lack thereof.

End Points

In each cohort, vital status and the development of CVD were ascertained during follow-up by use of death certificates, hospital medical records, and/or questionnaire surveys. All outcomes were

classified according to the International Classification of Diseases, 9th Revision (ICD-9). All events were recoded by coordinating center staff members. CVD was defined as the development of either stroke or myocardial infarction. Stroke was defined as an acute disturbance of focal neurological function with symptoms that lasted >24 hours or death caused by a stroke event (ICD-9 codes 430, 431, 433, 434, or 436). Myocardial infarction included both fatal and nonfatal myocardial infarction, which was diagnosed by use of an appropriate clinical history supported by ECG changes and/or elevations of cardiac enzymes or other biochemical markers of myocardial injury (ICD-9 410). Only the first event of the relevant outcome type was included in each analysis.

Statistical Analysis

The SAS software package for Windows, release 9.13 (SAS Institute, Inc, Cary, NC) was used to perform all statistical analyses. The incidence rate of each outcome for the GFR subgroups was calculated by the person-year method and adjusted for the age and sex distribution of the overall population enrolled in the CVD analysis by the direct method, in which the subgroups and study population were subdivided into the same set of age groups (defined by decade) and the age- and sex-specific incidence rates were calculated within each subgroup.¹⁸ The hazard ratios (HRs) and their 95% CIs for the development of events were estimated with the Cox proportional hazards regression model. The cohort effect was adjusted as a fixed effect by taking the study as a strata variable, assuming only proportional hazards within each study and not between studies.¹⁸ Heterogeneity across cohorts was examined with the Cochran Q test and the I^2 statistic.¹⁸ The risks of events according to blood pressure levels were also estimated with the Cox regression model. Trends in relationship between blood pressure levels and the risk of events were assessed by fitting models with a linear term for blood pressure categories according to kidney function status, and the heterogeneity of these relationships between kidney function status subgroups was estimated by the addition of an interaction term of a linear term for blood pressure levels and kidney function status to the relevant model. $P < 0.05$ was considered statistically significant in all analyses.

The authors had full access to and take full responsibility for the integrity of the data. All authors have read and agree to the manuscript as written.

Results

The characteristics of the 10 cohorts examined in the present study are shown in Table 1. Among all subjects, the mean age was 57.6 years, and the proportion of men was 38.0%. The mean value of serum creatinine was $78.6 \mu\text{mol/L}$, and the frequency of GFR $<60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ was 8.2%. During the average follow-up period of 7.4 years, a total of 727 subjects experienced CVD, 592 had strokes, 180 had myocardial infarctions, and a total of 2104 died.

Table 2 shows the baseline characteristics of the 23 033 subjects enrolled in the CVD analysis by sex. Their mean ages were 56.9 years for men and 58.2 years for women, and the frequency of GFR $<60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ was 5.2% for men and 10.1% for women. The frequencies of normal blood pressure, prehypertension, stage 1 hypertension, and stage 2 hypertension were 19.6%, 41.8%, 26.0%, and 12.6% for men and 24.3%, 41.0%, 24.3%, and 10.4% for women, respectively. Similar findings were observed in subjects enrolled in the analyses of stroke, myocardial infarction, and all-cause death.

The age- and sex-adjusted incidences of CVD and stroke increased with declining GFR levels in the overall population (Table 3); the differences were statistically significant between subjects with a GFR $\geq 90 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ and

Table 1. Characteristics of Included Cohort Studies

Regions	Cohort Name	No. of Population	Mean Age, y	Men, %	Mean sCr, $\mu\text{mol/L}^*$	GFR <60 , % †	Mean SBP/DBP, mm Hg	Mean Fasting Blood Glucose, mmol/L	Mean Total Cholesterol, mmol/L	Current Smoking, %	Follow-Up, y		No. of Events			
											Start-End	Mean	CVD	Stroke	MI	Death
Hokkaido	Tanno/Soubetsu	2066	60.1	43.9	89.6	19.4	133/78	5.1	5.0	25.9	1991–1999	5.5	120	93	27	136
Akita 2	Ikawa	2595	56.1	43.6	76.0	2.6	135/81	6.6	4.9	28.5	1985–1999	10.7	44	41	3	146
Ibaraki	Kyowa	4479	54.8	42.8	76.9	5.3	137/82	6.9	5.0	30.4	1985–1999	10.1	168	128	51	350
Niigata	Tokamachi	8480	58.0	33.1	79.6	7.7	127/73	NA	5.1	18.8	1993–2003	7.8	NA	NA	29	400
Osaka	Yao	3855	54.0	34.8	78.7	6.7	132/80	6.0	5.2	27.2	1985–1998	9.6	79	62	18	191
	Minami-takayasu															
Shiga 1	Shigaraki	2934	56.6	41.1	81.3	10.5	132/78	6.0	5.0	29.4	1982–2001	7.3	82	69	13	260
Hiroshima	Hiroshima	2222	72.1	28.7	84.0	23.8	136/78	6.2	5.6	15.4	1992–2000	3.6	73	63	12	350
Ehime	Ohzu	5300	59.5	33.9	76.9	6.2	130/76	5.3	5.3	15.2	1996–2003	5.5	99	89	10	184
Fukuoka 1	Hisayama	757	60.8	39.5	83.1	9.5	133/78	5.4	5.4	21.1	1990–2000	9.9	57	45	14	86
Kumamoto	...	2465	47.0	70.0	65.4	0.2	127/80	5.7	5.4	46.4	1999–2003	4.2	5	2	3	1
Total	...	35 153	57.6	38.0	78.6	8.2	131/78	6.0	5.2	24.5	1985–2003	7.4	727	592	180	2104

sCr indicates serum creatinine; SBP/DBP, systolic or diastolic blood pressure; MI, myocardial infarction; and NA, not available.

*Serum creatinine was measured by Jaffe's method in 8 cohorts, by enzymatic method in the Ehime cohort, and by either method in the Niigata cohort. The values of serum creatinine measured by the enzymatic method were corrected by the addition of 18.3 $\mu\text{mol/L}$.

$^\dagger\text{GFR}$ (unit: $\text{mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$) was estimated by the Modification of Diet in Renal Disease formula.

those with a $\text{GFR} < 90 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ (all $P < 0.01$). Subjects with a $\text{GFR} < 60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ showed a significantly higher age- and sex-adjusted incidence of myocardial infarction and all-cause mortality than those with a $\text{GFR} \geq 90 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ ($P < 0.001$). The age-adjusted incidences of CVD, stroke, and all-cause mortality were significantly higher in subjects with a $\text{GFR} < 60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ than in those with a $\text{GFR} \geq 90 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ in both sexes (all $P < 0.05$).

The risks of CVD, stroke, myocardial infarction, and all-cause death increased progressively with declining GFR

Table 2. Baseline Characteristics of the Study Population by Sex

Risk Factors	Men (n=9574)	Women (n=13 459)
Age, y	56.9 (11.1)	58.2 (11.4)
Serum creatinine, $\mu\text{mol/L}$	87.3 (16.7)	71.6 (13.6)
GFR, $\text{mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$	87.3 (20.2)	81.0 (19.1)
GFR levels ($\text{mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$), %		
≥ 90	39.0	27.1
60–89	55.9	62.8
< 60	5.2	10.1
Systolic blood pressure, mm Hg	133.6 (19.1)	132.3 (19.8)
Diastolic blood pressure, mm Hg	81.1 (11.5)	77.9 (11.0)
Blood pressure levels, %		
Normal	19.6	24.3
Prehypertension	41.8	41.0
Stage 1 hypertension	26.0	24.3
Stage 2 hypertension	12.6	10.4
Diabetes, %	9.5	5.2
Serum total cholesterol, mmol/L	5.0 (0.9)	5.4 (1.0)
Body mass index, kg/m^2	23.2 (3.0)	23.2 (3.3)
Current smoking, %	56.2	7.1

Values are means (SD) or frequencies.

levels in the overall population after adjustment for age and sex (Table 4). Even after adjustment for potential confounding factors, specifically age, sex, cohort, systolic blood pressure, diabetes, serum total cholesterol, body mass index, and current smoking status, the risks of CVD, myocardial infarction, and all-cause death were significantly higher in subjects with a $\text{GFR} < 60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ than in the overall population. There was no evidence of heterogeneity in these associations among study cohorts (all P for heterogeneity > 0.6 ; $Q = 2.46$, $I^2 = 0\%$ for CVD; $Q = 4.06$, $I^2 = 0\%$ for stroke; $Q = 3.75$, $I^2 = 0\%$ for myocardial infarction; and $Q = 1.14$, $I^2 = 0\%$ for all-cause death). Subjects with a $\text{GFR} < 60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ had a significantly greater risk of myocardial infarction and death in men and of CVD, stroke, and death in women.

The Figure shows the log-linear relationship between blood pressure levels at baseline and the hazard of CVD, stroke, and all-cause death regardless of kidney function status after adjustment for potential confounding factors (all P for trend < 0.01). There was no evidence of heterogeneity of the patterns in the association of blood pressure levels with the risk of outcomes between subgroups of kidney function status (all P for heterogeneity > 0.7). The age- and sex-adjusted HR of myocardial infarction increased in a log-linear fashion with increasing blood pressure levels in the normal, prehypertension, stage 1 hypertension, and stage 2 hypertension groups in subjects with a $\text{GFR} \geq 60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ (HR 0.56 [95% CI 0.33 to 0.95], 1.00 [reference], 1.60 [1.08 to 2.37], and 1.75 [1.06 to 2.87]; P for trend 0.03) and in those with a $\text{GFR} < 60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ (0.19 [0.02 to 1.47], 1.00 [reference], 1.72 [0.80 to 3.70], and 2.36 [1.02 to 5.44]; P for trend 0.04). The number of myocardial infarctions in subjects with normal blood pressure levels was too small to assess reliably for multivariate-adjusted analysis.

We also performed sensitivity analyses to assess the risk of CVD according to GFR levels estimated by the MDRD formula corrected according to the Japanese coefficient of 0.881.¹⁵ The correction shifted the GFR distribution to a

Table 3. Incidence Rate of CVD According to Kidney Function Status

GFR Levels, mL · min ⁻¹ · 1.73 m ⁻²	Overall				Men				Women			
	No. of Events	No. of Participants	PY at Risk	Incidence Rate per 1000 PY (95% CI)*	No. of Events	No. of Participants	PY at Risk	Incidence Rate per 1000 PY (95% CI)*	No. of Events	No. of Participants	PY at Risk	Incidence Rate per 1000 PY (95% CI)*
CVD												
GFR ≥90	105	7199	51 203	2.9 (2.1–3.6)	78	3672	23 964	4.4 (3.3–5.6)	27	3527	27 239	1.8 (0.9–2.8)
GFR 60–89	489	13 967	104 334	4.3 (3.9–4.7)†	245	5404	39 794	5.5 (4.8–6.2)	244	8563	64 540	3.5 (3.1–3.9)†
GFR <60	133	1867	12 013	6.5 (5.0–8.0)‡	49	498	3018	9.1 (5.7–12.5)‡	84	1369	8995	4.7 (3.6–5.8)‡
Stroke												
GFR ≥90	84	7206	51 315	2.2 (1.6–2.8)	61	3676	24 033	3.5 (2.5–4.5)	23	3530	27 281	1.4 (0.6–2.1)
GFR 60–89	404	14 003	104 808	3.5 (3.2–3.9)†	192	5433	40 160	4.2 (3.6–4.8)	212	8570	64 648	3.0 (2.6–3.4)†
GFR <60	104	1875	12 092	5.0 (3.7–6.4)‡	33	501	3048	6.6 (3.5–9.7)§	71	1374	9044	4.0 (3.0–5.0)‡
Myocardial infarction												
GFR ≥90	25	8350	60 807	0.6 (0.2–0.9)	21	4179	28 164	0.9 (0.4–1.4)	4	4171	32 643	0.4 (–0.2–0.9)
GFR 60–89	116	19 786	151 527	0.7 (0.6–0.8)	72	7345	54 855	1.1 (0.9–1.4)	44	12 441	96 672	0.4 (0.3–0.5)
GFR <60	39	2521	16 926	1.4 (0.9–1.9)‡	21	643	4039	2.4 (1.3–3.6)‡	18	1878	12 887	0.7 (0.4–1.1)
All-cause death												
GFR ≥90	289	8445	62 754	7.6 (6.4–8.7)	217	4225	29 119	11.4 (9.7–13.1)	72	4220	33 635	5.1 (3.5–6.6)
GFR 60–89	1388	20 280	161 168	7.0 (6.7–7.4)	809	7529	58 344	10.4 (9.7–11.1)	579	12 751	102 824	4.8 (4.4–5.2)
GFR <60	427	2649	18 935	12.9 (10.2–15.5)‡	184	681	4540	21.3 (14.9–27.7)‡	243	1968	14 395	7.3 (5.9–8.6)‡

PY indicates person-years.

*Incidence rates were adjusted for age by the direct standardized method. Overall results were additionally adjusted for sex.

† $P < 0.01$, ‡ $P < 0.001$, § $P < 0.05$ vs GFR ≥90 mL · min⁻¹ · 1.73 m⁻².

lower level. Consequently, more participants (21%) were assigned to the group whose GFR was <60 mL · min⁻¹ · 1.73 m⁻², and the age- and sex-adjusted risk of CVD among these subjects relative to those with a GFR ≥90 mL · min⁻¹ · 1.73 m⁻² was attenuated by 85% (95% CI 32% to 160%), although it was still significant. Similarly, a log-linear relationship between blood pressure levels and the risk of CVD was still observed in the subgroup whose GFR was <60 mL · min⁻¹ · 1.73 m⁻², even after correction with the Japanese coefficient (Data Supplement Figure).

Discussion

In the present study, we demonstrated a clear association between reduced GFR and high risk of CVD. To the best of our knowledge, this is the first overview of this issue in a Japanese community-based longitudinal study. Furthermore, the relationship between blood pressure levels at baseline and CVD risk was found to be strong and continuous, regardless of kidney function status.

There have been few studies showing the association of reduced GFR with an increased risk of CVD or mortality in the general Japanese population.^{6–8} The findings of the Hisayama study revealed that a GFR <60 mL · min⁻¹ · 1.73 m⁻² was a significant risk factor for the development of coronary heart disease in men and of CVD and stroke in women.⁶ In a large cohort study conducted by Irie et al,⁷ reduced GFR was strongly associated with mortality due to CVD or stroke. A report from NIPPON DATA 90 also showed an association between a GFR <30 mL · min⁻¹ · 1.73 m⁻² and a high risk of cardiovascular death.⁸ In the present study, we demonstrated a clear association between reduced GFR and the risks of CVD, stroke, myocardial infarction, and death in an overview of 10 Japanese cohort studies. These

results, therefore, highlight the importance of taking kidney function status into consideration in trying to reduce the burden of CVD in the general Japanese population.

There are several possible explanations for the association of reduced GFR with CVD.³ First, reduced GFR is associated with a high prevalence of traditional CVD risk factors, such as aging, hypertension, diabetes, smoking habits, and dyslipidemia.¹⁹ In the present study, reduced GFR was found to be a significant risk factor for the development of stroke after adjustment for demographic factors, but not after adjustment for potential traditional CVD risk factors, which suggests that an accumulation of traditional CVD risk factors in individuals with reduced GFR increases the risk of stroke. In contrast, the risks of CVD, myocardial infarction, and all-cause death in individuals with reduced GFR were also attenuated, although still significant, after adjustment for traditional CVD risk factors. Reduced GFR has been shown to be associated with increased levels of novel CVD risk factors, such as inflammation, asymmetric dimethylarginine, oxidative stress, and thrombogenic factors.^{19,20} Second, reduced GFR may be a marker of vascular disease; it is well recognized that renal arteriosclerosis and glomerular sclerosis are closely related to systemic atherosclerosis.²¹

In the present study, reduced GFR was associated with a high risk of stroke in men after adjustment for demographic factors but not after adjustment for potential confounding factors; however, this relationship was still observed in women even after adjustment for confounding factors. This sex difference may be a consequence of the effects of residual confounding factors, specifically, hypercoagulable states²² or gonadal steroids,²³ in women. Furthermore, the lack of a significant association between reduced GFR and a high risk of myocardial infarction is probably due to the relatively small number of events.

Table 4. Effects of Kidney Function on Development of CVD

	Age- and Sex-Adjusted*		Multivariate-Adjusted†	
	HR (95% CI)	P	HR (95% CI)	P
Overall				
CVD				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.41 (1.13–1.75)	0.002	1.24 (0.98–1.58)	0.07
GFR <60	2.26 (1.71–2.99)	<0.001	1.57 (1.14–2.15)	0.005
Stroke				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.40 (1.10–1.79)	0.007	1.24 (0.95–1.61)	0.11
GFR <60	2.06 (1.51–2.81)	<0.001	1.41 (0.99–2.00)	0.06
Myocardial infarction				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.32 (0.84–2.08)	0.22	1.26 (0.77–2.05)	0.35
GFR <60	3.35 (1.94–5.79)	<0.001	2.37 (1.29–4.34)	0.005
All-cause death				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.01 (0.88–1.15)	0.94	1.10 (0.96–1.27)	0.17
GFR <60	1.70 (1.44–2.00)	<0.001	1.65 (1.38–1.97)	<0.001
Men				
CVD				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.21 (0.93–1.58)	0.16	1.01 (0.75–1.35)	0.95
GFR <60	2.13 (1.45–3.11)	<0.001	1.47 (0.94–2.29)	0.09
Stroke				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.17 (0.86–1.57)	0.32	0.99 (0.71–1.38)	0.95
GFR <60	1.69 (1.08–2.65)	0.02	1.10 (0.64–1.89)	0.72
Myocardial infarction				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.25 (0.75–2.07)	0.39	1.05 (0.61–1.81)	0.85
GFR <60	3.95 (2.07–7.55)	<0.001	2.56 (1.24–5.27)	0.01
All-cause death				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	0.97 (0.83–1.14)	0.72	1.06 (0.90–1.25)	0.48
GFR <60	1.75 (1.42–2.16)	<0.001	1.73 (1.37–2.17)	<0.001
Women				
CVD				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.93 (1.28–2.92)	0.002	1.81 (1.17–2.79)	0.008
GFR <60	2.84 (1.79–4.52)	<0.001	1.97 (1.19–3.29)	0.009
Stroke				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	2.01 (1.28–3.14)	0.002	1.81 (1.14–2.89)	0.01
GFR <60	2.89 (1.75–4.79)	<0.001	1.98 (1.15–3.42)	0.01
Myocardial infarction				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.60 (0.55–4.60)	0.39	2.14 (0.63–7.24)	0.22
GFR <60	2.93 (0.93–9.23)	0.07	2.79 (0.74–10.56)	0.13
All-cause death				
GFR ≥90	1.00 (Reference)		1.00 (Reference)	
GFR 60–89	1.13 (0.87–1.47)	0.35	1.23 (0.94–1.62)	0.13
GFR <60	1.79 (1.34–2.38)	<0.001	1.68 (1.24–2.30)	<0.001

GFR was measured in mL · min⁻¹ · 1.73 m⁻².

*Sex was removed from model for the analysis stratified by sex.

†Estimates were adjusted for age, sex, cohort, systolic blood pressure, diabetes, serum total cholesterol, body mass index, and current smoking status. Sex was removed from model for the analyses stratified by sex.




Figure. Effects of blood pressure levels on the development of CVD according to kidney function status. Estimates were adjusted for age, sex, cohort, diabetes, serum total cholesterol, body mass index, and current smoking status. Solid boxes represent estimates on the risk of outcomes for each blood pressure level. Areas of the boxes are proportional to the number of events. "P for trend" tested the log-linear relationship between blood pressure levels at baseline and the risk of outcomes by kidney function status. "P for hetero." tested the heterogeneity of the association of blood pressure levels with the risk of outcomes between kidney function status subgroups. HT indicates hypertension; PYs, person-years.

In the present study, we demonstrated a clear log-linear association between blood pressure levels and the risks of CVD, stroke, and all-cause death, regardless of kidney function status. These findings are consistent with the results of other studies conducted in the general population.^{9,10} Recent publications of prospective cohort data suggest, however, that individuals with a reduced GFR and a systolic blood pressure below 120 mm Hg may be at increased risk of stroke or death.^{12,13} Other post hoc analyses of trials conducted on individuals with coronary heart disease²⁴ and with diabetic nephropathy²⁵ suggest an increased risk of coronary events at the lower achieved blood pressures. In the present study, however, no evidence of an increased risk of myocardial infarction was observed at the lower blood pressure level. One possible explanation for the J-curve association observed in the previous studies may be the phenomenon of reverse causality,²⁶ in which extensive vascular disease or subclinical cardiac dysfunction is associated with lower blood pressure levels and reduced GFR and is associated independently with a relatively high risk of CVD, rather than with any adverse effects of low blood pressure itself.

Several limitations of the present study should be noted. First, the generalizability of our findings to some populations at high risk for CVD may be limited. The participants excluded from the analysis due to missing baseline examination data or event data were likely to have a higher cardiovascular risk, because they were older (mean 63 years), had higher blood pressure levels (mean 138/80 mm Hg), and had a greater prevalence of diabetes (8.7%) than the study population. This bias has the potential to alter our findings, which may therefore be conservative. Second, the present GFR estimates, which were made with a simplified prediction equation, may not be sufficiently correct, which possibly could lead to a certain number of misclassifications of estimated kidney function status. Such misclassifications would weaken the association found in the present study, biasing the results toward the null hypothesis. Third, we were unable to obtain information regarding the use of antihypertensive drugs, medication compliance, or blood pressure control during the follow-up period. The lack of this information may reduce the accuracy of our findings to some extent. Fourth, the applicability of the present results to populations with severe kidney dysfunction is limited, because very few of our subjects (0.1%) had a GFR $<30 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$. Moreover, the absence of data on proteinuria in the present study makes it impossible to assess the effects of the earliest stages of kidney disease on the risk of CVD. Finally, creatinine measurement was conducted locally rather than at a central laboratory, which introduces a certain amount of variability that may reduce the reliability of the results.

In conclusion, the present findings suggest that a reduced GFR is associated significantly with a high risk of CVD in the general Japanese population. Furthermore, we observed a continuous relationship between blood pressure levels at baseline and the risk of CVD, regardless of kidney function status. The optimization of blood pressure control in individuals with kidney dysfunction is therefore likely to substantially reduce the burden of CVD in the general population.

Acknowledgments

The authors gratefully acknowledge the efforts of the JALS investigators, research coordinators, and committee members. A list of these individuals has been published previously¹⁴ and is provided in the Appendix (Data Supplement).

Sources of Funding

This study was supported by a research grant from Japan Arteriosclerosis Prevention Fund.

Disclosures

None.

References

- Chadban SJ, Briganti EM, Kerr PG, Dunstan DW, Welborn TA, Zimmet PZ, Atkins RC. Prevalence of kidney damage in Australian adults: the AusDiab kidney study. *J Am Soc Nephrol*. 2003;14(suppl 2):S131–S138.
- Coresh J, Astor BC, Greene T, Eknoyan G, Levey AS. Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey. *Am J Kidney Dis*. 2003;41:1–12.
- Sarnak MJ, Levey AS, Schoolwerth AC, Coresh J, Culleton B, Hamm LL, McCullough PA, Kasiske BL, Kelepouris E, Klag MJ, Parfrey P, Pfeffer M, Raij L, Spinosa DJ, Wilson PW; American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. Kidney disease as a risk factor for development of cardiovascular disease: a statement from the American Heart Association Councils on Kidney in Cardiovascular Disease, High Blood Pressure Research, Clinical Cardiology, and Epidemiology and Prevention. *Circulation*. 2003;108:2154–2169.
- Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med*. 2004;351:1296–1305.
- Manjunath G, Tighiouart H, Ibrahim H, MacLeod B, Salem DN, Griffith JL, Coresh J, Levey AS, Sarnak MJ. Level of kidney function as a risk factor for atherosclerotic cardiovascular outcomes in the community. *J Am Coll Cardiol*. 2003;41:47–55.
- Ninomiya T, Kiyohara Y, Kubo M, Tanizaki Y, Doi Y, Okubo K, Wakugawa Y, Hata J, Oishi Y, Shikata K, Yonemoto K, Hirakata H, Iida M. Chronic kidney disease and cardiovascular disease in a general Japanese population: the Hisayama Study. *Kidney Int*. 2005;68:228–236.
- Irie F, Iso H, Sairenchi T, Fukasawa N, Yamagishi K, Ikehara S, Kanashiki M, Saito Y, Ota H, Nose T. The relationships of proteinuria, serum creatinine, glomerular filtration rate with cardiovascular disease mortality in Japanese general population. *Kidney Int*. 2006;69:1264–1271.
- Nakamura K, Okamura T, Hayakawa T, Kadowaki T, Kita Y, Ohnishi H, Saitoh S, Sakata K, Okayama A, Ueshima H; the NIPPON DATA90 Research Group. Chronic kidney disease is a risk factor for cardiovascular death in a community-based population in Japan: NIPPON DATA90. *Circ J*. 2006;70:954–959.
- MacMahon S, Peto R, Cutler J, Collins R, Sorlie P, Neaton J, Abbott R, Godwin J, Dyer A, Stamler J. Blood pressure, stroke, and coronary heart disease: part 1, prolonged differences in blood pressure: prospective observational studies corrected for the regression dilution bias. *Lancet*. 1990;335:765–774.
- Lawes CM, Rodgers A, Bennett DA, Parag V, Suh I, Ueshima H, MacMahon S; Asia Pacific Cohort Studies Collaboration. Blood pressure and cardiovascular disease in the Asia Pacific region. *J Hypertens*. 2003;21:707–716.
- Blood Pressure Lowering Treatment Trialists' Collaboration. Effects of different blood-pressure-lowering regimens on major cardiovascular events: results of prospectively-designed overviews of randomised trials. *Lancet*. 2003;362:1527–1535.
- Weiner DE, Tighiouart H, Levey AS, Elsayed E, Griffith JL, Salem DN, Sarnak MJ. Lowest systolic blood pressure is associated with stroke in stages 3 to 4 chronic kidney disease. *J Am Soc Nephrol*. 2007;18:960–966.
- Kovesdy CP, Trivedi BK, Kalantar-Zadeh K, Anderson JE. Association of low blood pressure with increased mortality in patients with moderate to severe chronic kidney disease. *Nephrol Dial Transplant*. 2006;21:1257–1262.

14. Japan Arteriosclerosis Longitudinal Study Group. Japan Arteriosclerosis Longitudinal Study-ECC: rationale, design, and population characteristics. *Circ J*. 2008;72:1563–1568.
15. Imai F, Horio M, Nitta K, Yamagata K, Iseki K, Hara S, Ura N, Kiyohara Y, Hirakata H, Watanabe T, Moriyama T, Ando Y, Inaguma D, Narita I, Iso H, Wakai K, Yasuda Y, Tsukamoto Y, Ito S, Makino H, Hishida A, Matsuo S. Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. *Clin Exp Nephrol*. 2007;11:41–50.
16. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis*. 2002;39(suppl 1):S1–S266.
17. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Roccella EJ; Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. National Heart, Lung, and Blood Institute; National High Blood Pressure Education Program Coordinating Committee: seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 2003;42:1206–1252.
18. Woodward M. *Epidemiology: Study Design and Data Analysis*. 2nd ed. Boca Raton, Fla: Chapman & Hall/CRC Press; 2005.
19. Uhlig K, Levey AS, Sarnak MJ. Traditional cardiac risk factors in individuals with chronic kidney disease. *Semin Dial*. 2003;16:118–127.
20. Madore F. Uremia-related metabolic cardiac risk factors in chronic kidney disease. *Semin Dial*. 2003;16:148–156.
21. Keane WF, Kasiske BL, O'Donnell MP. Lipids and progressive glomerulosclerosis: a model analogous to atherosclerosis. *Am J Nephrol*. 1988;8:261–271.
22. Tracy RP, Arnold AM, Ettinger W, Fried L, Meilahn E, Savage P. The relationship of fibrinogen and factors VII and VIII to incident cardiovascular disease and death in the elderly: results from the cardiovascular health study. *Arterioscler Thromb Vasc Biol*. 1999;19:1776–1783.
23. Gabriel SR, Carmona L, Roque M, Sánchez GL, Bonfill X. Hormone replacement therapy for preventing cardiovascular disease in postmenopausal women. *Cochrane Database Syst Rev*. 2005;2:CD002229.
24. Messerli FH, Mancia G, Conti CR, Hewkin AC, Kupfer S, Champion A, Kolloch R, Benetos A, Pepine CJ. Dogma disputed: can aggressively lowering blood pressure in hypertensive patients with coronary artery disease be dangerous? *Ann Intern Med*. 2006;144:884–893.
25. Berl T, Hunsicker LG, Lewis JB, Pfeffer MA, Porush JG, Rouleau JL, Drury PL, Fismatjes E, Hricik D, Pohl M, Raz I, Vanhille P, Wiegmann TB, Wolfe BM, Locatelli F, Goldhaber SZ, Lewis EJ; the Collaborative Study Group. Impact of achieved blood pressure on cardiovascular outcomes in the Irbesartan Diabetic Nephropathy Trial. *J Am Soc Nephrol*. 2005;16:2170–2179.
26. Kalantar-Zadeh K, Block G, Humphreys MH, Kopple JD. Reverse epidemiology of cardiovascular risk factors in maintenance dialysis patients. *Kidney Int*. 2003;63:793–808.

CLINICAL PERSPECTIVE

There have been several studies reporting a strong association between reduced kidney function and cardiovascular risk. The findings, however, have been inconsistent in Asian populations, and there has been no attempt to date to review the evidence. Hence, we conducted an overview of individual participant data from Japanese community-based cohort studies to reliably assess the impact of reduced kidney function on cardiovascular risk in the general Japanese population. Our findings suggest a clear association between reduced kidney function and a 57% greater risk of cardiovascular disease in the Japanese population, as well as a log-linear relationship between blood pressure levels and cardiovascular risk in individuals with reduced kidney function. The optimization of blood pressure control in individuals with reduced kidney function is therefore likely to substantially reduce the burden of cardiovascular disease in the general population. Given that the prevalence of reduced kidney function is $\approx 10\%$ in the general population, we believe that these novel findings are significant in the areas of clinical and public health.

Go to <http://cme.ahajournals.org> to take the CME quiz for this article.

Impact of Kidney Disease and Blood Pressure on the Development of Cardiovascular Disease

An Overview From the Japan Arteriosclerosis Longitudinal Study

Toshiharu Ninomiya, MD, PhD; Yutaka Kiyohara, MD, PhD; Yosuke Tokuda, MS;
Yasufumi Doi, MD, PhD; Hisatomi Arima, MD, PhD; Akiko Harada, PhD;
Yasuo Ohashi, PhD; Hirotsugu Ueshima, MD, PhD;
for the Japan Arteriosclerosis Longitudinal Study Group

Background—Kidney disease is associated with an increased risk of cardiovascular disease (CVD); however, there have been few well-designed prospective studies of this issue in Asian populations. Recent epidemiological studies have suggested that a lower blood pressure level may be associated with an increased risk of CVD in individuals with kidney dysfunction.

Methods and Results—Using data from 10 community-based cohort studies in Japan, we conducted follow-up on a total of 30 657 individuals 40 to 89 years of age without preexisting CVD or kidney failure and examined the relationship between reduced glomerular filtration rate (GFR) and the risk of CVD. During an average 7.4-year follow-up, 727 individuals experienced CVD. The age- and sex-adjusted incidence of CVD increased significantly in subjects with GFR of 60 to 89 mL · min⁻¹ · 1.73 m⁻² (4.3 per 1000 person-years, *P*=0.002) and in those with a GFR <60 mL · min⁻¹ · 1.73 m⁻² (6.5, *P*<0.001) compared with those with a GFR ≥90 mL · min⁻¹ · 1.73 m⁻² (2.9). Even after adjustment for potential confounding factors, subjects with a GFR <60 mL · min⁻¹ · 1.73 m⁻² had a 57% (95% CI 14% to 115%) greater risk of CVD than those with a GFR ≥90 mL · min⁻¹ · 1.73 m⁻². The multivariate-adjusted hazard ratios of CVD increased in a log-linear manner with elevations in blood pressure levels, regardless of GFR levels (all *P* for trend <0.01).

Conclusions—Our findings suggest that a reduced GFR is a significant risk factor for CVD in the general Japanese population. Additionally, a log-linear association of blood pressure level with CVD risk was observed, without evidence of a J-curve association, regardless of GFR levels. (*Circulation*. 2008;118:2694-2701.)

Key Words: cardiovascular diseases ■ blood pressure ■ kidney ■ meta-analysis

Kidney disease is increasingly being recognized as a leading public health issue. Chronic kidney disease, most commonly defined by a reduction in glomerular filtration rate (GFR) or the presence of proteinuria, affects 10% to 15% of the adult population in Western countries^{1,2} and is associated with an increased risk of cardiovascular disease (CVD)³⁻⁵; however, there have been few well-designed large prospective studies in general Asian communities to date.⁶⁻⁸

Clinical Perspective p 2701

Blood pressure is an important determinant of the risk of CVD in the general population,^{9,10} in which it has been well established that treatment for high blood pressure prevents CVD.¹¹ Blood pressure is commonly elevated in individuals

with a reduced GFR,^{4,5} which suggests that lowering blood pressure may offer significant benefits in this population. Recent prospective cohort studies, however, have reported that the risk of stroke or death for individuals with a reduced GFR is greater among those with systolic blood pressure levels below 120 mm Hg than among those with higher levels.^{12,13} These data have raised concerns that lowering blood pressure may provide less benefit than previously believed, or may even be hazardous, in individuals with kidney dysfunction.

In the present report, we discuss the results of a pooling analysis from the Japan Arteriosclerosis Longitudinal Study—Existing Cohorts Combine (JALS-ECC), which is an overview of individual participant data from 21 community-based

Continuing medical education (CME) credit is available for this article. Go to <http://cme.ahajournals.org> to take the quiz.

From the Department of Environmental Medicine (T.N., Y.K., H.A.) and Department of Medicine and Clinical Science (Y.D.), Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan; Department of Biostatistics/Epidemiology and Preventive Health Sciences (Y.T., A.H., Y.O.), School of Health Sciences and Nursing, University of Tokyo, Tokyo, Japan; and Department of Health Science (H.U.), Shiga University of Medical Science, Otsu, Japan.

The online-only Data Supplement is available with this article at <http://circ.ahajournals.org/cgi/content/full/118/25/2694/DC1>.

Correspondence to Yutaka Kiyohara, MD, PhD, Department of Environmental Medicine, Graduate School of Medical Sciences, Kyushu University, Maidashi 3-1-1, Higashi-ku, Fukuoka, 812-8582 Japan. E-mail kiyohara@envmed.med.kyushu-u.ac.jp

© 2008 American Heart Association, Inc.

Circulation is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.108.792903

Downloaded from circ.ahajournals.org at KYUSHU UNIVERSITY on February 3, 2009

longitudinal observational studies in Japan.¹⁴ Our aims were to assess the impact of a reduction in GFR on the development of CVD in the general population and to examine the association of blood pressure with the risk of CVD in individuals with a reduced GFR.

Methods

Study Population

The rationale, study design, and methods of the JALS-ECC have been described elsewhere.¹⁴ In brief, cohort studies were eligible for inclusion in this project if they satisfied the following criteria: (1) Japanese population; (2) prospective cohort study; (3) at least 3000 person-years of follow-up; (4) date of birth, sex, height, weight, blood pressure, and serum total cholesterol recorded at baseline; and (5) date of death or the age at death recorded during a follow-up. Quality control of the collected cohort data were performed at the JALS Coordinating Center. The individual records of 66 691 participants in 21 cohort studies were included in the present project, with 82.7% of the participants from 17 community-based cohorts and 17.3% from 4 work-site-based cohorts. Permission to submit each collection of cohort data to the JALS Coordinating Center was obtained from the relevant institutional review boards for ethical issues.

Of the 21 cohort studies, 11 cohorts were excluded from the present analysis for the following reasons: 4 were work-site-based cohorts, 3 did not include creatinine data, 3 lacked many values for relevant variables, and 1 included no event data for either stroke or myocardial infarction. From the remaining 10 cohorts, we excluded participants less than 40 years of age and those 90 years of age or older, those with unavailable examination data at baseline or unavailable event data, those with a history of CVD, and those with an estimated GFR $<15 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$. For the analysis regarding CVD and stroke, the Niigata cohort was excluded, because information on stroke events was unavailable. A final total of 23 033 participants were enrolled in the CVD analysis, 23 084 in the stroke analysis, 30 657 in the myocardial infarction analysis, and 31 374 in the all-cause death analysis. The average follow-up period was 7.4 years.

Risk Factors

The JALS Coordinating Center requested individual participant data from the collaborating investigators. Serum creatinine was measured by Jaffe's method in 8 cohorts, by the enzymatic method in 1 cohort, and by both methods in 1 cohort. Serum creatinine values measured by the enzymatic method were corrected for Japanese subjects by the addition of $18.3 \mu\text{mol/L}$.¹⁵ GFR was estimated with the 4-variable Modification of Diet in Renal Disease study equation.³ In accordance with the National Kidney Foundation Kidney Disease Outcomes Quality Initiative guidelines,¹⁶ GFR levels were classified in the following ranges: ≥ 90 , 60 to 89, and $<60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$. Blood pressure was measured by a standard sphygmomanometer in all cohorts. Mean values were used in several cohorts that measured 2 or more blood pressure values. Blood pressure levels at baseline were classified into 4 categories (normal, prehypertension, stage 1 hypertension, and stage 2 hypertension) according to the criteria of the seventh report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure.¹⁷ Diabetes was defined as a fasting blood glucose level of $\geq 7.0 \text{ mmol/L}$, a casual blood glucose level of $\geq 11.1 \text{ mmol/L}$, current use of insulin or oral medication for diabetes, and/or a history of diabetes. Serum total cholesterol was determined enzymatically. Information on smoking habit was obtained through a standard questionnaire and classified as current habitual use or lack thereof.

End Points

In each cohort, vital status and the development of CVD were ascertained during follow-up by use of death certificates, hospital medical records, and/or questionnaire surveys. All outcomes were

classified according to the International Classification of Diseases, 9th Revision (ICD-9). All events were recorded by coordinating center staff members. CVD was defined as the development of either stroke or myocardial infarction. Stroke was defined as an acute disturbance of focal neurological function with symptoms that lasted >24 hours or death caused by a stroke event (ICD-9 codes 430, 431, 433, 434, or 436). Myocardial infarction included both fatal and nonfatal myocardial infarction, which was diagnosed by use of an appropriate clinical history supported by ECG changes and/or elevations of cardiac enzymes or other biochemical markers of myocardial injury (ICD-9 410). Only the first event of the relevant outcome type was included in each analysis.

Statistical Analysis

The SAS software package for Windows, release 9.13 (SAS Institute, Inc, Cary, NC) was used to perform all statistical analyses. The incidence rate of each outcome for the GFR subgroups was calculated by the person-year method and adjusted for the age and sex distribution of the overall population enrolled in the CVD analysis by the direct method, in which the subgroups and study population were subdivided into the same set of age groups (defined by decade) and the age- and sex-specific incidence rates were calculated within each subgroup.¹⁸ The hazard ratios (HRs) and their 95% CIs for the development of events were estimated with the Cox proportional hazards regression model. The cohort effect was adjusted as a fixed effect by taking the study as a strata variable, assuming only proportional hazards within each study and not between studies.¹⁸ Heterogeneity across cohorts was examined with the Cochran Q test and the I^2 statistic.¹⁸ The risks of events according to blood pressure levels were also estimated with the Cox regression model. Trends in relationship between blood pressure levels and the risk of events were assessed by fitting models with a linear term for blood pressure categories according to kidney function status, and the heterogeneity of these relationships between kidney function status subgroups was estimated by the addition of an interaction term of a linear term for blood pressure levels and kidney function status to the relevant model. $P < 0.05$ was considered statistically significant in all analyses.

The authors had full access to and take full responsibility for the integrity of the data. All authors have read and agree to the manuscript as written.

Results

The characteristics of the 10 cohorts examined in the present study are shown in Table 1. Among all subjects, the mean age was 57.6 years, and the proportion of men was 38.0%. The mean value of serum creatinine was $78.6 \mu\text{mol/L}$, and the frequency of GFR $<60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ was 8.2%. During the average follow-up period of 7.4 years, a total of 727 subjects experienced CVD, 592 had strokes, 180 had myocardial infarctions, and a total of 2104 died.

Table 2 shows the baseline characteristics of the 23 033 subjects enrolled in the CVD analysis by sex. Their mean ages were 56.9 years for men and 58.2 years for women, and the frequency of GFR $<60 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ was 5.2% for men and 10.1% for women. The frequencies of normal blood pressure, prehypertension, stage 1 hypertension, and stage 2 hypertension were 19.6%, 41.8%, 26.0%, and 12.6% for men and 24.3%, 41.0%, 24.3%, and 10.4% for women, respectively. Similar findings were observed in subjects enrolled in the analyses of stroke, myocardial infarction, and all-cause death.

The age- and sex-adjusted incidences of CVD and stroke increased with declining GFR levels in the overall population (Table 3); the differences were statistically significant between subjects with a GFR $\geq 90 \text{ mL} \cdot \text{min}^{-1} \cdot 1.73 \text{ m}^{-2}$ and