

		Motif Ia		Motif Ib	
		*		*	
<i>E. faecalis</i>	pMG2200	(22)	-FKEYLDYMERSE EA TRNEH	(66)	-----FSTYN-DYMSNPKK
<i>E. faecium</i>	pHT β	(32)	ELQKFVDYISRQ EA IRQDK	(82)	IKDLREMDKYI-DYMTTRKKA
<i>L. innocua</i>	pLI100	(18)	YWSNYIKYIDR EA VRNEH	(63)	EKE-RN-KKYI-DYMGPNPKK
<i>S. agalactiae</i>	NEM316	(18)	ANPQYVDY NR EA VA KIDE	(66)	QLNFER---YI-DYMNRSYA
<i>S. epidermidis</i>	ATCC12228	(15)	KFKGYLKYINDEKSNKANH	(54)	NLNLNSYSSYIIGYMKNNSI
<i>B. anthracis</i>	pXO2-84		-----	(64)	FNTTDDFEKTV-SYMGKRYA
<i>C. perfringens</i>	pCP13	(17)	KKFNFDYIDR EA TRKKN		-----
Consensus			Y EA		YI YM
		Motif II			
		*			
<i>E. faecalis</i>	pMG2200	(97)	YMKEYFEMAQENK S PLW Q LV F S F RNEWLIENNYLD P ET N Q L K T O		
<i>E. faecium</i>	pHT β	(128)	KIKESVILAK N NG S V M F Q D V IS F D N D L V L REGY N P E T N E L N E N		
<i>L. innocua</i>	pLI100	(97)	KYKDA F Q A E K N N S V M F Q H V I S F D N E L A E Q L Y D P K I G M L D E K		
<i>S. agalactiae</i>	NEM316	(109)	K N K L E--S A Y Q NG S L L W Q G V I S F D N A F L A E Q L Y D V A T G Q V D Q K		
<i>S. epidermidis</i>	ATCC12228	(106)	K L K D D F D A E K Q G C I N Y Q D I I S F D N D L I K N H L Y D A K T D E L N D		
<i>B. anthracis</i>	pXO2-84	(144)	E I K E L V G K A Q N K G S V V Y Q D V I S F D T D F L I E Q L Y D P V T D I L D E N		
<i>C. perfringens</i>	pCP13	(70)	K I K K D F D K A Q I N G S N M W Q V F S F D N F L A N G L Y D S R N G A L D E F		
Consensus			K K A G S Q V I S F D N F L Y D I E		
		Motif III			
		* * *			
<i>E. faecalis</i>	pMG2200	(149)	A I A-E L E K -K E L K G--E W T G A V H Y N T D N I H V H V G V E K N P T R E W I F Y K H		
<i>E. faecium</i>	pHT β	(180)	M G M G M Q E K -E E L V D P -- I W F A T I H R N T E H I H I H V T A M E R K N T R E I N E Y D G		
<i>L. innocua</i>	pLI100	(149)	S M K S F L R K -E G M E G S A- I W L A A I H K N T K H F H V H I S V T E P T P T R K F S N K R		
<i>S. agalactiae</i>	NEM316	(159)	M M P T T L Q K -E G L S D S A- F W G N I H I N T D N I H I F G L S E V E S N R E K I F Y Q P		
<i>S. epidermidis</i>	ATCC12228	(158)	K M I N R M I K D E N M N P Y Q T R W M A N I H Y D T D N I H I S T E L K N T R K I I T N G		
<i>B. anthracis</i>	pXO2-84	(196)	M M E Q L F -K D E Q I E N N G F W F A S I H R N T E H I H I F G I V E K E N R R K L V E V K V		
<i>C. perfringens</i>	pCP13	(122)	A M E E L S -K R E G F K D L -- T W S A L H Y N T D N I H V I A S V E I N P S R E R G K F K P		
Consensus			M K E W A I H N T I H H E R		

MOB_{MG} family 3His-Motif

WxxxxHxxTxxxHxHxxxxExxxxR

FIG. 4. Comparison of the N-terminal region of the deduced ORF44 protein of pMG2200 with putative relaxases found in sequence databases. The boldface letters indicated the amino acid residues conserved in each protein. The asterisks above the sequences show the key residues, Tyr, Ser, and His₃ (3His), in motifs I, II, and III, respectively. There are two motif III candidates (motifs Ia and Ib) in the most of the proteins. The GenBank accession numbers of the putative relaxases are as follows: for pXO2, NZ_ABJC1000063.1; for pCP13, NC_003042.1; for ATCC 12228, NC_004461.1; for NEM316, NC_004368.1; for pLI100, NC_003383.1; for pHT β , NC_007594.1.

insertion mutants (i.e., pAM7314, pAM7330, and pAM2125) of the 3' noncoding region in the E region (52). The 3' terminal border of the E region is mapped by the Tn917-lac insertion of pAM2125 and is located 371 bp from the stop codon of *traE1* (unpublished data). Further analyses of pMG2200 might provide clues that will allow elucidation of the regulation of the pheromone-responsive plasmids.

Concluding comments. The two pheromone-responsive conjugative plasmids pMG2200 (106.5 kbp) and pMG2201 (65.7 kbp) were isolated from VanB2-type *E. faecalis* isolates. This report describes the first case of the isolation and characterization of pheromone-responsive conjugative plasmid pMG2200 encoding the *vanB* resistance determinant. pMG2200 encoded vancomycin resistance and bacteriocin and responded to pheromone cCF10, and pMG2201 encoded erythromycin resistance and cytolysin (Hly/Bac) and responded to pheromone cAD1. Our results show that an *E. faecalis* strain can acquire these characteristics and that these characteristics provide a selective advantage for the organism by allowing it to obtain the pheromone-responsive plasmids encoding drug resistance or bacteriocins by conjugation with plasmid-bearing bacteria in patients in the clinical setting. The plasmid also conferred the cytolysin (Hly/Bac) function for pathogenesis (38, 39). The complete

nucleotide sequence of pMG2200 showed that pMG2200 consists of five major segments: (i) conjugative transposon Tn1549-like elements (33,812 bp) encoding the *vanB2*-type determinant, (ii) genes that regulate the pheromone response of pheromone-responsive plasmids, (iii) genes for UV resistance, (iv) the bacteriocin determinant, and (v) the origin of plasmid transfer. The genes corresponding to the pheromone-responsive regulatory genes, with the exception of the positive regulator *traE1* of plasmid pAD1, showed high levels of homology (100% amino acid identity) to those of pCF10. The data indicated that pMG2200 is a new type of pheromone-responsive plasmid which is a naturally occurring chimeric plasmid with regard to the negative regulatory gene *prgX* (*prgQ*) of pCF10 and the positive regulatory gene *traE1* of pAD1, resulting in a *prgX-prgQ-traE1* genetic organization. Using the chimeric plasmid, we showed that *traE1* is *cis* acting. The nucleotide sequence of the plasmid origin showed a high level of homology to a region within plasmid pAM α 1 of *E. faecalis* that is unrelated to the *oriT* region of pAM α 1, and the ORF corresponding to the putative relaxase showed homology to that of *E. faecium* conjugative plasmid pHT β (61, 64). These results indicate that the diversity within the genetic organization of housekeeping genes, such as the regulatory regions, origin of transfer, and plasmid replication in the pheromone-

responsive plasmids, could result from genetic recombination between different pheromone-responsive plasmids or between a pheromone-responsive and a non-pheromone-responsive plasmid.

To our knowledge, there has been only one report on sequence analysis of the conjugative transposon Tn1549 encoding the VanB gene cluster (30). The conjugative transposon Tn1549-like element encodes a vanB2-type resistance determinant that is almost completely identical to the Tn1546 transposon residing on pIP834 of *E. faecalis* E93/268 (30). There has been no report to date of a pheromone-responsive highly conjugative plasmid carrying the Tn1549-like element encoding the VanB gene cluster. Our report is the first to describe a pheromone-responsive plasmid carrying the Tn1549-like element encoding the VanB2 gene cluster.

ACKNOWLEDGMENTS

This work was supported by grants from the Japanese Ministry of Education, Culture, Sport, Science and Technology (Tokutei-ryoiki [Matrix of Infection Phenomena], Kiban [B], Kiban [C]) and the Japanese Ministry of Health, Labor and Welfare (H18-Shinko-11).

We thank E. Kamei for helpful advice on the manuscript.

REFERENCES

- Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local alignment search tool. *J. Mol. Biol.* 215:403-410.
- Arthur, M., C. Molinas, F. Depardieu, and P. Courvalin. 1993. Characterization of Tn1546, a Tn3-related transposon conferring glycopeptide resistance by synthesis of decapeptide peptidoglycan precursors in *Enterococcus faecium* BM4147. *J. Bacteriol.* 175:117-127.
- Carias, L. L., S. D. Rudin, C. J. Donskey, and L. B. Rice. 1998. Genetic linkage and cotransfer of a novel, vanB-containing transposon (Tn5382) and a low-affinity penicillin-binding protein 5 gene in a clinical vancomycin-resistant *Enterococcus faecium* isolate. *J. Bacteriol.* 180:4426-4434.
- Cetinikaya, Y., P. Falk, and C. G. Mayhall. 2000. Vancomycin-resistant enterococci. *Clin. Microbiol. Rev.* 13:686-707.
- Christie, P. J., and G. M. Dunny. 1986. Identification of regions of the *Streptococcus faecalis* plasmid pCF-10 that encode antibiotic resistance and pheromone response functions. *Plasmid* 15:230-241.
- Chung, J. W., and G. M. Dunny. 1992. Cis-acting, orientation-dependent, positive control system activates pheromone-inducible conjugation functions at distances greater than 10 kilobases upstream from its target in *Enterococcus faecalis*. *Proc. Natl. Acad. Sci. USA* 89:9020-9024.
- Clewell, D. B. 1981. Plasmids, drug resistance, and gene transfer in the genus *Streptococcus*. *Microbiol. Rev.* 45:409-436.
- Clewell, D. B. 2007. Properties of *Enterococcus faecalis* plasmid pAD1, a member of a widely disseminated family of pheromone-responding, conjugative, virulence elements encoding cytolysin. *Plasmid* 58:205-227.
- Clewell, D. B., and G. M. Dunny. 2002. Conjugation and genetic exchange in enterococci, p. 265-300. In M. S. Gilmore et al. (ed.), *The enterococci: pathogenesis, molecular biology, and antibiotic resistance*. American Society for Microbiology, Washington, DC.
- Clewell, D. B., S. E. Flannagan, and D. D. Jaworski. 1995. Unconstrained bacterial promiscuity: the Tn916-Tn1545 family of conjugative transposons. *Trends Microbiol.* 3:229-236.
- Clewell, D. B., P. K. Tomich, M. C. Gawron-Burke, A. E. Franke, Y. Yagi, and F. Y. An. 1982. Mapping of *Streptococcus faecalis* plasmids pAD1 and pAD2 and studies relating to transposition of Tn917. *J. Bacteriol.* 152:1220-1230.
- Courvalin, P. 2006. Vancomycin-resistance in gram-positive cocci. *Clin. Infect. Dis.* 42:525-534.
- Dahl, K. H., G. S. Simonsen, O. Olsvik, and A. Sundsfjord. 1999. Heterogeneity in the vanB gene cluster of genomically diverse clinical strains of vancomycin-resistant enterococci. *Antimicrob. Agents Chemother.* 43:1105-1110.
- De Boever, E. H., D. B. Clewell, and C. M. Francer. 2000. *Enterococcus faecalis* conjugative plasmid pAM373: complete nucleotide sequence and genetic analyses of sex pheromone response. *Mol. Microbiol.* 37:1327-1341.
- Diekema, D. J., B. J. Boots Miller, T. E. Vaughn, R. F. Woolson, J. W. Vankey, E. J. Ernst, S. D. Flach, M. M. Ward, C. L. J. Francis, M. A. Pfleger, and B. N. Doebberling. 2004. Antimicrobial resistance trends and outbreak frequency in United States hospitals. *Clin. Infect. Dis.* 38:78-85.
- Dunny, G. M., B. L. Brown, and D. B. Clewell. 1978. Induced cell aggregation and mating in *Streptococcus faecalis*: evidence for a bacterial sex pheromone. *Proc. Natl. Acad. Sci. USA* 75:3479-3483.
- Dunny, G. M., and D. B. Clewell. 1975. Transmissible toxin (hemolysin) plasmid in *Streptococcus faecalis* and its mobilization of noninfectious drug resistance plasmid. *J. Bacteriol.* 124:784-790.
- Dunny, G. M., R. A. Craig, R. Carron, and D. B. Clewell. 1979. Plasmid transfer in *Streptococcus faecalis*. Production of multiple sex pheromones by recipients. *Plasmid* 2:454-465.
- Dunny, G. M., and B. A. Leonard. 1997. Cell-cell communication in gram-positive bacteria. *Annu. Rev. Microbiol.* 51:527-564.
- Eaton, T. J., and M. J. Gasson. 2001. Molecular screening of *Enterococcus* virulence determinants and potential for genetic exchange between food and medical isolates. *Appl. Environ. Microbiol.* 67:1628-1635.
- Ehrenfeld, E. E., and D. B. Clewell. 1987. Transfer functions of the *Streptococcus faecalis* plasmid pAD1: organization of plasmid DNA encoding response to sex pheromone. *J. Bacteriol.* 169:3473-3481.
- Evers, S., and P. Courvalin. 1996. Regulation of VanB-type vancomycin resistance gene expression by VanS_{II}-VanS_{III}, two-component regulatory system in *Enterococcus faecalis* V583. *J. Bacteriol.* 178:1302-1309.
- Franca, M. V., and D. B. Clewell. 2002. Amplification of the tetracycline resistance determinant of pAMo1 in *Enterococcus faecalis* requires a site-specific recombination event involving relaxase. *J. Bacteriol.* 184:5187-5193.
- Franca, M. V., W. Haas, R. Wirth, E. Samberger, A. Mischoll-Silberhorn, M. S. Gilmore, Y. Ike, K. E. Weaver, F. Y. An, and D. B. Clewell. 2001. Completion of the nucleotide sequence of the *Enterococcus faecalis* conjugative virulence plasmid pAD1 and identification of a second transfer origin. *Plasmid* 46:117-127.
- Franca, M. V., A. Varsaki, M. P. Garcillan-Barcia, A. Latorre, C. Drinas, and F. de la Cruz. 2004. A classification scheme for mobilization regions of bacterial plasmids. *FEMS Microbiol. Rev.* 28:79-100.
- Franke, A. E., and D. B. Clewell. 1981. Evidence for a chromosome-borne resistance transposon (Tn916) in *Streptococcus faecalis* that is capable of "conjugal" transfer in the absence of a conjugative plasmid. *J. Bacteriol.* 145:494-502.
- Fujimoto, S., H. Hashimoto, and Y. Ike. 1991. Low cost device for electrotransformation and its application to the highly efficient transformation of *Escherichia coli* and *Enterococcus faecalis*. *Plasmid* 26:131-135.
- Fujimoto, S., H. Tomita, E. Wakamatsu, K. Tanimoto, and Y. Ike. 1995. Physical mapping of the conjugative bacteriocin plasmid pPD1 of *Enterococcus faecalis* and identification of the determinant related to the pheromone response. *J. Bacteriol.* 177:5574-5581.
- Fujita, N., M. Yoshimura, T. Komori, K. Tanimoto, and Y. Ike. 1998. First report of the isolation of high-level vancomycin-resistant *Enterococcus faecium* from a patient in Japan. *Antimicrob. Agents Chemother.* 42:2150.
- Garnier, F., S. Taouit, P. Glaser, P. Courvalin, and M. Galmann. 2000. Characterization of transposon Tn1549, conferring VanB-type resistance in *Enterococcus* spp. *Microbiology* 146:1481-1489.
- Gilmore, M. S., R. A. Segarra, M. C. Booth, C. P. Bogie, L. R. Hall, and D. B. Clewell. 1994. Genetic structure of the *Enterococcus faecalis* plasmid pAD1-encoded cytolysin toxin system and its relationship to lantibiotic determinants. *J. Bacteriol.* 176:7335-7344.
- Haas, W., B. D. Shepard, and M. S. Gilmore. 2002. Two-component regulator of *Enterococcus faecalis* cytolysin responds to quorum-sensing autoinduction. *Nature* 415:84-87.
- Hirt, H., D. A. Manias, E. M. Bryan, J. R. Klein, J. K. Marklund, J. H. Staddon, M. L. Paustian, W. Kapur, and G. M. Dunny. 2005. Characterization of the pheromone response of the *Enterococcus faecalis* conjugative plasmid pCF10: complete sequence and comparative analysis of the transcriptional and phenotypic responses of pCF10-containing cells to pheromone induction. *J. Bacteriol.* 187:1044-1054.
- Ike, Y., and D. B. Clewell. 1984. Genetic analysis of the pAD1 pheromone response in *Streptococcus faecalis*, using transposon Tn917 as an insertional mutagen. *J. Bacteriol.* 158:777-783.
- Ike, Y., and D. B. Clewell. 1992. Evidence that the hemolysin/bacteriocin phenotype of *Enterococcus faecalis* subsp. *zymogenes* can be determined by plasmids in different incompatibility groups as well as by the chromosome. *J. Bacteriol.* 174:8172-8177.
- Ike, Y., D. B. Clewell, R. A. Segarra, and M. S. Gilmore. 1990. Genetic analysis of the pAD1 hemolysin/bacteriocin determinant in *Enterococcus faecalis*: Tn917 insertional mutagenesis and cloning. *J. Bacteriol.* 172:155-163.
- Ike, Y., R. C. Craig, B. A. White, Y. Yagi, and D. B. Clewell. 1983. Modification of *Streptococcus faecalis* sex pheromones after acquisition of plasmid DNA. *Proc. Natl. Acad. Sci. USA* 80:5369-5373.
- Ike, Y., H. Hashimoto, and D. B. Clewell. 1984. Hemolysin of *Streptococcus faecalis* subspecies *zymogenes* contributes to virulence in mice. *Infect. Immun.* 45:528-530.
- Ike, Y., H. Hashimoto, and D. B. Clewell. 1987. High incidence of hemolysin production by *Enterococcus (Streptococcus) faecalis* strains associated with human parenteral infection. *J. Clin. Microbiol.* 25:1524-1528.
- Lauderdale, T. L., L. C. McDonald, Y. R. Shiao, P. C. Chen, H. Y. Wang, J. F. Lai, and M. Ho. 2002. Vancomycin-resistant enterococci from humans and retail chickens in Taiwan with unique VanB phenotype-vanA genotype incongruence. *Antimicrob. Agents Chemother.* 46:525-527.

41. Leclercq, R., E. Derlot, J. Duval, and P. Courvalin. 1988. Plasmid-mediated resistance to vancomycin and teicoplanin in *Enterococcus faecium*. N. Engl. J. Med. 319:157-161.
42. Martinez-Bueno, M., E. Valdivia, A. Galvez, and M. Maqueda. 2000. pSR6, a new theta-replicating plasmid from *Enterococcus faecalis*. Curr. Microbiol. 41:257-261.
43. Martone, W. J. 1998. Spread of vancomycin resistant enterococci: why did it happen in the United States? Infect. Control Hosp. Epidemiol. 19:539-545.
44. Morrison, D., N. Woodford, and B. Cookson. 1997. Enterococci as emerging pathogens of humans. J. Appl. Microbiol. 83(Suppl.):895-995.
45. Muscholl, A., D. Galli, G. Wanner, and R. Wirth. 1993. Sex pheromone plasmid pAD1-encoded aggregation substance of *Enterococcus faecalis* is positively regulated in *trans* by *traE1*. Eur. J. Biochem. 214:333-338.
46. Muscholl-Shilberhorn, A. B. 2000. Pheromone-regulated expression of sex pheromone plasmid pAD1-encoded aggregation substance depends on at least six upstream genes and a *cis*-acting, orientation-dependent factor. J. Bacteriol. 182:3816-3825.
47. Oana, K., Y. Kawakami, M. Ohnishi, M. Ishikawa, M. Hirota, M. Tozuka, K. Atarashi, K. Baba, K. Fujiki, M. Okazaki, T. Honda, and T. Hayashi. 2001. Molecular and epidemiological study of the first outbreak of *vanB* type vancomycin-resistant *Enterococcus faecalis* in Japan. Jpn. J. Infect. Dis. 54:17-22.
48. Ozawa, Y., K. Tanimoto, S. Fujimoto, H. Tomita, and Y. Ike. 1997. Cloning and genetic analysis of the UV resistance determinant (*uvr*) encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pAD1. J. Bacteriol. 179:7468-7475.
49. Ozawa, Y., K. Tanimoto, T. Nomura, M. Yoshinaga, Y. Arakawa, and Y. Ike. 2002. Vancomycin-resistant enterococci in humans and imported chickens in Japan. Appl. Environ. Microbiol. 68:6457-6461.
50. Patel, R., J. R. Uhl, P. Kohner, M. K. Hopkins, J. M. Steckelberg, B. Kline, and F. R. Cockerill III. 1998. DNA sequence variation within *vanA*, *vanB*, *vanC-1*, and *vanC-2/3* genes of clinical *Enterococcus* isolates. Antimicrob. Agents Chemother. 42:202-205.
51. Paulsen, I. T., L. Banerjee, G. S. A. Myers, K. E. Nelson, R. Sechadri, T. D. Read, D. E. Fouts, J. A. Eisen, S. R. Gill, J. F. Heidelberg, H. Tettelin, R. J. Dobson, L. Umayam, I. Brinkac, M. Beaman, S. Daugherty, R. T. DeBoy, S. Durkin, J. Kolonay, R. Madupu, W. Nelson, J. Vamathevan, B. Tran, J. Upton, T. Hansen, J. Shetty, H. Khouri, T. Utterback, D. Radunc, K. A. Kethum, B. A. Dougherty, and C. M. Fraser. 2003. Role of mobile DNA in the evolution of vancomycin-resistant *Enterococcus faecalis*. Science 299:2071-2074.
52. Pontius, L. T., and D. B. Clewell. 1992. Conjugative transfer of *Enterococcus faecalis* plasmid pAD1: nucleotide sequence and transcriptional fusion analysis of a region involved in positive regulation. J. Bacteriol. 174:3152-3160.
53. Quintilliani, R., and P. Courvalin. 1994. Conjugative transfer of the vancomycin resistance determinant *vanB* between enterococci involves the movement of large genetic elements from chromosome to chromosome. FEMS Microbiol. Lett. 119:359-364.
54. Rice, L. B., L. L. Carias, C. L. Donskey, and S. D. Rudin. 1998. Transferable, plasmid-dedicated *VanB*-type glycopeptide resistance in *Enterococcus faecium*. Antimicrob. Agents Chemother. 42:963-964.
55. Sahn, D. F., J. Kissinger, M. S. Gilmore, P. R. Murray, R. Mulder, J. Solliday, and B. Clarke. 1989. In vitro susceptibility studies of vancomycin-resistant *Enterococcus faecalis*. Antimicrob. Agents Chemother. 33:1588-1591.
56. Sambrook, J., and D. W. Russell. 2001. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
57. Semedo, T., M. A. Santos, P. Martins, M. F. S. Lopes, J. F. Marques, R. Tenreiro, and M. T. B. Crespo. 2003. Comparative study using type strains and clinical and food isolates to examine hemolytic activity and occurrence of the *cyt* operon in enterococci. J. Clin. Microbiol. 41:2569-2576.
58. Shiojima, M., H. Tomita, K. Tanimoto, S. Fujimoto, and Y. Ike. 1997. High-level plasmid-mediated gentamicin resistance and pheromone response of plasmids present in clinical isolates of *Enterococcus faecalis*. Antimicrob. Agents Chemother. 41:702-705.
59. Tomich, P. K., F. Y. Au, and D. B. Clewell. 1980. Properties of erythromycin-inducible transposon Tn917 in *Streptococcus faecalis*. J. Bacteriol. 141:1366-1374.
60. Tomita, H., and D. B. Clewell. 2000. A pAD1-encoded small RNA molecule, mD, negatively regulates *Enterococcus faecalis* pheromone response by enhancing transcription termination. J. Bacteriol. 182:1062-1073.
61. Tomita, H., S. Fujimoto, K. Tanimoto, and Y. Ike. 1996. Cloning and genetic organization of the bacteriocin 31 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pY117. J. Bacteriol. 178:3585-3593.
62. Tomita, H., S. Fujimoto, K. Tanimoto, and Y. Ike. 1997. Cloning and genetic analysis of the bacteriocin 21 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pPD1. J. Bacteriol. 179:7843-7855.
63. Tomita, H., and Y. Ike. 2005. Genetic analysis of transfer-related regions of the vancomycin resistance *Enterococcus* conjugative pHTB: identification of *oriT* and a putative relaxase gene. J. Bacteriol. 187:7727-7737.
64. Tomita, H., E. Kamei, and Y. Ike. 2008. Cloning and genetic analyses of the bacteriocin 41 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pY114: a novel bacteriocin complemented by two extracellular components (lysine and activator). J. Bacteriol. 190:2075-2085.
65. Tomita, H., C. Pierson, S. K. Lim, D. B. Clewell, and Y. Ike. 2002. Possible connection between a widely disseminated conjugative gentamicin resistance (pMG1-like) plasmid and the emergence of vancomycin resistance in *Enterococcus faecium*. J. Clin. Microbiol. 40:3326-3333.
66. Tomita, H., K. Tanimoto, S. Hayakawa, K. Moringa, K. Ezaki, H. Oshima, and Y. Ike. 2003. Highly conjugative pMG1-like plasmids carrying *Tn1546*-like transposons that encode vancomycin resistance in *Enterococcus faecium*. J. Bacteriol. 185:7024-7028.
67. Uttley, A. H., C. H. Collins, J. Naidoo, and R. C. George. 1988. Vancomycin-resistant enterococci. Lancet i:57-58.
68. Weaver, K. E., and D. B. Clewell. 1988. Regulation of the pAD1 sex pheromone response in *Enterococcus faecalis*: construction and characterization of *lacZ* transcriptional fusions in a key control region of the plasmid. J. Bacteriol. 170:4343-4352.
69. Weaver, K. E., and D. B. Clewell. 1990. Regulation of the pAD1 sex pheromone response in *Enterococcus faecalis*: effects of host strain and *traA*, *traB*, and C region mutants on expression of an E region pheromone-inducible *lacZ* fusion. J. Bacteriol. 172:2633-2641.
70. Wirth, R., F. Y. Au, and D. B. Clewell. 1986. Highly efficient protoplast transformation system for *Streptococcus faecalis* and a new *Escherichia coli*-*S. faecalis* shuttle vector. J. Bacteriol. 165:831-836.
71. Yagi, Y., and D. B. Clewell. 1980. Recombination-deficient mutant of *Streptococcus faecalis*. J. Bacteriol. 143:966-970.
72. Yu, H. S., S. Y. Seol, and D. T. Cho. 2003. Diversity of *Tn1546*-like elements in vancomycin-resistant enterococci isolated from humans and poultry in Korea. J. Clin. Microbiol. 41:2641-2643.
73. Zechner, E. L., F. de la Cruz, R. Eisenbrandt, A. M. Grabn, G. Koraimann, E. Lanka, G. Muth, W. Pansegrau, C. M. Thomas, B. M. Wilkins, and M. Zatyka. 2000. Conjugative-DNA transfer processes, p. 87-174. In C. M. Thomas (ed.), The horizontal gene pool. Bacterial plasmids and gene spread, Harwood Academic Publishers, The Netherlands.
74. Zheng, B., H. Tomita, Y. H. Xiao, S. Wang, Y. Li, Y., and Y. Ike. 2007. Molecular characterization of vancomycin-resistant *Enterococcus faecium* isolates from mainland China. J. Clin. Microbiol. 45:2813-2818.

Cloning and Genetic Analyses of the Bacteriocin 41 Determinant Encoded on the *Enterococcus faecalis* Pheromone-Responsive Conjugative Plasmid pYI14: a Novel Bacteriocin Complemented by Two Extracellular Components (Lysin and Activator)[†]

Haruyoshi Tomita,¹ Elizabeth Kamei,¹ and Yasuyoshi Ike^{1,2*}

Department of Bacteriology and Bacterial Infection Control¹ and Laboratory of Bacterial Drug Resistance,² Gunma University Graduate School of Medicine, Maebashi, Gunma, Japan

Received 5 July 2007/Accepted 4 January 2008

The conjugative plasmid pYI14 (61 kbp) was isolated from *Enterococcus faecalis* YI714, a clinical isolate. pYI14 conferred a pheromone response on its host and encoded bacteriocin 41 (*bac41*). Bacteriocin 41 (Bac41) only showed activity against *E. faecalis*. Physical mapping of pYI14 showed that it consisted of EcoRI fragments A to P. The clone pHT1100, containing EcoRI fragments A (12.6 kbp) and H (3.5 kbp), conferred the bacteriocin activity on *E. faecalis* strains. Genetic analysis showed that the determinant was located in a 6.6-kbp region within the EcoRI AH fragments. Six open reading frames (ORFs) were identified in this region and designated ORF7 (*bacL₁*), ORF8 (*bacL₂*), ORF9, ORF10, ORF11 (*bacA*), and ORF12 (*bacI*). They were aligned in this order and oriented in the same direction. ORFs *bacL₁*, *bacL₂*, *bacA*, and *bacI* were essential for expression of the bacteriocin in *E. faecalis*. Extracellular complementation of bacteriocin expression was possible for *bacL₁* and *-L₂* and *bacA* mutants. *bacL₁* and *-L₂* and *bacA* encoded bacteriocin component L and activator component A, respectively. The products of these genes are secreted into the culture medium and extracellularly complement bacteriocin expression. *bacI* encoded immunity, providing the host with resistance to its own bacteriocin activity. The *bacL₁*-encoded protein had significant homology with lytic enzymes that attack the gram-positive bacterial cell wall. Sequence data for the deduced *bacL₁*-encoded protein suggested that it has a domain structure consisting of an N-terminal signal peptide, a second domain with the enzymatic activity, and a third domain with a three-repeat structure directing the proenzyme to its cell surface receptor.

Bacteriocins are bacterial proteins or peptides which inhibit the growth of other bacteria that are closely related to the producer strain. They usually exhibit a relatively narrow spectrum of activity and are produced by a wide variety of gram-positive and gram-negative bacteria (27). Bacteriocin production is thought to provide the host strain with an ecological or other selective advantage over other strains.

Many *Enterococcus faecalis* clinical isolates produce a bacteriocin (3, 5), and the bacteriocin is frequently encoded on the *E. faecalis* pheromone-responding conjugative plasmid (6, 14, 21, 46). Several *E. faecalis* bacteriocins have been genetically and biochemically characterized (15, 35), including the β -hemolysin/bacteriocin (cytolysin) (6, 7, 18, 20, 22) and the peptide antibiotics AS-48 (33), bacteriocin 21 (47), and bacteriocin 31 (46), which are encoded by the *E. faecalis* conjugative plasmids pAD1 (58 kbp), pMB2 (58 kbp), pPD1 (59 kbp), and pYI17 (57.5 kbp), respectively.

A significant number of *E. faecalis* clinical isolates produce hemolysin/bacteriocin (10, 26), and more than 50% of the hemolytic clinical isolates carry transferable hemolysin/bacte-

riocin determinants (21, 26). The hemolysin/bacteriocin of pAD1 is associated with virulence in animal models (4, 25, 29), and this plasmid is considered to be a typical *E. faecalis* hemolysin/bacteriocin plasmid (21, 31). The mechanism of hemolysin/bacteriocin production in *E. faecalis* has been studied in detail with the hemolysin/bacteriocin determinant on this plasmid (16, 17, 18, 22, 39). The active hemolysin/bacteriocin is produced by extracellular complementation of the two CylL factors (i.e., CylL_L and CylL_S) and CylA.

Previously, we have shown that bacteriocins or bacteriocinogenic *E. faecalis* clinical isolates can be classified into five groups on the basis of their bacteriocin activity against *E. faecalis* FA2-2 and OG1-10, *Enterococcus hirae* 9790, *Streptococcus pyogenes*, *Streptococcus agalactiae*, *Streptococcus sanguinis*, *Streptococcus pneumoniae*, *Staphylococcus aureus*, and *Staphylococcus epidermidis* (46). *E. faecalis* FA-2-2 and OG1-10 and *E. hirae* have been chosen as representative enterococcal strains for the examination and classification of the bacteriocins produced by the clinical isolates in this study. Class 1 types produce the β -hemolysin/bacteriocin (cytolysin) and are active against a wide variety of gram-positive bacteria, including *S. aureus* (2, 15, 17, 24, 46). The β -hemolysin/bacteriocin (cytolysin) of pAD1 belongs to class 1. Class 2 is active against a broad spectrum of bacteria, including *E. faecalis*, other *Streptococcus* spp., and *S. aureus*. AS-48 and bacteriocin 21 belong to class 2. Class 3 is active against *E. faecalis* and *E. hirae*. Class 4 is active against *E. faecalis*, and class 5 is active against *E. hirae*. The YI717, YI718, and YI719 strains belong to class 3 and harbor plasmids pYI17 (57.5 kb), pYI18,

* Corresponding author. Mailing address: Department of Bacteriology and Bacterial Infection Control, Gunma University Graduate School of Medicine, Showa-machi 3-39-22, Maebashi, Gunma 371-8511, Japan. Phone: 81-27-220-7990. Fax: 81-27-220-7996. E-mail: yasuike@med.gunma-u.ac.jp.

[†] Supplemental material for this article may be found at <http://jfb.asm.org/>.

[‡] Published ahead of print on 18 January 2008.

and pYI19, respectively (46). These plasmids encode the same bacteriocin with respect to immunity to the bacteriocin activity. Bacteriocin 31 (Bac31), encoded on pYI17, is representative of the class 3 bacteriocins and is active against *E. faecalis* and *E. hirae*, as is the membrane-active class II bacteriocin of lactic acid bacteria (46). The Bac31 determinant consists of the structural gene *bacA* and the immunity gene *bacB*.

In this report, we describe the cloning and genetic analysis of the bacteriocin 41 determinant encoded on *E. faecalis* pheromone-responsive conjugative plasmid pYI14, which is a representative class 4 bacteriocin. We also describe the identification of the two functional domains that are required to produce the active bacteriocin by extracellular complementation of the two factors.

MATERIALS AND METHODS

Bacterial strains, plasmids, oligonucleotides, media, and reagents. The bacterial strains, plasmids, and oligonucleotides used in this study are listed in Table 1. *E. faecalis* strains were grown in Todd-Hewitt broth (THB; Difco Laboratories) at 37°C, unless otherwise noted. *Escherichia coli* strains were grown in Luria-Bertani (LB) broth. The following antibiotic concentrations were used for the selection of *E. faecalis*: erythromycin, 12.5 µg ml⁻¹; streptomycin, 250 µg ml⁻¹; kanamycin, 250 µg ml⁻¹; spectinomycin, 250 µg ml⁻¹; chloramphenicol, 20 µg ml⁻¹; rifampin, 25 µg ml⁻¹; fusidic acid, 25 µg ml⁻¹. The antibiotic concentrations used for the selection of *E. coli* were as follows: ampicillin, 100 µg ml⁻¹; kanamycin, 40 µg ml⁻¹; chloramphenicol 50 µg ml⁻¹; spectinomycin, 50 µg ml⁻¹. All antibiotics were obtained from Sigma Chemical Co. X-Gal (5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside) was obtained from Wako Pure Chemical Industries, Ltd., and was used at 40 µg ml⁻¹.

Conjugation experiments. Broth mating and solid-surface mating were performed as previously described (48, 49), with a donor/recipient ratio of 1:10. Broth matings (in THB) were carried out for 4 h, and solid-surface matings (on THB agar plates) were carried out overnight (16 h) at 37°C. Transfer frequencies were calculated as the number of transconjugants per donor cell (at the end of mating). Pheromone induction and detection of cell aggregation were performed as previously described (11, 12).

Soft-agar assay for bacteriocin production and immunity. The bacteriocin production assay was performed as described previously (22). The test for immunity to the bacteriocin was performed essentially as described previously (22).

Plasmid DNA methodology. Recombinant plasmids were generated in *E. coli* DH5α. Transformation of bacterial cells with plasmid DNA was achieved by electrotransformation as described previously (13). Plasmid DNA was purified from *E. coli* (38) or from *E. faecalis* as previously described (14). DNA fragments were purified from an agarose gel after electrophoresis with a Gene Clean II kit (Bio 101, Inc.). Recombinant DNA methodology, analyses of plasmid DNA with restriction enzymes, and agarose gel electrophoresis were carried out by standard methods (38). Restriction enzymes were purchased from New England Biolabs, Roche, Nippon Gene, and Takara Co., and reactions were carried out under the conditions recommended by the manufacturers. DNA ligations were performed with a DNA ligation kit from Takara. To end fill the endonuclease-digested DNA fragment for ligation, a DNA-blunting kit and Klenow enzyme were obtained from Takara and used according to the manufacturer's protocol (45).

Determination of the pYI14 restriction map. pYI14 plasmid DNA was digested with EcoRI, BamHI, KpnI, SphI, or XbaI or double digested with a combination of two of these restriction enzymes. Agarose gel electrophoresis analysis of the digested DNAs was performed to determine the cleavage sites within the plasmid. To determine the order of the EcoRI fragments of pYI14, a relational clone set was constructed as previously described (14, 46). After agarose gel electrophoresis of plasmid pYI14 DNA partially digested with EcoRI, fragments greater than 7 kb in size were eluted and used for cloning. The cloning vectors used were pBluescript-SK(+) and pAM401, and the host strain was *E. coli* DH5α.

DNA sequence analysis. Nucleotide sequence analysis was carried out as previously described (14). A deletion kit (Nippon Gene) was used. BamHI-E, BamHI-F, EcoRI-H, and the 2.1-kb fragment between BamHI-F and EcoRI-H were individually cloned into the pBluescript vector. The clones were used to construct a series of deletional clones. The resulting constructs were sequenced in both orientations with the *Taq* Dye primer and the *Taq* Big Dye terminator cycle sequencing kit (Applied Biosystems), a model 377 DNA sequencer, and a

310 gene analyzer (ABI Prism). A database search was performed with the BLASTn and tBLASTx programs of the National Center for Biotechnology Information, Bethesda, MD (1).

Generation of transposon (Tn5, mini-Tn7) insertion mutants. Insertion of Tn5 (Km^r) into the cloned plasmid DNA was performed as described elsewhere (47). Target plasmid pHT1100(pAM401 containing EcoRI fragments A and H) was introduced into *E. coli* K-12 TH688 (with Tn5 in the *thr* locus) (42) by electrotransformation. Transformants were spread onto selective plates containing kanamycin and chloramphenicol, and the plates were left at room temperature for 10 days. The bacteria that grew on the selective plates were pooled, and the plasmid DNA was then isolated and used to transform *E. coli* DH5α. The transformants were selected on plates containing kanamycin and chloramphenicol for the selection of Tn5-borne kanamycin resistance and plasmid-borne chloramphenicol resistance, respectively. The transformants were purified and examined to determine the location of Tn5 within the plasmid. The precise locations of Tn5 insertions were determined by DNA sequence analysis with a synthetic primer that hybridized to the end of Tn5. A GPS kit (NEB) was used to generate mini-Tn7 insertion mutants with plasmid pHT1100 according to the manufacturer's instructions.

PCR amplification and primers. PCR amplification was performed with the thermostable DNA polymerase Takara *Taq* (Takara Bio Inc.) and a Perkin-Elmer 9600 thermal cycler. PCR conditions varied according to the primers used and the size of the anticipated product. The custom primers used in this study were obtained from Invitrogen (Tokyo, Japan) and are listed in Table 1. Each of the amplified PCR products was trimmed by the appropriate restriction enzyme, purified with a QIAquick-spin column (Qiagen), and cloned into plasmid pAM401.

Nucleotide sequence accession number. The nucleotide sequence reported in this article is available from the DDBJ, EMBL, and GenBank nucleotide sequence databases under accession number AB271686.

RESULTS

Bacteriocinogenic *E. faecalis* strain and the pheromone-responsive bacteriocin plasmid. Four strains that were active against *E. faecalis* and were classified as class 4 bacteriocinogenic strains were isolated from clinical urine samples and were designated YI712, YI714, YI715, and YI716. YI712 harbored plasmid pYI12 (72 kb). YI714 and YI715 harbored plasmids pYI14 (61 kb) and pYI141 (48 kb) and plasmids pYI15 (61 kb) and pYI151 (48 kb), respectively. YI716 did not carry any plasmid. Each strain was used as a donor in mating experiments with plasmid-free recipient strain *E. faecalis* FA2-2 (Rif^r Fus^r) to determine whether these plasmids conferred bacteriocinogenic activity on the host. After incubating the broth mating cultures for 4 h, appropriately diluted mixtures were plated on an agar plate containing rifampin (25 µg/ml) and fusidic acid (25 µg/ml) to select for the recipient strains. After overnight incubation of the plates, a total of approximately 500 *E. faecalis* FA2-2 colonies were obtained from each mating and examined for bacteriocin production. Approximately 1 in 500 cells obtained from the mating experiments with each of the strains described above expressed bacteriocin activity against *E. faecalis* FA2-2. The bacteriocinogenic transconjugants of YI712, YI714, and YI715 harbored pYI12 (72 kb), pYI14 (61 kb), and pYI15 (61 kb), respectively. The same EcoRI restriction profiles were obtained for pYI14 and pYI15, implying that the two plasmids were identical. Each plasmid transferred between *E. faecalis* FA2-2 and *E. faecalis* OG1-10 at a frequency of about 10⁻³ per donor cell by broth mating. *E. faecalis* FA2-2(pYI12), FA2-2(pYI14), and FA2-2(pYI15) did not exhibit bacteriocin activity against *E. faecalis* OG1-10(pYI12), OG1-10(pYI14), or OG1-10(pYI15). These results imply that plasmids pYI12, pYI14, and pYI15 encoded the same bacteriocin with respect to the immunity character-

TABLE 1. Bacterial strains, plasmids, and oligonucleotides used in this study

Strain, plasmid, or oligonucleotide	Relevant features or sequence (5'-3') ^a	Reference(s), source, or generated plasmid(s)
Strains		
<i>E. faecalis</i>		
FA2-2	<i>rif^r fus</i>	7
JH2SS	<i>spc str</i>	44
OG1-10	<i>str</i> , derivative of OG1	12
OG1X	<i>str</i> , protease-negative derivative of OG1-10	23
YI712	pYI12(Bac)	This study
YI714	pYI14(Bac), pYI141 (48 kb); clinical isolate	This study
YI715	pYI15(Bac), pYI151 (48 kb); clinical isolate	This study
<i>E. coli</i> DH5 α		
	<i>endA1 recA1 gyrA96 thi-1 hsdR17 supE44 relA1 Δ(argE-lacZYA)U169</i>	Bethesda Research Laboratories
Plasmids		
pAM401	<i>E. coli-E. faecalis</i> shuttle plasmid; <i>cat tet</i>	50
pLZ12-Km	<i>E. coli-Streptococcus</i> shuttle plasmid; <i>aphA</i>	19
pBlueScript SKII(+)	<i>E. coli</i> cloning vector; Amp ^r	Stratagene
pPD1	Bac21, 59-kb conjugative plasmid from strain 39-5	14, 47
PMG326	pMW119 containing a 16.7-kbp EcoRI-SalI fragment of pPD1; pheromone-regulatory region	14, 41
pYI12	Bac41, 72-kb conjugation plasmid from YI712	This study
pYI14	Bac41, 61-kb conjugative plasmid from YI714	This study
pYI15	Bac41, 61-kb conjugative plasmid from YI715	This study
pHT1100	EcoRI-relational clone of pYI14; pAM401 containing EcoRI fragments A and H	This study
pHT1101	EcoRI-relational clone of pYI14; pAM401 containing EcoRI fragment A	This study
pHT1102	EcoRI-relational clone of pYI14; pAM401 containing EcoRI fragments H and M	This study
PMG1103	Derivative of pHT1100 with BamHI E fragment deleted	This study
PMG1104	Derivative of pHT1100 with BamHI F fragment deleted	This study
PMG1106	Derivative of pHT1100; BamHI site at 4.1 kbp blunted with Klenow enzyme	This study
PMG1108	Derivative of pHT1100; BamHI site at 6.3 kbp blunted with Klenow enzyme	This study
PMG1109	Derivative of pHT1100; KpnI site at 4.6 kbp blunted with DNA-blunting kit (Takara)	This study
PMG1105- <i>n</i>	Tn5 insertional derivatives of pHT1100	This study
PMG1107- <i>n</i>	Mini-Tn7 insertional derivatives of pHT1100 created with GPS kit (New England Biolabs)	This study
PMG1110	<i>baeL</i> ₁ and <i>baeL</i> ₂ ; pAM401 containing 2,932-bp EcoRI fragment amplified by PCR	This study
PMG1111	<i>baeA</i> ; pAM401 containing 2,836-bp SalI fragment amplified by PCR	This study
PMG1112	<i>baeI</i> ; pAM401 containing 777-bp BamHI fragment amplified by PCR	This study
PMG1113	<i>baeI</i> and ORF13; pAM401 containing 1,513-bp BamHI fragment amplified by PCR	This study
PMG1114	pLZ12-Km containing 10-kbp BglII fragment mapped from 1.7 kbp to 11.7 kbp	This study
PMG1115	Derivative of PMG1114; EcoRI fragment (8.5 kbp to vector region) deleted	This study
PMG1116	Derivative of PMG1114; three HindIII fragments (4.6- to 6.6-kbp region) deleted	This study
Oligonucleotides		
B9P2842F	cgc <u>gaa</u> tTC TAG CAA CCG AAA ACC ACG TTG G	pMG1110
B9P5773R	gcg <u>gaa</u> tTC ATT GCG CAG CAA ATC ATT GC	pMG1110
B9P6180F	aac gcg <u>tcg</u> ACA GGA ATT GAG ACA TAC GCT	pMG1111
B9P9015R	aac gcg <u>tcg</u> acT TCG TCA AAT CCA TTT CCC CTA	pMG1111
B9P8823F	ggc <u>gga</u> tcc GCA GCA GAA TTA GCA GGA GCG	pMG1112, pMG1113
B9P9599R	gcc <u>gga</u> tcc CAA AAG TCA TAC ATG ACC TCC	pMG1112
B9P10335R	gcc <u>gga</u> tcc CTG TAT AAA TCC ATA CTA CAC	pMG1113

^a Underlining indicates the following restriction endonuclease recognition sequences: GAATTC; EcoRI, GTCGAC; SalI, GGATCC; BamHI. Lowercase letters indicate incorporated tag sequences.

istic *E. faecalis* FA2-2 strains carrying pYI12, pYI14, or pYI15 were tested for bacteriocin production against the indicator strains *S. aureus* FDA209P, *E. faecalis* FA2-2 and OG1-10, *Enterococcus faecium* BM4105RF, *E. hirae* ATCC 9790, *Enterococcus*

durans ATCC 49135, *Enterococcus raffinosus* JCM8733, *Enterococcus gallinarum* BM4174, *S. agalactiae*, *S. pyogenes*, *Listeria monocytogenes*, and *Listeria denitrificans*. Each of the three bacteriocinogenic strains only showed bacteriocin activity against




FIG. 1. Physical map of pYI14 showing the locations of bacteriocin 41 determinants *bacL*₁, *bacL*₂, *bacA*, and *bacI*. Each value is the size of the fragment in kilobases.

E. faecalis. Plasmid pYI14, isolated from strain YI714, was used as the representative plasmid encoding the bacteriocin.

The donor *E. faecalis* OG1-10(pYI14) and recipient *E. faecalis* FA2-2 formed a mating aggregate in the mating mixture. When OG1-10(pYI14) cells were exposed to *E. faecalis* FA2-2 culture filtrate (pheromone) for 4 h at 37°C, the OG1-10(pYI14) cells showed aggregation. Agarose gel electrophoresis of the EcoRI restriction fragments of pYI14 DNA was carried out, and the DNA was transferred to a membrane for Southern hybridization. The membrane was hybridized with a DNA probe containing the pheromone response genes of the pheromone-responsive plasmid pPD1 or plasmid pMG326, which contains the putative surface exclusion protein gene and the N-terminal region of the aggregation substance gene of pPD1 (14, 41). Each probe hybridized to specific pYI14 EcoRI fragments (data not shown). These results indicated that plasmid pYI14 was a pheromone-responsive plasmid.

Restriction map of pYI14. To determine the order of the EcoRI fragments, a relational clone set was obtained. The order of EcoRI fragments was determined to be A-H-M-L-J-K-O-B-D-N-P-G-I-E-F-C (Fig. 1). Each clone was digested with BamHI, KpnI, SphI, and XbaI, and the cleavage sites were determined (Fig. 1). Restriction sites within the EcoRI A and H fragments were also confirmed by sequencing (see the supplemental material).

Bacteriocin activity of the cloned DNA fragment. To examine the bacteriocin activity of the relational clones, each clone was introduced into *E. faecalis* OG1-10 and the resulting transformant was examined for bacteriocin activity. *E. faecalis* OG1-10 carrying plasmid pHT1100, which contained the EcoRI A and H fragments (16.1 kb), exhibited the bacteriocin activity (Fig. 2). *E. faecalis* OG1-10 carrying either the EcoRI A (12.6 kb) or HM (4.6 kb) fragments (plasmids pHT1101 and pHT1102, respectively) did not exhibit bacteriocin activity (Fig. 2). *E. faecalis* OG1-10 carrying the EcoRI HM fragments

showed resistance to the bacteriocin activity of *E. faecalis* OG1-10(pYI14). These results indicated that the bacteriocin determinant of pYI14 is located on the EcoRI A and H (AH) fragments and the immunity gene (i.e., the gene for resistance to its own bacteriocin) is located on the EcoRI H fragment.

DNA sequence analysis. The EcoRI AH fragments were sequenced, and computer analysis was used to identify open reading frames (ORFs) within the sequence. Fifteen ORFs (ORF1 to ORF15) were located in the region spanning map positions 0 to 12 kbp, as indicated by the numerical scale shown in Fig. 2, where position 0 is the BamHI site located between BamHI fragments E and C and position 12 kbp is the EcoRI site located between the EcoRI H and M fragments (Fig. 1 and 2 and Table 2; see Fig. S1 in the supplemental material). Figure 2 shows the ORFs that have a deduced ribosome-binding site in the 20-base region upstream of the predicted start codon and the potential promoters for initiation of transcription.

Generation of Tn5 or mini-Tn7 insertion mutants. To examine the location of the bacteriocin determinant, mutants with altered bacteriocin expression were generated by Tn5 or mini-Tn7 insertion into pHT1100. The precise locations of Tn5 or mini-Tn7 insertions in the ORFs were determined by DNA sequence analysis (see Fig. S1 in the supplemental material), and the results are shown in Fig. 2 and Table 3. Tn5 insertions into ORF7, ORF8, and ORF11 resulted in defective bacteriocin activity in *E. faecalis* OG1-10. Insertion of mini-Tn7 into the C-terminal region of ORF11 also resulted in defective bacteriocin activity in *E. faecalis* OG1-10. *E. faecalis* OG1S carrying each of the insertion mutants showed resistance to the bacteriocin activity of *E. faecalis* OG1-10(pYI14), indicating that the mutant plasmids retained immunity to the bacteriocin.

Generation of deletion mutants by end filling after cleavage with a restriction enzyme. Mutant pHT1100 plasmids with BamHI fragment deletions within were also generated to examine the location of the bacteriocin determinant as described in Materials and Methods (Fig. 2) (47). Deletion mutant plasmids pMG1103 and pMG1104 possessed deletions of the 4.1-kbp BamHI E fragment between map positions 0 kb and 4.1 kb and the 2.2-kbp BamHI F fragment between map positions 4.1 kb and 6.3 kb, respectively. Plasmid pMG1103, which had a deletion in the amino-terminal region of ORF7 and had lost the six ORFs located upstream of ORF7, did not exhibit bacteriocin activity but retained immunity to the bacteriocin. Plasmid pMG1104, which had deletions within the carboxyl-terminal region of ORF7, ORF8, and ORF9 and the amino-terminal region of ORF10, did not exhibit bacteriocin activity but retained immunity to the bacteriocin. These results implied that the gene for immunity is located downstream of ORF11.

Generation of four-nucleotide insertion (deletion) mutants. Mutants with changes in ORF7 and ORF10 were generated to obtain mutants with in-frame changes in the determinant by blunt ending the recessed 3' terminus of the BamHI site or the prominent 3' terminus of the KpnI cleavage site within pHT1100 DNA that had been partially digested with these enzymes prior to ligation (Fig. 2) (45). Blunt ending the BamHI and KpnI sites resulted in the insertion of four nucleotides (5'-GATC-3') with the Klenow enzyme in the case of the BamHI site and the deletion of four nucleotides (5'-GTAC-3') with the T4 DNA polymerase DNA-blunting kit




FIG. 2. Physical maps of the 16.1-kb region containing EcoRI fragments A (12.6 kb) and H (3.5 kb) in pY114 (which is carried on pHT1100), transposon insertions, and subclones. The zero position of the numerical scale (top horizontal line) indicates the BamHI endonuclease recognition site located between the BamHI C and E fragments, and it runs in a clockwise direction on the physical map of Fig. 1. Thick horizontal arrows indicate the predicted ORFs and the direction of ORF transcription. The flags and hairpins below the ORFs indicate the potential promoter regions and inverted repeat sequences. The horizontal lines under the map represent the cloned pY114 DNA fragments in the derivative plasmids listed on the left. Small vertical bars at ends of the lines represent the endonuclease recognition sites for cloning. The dotted vertical lines represent the ends of the amplified PCR fragment of pY114 DNA used to clone the bacteriocin determinant. The endonuclease recognition sites incorporated for the cloning of the PCR products are indicated. Abbreviations of the endonuclease recognition sites: Eco, EcoRI; Ba, BamHI; Kp, KpnI; Bg, BglII; Hi, HindIII; Sa, Sall. Bac +, normal bacteriocin expression; Bac -, no bacteriocin expression; Imm +, resistance to bacteriocin 41; Imm -, sensitive to bacteriocin 41. The vertical lines with circular or triangular heads on the pHT1100 map show the points of transposon insertion. The circular heads indicate Tn5, and the triangular heads indicate mini-Tn7 and its orientations. The heads represent the levels of bacteriocin 41 expression in *E. faecalis* strains as follows: open heads, normal bacteriocin expression; black heads, no bacteriocin expression; gray heads, weak bacteriocin expression (Fig. 3A). The values on the insertions indicate the numbers of insertions and correspond to those shown in Table 3 (see also Fig. S1 in the supplemental material). The cross marks on the clones indicate the mutated endonuclease recognition sites (a four-base insertion or deletion). aa, amino acids.

(Takara) in the case of the KpnI site. The pMG1106 and pMG1109 mutants that resulted from the blunt ending of the BamHI site and KpnI sites in ORF7 did not exhibit bacteriocin activity but retained the immunity activity, indicating that ORF7 is essential for bacteriocin expression. The pMG1108 mutant, which resulted from the end filling of the BamHI site in ORF10, expressed both bacteriocin and immunity activity, suggesting that ORF10 is not essential for bacteriocin expression.

Subcloning of the bacteriocin determinant and generation of the derivative mutants. The 10.0-kb BglII fragment that is located between 1.7 kb and 11.7 kb on the map was cloned into shuttle vector pLZ12-Km (19) (Fig. 2), and the cloned plasmid

was designated pMG1114. pMG1114 expressed both bacteriocin activity and immunity, indicating that the bacteriocin determinant was located within the 10.0-kb BglII fragment. Deletion mutants pMG1115 and pMG1116 were generated from pMG1114. pMG1115 had a deletion of the 3.3-kbp EcoRI/BglII fragment between 8.4 kb and 11.7 kb on the map, which contains the C-terminal region of ORF11. pMG1115 did not express either the bacteriocin or immunity, indicating that ORF11 is necessary for bacteriocin activity. pMG1116 had a deletion of two HindIII fragments totaling 1.4 kb that were located between 5.2 kb and 6.6 kb on the map and contains the C-terminal region of ORF7 and all of ORF8, ORF9, and ORF10. pMG1116 did not express the bacteriocin but ex-

TABLE 2. ORFs encoded on the BamHI/EcoRI 11,952-bp-spanning region

ORF	Gene	Map location (bp)	Gene/protein size (base pairs/amino acids)	Homology	% Identity/similarity (amino acids)	Function
1		136-588	453/150	<i>pcfS</i> (<i>E. faecalis</i> pCF10)	98/100	Ssb ^a
2		602-754	153/50	EFB0044 (<i>E. faecalis</i> V583 pTEF2)	100/100	
3		766-1344	579/192	<i>pcfT</i> (<i>E. faecalis</i> pCF10)	89/90	Thermonuclease
4		1350-1670	321/106	<i>pcfU</i> (<i>E. faecalis</i> pCF10)	93/97	
5		1827-2009	183/60	Efae03001107 (<i>E. faecium</i>)	50/67	
6		2204-2920	717/238	Lipoprotein (<i>E. faecalis</i> V583)	31/45	
7	<i>bacl₁</i>	3058-4845	1,788/595	Lysozyme (<i>B. subtilis</i> bacteriophage B103)	37/52 (1-151)	Lysin (bacteriocin 41)
				Lysin (<i>S. agalactiae</i> prophage lambda Sa1)	46/63 (160-309)	
				Muramidase (<i>L. plantarum</i> WCFS1)	24/41 (318-577)	
8	<i>bacl₂</i>	5031-5666	636/211			Lysin expression
9		5689-6120	432/143	ORF50 (<i>S. pneumoniae</i> bacteriophage MM1)	31/51	Holin
10		6123-6650	528/175	EF0637 (<i>E. faecalis</i> V583)	27/45	
11	<i>bacl₃</i>	6693-8873	2,181/726	<i>ybfG</i> (<i>B. subtilis</i>)	41/56	Lysin activator
				<i>ykuG</i> (<i>B. subtilis</i>)	40/55	
12	<i>bacl₄</i>	8981-9523	543/180			Immunity
13		9590-10165	576/191			
14		10308-10640	333/110	EFB0057 (<i>E. faecalis</i> V583 pTEF2)	98/100	
15		11080-11781	702/233			

^a Ssb, single-stranded binding protein.

pressed immunity. Analysis of the insertion mutants and deletion mutants showed that ORF7, ORF8, ORF11, and ORF12 are necessary for bacteriocin expression.

Extracellular complementation of nonbacteriocinogenic mutants. Extracellular complementation experiments to express bacteriocin activity were performed with ORF7 or ORF8 and ORF11 mutant strains on soft agar plates containing the indicator strain. OG1-10(pMG1106) and OG1-10(pMG1109), which were ORF7 mutants prepared by blunt ending, were streaked in proximity to streaks of either OG1-10 carrying the pHT1100 derivatives of the Tn5 insertion mutants in ORF11 or OG1-10(pHT1101) with a deletion in ORF11. This experiment showed that there was complementation of the bacteriocin activity at the streak junction. When OG1-10(pHT1101) was streaked in proximity to streaks of OG1-10 carrying the pHT1100 derivatives of the Tn5 insertion mutants in ORF7 or ORF8 and the end-filled mutants of ORF7, complementation of the bacteriocin activity was observed at the streak junction. These results indicated that the mutants fell into one of two complementation groups. Representative results are shown in Fig. 3 and Table 4. The OG1-10(pHT1100), OG1-10(pHT1101), and OG1-10(pMG1106) strains were inoculated in proximity in soft agar containing the indicator strain (Fig. 3B). Bacteriolysis was observed around the wild-type strain and also between OG1-10(pHT1101) and the wild-type strain or OG1-10(pMG1106), respectively. Bacteriolysis was also observed surrounding OG1-10(pHT1101). Figure 3C shows the complementation activity that resulted from cross-streaking of OG1-10(pMG1106) and OG1-10(pHT1101) on soft agar containing the indicator strain. Bacteriolysis was observed at the junction of the two strains. Based on these observations, the two complementation substances were tentatively designated L (lysin) and A (activator). OG1-10(pHT1101) and the ORF11 mutants were presumed to be defective in bacteriocin component A synthesis and tentatively assigned an L⁻ A⁻ phenotype. The ORF11 gene was designated *bacl₄*. OG1-10(pMG1106) and the ORF7 and ORF8 mutants

were presumed to be defective in bacteriocin component L synthesis and tentatively assigned an L⁻ A⁺ phenotype. The ORF7 and ORF8 genes were designated *bacl₁* and *bacl₂*, respectively.

Cloning of component L, component A, and the immunity genes. The PCR product of each ORF was cloned to analyze its function in bacteriocin expression (Fig. 2). Cloned pMG1110, pMG1111, pMG1112, and pMG1113 contained ORF7/8 (*bacl₁* and *-L₂*), ORF11 (*bacl₄*), ORF12, and ORF12/13, respectively (Fig. 2). Each of the individually cloned fragments did not express bacteriocin activity. pMG1112 and pMG1113, which contained ORF12 and ORF12/13, expressed immunity to the bacteriocin activity, indicating that ORF12 was the immunity gene, and it was designated *bacl₃*.

Extracellular complementation between cloned L and A components. Cross streaks of strains carrying the two cloned fragments were made on bacteriocin assay plates. When OG1-10(pMG1110), which contained ORF7 (*bacl₁*) and ORF8 (*bacl₂*), was streaked across a preexisting streak of OG1-10(pMG1111), which contained ORF11 (*bacl₄*), a large area of bacteriolysis was observed around the two crossed strains (Table 4). Growth of the two strains was markedly inhibited. These data indicated that the product of each strain complemented to produce an active bacteriocin, but the two strains have no immunity to the bacteriocin; therefore, growth of the strains was inhibited by the bacteriocin.

DNA sequence analysis of ORFs located in the region containing the bacteriocin 41 determinant. A homology search of the 15 ORFs contained in the 12-kbp region was performed by BLAST against the protein databases, and the results are shown in Table 2 (1). ORF7 (*bacl₁*), ORF8 (*bacl₂*), ORF11 (*bacl₄*), and ORF12 (*bacl₃*) were essential for the expression of bacteriocin 41. *bacl₁* encoded a 595-amino-acid protein. Computer analysis suggested that the deduced *bacl₁*-encoded protein had a signal peptide sequence and that a potential signal peptidase processing site corresponding to the L-K-A sequence was located at positions 19 to 21 (Fig. 4A). Comparison

TABLE 3. Transposon insertion mutants of pHT1100 and bacteriocin expression

Insertion no. in Fig. 2	Plasmid(s) ^a	Transposon	Map position (kb) ^b	Insertion location	Bac ^c	Imm ^d
	pY114				++	+
	pHT1100				++	+
44	pMG1105-44	Tn5	-1.2	Upstream of ORF1	++	+
9	pMG1105-9	Tn5	-0.8	Upstream of ORF1	++	+
49	pMG1105-49	Tn5	-0.4	Upstream of ORF1	++	+
19	pMG1105-19	Tn5	0.7	ORF2	++	+
46	pMG1105-46	Tn5	1.0	ORF3	++	+
8, 186	pMG1105-8, -186	Tn5	1.7	Between ORF4 and ORF5	++	+
161	pMG1105-161	Tn5	1.9	ORF5	++	+
24	pMG1105-24	Tn5	2.6	ORF6	++	+
42	pMG1105-42	Tn5	3.2	ORF7 (<i>bacl</i> ₁)	-	+
30	pMG1105-30	Tn5	4.0	ORF7 (<i>bacl</i> ₁)	-	+
54, 105, 193	pMG1105-54, -105, -193	Tn5	4.5	ORF7 (<i>bacl</i> ₁)	-	+
105	pMG1105-105	Tn5	4.6	ORF7 (<i>bacl</i> ₁)	-	+
32	pMG1105-32	Tn5	4.7	ORF7 (<i>bacl</i> ₁)	-	+
25	pMG1105-25	Tn5	4.8	ORF7 (<i>bacl</i> ₁)	-	+
107	pMG1105-107	Tn5	5.2	ORF8 (<i>bacl</i> ₂)	-	+
188	pMG1105-188	Tn5	5.5	ORF8 (<i>bacl</i> ₂)	-	+
104, 121	pMG1105-104, -121	Tn5	6.9	ORF11 (<i>bacA</i>)	-	+
45	pMG1105-45	Tn5	7.1	ORF11 (<i>bacA</i>)	-	+
123	pMG1105-123	Tn5	7.3	ORF11 (<i>bacA</i>)	-	+
20	pMG1105-20	Tn5	7.3	ORF11 (<i>bacA</i>)	-	+
51	pMG1105-51	Tn5	7.7	ORF11 (<i>bacA</i>)	-	+
6	pMG1105-6	Tn5	8.1	ORF11 (<i>bacA</i>)	-	+
11	pMG1105-11	Tn5	8.4	ORF11 (<i>bacA</i>)	-	+
22	pMG1107-22	Mini-Tn7	8.7	ORF11 (<i>bacA</i>)	-	+
14, 101, 117	pMG1105-14, -101, -117	Tn5	9.5	ORF12 (<i>bacI</i>)	+	±
80	pMG1107-80	Mini-Tn7	9.5	Between ORF12 and ORF13	++	+
25	pMG1107-25	Mini-Tn7	9.6	ORF13	++	+
115, 153	pMG1105-115, -153	Tn5	10.4	ORF14	++	+
38, 39, 163	pMG1105-38, -39, -163	Tn5	10.7	Between ORF14 and ORF15	++	+
87	pMG1107-87	Mini-Tn7	11	Between ORF14 and ORF15	++	+
74	pMG1107-74	Mini-Tn7	11	Between ORF14 and ORF15	++	+
76	pMG1107-76	Mini-Tn7	11.1	Between ORF14 and ORF15	++	+
69	pMG1107-69	Mini-Tn7	11.1	ORF15	++	+
59	pMG1107-59	Mini-Tn7	11.2	ORF15	++	+
70	pMG1107-70	Mini-Tn7	11.3	ORF15	++	+
40	pMG1107-40	Mini-Tn7	11.4	ORF15	++	+
80	pMG1107-80	Mini-Tn7	11.4	ORF15	++	+
83	pMG1107-83	Mini-Tn7	11.5	ORF15	++	+

^a The host strain of the derivative was *E. faecalis* OG1S (OG1-10).

^b The map position is the distance from the junction between EcoRI fragments A and H. Minus values indicate the opposite direction.

^c Bac, bacteriocin expression. Symbols: ++, normal bacteriocin expression; +, weak bacteriocin expression (Fig. 3A); -, no bacteriocin expression.

^d Imm, immunity expression. Symbols: +, positive expression; -, no expression; ±, weak expression.

of the primary structure of the deduced amino acid sequence of the BacL₁ protein showed significant homology with the cell wall lytic enzymes found in gram-positive bacteria (Fig. 4A) (32). Of the 595 amino acid residues of the BacL₁ protein, the N-terminal 151 amino acid residues showed a high level of homology with the lysozyme encoded on *Bacillus subtilis* bacteriophage B103 (accession number Q37896) (37). The 150-amino-acid sequence from residue 160 to residue 309, which is located in the center of the *bacl*₁-encoded protein, showed a high level of homology with the N-terminal amino acid residues of the lysin encoded on the *S. agalactiae* prophage lambda Sa1 (accession number NP 687631) (43), and the C-terminal 260 amino acid residues showed a high level of homology with the C-terminal amino acid residues of the muramidase of *Lactobacillus plantarum* WCFS1 (accession number CAD64901) (30). The *bacl*₁-encoded protein harbored a three-repeat structure of an almost identical amino acid sequence (Fig. 4B). The three-repeat structure located at the C terminus of the

*bacl*₁-encoded protein corresponded to the homologous C-terminal region of the *L. plantarum* WCS1 muramidase, which is thought to be a choline-binding region (28, 51). The repeat structure was composed of three copies of an almost identical 74-amino-acid sequence. The first copy was located between amino acid residues 333 and 406, the second copy was located between amino acid residues 424 and 497, and the third copy was located between amino acid residues 520 and 593. *bacl*₂ encoded a 211-amino-acid protein and did not show any significant homology with other reported proteins. There was no obvious leader peptide with hydrophobic residues at the N-terminal peptide of the deduced *bacl*₂-encoded protein. *bacA* encoded a 726-amino-acid protein and showed a significant degree of homology with *ybfG* and *ykuG* of *B. subtilis*, but the function of these proteins is unknown (Fig. 5) (accession numbers CAB12014 for *YbfG* and CAA10870 for *YkuG*, respectively). The *bacA* protein had a putative signal peptide sequence, and a potential signal peptidase processing site corresponding to




FIG. 3. Bacteriocin expression assay by the soft-agar method with *E. faecalis* OG1-10 carrying the representative pY114 bacteriocin derivatives (A) and complementation assays (B and C). The indicator strain was *E. faecalis* OG1-10. The strains used are shown in Fig. 2 and Table 3. (A) 1, OG1-10(pHT1100) wild type; 2, OG1-10(pMG1105-14, a transposant of pHT1100::Tn5) (Tn5 inserted in the C-terminal region of ORF12); 3, OG1-10(pMG1106) in-frame *bacl*₁ mutant. (B) 1, OG1-10(pHT1100); 2, OG1-10(pHT1101) *bacl*₁ and *bacl*₂ deletion mutant; 3, OG1-10(pMG1106). (C) 1, OG1-10(pHT1101), 2, OG1-10(pMG1106). L, *bacl*₁ and *bacl*₂ expression; A, *bacl*₁ expression; I, immunity expression; +, positive expression; -, no expression; ±, weak expression.

the V-S-G sequence was located at positions 19 to 21 (Fig. 5). The *bacl*₁ protein contained a 60-amino-acid sequence corresponding to the putative peptidoglycan-binding domain, which was located between amino acids 81 and 140 in the *bacl*₁-encoded protein, suggesting that the BacA protein could be directed to the bacterial cell surface.

DISCUSSION

Bacteriocin 41 of strain YI714 was encoded on *E. faecalis* pheromone-responsive plasmid pY114 (61 kbp) and was only active against *E. faecalis*. The EcoRI AH fragments of pY114,

which conferred the bacteriocin activity, were cloned and used for genetic analysis of the bacteriocin determinant. Transposon insertion and deletion mutant analysis of the EcoRI AH fragments and further subcloning of the bacteriocin determinant showed that a 6.6-kb fragment of pY114 was the minimum-size fragment required for bacteriocin expression. The 6.6-kb region contained six ORFs, which were designated *bacl*₁, *bacl*₂, ORF9, ORF10, *bacl*₃, and *bacl*. All of the ORFs were oriented in the same direction and in that order. The insertion mutants were classified into one of two complementation classes for component L and component A. Each class showed extracellular complementation to produce the active bacteriocin. A series of PCR products containing the L-encoding region for component L, the A-encoding region for component A, and the immunity-encoding region for resistance to the bacteriocin were subcloned into *E. faecalis* OG1-10. The subclones for the L-encoding, A-encoding, and immunity-encoding regions contained *bacl*₁ and *bacl*₂, *bacl*₃, and *bacl*, respectively. The subclone containing *bacl*₁ and *bacl*₂ produced an L component capable of extracellular complementation with the A component for expression of bacteriocin activity, indicating that *bacl*₁ and *bacl*₂ were required for component L. These results indicated that of the ORFs within the 6.6-kb region, *bacl*₁, *bacl*₂, and *bacl*₃ are essential for the production of the active bacteriocin, and *bacl* is the immunity gene for resistance to the bacteriocin that is produced.

Tn5 insertions into *bacl*₁ or *bacl*₂ of the bacteriocin determinant did not result in a detectable polar effect on the expression of the downstream *bacl*₃ or *bacl* gene, and insertion into *bacl*₃ also did not result in a polar effect on the expression of *bacl*. Both component determinants and *bacl* were expressed when each of the determinants was cloned into vector plasmid pAM401 in either orientation within an *E. faecalis* OG1-10 background. These results suggested that a significant amount of transcription of the *bacl*₁, *bacl*₂, *bacl*₃, and *bacl* genes can occur from different promoters.

In the complementation experiment between the L⁺ A⁻ and L⁻ A⁺ strains, bacteriocin activity was observed around the L⁺ A⁻ strain. When the wild-type L⁺ A⁺ and mutant L⁺ A⁻ strains were inoculated in proximity to the bacteriocin assay, bacteriolysis was observed around the L⁺ A⁻ strain. The complementation experiment between the wild-type L⁺ A⁺ and mutant L⁻ A⁺ strains did not show any bacteriocin activ-

TABLE 4. Extracellular *trans*-complementation analysis of bacteriocin 41 activity^a

Plasmid(s)	Genotype	Phenotype ^b	pMG1105-42, pMG1105-25	pMG1106, pMG1109	pMG1105-107, pMG1105-188	pMG1105-104, pMG1105-11	pHT1101	pMG1110	pMG1111
pMG1105-42, pMG1105-25	<i>bacl</i> ₁ L ₂ ⁺ A ⁺ I ⁺	L ⁻ A ⁺ I ⁺	NT						
pMG1106, pMG1109	<i>bacl</i> ₁ L ₂ ⁺ A ⁺ I ⁺	L ⁻ A ⁺ I ⁺	C ⁻	NT					
pMG1105-107, pMG1105-188	<i>bacl</i> ₁ ⁺ L ₂ A ⁺ I ⁺	L ⁻ A ⁺ I ⁺	C ⁻	C ⁻	NT				
pMG1105-104, pMG1105-11	<i>bacl</i> ₁ ⁺ L ₂ ⁺ A ⁺ I ⁺	L ⁺ A ⁻ I ⁺	C ⁺	C ⁺	C ⁺	NT			
pHT1101	<i>bacl</i> ₁ ⁺ L ₂ ⁺ A ⁺ I	L ⁺ A ⁻ I ⁻	C ⁺	C ⁺	C ⁺	C ⁻	NT		
pMG1110	<i>bacl</i> ₁ ⁺ L ₂ ⁺ A ⁺ I	L ⁺ A ⁻ I ⁻	C ⁺	C ⁺	C ⁺	C ⁻	C ⁻	NT	
pMG1111	<i>bacl</i> ₁ L ₂ A ⁺ I	L ⁻ A ⁺ I ⁻	C ⁻	C ⁻	C ⁻	C ⁺	C ⁺	C ⁺	NT

^a C⁺; bacteriocin activity was detected at the intersection of the two strains by the soft-agar assay; C⁻; no bacteriocin activity; NT; not tested.

^b L, expression of lytic protein(s); A, activator expression; I, immunity expression.

A

putative processing site
↓
signal peptide

ORF7 (BacL1)	1	MNYSQKAILD CRYKSNFSLK AVAGRNGIL- -SIGVGHFTN EKHPIKPGMV ITESQATQIL RDDLNEHAAL ISKLLAIKAT	78
Lysozyme	1	MQISQAGINL IKSFEGLQLK AYKAVPTKHK YTYGYRHYGS DVSPRQ---V ITAKQAEQML RDOVQAFVGD VHKALKVSVT	77
ORF7 (BacL1)	79	QNOFDALVSF SRSKGLGFLP SSDIMHFTNN KEFNSAAREM KLVVYDIGSI KLPKLVERRN AETALYLEGA SGNEETTNRHA	158
Lysozyme	78	QNOFDALVSF AYNVGLGAFR SSSLLEYLNE GRITALAAEF PRWNKSGGKY YQG-LVNRRA QEQALFNSGT FKNV-----	150
ORF7 (BacL1)	159	-RIGFDVMIR WMEQKQAQHI TYSMDYRLGP NSYDCSSAVY FALKEAGFID PSTFPGNTDS LFGQLERVGW SQVFLVGGKY	247
Lysin (lambda Sa1)	1	MVINTEQATA WMAARQK-KV TYSMDYRNGP SBYDCSSSVY FALRSAGASD -NGMAVNTVEY EHDWLIKNGY VLI-AENTNW	77
ORF7 (BacL1)	250	HVQRGDFIW GIRGNSSGEL GHTGIFIDDK DNIHCTCGW DGNKCSINGI SVDNRDQVMV ASGRPPVTIY RFGGASKPYP	317
Lysin (lambda Sa1)	80	NAQRGDFIW GKRASAGAF GHTGMFV-DP DNIHCHNYG -----NSI TVNHRDEIHW YMGQPFVYAY RYARQKSNAX	149
ORF7 (BacL1)	318	GDSSGKSGDS -VNPSSAGVYF PSMRLPVSGD TDPNSPALDY YEAGQAIYVD SYVFANGYAW ISYVAGSGLR RYVAVGPDGD RTDTVWGTGF IN	408
Muramidase	554	GDEVGSVAKP DVVATSGGSR PTKTAKRSS PATSATTVGS YNAGDTVYNA GKVTINGQTW LRYMSYSGAQ HYVOISGEST STNVKQPVPT PQ	637
ORF7 (BacL1)	409	NTPSGSGST GSALSQVYFP SMRLPVSGDT DPNSPALAYY EAGQAIYDS YVFANGYAWI SYIAGSGLRR YVAVGPDGDR TDTVWGTGFF DN	500
Muramidase	638	-----SCSYRF TQTTAKNTP AGNAPSVTGY SAGDTVYNA KVTANGQTWL RYLSYSGAQH YVAV---SGN AAT-----	710
ORF7 (BacL1)	501	GGDPSQANP NSIGLVPKAG NFPVNRKLPV SAOTDPNSAA LDYYEAGQSI GYDSYIFANG YAMISYIAGS GLRHYVAVGP DDGRDTPVWG KGFN	595
Muramidase	711	-GNITSKPVT NSQG---AF RFTVTTNIRT APST---RASV VGEYNPGETV YVNGTVQAEQ YTWLRVLSRS GATHYVA	781

B

ORF7 (BacL1)	321	SGSKGDSVNP -SAGVYFPM RLPVSGDTP NSPALDYIE AQQAIYDYS VFANGYAWIS YVAGSGLRRY VAVGPDGDRT DTVWGTGFLN NTF	411
ORF7 (BacL1)	412	SGSGSTGSA -LSGVYFPM RLPVSGDTP NSPALAYE AQQAIYDYS VFANGYAWIS YIAGSGLRRY VAVGPDGDRT DTVWGTGFFD NGGDPGS	506
ORF7 (BacL1)	507	QAHFNSIGLV PKAGNFPVNR KLPVSAADTP NSPALDYIE AGQSIGYDYS IFANGYAWIS YIAGSGLRRY VAVGPDGDRT DTVWGTGFFN	595

FIG. 4. Comparison of the amino acid sequence of the predicted BacL₁ protein (ORF7) of bacteriocin 41 with the amino acid sequence of the cell wall lytic enzymes of gram-positive bacteria (A) and the repeat sequences found in the BacL₁ protein (B). Lysozyme, *B. subtilis* bacteriophage B103 (accession number Q37896); lysin, *S. galactiae* prophage lambda Sa1 (accession number NP 687631); muramidase, *L. plantarum* WCFS1 (accession number CAD64901).

ity. These results suggested that the activator of component A modified component L, that the activated component L possessed the bacteriocin activity, and also that an excess of component A existed in the extracellular medium.

The β -hemolysin/bacteriocin (cytolysin) determinant encoded on pAD1 consists of the eight genes *cylR2*, *cylR1*, *cylL₁*, *cylL₂*, *cylM*, *cylB*, *cylA*, and *cylI* (2, 8, 9, 17, 18, 39). CylL₁ and CylL₂ are the cytolysin structural subunits. The CylL₁ and CylL₂ proteins are modified posttranslationally by CylM₁ (2), and the modified CylL₁ and CylL₂ proteins are secreted via CylE₁, which is the ATP-binding exporter (16). The extracellular cytolysin precursors CylL₁ and CylL₂ are converted to the active cytolysin by CylA (2, 22). In an early study of the β -hemolysin/bacteriocin (cytolysin) determinant (22), two functional domains within the operon were identified and it was found that one region encodes the toxin precursor L component, which is now known to be encoded by CylL₁, CylL₂, CylM, and CylB, and the other region encodes an activator A component, which is now known to be encoded by CylA and CylI (2, 8, 9, 17, 18, 39). In the complementation experiment between the A component-producing strain or the wild-type strain and the L component-producing strain on blood agar plates, the β -hemolysis zone occurred around or along the L component-producing strain (22), indicating that the A component activates the L component extracellularly and that the activated L component possesses the β -hemolysin/bacteriocin activity and an excess of extracellular A component is present

in the culture medium of the wild-type strain (24). These observations are similar to the extracellular complementation observed between the L component-producing strain and the A component-producing strain for bacteriocin 41.

The deduced amino acid sequence encoded by *bacL₁* showed a high degree of homology with the cell wall lytic enzymes and murein hydrolases of lysozyme, lysine, and the muramidase of gram-positive bacteria (32). These enzymes cleave glycan chains either between the *N*-acetylmuramic acid and *N*-acetylglucosamine or at the alternative acetylglucosamine-muramic acid glycoside linkage (34). Sequence alignments of the murein hydrolases of the gram-positive bacteria show that most of these enzymes display a domain structure. In general, these enzymes harbor an N-terminal signal peptide, followed by a second domain containing the enzymatic activity. In addition, these proteins harbor repeat structures or cell wall-targeting structures that flank either the N- or C-terminal side of the enzymatic domain (40). The repeated domains direct the murein hydrolase to its receptor on the cell surface of gram-positive bacteria (51). Murein hydrolase is usually synthesized as a proenzyme, and after cleavage of the N-terminal signal peptide, the soluble proenzyme is secreted into the extracellular environment. The repeated domains or cell wall-targeting domains direct the proenzyme to its receptor on the bacterial cell surface. Proteolytic cleavage or activation of the proenzyme generates the mature enzyme (32).

Although the mechanism of activation or the precise mode of action of the *bacL₁*-encoded protein is not known, analysis

6. Clewell, D. B., and G. M. Dunny. 2002. Conjugation and genetic exchange in enterococci, p. 265–300. In M. S. Gilmore, D. B. Clewell, P. Courvalin, G. M. Dunne, B. E. Murray, and L. B. Rice, (ed.), *The enterococci: pathogenesis, molecular biology and antibiotic resistance*. ASM Press, Washington, DC.
7. Clewell, D. B., P. K. Tomich, M. C. Gawron-Burke, A. E. Franke, Y. Yagi, and F. Y. An. 1982. Mapping of *Streptococcus faecalis* plasmids pAD1 and pAD2 and studies relating to transposition of Tn917. *J. Bacteriol.* 152:1220–1230.
8. Coburn, P. S., L. E. Hancock, M. C. Booth, and M. S. Gilmore. 1999. A novel means of self-protection, unrelated to toxin activation, confers immunity to the bactericidal effects of the *Enterococcus faecalis* cytotoxin. *Infect. Immun.* 67:3339–3347.
9. Coburn, P. S., C. M. Pillar, B. D. Jett, W. Haas, and M. S. Gilmore. 2004. *Enterococcus faecalis* senses target cells and in response expresses cytotoxin. *Science* 306:2270–2272.
10. Coque, T. M., J. E. Patterson, J. M. Steckelberg, and B. E. Murray. 1995. Incidence of hemolysin, gelatinase, and aggregation substance among enterococci isolated from patients with endocarditis and other infections and from feces of hospitalized and community-based persons. *J. Infect. Dis.* 171:1223–1229.
11. Dunny, G. M., B. L. Brown, and D. B. Clewell. 1978. Induced cell aggregation and mating in *Streptococcus faecalis*: evidence for a bacterial sex pheromone. *Proc. Natl. Acad. Sci. USA* 75:3479–3483.
12. Dunny, G. M., R. A. Craig, R. L. Carron, and D. B. Clewell. 1979. Plasmid transfer in *Streptococcus faecalis*: production of multiple sex pheromones by recipients. *Plasmid* 2:454–465.
13. Fujimoto, S., H. Hashimoto, and Y. Ike. 1991. Low cost device for electrotransformation and its application to the highly efficient transformation of *Escherichia coli* and *Enterococcus faecalis*. *Plasmid* 26:131–135.
14. Fujimoto, S., H. Tomita, E. Wakamatsu, K. Tanimoto, and Y. Ike. 1995. Physical mapping of the conjugative bacteriocin plasmid pPD1 of *Enterococcus faecalis* and identification of the determinant related to the pheromone response. *J. Bacteriol.* 177:5574–5581.
15. Franz, C. M. A. P., M. J. van Belkum, W. H. Holzapfel, H. Abriouel, and A. Galvez. 2007. Diversity of enterococcal bacteriocins and their grouping in a new classification scheme. *FEMS Microbiol. Rev.* 31:293–310.
16. Gilmore, M. S., R. A. Segarra, and M. C. Booth. 1990. An HlyB-type function is required for expression of the *Enterococcus faecalis* hemolysin/bacteriocin. *Infect. Immun.* 58:3914–3923.
17. Gilmore, M. S., R. A. Segarra, M. C. Booth, C. P. Bogie, L. R. Hall, and D. B. Clewell. 1994. Genetic structure of the *Enterococcus faecalis* plasmid pAD1-encoded cytolytic toxin system and its relationship to lantibiotic determinants. *J. Bacteriol.* 176:7335–7344.
18. Haas, W., B. D. Shepard, and M. S. Gilmore. 2002. Two-component regulator of *Enterococcus faecalis* cytotoxin responds to quorum-sensing autoinduction. *Nature* 415:84–87.
19. Hanski, E., P. A. Horvitz, and M. G. Caparon. 1992. Expression of protein F, the fibronectin-binding protein of *Streptococcus pyogenes* JRS4, in heterologous streptococcal and enterococcal strains promotes their adherence to respiratory epithelial cells. *Infect. Immun.* 60:5119–5125.
20. Ike, Y., and D. B. Clewell. 1984. Genetic analysis of the pAD1 pheromone response in *Streptococcus faecalis*, using transposon Tn917 as an insertional mutagen. *J. Bacteriol.* 158:777–783.
21. Ike, Y., and D. B. Clewell. 1992. Evidence that the hemolysin/bacteriocin phenotype of *Enterococcus faecalis* subsp. *zymogenes* can be determined by plasmids in different incompatibility groups as well as by the chromosome. *J. Bacteriol.* 174:8172–8177.
22. Ike, Y., D. B. Clewell, R. A. Segarra, and M. S. Gilmore. 1990. Genetic analysis of the pAD1 hemolysin/bacteriocin determinant in *Enterococcus faecalis*: Tn917 insertional mutagenesis and cloning. *J. Bacteriol.* 172:155–163.
23. Ike, Y., R. A. Craig, B. A. White, Y. Yagi, and D. B. Clewell. 1983. Modification of *Streptococcus faecalis* sex pheromones after acquisition of plasmid DNA. *Proc. Natl. Acad. Sci. USA* 80:5369–5373.
24. Ike, Y., S. E. Flannagan, and D. B. Clewell. 1992. Hyperhemolytic phenomena associated with insertions of Tn916 into the hemolysin determinant of *Enterococcus faecalis* plasmid pAD1. *J. Bacteriol.* 174:1801–1809.
25. Ike, Y., H. Hashimoto, and D. B. Clewell. 1984. Hemolysin of *Streptococcus faecalis* subspecies *zymogenes* contributes to virulence in mice. *Infect. Immun.* 45:528–530.
26. Ike, Y., H. Hashimoto, and D. B. Clewell. 1987. High incidence of hemolysin production by *Enterococcus (Streptococcus) faecalis* strains associated with human parenteral infection. *J. Clin. Microbiol.* 25:1524–1528.
27. Jack, R. W., J. R. Tagg, and B. Ray. 1995. Bacteriocins of gram-positive bacteria. *Microbiol. Rev.* 59:171–200.
28. Jedrzejewski, M. K. 2001. Pneumococcal virulence factors: structure and function. *Microbiol. Mol. Biol. Rev.* 65:187–207.
29. Jett, B. D., H. G. Jensen, R. E. Nordquist, and M. S. Gilmore. 1992. Contribution of the pAD1-encoded cytotoxin to the severity of experimental *Enterococcus faecalis* endophthalmitis. *Infect. Immun.* 60:2445–2452.
30. Kleerebezem, M., J. Boekhorst, R. van Kranenburg, D. Molenaar, O. P. Kuipers, R. Leer, R. Turchini, S. A. Peters, H. M. Sandbrink, M. W. E. J. Fiers, W. Stiekema, R. M. Klein Lankhorst, P. A. Bron, S. M. Hoffer, M. N. Nierop Groot, R. Kerkhoven, M. de Vries, B. Ursing, W. M. de Vos, and R. J. Siezen. 2003. Complete genome sequence of *Lactobacillus plantarum* WCFS1. *Proc. Natl. Acad. Sci. USA* 100:1990–1995.
31. LeBlanc, D. J., L. N. Lee, D. B. Clewell, and D. Behnke. 1983. Broad geographical distribution of a cytotoxin gene mediating beta-hemolysis and bacteriocin activity among *Streptococcus faecalis* strains. *Infect. Immun.* 40:1015–1022.
32. López, R., and E. García. 2004. Recent trends on the molecular biology of pneumococcal capsules, lytic enzymes, and bacteriophage. *FEMS Microbiol. Rev.* 28:553–580.
33. Martínez-Bueno, M., M. Maqueda, A. Galvez, B. Samyn, J. V. Beeumen, J. Coyette, and E. Valdivia. 1994. Determination of the gene sequence and the molecular structure of the enterococcal peptide antibiotic AS-48. *J. Bacteriol.* 176:6334–6339.
34. Navarre, W. W., and O. Schneewind. 1999. Surface proteins of gram-positive bacteria and mechanisms of their targeting to the cell wall envelope. *Microbiol. Mol. Biol. Rev.* 63:174–229.
35. Nes, I. F., D. B. Diep, and H. Holo. 2007. Bacteriocin diversity in *Streptococcus* and *Enterococcus*. *J. Bacteriol.* 189:1189–1198.
36. Nilsen, T., I. F. Nes, and H. Holo. 2003. Enterolysin A, a cell wall-degrading bacteriocin from *Enterococcus faecalis* LMG 2333. *Appl. Environ. Microbiol.* 69:2975–2984.
37. Pecenkova, T., V. Benes, J. Paces, C. Vlcek, and V. Paces. 1997. Bacteriophage B103: complete DNA sequence of its genome and relationship to other *Bacillus* phages. *Gene* 199:157–163.
38. Sambrook, J., and D. W. Russell. 2001. *Molecular cloning: a laboratory manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
39. Segarra, R. A., M. C. Booth, D. A. Morales, M. M. Huycke, and M. S. Gilmore. 1991. Molecular characterization of the *Enterococcus faecalis* cytotoxin activator. *Infect. Immun.* 59:1239–1246.
40. Sheehan, M. M., J. L. Garcia, R. López, and P. Garcia. 1997. The lytic enzyme of the pneumococcal phage Dp-1: a chimeric lysin of intergeneric origin. *Mol. Microbiol.* 25:717–725.
41. Shiojima, M., H. Tomita, K. Tanimoto, S. Fujimoto, and Y. Ike. 1997. High-level plasmid-mediated gentamicin resistance and pheromone response of plasmids present in clinical isolates of *Enterococcus faecalis*. *Antimicrob. Agents Chemother.* 41:702–705.
42. Tanimoto, K., and T. Iino. 1985. Additional genes essential for replication of the mini-F plasmid from origin I. *Mol. Gen. Genet.* 198:358–359.
43. Tettelin, H., V. Masignani, M. J. Clesiewicz, J. A. Eisen, S. Peterson, M. R. Wessels, I. T. Paulsen, K. E. Nelson, I. Margarit, T. D. Read, L. C. Madoff, A. M. Wolf, M. J. Beanan, L. M. Brinkac, S. C. Daugherty, R. T. DeBoy, A. S. Durkin, J. F. Kolonay, R. Madupu, M. R. Lewis, D. Radune, N. B. Fedorova, D. Scanlan, H. Khouri, S. Mulligan, H. A. Carty, R. T. Cline, S. E. Van Aken, J. Gill, M. Scarselli, M. Mora, E. T. Jacobini, C. Brettoni, G. Galli, M. Mariani, F. Vegni, D. Maione, D. Rinaudo, R. Rappuoli, J. L. Telford, D. L. Kasper, G. Grandi, and C. M. Fraser. 2002. Complete genome sequence and comparative genomic analysis of an emerging human pathogen, serotype V *Streptococcus agalactiae*. *Proc. Natl. Acad. Sci. USA* 99:12391–12396.
44. Tomich, P. K., F. Y. An, and D. B. Clewell. 1980. Properties of erythromycin-inducible transposon Tn917 in *Streptococcus faecalis*. *J. Bacteriol.* 141:1366–1374.
45. Tomita, H., and D. B. Clewell. 2000. A pAD1-encoded small RNA molecule, mD, negatively regulates *Enterococcus faecalis* pheromone response by enhancing transcription termination. *J. Bacteriol.* 182:1062–1073.
46. Tomita, H., S. Fujimoto, K. Tanimoto, and Y. Ike. 1996. Cloning and genetic organization of the bacteriocin 31 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pY117. *J. Bacteriol.* 178:3585–3593.
47. Tomita, H., S. Fujimoto, K. Tanimoto, and Y. Ike. 1997. Cloning and genetic and sequence analysis of the bacteriocin 21 determinant encoded on the *Enterococcus faecalis* pheromone-responsive conjugative plasmid pPD1. *J. Bacteriol.* 179:7843–7855.
48. Tomita, H., and Y. Ike. 2005. Genetic analysis of transfer-related regions of the vancomycin resistance *Enterococcus* conjugative pHTB: identification of *oriT* and a putative relaxase gene. *J. Bacteriol.* 187:7727–7737.
49. Tomita, H., C. Pierson, S. K. Lim, D. B. Clewell, and Y. Ike. 2002. Possible connection between a widely disseminated conjugative gentamicin resistance (pMG1-like) plasmid and the emergence of vancomycin resistance in *Enterococcus faecium*. *J. Clin. Microbiol.* 40:3326–3333.
50. Wirth, R., F. Y. An, and D. B. Clewell. 1986. Highly efficient protoplast transformation system for *Streptococcus faecalis* and a new *Escherichia coli*-*S. faecalis* shuttle vector. *J. Bacteriol.* 165:831–836.
51. Wren, B. W. 1991. A family of clostridial and streptococcal ligand-binding proteins with conserved C-terminal repeat sequences. *Mol. Microbiol.* 5:797–803.

Identification and Characterization of Two Novel Methyltransferase Genes That Determine the Serotype 12-Specific Structure of Glycopeptidolipids of *Mycobacterium intracellulare*[†]

Noboru Nakata,^{1*} Nagatoshi Fujiwara,² Takashi Naka,³ Ikuya Yano,³
Kazuo Kobayashi,⁴ and Shinji Maeda⁵

Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, Tokyo, Japan¹; Department of Host Defense, Osaka City University Graduate School of Medicine, Osaka, Japan²; Japan BCG Laboratory, Tokyo, Japan³; Department of Immunology, National Institute of Infectious Diseases, Tokyo, Japan⁴; and Molecular Epidemiology Division, Mycobacterium Reference Center, The Research Institute of Tuberculosis, Japan Anti-Tuberculosis Association, Tokyo, Japan⁵

Received 23 August 2007/Accepted 5 November 2007

The *Mycobacterium avium* complex is distributed ubiquitously in the environment. It is an important cause of pulmonary and extrapulmonary diseases in humans and animals. The species in this complex produce polar glycopeptidolipids (GPLs); of particular interest is their serotype-specific antigenicity. Several reports have described that GPL structure may play an important role in bacterial physiology and pathogenesis and in the host immune response. Recently, we determined the complete structure of the GPL derived from *Mycobacterium intracellulare* serotype 7 and characterized the serotype 7 GPL-specific gene cluster. The structure of serotype 7 GPL closely resembles that of serotype 12 GPL, except for O methylation. In the present study, we isolated and characterized the serotype 12-specific gene cluster involved in glycosylation of the GPL. Ten open reading frames (ORFs) and one pseudogene were observed in the cluster. The genetic organization of the serotype 12-specific gene cluster resembles that of the serotype 7-specific gene cluster, but two novel ORFs (*orfA* and *orfB*) encoding putative methyltransferases are present in the cluster. Functional analyses revealed that *orfA* and *orfB* encode methyltransferases that synthesize O-methyl groups at the C-4 position in the rhamnose residue next to the terminal hexose and at the C-3 position in the terminal hexose, respectively. Our results show that these two methyltransferase genes determine the structural difference of serotype 12-specific GPL from serotype 7-specific GPL.

The *Mycobacterium avium* complex (MAC) consists of two species, *M. avium* and *Mycobacterium intracellulare*, which are opportunistic pathogens of humans and animals. Human exposure to the MAC is common because organisms of this complex are ubiquitous in the environment: they have been isolated from water, soil, plants, house dust, and other sources. In fact, the MAC is the most common cause of disease attributable to nontuberculous mycobacteria in humans (9). The majority of MAC infections are acquired environmentally, and person-to-person transmission is considered to be rare. The treatment of MAC infection is difficult because the organisms are often resistant to standard antituberculosis drugs.

Many antigenic or immunoregulatory glycolipids with structural diversity are expressed on the mycobacterial cell wall. These molecules are considered to be involved in bacterial virulence through host immune responses (5, 14, 22, 23). It is necessary to elucidate the molecular structure, biochemical characteristics, and biological functions of the lipid components to better understand the mechanisms of pathogenesis and drug resistance of the MAC. The most prominent feature

of the MAC is the presence of antigenic glycolipids, the glycopeptidolipids (GPLs), which are present on the cell surface (1). The standard method for differentiation of MAC strains is serologic typing based on the oligosaccharide (OSE) residue of the GPL. GPLs contain a tetrapeptide-amino alcohol core, D-phenylalanine-D-*allo*-threonine-D-alanine-L-alanine (D-Phe-D-*allo*-Thr-D-Ala-L-alanine), with an amido-linked 3-hydroxy or 3-methoxy C₂₆-to-C₃₄ fatty acid at the N terminus of D-Phe (4). The D-*allo*-Thr and terminal L-alanine are further linked with 6-deoxy-talose (6-d-Tal) and 3,4-di-O-methyl-rhamnose (3,4-di-O-Me-Rha), respectively. This core GPL is present in all species of the MAC and shows a common antigenicity (1). In the serotype-specific GPLs, a haptenic OSE is linked with the 6-d-Tal residue. To date, 31 distinct serotype-specific polar GPLs have been identified biochemically; the complete structures of GPLs are partly defined for serotype 1 to 4, 7, 8, 9, 12, 14, 17, 19 to 21, 25, and 26 GPLs (7, 10). On the other hand, it has been reported that serotype-specific GPLs participate in pathogenesis and immunomodulation in the host (2, 13). Modification of the GPL structure might play an important role not only in antigenicity but also in host immune responses and bacterial physiology (18). Recently, chemical synthesis of various haptenic OSEs was demonstrated, and the genes encoding glycosylation pathway enzymes for the biosynthesis of GPLs were identified and characterized (8, 12, 19, 21). However, genes responsible for serotype-specific glycosylation have yet to be analyzed for most of the serotypes.

* Corresponding author. Mailing address: Department of Microbiology, Leprosy Research Center, National Institute of Infectious Diseases, 4-2-1 Aoba-cho, Higashimurayama, Tokyo 189-0002, Japan. Phone: 81 (42) 391 8211. Fax: 81 (42) 394 9092. E-mail: n-nakata@nih.go.jp.

[†] Published ahead of print on 16 November 2007.




FIG. 1. Structures of serotype 7 and 12 GPLs. *O*-methyl groups specific to the serotypes are indicated by arrows.

In a previous study, we determined the complete structure of the GPL derived from *M. intracellulare* serotype 7 and characterized the serotype 7-specific gene cluster for GPL synthesis (10). The structure of serotype 7 GPL closely resembles that of serotype 12 GPL, except for *O* methylation (Fig. 1). In the present study, we determined the nucleotide sequence of the serotype 12-specific gene cluster involved in the glycosylation of the GPL and characterized two novel open reading frames (ORFs) encoding *O*-methyltransferases that determine the difference of serotype 12 GPL from serotype 7 GPL.

MATERIALS AND METHODS

Bacterial strains and construction of *M. intracellulare* cosmid library. *M. intracellulare* serotype 12 strain ATCC 35762 (NF 103), *M. intracellulare* serotype 7 strain ATCC 35847 (NF 027), and *M. intracellulare* serotype 7 strain NF 112 were used for this study. A cosmid library of *M. intracellulare* NF 103 was constructed as described previously (10). Briefly, genomic DNA of *M. intracellulare* NF 103 was prepared by mechanical disruption of bacterial cells in phosphate-buffered saline containing 50 mM EDTA, followed by phenol-chloroform extraction and precipitation with ethanol. Genomic DNA fragments randomly sheared to 30-kb to 50-kb fragments during the extraction process were fractionated and electroeluted from agarose gels. These DNA fragments were ligated to dephosphorylated arms of pYUB412 (XbaI-EcoRV and EcoRV-XbaI). After *in vitro* packaging using Gigapack III Gold extracts (Stratagene, La Jolla, CA), recombinant cosmids were introduced into *Escherichia coli* STBL2.

Isolation of cosmid clones carrying the GPL biosynthesis gene cluster and sequence analysis. PCR was used to isolate cosmid clones carrying the rhamnosyltransferase gene (*rtfA*), using primers *rtfA*-F (5'-TTTGGAGCGACGAGTTCATC-3') and *rtfA*-R (5'-GTGTAGTTGACACCGCCGAC-3'). The insert of cosmid clone 161 was sequenced using a kit (BigDye Terminator cycle sequencing kit,

version 3.1; Applied Biosystems, Foster City, CA) and a sequence analyzer (ABI Prism 310; Applied Biosystems). The putative function of each ORF was identified by similarity searches between the deduced amino acid sequences and those of known proteins, using BLAST (<http://www.ncbi.nlm.nih.gov/BLAST/>) and Frame-Plot (<http://www.nih.gov/jp-jun/cgi-bin/frameplot.pl>) with the DNASIS computer program (Hitachi Software Engineering, Yokohama, Japan).

Transformation of *M. intracellulare*. PCR was used to amplify and clone *orfA* and *orfB* into the plasmid vector pVVI6. *M. intracellulare* NF 027 and NF 112 were transformed with the resultant plasmids by electroporation. Primers used to amplify *orfA*, *orfB*, and *orfA-orfB* were *orfA*-F (5'-GCGGATCCAGTGTGCAGACGAGCGGAAC-3'), *orfA*-R (5'-GCGAATTCCTATCCGAGAAAAATAAAAG-3'), *orfB*-F (5'-GCGGATCCACTGTAGACTCCGCCACCAT-3'), and *orfB*-R (5'-GCGAATTCCTACACCTTCACGGCGGATC-3').

Preparation of GPLs and OSE moieties. GPL 7 and GPL 12 were purified from *M. intracellulare* NF 027 and NF 103, respectively. The preparation of GPLs was performed as described previously (10, 15, 17). Briefly, each strain was grown in Middlebrook 7H9 broth (Difco Laboratories, Detroit, MI) with 0.5% glycerol and 10% Middlebrook oleic acid-albumin-dextrose-catalase enrichment (Difco Laboratories) at 37°C for 2 to 3 weeks. The heat-killed bacteria were sonicated and extracted using chloroform-methanol (2:1 [vol/vol]). The extractable lipids were hydrolyzed with 0.2 N sodium hydroxide in methanol at 37°C for 2 h. After neutralization using 6 N hydrochloric acid, chloroform-methanol (2:1 [vol/vol]) and water were added. The organic phase containing alkaline-stable lipids was recovered and evaporated, with subsequent addition of acetone to remove any acetone-insoluble components. The supernatant was dried up. It was then treated using a Sep-Pak silica cartridge (Waters Corp., Milford, MA) with washing (chloroform-methanol [95:5 (vol/vol)]) and elution (chloroform-methanol [1:1 (vol/vol)]) for partial purification. The GPL was then purified completely by preparative thin-layer chromatography (TLC) with silica gel G (Uniplate; 20 cm × 20 cm × 250 μm; Analtech, Inc., Newark, DE). The TLC was developed repeatedly, using chloroform-methanol-water (60:16:2 [vol/vol/vol]), until a single spot was obtained. To prepare the OSE moiety, purified GPL was processed using β-elimination with alkaline borohydride, and then the carbohydrate chain moiety

TABLE 1. Similarity of Orfs in *M. intracellulare* serotype 12 strain ATCC 35762 to known protein sequences

Orf	Predicted molecular mass (Da)	Predicted pI	Similar protein	Identity (no. of matched amino acids/total no. of amino acids)	E value	GenBank accession no.
GtfB	45,830	6.87	Glycosyltransferase GtfB	412/418	0.0	BAF45360
Orf1	45,203	6.10	Putative glycosyltransferase	414/417	0.0	BAF45361
OrfA	28,904	7.42	Putative methyltransferase	182/224	5e-88	NP_218045
OrfB	29,930	5.15	Putative methyltransferase	102/204	1e-19	EAZ88812
Orf3	32,151	10.41	Putative glycosyltransferase	196/223	1e-108	BAF45363
Orf4	40,742	5.41	Putative aminotransferase	338/374	0.0	BAF45364
Orf5	35,812	5.26	Hypothetical protein	303/329	4e-162	BAF45365
Orf7	27,693	5.99	Putative metallophosphoesterase	223/241	1e-122	BAF45367
Tn	28,538	11.85	Putative transposase	213/255	6e-107	AAL61662
Orf8	80,044	9.16	Putative acyltransferase	689/747	0.0	BAF45368
Orf9	37,797	8.26	Putative glycosyltransferase	310/337	7e-169	BAF45369
DrrC	28,549	12.01	Daunorubicin resistance protein C	261/263	3e-141	BAF45370




FIG. 2. Comparison of genetic organization of GPL biosynthesis clusters. (a) *M. avium* strain A5 organization, based on the annotated sequence obtained from GenBank (accession no. AY130970). (b) *M. intracellulare* ATCC 35762 (NF 103), sequenced in this study. (c) *M. intracellulare* ATCC 35847 (NF 027), sequenced in our previous study (GenBank accession no. AB274811). The orientation of each gene is shown by the arrow direction. The black arrows represent mobile elements, and the gray arrow represents a pseudogene. Mutually homologous ORFs and sequences are indicated with dotted lines.

elongated from *D*-allo-Thr was released as described previously (10, 15). Briefly, GPL was treated with 5 mg/ml sodium borohydride or borodeuteride in 0.5 N sodium hydroxide-ethanol (1:1 [vol/vol]) at 60°C for 16 h, with stirring. The reaction mixture was decanted with Dowex 50W X8 beads (The Dow Chemical Company, Midland, MI). The supernatant was collected and evaporated under nitrogen to remove boric acid. The dried residue was partitioned into two layers, using chloroform-methanol (2:1 [vol/vol]) and water. The upper aqueous phase was recovered and evaporated. In these processes, the OSE was purified as an oligoglycosyl alditol.

MALDI-TOF MS and MALDI-TOF/TOF MS analyses. The molecular species of the intact GPLs were detected using matrix-assisted laser desorption/ionization-time-of-flight mass spectrometry (MALDI-TOF MS) with an Ultraflex II spectrophotometer (Bruker Daltonics, Billerica, MA). Each GPL was dissolved in chloroform-methanol (2:1 [vol/vol]) at a concentration of 1 mg/ml; 1 μ l of a sample was then applied directly to the sample plate, followed by the addition of 1 μ l of 10-mg/ml 2,5-dihydroxybenzoic acid in chloroform-methanol (1:1 [vol/vol]) as a matrix. The intact GPL was analyzed in the reflectron mode, with an accelerating voltage operating in positive mode at 20 kV (3). The OSE was analyzed by the fragment pattern with MALDI-TOF/TOF MS to determine the glycosyl composition. The OSE was dissolved with ethanol-water (3:7 [vol/vol]); the matrix was 10 mg/ml 2,5-dihydroxybenzoic acid in ethanol-water (3:7 [vol/vol]). The OSE and matrix were added to the sample plate by the same method as that for intact GPL. They were then analyzed in the lift-lift mode.

GC-MS analyses of alditol acetate derivatives. Gas chromatography (GC) and GC-MS analyses of partially methylated alditol acetate derivatives were performed to determine glycosyl compositions and linkage positions. Perdeuteromethylation was conducted using a modified procedure of Hakomori, as described previously (10, 11). Briefly, the dried OSE was dissolved with a mixture of dimethyl sulfoxide and sodium hydroxide, and deuteromethyl iodide was added. The reaction mixture was stirred at room temperature for 15 min, followed by the addition of water and chloroform. After centrifugation at 2,400 \times g for 15 min, the upper water layer was discarded. The chloroform layer was washed twice with water and evaporated completely. To prepare partially deuteromethylated alditol acetates, perdeuteromethylated OSE was hydrolyzed using 2 N trifluoroacetic acid at 120°C for 2 h, reduced with 10 mg/ml sodium borodeuteride at 25°C for 2 h, and acetylated with acetic anhydride at 100°C for 1 h (6, 10, 16). GC-MS was then performed using a benchtop ion-trap mass spectrometer (Trace DSQ GC/MS; Thermo Electron Corporation, Austin, TX) equipped with a fused capillary column (30 m; 0.25-mm internal diameter) (Equity-1 or SP-2380; Supelco, Bellefonte, PA). Helium was used as the carrier gas, and the flow rate was 1 ml/min. The SP-2380 column was used for the analysis of alditol acetate derivatives. The temperature program was started at 60°C, with an increase of 40°C/min to 260°C and a hold at 260°C for 25 min. The Equity-1 column was used for analysis of perdeuteromethylated alditol acetate derivatives. The temperature program was 80°C for 1 min, with an increase of 20°C/min to 180°C followed by an increase of 8°C/min to 280°C.

Nucleotide sequence accession number. The nucleotide sequence reported here has been deposited in the NCBI GenBank database under accession number AB353739.

RESULTS

Cloning and sequence of the serotype 12 GPL biosynthesis cluster. To isolate the serotype 12-specific GPL biosynthesis gene cluster, a genomic cosmid library of an *M. intracellulare* serotype 12 strain, NF 103, was constructed. DNA was extracted from each clone by boiling. Using colony PCR with *rftA* primers, the positive clone 161 was isolated from the *E. coli* transductants. Sequencing analysis revealed that cosmid clone 161 carried the DNA region from *gfbB* to *gfbR*. Ten ORFs and one pseudogene other than *gfbB* and *gfbR* were observed in the cluster (Table 1 and Fig. 2). The genetic organization between the *gfbB* and *gfbR* genes (15.6 kb) of *M. intracellulare* NF 103 (serotype 12) closely resembled that of




FIG. 3. TLC patterns of alkaline-stable lipids derived from *M. intracellulare* serotype 7 transformants. GPL 7 and GPL 12 were purified from *M. intracellulare* serotype 7 strain ATCC 35847 (NF 027) and serotype 12 strain ATCC 35762 (NF103). TLC was developed with a solvent system of chloroform-methanol-water (65:25:4 [vol/vol/vol]). Circled spots indicate prominent GPLs.

the same region of *M. intracellulare* NF 027 (serotype 7), except for three loci (Fig. 2). The first difference between them was an additional ORF encoding a transposase between *orf7* and *orf8* in NF 103 (Fig. 2). The second difference was that the *orf6* homologous sequence in NF 103 had a frame shift, indicating that this locus does not encode a protein. The third difference is that two novel ORFs (*orfA* and *orfB*) instead of *orf2* were found between *orf1* and *orf3* in NF 103.

Functional analysis of the two unique ORFs found in the serotype 12 GPL biosynthesis cluster. Based on sequence homology, *orfA* and *orfB* were able to encode methyltransferases responsible for producing serotype 12 GPLs (Table 1). We constructed three plasmids carrying *orfA* and/or *orfB* downstream of the *hsp-60* promoter to test this. These plasmids and a control vector plasmid were introduced individually into *M. intracellulare* serotype 7 (NF 027 and NF 112), and transformants were obtained. The GPLs produced from each transformant were analyzed.

The alkaline-stable lipids derived from six transformants of NF 027 and NF 112 in addition to the control strains (vector only) were developed by TLC, and the produced GPLs were compared to the spots of GPL 7 and GPL 12 (Fig. 3). The R_f values for GPLs synthesized in NF 027 transformed with *orfA* and NF 027 transformed with *orfA* and *orfB* (GPL 7-*orfA* and GPL 7-*orfAB*, respectively) were almost identical to that for GPL 12; the R_f value for the GPL synthesized in NF 027 transformed with *orfB* (GPL 7-*orfB*) was intermediate between those of GPL 7 and GPL 12, although the GPL synthesized in the control strain (GPL vector) was not changed from GPL 7. These results suggest that *orfA*, *orfB*, and *orfA-orfB* introduced into serotype 7 strain NF 027 were expressed and that they functioned for the modification of GPLs. We investigated the structural definition of these modified GPLs.

The GPLs produced in the transformants were purified using preparative TLC; their molecular weights were measured using MALDI-TOF MS (Fig. 4). The main molecularly related ions of GPL 7 and GPL 12 were detected as m/z 1,897 and 1,911, respectively, for $[M + Na]^+$ (Fig. 4a and b). The predominant m/z values were 1,911 for GPL 7-*orfA*, 1,897 for GPL 7-*orfB*, and 1,911 for GPL 7-*orfAB* (Fig. 4c to e). The molecular weight of GPL 7-*orfB* was the same as that of GPL 7, and those of GPL 7-*orfA* and GPL 7-*orfAB* were equal to that of GPL 12. Next, MALDI-TOF/TOF MS analysis was performed to determine the glycosyl pattern, using fragment ions of glycosyl cleavage. The fragment ions of the GPL vector (equal to GPL 7) showed m/z 254, 400, 546, and 692 for cleavage in turn from terminal 4*N*-acyl-hexose (Hex) and 336, 482, and 628 for cleavage in the opposite direction from 6-*d*-Tal (Fig. 5a). The fragment ions of GPL 7-*orfA*, m/z 414 and 642, were different from those of GPL 7, i.e., m/z 400 and 628, respectively; they demonstrated that the mass number of the sugar next to the terminal Hex increased 14 mass units (Fig. 5b). This result suggests that the second sugar from the terminal one was changed from Rha to *O*-methyl rhamnose (*O*-Me-Rha). Similarly, the fragment pattern of GPL 7-*orfAB* was identical to that of GPL 7-*orfA*, although that of GPL 7-*orfB* was the same as that of GPL 7 (Fig. 5c and d). Altogether, GPL 7-*orfAB* was predicted to have a modification of the *O*-Me position in the terminal Hex along with the substitution of *O*-Me-Rha for Rha in the sugar next to the terminal Hex; GPL




FIG. 4. MALDI-TOF MS spectra of GPLs derived from *M. intracellulare* serotype 7, serotype 12, and serotype 7 transformants. a.u., absorbance units.

7-*orfB* was modified only at the *O*-Me position in the terminal Hex.

GC-MS analyses of alditol acetate and perdeuteromethyl alditol acetate derivatives were performed to assign the linkage position of *O*-Me. As portrayed in Fig. 6a and b, the alditol acetate derivatives of the second sugar from the terminal 4*N*-acyl-Hex in GPL 7-*orfA* and GPL 7-*orfB* were assigned to 1,2,3,5-tetra-acetyl-4-*O*-methyl-rhamnitol (m/z 99, 131, 159, 201, and 261) and 1,2,3,4,5-penta-acetyl-rhamnitol (m/z 115, 157, 187, 217, 231, 289, and 303), respectively. The perdeuteromethyl alditol acetate derivatives of the terminal sugar in GPL 7-*orfA* and GPL 7-*orfB* were assigned to 3-*O*-deuteromethyl-1,5-di-*O*-acetyl-4-2'-*O*-deuteromethyl-propanoyl-deuteromethylamido-4,6-dideoxy-2-*O*-methyl-hexitol (m/z 105, 118, 165, 209, 222, 269, and 300) and 2-*O*-deuteromethyl-1,5-di-*O*-acetyl-4-2'-*O*-deuteromethyl-propanoyl-deuteromethyl-amido-4,6-dideoxy-3-*O*-methyl-hexitol (m/z 105, 121, 165, 206, 222, 266, and 300), respectively (Fig. 6c and d). In particular, the fragment ions of m/z 118 and 269 (Fig. 6c) versus m/z 121 and 266 (Fig. 6d) strongly indicated the different positions of linkages 2-*O*-Me and 3-*O*-Me. The alditol acetate and perdeuteromethyl alditol acetate derivatives in GPL 7-*orfAB* were detected with the same patterns of 4*N*-acyl-4,6-dideoxy-3-*O*-Me-Hex and 4-*O*-Me-Rha. According to these results, all OSE structures in GPLs derived from three serotype 7 transfor-




FIG. 5. Fragment patterns of MALDI-TOF/TOF MS spectra of OSEs in GPLs derived from *M. intracellulare* serotype 7, serotype 7 transformants. The MALDI-TOF/TOF MS spectra were acquired using 10 mg/ml 2,5-dihydroxybenzoic acid in ethanol-water (3:7 [vol/vol]) as the matrix; the molecularly related ions were detected as $[M + Na]^+$ in lift-lift mode. The assigned fragment patterns of glycosyl residues are depicted. a.u., absorbance units.

mantants were assigned as listed in Table 2. Altogether, the functions of the two genes were defined. The *orfA* product transfers a methyl group to the C-4 position of Rha next to the terminal sugar, and the *orfB* product transfers a methyl group to the C-3 position of the terminal sugar (Fig. 7). The results demonstrated that GPL 7 in the serotype 7 strain was changed completely to GPL 12 by introduction of the *orfA-orfB* gene cluster.

DISCUSSION

Nontuberculous mycobacteria, including the pathogenic species belonging to the MAC, have serotype-specific GPLs that are important components of the outer layer of the lipid-rich cell walls (5). Structural analyses of some serotype-specific GPLs derived from predominant clinical isolates have been

reported (20). We recently determined the complete structure of serotype 7 GPL and the nucleotide sequence of the serotype 7-specific GPL biosynthesis cluster (10). In this cluster, Orfs 1, 3, and 9 might engender transfer of the two molecules of L-Rha and the terminal Hex of serotype 7 GPL (10). Orfs 4, 5, 7, and 8 are homologous to an aminotransferase, a carbamoyl phosphate synthase protein, a metallophosphoesterase, and an acyltransferase, respectively, and possibly relate to the biosynthesis of 2'-hydroxypropanoylamido in the terminal Hex. Based on analysis of sequence homology, these ORFs are probably responsible for the glycosylation of serotype 7 GPL. Serotype 12 GPL has a similar structure to that of serotype 7 GPL, except for O methylation (Fig. 1). In the present study, we cloned the serotype 12 GPL biosynthesis cluster and analyzed its sequence. Although the genetic organization of the *gfb-to-drcC*




FIG. 6. Preparative GC-MS spectra of alditol acetate (a and b) and perdeuteromethylated alditol acetate (c and d) derivatives. The patterns of prominent fragment ions are presented. An SP-2380 column was used for the analysis of alditol acetate derivatives. The temperature program was started at 60°C, with an increase of 40°C/min to 260°C and a hold at 260°C for 25 min. An Equity-1 column was used for the perdeuteromethylated alditol acetate derivatives. The temperature program was 80°C for 1 min, with an increase of 20°C/min to 180°C followed by an increase of 8°C/min to 280°C.

region of the serotype 12 GPL biosynthetic cluster closely resembled that of serotype 7, significant differences were found in three loci (Fig. 2). The *M. intracellulare* serotype 12 strain NF 103 had one ORF encoding a transposase between *orf7* and *orf8* and had an *orf6* homologous sequence with frameshift inactivation. Orf6 in *M. intracellulare* serotype 7 exhibits sequence similarity to nucleotide sugar epimerases/dehydrogenases, but NF 112, one of the *M. intracellulare* serotype 7 isolates, had an interrupted *orf6* (10). These findings suggest that *orf6* is not involved in biosynthesis of either serotype 7 GPL or serotype 12 GPL. The most important difference between the two serotypes is that *M. intracellulare* serotype 12 had two unique ORFs, *orfA* and *orfB*, instead of *orf2* in *M. intracellulare* serotype 7. Actually, Orf2 in *M. intracellulare* se-

rotype 7 was assigned to a methyltransferase and might be responsible for synthesis of the *O*-methyl group at the C-2 position in the terminal Hex. That possibility suggests that the two unique ORFs for serotype 12 encode *O*-methyltransferases that produce the serotype 12-specific structure. NF 027 (serotype 7) transformed with *orfA* produced 4*N*-acyl-4,6-dideoxy-2-*O*-Me-Hex→4-*O*-Me-Rha→Rha→Rha→6-d-Tal, indicating that the product from *orfA* had activity to synthesize an *O*-methyl group at C-4 in L-Rha next to the terminal Hex (Table 2 and Fig. 7). NF 027 transformed with *orfB* produced 4*N*-acyl-4,6-dideoxy-3-*O*-Me-Hex→Rha→Rha→Rha→6-d-Tal, indicating that the product from *orfB* had activity to synthesize an *O*-methyl group at C-3 in the terminal Hex. NF 027 transformed with *orfA* and *orfB* produced serotype 12-specific GPL,

TABLE 2. Summarized structures of OSEs derived from serotype 7 transformants

GPL	Molecular weight of OSE	Fragment ions in MALDI-TOF/TOF MS	O-Methyl group		Structure of OSE
			Terminal sugar	Residue next to terminal sugar	
GPL 7 vector	859	254, 400, 546, 692	2- <i>O</i> -Met		4 <i>N</i> -acyl-4,6-dideoxy-2- <i>O</i> -Me-Hex→Rha→Rha→Rha→6-d-Tal
GPL 7-orfA	873	254, 414, 560, 706	2- <i>O</i> -Met	4- <i>O</i> -Met	4 <i>N</i> -acyl-4,6-dideoxy-2- <i>O</i> -Me-Hex→4- <i>O</i> -Me-Rha→Rha→Rha→6-d-Tal
GPL 7-orfB	859	254, 400, 546, 692	3- <i>O</i> -Met		4 <i>N</i> -acyl-4,6-dideoxy-3- <i>O</i> -Me-Hex→Rha→Rha→Rha→6-d-Tal
GPL 7-orfAB	873	254, 414, 560, 706	3- <i>O</i> -Met	4- <i>O</i> -Met	4 <i>N</i> -acyl-4,6-dideoxy-3- <i>O</i> -Me-Hex→4- <i>O</i> -Me-Rha→Rha→Rha→6-d-Tal