

- 8 Mullen AC, High FA, Hutchins AS, Lee HW, Villarino AV, Livingston DM *et al.* Role of T-bet in commitment of TH1 cells before IL-12-dependent selection. *Science* 2001; **292**: 1907–1910.
- 9 Afkarian M, Sedy JR, Yang J, Jacobson NG, Cereb N, Yang SY *et al.* T-bet is a STAT1-induced regulator of IL-12R expression in naive CD4+ T cells. *Nat Immunol* 2002; **3**: 549–557.
- 10 Mullen AC, Hutchins AS, High FA, Lee HW, Sykes KJ, Chodosh LA *et al.* Hlx is induced by and genetically interacts with T-bet to promote heritable T(H)1 gene induction. *Nat Immunol* 2002; **3**: 652–658.
- 11 Murakami K, Kojima T, Sakaki Y. Assessment of clusters of transcription factor binding sites in relationship to human promoter, CpG islands and gene expression. *BMC Genomics* 2004; **5**: 16.
- 12 Weinmann AS, Yan PS, Oberley MJ, Huang TH, Farnham PJ. Isolating human transcription factor targets by coupling chromatin immunoprecipitation and CpG island microarray analysis. *Genes Dev* 2002; **16**: 235–244.
- 13 Mao DY, Watson JD, Yan PS, Barsyte-Lovejoy D, Khosravi F, Wong WW *et al.* Analysis of Myc bound loci identified by CpG island arrays shows that Max is essential for Myc-dependent repression. *Curr Biol* 2003; **13**: 882–886.
- 14 Wells J, Yan PS, Cechvala M, Huang T, Farnham PJ. Identification of novel pRb binding sites using CpG microarrays suggests that E2F recruits pRb to specific genomic sites during S phase. *Oncogene* 2003; **22**: 1445–1460.
- 15 Jacquemin P, Lannoy VJ, Rousseau GG, Lemaigre FP. OC-2, a novel mammalian member of the ONECUT class of homeodomain transcription factors whose function in liver partially overlaps with that of hepatocyte nuclear factor-6. *J Biol Chem* 1999; **274**: 2665–2671.
- 16 Cho JY, Grigora V, Murphy TL, Murphy K. Identification of cooperative monomeric Brachyury sites conferring T-bet responsiveness to the proximal IFN-gamma promoter. *Int Immunol* 2003; **15**: 1149–1160.
- 17 Lee DU, Avni O, Chen L, Rao A. A distal enhancer in the interferon-gamma (IFN-gamma) locus revealed by genome sequence comparison. *J Biol Chem* 2004; **279**: 4802–4810.
- 18 Kispert A, Koschorz B, Herrmann BG. The T protein encoded by Brachyury is a tissue-specific transcription factor. *EMBO J* 1995; **14**: 4763–4772.
- 19 Lemaigre FP, Durvieux SM, Truong O, Lannoy VJ, Hsuan JJ, Rousseau GG. Hepatocyte nuclear factor 6, a transcription factor that contains a novel type of homeodomain and a single cut domain. *Proc Natl Acad Sci USA* 1996; **93**: 9460–9464.
- 20 Lannoy VJ, Burglin TR, Rousseau GG, Lemaigre FP. Isoforms of hepatocyte nuclear factor-6 differ in DNA-binding properties, contain a bifunctional homeodomain, and define the new ONECUT class of homeodomain proteins. *J Biol Chem* 1998; **273**: 13552–13562.
- 21 Vanhorenbeek V, Jacquemin P, Lemaigre FP, Rousseau GG. OC-3, a novel mammalian member of the ONECUT class of transcription factors. *Biochem Biophys Res Commun* 2002; **292**: 848–854.
- 22 Harada R, Berube G, Tamplin OJ, Denis-Larose C, Nepveu A. DNA-binding specificity of the cut repeats from the human cut-like protein. *Mol Cell Biol* 1995; **15**: 129–140.
- 23 Bettelli E, Sullivan B, Szabo SJ, Sobel RA, Glimcher LH, Kuchroo VK. Loss of T-bet, but not STAT1, prevents the development of experimental autoimmune encephalomyelitis. *J Exp Med* 2004; **200**: 79–87.
- 24 Lieberman LA, Banica M, Reiner SL, Hunter CA. STAT1 plays a critical role in the regulation of antimicrobial effector mechanisms, but not in the development of Th1-type responses during toxoplasmosis. *J Immunol* 2004; **172**: 457–463.
- 25 Clotman F, Jacquemin P, Plumb-Rudewicz N, Pierreux CE, Van der Smissen P, Dietz HC *et al.* Control of liver cell fate decision by a gradient of TGF beta signaling modulated by Onecut transcription factors. *Genes Dev* 2005; **19**: 1849–1854.
- 26 Vanhorenbeek V, Jenny M, Cornut JF, Gradwohl G, Lemaigre FP, Rousseau GG *et al.* Role of the Onecut transcription factors in pancreas morphogenesis and in pancreatic and enteric endocrine differentiation. *Dev Biol* 2007; **305**: 685–694.
- 27 Margagliotti S, Clotman F, Pierreux CE, Beaudry JB, Jacquemin P, Rousseau GG *et al.* The Onecut transcription factors HNF-6/OC-1 and OC-2 regulate early liver expansion by controlling hepatoblast migration. *Dev Biol* 2007; **311**: 579–589.
- 28 Remboutsika E, Yamamoto K, Harbers M, Schmutz M. The bromodomain mediates transcriptional intermediary factor 1alpha-nucleosome interactions. *J Biol Chem* 2002; **277**: 50318–50325.
- 29 Huang TH, Perry MR, Laux DE. Methylation profiling of CpG islands in human breast cancer cells. *Hum Mol Genet* 1999; **8**: 459–470.
- 30 Yamamoto K, Sonoda M, Inokuchi J, Shirasawa S, Sasazuki T. Polycomb group suppressor of zeste 12 links heterochromatin protein 1alpha and enhancer of zeste 2. *J Biol Chem* 2004; **279**: 401–406.

Supplementary Information accompanies the paper on Genes and Immunity website (<http://www.nature.com/gene>)

Mini-Review

Plants, endosymbionts and parasites

Absciscic acid and calcium signaling

Kisaburo Nagamune,¹ Liming Xiong,² Eduardo Chini³ and L. David Sibley^{4,*}

¹Graduate School of Life and Environmental Sciences; University of Tsukuba; Tsukuba, Ibaraki, Japan; ²Danforth Plant Science Center; St. Louis, Missouri USA; ³Department of Anesthesiology; Mayo Medical School; Mayo Clinic and Foundation; Rochester, Minnesota USA; ⁴Department of Molecular Microbiology; Washington University School of Medicine; St. Louis, Missouri USA

Key words: parasite, calcium, signaling, hormone, stress, chemotherapy

It was recently discovered that the protozoan parasite, *Toxoplasma gondii* produces and uses the plant hormone, abscisic acid (ABA), for communication. Following intracellular replication, ABA production influences the timing of parasite egress from the host cell. This density-dependent signal may serve to coordinate exit from the host cell in a synchronous manner by triggering calcium-dependent activation of motility. In the absence of ABA production, parasites undergo differentiation to the semi-dormant, tissue cyst. The pathway for ABA production in *T. gondii* may be derived from a relict endosymbiont, acquired by ingestion of a red algal cell. Although the parasite has lost the capacity for photosynthesis, the plant-like nature of this signaling pathway may be exploited to develop new drugs. In support of this idea, an inhibitor of ABA biosynthesis protected mice against lethal infection with *T. gondii*. Here, we compare the role of ABA in parasites to its activities in plants, where it is known to control development and stress responses.

Toxoplasma gondii: Ubiquitous Protozoan Parasite with a Plant-like Endosymbiont

T. gondii is found in a wide range of warm-blooded animals and infects about one third of world's human population.¹ While normally benign, infections in immunocompromised patients can result in severe disease.¹ This parasite has two main proliferative stages that contribute to infection in humans.² Tachyzoites replicate rapidly within a segregated vacuole, rupture out of the cell, and re-invade new cells. In contrast, bradyzoites encyst and grow slowly, mainly in long-lived cells of the central nervous system and musculature. The encysted form can survive for a long time and poses a risk due to reactivation following host immunosuppression. Once reactivated, parasites revert to the lytic phase characterized by rapid growth, egress and invasion of new host cells, thus causing considerable tissue damage.

T. gondii is a parasite in the phylum Apicomplexa, which includes a large number of obligate intracellular parasites. The causative agent of malaria, *Plasmodium* and a cause of serious gastrointestinal disease, *Cryptosporidium*, also belong to the Apicomplexa. Most apicomplexans contain a special organelle called the apicoplast, which was acquired by engulfment of an algal cell in a secondary endosymbiotic event.^{3,4} The original endosymbiont likely had four membranes; two of them derived from photosynthetic bacteria and two additional ones from the algal cell and progenitor host cell plasma membranes.⁵ This configuration persists in *T. gondii* (Fig. 1A), although some reports indicate only three membranes surround the malaria apicoplast.⁵ Debate about the origin of this endosymbiont was recently resolved by the discovery of a photosynthetic relative of the Apicomplexa and their sister taxa the dinoflagellates. Prototrophic branch of the tree is represented by the marine alga called *Chromera*, which contains an endosymbiont of red algal origin that retains key features seen in Apicomplexa and dinoflagellates.⁶ Since its engulfment, the endosymbiont has undergone a variety of rearrangements in different derived lineages to shuffle most of the genes to the nucleus. In the process, many pathways have been modified or entirely lost, for example the absence of photosynthesis among all apicomplexans.⁵

The presence of the apicoplast implies that *T. gondii* and related parasites have the potential to use plastid or algal-like pathways derived from the endosymbiont. Previous studies have highlighted the importance of various pathways for metabolism such as a fatty acid FAS II-like pathway, ferredoxin-ferredoxin-NADP⁺ reductase, heme biosynthesis and a DOXP isoprenoid biosynthesis pathway.⁵ These have been suggested as important targets for development of new therapeutics, which is especially attractive given their bacterial or plastid-like nature. However, acquisition of primitive signaling pathways had not been considered previously. Thus, the discovery of the ABA response pathway in *T. gondii* is notable as it suggests early branching alga may have contained metabolites that control signaling between cells, and these pathways may have been inherited in parasites.

Calcium, Motility and Signaling in *T. gondii*

T. gondii uses intracellular calcium as a signal to control protein secretion and to activate motility, two events that are crucial to the parasite's intracellular existence.⁷ Apicomplexan parasites undergo

*Correspondence to: L. David Sibley; Department of Molecular Microbiology; Washington University School of Medicine; 660 S. Euclid Avenue; St. Louis, Missouri 63110 USA; Email: sibley@wustl.edu

Submitted: 04/13/08; Accepted: 04/15/08

Previously published online as a *Communicative & Integrative Biology* Epublication: <http://www.landesbioscience.com/journals/cib/article/6106>

actin-based gliding on substrates and this form of motility powers cell invasion.⁸ Intracellular calcium stores are required for activating increases in cytosolic calcium, which induces protein secretion and facilitates cell invasion.⁹ Intracellular calcium has also been implicated in controlling egress of the parasite from infected cells.¹⁰ Pharmacological evidence suggests the presence of both ryanodine-responsive and IP_3 -receptor-like channels, although the genes encoding these activities have not been identified in apicomplexan parasites.¹¹ In searching the genome for calcium regulation pathways, it became clear that *T. gondii* lacks many animal-like pathways, yet has affinities with plants.¹² For example, *T. gondii* lacks conventional IP_3 channels but instead contains a gene that is similar to the two-pore calcium channel found in plants.^{12,13} Apicomplexans also contain a variety of calcium-dependent protein kinases, which are highly developed in plants but absent in animals.¹⁴

The second messenger cyclic ADP ribose (cADPR) evokes changes in cytosolic calcium in both plants¹⁵ and animals¹⁶ by stimulating calcium release from intracellular stores. *T. gondii* also uses the second messenger cADPR to stimulate release of internal calcium stores.¹⁷ The antagonist 8-Br-cADPR inhibits protein secretion and motility in the parasite,¹⁷ indicating this pathway is important for controlling cell invasion. Plants generate cADPR in response to the phytohormone ABA,¹⁸ leading to elevated cytosolic calcium that controls stomata closure in guard cells and induces gene expression.¹⁸ ABA has also been linked to generation of cADPR and induction of calcium signaling in other metazoans including hydra¹⁹ and sponges.²⁰ Recently, human granulocytes (myeloid cells) were found to produce ABA, which triggers calcium increases by a cADPR-dependent pathway.²¹ This prompted us to consider whether apicomplexan parasites might also contain such an ABA-dependent, cADPR pathway for controlling calcium release.

ABA Production and Signaling in *T. gondii*

Addition of exogenous ABA to extracellular *T. gondii* parasites strongly stimulated secretion of proteins from apical organelles called micronemes,²² a process necessary for parasite motility and cell invasion. Previous studies have shown that this pathway relies on intracellular calcium and is mediated by increases in cADPR.^{11,17} Addition of ABA also stimulated increases in cADPR and lead to a calcium-dependent induction of protein secretion.²² This activity was restricted to (\pm ABA) and was not stimulated by β -carotene, retinoic acid or (-) ABA.²²

Further evidence that *T. gondii* contains ABA was provided by ELISA using an antibody specific to ABA and by mass spectrometric (MS) analysis.²² Levels of ABA reached a peak just prior to natural egress, when parasite-containing vacuoles reach their maximum size. MS/MS analysis revealed the characteristic fragments for both *trans*-ABA and *cis*-ABA in extracts of *T. gondii* that were subjected to HPLC purification and gas chromatography and tandem MS/MS analysis.²² The levels of ABA in parasites ranged from 0.1–0.2 M during initial infection to ~4 M near the end of the infection cycle, when the density of parasites within the host cell is highest.




Figure 1. ABA production in *T. gondii*. (A) Cross-section through an intracellular *T. gondii* showing organelle structures. Insert illustrates four surrounding membranes from enlarged apicoplast. N, nucleus; A, apicoplast; AC, apical complex consisting of conoid and secretory organelles; PV, parasitophorous vacuole. (B) Model of ABA production by the parasite with in the PV. Accumulation of ABA (pink) leads to increases in intracellular calcium (blue) activating egress and triggering secretion of micronemes (MICs shown in green). Fluridone blocks ABA production, preventing egress and favoring differentiation of tissue cysts.

These levels are 10–40 times higher than that reported for human granulocytes,²¹ supporting the idea that ABA is produced by the parasite rather than concentrated from environmental sources. Remarkably, when exogenous ABA was added to late stage vacuoles, it induced premature egress, consistent with the idea that this signal mediates exit of the parasite from the host cell. Similar to plants, the production of ABA in *T. gondii* was inhibited by treatment with fluridone, which blocks ABA-biosynthesis at the step of phytoene desaturase²⁴ (Fig. 2). Disruption of ABA production by fluridone inhibited the egress of *T. gondii* from the host cell. The fluridone block was rescued by addition of exogenous ABA, supporting the idea that this signal controls nature egress. Collectively, these studies indicate that *T. gondii* produces ABA, elevating the second messenger cADPR, and increasing intracellular calcium to control motility and cell egress (Fig. 1B).

In plants, ABA also controls development by acting to increase dormancy in seeds and to prevent premature germination of embryos.²⁵ Transcript levels of ABA biosynthetic genes (see below) in plants are also upregulated during osmotic, cold and salt stress.²⁵ Thus, ABA has been recognized as a stress hormone, able to increase plant tolerance to stressful conditions. A role in the stress response does not appear to be directly analogous in parasites. Instead repression of ABA synthesis by fluridone induced the differentiation to bradyzoites *in vitro*.²² Dormancy in *T. gondii* is normally a stress-induced pathway that can be triggered by nutrient limitation.²⁶ Thus, differentiation to bradyzoites in the absence of ABA may simply reflect a default pathway for parasites remaining within a host cell that contains diminishing resources and which lacks a signal for egress. It is also possible that reduction in ABA synthesis may make parasites more susceptible to stressful conditions in the environment, thus triggering a differentiation process that would otherwise be blocked under high levels of ABA.

Fluridone was also effective in preventing lethal toxoplasmosis in the murine model, consistent with its ability to block ABA




Figure 2. Biosynthetic pathways for ABA production in higher plants and fungi. This is an abbreviated scheme showing only the major steps in the pathways. For further details consult.^{24,25,28} DOXP, deoxy-D-xylulose 5-phosphate; FPS, farnesyl diphosphate synthase; GGPS, geranylgeranyl diphosphate synthase; PSY, phytoene synthase; PDS, phytoene desaturase; ZEP, zeaxanthin epoxidase; NSY, neoxanthin synthase; NCED, 9-cis-epoxycarotenoid dioxygenase; ABA, abscisic acid.

production, delay egress and thus prevent spread of infection.²² Previous studies demonstrate the related compound norflorizone is also effective at inhibiting malaria *in vitro*,²⁷ although the biological role of ABA has not been studied in this system. Because fluridone is an effective herbicide that is relatively non-toxic to mammals, it may provide a good lead for the development of anti-parasitic compounds to combat infection.

Pathways for ABA Production in Plants, Fungi and Algae

The synthesis of ABA in higher plants occurs by an indirect pathway that is initiated by cleavage of carotenoids in the plastid and subsequent modification of the resulting xanthoxin products in the cytosol²⁴ (summarized in Fig. 2). The pathway is known from a variety of metabolic studies as well as mutants that are disrupted in specific steps of synthesis, largely from studies in *Arabidopsis*, maize and tomato.²⁴ In plastids, the β -carotene hydroxylated product zeaxanthin is converted to violaxanthin by a zeaxanthin epoxidase (ZEP), also known as *ABA1* in *Arabidopsis*. Mutants in this step are defective in production of ABA, demonstrating that ABA is derived from C_{40} carotenoids. The first committed step in the pathway to ABA is catalyzed by 9-cis-epoxycarotenoid dioxygenase (NCED), which cleaves the C_{40} substrate 9-cis-epoxycarotenoid to yield the C_{15} intermediate xanthoxin. NCED is related to dioxygenases in mammalian cells that generate vitamin A (retinol) from β -carotene obtained in the diet. NCED activity is impaired in the *viviparous 14* (*vp14*) mutant of maize, leading to premature seed germination.

Subsequent modification of xanthoxin by ABA2, a short-chain alcohol dehydrogenase/reductase, yields abscisic aldehyde. This product is then converted to ABA, by the action of ABA aldehyde oxidases (AAO), which require the molybdenum cofactor (MoCo) sulfuryase (ABA3)-activated MoCo factor for the catalytic activity. ABA is catabolized in plants to the inactive phaseic acid by the action of ABA 8'-hydroxylases. Fungi also produce ABA; however, they use a direct route from farnesyl pyrophosphate through a cyclized C_{15} intermediate known as α -ionylideneethanol²⁸ (Fig. 2).

The initial steps in ABA synthesis in higher plants take place in the plastids, attesting to their origin from algal endosymbionts. While it is not yet clear if the downstream enzymes are conserved in algae, there is compelling evidence for ABA biosynthesis in these organisms. ABA has been reported in a wide variety of algae including green (chlorophyta), red (rhodophyta), brown (heterokontophyta) and several cyanobacteria, with intracellular concentrations ranging from 1–40 M.²⁹ Radiolabeling studies indicate that ABA production by the green alga *Dunaliella salina* is enhanced by hypersalinity stress.³⁰ Cyanobacteria have also been reported to produce ABA under stress conditions.³¹ Furthermore, ABA influences stress responses and morphogenesis in the green alga *Haematococcus pluvialis*³² and induces differentiation and spore formation in the brown alga *Laminaria japonica*.³¹

While there is clear evidence for the production of ABA and physiological responses to this hormone in algae, the biosynthetic pathway is less well studied than in higher plants. Moreover, higher plants often contain gene families that encode ABA biosynthetic enzymes.²⁴ This gene redundancy may partly contribute to the 'leaky' phenotypes of certain ABA-deficient mutants (i.e., *vp14* in maize) that are only partially impaired in ABA production.²⁴ Additionally although most plants contain only single *ABA2* and *ABA3* genes, residual levels of ABA in mutants of these genes may reflect alternative shunt pathways that are not fully defined. Hence, the lack of close similarity between ABA biosynthetic genes in higher plants and algae may reflect gene divergence since the common origin of plastids.³³

Apicomplexans lack the mevalonate pathway but instead have a conserved DOXP-MEP pathway for synthesis of isoprenoids, that is thought to be derived from the plastid.³⁴ However, searching the *T. gondii* genome database (<http://toxodb.org>) does not provide strong support for a conserved carotenoid or ABA biosynthetic pathway. For example, *T. gondii* appears to lack clear orthologues for several steps in carotenoid biosynthesis including phytoene synthase (PSY) and NCED, at least based on searches using genes from higher plants. Weak orthologues for phytoene desaturase (PDS) and zeta carotene desaturase (ZDS) are found, and this is consistent with the inhibition of ABA production following treatment with fluridone.²² *T. gondii* contains multiple candidate genes for biosynthesis of ABA (i.e., *ABA1*, *ABA2* and *ABA3*). However, these genes have low homologies with higher plant genes, and do not allow unambiguous

assignments of the enzymes in this pathway. This may reflect a divergence of the pathways in parasite and/or similarity to algal like pathways, rather than those found in higher plants. Future analysis of this pathway in algae and parasites should provide further insight into the biosynthesis pathway and the role of this hormone outside of higher plants.

Outlook

Many apicomplexans retain an endosymbiont derived from ingestion of a red algal cell, giving rise to the possibility that plastid-like pathways have been retained in these parasites. Although apicomplexans have lost the means for photosynthesis, they appear to retain a primitive pathway for production of plant hormones such as ABA. Our studies reveal that *T. gondii* produces ABA, which likely originates from the DOXP isoprenoid pathway via carotenoid intermediates. However, apicomplexans appear to lack highly conserved enzymes found in this pathway in higher plants. Whether they contain a primitive or divergent pathway, or one that resembles the direct pathway found in fungi is presently unclear. In this regard, *Chromera*, an ancestrally related photosynthetic alveolate,⁶ may shed light on the pathways for carotenoid biosynthesis and downstream metabolites that existed in the ancestral algal cell that gave rise to the apicoplast.

In higher plants, the indirect pathway for ABA production begins in the plastid and is completed in the cytosol.²⁴ However, it is not known where ABA is synthesized, stored and accumulated in parasite-infected cells. Figure 1 shows ABA released into the vacuole; however, it is also possible it accumulates within the parasite or in the host cell. Since ABA is a weak acid (pKa 4.7) it tends to concentrate in compartments with neutral or basic pH and does not readily pass membranes. Accumulation in separate compartments may trigger different responses depending on the localization of specific mechanisms for detection.

Finally, it is not known how ABA is sensed in parasites. In this regard, the conservation of several recently described plant receptors for ABA in apicomplexan parasites is intriguing. The RNA-binding protein FCA is an ABA receptor that controls translational repression and flowering responses in plants.³⁵ The second of these is a G-protein-coupled receptor that functions as a plasma membrane receptor to control ABA responses in *Arabidopsis*.³⁶ Both of these receptors have putative orthologues with high BLAST hits in apicomplexan parasites (<http://ApiDB.org>). Downstream of such receptors, it is also not known if ABA induces transcriptional changes in parasites, similar to those described in plants, where this is a predominant mode of action.^{24,25} Thus, while our initial studies implicate ABA in calcium-triggered egress, it is likely this metabolite has many other influences on parasite biology that are yet to be discovered.

Acknowledgements

We are grateful to Boris Striepen, Sylvia Moreno, Geoff McFadden and Jan Zeevaert for helpful discussions. Work in the author's laboratories was supported by the NIH (L. David Sibley), AHA (Eduardo Chini), NSF (Liming Xiong) and MEXT (Kisaburo Nagamune).

References

- Peterson E. In: *Toxoplasma: Molecular and Cellular Biology*. Ajioka JW, Soldati D, eds. Norfolk, UK: Horizon Bioscience 2007; 37-58.
- Dubey JP, Lindsay DS, Speer CA. Structures of *Toxoplasma gondii* tachyzoites, bradyzoites, and sporozoites and biology and development of tissue cysts. *Clin Microbiol Rev* 1998; 11:267-99.
- Köhler S, Delwiche CF, Denny PW, Tilney LG, Webster P, Wilson RJ, Palmer JD, Roos DS. A plastid of probable green algal origin in Apicomplexan parasites. *Science* 1997; 275:1485-9.
- McFadden GI, Reith ME, Munholland J, Lang-Urnach N. Plastid in human parasites. *Nature* 1996; 381:482.
- Waller RF, McFadden GI. The apicoplast: a review of the derived plastid of apicomplexan parasites. *Curr Issues Mol Biol* 2005; 7:57-79.
- Moore RB, Obornik M, Janoušková J, Chudimský T, Vancová M, Green DH, Wright SW, Davies NW, Bolch CJ, Heimann K, Slapeta J, Hoegh-Guldberg O, Lagsdon JM, Carter DA. A photosynthetic alveolate closely related to apicomplexan parasites. *Nature* 2008; 451:559-63.
- Moreno SNJ, Docampo R. Calcium regulation in protozoan parasites. *Curr Opin Microbiol* 2003; 6:359-64.
- Sibley LD. Invasion strategies of intracellular parasites. *Science* 2004; 304:248-53.
- Lovett JL, Sibley LD. Intracellular calcium stores in *Toxoplasma gondii* govern invasion of host cells. *J Cell Sci* 2003; 116:3009-16.
- Black MW, Arrizabalaga G, Boothroyd JC. Ionophore-resistant mutants of *Toxoplasma gondii* reveal host cell permeabilization as an early event in egress. *Mol Cell Biol* 2000; 20:9399-408.
- Lovett JL, Marchesini N, Moreno SN, Sibley LD. *Toxoplasma gondii* microneme secretion involves intracellular Ca²⁺ release from IP₃/ryanodine sensitive stores. *J Biol Chem* 2002; 277:25870-6.
- Nagamune K, Sibley LD. Comparative genomic and phylogenetic analyses of calcium ATPases and calcium-regulated proteins in the Apicomplexa. *Mol Cell Evol* 2006; 23:1613-27.
- White PJ. Calcium channels in higher plants. *Biochem Biophys Acta* 2000; 1465:171-89.
- Harper JF, Harmon A. Plants, symbiosis and parasites: a calcium signalling connection. *Nat Rev Mol Cell Biol* 2005; 6:555-66.
- Allen GJ, Muir SR, Sanders D. Release of Ca²⁺ from individual plant vacuoles by both IP₃ and cyclic-ADP-ribose. *Science* 1995; 268:735-7.
- Guse AH. Cyclic ADP-ribose: a novel Ca²⁺-mobilising second messenger. *Cell Signal* 1999; 11:309-16.
- Chini EN, Nagamune K, Wetzl DM, Sibley LD. Evidence that the cADPR signaling pathway controls calcium-mediated secretion in *Toxoplasma gondii*. *Biochem J* 2005; 389:269-77.
- Wu Y, Kuzma J, Marchal E, Graeff R, Lee HC, Foster R, Chua NH. Abscisic acid signaling through cyclic ADP ribose in plants. *Science* 1997; 278:2126-30.
- Pucco S, Basile G, Bavestrello G, Bruzzone S, Corran C, Giovine M, Arillo A, Zocchi E. Abscisic acid signaling through cyclic ADP-ribose in hydroid regeneration. *J Biol Chem* 2004; 279:39783-8.
- Zocchi E, Carpaneto A, Corran C, Bavestrello G, Giovine M, Bruzzone S, Guida L, Franco L, Usai C. The temperature-signaling cascade in sponges involves a heat-gated cation channel, abscisic acid and cyclic ADP-ribose. *Proc Natl Acad Sci USA* 2001; 98:14859-64.
- Bruzzone S, Moreschi I, Usai C, Guida L, Damonte G, Sali A, Seari S, Millo E, De Flora A, Zocchi E. Abscisic acid is an endogenous cytokine in human granulocytes with cyclic ADP-ribose as second messenger. *Proc Natl Acad Sci USA* 2007; 104:5759-64.
- Nagamune K, Hicks LM, Lux B, Brossier F, Chini EN, Sibley LD. Abscisic acid controls calcium-dependent egress and development in *Toxoplasma gondii*. *Nature* 2008; 451:207-11.
- Finkelstein R, Reeves W, Aritizumi T, Steber C. Molecular aspects of seed dormancy. *Annu Rev Plant Biol* 2008; PMID: 18257111.
- Schwartz SH, Qin X, Zeevaert JAD. Elucidation of the indirect pathway of abscisic acid biosynthesis by mutants, genes, and enzymes. *Plant Physiol* 2003; 131:1591-601.
- Xiong L, Zhu JK. Regulation of abscisic acid biosynthesis. *Plant Physiol* 2003; 133:29-36.
- Weiss LM, Kim K. In: *Toxoplasma gondii* the model apicomplexan: perspectives and methods. Weiss LM, Kim K, eds. New York: Academic Press 2007; 341-66.
- Leef JL, Carlson PS. Carotenoid synthesis inhibiting herbicides and fatty acid synthesis oxime herbicides as anti-apicomplexa protozoan parasite agents. *Potomac Ltd. Prtn.* 1999; 1-8.
- Oritani T, Kiyota H. Biosynthesis and metabolism of abscisic acid and related compounds. *Nat Prod Rep* 2003; 20:414-25.
- Hirsch R, Hartung W, Gimmmler H. Abscisic acid content of algae under stress. *Botanica Acta* 1989; 102:326-34.
- Cowan AK, Rose PD. Abscisic acid metabolism in salt stressed cells of *Dunaliella salina*. *Plant Physiol* 1991; 97:798-803.
- Marsalek B, Zahradnickova H, Hronkova M. Extracellular abscisic acid produced by cyanobacteria under salt stress. *J Plant Physiol* 1992; 139:506-8.
- Kobayashi M, Hirai N, Kurimura Y, Ohigashi H, Tsuji Y. Abscisic acid-dependent algal morphogenesis in the unicellular green alga *Haematococcus pluvialis*. *Plant Growth Reg* 1997; 22:79-85.
- McFadden GI, van Dooren GG. Evolution: Red algal genome affirms a common origin of all plastids. *Curr Biol* 2004; 14:514-6.
- Ralph SA, van Dooren GG, Waller RF, Crawford MJ, Fraunholz MJ, Foth BJ, Tonkin CJ, Roos DS, McFadden GI. Metabolic maps and functions of the *Plasmodium falciparum* apicoplast. *Nature* 2004; 2:203-16.
- Razem FA, El-Keroumy A, Abrams SR, Hill RD. The RNA-binding protein FCA is an abscisic acid receptor. *Nature* 2006; 439:290-4.
- X Liu, Y Yue, B Li, Y Nie, W Li, WH Wu, L Ma A G protein-coupled receptor is a plasma membrane receptor for the plant hormone abscisic acid. *Science* 2007; 315:1712-6.

インド旅行で感染した腸管寄生原虫症の2例

東京都立墨東病院 感染症科

中村(内山) ふくみ・中村 造・古宮伸洋・大西健児

東京都健康安全センター 病原細菌研究科寄生虫

鈴木 淳

慶応義塾大学医学部 熱帯医学・寄生虫学教室

小林正規

Key Words : 腸トリコモナス, *Cryptosporidium hominis*, *Giardia intestinalis*, 遺伝子型, インド旅行

はじめに

下痢は発展途上国への旅行で経験する頻度が最も高い症状であり、1ヵ月の旅行で40～60%の人が経験すると報告されている。「海外、特に発展途上国への旅行に関連した感染性の下痢症」を旅行者下痢症と呼ぶ。今回、インド旅行で腸管原虫に複数感染していた旅行者下痢症の2例を経験し、原虫の遺伝子型分類を行ったので報告する。

症例1

患者は28歳の日本人女性で、3月1日～28日までインドを友人と旅行した。デリー、カルカッタ、

ダーズリン、ブッタガヤ、バラナシ、アグラ、ハリドワール、リシュケシュを訪れた。入国1週間後から下痢(軟便)が始まった。帰国後も軟便が続いていたが、4月6日に水様性下痢となったため、4月7日に当院外来を受診した。現地ではミネラルウォーターを飲んでいて、屋台で氷入りの飲み物やフルーツを食べたとのことである。蚊に多数回刺された。トラベラーズワクチンの接種なし。マラリア予防内服なし。身体所見では腹部の軽度の圧痛を認めるのみで全身状態は良好であった。検便で腸トリコモナス栄養体、ジアルジアの栄養体とシスト、クリプトスポリジウムのオーシストが検出された。メトロニダゾール750mg/日、7日間の治療を行っ

Two Cases of Traveler's Diarrhea Infected in India Due to Intestinal Protozoa Infection

Fukumi Nakamura-Uchiyama* Itaru Nakamura* Nobuhiro Komiya*
Kenji Ohnishi* Jun Suzuki** Seiki Kobayashi***

*Department of Infectious Diseases, Tokyo Metropolitan Bokutoh General Hospital

**Department of Microbiology, Tokyo Metropolitan Institute of Public Health

***Department of Tropical Medicine and Parasitology, School of Medicine, Keio University

論文請求先: 中村(内山) ふくみ 〒130-8575 東京都墨田区江東橋4-23-150 東京都立墨東病院 感染症科




図1 症例1・2から検出されたクリプトスポリジウムの形態と遺伝子型分類の結果




図2 症例1・2から検出されたジアルジアと腸トリコモナスの形態と遺伝子型分類の結果

た。原虫の遺伝子型解析の結果、*Cryptosporidium hominis* (Ia 2), *Giardia intestinalis* (Assemblage B) と分類された (図1, 2)。便培養で下痢起炎菌は検出されなかった。

メトロニダゾール治療終了直後の検便で腸トリコモナス、ジアルジアは陰性化していたが、クリプト

スポリジウムは残存していた。治療終了後1ヵ月目の検便でジアルジアのシストが少数検出され、メトロニダゾール1,500mg/日、7日間の治療を開始した。2回目の治療終了後にはジアルジアは陰性化していた。

症例 2

患者は30歳の女性で、症例1の友人でありともにインドを旅行した。同様に下痢が続いており、友人が寄生虫感染症と診断されたため検査目的で4月14日に当科外来を受診した。検便でジアルジアの栄養体とシスト、クリプトスポリジウムのオーシストが検出された。メトロニダゾール750mg/日、7日間の治療を行った。原虫の遺伝子型解析の結果、*C. canis*, *G. intestinalis* (Assemblage A) と同定された(図1, 2)。便培養で下痢起炎菌は検出されなかった。

メトロニダゾール治療終了直後の検便でジアルジアは陰性化しており、クリプトスポリジウムも陰性化していた。治療終了後1ヵ月目の検便でジアルジアのシストが多数検出された。メトロニダゾールで副作用が出現したためアルベンダゾール400mg/日、5日間の治療を開始した。2回目の治療終了後にはジアルジアは陰性化していた。

考 察

旅行者下痢症で病原体の判明したものうち、細菌性が85%と大部分を占め、原虫性は約10%、ウイルス性は約5%とされている。そのほとんどは軽症で、治療を行わなくとも1週間以内に90%の症例が改善する。しかし10日以上症状持続がある場合にはジアルジア症、クリプトスポリジウム症といった原虫感染の可能性が高まるため便の寄生虫検査を施行する必要がある¹⁾。本症例は典型的な原虫性の旅行者下痢症の2例である。同じ旅程で同様の生活パターンを過ごしていたにも関わらず、症例1では腸トリコモナス、ジアルジア、クリプトスポリジウムに感染し、症例2ではジアルジアとクリプトスポリジウムに感染していた。さらに遺伝子型の解析により、それぞれ*C. hominis* (Ia 2), *Giardia*

intestinalis (Assemblage B) と *C. canis*, *G. intestinalis* (Assemblage A) という別々の原虫に感染していたことは興味深い。

ジアルジアの治療にはチニダゾール、メトロニダゾール、フラゾリドン、キナクリン、アルベンダゾールが用いられるが²⁾、日本での第一選択はメトロニダゾールで、今回の2症例もまずは本薬で治療を行った。いずれも一度はジアルジアが陰性化したものの再燃を認めた。ジアルジアのメトロニダゾール治療抵抗性は以前から報告があり^{3) 4) 5)}、再燃率は高いものでは90%になるという²⁾。再燃後、症例1ではメトロニダゾール増量で、症例2ではアルベンダゾールで治療を行いジアルジアは陰性化している。今後も再々燃が起こらないかどうか、それぞれの薬剤に対する治療経過を観察していく予定である。

文 献

- 1) 古宮伸洋, 他 (2006) : 海外旅行帰国者の下痢. 成人病と生活習慣病, 36, 929-932.
- 2) Upcroft, P. *et al.* (2001) : Drug targets and mechanisms of resistance in the anaerobic protozoa. *Clin Microbiol Rev*, 14, 150-164.
- 3) Abboud, P. *et al.* (2001) : Successful treatment of metronidazole- and albendazole-resistant giardiasis with nitazoxanide in a patient with acquired immunodeficiency syndrome. *Clin Infect Dis*, 32, 1792-1794.
- 4) Boreham, P. F. L. *et al.* (1988) : Altered uptake of metronidazole *in vitro* by stocks of *Giardia intestinalis* with different drug sensitivities. *Trans R Soc Trop Med*, 82, 104-106.
- 5) Farthing, M. J. B. (1996) : Giardiasis. *Gastrointestrol Clin N Am*, 25, 493-515.

Original Article

Seroprevalence of *Entamoeba histolytica* Infection in Female Outpatients
at a Sexually Transmitted Disease Sentinel Clinic in Tokyo, Japan

Jun Suzuki*, Seiki Kobayashi¹, Ise Iku, Rie Murata, Yoshitoki Yanagawa and Tsutomu Takeuchi¹

Department of Microbiology, Tokyo Metropolitan Institute of Public Health, Tokyo 169-0073, and

*¹Department of Tropical Medicine and Parasitology, School of Medicine,
Keio University, Tokyo 160-8582, Japan*

Original Article

Seroprevalence of *Entamoeba histolytica* Infection in Female Outpatients at a Sexually Transmitted Disease Sentinel Clinic in Tokyo, Japan

Jun Suzuki*, Seiki Kobayashi¹, Ise Iku, Rie Murata, Yoshitoki Yanagawa and Tsutomu Takeuchi¹

Department of Microbiology, Tokyo Metropolitan Institute of Public Health, Tokyo 169-0073, and

¹Department of Tropical Medicine and Parasitology, School of Medicine, Keio University, Tokyo 160-8582, Japan

(Received December 13, 2007. Accepted February 19, 2008)

SUMMARY: From 2003 to 2006, we surveyed the seroprevalence of amoebic infection in female outpatients at a gynecologist's office, which was designated as a sexually transmitted disease sentinel clinic by the Tokyo Metropolitan Government, using an enzyme-linked immunosorbent assay (ELISA). The annual rate of anti-*Entamoeba histolytica* (HM-1:IMSSc16 strain; HM-1) antibody-positive cases as detected by ELISA increased during that period, and anti-*Chlamydia trachomatis* antibodies were detected in 60%, i.e., 24 of 40 anti-HM-1 antibody-positive individuals, suggesting sexual transmission of *E. histolytica*. We designed an ELISA with better sensitivity using the antigen extracted from the virulence-augmented *E. histolytica* strains (LHM-1 and LLA526 strains) by liver-passaging in hamsters. The average ratios of the S/N value (optical density [OD] of sample/OD of negative control) of ELISA with either the LHM-1 or LLA526 antigen and that of ELISA with the HM-1 antigen were significantly higher in intestinal amoebiasis cases with low S/N values than in amoebic liver abscess cases. In the present study of the seroprevalence of *E. histolytica* infection, the sera testing positive with low S/N values (<10) by ELISA with HM-1 antigen exhibited higher S/N values by ELISA using LHM-1 and LLA526 antigens. This modification of the antigen preparation for ELISA is expected to be effective in detecting anti-*E. histolytica* antibodies from such asymptomatic patients who have low antibody titers.

INTRODUCTION

In Japan, it was thought until the mid-1970s that amoebiasis was solely food borne and spread via food contaminated with cysts of *Entamoeba histolytica*. However, in the late 1970s, after amoebiasis was reported to have spread among men having sex with men (MSM) in large cities of the United States, it was recognized as a sexually transmitted disease (STD) (1,2). Within a few years, the suspected number of MSM having anti-*E. histolytica* antibodies along with anti-*Treponema pallidum* antibodies began to increase in densely populated cities in Japan (3,4).

In data provided by Japan's National Epidemiological Surveillance of Infectious Diseases, the number of notified cases with amoebiasis has been increasing annually; in 2006, 747 cases were reported, approximately 90% of which were male. However, with the spread of amoebiasis, the number of notified female cases has also increased at a slow but steady pace since 1999 (5,6).

In the present study, by detecting anti-*E. histolytica* (HM-1:IMSSc16 strain; HM-1) antibodies using an enzyme-linked immunosorbent assay (ELISA), we report the seroprevalence of amoebic infection in female outpatients who visited a gynecologist's office in Tokyo, Japan, from 2003 to 2006; this office was designated as an STD sentinel clinic by the Tokyo Metropolitan Government.

Moreover, in this study we attempted to design an ELISA with better sensitivity. This involved the use of the antigen

extracted from the virulence-augmented *E. histolytica* strains by liver-passaging in hamsters. The serum anti-*E. histolytica* antibody titers are low in a majority of asymptomatic amoebiasis cases. Practically, this serological method using LHM-1 and LLA526 antigens was tested on the anti-HM-1 antibody-positive sera in the present surveillance study.

MATERIALS AND METHODS

Study population: This study was conducted at a Tokyo gynecologist's office that was designated as an STD sentinel clinic by the Tokyo Metropolitan Government. We collected blood samples from 981 female outpatients between 2003 and 2006 (205 in 2003, 217 in 2004, 282 in 2005, and 277 in 2006) (Table 1). All individuals provided informed consent. Patient age was the only additional information. The anti-*E. histolytica* antibody-positive sera were examined for anti-*Chlamydia trachomatis* and anti-*T. pallidum* antibodies as indicators of STDs.

ELISA: *E. histolytica* antigens were prepared from axeni-

Table 1. Study samples in age categories from 2003 to 2006

Age	2003	2004	2005	2006	Total
<20	15	18	22	12	67
20-24	53	56	71	79	259
25-29	69	59	83	90	301
30-34	41	54	57	41	193
35-39	10	18	25	24	77
40-44	8	4	8	12	32
45-49	2	1	8	1	12
50<	1	2	5	7	15
unknown	6	5	3	11	25
Total	205	217	282	277	981

*Corresponding author: Mailing address: Division of Clinical Microbiology, Department of Microbiology, Tokyo Metropolitan Institute of Public Health, 3-24-1, Hyakunin-cho, Shinjuku-ku, Tokyo 169-0073, Japan. Tel: +81-3-3363-3231, Fax: +81-3-3368-4060, E-mail: Jun_Suzuki@member.metro.tokyo.jp

cally cultured *E. histolytica* (HM-1: ATCC no. 50527). The antigen was diluted with 0.05 M bicarbonate buffer to yield a concentration of 5 µg/mL. The diluted antigen (100 µL) was pipetted into each well of the microplate (Nunc-Immuno Module; Nunc Co., Roskilde, Denmark; Cat. no. 469078) and sensitized by incubation for 2 h at 37°C (7). After washing with a buffer (0.15 M phosphate buffer [PB] containing 0.05% Tween 20, pH 7.2; PB/T), 100 µL of the serum samples diluted 1:200 with a dilution buffer (PB/T containing 1% bovine serum albumin [BSA]) were pipetted into the microwells followed by incubation for 40 min at 37°C. The microplate was washed 3 times with PB/T after incubation, and 100 µL of 1:8,000 diluted peroxidase-conjugated anti-human IgG rabbit serum (ICN-Cappel Inc., Aurora, Ohio, USA; Cat. no. 55221) was added, followed by incubation for 40 min at 37°C. After washing with PB/T, the substrate solution comprising 0.03% 2,2'-azino-bis(3-ethylbenzo thiazoline-6-sulfonic acid) diammonium salt (ABTS; Sigma Co., St. Louis, Mo., USA; Cat. no. A1888), 0.01% H₂O₂ in 10 mL of 0.1 M Na₂HPO₄, and 10 mL of 0.1 M citric acid was added to each well. After 7 min, 50 µL of 1.25% NaF solution was added to arrest color development, and an ELISA reader (MTP-120) (Corona Electric Co., Ltd., Ibaragi, Japan) was used to measure the absorbance at 405 nm. The cut-off S/N optical density (OD) value, calculated using the average OD of negative sera from 5 healthy individuals, was set at 3.

Serological test for *C. trachomatis* and *T. pallidum* infections: Anti-*C. trachomatis* IgG and IgA antibodies were measured using a solid-phase enzyme immunoassay kit (Peptide-Chlamydia IgG and IgA; Ani LabSystems Ltd., Oy, Vantaa, Finland).

Nontreponemal anti-cardiolipin (CL) antibodies were detected using two kits (biologic false-positive tests for syphilis), i.e., slide test antigen (DS Pharma-Biomedicals, Osaka, Japan) and rapid plasma reagin (RPR) test (Sanko Junyaku, Tokyo, Japan). The sera that tested positive by these kits were retested using the *T. pallidum* passive hemagglutination (TPHA) kit (Fujirebio, Inc., Tokyo, Japan) for detection of anti-*T. pallidum* antibody. The OD values of the positive and negative control sera for quality control and the OD value of the positive control of the kits were measured during each run of all the serological tests.

ELISA with antigens from virulence-augmented amoebae: Based on the hypothesis that the amount of antigenic substances would also decrease simultaneously with the loss of virulence, we attempted to design an ELISA with better sensitivity in the following manner: (i) HM-1 and LA526 strains cultured axenically for 3 days in the TYI-S-33 medium were inoculated (dose, 1×10^6 amoebae/0.1 mL/head) into the left hepatic lobes of female Syrian golden hamsters (age, 3–4 weeks) (8). (ii) On the 6th day of inoculation, the hamsters were sacrificed and the livers dissected aseptically. The amoebic abscesses isolated from each of the livers were minced finely and crushed using scissors for medical use in the TYI-S-33 medium (9). (iii) After removing the tissue debris from the amoebic cell suspensions and washing twice in the TYI-S-33 medium by centrifugation ($175 \times g$ for 3 min), both *E. histolytica* strains were cultured axenically in the TYI-S-33 medium. They were named LHM-1 and LLA526. The long-term axenically cultured HM-1 was passaged 16 times through hamster liver due to the significant reduction in virulence, whereas the LA526 was passaged only once because it was newly isolated from the pus of a human amoebic liver abscess only 8 months earlier. (iv) LHM-1 and LLA526

were mass cultured within 2 weeks after their transfer into TYI-S-33 medium from the amoebic liver abscesses. The antigens were then harvested and washed twice in phosphate buffered saline (PBS) by centrifugation ($175 \times g$ for 3 min) and suspended in 5 mL of distilled water, followed by intermittent sonication (UH-150; SMT Co., Ltd., Tokyo, Japan) at 10 kHz for 5 min in an ice bath. (v) The sonicated suspensions were then centrifuged at $9,100 \times g$ for 30 min, and the protein concentrations of the aqueous soluble extracts were measured by Bradford's method (10). (vi) LHM-1 and LLA526 antigens were sensitized at a concentration of 0.5 µg/well according to the procedures described above.

Each serum sample was tested in triplicate for each of the three antigens—LHM-1, LLA526, and HM-1—and the average OD values were calculated. The sensitivity of ELISA for each of the three antigens was compared with the positive serum samples of 5 patients clinically diagnosed having amoebic liver abscesses and 5 mentally handicapped persons in a rehabilitation institution for the intellectually impaired in Japan, who were almost free from amoebiasis symptoms but positive for *E. histolytica* cysts on microscopy and for *E. histolytica* antigen when tested by using an *E. histolytica*-specific antigen detection kit (*E. histolytica* II kit; TechLab, Blacksburg, Va., USA). In each of the 10 human serum samples obtained as described above from the cases of amoebic liver abscess and asymptomatic cyst passers, the ratio was determined between the S/N values (OD value of serum sample [S]/average OD of negative sera from 5 healthy individuals [N]) of ELISA with the LHM-1 and HM-1 antigens and that between the S/N values of ELISA with the LLA526 and HM-1 antigens.

RESULTS

Seroprevalence of anti-*E. histolytica* antibodies in the female population: During the 4 years 2003 to 2006, in the 981 sera samples obtained from the study population, the seroprevalence of anti-*E. histolytica* (HM-1) antibodies increased every year. In 2005 and 2006, the annual positive rate was >5%; the average annual positive rate over the 4 years was 4.1% (40/981) (Table 2). In addition, 60%, i.e., 24/40 of these cases, were also positive for anti-*C. trachomatis* antibodies—an indicator of STDs. On the other hand, none of the cases were positive for anti-CL antibodies (a retest by the TPHA kit was not performed). The strong positive correlation between seropositivity for anti-*E. histolytica* and anti-*C. trachomatis* antibodies suggested sexual transmission of *E. histolytica* in the female population. The age range with the highest number of individuals positive for anti-*E. histolytica* antibodies was that of 25–29 years, with 11, and that of 30–

Table 2. Seroprevalence of anti-*Entamoeba histolytica* antibodies in the female outpatients from a gynecologist's office, Tokyo, Japan, by enzyme-linked immunosorbent assay from 2003 to 2006

Year	No. of samples	No. of positives	Positive rate %	No. of positives for anti-CT antibodies ¹⁾
2003	205	3	1.5	2
2004	217	8	3.7	6
2005	282	14	5.0	7
2006	277	15	5.4	9
Total	981	40	4.1	24

¹⁾ Number of positives for anti-*Chlamydia trachomatis* (CT) antibodies that were also positive for anti-*E. histolytica* antibodies.

Table 3. Age distribution of the female outpatients from a gynecologist's office with positive for anti-*Entamoeba histolytica* antibodies

Year	20-24	25-29	30-34	35-39	40-44	45-49	50<
2003	1	1 (1)		1 (1)			
2004		2 (1)	3 (3)	2 (2)			
2005	1	2 (1)	3 (1)		3	4 (3)	2 (2)
2006	3 (2)	6 (5)	3 (2)		2		1
Total	5 (2)	11 (8)	9 (6)	3 (3)	5	4 (3)	3 (2)

Number of positives for anti-*Chlamydia trachomatis* antibodies that were also positive for anti-*E. histolytica* antibodies are provided in parentheses.




Fig. 1. Ratio of the S/N values of ELISA with better sensitivity performed using the antigens from hamster liver-passaged LHM-1 and LLA526 strains to the S/N values of ELISA using the antigen from the HM-1:IMSSc16 (HM-1) strain in the clinical serum samples of amoebiasis (S/N values: OD of serum sample [S]/average OD of negative sera [N]). The mean of the ratios of triplicate ELISA are plotted. I: Samples ($n = 5$); anti-HM-1 antibody-positive sera from the clinical patients of amoebic liver abscess. The S/N values of ELISA with LHM-1 and LLA526 antigens did not increase significantly ($P > 0.05$ by t test). II: Samples ($n = 5$); anti-HM-1 antibody-positive sera from the mentally handicapped individuals admitted to a rehabilitation institution for the intellectually impaired in Japan. The S/N values of ELISA with LHM-1 and LLA526 antigens increased significantly ($P < 0.05$ by t test).




Fig. 2. Ratio of S/N values of ELISA performed using antigens from the LHM-1 and LLA526 strains to the S/N values of conventional ELISA performed using the antigen from the HM-1 strain in female outpatients from a gynecologist's office. The mean of the ratios of triplicate ELISA are plotted. I: Samples ($n = 16$); anti-HM-1 antibody-positive sera with high S/N values (≥ 10). The S/N values of ELISA with LHM-1 and LLA526 antigens did not increase significantly ($P > 0.05$ by t test). II: Samples ($n = 24$); anti-HM-1 antibody-positive sera with low S/N values (< 10). The S/N values by ELISA with LHM-1 and LLA526 antigens increased significantly ($P < 0.01$ by t test).

34 years, with 9. The number of individuals positive for anti-*C. trachomatis* antibodies correlated with the number of those positive for anti-*E. histolytica* antibodies (Table 3).

Comparison of S/N values of ELISA with three different antigens: Figure 1 shows the results of a pilot study in which ELISA with LHM-1 and LLA526 antigens was conducted using sera from 5 human cases each of amoebic liver abscess and asymptomatic cyst passers. The average ratio (1.324) between S/N values of ELISA with the LHM-1 and HM-1 antigens and that (1.254) between S/N values of ELISA with the LLA526 and HM-1 antigens increased significantly only in asymptomatic cases with low S/N values ($P < 0.05$ by t test) and not in amoebic liver abscess cases (1.048 and 1.006, respectively; $P > 0.05$ by t test).

The 40 anti-HM-1 antibody-positive sera as detected by ELISA were classified into two groups based on the magnitude of the S/N values (i.e., groups I and II with S/N values ≥ 10 and < 10 , respectively). The tendency of ELISA with LHM-1 and LLA526 antigens to yield significantly higher S/N values ($P < 0.01$ by t test) was also confirmed in seropositive cases from among the present study population with low S/N values (< 10) by ELISA using the HM-1 antigen (Figure 2).

DISCUSSION

In Japan, the MSM population is still thought to be a major high-risk group for STDs. However, our study provided evidence indicating that the seroprevalence of the *E. histolytica* infection in the female population of Tokyo is increasing annually.

In addition, the result that 60% of the female study population who were anti-*E. histolytica* antibody-positive were also positive for anti-*C. trachomatis* antibodies, an indicator of STD, along with the diversity of sexual behavior suggested that a major proportion of females positive for anti-*E. histolytica* antibodies may have been infected with *E. histolytica* by sexual transmission. We do not fully understand why none of the cases were positive for anti-CL antibodies in the female population, unlike the case in the MSM population (11,12). We are currently conducting further epidemiological studies on the route of *E. histolytica* infection in the female population.

The tendency of ELISA using the LHM-1 and LLA526 antigens to yield statistically higher S/N values ($P < 0.01$ by t test) was evident only in the positive cases with low S/N values (< 10) among the present female study population. The active antigenic substance that brought about this effect could not be identified in the present study. Despite the necessity of further evaluation, the improved ELISA is expected to be effective for detecting anti-*E. histolytica* antibodies from such asymptomatic patients who have low antibody titers. Moreover, the hamster liver-passaged *E. histolytica* may be applied as a sensitive antigen to other serodiagnostic methods, such as dot-ELISA (13) and immunofluorescence antibody tests (14).

Because of the public's indifference to STDs, the control of amoebiasis should start with efforts to raise public awareness of the risk of infection by sexual transmission. Also simpler and more sensitive mass examination methods should be developed, such as the newly designed ELISA using the antigen extracted from the virulence-augmented *E. histolytica* strains, which have a better sensitivity for the diagnosis of amoebiasis.

ACKNOWLEDGMENTS

A part of this work was supported by a Health Sciences Research Grant-in-Aid for Emerging and Re-emerging Infectious Diseases from the Ministry of Health, Labour and Welfare of Japan.

REFERENCES

1. Markell, E.K., Havens, R.F., Kuritsubo, R.A., et al. (1984): Intestinal protozoa in homosexual men of the San Francisco Bay area: prevalence and correlates of infection. *Am. J. Trop. Med. Hyg.*, 33, 239-245.
2. Schmerin, M.J., Gelston, A. and Jones, T.C. (1977): Amebiasis. An increasing problem among homosexuals in New York City. *JAMA*, 238, 1386-1387.
3. Takeuchi, T., Okuzawa, E., Nozaki, T., et al. (1989): High seropositivity of Japanese homosexual men for amebic infection. *J. Infect. Dis.*, 159, 808.
4. Nozaki, T., Motta, S.R., Takeuchi, T., et al. (1989): Pathogenic zymodemes of *Entamoeba histolytica* in Japanese male homosexual population. *Trans. R. Soc. Trop. Med. Hyg.*, 83, 525.
5. National Institute of Infectious Diseases and Tuberculosis and Infectious Diseases Control Division, Ministry of Health, Labour and Welfare (2007): Amebiasis in Japan, 2003-2006. *Infect. Agents Surveillance Rep.*, 28, 103'-104'.
6. Infectious Disease Surveillance Center, National Institute of Infectious Diseases: Annual Data Summary. Online at <<http://idsc.nih.gov/jp/idwr/ydata/index-e.html>>.
7. Kanwar, J.R. and Vinayak, V.K. (1991): A comparative efficacy of plate ELISA and dot ELISA to detect antiamebic antibodies in clinical patients. *Trop. Geogr. Med.*, 43, 261-265.
8. Diamond, L.S., Phillips, B.P. and Bartgis, I.L. (1974): A comparison of the virulence of nine strains of axenically cultivated *E. histolytica* in hamster liver. *Arch. Invest. Med.*, 5, 423-426.
9. Diamond, L.S., Harlow, D.R. and Cunnick, C.C. (1978): A new medium for the axenic cultivation of *Entamoeba histolytica* and other *Entamoeba*. *Trans. R. Soc. Trop. Med. Hyg.*, 72, 431-432.
10. Bradford, M.M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, 72, 248-254.
11. Takeuchi, T., Kobayashi, S., Asami, K., et al. (1987): Correlation of positive syphilis serology with invasive amebiasis in Japan. *Am. J. Trop. Med. Hyg.*, 36, 321-324.
12. Okusawa, E., Kobayashi, S., Miyahira, Y., et al. (1991): Study on clinical forms and demographic backgrounds of Japanese adult males with invasive amebiasis according to the history of sexual practice. *Jpn. Arch. Sex. Transm. Dis.*, 1, 153-156 (in Japanese).
13. Yamaura, H., Araki, K., Kiuchi, K., et al. (2003): Evaluation of dot-ELISA for serological diagnosis of amebiasis. *J. Infect. Chemother.*, 9, 25-29.
14. Ambroise-Thomas, P. and Truong, T.K. (1972): Fluorescent antibody test in amebiasis. *Am. J. Trop. Med. Hyg.*, 21, 907-912.

A SURVEY OF AMOEBIC INFECTIONS AND DIFFERENTIATION OF AN *ENTAMOEBIA HISTOLYTICA*-LIKE VARIANT (JSK2004) IN NONHUMAN PRIMATES BY A MULTIPLEX POLYMERASE CHAIN REACTION

Jun Suzuki, Seiki Kobayashi, Ph.D., Rie Murata, Hideo Tajima, D.V.M., Fumitaka Hashizaki, D.V.M., Yoshitoki Yanagawa, D.V.M., Ph.D., and Tsutomu Takeuchi, M.D., Ph.D.

Abstract: A pathogenic *Entamoeba histolytica*-like variant (JSK2004 strain) with genetic variations and a novel isoenzyme pattern isolated from a De Brazza's guenon in a Tokyo zoo in Japan has previously been documented. In this study, a multiplex polymerase chain reaction (PCR) assay that could distinguish the JSK2004-type *E. histolytica*-like variant (JSK04-Eh-V) from *E. histolytica* and *Entamoeba dispar* using three newly designed primer sets for amplifying each specific DNA fragment from their small-subunit ribosomal RNA genes was developed and established. Forty-seven primates (11 species) from the zoo were surveyed by multiplex PCR to assess the prevalence of JSK04-Eh-V infection, which was recognized in six individuals of four species, including an Abyssinian colobus monkey, a De Brazza's guenon (including the individual from whom JSK2004 was isolated), a white-faced saki, and a Geoffroy's spider monkey. In addition, the autopsied individuals of an Abyssinian colobus and Geoffroy's spider monkey that died of amoebic liver abscess were also evaluated. DNA samples were also analyzed for specific genotypes based on the nucleotide sequencing of two protein-coding (chitinase and serine-rich *E. histolytica* protein) genes and the protein-coding locus 1-2 that was used for fingerprinting of the *E. histolytica* strain. These studies indicated that the *E. histolytica*-like variant infection in this zoo was caused by the same type (i.e., JSK04-Eh-V). An axenic culture medium (yeast extract-iron-maltose-dihydroxyacetone-serum) was developed based on the yeast extract-iron-gluconic acid-dihydroxyacetone-serum medium, which is designed for axenic culture of *E. dispar*. This new medium could be used for axenically culturing *E. histolytica*, JSK04-Eh-V, and *E. dispar* in a single medium.

Key words: *Entamoeba histolytica*-like variant, multiplex PCR, nonhuman primates, zoo.

INTRODUCTION

Amoebiasis is a zoonotic protozoal infectious disease caused by *Entamoeba histolytica*. The estimated incidence of amoebiasis in humans is approximately 50 million per year, and it has caused nearly 70,000 human deaths.¹⁸ In 1997, *E. histolytica* was reclassified into two species: *E. histolytica* and *Entamoeba dispar* (nonpathogenic); earlier, this differentiation had been difficult to establish because of morphogenetic and phylogenetic similarities.¹⁹ This classification is based on the differences in the isoenzyme patterns (zymodemes), the detection of the *E. histolytica*-specific antigen, and *E. histolytica*- and *E. dispar*-specific DNA fragment

amplification by polymerase chain reaction (PCR).^{3,5}

In Japan, particularly during the last decade, *E. histolytica* infections have not been detected in nonhuman primates by PCR.^{12,14} However, three recent articles reported three different pathogenic *E. histolytica*-like variants showing subtle variations in the small-subunit ribosomal RNA (SSU rRNA) gene sequences isolated from cynomolgus monkey (*Macaca fascicularis*),¹⁵ rhesus monkey (*Macaca mulatta*),¹³ and De Brazza's guenon (*Cercopithecus neglectus*).¹² The JSK2004-type *E. histolytica* variant (JSK04-Eh-V)¹¹ has an SSU rRNA gene homology of 99.10% with *E. histolytica* and of 98.47% with *E. dispar*. Previously it was reported that the existing multiplex PCR² technique that targeted the specific region of the SSU rRNA gene sequence of *E. histolytica* did not yield the genomic DNA products of an axenic strain (JSK2004) of the *E. histolytica*-like variant from a De Brazza's guenon because of the variation in the nucleotide sequence of the gene.¹¹

In the present study, a new multiplex PCR assay that is capable of distinguishing the JSK04-Eh-V

From the Division of Clinical Microbiology, Department of Microbiology, Tokyo Metropolitan Institute of Public Health, 3-24-1, Hyakunin-cho, Shinjuku-ku, Tokyo 169-0073, Japan (Suzuki, Murata, Yanagawa); the Department of Tropical Medicine and Parasitology, School of Medicine, Keio University, 35 Shinanomachi, Shinjuku-ku, Tokyo 160-8582, Japan (Kobayashi, Takeuchi); and the Ueno Zoological Gardens, Ueno-park 9-83, Taito-ku, Tokyo 110-8711, Japan (Tajima, Hashizaki). Correspondence should be directed to Jun Suzuki (Jun.Suzuki@member.metro.tokyo.jp).

Table 1. Oligonucleotide primers used for polymerase chain reaction (PCR) assays in present study.

Primer name		Primer sequence (5' to 3')	Nucleotide position	Accession no.
EnthF	(forward)	ATG GCC AAT TCA TTC AAT GA	198–217	X65163
EnthR	(reverse)	TAC TTA CAT AAA GTC TTC AAA ATG T	648–672	X65163
EhvF	(forward)	ATT TTA TAC ATT TTG AAG ACT TTG CA	642–667	AB426549
EhvR	(reverse)	CTC TAA CCG AAA TTA GAT AAC TAC	1466–1489	AB426549
EhvR2	(reverse)	CAG ATT AAG AAA CAA TGC TTC TTC	1052–1075	AB426549
EntdF	(forward)	GTT AGT TAT CTA ATT TCG ATT AGA AC	1467–1492	AB282661
EntdR	(reverse)	ACA CCA CTT ACT ATC CCT ACC TA	1639–1661	AB282661
EchatF	(forward)	AGG ATT TGT TTT ATA ACA AGT TC	471–493	AF149912
EchatR	(reverse)	AAT AAC CTT TCT CCT TTT TCT ATC	660–685	AF149912
EhartF	(forward)	GTG AAG AGA AAG GAT ATC CAA AGT	221–244	AF149907
EhartR	(reverse)	ATA TCA TTT TCA ACT ACG AGC	623–643	AF149907
Chitinase	(forward)	GGA ACA CCA GGT AAA TGT ATA	466–487	U78319
Chitinase	(reverse)	TCT GTA TTG TGC CCA ATT	799–817	U78319
SREHP	(forward)	GCT AGT CCT GAA AAG CTT GAA GAA GCT G	258–286	M80910
SREHP	(reverse)	GGA CTT GAT GCA GCA TCA AGG T	784–806	M80910
R1 (locus1–2)	(forward)	CTG GTT AGT ATC TTC GCC TGT	1–21	AF276055
R2 (locus1–2)	(reverse)	CTT ACA CCC CCA TTA ACA AT	383–401	AF276055

from *E. histolytica* and *E. dispar* was designed, and the prevalence of JSK04-Eh-V infection in the primates of a zoo in Japan was surveyed using this assay. In addition, the identity of the JSK04-Eh-V strains in the zoo, determined by analyzing their polymorphic genotypes as a fingerprint for identifying the strain of *E. histolytica*, was investigated.

Moreover, the incidence of other amoebic infections in primates was investigated, and the first axenic culture medium that would support the growth of *E. histolytica*, JSK04-Eh-V, and *E. dispar* in a single medium was designed.

MATERIALS AND METHODS

Primates

In order to assess the prevalence of infection with JSK04-Eh-V, 47 captive individuals of 11 primate species from the Tokyo Zoo in Japan, where JSK2004 (JSK04-Eh-V) had been isolated from a De Brazza's guenon, were surveyed.¹¹ The primates, comprising 11 species, included three De Brazza's guenons (*Cercopithecus neglectus*), 11 Abyssinian colobus monkeys (*Colobus guereza*), two ring-tailed lemurs (*Lemur catta*), two mandrills (*Mandrillus sphinx*), one lesser slow loris (*Nycticebus pygmaeus*), two ruffed lemurs (*Varecia variegata*), one northern night monkey (*Aotus trivirgatus*), seven Geoffroy's spider monkeys (*Ateles geoffroyi*), 12 Japanese macaques (*Macaca fuscata*), five white-faced sakis (*Pithecia pithecia*), and one cotton-top tamarin (*Saguinus oedipus*). Each primate species was housed independently.

Microscopic examination and detection of the *E. histolytica*-specific antigen

Stool samples from each living individual were collected once daily for 3 days from 45 primates to obtain three samples per individual. Prior to performing the multiplex PCR, all stool specimens were examined microscopically after concentrating the *Entamoeba* cysts by the formalin-ether sedimentation technique.¹⁰ The specimens were also examined using an *E. histolytica*-specific antigen detection kit (*E. histolytica* II kit; TechLab, Blacksburg, Virginia 24060, USA). Tissue samples obtained from the primates that had liver abscesses were embedded in paraffin, stained with periodic acid-Schiff, and were examined for amoebae.

DNA preparation

Of the three stool specimens collected from each individual, the specimen that had the largest number of amoebic cysts was utilized for DNA preparation. The cysts were concentrated and partially purified using the modified formalin-ether sedimentation method, in which formalin was replaced with a phosphate-buffered solution (pH 7.4). Subsequently, the QIAamp® DNA stool mini-kit (Qiagen GmbH, Hilden 40724, Germany; catalog no. 51504) was used to isolate the genomic DNAs of amoebae. The genomic DNAs of the amoebae found in the two samples of abscesses obtained from the liver abscess of both the autopsied Abyssinian colobus monkey and Geoffroy's spider monkey that died of amoebic liver abscess; the two ref-



Figure 1. Polymerase chain reaction (PCR) products of the DNA samples from *Entamoeba histolytica*, JSK2004-type *E. histolytica*-like variant (JSK04-Eh-V), and *Entamoeba dispar* differentiated by multiplex PCR. Lane 1: mixture of the DNA templates of lanes 2, 3, and 4; Lane 2: DNA of HM-1:IMSScl6 (*E. histolytica*; length, 475 base pairs [bp]); Lane 3: DNA of JSK2004 clone 2 (JSK2004cl2; JSK04-Eh-V; length, 848 bp); Lane 4: DNA of SAW1734RclAR (*E. dispar*; length, 195 bp); Lane 5: Human *E. histolytica* DNA sample (pus from liver abscess); Lane 6: Human *E. histolytica* DNA sample (stool); Lane 7: JSK04-Eh-V DNA sample (pus from liver abscess) from a Geoffroy's spider monkey; Lane 8: JSK04-Eh-V DNA sample (stool) from a De Brazza's guenon; Lane 9: Human *E. dispar* DNA sample (stool); and Lane 10: *E. dispar* DNA sample (stool) from a De Brazza's guenon.

erence amoebic strains, HM-1:IMSS clone 6 (HM-1:IMSScl6 strain; *E. histolytica*) and SAW1734R clone AR (SAW1734RclAR strain; *E. dispar*), that were kindly supplied by Dr. Lois S. Diamond (National Institutes of Health); and the JSK2004 clone 2 (JSK2004cl2; JSK04-Eh-V) were isolated using the QIAamp DNA mini-kit (Qiagen GmbH; catalog no. 51304).

Primers for multiplex PCR

The primers for multiplex PCR were designed based on the two SSU rRNA gene sequences of HM-1:IMSScl6 (*E. histolytica*; GenBank accession no. X65163) and SAW1734RclAR (*E. dispar*; GenBank accession no. AB282661) and the previously reported sequence of JSK2004cl2 (JSK04-Eh-V; GenBank accession no. AB426549). The three primer sets that were designed—EnthF/EnthR for *E. histolytica*, EntdF/EntdR for *E. dispar*, and EHVf/EHVr for JSK04-Eh-V—are listed in Table 1.

Primer specificity was tested by conducting multiplex PCR on seven other intestinal parasitic protozoan and one nonprotozoan species: axenic trophozoites of *Entamoeba moshkovskii* (Laredo strain), *Entamoeba invadens* (IP-1 strain; ATCC no. 30994), and *Giardia intestinalis* (Portland-1 strain; ATCC no. 30888); cyst forms of *Escherichia coli* and *Cryptosporidium hominis*; culture form of *Blastocystis hominis* (nonprotozoan species) from human stool samples; and *E. coli*, *Entamoeba chat-*

toni, and *Entamoeba hartmanni* obtained from the stool samples of nonhuman primates.

Multiplex PCR

Amplification was performed in a reaction mixture (50 μ l) containing 100 ng of the DNA samples, 25 μ l of 2 \times Multiplex PCR Master Mix (Qiagen GmbH; catalog no. 206143), and 2 μ l of each primer at 10 mM. The touchdown method was used for thermal cycling. The cycling conditions were as follows: 15 min at 95°C followed by 40 cycles of denaturation at 94°C for 30 sec, annealing for 40 sec beginning at 61°C and ending at 56°C, and extension at 72°C for 1 min. The annealing temperature was lowered by 1°C after every four cycles until it reached 56°C, after which the same temperature was maintained until the end of the cycling process.

Semi-nested PCR for SSU rRNA

In cases in which a minimal PCR product from the DNA of JSK04-Eh-V was obtained, a semi-nested PCR using the primer set EHVf/EHVr2 was performed (Table 1). For this second PCR (semi-nested PCR), amplification was performed in a reaction mixture (50 μ l) containing 1 μ l of the first PCR product, 1.0 U of *exTaq* DNA polymerase (Takara Bio, Inc., Seta, Shiga 520-2134, Japan; catalog no. RR001A), 0.4 μ M of each primer, and 0.25 mM of deoxynucleoside triphosphate. The follow-

Table 2. The results of the surveillance for the prevalence of the JSK2004 type *Entamoeba histolytica*-like variant (JSK04-Eh-V) by a newly designed multiplex polymerase chain reaction (PCR) and microscopic examination in a zoo of Tokyo, Japan.

Primate species	Common name	No. of samples	Positive number by multiplex PCR (symptoms)				Antigen detector*	Microscopic examination†
			<i>Entamoeba histolytica</i>	JSK04-Eh-V	<i>Entamoeba dispar</i>			
Old World monkeys								
<i>Cercopithecus neglectus</i>	De Brazza's guenon	3	0	1 (Asymptomatic)	2	1	<i>E. coli</i> (3), <i>E. nana</i> (2)	
<i>Colobus guereza</i>	Abyssinian colobus	11	0	2 (ALA,‡ asymptomatic)	2	1	<i>E. coli</i> (8), <i>E. nana</i> (3), <i>G. intestinalis</i> (4)	
<i>Macaca fasciata</i>	Japanese macaque	12	0	0	2	0	<i>E. chattoni</i> (7), <i>E. coli</i> (2), <i>E. harmanni</i> (5)	
<i>Mandrillus sphinx</i>	Mandrill	2	0	0	0	0	<i>E. chattoni</i> (1), <i>E. nana</i> (1)	
<i>Lemur catta</i>	Ring-tailed lemur	2	0	0	0	0	<i>G. intestinalis</i> (2)	
<i>Varecia variegata</i>	Ruffed lemur	2	0	0	0	0	—	
<i>Nycticebus pygmaeus</i>	Lesser slow loris	1	0	0	0	0	—	
New World monkeys								
<i>Aotus trivirgatus</i>	Northern night monkey	1	0	0	0	0	<i>C. mesnili</i> (1)	
<i>Ateles geoffroyi</i>	Geoffroy's spider monkey	7	0	1 (Colitis)	1	0	—	
<i>Pithecia pithecia</i>	White-faced saki	5	0	2 (ALA,‡ asymptomatic)	1	2	<i>G. intestinalis</i> (1), <i>E. coli</i> (1), <i>E. nana</i> (1)	
<i>Saguinus oedipus</i>	Cotton-top tamarin	1	0	0	0	0	—	
Total		47	0	6	8	4		

* *E. histolytica*-specific antigen detection kit [*E. histolytica* II kit (TechLab)] was used for examination of stool samples.

† *E. coli*: *Entamoeba coli*; *E. nana*: *Endolimax nana*; *G. intestinalis*: *Giardia intestinalis*; *E. chattoni*: *Entamoeba chattoni*; *C. mesnili*: *Chilomastix mesnili*; *E. chattoni* and *E. harmanni* were identified by PCR. The numbers within the parentheses represent the number of positive cases.

‡ ALA, amebic liver abscess (death).




Figure 2. Differentiated polymerase chain reaction (PCR) products of the DNA samples of *Entamoeba chattoni* and *Entamoeba hartmanni* using PCRs. Lane 1: The product (length, 215 base pairs [bp]) from an *E. chattoni* DNA sample (stool) from a mandrill; Lane 2: The product (length, 215 bp) from an *E. chattoni* DNA sample (stool) from a Japanese macaque; Lane 3: The product (length, 423 bp) from an *E. hartmanni* DNA sample (stool) from a Japanese macaque.

ing cycling parameters were utilized: *Taq* activation at 94°C for 3 min; 35 cycles of denaturation at 94°C for 40 sec, annealing at 58°C for 40 sec, and extension at 72°C for 1 min; and extension at 72°C for 5 min.

PCR for *E. chattoni* and *E. hartmanni*

Entamoeba chattoni and *E. hartmanni* were identified by PCR assays using two primer sets (i.e., EchatF/EchatR¹⁶ and EhartF/EhartR, respectively) (Table 1). For *E. hartmanni*, a newly designed primer set based on its SSU rRNA sequence (GenBank accession no. AF149907) was used. These amplifications were performed in a reaction mixture (50 µl) containing 100 ng of the DNA sample, 1.0 U of *LATaq*DNA polymerase, 0.4 µM of each primer, and 0.25 mM of deoxynucleoside triphosphate. The following cycling parameters were utilized: *Taq* activation at 94°C for 3 min; 35 cycles of denaturation at 94°C for 40 sec, annealing at 55°C for 40 sec, and extension at 72°C for 1 min; and extension at 72°C for 5 min.

Polymorphic gene analysis

The genotyping of JSK04-Eh-V was reexamined to determine whether it would have the same genotype as JSK2004cl2. This was performed based on the nucleotide sequences of two protein-coding (chitinase and SREHP) genes and the protein-non-coding locus 1-2^{24,20} that was used as a fingerprint for identifying the *E. histolytica* strain. The primers used are shown in Table 1. These amplifications were performed in a reaction mixture (50 µl) containing 100 ng of the DNA sample, 1.0 U of *LATaq*DNA polymerase (Takara Bio, Inc.; catalog no. RR02AG), 0.4 µM of each primer, and 0.25 mM of deoxynucleoside triphosphate. The following cycling parameters were utilized: *Taq* activation at 94°C for 3 min; 35 cycles of denaturation at 94°C for 40 sec, annealing at 50°C (chitinase, SREHP, and locus 1-2) or 56°C (SSU rRNA) for 40 sec, and extension at 72°C for 1 min; and extension at 72°C for 5 min.

Sequence analysis

Sequence analysis was performed. The multiplex PCR products of SSU rRNA; the PCR products of chitinase, SREHP, and the locus 1-2 genes from the JSK04-Eh-V isolates; and the PCR products of SSU rRNA from *E. chattoni* and *E. hartmanni* were sequenced using the ABI Prism BigDye Terminator v3.1 Cycle Sequencing Ready Reaction Kit (Applied Biosystems, Foster City, California 94404, USA; catalog no. 4337455) and an ABI PRISM 3100 Genetic Analyzer.

Yeast extract-iron-maltose-dihydroxyacetone-serum (YIMDHA-S) medium

An axenic culture medium, namely, YIMDHA-S, based on the yeast extract-iron-gluconic acid-dihydroxyacetone-serum medium (YIGADHA-S)⁶ designed for the axenic culture of *E. dispar*, was designed for the isolation and culture of *E. histolytica*, the *E. histolytica*-like variant, and *E. dispar* in a single medium. YIGADHA-S differs from the existing YIMDHA-S medium in that gluconic acid, 0.5% in YIGADHA-S, is replaced with an equal concentration of maltose in YIMDHA-S. Another significant issue related to this culture system is that the growth of amoebae in YIMDHA-S is largely affected by the quality of the yeast extract. Accordingly, the effectiveness of several commercially available yeast extracts purchased from different manufacturers as an ingredient of YIMDHA-S was evaluated. Except for the standard stock obtained from BBL (Becton Dickinson Co., Cockeysville, Maryland 21030, USA; catalog no. 4311929; lot no. 100019DHJT), among all the yeast extracts test-

JSK2004c12	GGA	ACA	CCA	GGT	AAA	TGT	ATA	GGA	GAA	ACT	GTT	TGT	AAA	TGT	GGC	AGA	ACA	CAA	TAT	AAC
Samples*
EHMfas1
JSK2004c12	CCT	TGT	GTG	TGG	AAT	TTC	CTT	GAC	CTT	CCT	GAT	TGT	GAA	AAA	AAG	CCA	GGT	GAT	TTC	TTT
Samples
EHMfas1
JSK2004c12	GAG	AAG	TCA	CCA	GAT	TCT	TCT	GAA	TCT	AAG	CAT	GAA	TGT	TCT	GAA	ATT	AAA	CCA	GAT	TCT
Samples
EHMfas1
JSK2004c12	TCT	GAA	TCT	AAA	CAT	GAA	TCT	TCT	GAA	GTT	AAA	CCA	GAC	TGT	TCT	GAA	TCT	AAA	CAT	GAA
Samples
EHMfas1
JSK2004c12	TCT	TCT	GAA	GTT	AAA	CCA	GAT	TCT	TCT	GAA	TCT	AAG	CAT	GAA	TCT	TCT	GAA	GTT	AAA	CCA
Samples
EHMfas1
JSK2004c12	GAC	TCT	TCT	GAA	TCT	AAA	CAT	GAA	TCT	TCT	GAA	ATT	AAA	CCA	GAC	TCT	TCT	GAA	TCT	AAA
Samples
EHMfas1
JSK2004c12	CAT	GAA	TCT	TCT	GAG	CCA	GAA	GTT	AGT	GTC	CCA	AAG	AAA	ACA	GTT	GCT	TAT	TAT	ACT	AAT
Samples
EHMfas1
JSK2004c12	TGG	GCA	CAA	TAC	AGA	AG	(GenBank:AB426705)													
Samples	(GenBank:AB282755)													
EHMfas1	(GenBank:AB282755)													

Figure 3. Comparison of the chitinase sequences of the *Entamoeba histolytica*-like variant strain JSK2004c12 and *E. histolytica*-like variants detected from *Ateles geoffroyi* and *Pithecia pithecia* with the sequences of the reference *E. histolytica*-like variant strain isolated from *Macaca fascicularis*. JSK2004c12: The sequence of the chitinase gene of DNA from JSK2004c12 obtained from a De Brazza's guenon (GenBank accession no. AB426705). *Samples: The chitinase gene sequences of DNA samples from a Geoffroy's spider monkey (one pus sample from a liver abscess) and white-faced saki (two stool samples); EHMfas1: the chitinase gene sequences (GenBank accession no. AB282755) of the DNA of an isolate of another type of *E. histolytica*-like variant isolated from a cynomolgus monkey.¹⁵

ed, only that obtained from Merck (Merck KGaA, Darmstadt 64271, Germany; catalog no. 1.03753; lot no. VM510453 539) was effective for constant subculture of the axenic strains of *E. histolytica* (HM-1:IMSScl6), *E. dispar* (AS16 IR isolated from human samples and CYNO 09:TPC isolated from the cynomolgus monkey [*Macaca fascicularis*]),⁶ and JSK04-Eh-V (JSK2004) from the De Brazza's guenons that were subjected to a trial of axenic cultivation.

RESULTS

Specificity of multiplex PCR

Each PCR product from the genomic DNA of each of the axenic strains of *E. histolytica* (HM-1:IMSScl6), JSK04-Eh-V (JSK2004c12), and *E. dispar* (SAW1734clAR) analyzed by multiplex PCR was obtained independently; the lengths of the fragments were 475 base pairs (bp), 848 bp, and 195 bp, respectively (Fig. 1), which was confirmed by individual nucleotide sequencing. The findings of multiplex PCR were reproducible during the practical trials using amoebic DNAs isolated from the stool samples and liver abscesses of humans and nonhuman primates infected with *E. histolytica*,

JSK04-Eh-V, and *E. dispar* (Fig. 1). The specificity of multiplex PCR was examined by analyzing the templates of the DNA extracted from seven other intestinal parasitic protozoan and one nonprotozoan species, as previously described. None of the PCR products was observed on multiplex PCR examination of these parasites.

Sensitivity of multiplex PCR and semi-nested PCR

The sensitivity of multiplex PCR was found to be at least 200 cysts/100 mg of the stool sample for JSK04-Eh-V and 100 cysts/100 mg of the stool samples for *E. histolytica* and *E. dispar*. In an attempt to assess the sensitivity of the technique for mixed infections, 400 cysts or trophozoites of JSK04-Eh-V could be detected and differentiated in the presence of 100 cysts/100 mg of the stool samples for *E. histolytica* and *E. dispar*.

Semi-nested PCR using the primer set EHVf/EHVr2 detected 50 cysts/100 mg stool sample for JSK04-Eh-V. In one case of the white-faced saki, in which a minimal PCR product from the DNA of JSK04-Eh-V was obtained, semi-nested PCR yielded the 434-bp-long PCR product.

JSK2004c12	AAG	AAA	AAG	AAA	AAA	GTA	GCT	CAG	CAA	AAC	CAG	AAT	CAA	GTT	CAA	ACA	AAG	ATA	ATG	AAG				
EHMfas1				
P19				
JSK2004c12	ATG	AGG	AAG	ATG	AAG	ATG	AAG	ATG	ATG	AAG	AAG	ATG	AAG	ATG	AGA	ATG	AAA	AAG	CAA	GTT				
EHMfas1	...	---	---	-G				
P19	..	A	---	---	A	T	..	A	...	G				
JSK2004c12	CAA	GTG	ATA	AAT	CAG	AAG	CAA	GTT	CAA	GTG	ATA	AAT	CAG	A	---	---	---	---	AT	CAA	GCT			
EHMfas1	C	TA	ACA	AAC	CAG	A	G	...
P19	C
JSK2004c12	CAA	ATG	ATA	AAT	CAG	AAT	CAA	GCT	CAA	ATG	ATA	AAC	CAG	AAT	CAA	GCT	CAA	ATG	ATA	AAC				
EHMfas1	...	G	...	C	...	G	...	G	...	G	...	A	...	G	...	T	...	G	...	G				
P19	...	G	...	C	...	G	...	T	...	G	...	G	...	G	...	T	...	G	...	G				
JSK2004c12	CAG	AAG	CAA	GTT	CAA	GTG	ATA	AAT	CAG	AAG	CAA	GTT	CAA	GTG	ATA	AAC	CAG	ATA	ACA	AAC				
EHMfas1	...	---	---	---	---	-A	C	...	C				
P19	...	---	---	---	---	-A	C	...	C				
JSK2004c12	CAG	AAG	CAA	GTT	CAA	GTG	ATA	AAC	CAG	ATA	ACA	AAC	CAG	AAG	CAA	GTT	CAA	GTG	ATA	AAC				
EHMfas1				
P19				
JSK2004c12	CAG	ATA	ACA	AAC	CAG	AAG	CAA	GTT	CAA	GTG	ATA	AGC	CAG	ATA	ACA	AAC	CAG	AAG	CAA	GCT				
EHMfas1				
P19				
JSK2004c12	CAA	CTA	ATA	AAC	CAG	AAG	CAA	GTT	CAA	CTA	ATA	AAC	CAG	AAG	CAA	GTT	CAA	CTA	GTA	ATT				
EHMfas1				
P19				
JSK2004c12	CAA	-TG	ATA	AAT	CAG	AAA	GTA	GTT	CAG	ATA	ATG	ATA	ATA	ATA	(GenBank:AB426706)									
EHMfas1	...	A	(GenBank:AB197935)									
P19	...	A	(GenBank:AB282662)									

Figure 4. Comparison of the sequences in the SREHP of the *Entamoeba histolytica*-like variant JSK2004c12 strain with the reference *E. histolytica*-like variant strains isolated from *Macaca fuscicularis* and *Macaca mulatta*. JSK2004c12: The sequence of the serine-rich *E. histolytica* protein (SREHP) gene in the DNA of JSK2004c12 from a De Brazza's guenon (GenBank accession no. AB426706); EHMfas1: the sequences of the SREHP gene (GenBank accession no. AB197935) in the DNA of an isolate of another type of *E. histolytica*-like variant isolated from a cynomolgus monkey;¹⁵ P19: the sequence of the SREHP gene (GenBank accession no. AB282662) in the DNA of another type of *E. histolytica*-like variant (an established strain) isolated from a rhesus monkey.¹³




Figure 5. The growth kinetics of the axenically cultured HM-1:IMSScl6, JSK2004c12, AS 16 IR, and CYNO 09:TPC strains in the yeast extract-iron-maltose-dihydroxyacetone-serum (YIMDHA-S) medium. The mean numbers of amoebae in duplicate cultures have been plotted.

Prevalence of amoebic infections in the zoo

Since a pathogenic JSK04-Eh-V strain was isolated from De Brazza's guenons in the Tokyo zoo, the prevalence of JSK04-Eh-V infection was surveyed using multiplex PCR and was detected in six individuals of four primate species: Abyssinian colobus monkey; De Brazza's guenon, including the individual from which JSK2004 was isolated; white-faced saki; and Geoffroy's spider monkey. In addition, the autopsied samples of the Abyssinian colobus monkey and Geoffroy's spider monkey that died of amoebic liver abscess were positive for JSK04-Eh-V. Among the primates of the zoo, the infection rates with *E. histolytica*, *E. dispar*, and JSK04-Eh-V were 0% (0/47), 17% (8/47), and 13% (6/47), respectively (Table 2). Mixed infection with *E. dispar* and JSK04-Eh-V was not detected by multiplex PCR. The prevalence of the other species of amoebae examined microscopically (i.e., *E. coli* and *Endolimax nana*), were 30% (14/47) and 15%