

IDUs, including one female, declined to be interviewed (response rate of 98.5%). Compared to IDUs from the non-intervention zone, IDUs from the intervention zone were older [mean age, 30.81 versus 33.06 years, respectively; $t(417) = 2.61, p < 0.01$], were less likely to be of Fars ethnicity [54.5% versus 39.3%, respectively; $\chi^2(1, N = 419) = 9.63, p < 0.01$] and were more likely to be homeless [17.8% versus 59.2%; $\chi^2(1, N = 419) = 6.68, p < 0.01$]. However, IDUs from both neighborhoods were similar in terms of educational level, religious beliefs, and marital status (Table 1).

Drug Injection Characteristics

More than 99% of IDUs in both groups used heroin as their primary drug in the previous month. IDUs from the intervention zone had larger drug injecting networks ($M = 2.58; SD = 2.26$) than did IDUs recruited from the non-intervention zone [$M = 2.05, SD = 0.89, t(414) = 3.09, p < 0.01$]. In addition, IDUs from the intervention zone reported more frequent drug injection in the previous month ($M = 110.26, SD = 48.43$) compared to the control [$M = 92.61, SD = 47.19, t(415) = 3.76, p < 0.01$]. However, in the bivariate analyses, IDUs from both neighborhoods were similar in terms of the duration of lifetime injecting practices, the proportion of respondents who had ever shared a needle/syringe for drug injection, those who had been treated by a physician for drug addiction, and those who had shared a drug injection in prison (Table 1).

Although more than 67% of IDUs recruited from the intervention zone had received at least one needle or syringe from the NSP in the previous month, none of the participants from the non-intervention zone had received sterile needle/syringes from any NSP. The majority of the IDUs from the intervention zone reported that they obtained sterile needles/syringes primarily from reliable sources such as pharmacies (49%) and the NSP (47%), and only 3% turned to less reliable sources such as friends or drug dealers. In contrast, up to 30% of IDUs from the non-intervention zone reported that they obtained needles/syringes primarily from friends or drug dealers in the previous month. Compared to the IDUs from the non-intervention zone, a significantly higher proportion of respondents from the intervention zone reported that they had easy (51% versus 90%; OR, 8.90; 95% CI, 5.21–15.21) and regular (57% versus 96%; OR, 18.26; 95% CI, 8.56–38.94) access to needle/syringes during the previous month. Most importantly, a smaller proportion of IDUs from the intervention zone reported sharing a drug injection in the previous month (21%) than did those recruited from the non-intervention zone (40%; OR, 0.40; 95% CI, 0.26–0.62). This association between shared needle/syringe

use and frequenting a neighborhood with an active needle/syringe program remained significant in the multivariate analysis, which incorporated all of the variables mentioned in the data analyses (adjusted OR, 0.24; 95% CI, 0.13–0.45; Table 1).

Discussion

Needle and syringe programs constitute one of the main components of the government's comprehensive HIV prevention package, and are becoming increasingly accessible to community-based IDUs in Iran. However, there was little evidence concerning the effectiveness of these interventions in reducing injection-related risk behaviors among IDUs. Through this study, we investigated the potential impact of one NSP on drug-related risk behaviors in community-based IDUs in Tehran by comparing needle and syringe sharing practices among IDUs in two neighborhoods, one with and one without an NSP.

We found that a significantly smaller proportion of IDUs from the neighborhood with an NSP reported shared needle and syringe use in the previous month as compared to IDUs from a neighborhood without an intervention program. Our results showed that access to an NSP provided IDUs with better sources for sterile needles/syringes, free of charge, and with easier and more regular access to needles/syringes, which may have been the basis for the finding that IDUs from the intervention zone shared needles/syringes less frequently than did those in a non-intervention zone. The importance of these results only increases if we consider that IDUs from the intervention zone were living in a more vulnerable context than IDUs from the non-intervention zone: as mentioned earlier, 59.2% of the participants were homeless at the time of interview, compared to only 17.8% in the non-intervention zone. Subsequently, IDUs from the area with an NSP had larger drug injecting networks, and more frequent drug injection in the previous month, as compared to the non-intervention group. As may be conjectured, the general vulnerable context of the intervention zone is not a result of the NSP there, but is a known characteristic of that neighborhood, and is one of the primary reasons why health policy managers in Tehran decided to start community-based HIV prevention interventions for drug users there (Vazirian et al. 2005). Although IDUs from this neighborhood with an active NSP showed background characteristics that should have increased the risk of shared drug injection (Tyndall et al. 2003), shared drug injection was reportedly less in the previous month, as compared to the non-intervention group.

The effectiveness of NSPs has been researched extensively in many developed countries (Hartgers et al. 1989;

Bluthenthal et al. 2000; Cox et al. 2000; Gibson et al. 2002; Power and Nozhkina 2002; Wodak and Cooney 2005), and some information exists regarding Asian countries (Gray 1998; Quan et al. 1998; Chen and Liao 2005). Our findings indicated that the NSP in Tehran may have allowed the IDUs from a neighborhood with an active program to better avoid shared drug injection, as compared to participants from a neighborhood lacking this intervention.

In our previous biobehavioral study conducted in the same neighborhood with an active NSP in 2004, we found that over 23% of IDUs were infected with HIV and that infection was associated with a history of shared drug injection while in prison and with multiple incarcerations (Zamani et al. 2006). The risk of HIV transmission in association with shared drug injection in Iranian prisons has also been documented in other studies (Razzaghi et al. 2000; Razzaghi and Rahimi 2005; Zamani et al. 2005). Although the Iran Prison Organization has increased the scope of HIV prevention programs for drug-using inmates, concern remains over the possibility of HIV transmission from ex-prisoner IDUs to their community-based injecting and sexual networks in Tehran. This study clearly demonstrated that up to one-third, or more, of community-based IDUs recruited from two neighborhoods had a history of shared drug injection inside prison and it is very likely that many participating IDUs with a history of shared drug injection may have contracted HIV during incarceration, potentially providing infection routes into community needle/syringe sharing and sexual networks. Consequently, it is very important that comprehensive HIV prevention measures, including NSPs, become widely available for community-based IDUs in order to best control the epidemic among IDUs and prevent further transmission of the infection to broader populations.

Some limitations to this study should be noted. Although a considerable effort was made to recruit comparable IDUs from both neighborhoods, there is a possibility that our comparison was biased because of unknown background characteristics. The cross-sectional design of the study precludes us from drawing causal relationships between access to an NSP and not participating in shared drug injection. Although interviews were conducted by independent interviewers, outreach workers affiliated to the NGO that provides the NSP in the intervention zone assisted in participant recruitment. Because the potential participants were first approached by one of the outreach workers, it is possible that IDUs who were active in the NSP were over-represented in the study population. Therefore, the proportion of IDUs exposed to the NSP (67%) should not be used as an indication of the program's overall coverage in that neighborhood. We relied primarily on self-reported risk behaviors, which are vulnerable to bias through faulty recall ability or social

desirability (Latkin et al. 1993). Shrestha et al. (2006) recently reported on the limitations of relying on self-reported sharing of needles and syringes, rather than on other, more objective, measures.

This study is among the first in Iran to examine the impact of NSP on drug-related risk behaviors in community-based IDUs. The findings of this study indicate that access to an NSP may reduce sharing practices among IDUs in Tehran, Iran. Because the HIV epidemic is currently expanding among this population, it is essential that comprehensive prevention measures, including NSPs, are made available to both incarcerated IDUs and those in the community. Future research can investigate barriers to NSPs among IDUs in neighborhoods in Tehran lacking an NSP, as well as other cities with emerging HIV epidemics among IDUs where prevention interventions have not yet been established.

Acknowledgments The authors would like to express their sincere appreciation to the participants of this study for sharing their experiences. We would also like to extend our special thanks to the staff members of the Persepolis Society and the National Center for Addiction Studies at Tehran University of Medical Sciences, and the survey team, in particular Drs. Mehran Boroumand, Mohammad Sajjadi-Far, and Seyed Reza Hosseini. We also wish to acknowledge Dr. Lisa Maher and Ms. Jane Koerner for their critical comments. This research was supported jointly by the Fujiwara Foundation in Japan and the Tehran University of Medical Sciences and Health Services, Iran.

References

- Bluthenthal, R. N., Kral, A. H., Gee, L., Erringer, E. A., & Edlin, B. R. (2000). The effect of syringe exchange use on high-risk injection drug users: A cohort study. *AIDS, 14*, 605–611. doi: 10.1097/00002030-200003310-00015.
- Center for Disease Management. (2006). *AIDS/HIV Surveillance Report from the Center for Disease Management, Ministry of Health and Medical Education, Iran*. Tehran: Center for Disease Management.
- Chen, H. T., & Liao, Q. (2005). Pilot study of the NGO-based relational intervention model for HIV prevention among drug users in China. *AIDS Education and Prevention, 17*, 503–514. doi:10.1521/acap.2005.17.6.503.
- Cox, G. M., Lawless, M. C., Cassin, S. P., & Geoghegan, T. W. (2000). Syringe exchanges: A public health response to problem drug use. *Irish Medical Journal, 93*, 143–146.
- Gibson, D. R., Brand, R., Anderson, K., Kahn, J. G., Perales, D., & Gudyish, J. (2002). Two- to sixfold decreased odds of HIV risk behavior associated with use of syringe exchange. *Journal of Acquired Immune Deficiency Syndromes, 31*, 237–242.
- Gray, J. (1998). Harm reduction in the hills of northern Thailand. *Substance Use & Misuse, 33*, 1075–1091.
- Hartgers, C., Buning, E. C., van Santen, G. W., Verster, A. D., & Coutinho, R. A. (1989). The impact of the needle and syringe-exchange programme in Amsterdam on injecting risk behaviour. *AIDS, 3*, 571–576. doi:10.1097/00002030-198909000-00003.
- Iran Prison Organization. (2006). *An overview on HIV/AIDS in prisons of The Islamic Republic of Iran [In Persian] by Health and Treatment Headquarter*. Tehran: Iran Prison Organization.

- Latkin, C. A., Vlahov, D., & Anthony, J. C. (1993). Socially desirable responding and self-reported HIV infection risk behaviors among intravenous drug users. *Addiction*, *88*, 517–526. doi:10.1111/j.1360-0443.1993.tb02058.x.
- Power, R., & Nozhkina, N. (2002). The value of process evaluation in sustaining HIV harm reduction in the Russian Federation. *AIDS*, *16*, 303–304. doi:10.1097/00002030-200201250-00028.
- Quan, V. M., Chung, A., & Abdul-Quader, A. S. (1998). The feasibility of a syringe-needle-exchange program in Vietnam. *Substance Use & Misuse*, *33*, 1055–1067.
- Razzaghi, E., Nassirimanesh, B., Afshar, P., Ohiri, K., Claeson, M., & Power, R. (2006a). HIV/AIDS harm reduction in Iran. *Lancet*, *368*, 434–435. doi:10.1016/S0140-6736(06)69132-0.
- Razzaghi, E. M., & Rahimi, M. A. (2005). *Injection drug use in Tehran*. Tehran: Tehran University of Medical Sciences Press.
- Razzaghi, E. M., Rahimi M. A., Green, T. C., & Khoshnood, K. (2006b). Profiles of risk: A qualitative study of injecting drug users in Tehran, Iran. *Harm Reduction Journal*, *3*, article 12. Accessed 30 Oct 30 2007, at <http://www.harmreductionjournal.com/content/3/1/12>.
- Razzaghi, E. M., Rahimi, M. A., Hoseini, M., Madani, S., & Mohammad, K. (2000). *Rapid assessment of drug misuse in Iran, Tehran*. Tehran: Iranian Welfare Organization Press.
- Shrestha, S., Smith, M. W., Broman, K. W., Farzadegan, H., Vlahov, D., & Strathdee, S. A. (2006). Multiperson use of syringes among injection drug users in a needle exchange program: A gene-based molecular epidemiologic analysis. *Journal of Acquired Immune Deficiency Syndromes*, *43*, 335–343. doi:10.1097/01.qai.0000230528.25083.0b.
- Tyndall, M. W., Currie, S., Spittal, P., Li, K., Wood, E., O'Shaughnessy, M. V., et al. (2003). Intensive injection cocaine use as the primary risk factor in the Vancouver HIV-1 epidemic. *AIDS*, *17*, 887–893. doi:10.1097/00002030-200304110-00014.
- UNAIDS. (2007). Country situation analysis-Islamic Republic of Iran. Accessed 30 October 2007, at http://www.unaids.org/en/Regions_Countries/Countries/Iran_Islamic_Republic_of.asp.
- Vazirian, M. (2003). Review of drug demand reduction programs in Iran: Advice for development and strategic planning [in Persian]. *Social Welfare Quarterly*, *9*, 145–201.
- Vazirian, M., De Kort, G., & Nassirimanesh, B. (2006). Young people and drugs—towards a comprehensive health promotion policy—Tehran report. Asian Harm Reduction Network, UNODC, Ministry of Health and Medical Education, Iran.
- Vazirian, M., Nassirimanesh, B., Zamani, S., Ono-Kihara, M., Kihara, M., Mortazavi Ravari, S., & Gouya, M. M. (2005). Needle and syringe sharing practices of injecting drug users participating in an outreach HIV prevention program in Tehran, Iran: A cross-sectional study. *Harm Reduction Journal*, *2*, article 19. Accessed 30 October 2007, at <http://www.harmreductionjournal.com/content/2/1/19>.
- WHO. (2004). Best practice in HIV/AIDS prevention and care for injecting drug abusers: The triangular clinic in Kermanshah, Islamic Republic of Iran. Cairo: Regional Office for the Eastern Mediterranean, World Health Organization. Accessed 30 October 2007, at <http://www.emro.who.int/asd/PDF/STD-052.pdf>.
- Wodak, A., & Cooney, A. (2005). Effectiveness of sterile needle and syringe programmes. *The International Journal of Drug Policy*, *16*, S31–S44. doi:10.1016/j.drugpo.2005.02.004.
- Zamani, S., Kihara, M., Gouya, M. M., Vazirian, M., Nassirimanesh, B., Ono-Kihara, M., et al. (2006). High prevalence of HIV infection associated with incarceration among community-based injecting drug users in Tehran, Iran. *Journal of Acquired Immune Deficiency Syndromes*, *42*, 342–346. doi:10.1097/01.qai.0000219785.81163.67.
- Zamani, S., Kihara, M., Gouya, M. M., Vazirian, M., Ono-Kihara, M., Razzaghi, E. M., et al. (2005). Prevalence of and factors associated with HIV-1 infection among drug users visiting treatment centers in Tehran, Iran. *AIDS*, *19*, 709–716. doi:10.1097/01.aids.0000166094.24069.72.

Original research article

Unintended pregnancy and its risk factors among university students in eastern China

Qiaoqin Ma^{a,b,*}, Masako Ono-Kihara^b, Liming Cong^a, Guozhang Xu^c, Xiaohong Pan^a,
Saman Zamani^b, Shahrzad Mortazavi Ravari^b, Masahiro Kihara^b

^aCenter for Disease Control and Prevention of Zhejiang Province, Hangzhou 310009, China

^bDepartment of Global Health and Socio-epidemiology, Kyoto University School of Public Health, Kyoto 606-8501, Japan

^cCenter for Disease Control and Prevention of Ningbo Municipality, Ningbo 315010, China

Received 13 August 2007; revised 30 September 2007; accepted 15 October 2007

Abstract

Background: The study was conducted to determine the prevalence of unintended pregnancy, induced abortion and contraceptive use, and factors associated with unintended pregnancy among Chinese university students.

Study Design: A self-administered questionnaire survey with cross-sectional design was administered among students in two universities in Ningbo, China, in November–December 2003. Sociodemographic and behavioral factors associated with unintended pregnancy were identified in both genders using univariate and multivariate analyses.

Results: Of sexually active students, 10.6% of male and 11.6% of female students reported their partner or they had a history of pregnancy; 10.0% of male and 11.3% of female students reported their partner or they had a history of induced abortion. The risk factors for unintended pregnancy identified among males by multivariate analysis were older age [odds ratio (OR), 2.12; 95% confidence interval (CI), 1.15–3.88], initiation of sexual activity before high school (OR, 2.45; 95% CI, 1.36–4.44), reported lack of condom use in first sexual activity (OR, 1.71; 95% CI, 1.10–2.64), multiple sexual partners (OR, 1.54; 95% CI, 1.06–2.23), and often used condoms during their lifetime (OR, 1.97; 95% CI, 1.01–3.81). The identified risk factors among females were initiation of sexual activity before high school (OR, 5.12; 95% CI, 1.49–17.68), non-consensual sexual intercourse as first sex (OR, 1.77; 95% CI, 1.08–2.90), multiple partners (OR, 2.75; 95% CI, 1.61–4.71), and sometimes/never (OR, 3.02; 95% CI, 1.16–7.87) or often (OR, 3.92; 95% CI, 1.43–10.73) used condoms during their lifetime.

Conclusion: The high prevalence of unintended pregnancies and induced abortions in this population indicates a need for better and targeted sex education and family planning services.

© 2008 Elsevier Inc. All rights reserved.

Keywords: Unintended pregnancy; Abortion; Contraceptive use; University; Students

1. Introduction

During the past two decades, China has experienced dramatic social change, associated with rapid economic growth and reform. Traditional attitudes toward sex, marriage and family have changed considerably. Young people, including university students, consider premarital sex more acceptable, and their sexual behavior and attitudes have become more open [1,2].

Many studies in developing countries have shown that young people do not receive sufficient information about contraception and disease prevention, which often results in misconceptions about reproduction [3]. The sexual and reproductive health of young people could become a significant public health problem in China because of the real and perceived increase in sexual activity, premarital pregnancies, abortions and sexually transmitted infections (STIs) among young people. Research on sexuality and the sexual behavior of Chinese university students in the last decade has shown that 9–16% of students report that they are sexually experienced [4–6]. In 2005, the Chinese Ministry of Education lifted the ban on the marriage of university students [7]; the sexual activity among Chinese university

* Corresponding author. Center for Disease Control and Prevention of Zhejiang Province, Hangzhou 310009, China. Tel.: +86 571 8723 5170; fax: +86 571 8723 5048.

E-mail address: qiaoqinma@yahoo.com.cn (Q. Ma).

students will increase rapidly as a result of this policy and the continuing rapid social and economic change in China. However, thus far, little attention has been given to reproductive health issues, including contraception, unintended pregnancies and induced abortions in this population. The studies on unintended pregnancies and induced abortions among unmarried young people in China have focused mainly on women attending premarital examinations [8] and those seeking to terminate a pregnancy at clinics and hospitals [9–12]. Thus, there is little literature documenting other aspects of sexual behavior in university students.

In the present study, we primarily report data on unintended pregnancies and the associated factors among students at two universities in an eastern Chinese city to assess the determinants of the pregnancies and the effectiveness of current contraceptive services and education in meeting the needs of students.

2. Materials and methods

2.1. Setting and participants

This study was conducted in Ningbo Municipality, Zhejiang Province. Ningbo, a large city with a population of 5.5 million on the eastern coast of China, is one of the most economically developed areas in the country. The city is home to two universities, both of which participated in the study. All students in Grades I to IV at both universities were recruited. Of the 29,409 possible respondents, 22,493 responded validly, giving a final response rate of 76.5%. Of the 22,493 respondents, 17.6% (1,981/11,255) of males and 8.6% (963/11,238) of females were sexually active. Only sexually active students who responded to the question about a history of pregnancy were included in the analysis. Thus, the final sample comprised 1835 males and 848 females. We considered all the pregnancies reported here to be unintended because, at the time of the study, the Ministry of Education prohibited all university students from marrying, and pregnant students could have been discharged from a university for breaching regulations.

2.2. Data collection

The instrument was developed through reviews of Chinese and international literature and was modified using the results of qualitative studies that included in-depth interviews with 11 students (three males and three females from one university, and three males and two females from the other) and interviews with four focus groups (one for each gender at each university) comprising a total of 21 students recruited by university counselors from the two universities. The revised instrument was pilot-tested at one of the universities in a group of 50 students randomly selected from various grades and faculty members of both genders. The instrument's reliability was evaluated in 89 of 160 college students recruited from another city who could be matched between the two tests with a 1-week interval. More

than 70% of the categorical variables had Kappa statistics over 0.4 (all $p < .05$), and the coefficient of correlation for discrete variables varied from 0.63 to 0.93 (all $p < .05$).

For the main survey, students were asked by university staff and student leaders to come to classrooms at specified times outside lecture hours in order to complete the self-administered questionnaire. Field teams consisted of trained staff from the local Centers for Disease Control and Prevention and counselors from each university. Data were collected from November to December 2003.

2.3. Ethical considerations

Those responsible for institutional review at Zhejiang Province's Center for Disease Control and Prevention, the two universities involved and the Ningbo Education Board approved the study protocol. All students were advised of the study's purpose and were told that nonparticipation would cause no disadvantage to them. The students were assured that their privacy and confidentiality would be strictly protected because no personal names or other identifiers were to be included in the questionnaire and the data were to be presented only in an aggregate manner.

2.4. Statistical analysis

For the analyses, SPSS for Windows (Version 12.01; SPSS, Chicago, IL) was used. We first determined which factors were associated with a history of pregnancy in themselves or a partner in univariate analyses. Variables significant in the univariate analyses ($p < .05$) other than condom and oral contraceptive (OC) use during the most recent year were entered in a multivariate logistic regression model that was performed by a backward stepwise logistic regression, with p value $> .10$ as a criterion for removal of a variable from the model.

3. Results

Of the sexually active students, 11.6% females reported a history of pregnancy, and 10.6% males reported that a partner had such a history. A history of induced abortion was reported by 11.3% females, and 10.0% males reported this history in a partner (Table 1). Up to 33.7% females and 22.5% males thought they or a partner could possibly or highly possibly become pregnant by sexual intercourse, with females having significantly higher risk awareness than males ($\chi^2 = 53.271$, $p < .001$).

The majority of the students, male and female, were over 19 years old, were from urban areas and had a middle-class family economic status (Table 2). Among males, 66% had ever smoked cigarettes; this rate was only 13% in females. In both genders, over 79% and 66% had ever consumed alcoholic beverages and danced, respectively. According to the univariate analysis, having ever smoked cigarettes and dancing in women and older age and having ever smoked cigarettes in men were related to a history of pregnancy in the

Table 1
Reproductive health and risk awareness among sexually active students

	Male (n=1835)		Female (n=848)	
	n	%	n	%
History of unintended pregnancy				
Yes	194	10.6	98	11.6
No	1546	84.3	725	85.5
Not sure	95	5.2	25	2.9
History of induced abortion				
Yes	184	10.0	96	11.3
No	1536	83.7	729	86.0
Not sure	105	5.7	20	2.4
Risk awareness for pregnancy				
Not-low possible	5	72.9	496	58.8
Some-high possible	413	22.5	286	33.7
Not sure	84	4.6	63	7.4

female or the male's partner. Hometown area, family economic status, and alcohol beverage consumption were unrelated to pregnancy in both males' partners and females.

The rate of condom use during the first sex experience was 30.0% for males and 26.6% for the females' partners, whereas the condom use rates during the most recent sexual encounter were 48.3% and 46.0% for males and females, respectively (Table 3). Among the students, 14.5% males and 18.9% females reported that they or their partner always used a condom throughout their lifetime, and among these "always" users, 4.6% males and 3.2% females reported that their partner or they had a history of pregnancy. The reported pregnancy rate was similar between those who often used condoms and those who sometimes/never used one, in both genders. Oral contraceptive use during the most recent sex was 26.5% for the males' partners and 29.2% for females.

With regard to risky behaviors associated with pregnancy, students who initiated sex at a younger age (vs. those who initiated later), students who did not report using a condom during first sex (vs. those who did), students who had multiple sexual partners (vs. those with only one), students who often or sometimes/never used condoms throughout their lifetime (vs. those who always did), students who reported having been diagnosed with an STI (vs. those with no such history) and students who often or sometimes/never used condoms and OCs during the most recent year (vs. those who always did) were more likely to report having a history of pregnancy during their lifetime, for both males' partners and female students. Non-consensual sex as a first sex experience was related to pregnancy among women. Condom use and contraceptive use during the most recent sex for both males and females were not associated with a history of pregnancy in males' partners or females.

Multivariate logistic regression analysis for men showed that older age [odds ratio (OR), 2.12; 95% confidence interval (CI), 1.15–3.88], initiation of sexual activity before high school (OR, 2.45; 95% CI, 1.36–4.44), reported lack of condom use at first sex (OR, 1.71; 95% CI, 1.10–2.64), multiple sex partners (OR, 1.54; 95% CI, 1.06–2.23) and often condom use during their lifetime (OR, 1.97; 95% CI, 1.01–3.81) remained as risk factors for pregnancy in their partners (Table 4). Multivariate logistic regression analysis for women revealed that initiation of sexual activity before high school (OR, 5.12; 95% CI, 1.49–17.68), non-consensual sexual intercourse at first sex (OR, 1.77; 95% CI, 1.08–2.90), multiple sexual partners (OR, 2.75; 95% CI, 1.61–4.71) and sometimes/never (OR, 3.02; 95% CI, 1.16–7.87) or often (OR, 3.92; 95% CI, 1.43–10.73)

Table 2
Sociodemographic characteristics and factors for unintended pregnancy among sexually active students

	Male (n=1835)			Female (n=848)		
	Total n* (%)	Pregnancy n (%)	COR (95% CI)	Total n* (%)	Pregnancy n (%)	COR (95% CI)
Age						
≤19	257 (14.1)	16 (6.2)	1	186 (22.0)	19 (10.2)	1
≥20	1567 (85.9)	176 (11.2)	1.91 (1.12–3.24)	660 (78.0)	77 (11.7)	1.16 (0.68–1.97)
Hometown area						
Rural	434 (23.7)	44 (10.1)	1	165 (19.5)	17 (10.3)	1
Urban	1394 (76.3)	149 (10.7)	1.06 (0.74–1.51)	681 (80.5)	81 (11.9)	1.18 (0.68–2.04)
Family's economic status						
Rich	231 (12.6)	32 (13.9)	1.47 (0.73–2.98)	87 (10.3)	14 (16.1)	0.96 (0.36–2.59)
Between	1478 (80.7)	149 (10.1)	1.03 (0.55–1.91)	719 (84.8)	77 (10.7)	0.60 (0.26–1.40)
Poor	122 (6.7)	12 (9.8)	1	42 (5.0)	7 (16.7)	1
Cigarette smoking						
Never	618 (33.7)	44 (7.1)	1	737 (87.0)	78 (10.6)	1
Ever	1216 (66.3)	150 (12.3)	1.84 (1.29–2.61)	110 (13.0)	20 (18.2)	1.88 (1.10–3.22)
Alcoholic drinking						
Never	122 (6.6)	12 (9.8)	1	175 (20.6)	13 (7.4)	1
Ever	1713 (93.4)	182 (10.6)	1.09 (0.59–2.02)	673 (79.4)	85 (12.6)	1.80 (0.98–3.31)
Dancing						
Never	619 (33.8)	71 (11.5)	1	240 (28.3)	16 (6.7)	1
Ever	1215 (66.2)	123 (10.1)	0.87 (0.64–1.19)	608 (71.7)	82 (13.5)	2.18 (1.25–3.81)

COR, crude odds ratio.

* Total may not add up to 1835 for male and 848 for female due to missing data.

Table 3
Sexual behavior characteristics and factors for unintended pregnancy among sexually active students

	Male (n=1835)			Female (n=848)		
	Total n ^a (%)	Pregnancy n (%)	COR (95% CI)	Total n ^a (%)	Pregnancy n (%)	COR (95% CI)
Period of first sex						
Before high school	109 (6.0)	28 (25.7)	3.49 (2.16–5.64)	18 (2.1)	10 (55.6)	12.31 (4.68–32.39)
High school	670 (36.8)	72 (10.7)	1.22 (0.88–1.68)	186 (22.0)	28 (15.1)	1.75 (1.08–2.83)
University	1043 (57.2)	94 (9.0)	1	640 (75.8)	59 (9.2)	1
Condom use during first sex						
Yes	547 (30.0)	40 (7.3)	1	225 (26.6)	15 (6.7)	1
No/not sure	1276 (70.0)	154 (12.1)	1.74 (1.21–2.50)	621 (73.4)	82 (13.2)	2.13 (1.20–3.78)
Consent at first sex						
Consent	1748 (96.7)	185 (10.6)	1	525 (63.0)	50 (9.5)	1
Nonconsent	59 (3.3)	6 (10.2)	0.96 (0.41–2.26)	308 (37.0)	45 (14.6)	1.63 (1.06–2.50)
No. of sexual partner during their lifetime						
1	1076 (63.7)	86 (8.0)	1	640 (78.9)	49 (7.7)	1
≥2	614 (36.3)	90 (14.7)	1.98 (1.44–2.71)	171 (21.1)	43 (25.1)	4.05 (2.58–6.37)
Condom use during their lifetime						
Always	261 (14.5)	12 (4.6)	1	156 (18.9)	5 (3.2)	1
Often	532 (29.6)	65 (12.2)	2.89 (1.53–5.45)	197 (23.9)	28 (14.2)	5.00 (1.88–13.29)
Sometimes/never	1002 (55.8)	117 (11.7)	2.74 (1.49–5.05)	471 (57.2)	65 (13.8)	4.84 (1.91–12.24)
History with diagnosed STI						
Yes	25 (1.4)	8 (32.0)	4.11 (1.75–9.67)	14 (1.8)	6 (42.9)	6.25 (2.12–18.46)
No	1704 (98.6)	175 (10.3)	1	775 (98.2)	83 (10.7)	1
Condom use during most recent year ^b						
Always	202 (15.1)	11 (5.4)	1	105 (19.4)	5 (4.8)	1
Often	363 (27.1)	48 (13.2)	2.65 (1.34–5.22)	114 (21.0)	19 (16.7)	4.00 (1.43–11.14)
Sometimes/never	775 (57.8)	107 (13.8)	2.78 (1.47–5.28)	323 (59.6)	49 (15.2)	3.58 (1.39–9.23)
OC use during most recent year ^b						
Always	104 (7.6)	6 (5.8)	1	81 (14.3)	5 (6.2)	1
Often	208 (15.2)	26 (12.5)	2.33 (0.93–5.86)	89 (15.8)	15 (16.9)	3.08 (1.07–8.91)
Sometimes/never	1052 (77.1)	136 (12.9)	2.43 (1.04–5.64)	395 (69.9)	59 (14.9)	2.67 (1.04–6.88)
Condom use during most recent sex						
Yes	836 (48.3)	76 (9.1)	1	368 (46.0)	37 (10.1)	1
No	894 (51.7)	104 (11.6)	1.32 (0.96–1.80)	432 (54.0)	56 (13.0)	1.33 (0.86–2.07)
OC use during most recent sex						
Yes	454 (26.5)	44 (9.7)	1	224 (29.2)	20 (8.9)	1
No	1257 (73.5)	135 (10.7)	1.12 (0.78–1.61)	543 (70.8)	73 (13.4)	1.58 (0.94–2.67)

^a Total may not add up to 1835 for male and 848 for female due to missing data.

^b Students who were sexually active in the most recent year: n=1403 for male and 585 for female; total may not be so large due to missing data.

condom use during their lifetime remained as risk factors for pregnancy. Having smoked cigarettes among males and having danced among females were also associated with unintended pregnancy.

4. Discussion

To our knowledge, these are the first data regarding the reproductive health of Chinese university students. The high prevalence of pregnancy and abortion found in this study indicates that sex education and contraceptive services for young people in China do not seem to be adequate.

In China, family planning programs are now widely available, free of charge. However, these services are used primarily by adults and rarely by unmarried young people. Although the acceptance of premarital sex is increasing in China, unmarried youth still face the risk of being stigmatized by their search for contraceptive services, and the lack of counseling and privacy in these services prevents

many young people from seeking contraception [13]. Further, family planning workers are often ambivalent about the provision of sexual and reproductive health information to unmarried young people [14], and parents and teachers still avoid discussing sensitive topics such as premarital sex, contraception, and abortion with young people [15]. These barriers may cause young people to have lower risk awareness and lower use of contraception, resulting in the high rate of pregnancy seen in this study. In unmarried young women, especially young students, being pregnant is still considered unacceptable in Chinese culture. Data from China have shown that the primary reason for unmarried women to undergo induced abortions is simply that they are unmarried [8,12], and this could be true in the present study as well.

When used consistently and correctly, condoms offer considerable protection against unintended pregnancy as well as STIs, including HIV infection. The condom is the most popular method of contraception among young Chinese [5,13,16,17]. However, our data revealed that

Table 4
Factors associated with unintended pregnancy among sexually active students by genders

Risk factors	Male	Female
	AOR (95% CI)	AOR (95% CI)
Age		
≤19	1	
≥20	2.12 (1.15–3.88)	
Period of first sex		
Before high school	2.45 (1.36–4.44)	5.12 (1.49–17.68)
High school	1.05 (0.71–1.55)	1.12 (0.63–1.99)
University	1	1
Consent at first sex		
Consent		1
Nonconsent		1.77 (1.08–2.90)
Condom use during first sex		
Yes	1	
No/not sure	1.71 (1.10–2.64)	
No. of sexual partner during their lifetime		
1	1	1
≥2	1.54 (1.06–2.23)	2.75 (1.61–4.71)
Condom use during their lifetime		
Always	1	1
Often	1.97 (1.01–3.81)	3.92 (1.43–10.73)
Sometimes/never	1.49 (0.77–2.88)	3.02 (1.16–7.87)

Ever cigarette smoking among males and ever dancing among females were associated with unintended pregnancy as well. AOR, adjusted odds ratio.

"always used a condom during their lifetime" was true for less than 20% of sexually active males and of females' partners. In addition, of those "always" condom users, 4.6% of males and 3.2% of females reported that their partner or they had a history of pregnancy. This rate was significantly lower than that for "not always" users and may be attributable to condom use failures among young people [10,18], suggesting that sex education should address the correct use of a condom while also emphasizing the importance of consistent condom use for the prevention of unintended pregnancy and STIs.

We found that the female partners of male students who used a condom in first sex had a significantly lower pregnancy rate. It may be that those who used condoms during first sex were more likely to also use them later [19,20], highlighting the importance of condom use in first sex. Our data also showed that the condom use rate increased greatly between first sex and most recent sex, for both males and females' partners. Thus, it appeared that condom use increased as sex life become longer. Studies in China have shown that OC use during first sex was 4.1% for males' partners and 6.3% for females among university students [16], and 4.9% for both genders among unmarried persons [13]. Our findings indicated that the OC use rate during the most recent sex was 26.5% for males' partners and 29.5% for females. Studies in other countries have reported that the likelihood of using contraception increases with age in sexually active young people [21–24]. If this pattern applies in China, then students who are at an early stage of their sex life would be less likely to use condoms and OCs and thus

would be more vulnerable to pregnancy and STIs than those further along in their sexual life.

Condoms, OCs, periodic abstinence and withdrawal are the most commonly used contraceptive methods among university students [5,16] and unmarried young people [13,17]. It is possible that some of the students who did not use condoms or OCs may have used periodic abstinence or withdrawal. However, both of these methods could lead to contraceptive failure [10,11]. The use of these two methods could have contributed to the unintended pregnancies reported here as well.

We found that young people who commenced sexual activity before high school were more likely to have experienced an unintended pregnancy than those who initiated sex later; they may have been less aware of the risks for pregnancy and thus used contraception less often. Indeed, it has been reported in studies in China and elsewhere that unmarried young people who initiate sex at early ages use contraception less often than do later initiators [10,21,25,26]. This pattern may explain the high rate of pregnancy among early initiators in our study. Further, earlier initiation of sexual intercourse exposes students to an extended period during which they are at risk for becoming pregnant.

It is a major concern that having multiple partners was a predictor of pregnancy in both genders. Compared with students with one partner, students with more than one partner tended to be less likely to use an effective contraceptive method. It is suggested that those students who engaged in multiple sexual partnership without valid protection also put themselves at great risk for STIs. Students who had become pregnant were evidently at greater risk for STIs because they clearly were not effectively protected with a barrier method, mainly condoms, during sex.

For women, non-consensual sexual intercourse at first sex was related to pregnancy in the multivariate analysis. Chinese studies have reported that unmarried women with a history of non-consensual sex were more likely to have initiated sex earlier, to have had multiple partners, to have undergone induced abortions and to have contracted STIs [27,28]. Non-consensual sex was also associated with unintended pregnancies in India and America [29–31]. Given that 37% of females reported that their first sex was non-consensual, the actual history of non-consensual sex over a lifetime could be much higher in females. Reproductive health education should clearly target non-consensual sex as well as its associated contexts and consequences.

This study has some limitations. The data presented here were derived from a sexual behavior survey of university students. Determining the prevalence of and risk factors for unintended pregnancies in students were not the primary objectives, and thus some important information, including OC use at first sex and over a lifetime, and non-consensual sex over a lifetime, was not provided. Second, reporting bias could be an issue because the information on sexual behavior was excluded for those sexually active students

who did not respond to the question about a history of pregnancy. Third, no cause-effect relationship can be established because of the cross-sectional design of the study. Further research is needed to explore unintended pregnancy and how sexual behaviors and lifestyle choices, such as cigarette smoking and dancing, affect the risk of unintended pregnancy in this population.

Sexually active university students in China have a high risk of unintended pregnancies and induced abortions. Sex education programs conducted at Chinese universities should focus on not only HIV/STI prevention but also avoidance of unintended pregnancies, explaining the possible risk factors associated with unprotected sex and unintended pregnancies. A concerted effort that includes increased awareness of reproductive health, the appropriate introduction of contraceptive information, the availability of student-friendly family planning services and the facilitation of access to contraception for young people is urgently required to meet the reproductive health needs of this population.

Acknowledgments

The Japanese Foundation for AIDS Prevention supported this research. The authors thank the research team, the students, and the two universities involved for their support and cooperation.

References

- Zhang K, Li D, Li H, Beck EJ. Changing sexual attitudes and behavior in China: implications for the spread of HIV and other sexually transmitted diseases. *AIDS Care* 1999;11:581–9.
- Higgins LT, Zheng M, Liu YL, Sun CH. Attitudes to marriage and sexual behaviors: a survey of gender and culture differences in China. *Sex Role* 2002;46:75–89.
- World Health Organization, UNICEF. A picture of health: A review and annotated bibliography of the health of young people in developing countries. Geneva: WHO; 1995.
- Li A, Wang A, Xu B. University students' attitudes toward premarital sex and their sexual activity in Beijing. *Sexology (in Chinese)* 1998;7:19–24.
- Research Group on Sex Education among University Students. Report on sexual behavior survey among Chinese university students in 2000. *Youth Study (in Chinese)* 2001;12:31–9.
- Zhang L, Gao X, Dong Z, et al. Premarital sexual activities among students in a university in Beijing, China. *Sex Transm Dis* 2002;29:212–5.
- China Ministry of Education (CMOE). No. 21 order of Ministry of Education of the People's Republic of China: Regulation for management of students in higher education institutions. Beijing: CMOE; 2005.
- Qian X, Tang S, Garner P. Unintended pregnancy and induced abortion among unmarried women in China: a systematic review. *BMC Health Serv Res* 2004;4:1.
- Wu J, Wang L, Rauyajin O, Good S. Contraceptive use behavior among never married young women who are seeking pregnancy termination in Beijing. *Chin Med J* 2002;115:851–5.
- Cheng Y, Gao X, Li Y, Li S, Qu A, Kang B. Repeat induced abortions and contraceptive practices among unmarried young women seeking an abortion in China. *Int J Gynaecol Obstet* 2004;87:199–202.
- Cheng Y, Zhu W, Li Z, Zhang Y, Wang A. Contraceptive practices of women requesting termination of pregnancy: a study from China. *Contraception* 1997;55:15–7.
- Luo L, Wu SZ, Chen XQ, Li MX, Pullum TW. Induced abortion among unmarried women in Sichuan province, China. *Contraception* 1995;51:59–63.
- Zhang LY, Jejeebhoy S, Shah IH, Zhang LH, Hsia J, Im-em W. Access to contraceptive services among unmarried young people in the north-east of China. *Eur J Contracept Reprod Health Care* 2004;9:147–54.
- Tu X, Cui N, Lou C, Gao E. Do family-planning workers in China support provision of sexual and reproductive health services to unmarried young people? *Bull WHO* 2004;82:274–80.
- Huang J, Bova C, Fennie KP, Rogers A, Williams AB. Knowledge, attitudes, behaviors, and perceptions of risk related to HIV/AIDS among Chinese university students in Hunan, China. *AIDS Patient Care STDS* 2005;19:769–77.
- Li AL, Li LM, Ma CH, Ye GJ. The analysis on contraceptive knowledge, attitudes, and behavior of university students in Beijing. *Chin J Fam Plann* 2000;59:112–5.
- Che Y, Cleland J. Contraceptive use before and after marriage in Shanghai. *Stud Fam Plann* 2003;34:44–52.
- Fu H, Darroch JE, Haas T, Ranjit N. Contraceptive failure rates: new estimates from the 1995 National Survey of Family Growth. *Fam Plann Perspect* 1999;31:56–63.
- Shafiq T, Stovel K, Davis R, Holmes K. Is condom use habit forming? Condom use at sexual debut and subsequent condom use. *Sex Transm Dis* 2004;31:366–72.
- Klavns I, Rodrigues LC, Wellings K, Weiss HA, Hayca R. Increased condom use at sexual debut in the general population of Slovenia and association with subsequent condom use. *AIDS* 2005;19:1215–23.
- Santelli JS, Warren CW, Lowry R, Sogolow E, Collins J, Kann L. The use of condoms with other contraceptive methods among young men and women. *Fam Plann Perspect* 1997;29:261–7.
- Svare EI, Kjaer SK, Poll P, Bock JE. Determinants for contraceptive use in young, single, Danish women from the general population. *Contraception* 1997;19:287–94.
- Blanc AK, Way AA. Sexual behavior and contraceptive knowledge and use among adolescents in developing countries. *Stud Fam Plann* 1998;29:106–16.
- Saleem S, Bobak M. Women's autonomy, education and contraception use in Pakistan: a national study. *Reprod Health* 2005;2:8.
- Manzini N. Sexual initiation and childbearing among adolescent girls in KwaZulu Natal, South Africa. *Reprod Health Matters* 2001;9:44–52.
- Bender SS, Kosunen E. Teenage contraceptive use in Iceland: a gender perspective. *Public Health Nurs* 2005;22:17–26.
- Wu J, Wang L, Zhao G, Zhang X. Sexual abuse and reproductive health among unmarried young women seeking abortion in China. *Int J Gynaecol Obstet* 2006;92:186–91.
- Yimin C, Shouqing L, Arzhu Q, Yuke Z, Jianhua W, Jinxin Z. Sexual coercion among adolescent women seeking abortion in China. *J Adolesc Health* 2002;31:482–6.
- Jejeebhoy SJ. Adolescent sexual and reproductive behavior: a review of the evidence from India. *Soc Sci Med* 1998;46:1275–90.
- Stock JL, Bell MA, Boyer DK, Connell FA. Adolescent pregnancy and sexual risk-taking among sexually abused girls. *Fam Plann Perspect* 1997;29:200–3, 227.
- Wingood GM, DiClemente RJ. Child sexual abuse, HIV sexual risk, and gender relations of African-American women. *Am J Prev Med* 1997;3:380–4.

Research article

Open Access

Sexual behaviors and their correlates among young people in Mauritius: a cross-sectional study

Yumiko H Nishimura*¹, Masako Ono-Kihara¹, Jagdis C Mohith²,
Renaud NgManSun³, Takayuki Homma⁴, Ralph J DiClemente⁵, Delia L Lang⁵
and Masahiro Kihara¹

Address: ¹Department of Global Health and Socio-epidemiology, Kyoto University School of Public Health, Yoshida-Konoe-cho, Sakyo-ku, Kyoto-shi, Kyoto, 606-8501, Japan, ²Mauritius Institute of Health, Powder Mill, Pamplemousses, Mauritius, ³Indian Ocean Commission, Farquhar Avenue, Quatre Bornes, Mauritius, ⁴Division of Pharmaceutical Sciences, Graduate School of Natural Science and Technology, Kanazawa University, Kakuma-machi, Kanazawa, 920-1192, Japan and ⁵Department of Behavioral Sciences and Health Education, Rollins School of Public Health of Emory University, 1518 Clifton Road, Atlanta, Georgia 30322, USA

Email: Yumiko H Nishimura* - yumiko@pbh.med.kyoto-u.ac.jp; Masako Ono-Kihara - okmasako@pbh.med.kyoto-u.ac.jp; Jagdis C Mohith - mihealth@intnet.mu; Renaud NgManSun - renaud.ngmansun@coi-ioc.org; Takayuki Homma - thomma@p.kanazawa-u.ac.jp; Ralph J DiClemente - rdiclem@sph.emory.edu; Delia L Lang - dlang2@sph.emory.edu; Masahiro Kihara - poghse@pbh.med.kyoto-u.ac.jp

* Corresponding author

Published: 5 October 2007

Received: 4 April 2007

Accepted: 5 October 2007

BMC International Health and Human Rights 2007, 7:8 doi:10.1186/1472-698X-7-8

This article is available from: <http://www.biomedcentral.com/1472-698X/7/8>

© 2007 Nishimura et al.; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Little is known about the HIV/AIDS epidemic in the Indian Ocean region, including Mauritius. National records suggest a prevalence of HIV in Mauritius of < 1% in the general population, which is one of the lowest prevalence rates in southern Africa. However, HIV-positive cases have been increasing recently in Mauritius. We conducted a cross-sectional survey in January 2003 to assess the prevalence of HIV-related sexual behaviors and their correlates among young people aged 15–24 years in Mauritius.

Methods: We identified 1200 participants using two-stage cluster sampling. Demographic, social, sexual, and knowledge of HIV/AIDS data were obtained in face-to-face interviews using a structured questionnaire administered by trained interviewers. The prevalence of sexual behaviors was described in relation to gender, and the correlates of ever having had sex and nonuse of condom at last sex were analyzed using logistic regression.

Results: In the target population, 30.9% of males and 9.7% of females reported a history of sexual intercourse. Of the currently sexually active participants, 50.6% of men and 71.2% of women did not use condoms at their last sexual encounter. Logistic regression revealed that work experience and marijuana use were significantly associated with men's sexual experience, whereas being out of school and drinking experience were significantly associated with women's sexual experience. For both men and women, being Christian and visiting nightclubs were associated with having ever had sexual intercourse ($P < 0.05$). In addition, not using a condom at the first sexual encounter and lack of exposure to a nongovernmental organization (NGO) dealing with HIV/AIDS were associated with the nonuse of condoms at the last sexual encounter ($P < 0.05$).

Conclusion: Young people in Mauritius are at risk of a future HIV epidemic because behaviors predisposing to HIV infection are prevalent among sexually experienced youth. A focused prevention program targeting young people should be reinforced as part of the National AIDS Control Program, taking into account the predictors of sexual behaviors identified here.

Background

Mauritius is an island country with a population of 1.2 million, located in the Indian Ocean about 900 km east of Madagascar. The country is noted for its current rapid economic growth and the cultural diversity of its inhabitants, who came from Asia, Africa, and Europe. Little information is available about the status of the HIV/AIDS epidemic on Mauritius and in the Indian Ocean region [1,2].

National records report 374 cumulative HIV/AIDS cases (238 males and 136 females) among Mauritians at the end of 2002 since the first AIDS case was reported in 1987. Of these cases, almost 70% occurred through heterosexual contact and nearly 15% through either heterosexual contact or injecting drug use (IDU), although IDU-related cases have increased rapidly as of 2003. The peak age of reported cases up to 2002 was 25–39 years for men and 20–29 years for women. Based on HIV antibody tests for blood donors and pregnant women, the estimated HIV prevalence in the general population of Mauritius is < 1%, which is one of the lowest prevalence in the African region. However, concerns about an HIV epidemic has been growing in this country because the number of reported HIV/AIDS cases continued to increase during the last decade, and suddenly doubled from 50 cases in 2000 to 98 cases in 2002 [3]. In addition, Mauritius hosts more than 700,000 tourists annually, mainly from Europe, South Africa, and India, where HIV epidemics are more pronounced than in Mauritius. Tourists can potentially transmit HIV and sexually transmitted diseases (STDs) from country to country [4]. Growing number of people from overseas is causing concerns about a potential HIV epidemic in Mauritius.

The government of Mauritius launched the National HIV/AIDS Strategic Plan in 2001 and identified 12 strategic objectives [3]. After a reduction in new STI/HIV infections among groups with high-risk behaviors (sex workers, IDUs, men who have sex with men, and prison inmates), reducing vulnerability among the youth and children was the third strategic objective of the plan. About 17% of reported HIV/AIDS up to 2001 were among those aged 15–24. Although more cases were reported among those aged 25–39 for men and 20–29 for women, actual disease contraction must have occurred when these people were younger because people are typically unaware of their infection for several years. Prevention of HIV/AIDS among young people is thus one of the country's priorities.

Several studies have examined the sexual behavior of young people in Mauritius. One study of female Economic Processing Zone workers in 1994 revealed that 68% of young female workers reported some sort of intimate behavior with their boyfriends, although only 7%

reported full penetration (*i.e.*, penile-vaginal sex) [5]. A nationwide survey of male and female adolescent behavior using two-stage clustered sampling was conducted as part of the Youth Profile of the Republic of Mauritius in 1996. According to the survey, 43% of never-married males and 11% of never-married females aged 18–25 years reported having experienced sexual intercourse, and the mean age at first sexual intercourse was approximately 18 years [6]. Other surveys were undertaken to assess the knowledge, attitudes, behaviors, and practices (KABP) related to HIV/AIDS among individuals aged 15–49 years [7–9].

Although some studies have yielded useful information, they have several limitations in aiding the design of HIV prevention programs for young people. For example, some of these studies targeted subgroups of young people such as female workers or people approached using convenient sampling methods, thus limiting the generalizability of the results. Other studies included populations other than young people, yielding limited information on young people. In addition, the studies assessed the sexual behaviors of people in terms of certain demographic characteristics, and did not adequately assess them in relation to social and cultural factors, although these factors are undoubtedly important determinants of HIV-associated behavior and are important in designing prevention programs [10–12]. Most of the studies were conducted in the mid-1990s; thus, the information needs to be updated because the prevalence of HIV has increased markedly in the ensuing years.

We designed our study to assess the current HIV-related sexual behaviors of unmarried young people and to investigate the correlates of sexual debut and condom nonuse. More broadly, we assessed the possible influence in the lives of young people of changing Mauritian society brought about by modernization and globalization, *e.g.*, educational attainment and exposure to a variety of socializing items, in connection with sexual behavior. Our ultimate objective was to assist the National AIDS Control program of Mauritius to monitor, evaluate, and reformulate its HIV prevention program for young people.

Methods

Sampling frame

The target population was never-married men and women aged 15–24 years living on the island of Mauritius. This population consisted of approximately 173,200 people, which was about 15% of the total population of Mauritius in 2000 [13].

A two-stage cluster sampling method was used to select respondents. The enumeration areas (EAs) of the Central Statistics Office were used as the sampling frame for the

first stage. Each EA consisted of approximately 300–600 people comprising 75–150 households, and those in the capital city of Port Louis and Plaines Wilhems District were defined as being from urban areas (43% of the total population). In the first stage, 50 EAs from a total of 3472 were selected using the probability proportional to size (PPS) method [14,15]. All never-married persons aged 15–24 years in selected areas were listed during the preparatory fieldwork for a total of 2592 persons. In the second stage, 12 males and 12 females from each EA were selected using a computer-generated random numbers algorithm. Using this multistage procedure, 600 males and 600 females were chosen to participate in the survey. The sample size was chosen to allow us to detect a 10% difference in sexual experience rate between in-school and out-of-school students with sufficient statistical significance ($\alpha = 0.05$, two-tailed) and a power of 0.8, as well as to adjust for the confounding effect of multiple factors.

Of the young people selected, 59 respondents (4.9%) did not participate. The primary reasons for nonparticipation were "moved away," "gone abroad," "refused (by himself/herself or parents)," and "married." Consequently, 1141 young people (575 men and 566 women) completed the survey interviews, yielding a response rate of 95.1%. Comparisons of participants and nonparticipants showed no marked difference in sociodemographic characteristics with respect to gender, religion, or age.

Instrument development

The research questionnaire was developed based on the Family Health International (FHI) HIV/AIDS/STD Behavioral Surveillance Surveys (BSS) Questionnaire for youth [15]. This instrument includes questions that can assess internationally agreed-upon indicators for young people's behaviors in relation to HIV/AIDS [16].

A multiphase process was used to develop the research instrument to ensure that it was developmentally, culturally, and linguistically appropriate. First, seven focus group discussions were conducted in youth centers to tailor the instrument culturally and linguistically for Mauritian youth. Second, persons working in nongovernmental organizations (NGOs) that deal with HIV/AIDS and drug use were invited to review the instrument and make recommendations for its modification. Third, the English instrument was translated into Creole and back-translated into English to confirm the accuracy of the Creole translation. Finally, the instrument was pilot-tested with 50 young people during preparatory fieldwork. Fieldworkers reported the items for which participants had difficulty in understanding the phrase or terms of the question after the pretest interview. The questionnaire was modified based on the fieldworkers' comments and finalized in January 2003.

The final version of the instrument included the following domains of interest: sociodemographic characteristics (19 items), social life (15 items), knowledge and attitude about condoms (10 items), sexual experience (22 items), and knowledge of HIV/AIDS and exposure to an HIV/AIDS program (30 items). A complete questionnaire may be obtained from the first author.

The sociodemographic variables included sex, age, religion, educational background, and work experience. Religion represents racial/ethnic background of the participants. Usually, Hindus represent people whose ancestors came from India, Muslims, from India and the Middle East, and Christians, from Africa, Europe, and China. Concerning social life, questions were asked regarding respondents' mobile phone use, nightclub visits, pornographic film viewing, Internet use, and alcohol and other drug use. Items concerning knowledge and attitude about condoms included questions related to obtaining and the ability to use condoms.

In the section on sexual behavior, young people were first asked about lifetime sexual experiences. Sexual intercourse was defined as vaginal or anal penetration. Those who reported a history of sexual intercourse responded to further questions, including the age of first sexual intercourse, condom use at the first sexual encounter, age of the partner, number of lifetime sexual partners, and sexual activity in the previous 12 months. Those who were sexually active in the previous 12 months were also asked questions regarding the use of condoms at the last sexual encounter and the frequency of condom use with both commercial and noncommercial partners. A commercial partner was defined as a partner with whom a respondent had sex in exchange for money; any partner other than this was defined as a noncommercial partner.

The section on HIV/AIDS knowledge contained 11 items, including modes of HIV transmission and measures for preventing sexual transmission. Items on the exposure to HIV/AIDS prevention programs had questions on exposure to posters on HIV/AIDS and knowledge of government programs and NGOs.

Instrument validation

To examine the instrument's reliability, 45 items extracted from the original survey were retested 1 week later for 48 randomly selected respondents. Overall, a high test-retest correlation was observed between items (Pearson $r = 0.85$ – 1.00 [median = 0.96] for continuous variables, kappa = 0.26– 1.00 [median = 0.71] for categorical variables) [17]. The internal consistency was satisfactory ($\alpha = 0.68$) for 11 items of knowledge. In addition, a random response technique was administered to all subjects to assess the validity of the interview-based response on sex-

ual experience in a subsequent procedure [18-20]; at the end of interview, respondents were asked to answer one of two "yes/no" questions, (1) "Is your birthday on the 1st to 10th of the month?" or (2) "Have you ever had sexual intercourse?" by randomly picking up a piece of paper that contained one of these questions. The interviewer did not know which question they answered. Under this less intimidating circumstance, 28.4% male and 7.4% female were estimated to have answered "yes" to question 2. These prevalence rates were comparable to those obtained in the face-to-face interviews (30.9% for men and 9.7% for women).

Data collection

A team of 50 trained fieldworkers under the direct supervision of five superiors conducted the fieldwork. Data were collected over a 3-week period in January and February 2003. Fieldworkers visited the homes of the respondents and interviewed them in their native language, Creole, using the structured questionnaire. Due to the sensitive nature of some questions, the gender of the interviewers was matched to that of the respondents. The interviews took 20-30 min to complete on average. The decision to use face-to-face interviews was made by taking into account the local language environment of Mauritius where the first language (Creole) is a highly conversation-based language, and French or English is not understood by people with a low educational attainment.

Statistical analysis

Statistical analyses were performed using SPSS Complex Samples 12.0J (SPSS Japan, Tokyo, Japan) and SUDAAN 8.0.2 (Research Triangle Institute, Research Triangle Park, NC, USA) to account for the clustered sampling design. First, we assessed the prevalence of HIV-related sexual behaviors. Subsequently, using "being sexually experienced" and "condom nonuse at last sex" as the outcome variables, bivariate associations with dichotomous correlates were analyzed using cross-tabulation. Finally, variables that achieved a statistically significant association ($P < 0.10$) in the bivariate analyses and variables that were epidemiologically important were entered into logistic regression models to obtain an adjusted odds ratio for each association.

Ethical issues

The study protocol was reviewed and approved by the Kyoto University Graduate School and Faculty of Medicine Ethics Committee and Mauritius Ministry of Health and Quality of Life. Before each interview, the interviewer explained the objectives and methodology of the study to each participant; the interviewer signed the survey document when verbal informed consent was obtained from the participant.

Results

Description of the sample

The mean age for men was 19.2 years ($SE = 0.14$) and that for women was 18.6 years ($SE = 0.11$). The median ages were 19 years for men and 18 years for women, respectively. Religious affiliations comprised approximately 55% Hindu, 20% Muslim, and 25% Christian. Nearly half of the participants were still attending school and lived in urban areas at the time of the survey. For men, 45.8% were employed full time, whereas 57.4% of women had not worked in the previous 12 months (Table 1).

Sexual behavior

The proportion of people who had ever had penetrative sexual intercourse differed significantly between men (30.9%; 95% CI, 25.8-36.5) and women (9.7%; 95% CI, 6.5-14.0; $P = 0.001$). The mean age of first sex among those sexually experienced was 17.4 years for males (median = 17) and 17.8 years for females (median = 18). The mean age of the first partner was 21.4 years for females (median = 21), indicating that women tend to have an older first partner. Significantly more men (65.0%) than women (32.0%) reported having had more than one sexual partner ($P = 0.001$). Condom use during sexual intercourse was low for both men and women, and only 35.7% of men and 24.5% of women reported having used a condom consistently with their noncommercial sex partner during the 12 months before completing the survey (Table 2). Only 23 male respondents reported sex with a commercial partner.

Correlates of sexual experience

The correlates of sexual experience in terms of sociodemographic characteristics and social life indicators such as exposure to nightclubs and drug use were analyzed by gender (Table 3). Logistic regression results indicate that for both sexes, non-Christians (i.e., Hindus and Muslims) had a significantly lower adjusted odds ratio (AOR) of being sexually experienced (AOR = 0.27 for men and 0.15 for women). Men who had ever worked, ever visited nightclubs, or had a history of using marijuana had a significantly higher likelihood of being sexually experienced (AOR = 3.14, 3.07, and 3.09, respectively). Although not statistically significant in the logistic regressions, most social life factors among men, i.e., having a mobile phone, a history of watching pornographic films, and drinking alcohol, were associated with being sexually experienced in bivariate analyses. Women who were not in school, had ever visited nightclubs, or reported a history of alcohol use in the 12 months before the survey had a significantly higher likelihood of being sexually experienced in logistic regression analyses (AOR = 4.28, 3.14, and 2.81, respectively).

Table 1: Demographic characteristics of participants by gender

	Male (n = 575) %	Female (n = 566) %
Age group		
15-19	53.1	63.3
20-24	46.9	36.7
Mean age	19.2 (SE 0.14)	18.6 (SE 0.11)
Religion		
Hindu	55.7	52.6
Muslim	20.5	20.6
Christian	22.8	25.8
Other/No religion	1.0	1.0
Education status		
Currently attending school	41.2	48.5
Completed > 3 years in secondary	39.6	38.2
Completed ≤ 3 years in secondary	19.2	13.0
Never attended school	0	0.3
Geographic region of residence		
Urban	42.8	42.9
Rural	57.2	57.1
Work experience in the previous 12 months		
Full time	45.8	29.9
Part time	19.5	12.7
Did not work	34.7	57.4
Monthly personal expenditure (in Rupees) ^a		
High	65.4	48.7
Low	34.6	51.3
Mean expenditure	1662.5 (SE 75.68)	1307.8 (SE 93.62)

SE, standard error.

a. Median split of total sample spending pattern: "Low," < 1000; "High," ≥ 1000; n = 574 for males and n = 561 for females.

Correlates of condom nonuse at the last sexual encounter

Of those who had sex with noncommercial partners in the last 12 months (n = 123 for men and n = 40 for women), more than 50% of men and 70% of women did not use a condom at the last sexual encounter. The correlates of condom nonuse at the last sexual encounter were analyzed in terms of sociodemographic characteristics, condom attitudes, sexual behaviors, HIV/AIDS knowledge,

and exposure to an HIV/AIDS program (Table 4). Male and female data were combined for condom nonuse analyses because prior bivariate analyses stratified by gender indicated that the direction and magnitude of all independent variables were similar for men and women. In the logistic regression analyses, condom nonuse at the first sexual encounter was found to be the most significant correlate of nonuse at the last sexual encounter (AOR =

Table 2: Prevalence of sexual behavior by gender

	Male % (95% CI)	Female % (95% CI)	P ^d (χ ²) ^e
Ever had sex ^a	30.9(25.8-36.5)	9.7(6.5-14.0)	0.001 (51.53***)
Mean age at first sexual encounter ^b	17.4 (SE 0.20)	17.8 (SE 0.33)	0.362
Mean age of first partner ^b	18.1 (SE 0.25)	21.4 (SE 0.58)	0.001
Used condom at first sexual encounter ^b	41.1 (31.9-51.1)	32.0 (18.7-49.0)	0.277 (1.21)
Had multiple sexual partners ^b	65.0 (56.8-72.5)	32.0 (21.5-44.8)	0.001 (14.83***)
Had sex in the past 12 months ^b	71.3 (63.5-78.1)	75.1 (57.4-87.1)	0.688 (0.16)
Used a condom at last sexual encounter ^c	49.4 (38.2-60.6)	28.8 (12.1-54.2)	0.079 (3.23)
Always used a condom ^c	35.7 (25.6-47.4)	24.5 (10.1-48.4)	0.274 (1.23)

CI, confidence interval; SE, standard error.

a. Among total respondents (n = 575 for males and n = 566 for females).

b. Among people who had ever had sex (n = 183 for males and n = 55 for females).

c. Among people who had sex with noncommercial sex partners in the past 12 months (n = 123 for males and n = 40 for females).

d. Chi-square test for the percentage and t-test for the mean.

e. **P < 0.01.

Table 4: Bivariate and multivariate association between dichotomous correlates and engaging in sex without a condom at the last sexual encounter with a noncommercial partner

Correlate			Bivariate		Multivariate *		n = 161
	n	Percent reporting condom nonuse at last sexual encounter	Prevalence ratio (95% CI)	P (χ^2) ^b	Adjusted odds ratio (95% CI)	P	
Sociodemographic							
Gender							
Female	40	71.2	1.41	0.079	3.09	0.056	
Male	123	50.6	(0.94–2.11)	(3.23)	(0.97–9.85)		
Age group (years)							
Older (20–24)	104	61.6	1.45	0.034	2.68	0.051	
Younger (15–19)	59	42.5	(1.00–2.11)	(4.78 ^{**})	(1.00–7.23)		
Geographic area							
Urban	66	45.0	0.73	0.065	0.95	0.893	
Rural	97	62.0	(0.53–0.99)	(3.56)	(0.43–2.11)		
Currently in school							
No	127	58.1	1.42	0.051	0.99	0.984	
Yes	36	40.9	(0.97–2.09)	(4.02)	(0.27–3.56)		
Work experience							
Ever worked	132	55.8	1.16	0.520	N/A	N/A	
Never worked	31	48.2	(0.74–1.81)	(0.42)			
Condom attitudes							
Perceived difficulty in obtaining a condom							
High	23	80.4	1.58	0.017	4.17	0.084	
Low	138	50.9	(1.18–2.12)	(6.17 ^{**})	(0.82–21.14)		
Self-efficacy of using a condom with a new partner							
Low	28	88.5	1.83	0.001	3.11	0.066	
High	133	48.3	(1.46–2.30)	(27.38 ^{***})	(0.93–10.41)		
Confidence to refuse sex without condom							
Low	86	60.4	1.28	0.111	N/A	N/A	
High	75	47.2	(0.95–1.72)	(2.64)			
Sexual behaviors							
Used a condom at first sexual encounter							
No	103	78.3	3.75	0.001	12.92	0.001	
Yes	60	20.9	(2.34–6.01)	(38.64 ^{***})	(4.96–33.62)		
Had more than one sexual partner							
Yes	111	54.4	0.99	0.958	N/A	N/A	
No	52	54.9	(0.74–1.34)	(0.00)			
Knowledge of HIV prevention methods							
Effectiveness of condom in HIV prevention							
Low	20	79.1	1.54	0.029	1.94	0.398	
High	143	51.4	(1.13–2.10)	(5.06 ^{**})	(0.41–9.19)		
Effectiveness of having one faithful partner in HIV prevention							
Low	60	58.7	1.13	0.454	N/A	N/A	
High	103	52.0	(0.83–1.54)	(0.57)			
Exposure to HIV program							
Ever heard of PILS (NGO)							

Table 4: Bivariate and multivariate association between dichotomous correlates and engaging in sex without a condom at the last sexual encounter with a noncommercial partner (Continued)

No	66	66.0	1.42	0.018	3.27	0.010
Yes	97	46.5	(1.08–1.86)	(5.98*)	(1.35–7.90)	

NI/A, not applicable; 95% CI, 95% confidence interval; NGO, nongovernmental organization.

a. Three basic sociodemographic variables (gender, age group, and geographic area) and variables that attained $P < 0.10$ in the bivariate analyses were entered into the logistic regression model.

b. * $P < 0.05$. ** $P < 0.01$.

12.92). Moreover, those who had not been exposed to HIV prevention programs by NGOs that promoted HIV prevention among young people had a significantly higher AOR (3.27). Those who were female and older had higher likelihoods (AOR = 3.09 and 2.68, respectively) of not using a condom at the last sexual encounter, although these factors were not statistically significant in logistic regression analyses.

Discussion

We described the results of a survey on HIV-related sexual behaviors among a population-based probability sample of young people on the island of Mauritius. Among unmarried young people aged 15–24 years, 30.9% of males and 9.7% of females reported a history of penetrative sexual intercourse. Of these sexually active people, behaviors that increase vulnerability to HIV infection were prevalent, e.g., having multiple partners (males), having sex without a condom, and having an older partner (females) [21–23].

The proportion of sexually active unmarried youth in Mauritius, i.e., 30.9% of men and 9.7% of women, appears to be relatively low compared to that in other African countries. Data from 17 national surveys conducted in the past 5 years among young people aged 15–24 years from diverse countries in sub-Saharan Africa indicate that, on average, about 38% of men and 30% of women report having had premarital sex in the previous year [24]. As prevalence of ever having had sex in these countries should be higher than that of sex in previous year, we can say that sexual activity of Mauritian youth is comparatively much lower than that of other African countries. In addition, when compared with data from the 1996 Mauritius Youth profile study, we did not observe a dramatic change in the proportion of sexually experienced unmarried youth. This relatively low prevalence of sexually active youth may be due at least in part to the predominance of religions with conservative sex norms and may partly explain how Mauritius has managed to maintain a stable low level of HIV infection in the 1990s, despite the devastating HIV epidemic in neighboring Africa.

Nevertheless, these findings do not leave room for complacency. Among sexually experienced Mauritian youths, 50.6% of male and 71.2% of female respondents did not use condoms at the last sexual encounter with a noncom-

mercial partner. Although HIV infection risks may not be very high for young people as a whole, there certainly exist a group of youth who have unprotected sex and are vulnerable to HIV infection. Therefore, the HIV program for youth in Mauritius should aim to maintain the low prevalence of sexually active youth and to promote safe sexual behavior among those who are sexually active.

To achieve these goals, we need to identify the predictors of sexual behavior to effectively and efficiently target HIV prevention to this vulnerable subpopulation. We identified several factors related to the sexual behaviors of young people in Mauritius. For men, having ever worked and marijuana use were significantly associated with having ever had sex. In contrast, being out of school and drinking were associated with women's sexual experience. In addition, being Christian and visiting nightclubs were significantly associated with sexual experience in both sexes. Furthermore, among sexually experienced young people, the use of condom at first sex and exposure to an NGO's HIV program were significantly related to condom use at last sexual encounter. These findings suggest possible directions for effective youth HIV prevention programs in Mauritius. We grouped them into three possible approaches: (1) out-of-school and in-school approaches, (2) nightclub and popular site approaches, and (3) religion and social status approaches.

Out-of-school and in-school approaches

Prevention efforts are needed outside the school system because the young people that were more sexually experienced were men who have ever worked or women who are not in school. Two channels currently exist in Mauritius to reach young people outside the school system: (1) peer educators trained by the AIDS unit of the Ministry of Health and (2) youth leaders attached to the youth centers of the Ministry of Youth and Sports. These resources should be reinforced to reach out-of-school youth to provide better reproductive health services. Furthermore, one of our most important findings was that exposure to an NGO program was associated with safer sexual behavior in young people. Because these activities are easily modifiable, a great opportunity may exist for the promotion of safer sex by reinforcing the activities of NGOs through peer educators or youth leaders.

At the same time, to maintain or reduce the number of out-of-school youth, enrollment in secondary and tertiary education should be promoted in a long-term perspective. In Mauritius, the enrollment ratio for primary school is 100%, but falls to 60% for secondary school, and 4.4% for post-secondary school [25]; there is therefore room to promote secondary and tertiary education. By doing so, young people may delay their sexual debut by having greater access to information and skills, enhanced future orientation and aspirations, and by delaying exposure to the workplace where they encounter older people.

Reproductive health education needs to be promoted in schools because it is important for all young people, including the sexually inexperienced, to have the requisite knowledge and skills to use condoms beginning with their first sexual encounter. Our results indicate that condom nonuse at the last sexual encounter is significantly associated with condom nonuse at the first sexual encounter, i.e., those who used a condom at their first sexual encounter were much more likely to report using a condom at their most recent sexual encounter. The same trends were found among adolescents from other parts of the world [26,27]. These results suggest that HIV prevention programs emphasizing appropriate and consistent condom use should target young people before the onset of sexual activity. In schools, where most students are sexually inexperienced, information on the risks accompanying sexual behavior should be provided to all students so that they can make informed future decisions.

Nightclub and popular site approaches

Nightclub visitation was correlated with other variables assessing popular sensationseeking behavior, such as possession of a mobile phone, watching pornographic films, alcohol intake, and marijuana use, in bivariate analysis. Although only nightclub visitation was significantly associated with sexual behavior in both men and women in multivariate analyses, the effects of these other items should be considered in the HIV prevention program. Media campaigns using various channels are potentially one of the most effective approaches to convey HIV prevention messages to these young people who seek out and are exposed to risks. As suggested in previous studies, role model stories printed in youth-attracting flyers may be a useful means to convey HIV prevention messages, even to hard-to-reach youths, because they can readily be distributed at popular sites where young people congregate [28,29].

In addition, HIV prevention interventions, including education on the risks of drinking and drug use, could be explored in nightclubs because we identified nightclubs as an important place where sexually experienced young men and women congregate. The identification of specific

access points will help the national AIDS control program to effectively use available resources for targeted interventions. Ford and Inman [30] documented successful interventions to promote "safer sex" among young people in nightclubs in southwest England. The event was organized in a way that appeals to young people's sense of humor: competitions run by disc jockeys, free condoms, and "safer sex" materials. Although gaining the cooperation of venue owners may be challenging, such events could be considered in nightclubs in Mauritius to effectively reach young people with high risk factors for contracting HIV.

The effect of watching pornographic films should also be cautiously assessed because the bivariate analysis, but not logistic regression, showed this to be significantly associated with sexual experiences in young men. In fact, 70.6% of men, as compared to 12.9% of women, have reported ever having watched pornographic films. These films, most of which are from overseas, include unprotected sexual behaviors. A recent cohort study in the United States indicated that exposure to sexual content in the media accelerated adolescents' sexual activity and increased their risk of engaging in early sexual intercourse [31]. Similarly, a study in China reported that younger students are more exposed to pornographic media than are older students [32]. Taking these findings and the recent rapid increase in Internet use in Mauritius into account, attention should be given to trends related to the excessive exposure of young men to sexual media content. Video rental stores, of which there are many on the island, are potential intervention sites to address the watchers of pornographic films.

Religion and social status approaches

Christian youth reported more sexual experience than those in other religious groups. This may be partly related to their higher exposure to nightclubs, alcohol, and marijuana use because a relatively higher proportion of Christians live in urban areas compared to Muslims or Hindus. In this context, the popular site approaches (above) should be implemented, focusing on Christian youth.

Given disproportionate differences in sexual experience rates among religious groups, religious gatherings should also be considered as potential places for HIV prevention sessions to address sexual norms and reproductive health [33]. Religious groups in Mauritius are still unwilling to promote condoms as a means of protection, and would rather stress abstinence and fidelity. Some religious people, however, are more involved in the fight against poverty, the commercial sex trade, and substance abuse, and through these related issues, they also deal with HIV/AIDS. Greater participation of religious leaders in HIV/AIDS programs may be facilitated through greater involve-

ment of people living with HIV/AIDS (PLWHA), as suggested by work in Trinidad, where the religious profile is as diverse as in Mauritius [34]. In Trinidad, personal interaction with PLWHA was positively associated with the involvement of religious leaders in HIV/AIDS initiatives.

Finally, it is needless to say that National AIDS Control program of Mauritius should swiftly act on HIV prevention for injecting drug users given the contemporary trends of HIV spread in this group. At the same time, with regard to young people, our findings suggest that early and focused programs are needed. Evidence from Cambodia indicates that young people are more willing to adopt safer behaviors than older people when provided with the appropriate services [35]. Focused programs specifically targeting young people in general, as well as at-risk populations, such as sex workers and drug users, will have maximum impact and require the least resources when implemented at this early stage of the HIV epidemic. In particular, young women warrant special attention because the results indicate that women tended to have older sexual partners, which means that they may be more exposed to HIV than men [22]. In Africa, young people, especially younger women, are the most seriously affected by HIV because of their social and biological vulnerability [36]. Precautions are needed to protect younger women in Mauritius to avoid a situation similar to that on the African continent. The National AIDS Control Program of Mauritius should take every opportunity to intervene before the HIV epidemic escalates.

Our study is not without limitations, which include face-to-face interviews as the only modality of data collection. Although under-reporting might not have been completely avoided, the validity of the responses concerning sexual experience was assessed using a random response technique and considerable effort was made to put the respondents at ease, particularly for sensitive questions. However, self-report information may be subject to reporting errors and biases. Furthermore, the study was based on cross-sectional data for which the direction of causal relationships cannot be determined. Future studies need to assess the effectiveness of HIV prevention interventions in the context of young people's lives in Mauritius.

Conclusion

Premarital sexual intercourse was reported by 30.9% of male and 9.7% of female young people in Mauritius. Of these sexually active young people, behaviors that increase the vulnerability to HIV infection were prevalent and were associated not only with their culture, which was represented by religion, but also with the influence of a changing society such as school enrolment, NGOs, night-clubs, and pornographic films. Taking these predictors of

sexual behaviors into account, a focused HIV prevention program for young people needs to be reinforced in Mauritius to minimize the future consequences of the epidemic, even though HIV levels are currently low.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

YHN, MOK, and MK conceived the study, performed and interpreted the analyses, and wrote the first draft of the manuscript. JCM and RN supervised and coordinated the data collection in the field. TH, RJD, and DLL provided advice on the statistical data analysis and interpretation of results. All authors read and approved the final manuscript.

Acknowledgements

We are grateful to all the respondents who participated in our study. We thank the study team and all the staff and colleagues of the Mauritius Institute of Health, Ministry of Health and Quality of Life, Ministry of Youth and Sports, WHO, PILS, MFPA, and NATReSA of Mauritius for their assistance and input. We also thank the researchers and the colleagues from the Kyoto University School of Public Health and the Rollins School of Public Health of Emory University for their comments and advice. This study was supported by the associate guest researcher program (In fiscal year 2002) of the Japan International Cooperation Agency. A scholarship from the Ishizaka Foundation (fiscal year 2003) allowed the first author to study in the United States during the data analysis stage of the study.

References

- Matteelli A, Pizzocolo C, Bakar HA, Toyb M, Pyndiah N, Udonwa NE, Gedeon AG, Farina C, Caligaris S, Castelli F, Carosi G: **STI epidemics in the Indian Ocean region: can the phase be assessed?** *Sex Transm Infect* 2002, **78** Suppl 1:i121-4.
- Andriamahena R, Raveloajona B, Rarivoarilala E, Ravaoarimalala C, Andriamadiana J, Andriamahafazy B, May JF, Behets F, Rasamindrakotroka A: **[AIDS in Madagascar. I. Epidemiology, projections, socioeconomic impact, interventions].** *Bull Soc Pathol Exot* 1998, **91**(1):68-70.
- Ministry of Health and Quality of Life: **National HIV/AIDS Strategic Plan 2001-2005, Mauritius.** Mauritius : 2001.
- Bellis MA, Hughes K, Thomson R, Bennett A: **Sexual behaviour of young people in international tourist resorts.** *Sex Transm Infect* 2004, **80**(1):43-47.
- Schensul SL, Schensul JJ, Oodit G, Bhowan U, Ragobur S: **Sexual intimacy and changing lifestyles in an era of AIDS: Young women workers in Mauritius.** *Reproductive Health Matters* 1994, **3**:83-93.
- Mauritius Institute of Health: **Study on youth profile in the republic of Mauritius.** Mauritius : 1996.
- Mauritius Institute of Health: **Report on knowledge, attitudes, beliefs and practices related to HIV/AIDS in republic of Mauritius.** Mauritius : 1998.
- Ministry of Health, University of Mauritius, WHO: **Knowledge attitudes beliefs practices related to AIDS, KABP survey 1989 Mauritius.** Mauritius : 1989.
- Ministry of Health, University of Mauritius, WHO, UNICEF: **Rodrigues STD/AIDS Knowledge attitudes beliefs practices: a qualitative KABP study.** Mauritius : 1992.
- Crosby RA, Holtgrave DR, DiClemente RJ, Wingood GM, Gayle JA: **Social capital as a predictor of adolescents' sexual risk behavior: a state-level exploratory study.** *AIDS Behav* 2003, **7**(3):245-252.

11. DiClemente RJ, Wingood GM, Crosby R, Cobb BK, Harrington K, Davies SL: **Parent-adolescent communication and sexual risk behaviors among African American adolescent females.** *Journal of Pediatrics* 2001, **139**(3):407-412.
12. Ono-Kihara M, Kihara M, Yamazaki H: **Sexual practice and the risk for HIV/STDs infection of youth in Japan.** *Japan Medical Association Journal* 2002, **45**(12):520-525.
13. Central Statistics Office, Ministry of Economic Development Financial Services and Corporate Affairs: **2000 Housing and population census Republic of Mauritius, Volume II: Demographic and fertility characteristics.** Mauritius ; 2001.
14. Bennett S, Woods T, Lyanage WM, Smith DL: **A simplified general method for cluster-sample surveys of health in developing countries.** *World Health Statistics Quarterly* 1991, **44**:98-106.
15. Family Health International: **Behavioral Surveillance Surveys BSS Guidelines for repeated behavioral surveys in populations at risk of HIV.** Arlington ; 2000.
16. UNAIDS: **National AIDS programmes, A guide to monitoring and evaluation.** Geneva ; 2000.
17. Brener ND, Kann L, McManus T, Kinchen SA, Sundberg EC, Ross JG: **Reliability of 1999 Youth Risk Behavior Survey Questionnaire.** *Journal of Adolescent Health* 2002, **31**:336-342.
18. Dare OO, Cleland JG: **Reliability and validity of survey data on sexual behaviour.** *Health Transit Rev* 1994, **4** Suppl:93-110.
19. Fisher M, Kupferman LB, Lesser M: **Substance Use in a School-Based Clinic Population: Use of the Randomized Response Technique to Estimate Prevalence.** *Journal of Adolescent Health* 1992, **13**:281-285.
20. Kolata G: **How to ask about sex and get honest answers.** *Science* 1987, **236**(4800):382.
21. Ford K, Lepkowski JM: **Characteristics of sexual partners and STD infection among American adolescents.** *Int J STD AIDS* 2004, **15**(4):260-265.
22. Miller KS, Clark LF, Moore JS: **Sexual initiation with older male partners and subsequent HIV risk behavior among female adolescents.** *Fam Plann Perspect* 1997, **29**(5):212-214.
23. Vanoss Marlin B, Coyle KK, Gomez CA, Carvajal SC, Kirby DB: **Older boyfriends and girlfriends increase risk of sexual initiation in young adolescents.** *J Adolesc Health* 2000, **27**(6):409-418.
24. USAID, UNICEF, CDC, UNAIDS, WHO, US Census Bureau, Family Health International, MEASURE Evaluation, The Synergy Project, MEASURE DHS+: **HIV/AIDS Survey Indicators Database.** 2006 [<http://www.measuredhs.com/hivdata/>].
25. Central statistics office, Ministry of Economic Development Financial Services and cooperate Affairs: **Mauritius in figures 2000.** 2000.
26. St.Lawrence JS: **African-American Adolescents' Knowledge, Health-Related Attitudes, Sexual Behavior, and Contraceptive Decisions: Implications for the Prevention of Adolescent HIV Infection.** *Journal of Consulting and Clinical Psychology* 1993, **61**(1):104-112.
27. Svare EI, Kjaer SK, Poll P, Bock JE: **Determinants for contraceptive use in young, single, Danish women from the general population.** *Contraception* 1997, **55**(5):287-294.
28. Bond L, Bowden-Proctor J, Lauby J, Walls C, Woll M: **Developing non-traditional print media for HIV prevention: role model stories for young urban women.** *Am J Public Health* 1997, **87**(2):289-290.
29. Corby NH, Enguidanos SM, Kay LS: **Development and use of role model stories in a community level HIV risk reduction intervention.** *Public Health Rep* 1996, **111** Suppl 1:54-58.
30. Ford N, Inman M: **Safer sex in tourist resorts.** *World Health Forum* 1992, **13**(1):77-80.
31. Brown JD, L'Engle KL, Pardun CJ, Guo G, Kenneavy K, Jackson C: **Sexy media matter: exposure to sexual content in music, movies, television, and magazines predicts black and white adolescents' sexual behavior.** *Pediatrics* 2006, **117**(4):1018-1027.
32. Ma Q, Ono-Kihara M, Cong L, Xu G, Zamanli S, Ravari SM, Kihara M: **Sexual behavior and awareness of Chinese university students in transition with implied risk of sexually transmitted diseases and HIV infection: a cross-sectional study.** *BMC Public Health* 2006, **6**:232.
33. Agadjanian V: **Gender, religious involvement, and HIV/AIDS prevention in Mozambique.** *Soc Sci Med* 2005, **61**(7):1529-1539.
34. Genrich GL, Brathwaite BA: **Response of religious groups to HIV/AIDS as a sexually transmitted infection in Trinidad.** *BMC Public Health* 2005, **5**:121.
35. MAP Monitoring the AIDS pandemic: **AIDS in Asia: Face and Facts: A comprehensive analysis of the AIDS epidemics in Asia.** 2004.
36. UNAIDS, WHO: **AIDS Epidemic Update.** Geneva ; 2004.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1472-698X/7/8/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime.

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

