


- ³⁰ Kankasa, C et al. Routine and Universal Counseling and Testing Among Hospitalized Children at University Teaching Hospital, Lusaka, Zambia. PEPFAR Implementers Meeting, Durban, South Africa, June 12-15, 2006 (Abstract 215).
- ³¹ Corneli A et al. Patient and Provider Perspectives on Improving Access to HIV Testing and Counselling for TB Patient in Kinshasa, Democratic Republic of Congo (DRC). 3rd IAS Conference on HIV Pathogenesis and Treatment, Rio De Janeiro, July 24-27, 2005 (Abstract No.TuPe7.1C19).
- ³² Nawavvu C et al. Routine HIV Testing for Children: Challenges and Lessons Learned. PEPFAR Implementers Meeting, Durban, South Africa, June 12-15, 2006 (Abstract 223).
- ³³ Homsy J et al. Routine Intrapartum HIV Counseling and Testing for Prevention of Mother-to-Child Transmission of HIV in a Rural Ugandan Hospital. *Journal of acquired immune deficiency syndromes*. June 2006, 42:149-154.
- ³⁴ Semafumu E, Ngabirano T. Building on the Success of STD Programmes to Increase Access to Prevention and Care for HIV/AIDS: The Case for Routine Testing and Counseling in STD Patients. PEPFAR Implementers Meeting, Durban, South Africa, June 12-15 2006 (Abstract 69).
- ³⁵ Andia I. Evolving Clinical Picture Secondary to Routine HIV Testing and Early Linkage to Care at the HIV Clinic at Mbarara Regional Referral Hospital. PEPFAR Implementers Meeting, Durban, South Africa, June 12-15, 2006 (Abstract 195).
- ³⁶ Steen TW et al. Two and a Half Years of Routine HIV Testing in Botswana. *Journal of acquired immune deficiency syndromes*. 2007, 44(4):484-8.
- ³⁷ Op. cit number 24
- ³⁸ Greenwald JL. Routine Rapid HIV Testing in Hospitals: Another Opportunity for Hospitalists to Improve Care. *Journal of Hospital Medicine*. 2006,1(2): 106 -112.
- ³⁹ OP. cit. number 29
- ⁴⁰ Medley A et al. Rates, barriers and outcomes of HIV serostatus disclosure among women in developing countries: implications for prevention of mother-to-child transmission programmes. Geneva: *Bulletin of the World Health Organization*. 2004, 82(4): 299-307.
- ⁴¹ Maman S et al. High rates and positive outcomes of HIV-serostatus disclosure to sexual partners: Reasons for cautious optimism from a voluntary counseling and testing clinic in Dar es Salaam, Tanzania. *AIDS and behaviour*. 2003,7(4):373-82.
- ⁴² Gielen AC et al. Women's lives after an HIV-positive diagnosis: disclosure and violence. *Maternal and Child Health Journal*, 2000, 4(2):111-20.
- ⁴³ Gaillard P et al. Vulnerability of women in an African setting: Lessons for mother-to-child HIV transmission prevention programmes. *AIDS*. 2002,16(6):937-9.

- ⁴⁴ Semraua K et al. Women in couples antenatal HIV counseling and testing are not more likely to report adverse social events. *AIDS*, 2005, 19:603–609.
- ⁴⁵ Passin WF et al. A systematic review of HIV partner counseling and referral services: client and provider attitudes, preferences, practices, and experiences. *Sexual Transmitted Disease*. 2006, 33(2):1-9.
- ⁴⁶ Op. cit. number 39
- ⁴⁷ USAID/Synergy. *Women's Experiences with HIV Serodisclosure in Africa: Implications for VCT and PMTCT*. Meeting Report. Washington DC: USAID, March 2004.
- ⁴⁸ World Health Organization Regional Office for South-East Asia. *Voluntary HIV counselling and testing: manual for training of trainers parts 1 & 2*. 2004.
- ⁴⁹ World Health Organization Regional Office for Africa. *Regional HIV/AIDS Voluntary Counselling and Testing Guidelines*. Brazzaville, 2005
- ⁵⁰ WHO/UNAIDS. Provider-Initiated HIV Testing and Counselling in Clinical Settings: Operational Recommendations. A Meeting Report. WHO/UNAIDS consultation meeting, 3-4 July 2006
- ⁵¹ WHO. *The Right to Know: New Approaches to HIV Testing and Counselling*. Geneva: World Health Organization, 2003 (WHO/HIV/2003.08).
- ⁵² Op. cit. number 4
- ⁵³ Some of these terms were proposed in earlier drafts of this document, and the term "routine offer" was used in the WHO/UNAIDS Policy Statement on HIV Testing and Counselling. The policy Statement will be updated to reflect the terminology used in this document.
- ⁵⁴ A fourth epidemic scenario, hyperendemic epidemic, has been proposed for HIV programme planning purposes in countries with HIV prevalence greater than 15%. The recommendations made for generalized epidemics in this document would also apply to hyperendemic epidemics. See: *Practical guidelines for intensifying HIV prevention: towards universal access*. UNAIDS. 2007.
- ⁵⁵ WHO. *Case Definitions of HIV for Surveillance and Revised Clinical Staging and Immunological Classification of HIV-Related Disease in Adults and Children*. Geneva: WHO, 2006
- ⁵⁶ Horwood C., Liebeschutz S., Blauuw D., Cassol S. And Qazi S. Diagnosis of pediatric HIV infection in a primary health care setting with a clinical algorithm. *Bulletin of the World Health Organization* .2003, 81 (12).
- ⁵⁷ WHO/UNAIDS. *Technical Consultation on Male Circumcision and HIV Prevention: Research Implications for Policy and Programming. Conclusions and Recommendations*. World Health Organization and UNAIDS. Geneva: 28 March 2007.

- ⁵⁸ Bulterys M. et al. Rapid HIV testing during labour: a multicenter study. *JAMA*. 2004, 292:219-223
- ⁵⁹ WHO. *Antiretroviral drugs for treating pregnant women and preventing HIV infection in infants in resource-limited settings: towards Universal Access. Recommendations for a public health approach*. Geneva: WHO, 2006.
- ⁶⁰ For more information see: *Toolkit for Targeted HIV/AIDS Prevention and Care in Sex Work Settings*, WHO 2005; *Policy and Programming Guide for HIV/AIDS Prevention and Care Among Injecting Drug Users*, WHO 2005; *Advocacy guide: HIV/AIDS prevention among injecting users: workshop manual*. WHO/UNAIDS 2004
- ⁶¹ M-L. Newell et al. Mortality of infected and uninfected infants born to HIV-infected mothers in Africa: a pooled analysis. *Lancet*. Volume 364, Issue 9441, 2 October 2004-8 October 2004, Pages 1236-1243
- ⁶² Lewis DK, Callaghan M, Phiri K, et al. Prevalence and indicators of HIV and AIDS among adults admitted to medical and surgical wards in Blantyre, Malawi. *Trans R Soc Trop Med Hyg* 2003; 97: 91-96.
- ⁶³ Surveillance of HIV among TB patients is a sensitive indicator of the spread of HIV into the general population. Information about HIV prevalence in TB patients is essential to support the scale-up of comprehensive HIV treatment, care and support to HIV-positive TB patients.
- ⁶⁴ Op. cit. number 40
- ⁶⁵ WHO. *Addressing violence against women in the context of HIV testing and counselling - strategies and recommendations*. WHO meeting report January 16-18, 2006. Geneva: (forthcoming)
- ⁶⁶ WHO. *TB/HIV - A Clinical Manual*. Second edition. WHO, 2004. WHO/HTM/TB/2004.329
- ⁶⁷ *International guidelines on HIV/AIDS and human rights*, UNAIDS/OHCHR, 2006 consolidated version
- ⁶⁸ UNAIDS Best Practice Collection, Key Material, UNAIDS/WHO, Geneva, 2000.
- ⁶⁹ Family Health International, *Voluntary counselling and testing: a reference guide – responding to the needs of young people, children, pregnant women and their partners*. FHI, 2001
- ⁷⁰ *Guidelines for Assuring the Accuracy and Reliability of HIV Rapid Testing: Applying a Quality System Approach*. CDC/WHO, 2005



Photograph: Gideon Mendel/The International HIV/AIDS Alliance/Corbis

For more information, contact:

World Health Organization
Department of HIV/AIDS

20, avenue Appia
1211 Geneva 27
Switzerland

E-mail: hiv-aids@who.int

www.who.int/hiv

618

ISBN 978 92 4 159556 8



9 789241 595568



