

Hepatitis C Virus Infection Can Present with Metabolic Disease by Inducing Insulin Resistance

Kazuhiko Koike

Department of Infectious Diseases, Internal Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

Key Words

Diabetes · Hepatitis C virus · Insulin resistance · Insulin receptor substrate · Transgenic mouse

Abstract

Although hepatitis C virus (HCV) targets the liver, it has become increasingly evident that HCV can induce diseases of many organs. Recently, much attention is drawn to metabolic disorders in HCV infection. First, hepatic steatosis and derangement in lipid metabolism have been found characteristic of HCV infection, and later on, a correlation was noted between HCV infection and diabetes as well as insulin resistance. We have demonstrated that HCV by itself can induce insulin resistance through disturbing the insulin signaling pathway by HCV proteins. The fact that HCV infection induces insulin resistance by the virus itself may influence the progression of chronic liver disease and open up novel therapeutic approaches. In conclusion, towards the future, HCV infection needs to be viewed not only as a liver disease but also as a metabolic disease.

Copyright © 2006 S. Karger AG, Basel

Introduction

Hepatitis C virus (HCV) infects approximately 1.8 million people in Japan alone and as many as 200 million over the world and induces liver disease ranging from

chronic hepatitis through cirrhosis to hepatocellular carcinoma (HCC) [1, 2]. It has been noticed soon after the discovery that the infection with HCV does not exclusively involve the liver. In fact, type II cryoglobulinemia [3] and membranoproliferative glomerulonephritis [4] frequently occur in patients infected with HCV. Furthermore, strong associations of HCV infection with Sjogren's syndrome [5] and lichen planus [6] have been noted, which is verified in the animal model [7]. In addition, the relation between HCV infection and B cell lymphoma has attracted attention especially in Europe [8].

Recently, there have been increasing lines of evidence to indicate metabolic disturbances in HCV infection which, in turn, would influence the pathogenesis of chronic hepatitis C. The discovery of HCV in 1989 [9] enabled a comparison between chronic hepatitis C and other chronic liver diseases. As shown in the results, it has been repeatedly reported that steatosis is significantly associated with chronic hepatitis C [10, 11]. Steatosis in HCV infection is reproduced in animal models [12-14] to reinforce a pathologic role of HCV. Furthermore, patients infected with HCV have abnormalities in serum lipids, such as hypocholesterolemia and abnormal levels of apolipoproteins in serum [15, 16]; they are rectified in sustained virological responders to interferon (IFN) [16]. Thus, the association between HCV infection and a derangement in lipid metabolism has become increasingly strong, both in patients and experimental systems in animals. Finally, patients with chronic hepatitis C accompanied by severe steatosis develop hepatic fibrosis with an

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2006 S. Karger AG, Basel
0300-5526/06/0492-0051\$23.50/0

Accessible online at:
www.karger.com/int

Kazuhiko Koike, MD
Department of Infectious Diseases, Internal Medicine
Graduate School of Medicine, University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655 (Japan)
Fax +81 3 5800 8799, E-Mail kkoike-ky@umin.ac.jp

increased velocity [17]. All in all, we could say that abnormal lipid metabolism in HCV infection is deeply involved in the pathogenesis of hepatitis C.

HCV Infection and Diabetes

Diabetes is suggested as another metabolic disease in association with HCV infection. In 1994, Allison et al. [18] reported an epidemiological link between diabetes and HCV infection. However, doubts were cast on the association in view of a decreased glucose tolerance in advanced chronic hepatitis as well as an increased opportunity for HCV infection in diabetics who frequently receive determination of blood sugar. Several reports from the same group and others followed along this line. The trend to accept the solid association between diabetes and HCV infection seems to have been triggered in the United States by the population study by Metha et al. [19].

However, the association between diabetes and HCV infection is blemished by factors responsible for decreased glucose tolerance, such as advanced cirrhosis, obesity and ageing common in patients with hepatitis C; they make it difficult to prove this association. Hence, there is a need to evaluate the association by basic studies in experimental systems.

HCV Infection Induces Insulin Resistance

We set out to demonstrate the association between HCV infection and diabetes using the animal model. Mice transgenic for the HCV core gene were employed to this end [12, 13]. These mice are engineered to have the HCV core gene of genotype 1b in the absence of other viral genes. They express HCV core protein of the expected size in the liver, in levels comparable with those of patients with chronic hepatitis C (fig. 1). Half of them develop HCC later during their lives [13]. These transgenic mice were fed with their normal littermates, and the glucose metabolism was compared between them [20].

Although mice transgenic for the core gene did not develop overt diabetes, they had markedly elevated serum levels of insulin. Plasma glucose levels were somewhat higher in transgenic mice than in their normal littermates, both in the fast and after ample feeding, with no significant differences between them. In remarkable contrast, serum insulin levels were significantly higher in transgenic than in normal mice in both conditions (fig. 2). Since such a combination of normal glucose levels and




Fig. 1. Expression of HCV core gene in transgenic mouse. It carries the core gene of HCV genotype 1b alone and expresses the core protein of expected size in the liver, at levels similar to those in human patients. Mice eventually develop HCC later in their lives.

hyperinsulinemia points to insulin resistance, glucose and insulin tolerance tests were conducted.

Mice transgenic for the HCV core gene exhibited glucose levels a little higher than those in normal littermates, without any significant differences between them. In insulin tolerance tests, glucose levels were significantly higher in transgenic than in normal mice, both 40 and 60 min after they were injected with insulin intraperitoneally (fig. 3). These results indicate suppression of the activity of insulin to decrease blood glucose levels for inducing insulin resistance in core-transgenic mice. Since only the HCV core gene had been incorporated into these transgenic mice, HCV core protein was able to induce insulin resistance *in vivo*.




Fig. 2. Altered homeostasis of glucose in mice transgenic for the HCV core gene. Body weight of 2-month-old mice (a), plasma glucose levels in fasting or fed mice (b) and serum insulin levels in fasting or fed mice (c) are shown. Values represent means \pm SE. NS = Not significant statistically; nTg = nontransgenic mice; Tg = transgenic mice.




Fig. 3. Insulin resistance in transgenic mice. Glucose tolerance in mice after overnight fasting (a). *D*-Glucose (1 g/kg body weight) was given intraperitoneally to conscious mice, and plasma glucose levels were determined at time points indicated. **b** Insulin tolerance in mice fasted overnight. Human insulin (1 U/kg body weight) was injected intraperitoneally, and glucose concentrations were determined sequentially. Values were normalized to the baseline glucose concentration at the time of insulin administration. NS = Not significant statistically; nTg = non-transgenic mice; Tg = transgenic mice.

By what mechanism does insulin resistance arise in this animal model? The insulin resistance is considered to involve two factors, namely central and peripheral insulin resistances (table 1) [21]. The hyperinsulinemic-euglycemic clamp method was employed to differentiate between them. In this method, hepatic glucose production (HGP) is calculated on the basis of amounts of glu-

cose required to keep plasma glucose levels within a certain range at serum insulin levels higher than physiological ones. In normal control mice, HGP was suppressed by 60% by the administration of insulin, in contrast to core-transgenic mice in which there was no appreciable suppression of HGP by insulin (fig. 4). These results indicate a hepatic (central) origin of the insulin resistance

Table 1. Two types of insulin resistance

Type	Mechanism
Peripheral	A shortage of insulin action in the muscle due to deficit in the insulin-induced uptake of glucose into muscles
Central	A shortage of insulin action in the liver due to deficit in the insulin-induced suppression of glucose production in hepatocytes

in transgenic mice. For further confirmation, an uptake of glucose into muscle was determined. There were no differences in the uptake in response to administration of insulin between normal and transgenic mice. Therefore, the insulin resistance in mice transgenic for the HCV core gene is central and hepatic.

HCV Core Protein Suppresses the Transduction of Insulin Signaling in Hepatocytes

Next, we evaluated how insulin resistance elicits in mice transgenic for the HCV core gene. For this purpose, liver homogenate was immunoblotted with antiphosphotyrosine and antiphosphoserine antibodies after insulin receptor substrate (IRS)-1 and IRS-2 had been immunoprecipitated. Tyrosines in IRS-1 were weakly phosphorylated both in normal and transgenic mice before they received insulin, with no differences between them. However, after the administration of insulin, the phosphorylation of tyrosines in IRS-1 increased in normal but not in transgenic mice (fig. 5). Obtained results suggested disturbance in tyrosine phosphorylation as one of the factors responsible for insulin resistance in the liver. There were no differences in phosphorylation of serines in IRS-1 or tyrosines in IRS-2 between normal and transgenic mice. Combined, they provided experimental evidence for the development of insulin resistance by the presence of HCV in the liver that would occur by disturbing the transduction of insulin signaling in hepatocytes (fig. 6).

There remains a possibility for the HCV core protein to directly prohibit phosphorylation of tyrosines, or else, it might inhibit tyrosine phosphorylation via certain cytokines. In our extensive searches for the expression of cytokines in the liver of transgenic mice, only TNF- α and IL-1 β have been found with an increased expression [22]. Therefore, for the purpose of evaluating the role of




Fig. 4. Characterization of glucose metabolism in transgenic mice. Glucose production in the liver was calculated using the hyperinsulinemic-euglycemic clamp method. NS = Not significant statistically; nTg = nontransgenic mice; Tg = transgenic mice.




Fig. 5. Phosphorylation of tyrosine in IRS-1 in response to insulin stimulation. Liver tissues from control mice, transgenic mice with or without anti-TNF- α antibody treatment, were analyzed before and 20 as well as 40 min after administration of insulin. Samples were subjected to immunoprecipitation with anti-IRS-1 antibody and then immunoblotted with indicated antibodies. Experiments were performed in triplicate, and a representative picture is exhibited. Immunoblotting with antiphosphotyrosine (PY) antibody (lane a) did not augment phosphorylation of tyrosine in IRS-1 after stimulation with insulin in the core gene transgenic mice (Tg), in contrast to tyrosine phosphorylation markedly enhanced in control mice (nTg). Insulin-stimulated tyrosine phosphorylation was restored 40 min after treatment with anti-TNF- α antibody. Note differences in the intensity of bands 40 min after the administration of insulin (arrowheads). Immunoblotting with anti-IRS-1 antibody (lane b) served as control for the IRS-1 load.




Fig. 6. A proposed mechanism for insulin resistance in HCV infection. HCV itself or elevated levels of cytokines such as TNF- α may inhibit tyrosine phosphorylation of IRS-1 in the liver, suppress intracellular transduction of insulin signal and lead to insulin resistance. PKC = Protein kinase C; MAPK = mitogen-activated protein kinase.

TNF- α in insulin resistance in transgenic mice, serum insulin was determined and an insulin tolerance test performed after they had received anti-TNF- α intraperitoneally. Pretreatment with anti-TNF- α partially improved insulin resistance in mice transgenic for the HCV core gene. Albeit a direct anti-insulin activity of core protein and direct or indirect factors for insulin resistance are not to be excluded, high levels of TNF- α in the liver would be one of the factors for expression of insulin resistance in this mouse model.

Insulin Resistance in Patients with Chronic Hepatitis C

Concurrently with our report in experimental systems, Aytug et al. [23] investigated insulin signaling in biopsied liver specimens from patients with chronic hepatitis C. Specifically, they evaluated changes in IRS-1, IRS-2 and phosphatidylinositol (PI)3 kinase levels in the liver of patients. With insulin stimulation of biopsied liver samples, insulin receptor proteins and IRS-1 increased, while phosphorylation of tyrosines in IRS-1 decreased to one half of the baseline value, along with a diminished activity for PI3 kinase associated with IRS-1, in patients with chronic hepatitis C. The authors went on to propose a possibility for disturbed transduction of the insulin sig-

nal pathway in the liver to induce insulin resistance in patients with chronic hepatitis C [23]. Their report is quite intriguing in that it opens up the way for evaluating an association between HCV infection and insulin resistance in clinical samples at the molecular level.

The results of Aytug et al. [23] inadvertently coincide with ours in analyzing the mechanism of insulin resistance with the experimental system in mice (*vide supra*). They unanimously incriminate impaired tyrosine phosphorylation in IRS-1 in the induction of insulin resistance by HCV infection. It struck us as a surprise that the mechanism of insulin resistance induced by HCV infection has been in agreement between clinical samples and experimental animals, in spite of hepatic IRS-2 that was preferred to IRS-1 for its role in development of insulin resistance in former studies [24]. HCV infection is peculiar in that IRS-1 weighs heavier than IRS-2 in the induction of hepatic insulin resistance.

Although our data strongly indicate a hepatic character of insulin resistance in HCV infection, they by no means exclude roles of other factors in the induction of this resistance. There is little expression of the HCV core gene in muscles of our animal model; it is not known if HCV infects muscular cells in patients with chronic hepatitis C. Factors not intrinsic to the liver would have to be evaluated to sort this out, including dysfunction of mitochondria for induction of insulin resistance [25].

Insulin Resistance for Advanced Hepatic Fibrosis

Insulin resistance in HCV infection may have an additional significant clinical implication. In 260 patients with chronic hepatitis C, Hui et al. [26] have tried to establish the relationship between liver histology and indicators of glucose metabolism, as well as insulin resistance represented by the homeostasis model assessment of insulin resistance. They have found that insulin resistance already exists in hepatitis C patients with stage 0 or I fibrosis in the liver. This indicates that insulin resistance in HCV infection is not attributable to advanced liver disease. In their study, independent predictors of insulin resistance in HCV infection were body mass index, non-response to antiviral treatment, intensity of portal inflammation and infection with HCV genotype 3 [26]. Furthermore, the homeostasis model assessment of insulin resistance was a significant and independent predictor of the stage and velocity of hepatic fibrosis. The results of the study are of much importance, because they implicate a role of insulin resistance and hyperinsulinemia by inference, in promoting the progression of hepatic fibrosis. Insulin has been proven as an aggravating factor not only in atherosclerosis, but also in systemic inflammation and fibrosis. The liver is no exception to this.

References

- Saito I, Miyamura T, Ohbayashi A, Harada H, Katayama T, Kikuchi S, Watanabe Y, Koi S, Onji M, Ohta Y: Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc Natl Acad Sci USA* 1990;87:6547-6549.
- Simonetti RG, Camma C, Fiorello F, Cottone M, Rapicetta M, Marino L, Fiorentino G, Craxi A, Ciccaglione A, Giuseppetti R: Hepatitis C virus infection as a risk factor for hepatocellular carcinoma in patients with cirrhosis. A case-control study. *Ann Intern Med* 1992; 116:97-102.
- Agnetto V, Chung RT, Kaplan LM: A role for hepatitis C virus infection in type II cryoglobulinemia. *N Engl J Med* 1992;327:1490-1495.
- Johnson RJ, Gretch DR, Yamabe H, Hart J, Bacchi CE, Hartwell P, Couser WG, Corey L, Wener MH, Alpers CE: Membranoproliferative glomerulonephritis associated with hepatitis C virus infection. *N Engl J Med* 1993;328: 465-470.
- Haddad J, Deny P, Munz-Gotheil C, Ambrosini JC, Trinchet JC, Pateron D, Mal F, Callard P, Beaugrand M: Lymphocytic sialadenitis of Sjogren's syndrome associated with chronic hepatitis C virus liver disease. *Lancet* 1992; 339:321-323.
- Pawlotsky JM, Benchiki H, Pellet C, Duval J, Dhumeaux D, Revuz J, Bagot M: Lichen planus and hepatitis C virus (HCV)-related chronic hepatitis: Evaluation of HCV genotypes. *Br J Dermatol* 1995;133:666-667.
- Koike K, Moriya K, Ishibashi K, Yotsuyanagi H, Shintani Y, Fujie H, Kurokawa K, Matsuura Y, Miyamura T: Sialadenitis histologically resembling Sjogren syndrome in mice transgenic for hepatitis C virus envelope genes. *Proc Natl Acad Sci USA* 1997;94:233-236.
- Gisbert JP, Garcia-Buey L, Pajares JM, Moreno-Otero R: Prevalence of hepatitis C virus infection in B-cell non-Hodgkin's lymphoma: Systematic review and meta-analysis. *Gastroenterology* 2003;125:1723-1732.
- Choo QL, Kuo G, Weiner AJ, Overby LR, Bradley DW, Houghton M: Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 1989; 244:359-362.
- Bach N, Thung SN, Schaffner F: The histological features of chronic hepatitis C and autoimmune chronic hepatitis: A comparative analysis. *Hepatology* 1992;15:572-577.
- Lefkowitz JH, Schiff ER, Davis GL, Perrillo RP, Lindsay K, Bodenheimer HC Jr, Balart LA, Ortego TJ, Payne J, Dienstag JL, Gibas A, Jacobson IM, Tamburro CH, Carey W, O'Brien C, Sampliner R, van Thiel DH, Feit D, Albrecht J, Meschivitz C, Sanghvi V, Vaughan RD: Pathological diagnosis of chronic hepatitis C: A multicenter comparative study with chronic hepatitis B. *Gastroenterology* 1993; 104:595-603.
- Moriya K, Yotsuyanagi H, Shintani Y, Fujie H, Ishibashi K, Matsuura Y, Miyamura T, Koike K: Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J Gen Virol* 1997;78:1527-1531.
- Moriya K, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, Ishibashi K, Matsuura Y, Kimura S, Miyamura T, Koike K: The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065-1067.

Conclusions: Hepatitis C Viewed as a Metabolic Disease and Outlook for Therapeutic Strategies in the Future

We have demonstrated that HCV per se induces insulin resistance in the animal model. Superimposed high-fat diet and obesity may lead to overt diabetes. Since insulin resistance accelerates the progression of chronic hepatitis C, it would naturally influence the development of HCC. Although the association has not been established between nonalcoholic steatohepatitis and HCC, it needs to be energetically pursued in view of the histological homology of nonalcoholic steatohepatitis to chronic hepatitis C. Drugs for improving glucose metabolism and insulin resistance need to be kept in store in the treatment of hepatitis C patients who have failed to respond to antivirals, because they may well prevent progression of fibrosis and development of HCC in such patients. Traditional 'high-protein and high-calorie' diet, especially advocated in Japan after World War II, is obviously detrimental, except in some patients with advanced cirrhosis. Consultation on the dietary habit with hepatitis C patients should include iron restriction [27] as well as weight control, because high-calorie intakes are likely to accelerate hepatic fibrosis by aggravating insulin resistance.

- 14 Lerat H, Honda M, Beard MR, Loesch K, Sun J, Yang Y, Okuda M, Gosert R, Xiao SY, Weinman SA, Lemon SM: Steatosis and liver cancer in transgenic mice expressing the structural and nonstructural proteins of hepatitis C virus. *Gastroenterology* 2002;122:352-365.
- 15 Moriya K, Sintani Y, Fujie H, Miyoshi H, Tsutsumi T, Yotsuyanagi H, Yasuda K, Iino S, Kimura S, Koike K: Serum lipid profile of patients with genotype 1b hepatitis C viral infection in Japan. *Hepatol Res* 2003;25:369-374.
- 16 Naeem M, Bacon BR, Mistry B, Britton RS, Di Bisceglie AM: Changes in serum lipoprotein profile during interferon therapy in chronic hepatitis C. *Am J Gastroenterol* 2001;96:2468-2472.
- 17 Lonardo A, Adinolfi LE, Loria P, Carulli N, Ruggiero G, Day CP: Steatosis and hepatitis C virus: Mechanisms and significance for hepatic and extrahepatic disease. *Gastroenterology* 2004;126:586-597.
- 18 Allison ME, Wreghitt T, Palmer CR, Alexander GJ: Evidence for a link between hepatitis C virus infection and diabetes mellitus in a cirrhotic population. *J Hepatol* 1994;21:1135-1139.
- 19 Mehta SH, Brancati FL, Sulkowski MS, Strathdee SA, Szklo M, Thomas DL: Prevalence of type 2 diabetes mellitus among persons with hepatitis C virus infection in the United States. *Ann Intern Med* 2000;133:592-599.
- 20 Shintani Y, Fujie H, Miyoshi H, Tsutsumi T, Tsukamoto K, Kimura S, Moriya K, Koike K: Hepatitis C virus infection and diabetes: Direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;126:840-848.
- 21 Kahn BB: Type 2 diabetes: When insulin secretion fails to compensate for insulin resistance. *Cell* 1998;92:593-596.
- 22 Tsutsumi T, Suzuki T, Moriya K, Yotsuyanagi H, Shintani Y, Fujie H, Matsuura Y, Kimura S, Koike K, Miyamura T: Alteration of intrahepatic cytokine expression and AP-1 activation in transgenic mice expressing hepatitis C virus core protein. *Virology* 2002;304:415-424.
- 23 Aytug S, Reich D, Sapiro LE, Bernstein D, Begum N: Impaired IRS-1/PI3-kinase signaling in patients with HCV: A mechanism for increased prevalence of type 2 diabetes. *Hepatol* 2003;38:1384-1392.
- 24 Suzuki R, Tobe K, Aoyama M, Inoue A, Sakamoto K, Yamauchi T, Kamon J, Kubota N, Terauchi Y, Yoshimatsu H, Matsuhisa M, Nagasaka S, Ogata H, Tokuyama K, Nagai R, Kadowaki T: Both insulin signaling defects in the liver and obesity contribute to insulin resistance and cause diabetes in *Irs2(-/-)* mice. *J Biol Chem* 2004;279:25039-25049.
- 25 Petersen KF, Befroy D, Dufour S, Dziura J, Ariyan C, Rothman DL, DiPietro L, Cline GW, Shulman GI: Mitochondrial dysfunction in the elderly: Possible role in insulin resistance. *Science* 2003;300:1140-1142.
- 26 Hui JM, Sud A, Farrell GC, Bandara P, Byth K, Kench JG, McCaughan GW, George J: Insulin resistance is associated with chronic hepatitis C virus infection and fibrosis progression. *Gastroenterology* 2003;125:1695-1704.
- 27 Kato J, Kobune M, Nakamura T, Kuroiwa G, Takada K, Takimoto R, Sato Y, Fujikawa K, Takahashi M, Takayama T, Ikeda T, Niitsu Y: Normalization of elevated hepatic 8-hydroxy-2'-deoxyguanosine levels in chronic hepatitis C patients by phlebotomy and low iron diet. *Cancer Res* 2001;61:8697-8702.



Review

Oxidative stress and hepatitis C viral infection

Kazuhiko Koike*, Hideyuki Miyoshi

Department of Internal Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

Received 17 October 2005; received in revised form 4 November 2005; accepted 4 November 2005

Available online 20 December 2005

Abstract

The involvement of oxidative stress in the pathogenesis of hepatitis and hepatocellular carcinoma has been strongly suggested. Oxidative stress is produced by inflammatory processes that occur in hepatitis via immunological mechanisms. In addition, in hepatitis C virus (HCV) infectious disease, some role has been assigned to viral proteins in the induction of oxidative stress. In the presence of hepatic steatosis, insulin resistance and increased levels of some cytokines, all of which are also induced by viral protein expression, oxidative stress is enhanced in HCV infection. In this sense, the role of oxidative stress in the progression of chronic hepatitis and hepatocarcinogenesis is greater in hepatitis C than in other types of hepatitis such as hepatitis B or autoimmune hepatitis. The additive effects of oxidative stress caused by the inflammatory process and that induced by HCV proteins may, furthermore, exert synergistic effects with alterations in intracellular signaling systems such as mitogen-activated protein kinases (MAPK), which are also induced by HCV proteins. These synergistic effects may be responsible for rare characteristics, that is, the high incidence and multicentric nature of hepatocarcinogenesis in HCV infection.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Oxidative stress; Hepatitis C virus; Hepatocarcinogenesis; Lipid peroxidation; Steatosis; Insulin resistance

Contents

1. Introduction	66
2. Oxidative stress, reactive oxygen, and the liver	66
2.1. Oxidative stress and reactive oxygen	66
2.2. Antioxidation system and oxidative stress markers	66
3. Viral infection and oxidative stress	67
3.1. ROS production associated with viral infection	67
3.2. Nitric oxide production associated with viral infection	67
4. Oxidative stress caused by viral protein	68
5. Relationship of HCV infection with insulin resistance	68
6. Relationship of HCV infection with hepatic steatosis	69
7. Iron and reactive oxygen	69
8. Interactions with alcohol	69
9. Hepatocarcinogenesis and oxidative stress	69
10. Conclusions	70
References	71

* Corresponding author at: Department of Infectious Diseases, Internal Medicine, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan. Tel.: +81 3 5800 8800; fax: +81 3 5800 8799.

E-mail address: kkoike-iky@umin.ac.jp (K. Koike).

1. Introduction

There are approximately 200 million people infected with hepatitis C virus (HCV) worldwide, of which about 1.8 million are in Japan. It is one of the most serious causes of liver disease. It was reported that approximately 70% of those with HCV infection suffer from persistent infection, causing active or inactive chronic hepatitis and that about 30% of patients with chronic hepatitis are assumed to develop cirrhosis within their lifetime. Once HCV infection develops into cirrhosis, hepatocellular carcinoma (HCC) develops at an annual rate of 5–7% [1]. The strong association of oxidative stress with HCV infection has been demonstrated recently and it has become possible to explain at least part of the clinical progression of the disease. The pathogenesis of chronic hepatitis C is not merely ascribed to inflammation caused by viral infection, but the role of viral proteins in the pathogenesis was also reported [2–4]. Of proteins constituting HCV, the core protein, in particular, has various functions with respect to host cells [5] and is closely related to oxidative stress. In this overview, the relationship between HCV infection and oxidative stress is reviewed focusing on the pathological effect of the core protein of HCV, and the significance of oxidative stress in the pathogenesis of liver disease will be discussed.

2. Oxidative stress, reactive oxygen, and the liver

2.1. Oxidative stress and reactive oxygen

The main source of reactive oxygen species (ROS) in hepatocytes is the mitochondria. Outside of hepatocytes, ROS also originate from nicotinamide adenine dinucleotide phosphate

(NADPH) oxidase and xanthine oxidase in Kupffer cells and inflammatory cells. Several percent of consumed oxygen is constantly converted into ROS in the mitochondria accompanied by oxygen consumption in the electron transport system (ETS, Fig. 1). Hepatocytes contain many mitochondria and therefore have a high ROS production. Generated ROS are very unstable and highly reactive, and attack biomolecules such as DNA, lipids, and proteins. The liver not only produces much ROS but is also the center of the anti-oxidative effect in the form of protein synthesis. Oxidative stress refers to the oxidation-reaction-dominant state of the living body induced by an imbalance between the oxidation reaction caused by ROS and the anti-oxidation reaction. Main ROS include superoxide (O_2^-), hydrogen peroxide (H_2O_2) and the hydroxyl radical (HO^\bullet). ROS are mainly produced from O_2^- and converted into stable H_2O_2 through dismutation reaction. H_2O_2 is converted into highly reactive HO^\bullet in the presence of a transition metal.

2.2. Antioxidation system and oxidative stress markers

Antioxidants include glutathione (GSH), thioredoxin (TRX), vitamin E, vitamin C, and β -carotene. Reactive oxygen elimination enzymes include superoxide dismutase (SOD), glutathione peroxidase (GPx), and catalase. SOD is induced by oxidative stress and dismutates O_2^- to H_2O_2 and oxygen. GSH is a compound belonging to the SH group and is highly abundant in the living body, and the SH group provides electrons to free radicals to stabilize the radicals. GSH exists in a reduced form in cells. Because it is converted into dimeric oxidized glutathione (GSSG) and becomes stable after donating electrons, GSSG prevents free radicals from continuously scrambling for electrons. GPx decomposes H_2O_2 into water




Fig. 1. The electron transfer system (ETS) of the mitochondrion. Most of the oxygen, consumed by mammalian cells, is converted to water via the mitochondrial ETS. However, up to 5% of the electrons entering the mitochondrial ETS can become uncoupled and singly leak out onto oxygen to form superoxide. Therefore, if there is impairment in the mitochondrial ETS function, it can be a cause of the overproduction of reactive oxygen species (ROS). TCA, tricarboxylic acid; NADH, nicotinamide adenine dinucleotide phosphate; FMN, flavine mononucleotide; CoQ, coenzyme Q; cyt c, cytochrome c.




Fig. 2. Generation and scavenging of oxidative stress. SOD, super-oxide dismutase; GSH, reduced glutathione; GPx, glutathione peroxidase; TRX, thioredoxin.

and oxygen with GSH as an electron donor and reduces lipid peroxide to become neutralized. GSSG is converted back to GSH when glutathione reductase transfers an electron from NADPH to GSSG. Catalase in peroxisomes also decomposes H_2O_2 to water and oxygen. TRX is also a protein induced by oxidative stress, and is reduced via the S-S binding of the substrate protein by two SH groups in TRX and acts on the H_2O_2 elimination system via peroxiredoxins (Fig. 2).

ROS cause various forms of cellular damage. 4-Hydroxy-2-nonenal (HNE) and malondialdehyde (MDA) are the peroxidation reaction products of lipids, and 8-hydroxydeoxyguanosine (8-OHdG) is the product of DNA base modification (Fig. 3). These products serve as oxidative stress markers.




Fig. 3. Representative aldehyde species generated via lipid peroxidation reaction.

3. Viral infection and oxidative stress

3.1. ROS production associated with viral infection

Upon viral infection, ROS are produced by NADPH oxidase and xanthine oxidase in neutrophils and macrophages. In particular, NS3, one of the non-structural proteins of HCV, was reported to induce ROS production by NADPH oxidase in neutrophils [6]. Furthermore, in viral hepatitis, ROS are also produced in hepatocytes through the release of inflammatory cytokines such as TNF- α and IL-1 β from inflammatory cells. Increased hepatic or serum 8-OHdG, HNE and MDA levels are observed in chronic hepatitis C, indicating an increase in ROS production [7–13]. Findings that indicate an increase in the activity of the ROS elimination system including decreased hepatic and blood GSH levels, an increased GSSG/GSH + GSSG ratio, and an increased serum TRX level have been reported [13–16]. The findings of markedly decreased HNE level following viral eradication with interferon [12] and decreased serum ALT and TRX levels following the administration of vitamin E, an antioxidant [17], also demonstrated that oxidative stress plays an important role in chronic hepatitis C.

3.2. Nitric oxide production associated with viral infection

In the presence of an inflammation, inducible nitric oxide synthase (iNOS) is induced in macrophages and hepatocytes by TNF- α and IFN- γ [18–20]. Other investigators reported that protein kinase (PKR) activated by double-stranded RNA formed during virus reproduction in turn activates the transcripts of NF- κ B and IRF-1 to induce iNOS [21]. In the case of HCV, it was reported that its constituent proteins (E2 and non-structural (NS) protein 5A) inhibit PKR activity [22,23], but iNOS induction by viral RNA via PKR is also suspected. Indeed, iNOS synthesis correlates with intrahepatic viral load in chronic hepatitis C [24].

NO is generally synthesized as a non-specific defense reaction to infectious diseases; however, in viral infection, antiviral activity may be present or absent in various viral types [20]. NO is reported to exhibit no antiviral activity against a tick-borne encephalitis virus (TBE-V), flavivirus [25], and NO may also have no antiviral activity against HCV. On the contrary, NO causes cellular damage upon its reaction to O_2 or simultaneously produced O_2^- (reactive nitrogen species, RNS). Upon reaction to O_2^- , in particular, NO acts as a strong oxidant with the generation of peroxynitrous acid ($ONOO^-$), and $ONOO^-$ also produces nitrotyrosine through the nitration of aromatic amino acid residues in the presence of a transition metal. Nitrotyrosine accumulation was observed in correlation to inflammation severity in chronic hepatitis C tissue [26]; suggesting that the production of both NO and ROS increased. ROS and RNS are produced as defense factors for biological viral clearance, but these factors also have cytotoxic effects that

are assumed to contribute to the exacerbation of the disease state.

4. Oxidative stress caused by viral protein

The HCV genome comprises the genes of four structural proteins and six non-structural proteins (Fig. 4), and it has been reported that at least two viral proteins cause oxidative stress in cells. The core protein, a structural protein, was found to have various actions, including the induction of oxidative stress and accumulation of lipids, in experimental studies using cultured cells and transgenic mice [2,27]. Experiments using mice transgenic for the core gene showed an increased ROS production, an increased intrahepatic catalase activity, a decreased intrahepatic GSH level and a decreased GSH/GSH – GSSG ratio indicating an antioxidant effect inhibition, although there was no increase in serum ALT level nor a histological finding of hepatitis. Increased levels of intrahepatic peroxide lipids in the core gene transgenic mice with aging as compared with those in the control mice also indicate increased oxidative stress. As a mechanism underlying oxidative stress induction by the core protein, mitochondrial damage is considered. Morphological abnormalities of the mitochondria were observed in the core gene transgenic mouse liver [2], and an increased ROS production caused by damage of the mitochondrial electron transport system was noted in core-protein-expressing cells [27]. Mitochondrial DNA, which has no protective proteins such as histone, is susceptible to damage by ROS [28,29]. Mitochondrial DNA in the core gene transgenic mice showed damage as early as 3-months old. This mitochondrial damage disrupts the synthesis of proteins constituting the electron transport system complex and could also increase oxidative stress caused by damage of the electron transport system.

A study using a cell culture system demonstrated that non-structural protein 5A (NS5A) also causes oxidative stress. NS5A induces endoplasmic reticulum calcium release via

endoplasmic reticulum stress, and this leads to an increased ROS production in the mitochondria [4]. Although the effect of NS5A has not been confirmed yet by other study groups, HCV has the direct action of increasing intracellular ROS production via its proteins, separate from oxidative stress induction by inflammation caused by viral infection. A report that oxidative stress was also observed in HCV carriers with a normal ALT level [13] indicates that it is caused by a direct oxidative stress induction without being mediating inflammatory reactions.

5. Relationship of HCV infection with insulin resistance

The relationship of HCV infection with insulin resistance and type 2 diabetes has been suggested epidemiologically [30–32]. Insulin resistance was also observed in core gene transgenic mice before the onset of hepatic steatosis [33]. A disrupted tyrosine phosphorylation of the insulin receptor substrate (IRS-1) was observed in the liver of these transgenic mice. The analysis of hepatic tissues in patients with chronic hepatitis C not complicated by diabetes showed that insulin receptor and IRS-1 expression levels are elevated in patients with HCV infection, whereas the tyrosine phosphorylation of IRS-1 induced by insulin is inhibited. An excessive oxidative stress may be another potential cause of this insulin resistance. Oxidative stress indirectly blocks the phosphorylation of tyrosine residues of insulin receptors and IRS-1 and inhibits insulin signaling [34].

These reported results thus indicate an insulin signaling disorder in the liver infected with HCV [35]. There has been no report to date directly proving that hepatic insulin signaling disorder in patients with HCV infection is attributable to oxidative stress. However, because diabetes, which is the state of having abnormally high blood sugar levels that cannot be self-regulated by individual organisms, also induces oxidative stress [34], the close relationship between insulin resistance or diabetes and oxidative stress as the cause and the




Fig. 4. Structure of hepatitis C virus genome. The genome of HCV consists of two parts, structural and non-structural regions. The former comprises the core and envelope regions, and the latter consists of NS2 to NS5a, which regions chiefly code enzymes necessary for viral replication. NS, non-structural; ISDR, interferon sensitivity-determining region.

result, respectively, is a very interesting issue to investigate in the future.

6. Relationship of HCV infection with hepatic steatosis

Hepatic steatosis is frequently observed in patients with HCV infection. The relationships of HCV infection with intrahepatic viral loads and core protein levels, different prevalence of hepatic steatosis by viral genotype [higher incidence for genotype 3a], and improved steatosis following viral eradication were reported [36–38]. It is presumed from these reports that HCV itself causes hepatic steatosis. A similar hepatic steatosis caused even by the core protein alone was observed in a study using an expression system in cultured cells and transgenic mice, and it was thus suggested that the core protein plays a significant role in hepatic steatosis as the direct action of HCV [39,40]. Hyperinsulinemia induced by insulin resistance mentioned above causes the overloading of the liver with fatty acids from fat cells, and mitochondrial damage inhibits the β -oxidation of fatty acids [41]. Furthermore, the core protein was reported to inhibit microsomal triglyceride transfer protein (MTP) activity that is required when neutral fat is released as very low-density lipoproteins (VLDLs) [42]. All these actions could cause hepatic steatosis. In the liver of non-alcoholic steatohepatitis (NASH) patients, it was reported that β -oxidation in the mitochondria and peroxisomes or the metabolism of fatty acids by cytochrome P450 2E1 (CYP2E1) in microsomes is promoted under an excessive load of fatty acids, resulting in ROS production [43,44]. In HCV infection as well, intrahepatic fat accumulation possibly increases ROS production as in NASH. Because hepatic steatosis in chronic hepatitis C was reported to be a factor for disease progression [45–47], increased oxidative stress associated with hepatic steatosis is presumably involved in disease progression.

7. Iron and reactive oxygen

The iron content in the liver and spleen is high, and transition metals facilitate electron transfer and play an important role in the production of free radicals. Iron in combination with transferrin and ferritin is stable, but an unstable iron ion is freed when ferritin is decomposed by lysosomes [48]. ROS additionally promote iron release from ferritin [49]. A free iron ion catalyzes changes from relatively poor reactive O_2^- and H_2O_2 to a highly reactive HO^\bullet (Fenton reaction) [50,51]. HO^\bullet oxidizes membrane phospholipids, which compose cells and intracellular organelles, and iron forms radicals from produced peroxide lipids, thereby enhancing lipid peroxidation. Iron site-specifically combines with DNA and promotes DNA damage caused by ROS. Iron also increases ROS production by CYP2E1 [52]. A report that an enhanced peroxidation of intrahepatic lipids is attenuated by exsan-

guination in hemochromatosis also supports the involvement of iron in oxidative stress [53].

An excessively high iron content in the liver was observed in chronic hepatitis C [8,54]. Other investigators reported that iron removal therapy by exsanguination of chronic hepatitis C patients significantly improves serum ALT level without affecting viral load [55–57]. Another study showed that hepatic impairment is exacerbated following the administration of iron to chimpanzees with chronic hepatitis C [58]. Furthermore, oxidative stress is decreased by the iron removal therapy for chronic hepatitis C using intrahepatic 8-OHdG level as an index [57]. The above-mentioned reports show the close relationships of chronic hepatitis C with iron metabolism and oxidative stress.

8. Interactions with alcohol

Alcohol metabolism plays an important role in ROS production. Mainly alcohol dehydrogenase (ADH) in the cytosol and CYP2E1 (microsomal ethanol-oxidizing system) in microsomes are responsible for alcohol metabolism in the liver. When alcohol dehydrogenase oxidizes ethanol to acetaldehyde, the reduction from NAD^+ to NADH simultaneously occurs. NADH accumulation causes stress on the mitochondrial electron transfer system, leading to an increased production of ROS [59]. NADH also inhibits xanthine dehydrogenase activity, and xanthine is thereby oxidized by xanthine oxidase with the production of ROS [60]. CYP2E1 is induced by chronic alcohol intake and ROS are produced when CYP2E1 oxidizes ethanol to acetaldehyde [52,61].

There is no significant difference in hepatic peroxide level between core gene transgenic mice at 3–6-months old and control transgenic mice, but hepatic peroxide level significantly increases following the administration of a low dose of alcohol in the core gene transgenic mice [2]. ROS production increases upon glutathione reduction in HepG2 cells, with the co-expression of the core protein and CYP2D1, the latter of which is induced by alcohol [62]. These findings show that the core protein and alcohol in combination increase oxidative stress. Indeed, it was reported that alcohol intake plays a role in promoting the progression of chronic hepatitis C [63,64] and that increased levels of oxidative stress markers such as HNE and lipid hydroperoxide also support these findings [65]. From the viewpoint of oxidative stress also, HCV infection and alcohol intake are both considered to promote hepatic impairment.

9. Hepatocarcinogenesis and oxidative stress

It has been demonstrated that oxidative stress plays a key role in carcinogenesis [66,67]. Animal experiments using hepatocarcinogenesis models with the administration of a chemical substance (diethyl-nitrosamine, peroxisome proliferators) and with the administration of a choline-deficient

amino acid diet also indicates the involvement of oxidative stress [68–72]. In Long Evans Cinnamon (LEC) rats, an animal model that spontaneously develops heritable hepatitis and HCC caused by an abnormal copper accumulation, a congenitally decreased glutathione peroxidase expression level was reported, and the close relationship between oxidative stress and hepatocarcinogenesis was indicated [73].

The epidemiological relationship between HCV infection and HCC is evident [74,75], but the mechanism underlying this relationship has not been fully elucidated yet. Among postulated hypotheses on the mechanism of HCV-associated hepatocarcinogenesis, that of the involvement of the viral protein, in particular, the core protein of HCV is attractive: HCC develops in core gene transgenic mice, and carcinogenesis starts with well-differentiated carcinoma with an excessively high fat content, similar to hepatocarcinogenesis in human chronic hepatitis C, and poorly differentiated carcinoma with a low fat content develops in the form of “nodules in nodules” [76]. Because oxidative stress is increased in the core gene transgenic mice as mentioned above, it is assumed that oxidative stress plays an important role in hepatocarcinogenesis in chronic hepatitis C. Because the development of HCC is also observed in transgenic mice carrying the full-length HCV protein gene, the non-structural protein may have an additive effect to the effect of the structural proteins including the core protein, contributing to hepatocarcinogenesis [77]. NS5A, which was also reported to induce ROS production [4], may also contribute to hepatocarcinogenesis, although ROS induction by NS5A is not unequivocally confirmed yet.

Mitochondrial DNA has no potent protective proteins such as histone and is near the electron transport system, the major ROS production site. Hence, it is 10 to 15 times more susceptible to mutation caused by ROS than nuclear DNA [28,29]. In an investigation of mitochondrial DNA mutation in the human normal liver, both cancerous and non-cancerous liver tissues in patients with HCC showed very high incidences of DNA mutations [78]; thus, a relationship between oxidative stress persistence and hepatocarcinogenesis is suggested.

In the core protein expression system in the hepatic tissue and cultured cells of core gene transgenic mice, the activation of transcription factor AP-1 via mitogen-activated protein (MAP) kinase was observed [79–83]. The activation of the transcription factors AP-1, NF- κ B, and signal transducer and activator of transcription (STAT) 3 by NS5A were also reported [4,84]. The activation of these transcription factors may facilitate cell proliferation, contributing to tumorigenic transformation.

It was also reported that ROS facilitate apoptosis via c-Jun N-terminal kinase (JNK)/p38 MAP kinase or by directly attacking the mitochondria. Apoptosis is a protective mechanism of the host against viral infection and carcinogenesis. Some reports stated that the core protein facilitates apoptosis [85–88], whereas other reports stated that the core protein inhibits apoptosis [89–92]; thus, no fixed view has yet been established. If it indeed inhibits apoptosis, it is assumed that this inhibition proceeds by maintaining oxidative stress and

that the core protein has a beneficial effect against carcinogenesis and persistent viral infection.

In HCV infection, viral proteins such as the core protein and, possibly, NS5A protein induce oxidative stress, intracellular signaling, and transcription factors, which are not reflected in blood ALT level, contributing to the progression of carcinogenesis. Carcinogenesis, however, is slow as is observed in humans and core gene transgenic mice, the latter of which developed HCC in the latter half of their life. Recently, Okanoue et al. reported a long-term follow-up study of subjects with persistent HCV infection who had persistently normal ALT levels (PNAL) [93]. In their study, serum thioredoxin levels were not elevated in those with PNAL compared to those with chronic hepatitis. This may apparently seem contradictory to the results of our above-mentioned animal model studies. However, we should realize that anti-oxidant system is also instrumental in the liver. In these relatively younger people with PNAL than those with CH [93], active anti-oxidant system may erase the apparent elevation of ROS. Such a phenomenon was described in a mouse model by Moriya et al. [2], in which ROS was apparently normal in young core gene transgenic mice with the activation of catalase and reduction of GSH. Clinically, the presence of inflammation is thought to facilitate the process of hepatocarcinogenesis.

10. Conclusions

A very close pathological relationship between oxidative stress and HCV infection is observed, as shown by the above overview of relevant publications and discussion. The causes of oxidative stress in HCV infection are considered to include various factors such as mitochondrial damage, endoplasmic reticulum stress, iron accumulation, and lipid accumulation in the liver. Various study results demonstrated that even only viral proteins, mainly the HCV core protein, cause oxidative stress. When inflammation via immunoreactions to viral infection is added to oxidative stress, ROS production is expected to further increase, leading to a state in which the anti-oxidation system cannot cope with. In this sense, inflammation in chronic hepatitis C is considered to be qualitatively different from inflammation observed in other types of hepatitis such as autoimmune hepatitis or hepatitis B [94] (Fig. 5). As a treatment of chronic hepatitis C, the eradication of the virus is ideal. If it is not possible, however, the control of factors that exacerbate oxidative stress, such as inflammation via immune reaction and alcohol, and the relief of oxidative stress by the iron removal therapy and the administration of an anti-oxidation agent are considered to delay the progression of chronic hepatitis.

The development of such new anti-oxidation agents is being awaited. In further studies on the development of new therapies for hepatitis C and control methods for hepatocarcinogenesis in the future, the importance of those focusing on oxidative stress is expected to markedly increase.




Fig. 5. Oxidative stress and hepatocarcinogenesis in various types of hepatitis (hypothesis). Oxidative stress is generated in all types of hepatitis via inflammation accompanied by continual cell death and regeneration. In HCV infection, HCV itself causes the production of oxidative stress in a synergy with inflammation. In this sense, the quality of "inflammation" in HCV infection may be different from that in other types of hepatitis. Additional impact of HCV proteins on the intracellular signal transduction would provoke the development of HCC. These may explain the conspicuous properties of HCC development.

References

- [1] Ikeda K, Saitoh S, Suzuki Y, et al. Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: a prospective observation of 2215 patients. *J Hepatol* 1998;28:930–8.
- [2] Moriya K, Nakagawa K, Santa T, et al. Oxidative stress in the absence of inflammation in a mouse model for hepatitis C virus-associated hepatocarcinogenesis. *Cancer Res* 2001;61:4365–70.
- [3] Koike K, Moriya K, Ishibashi K, et al. Sialadenitis histologically resembling Sjögren syndrome in mice transgenic for hepatitis C virus envelope genes. *Proc Natl Acad Sci USA* 1997;94:233–6.
- [4] Gong G, Waris G, Tanveer R, Siddiqui A. Human hepatitis C virus NS5A protein alters intracellular calcium levels, induces oxidative stress, and activates STAT-3 and NF- κ B. *Proc Natl Acad Sci USA* 2001;98:9599–604.
- [5] Suzuki R, Suzuki T, Ishii K, Matsuura Y, Miyamura T. Processing and functions of hepatitis C virus proteins. *Intervirology* 1999;42:145–52.
- [6] Bureau C, Bernad J, Chauouche N, et al. Non-structural 3 protein of hepatitis C virus triggers an oxidative burst in human monocytes via activation of NADPH oxidase. *J Biol Chem* 2001;276:23077–83.
- [7] Shimoda R, Nagashima M, Sakamoto M, et al. Increased formation of oxidative DNA damage, 8-hydroxydeoxyguanosine, in human livers with chronic hepatitis. *Cancer Res* 1994;54:3171–2.
- [8] Farinati F, Cardin R, Maria ND, et al. Iron storage, lipid peroxidation and glutathione turnover in chronic anti-HCV positive hepatitis. *J Hepatol* 1995;22:449–56.
- [9] De Maria N, Calantoni A, Fagivoli S, et al. Association between reactive oxygen species and disease activity in chronic hepatitis C. *Free Radic Biol Med* 1996;21:291–5.
- [10] Romero MJ, Bosch-Morell F, Romero B, Rodrigo JM, Serra MA, Romero FJ. Serum malondialdehyde: possible use for the clinical management of chronic hepatitis C patients. *Free Radic Biol Med* 1998;25:993–7.
- [11] Farinati F, Cardin R, Degan P, et al. Oxidative DNA damage in circulating leukocytes occurs as an early event in chronic HCV infection. *Free Radic Biol Med* 1999;27:1284–91.
- [12] Kageyama F, Kobayashi Y, Kawasaki T, Toyokuni S, Uchida K, Nakamura H. Successful interferon therapy reverses enhanced hepatic iron accumulation and lipid peroxidation in chronic hepatitis C. *Am J Gastroenterol* 2000;95:1041–50.
- [13] Vendemiale G, Grattagliano I, Portincasa P, Serviddio G, Palasciano G, Altomare E. Oxidative stress in symptom-free HCV carriers: relation with ALT flare-up. *Eur J Clin Invest* 2001;31:54–63.
- [14] Barbaro G, Di Lorenzo G, Ribersani M, et al. Serum ferritin and hepatic glutathione concentrations in chronic hepatitis C patients related to the hepatitis C virus genotype. *J Hepatol* 1999;30:774–82.
- [15] Sumida Y, Nakashima T, Yoh T, et al. Serum thioredoxin levels as an indicator of oxidative stress in patients with hepatitis C virus infection. *J Hepatol* 2000;33:616–22.
- [16] Jain SK, Pemberton PW, Smith A, et al. Oxidative stress in chronic hepatitis C: not just a feature of late stage disease. *J Hepatol* 2002;36:805–11.
- [17] Mahmood S, Yamada G, Niiyama G, et al. Effect of vitamin E on serum aminotransferase and thioredoxin levels in patients with viral hepatitis C. *Free Radic Res* 2003;37:781–5.
- [18] Schwarz KB. Oxidative stress during viral infection: a review. *Free Radic Biol Med* 1996;21:641–9.
- [19] Majano PL, Garcia-Monzon C, Lopez-Cabrera M, et al. Inducible nitric oxide synthase expression in chronic viral hepatitis. *J Clin Invest* 1998;101:1343–52.
- [20] Akaike T. Role of free radicals in viral pathogenesis and mutation. *Rev Med Virol* 2001;11:87–101.
- [21] Uetani K, Der SD, Zamanian-Daryoush M, et al. Central role of double-stranded RNA-activated protein kinase in microbial induction of nitric oxide synthase. *J Immunol* 2000;165:988–96.
- [22] Gale Jr MJ, Korth MJ, Tang NM, et al. Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the non-structural 5A protein. *Virology* 1997;230:217–27.
- [23] Taylor DR, Shi ST, Romano PR, Barber GN, Lai MM. Inhibition of the interferon-inducible protein kinase PKR by HCV E2 protein. *Science* 1999;285:107–10.
- [24] Mihm S, Fayyazi A, Ramadori G. Hepatic expression of inducible nitric oxide synthase transcripts in chronic hepatitis C virus infection.

- tion: relation to hepatic viral load and liver injury. *Hepatology* 1997;26:451–8.
- [25] Kreil TR, Eibl MM. Nitric oxide and viral infection: no antiviral activity against a flavivirus in vitro, and evidence for contribution to pathogenesis in experimental infection in vivo. *Virology* 1996;219:304–6.
- [26] Garcia-Monzon C, Majano PL, Zubia I, Sanz P, Apolinario A, Moreno-Otero R. Intrahepatic accumulation of nitrotyrosine in chronic viral hepatitis is associated with histological severity of liver disease. *J Hepatol* 2000;32:331–8.
- [27] Okuda M, Li K, Beard MR, et al. Mitochondrial injury, oxidative stress, and antioxidant gene expression are induced by hepatitis C virus core protein. *Gastroenterology* 2002;122:366–75.
- [28] Shigenaga MK, Hagen TM, Ames BN. Oxidative damage and mitochondrial decay in aging. *Proc Natl Acad Sci USA* 1994;91:10771–8.
- [29] Lee CM, Weindruch R, Aiken JM. Age-associated alterations of the mitochondrial genome. *Free Radic Biol Med* 1997;22:1259–69.
- [30] Caronia S, Taylor K, Pagliaro L, et al. Further evidence for an association between non-insulin-dependent diabetes mellitus and chronic hepatitis C virus infection. *Hepatology* 1999;30:1059–63.
- [31] Alexander GJM. An association between hepatitis C virus infection and type 2 diabetes mellitus: what is the connection? *Ann Intern Med* 2000;133:650–2.
- [32] Mehta SH, Brancati FL, Sulkowski MS, Strathdee SA, Szklo M, Thomas DL. Prevalence of type 2 diabetes mellitus among persons with hepatitis C virus infection in the United States. *Ann Intern Med* 2000;133:592–9.
- [33] Shintani Y, Fujie H, Miyoshi H, et al. Hepatitis C virus infection and diabetes: direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;126:840–8.
- [34] Evans JL, Goldfine ID, Maddux BA, Grodsky GM. Are oxidative stress-activated signaling pathways mediators of insulin resistance and β -cell dysfunction? *Diabetes* 2003;52:1–8.
- [35] Aytug S, Reich D, Sapiro LE, Bernstein D, Begum N. Impaired IRS-1/PI3-kinase signaling in patients with HCV: a mechanism for increased prevalence of type 2 diabetes. *Hepatology* 2003;38:1384–92.
- [36] Fujie H, Yotsuyanagi H, Moriya K, et al. Steatosis and intrahepatic hepatitis C virus in chronic hepatitis. *J Med Virol* 1999;59:141–5.
- [37] Rubbia-Brandt L, Quadri R, Abid K, et al. Hepatocyte steatosis is a cytopathic effect of hepatitis C virus genome 3. *J Hepatol* 2000;33:106–15.
- [38] Kumar D, Farrell GC, Fung C, George J. Hepatitis C virus genotype 3 is cytopathic in hepatocytes. Genotype-specific reversal of hepatic steatosis after sustained response to antiviral therapy. *Hepatology* 2002;36:1266–72.
- [39] Moriya K, Yotsuyanagi H, Shintani Y, et al. Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J Gen Virol* 1997;78:1527–31.
- [40] Barba G, Harper F, Harada T, et al. Hepatitis C virus core protein shows a cytoplasmic localization and associates to cellular lipid storage droplets. *Proc Natl Acad Sci USA* 1997;94:1200–5.
- [41] Fromenty B, Pessayre D. Inhibition of mitochondrial β -oxidation as a mechanism of hepatotoxicity. *Pharmacol Ther* 1995;67:101–54.
- [42] Perlemuter G, Sabile A, Letteron P, et al. Hepatitis C virus core protein inhibits microsomal triglyceride transfer protein activity and very low density lipoprotein secretion: a model of viral-related steatosis. *FASEB J* 2002;16:185–94.
- [43] Day CP, James OFW. Steatohepatitis: a tale of two "hits"? *Gastroenterology* 1998;114:842–5.
- [44] Weltman MD, Farrell GC, Hall P, Ingelman-Sundberg M, Liddle C. Hepatic cytochrome P450 2E1 is increased in patients with non-alcoholic steatohepatitis. *Hepatology* 1998;27:128–33.
- [45] Czaja AJ, Carpenter HA, Santrach PJ, Moore SB. Host- and disease-specific factors affecting steatosis in chronic hepatitis C. *J Hepatol* 1998;29:198–206.
- [46] Hourigan LF, Macdonald GA, Purdie D, et al. Fibrosis in chronic hepatitis C correlates significantly with body mass index and steatosis. *Hepatology* 1999;29:1215–9.
- [47] Serfaty L, Poujol-Robert A, Carbonell N, Chazouilleres O, Poupon RE, Poupon R. Effect of the interaction between steatosis and alcohol intake on liver fibrosis progression in chronic hepatitis C. *Am J Gastroenterol* 2002;97:1807–12.
- [48] Radisky DC, Kaplan J. Iron in cytosolic ferritin can be recycled through lysosomal degradation in human fibroblasts. *Biochem J* 1998;336:201–5.
- [49] Kakhlon O, Cabantchik ZI. The labile iron pool: characterization, measurement, and participation in cellular processes. *Free Radic Biol Med* 2002;33:1037–46.
- [50] Halliwell B, Gutteridge JMC. Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J* 1984;219:1–14.
- [51] Aust SD, Morehouse LA, Thomas CE. Roles of metals in oxygen radical reactions. *Free Radic Biol Med* 1985;1:3–25.
- [52] Cederbaum AI. Iron CYP2E1-dependent oxidative stress and toxicity. *Alcohol* 2003;30:115–20.
- [53] Houghum K, Ramm G, Crawford DH, Witzum JL, Powell LW, Chojkier M. Excess iron induces hepatic oxidative stress and transforming growth factor β 1 in genetic hemochromatosis. *Hepatology* 1997;26:605–10.
- [54] Bonkovsky H, Banner BF, Rothman AL. Iron and chronic viral hepatitis. *Hepatology* 1997;25:759–68.
- [55] Hayashi H, Takikawa T, Nishimura N, Yano M, Isomura T, Sakamoto N. Improvement of serum aminotransferase levels after phlebotomy in patients with chronic active hepatitis C and excess hepatic iron. *Am J Gastroenterol* 1994;89:986–8.
- [56] Sartori M, Andorno S, Rigamonti C, Boldorini R. Chronic hepatitis C treated with phlebotomy alone: biochemical and histological outcome. *Dig Liver Dis* 2001;33:157–62.
- [57] Kato J, Kobune M, Nakamura T, et al. Normalization of elevated hepatic 8-hydroxy-2'-deoxyguanosine levels in chronic hepatitis C patients by phlebotomy and low iron diet. *Cancer Res* 2001;61:8697–702.
- [58] Bassett SE, Bisceglie AMD, Bacon BR, et al. Effects of iron loading on pathogenicity in hepatitis C virus-infected chimpanzees. *Hepatology* 1999;29:1884–92.
- [59] Cunningham CC, Bailey SM. Ethanol consumption and liver mitochondrial function. *Biol Signals Recept* 2001;10:271–82.
- [60] Kato S, Kawase T, Alderman J, et al. Role of xanthine oxidase in ethanol-induced lipid peroxidation in rats. *Gastroenterology* 1990;98:203–10.
- [61] Ingelman-Sundberg M, Johansson I. Mechanisms of hydroxyl radical formation and ethanol oxidation by ethanol-inducible and other forms of rabbit liver microsomal cytochrome P-450. *J Biol Chem* 1984;259:6447–58.
- [62] Wen F, Abdalla MY, Aloman C, et al. Increased prooxidant production and enhanced susceptibility to glutathione depletion in HepG2 cells co-expressing HCV core protein and CYP2E1. *J Med Virol* 2004;72:230–40.
- [63] Ostapowicz G, Watzon KJR, Locarnini SA, Desmond PV. Role of alcohol in the progression of liver disease caused by hepatitis C virus infection. *Hepatology* 1998;27:1730–5.
- [64] Hassan MM, Hwang LY, Hatten CJ, et al. Risk factors for hepatocellular carcinoma: synergism of alcohol with viral hepatitis and diabetes mellitus. *Hepatology* 2002;36:1206–13.
- [65] Rigamonti C, Mottaran E, Reale E, et al. Moderate alcohol consumption increases oxidative stress in patients with chronic hepatitis C. *Hepatology* 2003;38:42–9.
- [66] Cerutti PA, Trump BF. Inflammation and oxidative stress in carcinogenesis. *Cancer Cells* (Cold Spring Harbor) 1991;3:1–7.
- [67] Dreher D, Junod AF. Role of oxygen free radicals in cancer development. *Eur J Cancer* 1996;32:30–8.

- [68] Perera MI, Betschart JM, Virji MA, Katyal SL, Shinozuka H. Free radical injury and liver tumor promotion. *Toxicol Pathol* 1987;15:51–9.
- [69] Reddy JK, Rao MS. Oxidative DNA damage caused by persistent peroxisome proliferation: its role in hepatocarcinogenesis. *Mutat Res* 1989;214:63–8.
- [70] Nakae D, Kobayashi Y, Akai H, et al. Involvement of 8-hydroxyguanine formation in the initiation of rat liver carcinogenesis by low dose levels of *N*-nitrosodiethylamine. *Cancer Res* 1997;57:1281–7.
- [71] Nakae D, Kotake Y, Kishida H, et al. Inhibition by phenyl *N*-tert-butyl nitron on early phase carcinogenesis in the livers of rats fed a choline-deficient, L-amino acid-defined diet. *Cancer Res* 1998;58:4548–51.
- [72] Floyd RA, Kotake Y, Hensley K, Nakae D, Konishi Y. Reactive oxygen species in choline deficiency induced carcinogenesis and nitron inhibition. *Mol Cell Biochem* 2002;234–235:195–203.
- [73] Suemizu H, Yoshimura S, Takeichi N, Moriuchi T. Decreased expression of liver glutathione peroxidase in Long-Evans cinnamon mutant rats predisposed to hepatitis and hepatoma. *Hepatology* 1994;19:694–700.
- [74] Saito I, Miyamura T, Ohbayashi A, et al. Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc Natl Acad Sci USA* 1990;87:6547–9.
- [75] Simonetti RG, Camma C, Fiorello F, et al. Hepatitis C virus infection as a risk factor for hepatocellular carcinoma in patients with cirrhosis. *Ann Intern Med* 1992;116:97–102.
- [76] Moriya K, Fujie H, Shintani Y, et al. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065–7.
- [77] Lerat H, Honda M, Beard MR, et al. Steatosis and liver cancer in transgenic mice expressing the structural and non-structural proteins of hepatitis C virus. *Gastroenterology* 2002;122:352–65.
- [78] Nishikawa M, Nishiguchi S, Shiomi S, et al. Somatic mutation of mitochondrial DNA in cancerous and non-cancerous liver tissue in individuals with hepatocellular carcinoma. *Cancer Res* 2001;61:1843–5.
- [79] Shrivastava A, Manna SK, Ray R, Aggarwal BB. Ectopic expression of hepatitis C virus core protein differentially regulates nuclear transcription factors. *J Virol* 1998;72:9722–8.
- [80] Kato N, Yoshida H, Kioko Ono-Nita S, et al. Activation of intracellular signaling by hepatitis B and C viruses: C-viral core is the most potent signal inducer. *Hepatology* 2000;32:405–12.
- [81] Erhardt A, Hassan M, Heintges T, Haussinger D. Hepatitis C virus core protein induces cell proliferation and activates ERK, JNK, and p38 MAP kinases together with the MAP kinase phosphatase MKP-1 in a HepG2 Tet-Off cell line. *Virology* 2002;292:272–84.
- [82] Tsutsumi T, Suzuki T, Moriya K, et al. Alteration of intrahepatic cytokine expression and AP-1 activation in transgenic mice expressing hepatitis C virus core protein. *Virology* 2002;304:415–24.
- [83] Tsutsumi T, Suzuki T, Moriya K, et al. Hepatitis C virus core protein activates ERK and p38 MAPK in cooperation with ethanol in transgenic mice. *Hepatology* 2003;38:820–8.
- [84] Qadri I, Iwahashi M, Capasso JM, Hopken MW, Flores S, Schaack J. Induced oxidative stress and activated expression of manganese superoxide dismutase during hepatitis C virus replication: role of JNK, p38 MAPK and AP-1. *Biochem J* 2004;378:919–28.
- [85] Ruggieri A, Harada T, Matsuura Y, Miyamura T. Sensitization to Fas-mediated apoptosis by hepatitis C virus core protein. *Virology* 1997;229:68–76.
- [86] Zhu N, Khoshnan A, Schneider R, et al. Hepatitis C virus core protein binds to the cytoplasmic domain of tumor necrosis factor (TNF) receptor 1 and enhances TNF-induced apoptosis. *J Virol* 1998;72:3691–7.
- [87] Hahn CS, Cho YG, Kang BS, Lester IM, Hahn YS. The HCV core protein acts as a positive regulator of fas-mediated apoptosis in a human lymphoblastoid T cell line. *Virology* 2000;276:127–37.
- [88] Honda M, Kaneko S, Shimazaki T, et al. Hepatitis C virus core protein induces apoptosis and impairs cell-cycle regulation in stably transformed Chinese hamster ovary cells. *Hepatology* 2000;31:1351–9.
- [89] Ray RB, Meyer K, Steele R, Shrivastava A, Aggarwal BB, Ray R. Inhibition of tumor necrosis factor (TNF- α)-mediated apoptosis by hepatitis C virus core protein. *J Biol Chem* 1998;273:2256–9.
- [90] Marusawa H, Hijikata M, Chiba T, Shimotohno K. Hepatitis C virus core protein inhibits Fas- and tumor necrosis factor α -mediated apoptosis via NF- κ B activation. *J Virol* 1999;73:4713–20.
- [91] Otsuka M, Kato N, Taniguchi H, et al. Hepatitis C virus core protein inhibits apoptosis via enhanced Bcl-xL expression. *Virology* 2002;296:84–93.
- [92] Sacco R, Tsutsumi T, Suzuki R, et al. Antiapoptotic regulation by hepatitis C virus core protein through up-regulation of inhibitor of caspase-activated DNase. *Virology* 2003;317:24–35.
- [93] Okanoue T, Makiyama A, Nakayama M, et al. A follow-up study to determine the value of liver biopsy and need for antiviral therapy for hepatitis C virus carriers with persistently normal serum aminotransferase. *J Hepatol* 2005;43:599–605.
- [94] Koike K. Molecular basis of hepatitis C virus-associated hepatocarcinogenesis: lessons from animal model studies. *Clin Gastroenterol Hepatol* 2005;3:S132–5.

Editorial

Oxidative stress and apoptosis in hepatitis C: the core issue

Article on page 257

Hepatitis C virus core protein inhibits deoxycholic acid-mediated apoptosis despite generating mitochondrial reactive oxygen species

HARA Y, HINO K, OKUDA M, et al.

Extra- as well as intracellular stimuli elicit a wide range of responses, such as cell proliferation, differentiation, survival and apoptosis, via the regulation of intracellular signaling. Recent studies have revealed that stress-responsive signal transduction pathways are stringently regulated by the intracellular redox state.¹ The redox state of the cell is determined by the delicate balance between the levels of oxidizing and reducing equivalents, including reactive oxygen species (ROS) and endogenous antioxidants. The production of ROS, a representative of oxidative stress, fluctuates in response to alterations in both external and internal environments and, in turn, triggers specific signaling cascades, such as mitogen-activated protein kinases, which determine cell survival or death. Thus, ROS are profoundly involved in cell death or apoptosis.

In the liver, ROS are also key cytotoxic and signaling mediators in the pathophysiology of liver diseases, including viral hepatitis, in which hepatocytes and resident and infiltrating phagocytes can generate ROS. While ROS are able to cause cell death through massive lipid peroxidation, they also act to modulate signal transduction pathways by affecting redox-sensitive enzymes, transcription factors, and organelles, including mitochondria and endoplasmic reticulum. ROS, thus, directly regulate apoptotic and necrotic cell death.² In addition, ROS have indirect effects on the pathophysiology of cell death by supporting protease activity via inactivation of antiproteases.

In hepatitis C virus (HCV) infection, both ROS and apoptosis are closely involved in the process of progressive liver diseases from chronic hepatitis to cirrhosis and hepatocellular carcinoma (HCC).^{3–7} ROS are assumed to play a major role in the pathogenesis of chronic hepatitis, which is characterized by continual

cell death followed by regeneration. In this condition, the viral proteins of HCV, the core and nonstructural (NS) 5A proteins, have been shown to play a role in inducing ROS as well as in modulating apoptosis of hepatocytes.

The HCV genome comprises the genes of four structural proteins and six nonstructural proteins, and at least two of these viral proteins have been reported to cause oxidative stress in cells. The core protein, a structural protein, has been found to have various actions, including the induction of oxidative stress and the accumulation of lipids, in experimental studies using cultured cells and transgenic mice.^{7,8} Experiments using mice transgenic for the core gene showed increased ROS production, increased intrahepatic catalase activity, a decreased intrahepatic glutathione (GSH) level, and a decreased GSH/GSH-GSSG (dimeric oxidized glutathione) ratio, indicating inhibition of antioxidation effects, although there was neither an increase in the serum alanine aminotransferase (ALT) level nor a histological finding of hepatitis.⁷ Increased levels of intrahepatic peroxide lipids in the core gene transgenic mice with aging, compared with levels in control mice, also indicate an increase in oxidative stress. One possible mechanism underlying oxidative stress induction by the core protein is mitochondrial damage. Morphological abnormalities of the mitochondria have been observed in core gene transgenic mouse liver,⁷ and increased ROS production caused by damage to the mitochondrial electron transport system has been noted in core protein-expressing cells.⁸ Mitochondrial DNA, which has no protective proteins such as histone, is susceptible to damage by ROS. Mitochondrial DNA damage in the core gene transgenic mice appeared when they were as young as 3 months old. This mitochondrial damage disrupts the synthesis of proteins constituting the electron transport system complex and might also increase oxidative stress caused by damage to the electron transport system.

Reprint requests to: K. Koike




Fig. 1. Relationships among oxidative stress, apoptosis, and HCV proteins

A study using a cell culture system demonstrated that the NS5A protein also causes oxidative stress. NS5A induces the endoplasmic reticulum to release calcium by causing stress to the endoplasmic reticulum, and this release leads to increased ROS production in the mitochondria.⁹ Although the effect of NS5A has not yet been confirmed by other study groups, HCV has the direct action of increasing intracellular ROS production via its proteins, separate from the induction of oxidative stress as a result of inflammation caused by viral infection (Fig. 1). A report that oxidative stress is also observed in HCV carriers with a normal ALT level¹⁰ indicates that it is induced directly, without any mediating inflammatory reactions being necessary.

In contrast to the production of ROS by the core protein, which is now quite evident, the role of HCV core protein in apoptosis is rather controversial. Regarding the HCV-induced apoptotic mechanism, the HCV core protein may have a regulatory function in modulating apoptosis, either by enhancing or inhibiting it. In particular, the core protein exhibits both proapoptotic and antiapoptotic actions, depending on experimental conditions and the type of cells used,¹¹⁻¹⁴ whereas both the NS3 and the NS5A proteins have antiapoptotic effects (Fig. 1).¹⁵ Modulation of apoptosis may involve binding of the core protein to the intracellular signal transducing portion of death receptors such as TNF- α , Fas, or lymphotoxin- $\alpha\beta$. Thus, HCV proteins may modulate hepatocyte apoptosis by indirect rather than by direct mechanisms. The real role of the core protein in the apoptotic process is, thus, not defined yet.

In the current issue of *Journal of Gastroenterology*, Hara et al.¹⁶ tried to elucidate this core issue of HCV pathogenesis by separating the two properties of the HCV core protein with cultured cells, Huh-7 and HeLa. They confirmed that the core protein induced ROS, which was followed by activation of the scavenging system and insults to the cellular DNA, as shown previously.⁶ In the study by Hara et al.,¹⁶ the core protein inhibited the proapoptotic action of deoxycholic acid (DCA), which is known to cause both ROS production and apoptosis. Thus, the core protein seems to act to

oppose the proapoptotic function of ROS, which ROS are also induced by the core protein itself. Such apparently opposing actions of the core protein, the production of ROS and the inhibition of apoptosis, might explain the mode of hepatocarcinogenesis in HCV infection: hepatocytes with ROS-induced DNA damage may evade apoptosis by another effect of the core protein that inhibits apoptosis. Such a mechanism, similar to one previously postulated by other researchers,³ in which both ROS production and mitogen-activated protein kinase activation are ascribed to the core protein, may clarify how cells with DNA damage can survive and develop into buds of HCC. Regrettably, the current study was done using DCA as an agent to induce both apoptosis and ROS, making it difficult to interpret the authentic role of the core protein in the execution of such biological functions. As noted above, the effect of the core protein in apoptosis varies depending on the system used. Therefore, further studies using different systems may be necessary to bring a conclusion to the core issue of HCV-induced pathogenesis associated with the multipotential HCV core protein.

Kazuhiko Koike, MD, PhD

Department of Infectious Diseases, Internal Medicine, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

References

- Schattenberg JM, Singh R, Wang Y, Lefkowitz JH, Rigoli RM, Scherer PE, et al. JNK1 but not JNK2 promotes the development of steatohepatitis in mice. *Hepatology* 2006;43:163-72.
- Rahman I, Biswas SK, Jimenez LA, Torres M, Forman HJ. Glutathione, stress responses, and redox signaling in lung inflammation. *Antioxid Redox Signal* 2005;7:42-59.
- Koike K. Molecular basis of hepatitis C virus-associated hepatocarcinogenesis: lessons from animal model studies. *Clin Gastroenterol Hepatol* 2005;3:S132-5.
- Tanikawa K. Pathogenesis and treatment of hepatitis C virus-related liver diseases. *Hepatobiliary Pancreat Dis Int* 2004;3:17-20.
- Farinati F, Cardin R, De Maria N, Della Libera G, Marafin C, Lecis E, et al. Iron storage, lipid peroxidation and glutathione turnover in chronic anti-HCV positive hepatitis. *J Hepatol* 1995; 22:449-56.
- Kitase A, Hino K, Furutani T, Okuda M, Gondo T, Hidaka I, et al. In situ detection of oxidized n-3 polyunsaturated fatty acids in chronic hepatitis C: correlation with hepatic steatosis. *J Gastroenterol* 2005;40:617-24.
- Moriya K, Nakagawa K, Santa T, Shintani Y, Fujie H, Miyoshi H, et al. Oxidative stress in the absence of inflammation in a mouse model for hepatitis C virus-associated hepatocellular carcinogenesis. *Cancer Res* 2001;61:4365-70.
- Okuda M, Li K, Beard MR, Showalter LA, Scholle F, Lemon SM, et al. Mitochondrial injury, oxidative stress, and antioxidant gene expression are induced by hepatitis C virus core protein. *Gastroenterology* 2002;122:366-75.
- Gong G, Waris G, Tanveer R, Siddiqui A. Human hepatitis C virus NS5A protein alters intracellular calcium levels, induces

oxidative stress, and activates STAT-3 and NF- κ B. *Proc Natl Sci U S A* 2001;98:9599-604.

10. Vendemiale G, Grattagliano I, Portincasa P, Serviddio G, Palasciano G, Altomare E. Oxidative stress in symptom-free HCV carriers: relation with ALT flare-up. *Eur J Clin Invest* 2001;31:54-63.
11. Ruggieri A, Harada T, Matsuura Y, Miyamura T. Sensitization to Fas-mediated apoptosis by hepatitis C virus core protein. *Virology* 1997;229:68-76.
12. Ray RB, Meyer K, Steele R, Shrivastava A, Aggarwal BB, Ray R. Inhibition of tumor necrosis factor (TNF- α)-mediated apoptosis by hepatitis C virus core protein. *J Biol Chem* 1998;273:2256-9.
13. Zhu N, Khoshnan A, Schneider R, Matsumoto M, Dennert G, Ware C, et al. Hepatitis C virus core protein binds to the cytoplasmic domain of tumor necrosis factor (TNF) receptor 1 and enhances TNF-induced apoptosis. *J Virol* 1998;72:3691-7.
14. Marusawa H, Hijikata M, Chiba T, Shimotohno K. Hepatitis C virus core protein inhibits Fas- and tumor necrosis factor α -mediated apoptosis via NF- κ B activation. *J Virol* 1999;73:4713-20.
15. Macdonald A, Harris M. Hepatitis C virus NS5A: tales of a promiscuous protein. *J Gen Virol* 2004;85:2485-502.
16. Hara Y, Hino K, Okuda M, Furutani T, Hidaka I, Yamaguchi Y, et al. Hepatitis C virus core protein inhibits deoxycholic acid-mediated apoptosis despite generating mitochondrial reactive oxygen species. *J Gastroenterol* 2006;41:257-68.



RAPID COMMUNICATION

Risk factors for retinopathy associated with interferon α -2b and ribavirin combination therapy in patients with chronic hepatitis C

Chiaki Okuse, Hiroshi Yotsuyanagi, Yoshihiko Nagase, Yuhtaro Kobayashi, Kiyomi Yasuda, Kazuhiko Koike, Shiro Iino, Michihiro Suzuki, Fumio Itoh

Chiaki Okuse, Yoshihiko Nagase, Yuhtaro Kobayashi, Michihiro Suzuki, Fumio Itoh, Division of Gastroenterology and Hepatology, Department of Internal Medicine, St. Marianna University School of Medicine, Kawasaki, Kanagawa, 216-8511, Japan
Hiroshi Yotsuyanagi, Kazuhiko Koike, Department of Infectious Diseases, Internal Medicine, Faculty of Medicine, University of Tokyo, 7-3-1, Hongo, Bunkyo, Tokyo 113-8655, Japan
Kiyomi Yasuda, Shiro Iino, Center for Liver Diseases, Kiyokawa Hospital, Minamiasagaya, Suginami, Tokyo, 166-0004, Japan
Correspondence to: Dr. Hiroshi Yotsuyanagi, Department of Infectious Diseases, Internal Medicine, Faculty of Medicine, University of Tokyo, 7-3-1, Hongo, Bunkyo, Tokyo 113-8655, Japan. hyotsu-ty@umin.ac.jp
Telephone: +81-3-58008720 Fax: +81-3-58008796
Received: 2006-01-13 Accepted: 2006-02-18

CONCLUSION: Retinopathy associated with combination therapy of interferon α -2b and ribavirin tends to develop in patients with hypertension.

© 2006 The WJG Press. All rights reserved.

Key words: Retinopathy; Ribavirin; Chronic hepatitis C; Interferon

Okuse C, Yotsuyanagi H, Nagase Y, Kobayashi Y, Yasuda K, Koike K, Iino S, Suzuki M, Itoh F. Risk factors for retinopathy associated with interferon α -2b and ribavirin combination therapy in patients with chronic hepatitis C. *World J Gastroenterol* 2006; 12(23): 3756-3759

<http://www.wjgnet.com/1007-9327/12/3756.asp>

Abstract

AIM: To elucidate the frequency and risk factors for retinopathy in patients with chronic hepatitis C who are treated by interferon-ribavirin combination therapy.

METHODS: We prospectively analyzed 73 patients with histologically confirmed chronic hepatitis C, who underwent combination therapy for 24 wk. Optic fundi were examined before, and 2, 4, 12 and 24 wk after the start of combination therapy.

RESULTS: Fourteen patients (19%) developed retinopathy, which was initially diagnosed by the appearance of a cotton wool spot in 12 patients. Retinal hemorrhage was observed in 5 patients. No patient complained of visual disturbance. Retinopathy disappeared in 9 patients (64%) despite the continuation of combination therapy. However, retinopathy persisted in 5 patients with retinal hemorrhage. A comparison of the clinical background between the groups with and without retinopathy showed no significant differences in age, gender, viral genotype, RNA level, white blood cell count, platelet count, prothrombin time, complications by diabetes mellitus or hypertension, or pretreatment arteriosclerotic changes in the optic fundi. However, multiple logistic regression analysis revealed that complication by hypertension was observed with a high frequency in the group with retinopathy ($P = 0.004$, OR = 245.918, 95% CI = 5.6-10786.2).

INTRODUCTION

Chronic hepatitis C, which affects more than 170 million people in the world^[1], may eventually lead to cirrhosis and/or hepatocellular carcinoma. The main treatment for this intractable disease is interferon administration. Published guidelines recommend interferon-ribavirin combination therapy as a first-line treatment^[2]. Interferon is also used in the treatment of other viral and neoplastic diseases.

Various adverse effects have been reported due to use of interferon^[3]. An influenza-like syndrome, characterized by fever, chills, myalgias, arthralgias, and headache, is the most common adverse effect. Toxicities of the central nervous, hematopoietic, gastrointestinal, urinary, cardiovascular, musculoskeletal and endocrine systems have also been described. However, ocular toxicity was not reported before the use of interferon for chronic hepatitis^[4].

After the introduction of interferon for the treatment of hepatitis, retinal complications have been reported. Hayakawa *et al* showed that 17 of 43 patients developed retinopathy during interferon monotherapy. They also showed that the prevalence of retinopathy was higher in patients with diabetes^[4]. Subsequently, several papers have shown that a substantial proportion of patients undergoing interferon monotherapy develop retinopathy^[5-7]. However, the prevalence of retinopathy is variable, which is