



WJG

World Journal of Gastroenterology®

Indexed and Abstracted in:

Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®) and Journal Citation Reports/Science Edition, *Index Medicus*, MEDLINE and PubMed, Chemical Abstracts, EMBASE/Excerpta Medica, Abstracts Journals, *Nature Clinical Practice Gastroenterology and Hepatology*, CAB Abstracts and Global Health.
ISI JCR 2003-2000 IF: 3.318, 2.532, 1.445 and 0.993.

Volume 14 Number 13

April 7, 2008

World J Gastroenterol

2008 April 7; 14(13): 1969-2132

Online Submissions

wjg.wjgnet.com

www.wjgnet.com

Printed on Acid-free Paper

世界胃肠病学杂志

A Weekly Journal of Gastroenterology and Hepatology



National Journal Award
2005

World Journal of Gastroenterology®

Weekly Established in October 1995

Volume 14 Number 13

April 7, 2008



百世登
Baishideng™

Contents

EDITORIAL	1969	Wireless capsule endoscopy <i>Mata A, Lluch J, Bordas JM</i>
	1972	Current view of the immunopathogenesis in inflammatory bowel disease and its implications for therapy <i>Torres MI, Rios A</i>
REVIEW	1981	Role of endoscopy in the management of acute diverticular bleeding <i>Philichos C, Bobotis E</i>
TOPIC HIGHLIGHT	1984	Endoscopic mucosal resection in the upper gastrointestinal tract <i>Ahmadi A, Drozanov P</i>
GASTRIC CANCER	1990	Relationship between cell adhesion molecules expression and the biological behavior of gastric carcinoma <i>Chu YQ, Ye ZY, Tao HQ, Wang YY, Zhao ZS</i>
LIVER CANCER	1997	Long-term outcome of percutaneous ethanol injection therapy for minimum-sized hepatocellular carcinoma <i>Taniguchi M, Kim SR, Imoto S, Ikawa H, Ando K, Mita K, Fuki S, Sasase N, Matsuoaka T, Kudo M, Hayashi Y</i>
	2003	Anti-cancer and anti-angiogenic effects of curcumin and tetrahydrocurcumin on implanted hepatocellular carcinoma in nude mice <i>Yoyungnoen P, Wirachwong P, Changtam C, Suksamrarn A, Patumraj S</i>
VIRAL HEPATITIS	2010	Serial changes in expression of functionally clustered genes in progression of liver fibrosis in hepatitis C patients <i>Takahara Y, Takahashi M, Zhang QW, Wagatsuma H, Mori M, Tamori A, Shiomi S, Nishiguchi S</i>
BASIC RESEARCH	2023	Pancreatic stellate cells promote proliferation and invasiveness of human pancreatic cancer cells <i>via</i> galectin-3 <i>Jiang HB, Xu M, Wang XP</i>
CLINICAL RESEARCH	2029	Probiotic intervention has strain-specific anti-inflammatory effects in healthy adults <i>Kekkonen RA, Lunnela N, Karjalainen H, Latvala S, Tynkkyinen S, Järvenpää S, Kuittinen H, Julkunen I, Vapaatalo H, Korpela R</i>
RAPID COMMUNICATION	2037	A combination therapy of ethanol injection and radiofrequency ablation under general anesthesia for the treatment of hepatocellular carcinoma <i>Kurokohchi K, Watanabe S, Yoneyama H, Deguchi A, Masaki T, Himoto T, Miyoshi H, Mohammad HIS, Kitamuka A, Taminato T, Kuriyama S</i>

- 2044 Serum type IV collagen level is predictive for esophageal varices in patients with severe alcoholic disease
Mumori S, Searashi Y, Matsushima M, Hashimoto K, Uetake S, Matsudaira H, Ito S, Nakajima H, Tajiri H
- 2049 Early effects of Lansoprazole orally disintegrating tablets on intragastric pH in CYP2C19 extensive metabolizers
Yamagishi H, Koike T, Ohara S, Horii T, Kikuchi R, Kobayashi S, Abe Y, Iijima K, Imatani A, Suzuki K, Hishinuma T, Goto J, Shimosegawa T
- 2055 p16 promoter hypermethylation: A useful serum marker for early detection of gastric cancer
Abbaszadegan MR, Moaven O, Sima HR, Ghafarzadegan K, Arabi A, Forghani MN, Raziee HR, Mashhadnejad A, Jafarzadeh M, Esmaili-Shandi: E, Dadkhah E
- 2061 Prospective evaluation of small bowel preparation with bisacodyl and sodium phosphate for capsule endoscopy
Franke A, Hummel F, Knebel P, Antoni C, Böcker U, Singer MV, Löhr M
- 2065 Effect of Prometheus liver assist system on systemic hemodynamics in patients with cirrhosis: A randomized controlled trial study
Deitloff T, Tofteng F, Frederiksen HJ, Hojskov M, Hansen BA, Larsen FS
- 2072 Ultrasonography in differentiation between chronic viral hepatitis and compensated early stage cirrhosis
Iliopoulos P, Vlychou M, Karatza C, Yarmenitis SD, Repanti M, Tsamis I, Tepetes K
- 2080 Endoscopic band ligation and endoscopic hemoclip placement for patients with Mallory-Weiss syndrome and active bleeding
Cho YS, Chae HS, Kim HK, Kim JS, Kim BW, Kim SS, Han SW, Choi KY
- 2085 Effects of honey as a scolicidal agent on the hepatobiliary system
Kilicoglu B, Kismet K, Kilicoglu SS, Erel S, Gencay O, Sorkun K, Erdemli E, Akhan O, Akkus MA, Sayek I
- 2089 Predictive factors for early aspiration in liver abscess
Khan R, Hamid S, Abid S, Jafri W, Abbas Z, Islam M, Shah H, Beg S
- 2094 Enhancement of CD4⁺ T cell activities and modulation of Th1/Th2 lineage development in irradiated tumor-bearing rats treated with male zooid of *Antheraea pernyi* extracts
Zhao WH, Li L, Zhang B, Zhang WD, Zeng M, Tang JD, Zhang HY, Li S
- 2100 Effect of Oxymatrine on the TGFbeta-Smad signaling pathway in rats with CCl₄-induced hepatic fibrosis
Wu XL, Zeng WZ, Jiang MD, Qin JP, Xu H
- 2106 Intraperitoneal administration of gonadotropin-releasing hormone-PE40 induces castration in male rats
Yu L, Zhang ZF, Jing CX, Wu FL
- 2110 Expression of connective tissue growth factor in tumor tissues is an independent predictor of poor prognosis in patients with gastric cancer
Lin LY, Han YC, Wu SH, Lv ZH

CASE REPORT

- 2115 Endoscopic ultrasonography-guided trucut biopsy for the preoperative diagnosis of peripancreatic castleman's disease: A case report
Rhee KH, Lee SS, Huh JR

Contents

	2118	Primary rectal signet ring cell carcinoma with peritoneal dissemination and gastric secondaries <i>Sim HL, Tan KY, Poon PL, Cheng A</i>
	2121	Diagnosis and treatment of Gardner syndrome with gastric polyposis: A case report and review of the literature <i>Gu GL, Wang SL, Wei XM, Bai L</i>
BOOK REVIEW	2124	Rome III: The functional gastrointestinal disorders, third edition, 2006 <i>Mostafa R</i>
LETTERS TO THE EDITOR	2126	Role of <i>ABCC2</i> common variants in intrahepatic cholestasis of pregnancy <i>Sookkian S, Castaño G, Pirota CJ</i>
ACKNOWLEDGMENTS	2128	Acknowledgments to Reviewers of <i>World Journal of Gastroenterology</i>
APPENDIX	2129	Meetings
	2130	Instructions to authors
FLYLEAF	I-V	Editorial Board
INSIDE BACK COVER		Online Submissions
INSIDE FRONT COVER		Online Submissions

RESPONSIBLE EDITORS FOR THIS ISSUE

Assistant Editor: *Yao Jiang* Review Editor: *Jing Zhu* Electronic Page Editor: *Ying Liu*
Editor-in-Charge: *Jian-Xu Chang* Copy Editor: *Piero Invernizzi* MD PhD Associate Senior Editor: *Yi Liu*
Layout Editor: *Lian-Sheng Ma*

<p>NAME OF JOURNAL <i>World Journal of Gastroenterology</i></p> <p>RESPONSIBLE INSTITUTION Department of Science and Technology of Shanxi Province</p> <p>SPONSOR Taiyuan Research and Treatment Center for Digestive Diseases, 77 Shuangta Xijie, Taiyuan 030001, Shanxi Province, China</p> <p>EDITING Editorial Board of <i>World Journal of Gastroenterology</i>, 77 Shuangta Xijie, Taiyuan 030001, Shanxi Province, China Telephone: +86-351-4078656 E-mail: wjg@wjgnet.com</p> <p>PUBLISHING Editorial Department of <i>World Journal of Gastroenterology</i>, 77 Shuangta Xijie, Taiyuan 030001, Shanxi Province, China Telephone: +86-351-4078656 E-mail: wjg@wjgnet.com http://www.wjgnet.com</p> <p>PRINTING Beijing Kezhan Printing House</p> <p>OVERSEAS DISTRIBUTOR Beijing Bureau for Distribution of Newspapers and Journals (Code No. 82-261) China International Book Trading Corporation PO Box 399, Beijing, China (Code No. M4481)</p> <p>PUBLICATION DATE April 7, 2008</p> <p>EDITOR-IN-CHIEF Lian-Sheng Ma, Taiyuan</p>	<p>SUBSCRIPTION RMB 50 Yuan for each issue, RMB 2400 Yuan for one year</p> <p>CSSN ISSN 1007-9327 CN 14-1219/R</p> <p>HONORARY EDITORS-IN-CHIEF Ke-Ji Chen, Beijing Li-Fang Chou, Taipei Zhi-Qiang Huang, Beijing Shun-Jiang Huang, Taipei Min-Liang Kuo, Taipei Nicholas F LaRusso, Rochester Jie-Shou Li, Nanjing Geng-Tao Liu, Beijing Lein-Ray Mo, Taiwan Bo-Rong Pan, Xi'an Fu-Zu Qiu, W'nan Eamonn M Quigley, Cork David S Rampton, London Rudi Schmid, Basel Nicholas J Talley, Rochester Guido NJ Tytgat, Amsterdam H-P Wang, Taipei Jaw-Ching Wu, Taipei Meng-Chao Wu, Shanghai Ming-Shiang Wu, Taipei Jia-Yu Xu, Shanghai Ts-Sen Yeh, Taiyuan</p> <p>ASSOCIATE EDITORS-IN-CHIEF Gianfranco D Alpini, Trieste Bruno Annibale, Roma Roger William Chapman, Oxford Chi-Hin Cho, Hong Kong Alexander L Gerbes, Mainz Shou Dong Lee, Taipei Walter Edwin Longo, New Haven You-Yong Lu, Beijing Masao Omata, Tokyo Henry HX Xia, Hainan</p> <p>EXECUTIVE VICE DIRECTOR Ye Li, Beijing</p>	<p>DEPUTY DIRECTOR Jian-Zhong Zhang, Beijing</p> <p>LANGUAGE EDITORS Director: Jing-Yun Ma, Beijing Deputy Director: Xian-Lin Wang, Beijing</p> <p>MEMBERS Gianfranco D Alpini, Trieste BS Anand, Houston Richard B Bezzi, Padova Giuseppe Chionigi, Volterra John Frank Di Mari, Texas Shannon S Glaser, Tampa Mario Gualandi, Milano Martin Hensenberg, Bonn Atif Iqbal, Omaha Manoj Kumar, Nagpur Patricia F Lalor, Birmingham Ming Li, New Orleans Margaret Lutzke, Chicago Jing-Yun Ma, Beijing Daniel Markovich, Bratislava Sabine Mehr, Göttingen Francesco Negro, Gendin Bernardino Rampone, Siena Richard A Rippe, Chapel Hill Stephen E Roberts, Swansea Ross C Smith, Sydney Seng-Lai Tan, Seattle Xian-Lin Wang, Beijing Eddie Wise, Karlsruhe Daniel Lanslay Worthley, Bedford</p> <p>NEWS EDITOR Lian Zhu, Berkeley</p> <p>COPY EDITORS Gianfranco D Alpini, Trieste Sujit Kumar Bhattacharya, Kolkata Filip Bract, Sydney Kirsteen N Browning, Baten Range Radha K Dhirman, Chandigarh John Frank Di Mari, Texas Shannon S Glaser, Tampa</p>	<p>Martin Hensenberg, Bonn Eberhard Huidt, Berlin Patricia F Lalor, Birmingham Ming Li, New Orleans Margaret Lutzke, Chicago MI Torres, John Sri Prakash Murti, Allahabad Giovanni Monteleone, Rome Giovanni Musso, Torino Valerio Nobili, Rome Osman Cavit Ozdogan, Istanbul Francesco Perri, San Gimignano, Firenze Thierry Piche, Nice Bernardino Rampone, Siena Richard A Rippe, Chapel Hill Ross C Smith, Sydney Daniel Lanslay Worthley, Bedford George Y Wu, Farmington Jian Wu, Sacramento</p> <p>COPYRIGHT © 2008 Published by WJG. All rights reserved; no part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise without the prior permission of WJG. Authors are required to grant WJG an exclusive licence to publish.</p> <p>SPECIAL STATEMENT All articles published in this journal represent the viewpoints of the authors except where indicated otherwise.</p> <p>INSTRUCTIONS TO AUTHORS Full instructions are available online at http://www.wjgnet.com/wjg/help/instructions.jsp. If you do not have web access please contact the editorial office.</p> <p>ONLINE SUBMISSION http://wjg.wjgnet.com</p>
--	--	--	--

ISI journal citation reports

2003 – GASTROENTEROLOGY AND HEPATOLOGY

Rank	Abbreviated journal title	ISSN	2003 Total cites	Impact factor	Immediacy index	2003 Articles	Cited Half-life
1	GASTROENTEROLOGY	0016-5085	46 174	12.718	2.810	316	6.9
2	HEPATOLOGY	0270-9139	30 844	9.503	1.574	296	5.5
3	SEMIN LIVER DIS	0272-8087	2 524	6.524	0.810	42	4.7
4	GUT	0017-5749	20 612	5.883	1.147	307	6.5
5	J HEPATOL	0168-8278	11 111	5.283	0.879	281	5.0
6	LIVER TRANSPLANT	1527-6465	2 944	4.242	0.621	211	2.9
7	AM J GASTROENTEROL	0002-9270	17 923	4.172	0.678	391	5.1
8	ALIMENT PHARM THERAP	0269-2813	60 32	3.529	0.573	372	3.9
9	AM J PHYSIOL-GASTR L	0193-1857	11 047	3.421	0.634	268	5.9
10	GASTROINTEST ENDOSC	0016-5107	10 047	3.328	1.798	331	4.8
11	WORLD J GASTROENTERO	1007-9327	2 387	3.318	0.345	632	2.4
12	J VIRAL HEPATITIS	1352-0504	1 320	3.258	0.386	70	4.0
13	ENDOSCOPY	0013-726X	4 238	3.227	0.349	152	5.5
14	INFLAMM BOWEL DIS	1078-0998	1 278	3.023	0.625	40	3.8
15	HELICOBACTER	1083-4389	798	2.624	0.426	68	3.6
16	NEUROGASTROENT MOTIL	1350-1925	848	2.500	0.333	66	3.7
17	DIS COLON RECTUM	0012-3706	8 130	2.343	0.191	256	7.6
18	SCAND J GASTROENTERO	0036-5521	7 271	2.140	0.118	238	7.6
19	LIVER	0106-9543	1 337	2.076		0	5.4
20	J GASTROINTEST SURG	1091-255X	1 236	1.881	0.430	142	3.6
21	PANCREAS	0885-3177	2 271	1.855	0.364	140	5.8
22	INT J COLORECTAL DIS	0179-1958	1 167	1.848	0.373	83	6.4
23	GASTROENTEROL CLIN N	0889-8553	1 517	1.684	0.019	54	6.7
24	PANCREATOLOGY	1424-3903	240	1.596	0.115	52	2.2
25	EUR J GASTROEN HEPAT	0954-691X	3 163	1.578	0.367	177	4.6
26	J CLIN GASTROENTEROL	0192-0790	3 215	1.564	0.484	155	7.2
27	J GASTROEN HEPATOL	0815-9319	3 171	1.530	0.254	193	4.5
28	DIGEST LIVER DIS	1590-8658	637	1.463	0.151	159	2.5
29	J PEDIATR GASTR NUTR	0277-2116	3 963	1.402	0.257	183	6.1
30	DIGESTION	0012-2823	2 193	1.399	0.091	55	7.7
31	DIGEST DIS SCI	0163-2116	9 149	1.387	0.144	347	7.9
32	CAN J GASTROENTEROL	0835-7900	813	1.265	0.134	82	3.9
33	J GASTROENTEROL	0944-1174	1 625	1.179	0.319	191	4.5
34	DIGEST DIS	0257-2753	534	1.151	0.031	32	5.7
35	Z GASTROENTEROL	0044-2771	918	1.076	0.112	116	5.4
36	ABDOM IMAGING	0942-8925	1 190	0.996	0.194	124	5.0
37	BEST PRACT RES CL GA	1521-6918	342	0.992	0.000	66	3.4
38	HEPATOL RES	1386-6346	406	0.991	0.168	155	2.6
39	GASTROEN CLIN BIOL	0399-8320	1 497	0.884	0.235	153	6.3
40	HEPATO-GASTROENTEROL	0172-6390	3 886	0.837	0.044	544	5.0
41	DIS ESOPHAGUS	1120-8694	354	0.809	0.027	75	4.7
42	ACTA GASTRO-ENT BELG	0001-5644	345	0.670	0.065	31	5.2
43	DIGEST SURG	0253-4886	548	0.619	0.114	79	4.2
44	CURR OPIN GASTROEN	0267-1379	300	0.598	0.050	60	3.9
45	REV ESP ENFERM DIG	1130-0108	255	0.348	0.211	57	6.0
46	CI IIR GASTROENTEROL	0177-9990	74	0.157	0.039	76	
47	LIVER INT	1478-3231	4		0.053	75	

Serial changes in expression of functionally clustered genes in progression of liver fibrosis in hepatitis C patients

Yoshiyuki Takahara, Mitsuo Takahashi, Qing-Wei Zhang, Hiroataka Wagatsuma, Maiko Mori, Akihiro Tamori, Susumu Shiomi, Shuhei Nishiguchi

Yoshiyuki Takahara, Mitsuo Takahashi, Qing-Wei Zhang, Hiroataka Wagatsuma, Maiko Mori, Exploratory & Applied Pharmaceutical Research Department, Pharmaceutical Company, Ajinomoto Co., Inc., 1-1 Suzuki-cho, Kawasaki-ku, Kawasaki 210-8681, Japan

Akihiro Tamori, Susumu Shiomi, Department of Nuclear Medicine, Graduate School of Medicine, Osaka City University, Osaka, Japan

Shuhei Nishiguchi, Division of Hepatobiliary and Pancreatic Diseases, Department of Internal Medicine, Hyogo College of Medicine, Nishinomiya, Japan

Correspondence to: Yoshiyuki Takahara, Exploratory & Applied Pharmaceutical Research Department, Pharmaceutical Company, Ajinomoto Co., Inc., 1-1 Suzuki-cho, Kawasaki-ku, Kawasaki 210-8681, Japan. yoshiyukitakahara@gmail.com
Telephone: +81-44-2105822

Received: April 23, 2007 Revised: July 18, 2007

metabolism showed altered expression in the late phase of fibrosis (F3/F4). Therefore, molecular networks showing serial changes in gene expression are present in liver fibrosis progression in hepatitis C patients.

CONCLUSION: Analysis of gene expression profiles from a perspective of functional categories or molecular networks provides an understanding of disease and suggests new diagnostic methods. Selected marker genes have potential utility for biological identification of advanced fibrosis.

© 2008 WJG. All rights reserved.

Key words: Hepatitis C; Liver fibrosis; Marker gene; Gene expression; RT-PCR; Molecular network; Metabolism; Transcription factor; Diagnosis

Peer reviewer: Richard A Rippe, Dr, Department of Medicine, The University of North Carolina at Chapel Hill, Chapel Hill, NC 27599-7038, United States

Takahara Y, Takahashi M, Zhang QW, Wagatsuma H, Mori M, Tamori A, Shiomi S, Nishiguchi S. Serial changes in expression of functionally clustered genes in progression of liver fibrosis in hepatitis C patients. *World J Gastroenterol* 2008; 14(13): 2010-2022
Available from: URL: <http://www.wjnet.com/1007-9327/14/2010.asp>
DOI: <http://dx.doi.org/10.3748/wjg.14.2010>

Abstract

AIM: To investigate the relationship of changes in expression of marker genes in functional categories or molecular networks comprising one functional category or multiple categories in progression of hepatic fibrosis in hepatitis C (HCV) patients.

METHODS: Marker genes were initially identified using DNA microarray data from a rat liver fibrosis model. The expression level of each fibrosis associated marker gene was analyzed using reverse transcription-polymerase chain reaction (RT-PCR) in clinical biopsy specimens from HCV-positive patients ($n = 61$). Analysis of changes in expression patterns and interactions of marker genes in functional categories was used to assess the biological mechanism of fibrosis.

RESULTS: The profile data showed several biological changes associated with progression of hepatic fibrosis. Clustered genes in functional categories showed sequential changes in expression. Several sets of clustered genes, including those related to the extracellular matrix (ECM), inflammation, lipid metabolism, steroid metabolism, and some transcription factors important for hepatic biology showed expression changes in the immediate early phase (F1/F2) of fibrosis. Genes associated with aromatic amino acid (AA) metabolism, sulfur-containing AA metabolism and insulin/Wnt signaling showed expression changes in the middle phase (F2/F3), and some genes related to glucose

INTRODUCTION

Liver fibrosis is caused by liver disorders such as hepatitis C, hepatitis B, alcoholic hepatitis and non-alcoholic hepatitis. Fibrosis progresses gradually and finally disrupts liver structure and function over several decades, leading to fatal diseases such as cirrhosis and hepatocellular carcinoma (HCC). Classification of fibrosis progression is usually based on histological criteria using the METAVIR scoring system^[1], which includes five stages: F0 (no fibrosis), F1, F2, F3, and F4 (cirrhosis). Such a classification is essential in decisions regarding treatment of liver fibrosis. Prominent subjective symptoms do not occur from F1 to F3, but patients begin to be aware of symptoms after F4. However, the F4 stage of cirrhosis is almost incurable and diagnosis of fibrosis at an earlier stage is desirable. The biology after F4 (cirrhosis) has been well studied, due to the interest in diagnosis and therapy for hepatocellular carcinoma (HCC).

but progression to HCC may begin in the early stage of fibrosis in hepatitis C.¹²

Prevention of HCC and inhibition of fibrosis in the early phase is important, but the detailed hepatic biological changes corresponding to the F stage are unclear. To understand the background of the early stage of fibrosis, we previously identified genes that can be used as markers of biological changes in progression of hepatic fibrosis and diagnosis of fibrotic progression; these data were obtained from DNA microarray data from an experimental DMN (dimethylnitrosamine)-treated rat model of hepatic fibrosis.¹³ This work led to marker genes that were arranged in functional categories related to fibrosis, based on genes associated with hepatic cell types such as Kupffer cells, hepatic stellate cells, and hepatocytes.

These marker genes give information on cell-specific and time-dependent behavior of each hepatic cell in fibrogenesis. In the current work, the behavior of marker genes associated with a particular F stage was analyzed using RT-PCR in clinical biopsy specimens from hepatitis C patients. This profile data revealed several biological changes in progression of hepatic fibrosis. Since many functionally clustered genes showed similar changes in expression, we propose serial expression changes in molecular networks associated with liver fibrosis progression in hepatitis C patients. Many functionally clustered genes showed large changes in expression in the early stage of fibrosis, suggesting the importance of therapy at this early stage. Alteration of gene expression also suggested qualitative changes in biological status in the transition from F3 to F4, which is a suspected risk factor for development of HCC. We conclude that analysis of gene expression profiles from a perspective of molecular networks provides improved understanding of disease and indicates potential methods of diagnosis.

MATERIALS AND METHODS

Patients in the clinical study

All patients were recruited from the Osaka City University Hospital (Osaka, Japan). Sixty-one patients with seropositive results in diagnosis using the third-generation hepatitis C virus enzyme-linked immunosorbent assay (Lumipulse II Ortho HCV, Ortho-Clinical Diagnostics, Tokyo, Japan) and positive serum HCV-RNA were included in the study. Informed consent was obtained from all patients. Liver biopsies were performed on all patients enrolled in the study, and the histological features of the liver specimens were analyzed and graded using the METAVIR scoring system¹⁴. The liver fibrotic stage (F stage) and inflammatory activity (A grade) were determined histologically; at least four subjects were found to be in each F stage classification. Determination for chymase I exceptionally has been done with three subjects in F4 stage due to the lack of appropriate samples.

Part of the biopsy sample from each patient was immediately immersed in RNAlater (QIAGEN, The Netherlands) to inhibit RNAase and then kept at 4°C overnight before being transferred to another tube and frozen at -80°C.

RT-PCR analysis

Total RNA was extracted from liver biopsy samples using

an ISOGEN kit (Nippongene) and reverse transcribed using a High Capacity cDNA Archive Kit (ABI, Foster City, CA), in each case according to the manufacturer's instructions. The total RNA in the final reaction mixture was 10 ng/ μ L. Real-time PCR was performed on an Applied Biosystems 7500 Real-Time PCR System (ABI) data collection system, and analyses were performed using the accompanying software. RT-PCR was performed using 0.8 μ L cDNA in each well, with a final concentration of 1X the probes of the TaqMan[®] Gene Expression Assay and 1X the Taqman Universal PCR Master Mix (ABI). The final reaction volume was 20 μ L. Each sample was analyzed in duplicate. The thermal cycler conditions were 2 min at 50°C and 10 min at 95°C, followed by 40 cycles of 15 s at 95°C and 1 min at 60°C.

Data were analyzed using the comparative CT method, in which the expression level of a target gene is normalized relative to an endogenous reference. GAPDH was used as the endogenous reference in all experiments. The target Ct and endogenous control Ct were calculated for each sample, and the target gene expression level was then calculated using the formula $2^{(4-C_T)}$. The average of duplicate measurements was obtained, and the relative expression of each gene in a sample was calculated by setting the expression of GAPDH equal to 1000. PCR fluorogenic probes for all the target genes and the endogenous reference were purchased as TaqMan[®] Gene Expression Assays (ABI).

Statistical analysis

A Kruskal-Wallis test was applied to select marker genes with statistically significant changes ($P < 0.05$) in expression level during the fibrosis progression. This calculation was performed using SPSS (SPSS Inc., IL, USA). The gene expression data were subjected to hierarchical clustering analysis using Genowiz[™] software (Ocimum Biosolutions).

Pathway analysis of PCR data

The behavior and relationships of marker genes in pathways associated with lipid metabolism were analyzed with bioSpace Explorer, a system for analysis of expression profile data. This system was produced collaboratively by Pharmafrontier Co. Ltd. and World Fusion Co. Ltd. to examine molecular interactions in expression profile data, using both manual and computational text mining.

RESULTS

Expression behavior of marker genes

All marker genes determined in this study are listed in Table 1. For each gene, since the quantitative limitation of biopsy specimens resulted in a difference in sample number for each probe, the type of samples, indicating the number of biopsy specimens, is listed in Table 2. Marker genes were selected as representative members of functional categories or molecular networks based on their expression changes in an experimental hepatic fibrosis model¹⁵. Marker genes with statistically significant changes in expression are listed in Table 1. Genes that showed statistically insignificant changes in expression are listed with the gene name only in Table 1. Marker genes that showed statistically significant changes by *t* test during a

Table 1 Expression profiles of marker genes with statistically significant changes during fibrosis progression

Functional category	Description	Gene name	Expression at F1	Group	Expression ratio			Serial	Type	
					F2/F1	F3/F2	F4/F3			
ECM (or other HSC marker)	Decorin	DCN	217.6	2	1.2	1.1	1.1	1	1	
	Matrix metalloproteinase 2	MMP2	11.7	2	1.5	1.2	0.9	2	1	
	Hyaluronan-mediated motility receptor	HMMR	1.2	2	2.6	0.6	1.1	3	1	
	Lysyl oxidase	LOX	1.0	2	1.4	1.2	0.9	4	1	
	Lysyl oxidase-like 1	LOXL1	0.8	2	1.8	1.4	1.0	5	1	
	Tropoelastin 1	TPM1	35.0	2	1.8	1.5	1.1	6	1	
	Prion	PRNP	19.4	3	1.2	0.7	1.0	7	1	
	Collagen, type I, alpha 1	COL1A1	16.2	2	1.3	2.2	0.6	8	1	
	Collagen type III alpha 1	COL3A1	143.9	2	1.3	1.5	0.8	9	1	
	Collagen type alpha 1	COL4A1	16.8	2	1.3	1.9	0.6	10	1	
	Humican	LUM	22.8	2	1.2	2.0	0.9	11	1	
	Staloprotein	SPP1	13.2	2	1.2	2.2	1.4	12	1	
	Glypican 3	GPC3	23.5	2	1.8	2.5	1.0	13	1	
	Proline 4-hydroxylase, alpha polypeptide I	P4HA1	34.0	1	1.1	0.4	1.2	14	1	
	Insignificant change: MGP, BGN, TAGLN, LGALS1, EDG2, EDG5, TNNT2									
	Inflammation (or apoptosis)	Lysozyme	LYZ	185.3	2	2.1	0.8	1.0	15	2
		TGF beta	TGFB1	51.9	3	1.3	0.8	0.9	16	1
		TGF beta 3	TGFB3	3.3	3	1.4	0.9	0.9	17	3
		TNF	TNF	2.6	3	1.7	0.6	1.2	18	3
		Natural killer cell proteinase 1	GZMB	1.7	3	1.6	0.5	1.1	19	1
IL1 beta		IL1B	2.6	3	1.5	0.7	1.1	20	3	
Hemopoietic cell kinase		HCK	22.4	3	1.3	0.6	0.9	21	4	
Interleukin 6 receptor		IL6R	142.9	1	0.9	0.7	1.0	22	3	
BCL2-related ovarian killer		BOK	205.9	1	0.9	0.7	1.1	23	5	
Caspase 2		CASP2	1.3	1	1.2	0.7	0.8	24	5	
Chymase 1, mast cell		CMA1	0.2	2	0.7	1.8	2.2	25	6	
Insignificant change: LTBP1, LBP, TNFRSF1B, DEFB1, IL1RN, S100A8, BRIC3, CARD12, CASP1, CASP4, CASP8, PAWR, CD19, CD32, MS4A1, CD37, TRA*										
Growth factor		Growth hormone receptor	GHR	102.3	1	0.8	0.9	0.9	26	4
		IGF1	IGF1	56.5	1	0.9	0.7	1.1	27	1
Insignificant change: PTN, PST, PRLR										
Insulin/ Wnt signal	Cyclin D1	CCND1	190.6	2	1.7	1.0	1.0	28	5	
	Forkhead box M1	FOXM1	0.6	3	3.7	0.8	1.1	29	4	
	Gap junction protein, alpha 1, 43 kDa (connexin 43)	GJA1	4.0	3	2.4	0.5	1.0	30	5	
	V-akt murine thymoma viral oncogene homolog 1	AKT1	77.6	1	0.9	0.9	1.0	31	4	
	V-akt murine thymoma viral oncogene homolog 2	AKT2	59.6	1	0.9	0.7	1.0	32	7	
	Catenin (cadherin-associated protein), beta 1, 86 kDa	CTNNB1	159.0	1	1.1	0.7	0.8	33	5	
	Catenin, beta interacting protein 1	CTNNBIP1	24.0	1	1.1	0.7	0.8	34	5	
	Glycogen synthase kinase 3 beta	GSK3B	44.1	1	1.1	0.8	0.8	35	5	
	Dishevelled, dsh homolog 1 (Drosophila)	DVL1	12.2	1	1.1	0.7	0.9	36	7	
	Membrane-bound transcription factor peptidase, site 1	MBTPS1	72.4	1	0.9	0.7	1.0	37	7	
	Membrane-bound transcription factor peptidase, site 2	MBTPS2	13.2	1	0.9	0.7	1.0	38	7	
	Tribbles homolog 3 (Drosophila)	TRUB3	18.4	1	1.1	0.9	0.4	39	7	
	Insignificant change: GSK3A, INSIG1, INSIG2, PRKCB1, PRKCD									
	Others signal	Regucalcin (senescence marker protein-30)	RGN	697.8	1	0.9	0.8	0.8	56	4
	Insignificant change: DAB2, PMP22, S100A10, LCN2									
	Transcription factors	CCAAT/enhancer binding protein (C/EBP), alpha	CEBPA	352.2	1	0.7	1.2	0.7	40	4
		Retinoid X receptor, alpha	RXRA	409.6	1	0.7	1.0	0.8	41	4
		Hepatocyte nuclear factor 4, alpha	HNF4A	708.7	1	0.7	1.1	0.9	42	4
Transcription factor 1 (HNF1)		TCF1	26.7	1	0.8	0.9	0.8	43	4	
Nuclear receptor subfamily 0, group B, member 2		NR0B2	166.7	1	0.7	0.9	0.8	44	4	
Peroxisome proliferative activated receptor, alpha		PPARA	62.9	1	0.7	0.8	0.9	45	4	
Inhibitor of DNA binding 1 (splice variation)		ID1	463.4	3	1.6	0.5	1.4	46	1	
AE binding protein 1		AEBP1	24.6	2	1.4	1.2	1.0	47	4	
Nuclear receptor subfamily 1, group H, member 2		NR1H2 (LXRβ)	9.0	1	1.2	0.7	1.0	48	7	
Nuclear receptor subfamily 1, group H, member 3		NR1H3 (LXRA)	27.8	1	1.0	0.7	0.8	49	7	
Nuclear receptor subfamily 1, group H, member 4		NR1H4 (FXR)	107.0	1	1.0	0.7	0.9	50	7	
c/EBPbeta		CEBPB	274.3	1	0.9	0.7	0.9	51	2	
Upstream transcription factor 2, c-fos interacting		USF2	323.7	1	0.9	0.7	0.9	52	4	
Estrogen-related receptor alpha		ESRRA	79.6	1	1.2	0.6	0.8	53	7	
C-met		MET	73.5	1	1.0	0.7	1.1	54	3	
Upstream transcription factor 1		USF1	30.8	1	1.0	0.9	0.8	55	4	
Insignificant change: ONECUT1, JUNB, NR3C1, PFARG, PPARGC1B, PPARGC1A, SREBF2, FHL2										
Transporter		Solute carrier family 6, member 6	SLC6A6	3.7	3	2.5	0.6	1.0	57	5
	Solute carrier family 7, member 1	SLC7A1	0.9	3	2.3	0.7	1.3	58	5	
	Solute carrier family 38, member 2 Alanine-transporter	SLC38A2	68.6	3	1.3	0.6	1.0	59	5	
	Solute carrier family 25 member 15	SLC25A15	85.7	1	1.1	0.7	0.9	60	5	
	Solute carrier family 7, member 7	SLC7A7	7.1	1	1.1	0.6	1.0	61	5	
Solute carrier family 17 (sodium phosphate), member 1	SLC17A1	23.6	1	0.9	0.7	1.0	62	4		

		Insignificant change: SLC38A3, ABCB1, SLC15A4								
Redox	Catalase	CAT	1977.7	1	0.8	0.9	1.0	63	4	
	Paraoxonase 1	PON1	191.1	1	0.8	0.7	0.9	64	4	
Blood coagulation	Coagulation factor X	F10	187.5	1	0.8	1.0	0.9	65	4	
	Angiotensinogen	AGT	1958.4	1	0.8	1.0	1.0	66	4	
	Fibrinogen, A alpha polypeptide	FGA	8337.5	1	0.6	1.0	0.9	67	4	
	Plasminogen	PLG	6156.1	1	0.8	0.8	1.0	68	4	
	Plai 1	SERPINE1	11.6	3	0.9	2.9	0.5	69	1	
Lipid metabolism	Acyl-Coenzyme A oxidase 2, branched chain	ACOX2	122.2	1	0.8	0.9	0.9	70	4	
	L-3-hydroxyacyl-Coenzyme A dehydrogenase, short chain	HADHSC	139.8	1	0.8	0.9	0.9	71	4	
	Acyl-CoA synthetase long-chain family member 1	ACSL1	2800.8	1	0.8	0.9	0.9	72	4	
	Acyl-Coenzyme A oxidase 1, palmitoyl	ACOX1	357.7	1	0.7	0.9	1.0	73	4	
	Carnitine O-octanoyltransferase	CROT	10.3	1	0.7	0.8	1.1	74	4	
	2,4-dienoyl CoA reductase 2, peroxisomal	DECR2	135.2	1	0.8	0.9	0.9	75	4	
	Acetyl-Coenzyme A acyltransferase 2	ACAA2	1167.7	1	0.8	0.8	1.1	76	4	
	Acetyl-Coenzyme A acetyltransferase 1	ACAT1	714.2	1	0.7	1.1	0.8	77	4	
	Acyl-CoA synthetase long-chain family member 5	ACSL5	57.3	1	0.8	0.9	1.0	78	4	
	Dodecenoyl-Coenzyme A delta isomerase	DCI	327.2	1	0.9	0.8	0.9	79	4	
	Enoyl Coenzyme A hydratase, short chain, 1, mitochondrial	ECH51	324.3	1	0.7	1.0	0.9	80	4	
	Hydroxyacyl-Coenzyme A dehydrogenase, type II	HADH2	422.5	1	0.7	0.9	0.9	81	4	
	Hydroxyacyl-Coenzyme A dehydrogenase, beta subunit	HADHB	398.8	1	0.8	0.9	1.1	82	4	
	Lipase, hepatic	LIPC	561.5	1	0.5	1.5	0.9	83	4	
	Hydroxyacyl-Coenzyme A dehydrogenase, alpha subunit	HADHA	103.2	1	0.9	0.9	1.0	84	4	
	Palmitoyl-protein thioesterase 1	PPT1	82.1	1	1.0	0.8	0.8	85	4	
	Fatty acid synthase	FASN	105.3	3	0.8	1.8	0.6	86	4	
	Peroxisomal D3, D2-enoil-CoA isomerase	PECI	292.8	1	0.8	0.8	0.8	87	4	
	Acyl-CoA synthetase long-chain family member 4	ACSL4	5.8	2	1.1	1.4		88	4	
	Insignificant change: BHADH, ACAA1, CPT1A, ACADM, ACACA, CPT2									
Steroid (or drug) metabolism	Aldo-keto reductase family 1, member D1	AKR1D1	248.3	1	0.6	0.8	0.6	89	4	
	HMT1 hnRNP methyltransferase-like 2	HRMT1L2	6.6	1	0.8	1.0	0.8	90	4	
	Hydroxysteroid (11-beta) dehydrogenase 1	HSD11B1	1064.6	1	0.7	0.8	0.8	91	4	
	Hydroxysteroid (17-beta) dehydrogenase 4	HSD17B4	66.7	1	0.8	0.9	0.9	92	4	
	Steroid-5-alpha-reductase, alpha polypeptide 1	SRD5A1	74.3	1	0.7	0.8	0.9	93	4	
	UDP glycosyltransferase 2 family, polypeptide B7	UGT2B7	530.3	1	0.8	0.8	0.9	94	4	
	Sulfotransferase family 1E, estrogen-preferring, member 1	SULT1E1	52.8	1	0.5	0.5	1.4	95	7	
	Aldo-keto reductase family 1, member C4	AKR1C4	137.1	1	0.9	0.6	1.2	96	4	
	Hydroxysteroid (17-beta) dehydrogenase 2	HSD17B2	496.9	1	1.0	0.8	1.1	97	4	
	Sulfotransferase family, cytosolic, 2A, member 1	SULT2A1	689.0	1	0.9	0.8	0.9	98	4	
	Hydroxysteroid (17-beta) dehydrogenase 8	HSD17B8	72.2	1	0.7	0.7	1.0	99	7	
	Steroid sulfatase (microsomal), arylsulfatase C, isozyme S	STS	10.9	1	1.0	1.0	0.6	100	4	
	Emopamil binding protein (sterol isomerase)	EBP	197.0	1	1.0	0.7	0.7	101	5	
	Farnesyl-diphosphate farnesyltransferase 1	FDFT1	252.6	1	0.7	0.9	0.7	102	5	
	Insignificant change: HSD17B2, HSD3B1, LCMT1, SULT2A1, HMGCR, DHCR7, CBS2									
	Bile acid metabolism	Sterol O-acyltransferase 1	SOAT1	9.1	1	1.0	0.8	0.9	103	5
		Alcohol dehydrogenase 1C (class I), gamma polypeptide	ADH1C	312.7	1	0.7	0.6	1.1	104	5
		Alcohol dehydrogenase, iron containing, 1	ADHFE1	112.7	1	0.7	0.7	0.8	105	5
		Cytochrome P450, family 7, subfamily A, polypeptide 1	CYP7A1	104.1	1	0.7	0.6	2.9	106	5
	Prostenoind	Arachidonate 5-lipoxygenase-activating protein	ALOX5AP	7.9	2	1.7	0.7	1.0	107	3
Leukotriene B4 receptor 2		LTBR2	2.5	1	1.1	0.7	1.0	108	3	
Insignificant change: LTA4H, CYSLTR1, CYSLTR2, LTC4S, PPT1										
Aromatic amino acid metabolism	Dopa decarboxylase	DDC	51.4	1	1.0	0.8	0.8	109	4	
	Monoamine oxidase B	MAOB	426.6	1	0.8	0.8	1.0	110	4	
	Kynurenine 3-monooxygenase	KMO	51.7	1	1.0	0.7	1.0	111	4	
	Kynureninase	KYNU	63.3	1	0.9	0.7	0.8	112	4	
	Tyrosine aminotransferase	TAT	658.2	1	0.9	0.5	1.3	113	4	
	GTP cyclohydrolase 1	GCH1	49.6	1	0.9	0.7	0.9	114	4	
Insignificant change: HPD										
Sulfur-containing amino acid metabolism	MAT2	MAT2B	111.7	1	1.1	0.7	1.1	115	3	
	Cystathionase (cystathionine gamma-lyase)	CTH	112.0	1	1.0	0.7	1.1	116	3	
	Cystathionine-beta-synthase	CBS	350.4	1	1.1	0.7	0.9	117	3	
	Betaine-homocysteine methyltransferase	BHMT	1032.1	1	1.0	0.7	1.0	118	3	
	Methionine adenosyltransferase 1, alpha	MAT1A	547.6	1	1.0	0.5	1.1	119	2	
	Cysteine dioxygenase, type I	CDO1	84.4	1	0.9	0.7	1.0	120	1	
	Glutamate-cysteine ligase, catalytic subunit	GCLC	175.9	1	1.1	0.6	1.1	121	1	
	Glutathione S-transferase A1	GSTA1	2812.3	1	1.0	0.7	1.0	122	1	
	Alanyl (membrane) aminopeptidase	ANPEP	501.9	1	0.9	0.8	0.9	124	5	
	Bile acid Coenzyme A: amino acid N-acyltransferase	BAAT	284.4	1	1.0	0.6	1.0	125	5	
	Glutathione synthetase	GSS	70.1	1	1.1	0.8	0.8	126	5	
	Lactate dehydrogenase A	LDHA	716.3	1	0.8	0.7	1.0	127	5	
	Mercaptopyruvate sulfurtransferase	MPST	684.3	1	0.9	0.7	0.9	128	5	
	Serine dehydratase	SDS	440.4	1	0.9	0.4	1.0	129	5	
	Methionine adenosyltransferase II, alpha	MAT2A	122.9	1	1.0	0.9	0.8	130	5	
Insignificant change: GGT1, GSR, MTR, DNMMT1, CSAD, GCLM, LDHB										

Energy source amino acid metabolism	Phosphoenolpyruvate carboxykinase 2 (mitochondrial)	PCK2	791.8	1	0.8	0.9	1.0	131	4	
	Alanine-glyoxylate aminotransferase	AGXT	3352.3	1	0.7	0.9	1.0	132	4	
	Alanine-glyoxylate aminotransferase 2	AGXT2	122.8	1	0.6	0.8	1.0	133	4	
	Aldehyde dehydrogenase 2 family (mitochondrial)	ALDH2	2027.7	1	0.7	0.9	0.9	134	4	
	Aldehyde dehydrogenase 9 family, member A1	ALDH9A1	160.9	1	0.8	0.8	0.9	135	4	
	Pyruvate kinase, liver and RBC	PKLR	211.3	1	0.6	1.1	0.7	136	4	
	Aldehyde dehydrogenase 3 family, member A2	ALDH3A2	237.4	1	0.9	0.8	1.1	137	4	
	Phosphoenolpyruvate carboxykinase 1 (soluble)	PCK1	3166.8	1	1.0	0.5	1.2	138	4	
	Dihydropyrimidine dehydrogenase	DLD	72.3	1	0.9	0.8	0.9	139	4	
	Glutaminase 2 (liver, mitochondrial)	GLS2	133.8	1	0.8	0.7	1.0	140	4	
	Glutamate-ammonia ligase	GLUL	6.2	1	1.0	0.6	1.2	141	4	
	Glutamic-oxaloacetic transaminase 1, soluble	GOT1	388.2	1	0.9	0.7	0.8	142	4	
	Glutamic-pyruvate transaminase	GPT	298.8	1	0.8	0.6	1.1	143	4	
	Pyruvate carboxylase	PC	27.8	1	0.7	0.7	0.9	144	4	
	Phosphoglucomutase 1	PGM1	218.8	1	0.8	0.8	1.0	145	4	
	Pyruvate dehydrogenase kinase, isoenzyme 2	PDK2	24.0	1	0.9	0.8	0.9	146	7	
	Pyruvate dehydrogenase kinase, isoenzyme 4	PDK4	85.6	1	0.7	0.5	1.4	147	7	
	Fumarylacetoacetate hydrolase	FAH	85.8	1	0.9	0.9	0.8	148	4	
	Malic enzyme 1, NADP (+)-dependent, cytosolic	ME1	4.9	2	1.0	1.0	1.3	149	4	
	Insignificant change: ALDOA, ASNS, GOT2, MGCC33309, PDHB, PDK1									
	Glucose metabolism	Phosphorylase, glycogen, liver	PyGL	37.2	1	0.8	1.0	1.0	150	4
		Aldolase B, fructose-bisphosphate	ALDOB	13141.8	1	0.7	0.8	0.9	151	4
		Hexokinase 3 (white cell)	HK3	3.3	1	1.2	0.6	0.9	152	4
		Glycogen synthase 2 (liver)	GYS2	276.6	1	0.9	0.6	0.9	153	7
		Sterol regulatory element binding transcription factor 1	SREBF1	10.6	2	1.1	1.2	0.9	154	4
		Glycerol-3-phosphate dehydrogenase 1 (soluble)	GPD1	130.8	1	0.8	1.0	0.8	155	4
		Ketohexokinase (fructokinase)	KHK	411.8	1	0.7	1.0	0.8	156	4
Glucokinase (hexokinase 4) regulator		GCKR	204.8	1	0.7	1.0	0.8	157	4	
Aldolase C, fructose-bisphosphate		ALDOC	36.8	1	0.7	0.9	0.6	158	4	
Insignificant change: GCK, PFKFB1, G6PC, HK2										
Urea cycle		Carbamoyl-phosphate synthetase 1, mitochondrial	CPS1	1230.8	1	0.7	0.9	0.9	159	7
		Ornithine aminotransferase (gyrate atrophy)	OAT	94.8	1	0.5	0.6	1.6	160	7
Insignificant change: OTC										

Liver biopsy samples were analyzed by RT-PCR using a TaqMan® Gene Expression Assay probe, as described in the Materials and Methods. The genes were selected from DNA microarray data from a rat fibrosis model presented in a previous paper [3]. Genes in the ECM category (or other HSC markers) were selected as marker genes for HSC. Genes in the inflammation category were selected as markers of inflammatory cells in liver. Other genes were selected as markers of hepatocytes. Expression profiles of marker genes that showed statistically significant changes during fibrosis by a Kruskal-Wallis test are listed. Genes that showed statistically insignificant changes in expression are listed with the gene name only in Table 1. Columns from left to right indicate functional category, gene description, gene name, expression intensity at F1 stage, group classification (expression of marker genes in group 1 decreased almost linearly along with the F stages, expression of marker genes in group 2 increased almost linearly along with the F stages, and expression of marker genes in group 3 had a peak in the middle of the F stages), expression ratio between neighbouring F stages (a decreased ratio is shown in light gray and an increased ratio in deep gray), serial gene number and types of samples, which indicates the number of biopsy specimens, as shown in the Table 2. Actual alteration behavior of some characteristic genes are shown in Figure 1.

Table 2 Types of samples indicating the number of biopsy specimens

Type of sample	Number of samples			
	F1	F2	F3	F4
1	25	13	14	9
2	23	13	14	8
3	22	12	14	8
4	12	6	9	4
5	7	4	5	4
6	6	4	4	3
7	5	4	5	5

Quantitative limitation of biopsy specimens resulted in a difference in sample number for each probe, according to the interest in the particular gene.

transition to an adjacent F stage are similar to the selected genes in Table 1. An additional four marker genes selected in the *t* test analysis were added to the genes in Table 1. Over half of the selected marker genes from the DNA microarray data from the animal model showed statistically significant changes in the human samples, showing the effectiveness of the animal model for selection of genes

of significance in humans.

Genes showing changes in expression were roughly divided into three groups: genes in group 1 showed an almost linear decrease in expression along with an increase in F stage; genes in group 2 showed an almost linear increase with an increase in F stage; and genes in group 3 showed a peak in the middle of the F stage scale. Almost all genes in one category belonged to one or two groups, suggesting that genes in one category showed similar changes in expression during progression of fibrosis. The expression ratio between F stages is also shown in Table 1, with the peak ratio shown in bold. The peak for almost all genes in a given category occurred at the transition to the same F stage, again suggesting that genes in one category underwent changes in expression under similar mechanistic control.

Expression changes in gene clusters in the early phase (F1 to F2) of fibrosis

The functional categories showing a peak change in expression ratio in the early phase of fibrosis were inflammation, ECM, blood coagulation, lipid metabolism, half of the




Figure 1 Clusters of genes in functional categories. The characteristic behavior of gene clusters for each functional category is shown separately in Figure 1A-I. The numbers in each figure refer to the serial number of genes in Table 1. The expression ratio relative to the F1 stage is plotted in each graph. A: Inflammation gene cluster with a peak at F2; B: ECM gene cluster showing increased expression along with fibrosis progression; C: Insulin/Wnt signaling gene cluster with increased expression in the early phase of fibrosis; D: Blood coagulation factor gene cluster showing decreased expression along with fibrosis progression; E: Lipid metabolism gene cluster showing decreased expression along with fibrosis progression; F: Sulfur-containing amino acid metabolism gene cluster showing a synchronous decrease in expression from F2 to F3; G: Two transcription factor gene clusters showing sequential decreases in expression; H: Insulin/Wnt signaling gene cluster showing decreased expression from F2 to F3; I: A gene cluster showing a peak or bottom at F3.

genes in steroid metabolism, half of those in energy source amino AA metabolism, and half of transcription factors.

Expression of marker genes in the inflammation and ECM categories started to increase in the early phase. Expression

Table 3 Classification of gene clusters based on serial alteration of gene expression along with the fibrosis progression

F1-F2	F2-F3	F3-F4
Inflammation Wound-healing (ECM) Blood coagulation		
Transcription factors (cluster 1)	Transcription factors (cluster 2) Insulin/Wnt signal	
Lipid metabolism		
Steroid metabolism (cluster 1)	Steroid metabolism (cluster 1) Bile acid metabolism	
Energy source AA metabolism (cluster 1)	Energy source AA metabolism (cluster 1) Aromatic AA metabolism Sulfur-containing AA metabolism Glucose metabolism (cluster 1)	Glucose metabolism (cluster 2)

Gene clusters are indicated under the name of gene category. The pattern of serial alteration for gene cluster was determined based on the maximum changing point of the expression ratio shown in Table 1.

of marker genes related to inflammation, such as LYZ, GZMB, IL1B, TNF and TGFB1, occurred in clusters and reached a peak at fibrotic stage F2, as shown in Figure 1A, suggesting that inflammatory events are particularly active at the F2 stage. However, histological classification of inflammatory activity shows a tendency for an increase in inflammation that is correlated with an increase in F stage; therefore, the conclusion regarding inflammatory events based on expression of marker genes appears to differ from that based on histological classification. Gene expression in the ECM category also increased until F3 or F4, as shown in Figure 1B; expression of such genes might indicate an inflammatory response for wound healing. Increased expression of some clustered genes related to the cell cycle, CCND1, FOXM1 and GJA1 (Connexin 43), was also found at an early stage, as shown in Figure 1C, and might reflect a response to hepatic cell injury.

Almost all other genes were linearly down-regulated. Several genes related to blood coagulation, i.e. F10, AGT, FGA and PLG, were down-regulated as a cluster in the early phase, as shown in Figure 1D; this down-regulation may be linked to prolongation of the blood coagulation time in cirrhosis. An early response of many clustered genes associated with lipid metabolism was also found, as shown in Figure 1E. Expression of these genes decreased consecutively during fibrosis and the early response of genes affecting lipid metabolism is of interest.

Expression changes in gene clusters in the middle phase (F2 to F3) of fibrosis

The peak change in the expression ratio of marker genes in metabolism of sulfur-containing AA and aromatic AA was delayed, compared to genes associated with other kinds of metabolism. Marker genes for sulfur-containing AA metabolism decreased remarkably as a cluster in the phase from F2 to F3, as shown in Figure 1F. Decreased expression of marker genes for steroid metabolism, energy

source AA, and transcription factors were separable into two groups: early-response and middle-response genes.

All the down-regulated transcription factors, including TCF1 (HNF-1), HNF4A (HNF-4), CEBPA (C/EBP alpha), CEBPB (C/EBP beta), PPARA (PPAR alpha), RXRA (RXR alpha), NR1H3 (LXRA), NR1H2 (LXRB), NR1H4 (FXR), USF-1, USF-2, and NR0B2 are important in hepatic metabolism and other regulatory mechanisms. Alteration of expression of these genes might be related to abnormal expression of metabolic enzymes. Two clusters of transcription factors were clearly distinguishable based on their expression pattern, as shown in Figure 1G: the first cluster, including HNF-4, C/EBPA, RXR, TCF1 (HNF1), PPARA, and NR0B2, which showed altered expression in the early phase, might influence expression of the second cluster, including CEBPB, NR1H2(LXRB), NR1H3(LXRA), NR1H4(FXR), ESRRB and USF2, which showed altered expression in the middle phase of fibrosis.

A cluster of genes associated with insulin/Wnt signaling were down-regulated, as shown in Figure 1H, suggesting a common regulatory mechanism. These expression changes are likely to be related to changes in expression of transcription factors and genes in metabolic networks. The down-regulated genes included GSK3B and CTNNB1 (catenin beta 1), which participates in Wg/Wnt signaling for regulation of cell proliferation and differentiation^[4]; GJA1 (connexin 43), which forms gap junctions and is regulated by Wg/Wnt signaling^[9]; and FOXM1, which is associated with cell proliferation^[8] and liver regeneration^[7]. All of these genes had peak expression at F2 in a clustered manner, as shown in Figure 1C and G. Enhancement of cell proliferation for wound healing might be linked with a peak of inflammation at F2, and expression of genes such as CCND1, GJA1 and FOXM1 in the downstream part of the insulin/Wnt pathway were altered ahead of genes involved in insulin/Wnt signaling. The relationship between these genes requires further study.

Expression changes in gene clusters in the late phase (F3 to F4) of fibrosis

Few genes showed altered expression in the late phase of fibrosis, but a cluster of genes in the glucose metabolism category showed decreased expression. It was also of interest that expression of several genes reversed direction or abruptly altered in the late phase, as shown in Figure 1I. These results suggest different biological changes from the start of the late stage in the transition from F3 to F4.

Serial expression changes for the functional categories are summarized in Table 3 and these data indicate associations between clustered genes in one category and inter-category relationships.

Down-regulated individual marker genes (group 1)

Regeneration of hepatic cells is suppressed during fibrosis and such suppression is thought to then cause further fibrosis. IGF1, GHR and IL6R (inflammation) support the regeneration of hepatocytes and down-regulation of the expression of these genes may be linked directly to formation of fibrosis. Down-regulation of PON1, which associates with HDL (high-density lipoprotein)

and regulates the cellular redox state, and PPT1, which is known as a lysosomal hydrolase of long chain fatty acyl CoA and has a role in maintenance of synaptic function, may be related to mitochondrial damage, as we and others have suggested^[10,11]. RGN, a calcium-binding protein that plays a pivotal role in maintaining cell homeostasis and function, was also down-regulated. Down-regulation of these genes may impair liver function. TRIB3 inhibits Akt/PKB activation by insulin^[12], and this gene was suddenly down-regulated from the F3 to F4 phase, suggesting new conditions in the insulin signaling pathway in the transition from F3 to F4.

Up-regulated individual marker genes (group 2)

Most inflammatory marker genes showed peak expression in the middle phase of fibrosis, as shown in Figure 1A, but CMA1 (chymase 1), which is produced by mast cells, underwent a linear increase in expression with progression of fibrosis. This is of note, since chymase has been reported to be involved in chronic hepatic fibrosis in autoimmune hepatitis and primary biliary cirrhosis^[13], and mast cells may have a special role in fibrogenesis. The role of mast cells in chronic inflammation, however, deserves further study, because of the shortage of determined sample and low expression of CMA1.

Individual marker genes with peak expression in the middle phase of fibrosis (group 3)

The only hepatic stellate cell (HSC)-specific marker gene to show peak expression at the F2 stage was PRNP, which is reported to be a marker for the early phase of HSC activation. The amino acid transporters SLC38A2 (ATA2), SLC6A6 (TAUT) and SLC7A1 (CAT-1) showed peak expression at F2, which may also suggest enhancement of cell proliferation at this stage of fibrosis. Increased expression of SLC38A2, which preferentially transports alanine, has been reported in regeneration of hepatocytes^[12], since hepatocytes require alanine as an energy source^[13]. Based on our results, down-regulation of SLC38A2 in the late phase of fibrosis suggests that utilization of alanine as an energy source decreases at this stage of fibrosis. Up-regulation of SLC6A6, a taurine transporter, in the early phase of fibrosis can be understood as protective behavior against injury of hepatocytes^[14]. FASN, a marker gene for fatty acid metabolism, showed peak expression at the F3 stage of fibrosis. We have observed suppression of biosynthesis and degradation of fatty acids in the liver in a CCl₄-induced cirrhotic rat model, and clinical results show temporal enhancement of fatty acid biosynthesis before subsequent suppression in an advanced phase of fibrosis; this may be a compensative action related to suppression of other forms of energy metabolism in fibrosis, as we have previously suggested^[8].

Candidate marker genes for diagnosis of fibrosis and discrimination of fibrotic stages

Diagnosis in the early stage of fibrosis may be important for monitoring progression of fibrosis and hepatic biological changes. Candidate marker genes at each step of fibrosis were selected based on an expression change

ratio > 1.5 and an intensity > 10. Up-regulation of LYZ and down-regulation of FGA, OAT and AGXT2 were noteworthy in the transition from F1 to F2, and up-regulation of genes in the ECM category and down-regulation of genes in metabolism of energy source AA, aromatic AA, steroids and sulfur-containing AA occurred in the transition from F2 to F3. Diagnosis of the late stage of fibrosis (the transition from F3 to F4) is important because of the risk of tumorigenesis. Some genes showed a reversal of expression in the F3 to F4 transition, suggesting that biological changes in the stage from F3 to F4 are qualitatively different from those at earlier stages. Remarkable down-regulation of TRIB3, which inhibits insulin signaling and NF κ B signaling, was noteworthy in the F3 to F4 transition. The reversal in expression of FASN in this phase may indicate changes in lipid metabolism and that of OAT indicates changes in ornithine metabolism or the urea cycle. The early increase in collagen expression began to decline in the F3 to F4 stage and the similar decline of SERPINE1 (Pai1) expression may reflect increased fibrolysis. Consecutive analysis of marker molecules in plasma will be important for monitoring progression of fibrosis at each step, and the data in this paper provides useful information for the selection of serum markers and interpretation of changes in the levels of these markers.

Coordinated regulation of functional categories

All genes in Figure 1 were subjected to hierarchical clustering analysis. Statistical clustering of the expression ratio between neighboring F stages for the genes in Table 1 was combined with functional categories, as shown in Figure 2. Functional categories clearly corresponded to the statistical clusters, suggesting coordinated regulation of genes in one functional category. Overlap of functional categories in statistical clusters suggested that these categories might be regulated by correlated mechanisms. In contrast, separation of members of a category into multiple positions of a statistical cluster suggests that the functional category may be divided into subgroups with respect to regulation.

DISCUSSION

Biological interpretation of changes in gene expression

Changes in expression of hepatic cell-specific marker genes reflects biological changes in the progression of hepatic fibrosis. Shimizu *et al.*^[5] reported co-localization of chymase with fibrotic tissue, and we have reported increased expression of marker genes for mast cells, including chymase, in progression of fibrosis in a DNM-induced rat fibrosis model. The results reported here also show correlation of the expression of chymase with the stage of fibrosis. Therefore, these results suggest that a certain cell population producing chymase has an important role in the pathogenesis of fibrosis.

Many marker genes related to inflammation showed peak expression at F2. Inflammation has been reported to induce activation of HSCs^[15], and we have shown a peak in activated inflammatory cells in a DNM-induced fibrosis model. These results suggest that a temporal




Figure 2 Comparison between statistical clustering analysis and functional cluster. All genes in Figure 1 were subjected to hierarchical clustering analysis using Genowiz™. Statistical clustering of the expression ratio between neighboring F stages for the genes in Table 1 was combined with functional categories. The color coding in the center column shows the expression ratio between neighboring F stages for the genes in Table 1; Ratio increases from green to red. The left dendrogram shows the result of hierarchical clustering by Genowiz™ software. The numbers on the right of gene names indicate the subgraph number in Figure 1, which represents functional categories, and the same numbers are tied with color bars.

strong inflammatory reaction is required for initiation of auto-stimulatory activation of HSCs. In the DMN-induced fibrosis model, inflammation was suppressed after it reached a peak, but weaker inflammation might still be sufficient to maintain activation of HSCs.

Expression of HSC-specific marker genes such as COL1A1, LUM, TAGLN, MGP, SERPINE1 (PAI-1) and LOX did not increase or decrease continuously from F3 to F4, as shown in Figure 1B and 1I, whereas expression of DCN, BGN and GPC3 increased comparatively from F3 to F4, as shown in Figure 1B. This suggests some changes in the behavior of HSCs between F3 and F4, and it has been hypothesized that the perpetuation phase^[11] may change to a new phase in fibrosis of stage F4. Therefore, profiles of HSC-specific marker genes may be useful for

indication of the phase of fibrosis.

A cluster of down-regulated genes associated with sulfur-containing AA metabolism is shown in Figure 1F. Changes in expression of the clustered genes occurred simultaneously between the F2 and F3 stages; this transition may reflect a biological change causing fibrotic progression, and the change in expression of the clustered genes seems to suggest an increased demand for glutathione^[18]. Abnormalities in metabolism of sulfur-containing AA in thioacetamide-induced cirrhosis in rat liver have been found in a proteomics analysis^[19], and similar abnormalities were also reported based on hepatic gene expression changes in patients with alcoholic hepatitis^[20].

There is only limited information on altered expression of transcription factors in fibrotic liver, but down-regulation

of HNF-4 in human cirrhosis^[21] and of PPARs in hepatitis C virus genotype 3^[22] have been reported. C/EBP alpha and C/EBP beta regulate proliferation of hepatocytes^[23] and glucose and lipid homeostasis^[24-27], and HNF-1 and HNF 4 broadly regulate hepatic functions such as carbohydrate metabolism^[28], lipid metabolism^[29], bile acid metabolism and HDL-cholesterol metabolism^[30]; expression of HNF-1 is also regulated by HNF-4^[31]. USF1 and USF2 have been reported as glucose signals^[32,33], and the USF1 and USF2 homodimers and the USF1-USF2 heterodimer regulate expression of liver-specific genes such as apolipoprotein A2 and pyruvate kinase. HNF-4 and USF2a bind to the enhancer sequence cooperatively^[34]. HNF-4 also regulates PPAR alpha^[35], which in turn regulates glucose^[36], lipid^[37] and cholesterol metabolism^[38]. RXR alpha regulates lipid, bile acid and cholesterol homeostasis^[39], and LXR alpha, LXR beta and FXR are associated with lipid^[40,41], bile acid and cholesterol homeostasis^[40,42]. RXR and FXR form heterodimers with other transcription factors, including other members of the same family or with PPAR alpha^[40,43,44]. NR0B2 (SHP) regulates cholesterol metabolism^[45], glucose metabolism^[46] and bile acid synthesis^[46], and interacts with LXR^[47]. AKT1 and AKT2 are important kinases in the pathway of insulin regulation of glucose homeostasis^[48] and in fatty acid synthesis^[49]. Hence, the altered expression of these transcription factors may relate to altered expression of metabolic enzymes in glucose and lipid metabolism in the fibrotic liver of hepatitis C patients.

The continuous increase in expression of cyclin D1 (CCND1) correlated with F stage, as shown in Figure 1C, and appears to be important for hepatic tumorigenesis. The association of Wnt signaling^[50,51] and cyclin D^[52] with tumorigenesis is well known. Catenin beta 1 (CTNNB1) regulates cyclin D expression^[53] and is itself regulated through phosphorylation by GSK3B^[54] or other kinases^[55]. Down-regulation of expression of CTNNB1 occurred in advanced fibrotic stages, as shown in Figure 1H, and beta interacting protein 1 (CTNNBIP1), which is a negative regulator of CTNNB1, was also down-regulated, as also shown in Figure 1H. From this perspective, it was interesting that alterations in expression of genes associated with insulin signaling, including GSK3B, CTNNB1, CTNNBIP1 and downstream genes such as CCND1 and GJA1 (connexin 43), were clustered, as shown in Figures 1C and H. Suppression of insulin signaling has been reported in cirrhosis^[56]; however, the response of downstream genes in this pathway was inconsistent with suppression of the insulin signal. Therefore, further studies will be necessary to clarify whether this inconsistency arises from differences between expression levels and the activity of the protein products of these genes, or if another signal^[57] is involved in the Wnt and insulin signaling pathways.

The bioSpace Explorer is a system for analysis of DNA microarray data that may allow an understanding of the molecular relationships underlying the above results. The network of genes from previous reports revealed integral relationships among insulin/Wnt signal, CEBPA, PPARA, RXRA, glucose metabolism, lipid synthesis and lipid metabolism. An initial version of bioSpace Explorer was constructed based on molecular networks related to lipid metabolism, with entries showing relationships between

molecules *via* a line between the molecules. This bird's eye view of the pathway including lipid metabolism with the input PCR data is illustrated in Figure 3; molecules related to inflammation were up-regulated and many genes related to lipid metabolism were down-regulated.

Hypothetical causes of biological changes in progression of fibrosis

Biological changes in fibrosis can be summarized as follows. Initially, Kupffer cells or other inflammatory cells are activated in the transition from F1 to F2. This event immediately influences production of ECM and cell cycle genes for wound-healing^[58]. Blood coagulation is quickly suppressed in moving from F1 to F2, as shown by down-regulation of coagulation factor genes and up-regulation of the inhibitor, PAI-1. Several transcription factor genes are also immediately influenced, probably due to inflammation, as shown by the down-regulation of CEBPA, HNF4A, TCF1 and NR0B2. Expression of many genes associated with lipid metabolism also changed quickly in the transition from F1 to F2. Down-regulation of these genes may be controlled by down-regulation of the transcription factors, especially RXRA, PPARA, LXRs and FXR. Some genes related to steroid metabolism also responded quickly for control of inflammation.

Expression of genes associated with sulfur-containing AA metabolism and aromatic AA metabolism changed simultaneously in the transition from F2 to F3. The first type of metabolism relates to the redox state^[59] and the second is associated with production of active metabolites such as catecholamines and serotonin. Such important biological states are controlled to maintain homeostasis through several mechanisms^[60] and this may explain the delayed change in expression of these genes. The expression of many genes related to intracellular signaling, including insulin/Wnt signaling, also changed simultaneously in the transition from F2 to F3. This delayed change may also reflect compensative action for hepatic cellular defects on metabolism for energy supply and/or hepatic cellular proliferation. The main molecules in fibrosis, such as collagens, increase in expression from F2 to F3 and cause development of fibrosis through activation of HSCs through a stimulatory cycle involving inflammatory cells, HSCs and hepatocytes, as described previously^[61].

Some quantitative biological changes started in the transition from F3 to F4. Down-regulation of genes associated with sugar metabolism and fatty acid synthesis at this stage might induce persistent defects in energy storage and supply to the liver. The liver transits into an inescapable negative cycle between this defect in the hepatic energy state and mitochondrial damage in cirrhosis. This negative cycle will be discussed in a future paper describing DNA microarray analysis of an animal model of cirrhosis.

Coordinated regulation of functional categories

Statistical cluster analysis showed coordinated regulation of functional categories in liver fibrosis. These regulatory mechanisms can be examined prospectively using bioSpace Explorer, and these results will be discussed in a future paper. In the current work, the clinical gene expression profiles assembled using RT-PCR, using genes




Figure 3 Molecular network associated with lipid metabolism. Gene expression changes in pathways related to lipid metabolism, illustrated with bioSpace Explorer (a system for analysis of DNA microarray data for lipid metabolism; PharmaFrontier Co. Ltd / World Fusion Co. Ltd; see texts for details). The bird's-eye view of the lipid metabolism is displayed. The up-regulated and down-regulated gene expression ratios at F3 vs F1 in Table 1 are displayed in red and green, respectively, with the color gradation proportional to the ratio. Genes in Table 1 that did not show a statistically significant change in expression are indicated with a blue circle with gray background. An entry with a gray background only indicates no input data. Most of the entity names in Figure 2 are the same as the gene name in Table 1, but the names 'C20orf97', 'SCEH1', 'Acaa2', and 'ACSS5' in Figure 2 refer to 'TRIB3', 'ECHS1', 'ACAA2', and 'ACSL5', respectively, in Table 1. These differences are due to the software used in bioSpace Explorer.

originally selected based on DNA microarray data from an experimental animal model, provided an improved understanding of disease and suggested new methods of diagnosis. Therefore, statistical analysis and functional clustering or network analysis of the transcriptome, alone or in combination, can provide an overview of a dynamic biological system.

ACKNOWLEDGMENTS

We thank Ms. Tomoko Kobayashi of Pharmafrontier Co. Ltd. and Mr. Ken Osaki of World Fusion Co. Ltd for statistical analysis and operation of the bioSpace Explorer for analysis of DNA microarray data for lipid metabolism.

COMMENTS

Background

Information from clinical specimens is very important, but there is a limitation in sample preparation. With DNA microarray, it is difficult to determine gene expression profile from a small amount of sample such as a clinical needle biopsy sample. RT-PCR with TaqMan probe can make it possible with high quality. We have to get maximum information with a minimum number of TaqMan probes

because the amount of samples is limited and probes are expensive. Selection of probes is a key factor. We extensively determined gene expression profiles from animal models of liver fibrosis with DNA microarray (WJG 2006; 12: 6473). The gene marker sets were arranged to show the change of gene expression of a molecular network or functionally clustered gene sets. In this paper, the selected gene marker sets effectively showed the dynamic behavior of global gene network change during liver fibrosis.

Research frontier

Dynamic behavior of genome-wide genes expression is now measured with DNA microarray. This technology must be applied to clinical samples. Such information can greatly advance the study of disease pathogenesis, diagnosis and therapy. One problem is the interpretation of huge expression profile data from DNA microarray. Advanced technology of computational text-mining has recently shown the genome-wide molecular networks or functional molecular clusters. This genome-wide network is going to be applied to the analysis of DNA microarray data. When expression profile data are arranged as a change in their networks of functional clusters, these huge data are expected to suggest effectively the biological meaning in terms of broad aspects of research interest. Therefore, we are developing an analysis system genome-wide molecular network which was made possible by a combination system based on both computational and manual text-mining. This system was partially applied on the interpretation of gene expression profile in this paper. Prevention therapy for individual patients at an early stage is required because genomic polymorphism is going to reveal the personal risk of diseases. The biological background of progression to disease onset must be understood for development of prevention therapy. Prediction of liver cancer risk is going to be tied in the early

stage of liver fibrosis before cirrhosis. The accumulation process of hepatic stress which leads to onset of liver cancer has to be elucidated. For example, there is a question why BCAA (branched chain amino acids), which improves hepatic metabolism, reduces onset of liver cancer.

Innovations and breakthrough

We have already prepared the gene marker sets which can show the change of molecular networks or functionally clustered genes by DNA microarray experiment on liver fibrosis of animal models. The gene marker sets were linked to biological events in each hepatic cell such as hepatocytes, immune cells and stellate cells. Application of gene marker sets and RT-PCR with TaqMan probe technology on clinical specimens successfully showed the serial change of gene expression in molecular network or functionally clustered gene sets in the progression of liver fibrosis. It was proved that network analysis is a powerful tool for biological research. Our sequential approach (animal model/DNA microarray → selection of appropriate gene marker sets in molecular networks → clinical samples/RT-PCR → analysis functionally clustered gene markers in molecular networks → analysis of the relation between molecular networks) can effectively advance clinical research. Serial change of clustered gene expression during liver fibrosis progression, which was made clear in this paper, will reveal the molecular mechanism of many symptoms before and after the onset of cirrhosis and liver cancer.

Application

Gene marker sets and RT-PCR on clinical specimens as well as analytical methods with genome-wide gene networks can be applied to get the information of dynamic biological progression on various diseases. Serial change of clustered gene expression during liver fibrosis progression can provide a method of diagnosis and therapy in liver fibrosis. For example, the question why BCAA, which improves hepatic metabolism, reduces onset of liver cancer, will be solved.

Terminology

Text-mining: all published information about molecular interaction and their function are collected and arranged to make new valuable information such as gene-wide molecular networks; TaqMan-PCR probe: Applied Biosystems offers more than 700,000 TaqMan® Gene Expression Assays, the most comprehensive set of pre-designed Real-Time PCR assays available. All TaqMan® Gene Expression Assays have been designed using validated bioinformatics pipeline of Applied Biosystems, and run with the same PCR protocol, eliminating the need for primer design or PCR optimization.

Peer review

This paper revealed that metabolic deficiency occurs before the onset of cirrhosis. It had already been found in animal models with hepatic toxic substances in the preceding paper. Metabolic deficiency in hepatitis which was caused by a virus was found to be the same as animal models in this paper. Gene marker sets, selected from analysis of animal models, and analysis methods using molecular networks can lead to success in finding the serial change of functionally clustered genes expression during liver fibrosis progression. A set of appropriate gene markers in each network was a key to analysis. The sequential approach (animal model/DNA microarray → appropriate gene marker sets in molecular networks → clinical samples/RT-PCR → analysis functionally clustered gene markers in molecular network → analysis of the relation between molecular networks) is useful to elucidate the molecular mechanism of disease.

REFERENCES

- Intraobserver and interobserver variations in liver biopsy interpretation in patients with chronic hepatitis C. The French METAVIR Cooperative Study Group. *Hepatology* 1994; 20: 15-20
- Gobel T, Vorderwulbecke S, Hauck K, Fey H, Haussinger D, Erhardt A. New multi protein patterns differentiate liver fibrosis stages and hepatocellular carcinoma in chronic hepatitis C serum samples. *World J Gastroenterol* 2006; 12: 7604-7612
- Takahara Y, Takahashi M, Wagatsuma H, Yokoya F, Zhang QW, Yamaguchi M, Aburatani H, Kawada N. Gene expression profiles of hepatic cell-type specific marker genes in progression of liver fibrosis. *World J Gastroenterol* 2006; 12: 6473-6499
- Dierick H, Bejsovec A. Cellular mechanisms of wingless/Wnt signal transduction. *Curr Top Dev Biol* 1999; 43: 153-190
- Ai Z, Fischer A, Spray DC, Brown AM, Fishman GI. Wnt-1 regulation of connexin43 in cardiac myocytes. *J Clin Invest* 2000; 105: 161-171
- Wang IC, Chen YJ, Hughes D, Petrovic V, Major ML, Park HJ, Tan Y, Ackerson T, Costa RH. Forkhead box M1 regulates the transcriptional network of genes essential for mitotic progression and genes encoding the SCF (Skp2-Cks1) ubiquitin ligase. *Mol Cell Biol* 2005; 25: 10875-10894
- Wang X, Bhattacharyya D, Dennewitz MB, Kalinichenko VV, Zhou Y, Lepe R, Costa RH. Rapid hepatocyte nuclear translocation of the Forkhead Box M1B (FoxM1B) transcription factor caused a transient increase in size of regenerating transgenic hepatocytes. *Gene Expr* 2003; 11: 149-162
- Ferre N, Marsillach J, Camps J, Mackness B, Mackness M, Riu F, Coll B, Tous M, Joven J. Paraoxonase-1 is associated with oxidative stress, fibrosis and FAS expression in chronic liver diseases. *J Hepatol* 2006; 45: 51-59
- Kim SJ, Zhang Z, Lee YC, Mukherjee AB. Palmitoyl-protein thioesterase-1 deficiency leads to the activation of caspase-9 and contributes to rapid neurodegeneration in INCL. *Hum Mol Genet* 2006; 15: 1580-1586
- Du K, Herzig S, Kulkarni RN, Montminy M. TRB3: a tribbles homolog that inhibits Akt/PKB activation by insulin in liver. *Science* 2003; 300: 1574-1577
- Satomura K, Yin M, Shimizu S, Kato Y, Nagano T, Komeichi H, Ohsuga M, Katsuta Y, Aramaki T, Omoto Y. Increased chymase in livers with autoimmune disease: colocalization with fibrosis. *J Nippon Med Sch* 2003; 70: 490-495
- Fowler FC, Banks RK, Mailliard ME. Characterization of sodium-dependent amino acid transport activity during liver regeneration. *Hepatology* 1992; 16: 1187-1194
- Freeman TL, Ngo HQ, Mailliard ME. Inhibition of system A amino acid transport and hepatocyte proliferation following partial hepatectomy in the rat. *Hepatology* 1999; 30: 437-444
- Warskulat U, Borsch E, Reinehr R, Heller-Stilb B, Monnighoff I, Buchczyk D, Donner M, Flögel U, Kappert G, Soboll S, Beer S, Pfeiffer K, Marschall HU, Gabrielsen M, Amiry-Moghaddam M, Ottersen OP, Dienes HP, Haussinger D. Chronic liver disease is triggered by taurine transporter knock-out in the mouse. *FASEB J* 2006; 20: 574-576
- Shimizu S, Satomura K, Aramaki T, Katsuta Y, Takano T, Omoto Y. Hepatic chymase level in chronic hepatitis: colocalization of chymase with fibrosis. *Hepatology Res* 2003; 27: 62-66
- Baroni GS, Pastorelli A, Manzin A, Benedetti A, Marucci L, Solfrosi L, Di Sario A, Brunelli E, Orlandi F, Clementi M, Macarri G. Hepatic stellate cell activation and liver fibrosis are associated with necroinflammatory injury and Th1-like response in chronic hepatitis C. *Liver* 1999; 19: 212-219
- Gressner AM. Transdifferentiation of hepatic stellate cells (Ito cells) to myofibroblasts: a key event in hepatic fibrogenesis. *Kidney Int Suppl* 1996; 54: S39-S45
- Wu G, Fang YZ, Yang S, Lupton JR, Turner ND. Glutathione metabolism and its implications for health. *J Nutr* 2004; 134: 489-492
- Low TY, Leow CK, Salto-Tellez H, Chung MC. A proteomic analysis of thioacetamide-induced hepatotoxicity and cirrhosis in rat livers. *Proteomics* 2004; 4: 3960-3974
- Lee TD, Sada MR, Mendler MH, Bottiglieri T, Kanel G, Mato JM, Lu SC. Abnormal hepatic methionine and glutathione metabolism in patients with alcoholic hepatitis. *Alcohol Clin Exp Res* 2004; 28: 173-181
- Berasain C, Herrero JI, Garcia-Trevijano ER, Avila MA, Esteban JI, Mato JM, Prieto J. Expression of Wilms' tumor suppressor in the liver with cirrhosis: relation to hepatocyte nuclear factor 4 and hepatocellular function. *Hepatology* 2003; 38: 148-157
- de Gottardi A, Paziienza V, Fugnale F, Rubbia-Brandt L, Juge-Aubry CE, Meier CA, Hadengue A, Negro F. Peroxisome proliferator-activated receptor-alpha and -gamma mRNA levels are reduced in chronic hepatitis C with steatosis and genotype 3 infection. *Aliment Pharmacol Ther* 2006; 23: 107-114
- Greenbaum LE, Cressman DE, Haber BA, Taub R. Coexistence of C/EBP alpha, beta, growth-induced proteins and DNA synthesis in hepatocytes during liver regeneration. Implications

- for maintenance of the differentiated state during liver growth. *J Clin Invest* 1995; 96: 1351-1365
- 24 Arizmendi C, Liu S, Croniger C, Poli V, Friedman JE. The transcription factor CCAAT/enhancer-binding protein beta regulates gluconeogenesis and phosphoenolpyruvate carboxykinase (GTP) gene transcription during diabetes. *J Biol Chem* 1999; 274: 13033-13040
- 25 Gautier-Stein A, Mithieux G, Rajas F. A distal region involving hepatocyte nuclear factor 4alpha and CCAAT/enhancer binding protein markedly potentiates the protein kinase A stimulation of the glucose-6-phosphatase promoter. *Mol Endocrinol* 2005; 19: 163-174
- 26 Qiao L, MacLean PS, You H, Schaack J, Shao J. Knocking down liver ccaat/enhancer-binding protein alpha by adenovirus-transduced silent interfering ribonucleic acid improves hepatic gluconeogenesis and lipid homeostasis in db/db mice. *Endocrinology* 2006; 147: 3060-3069
- 27 Wang ND, Finegold MJ, Bradley A, Ou CN, Abdelsayed SV, Wilde MD, Taylor LR, Wilson DR, Darlington GJ. Impaired energy homeostasis in C/EBP alpha knockout mice. *Science* 1995; 269: 1108-1112
- 28 Pontoglio M. Hepatocyte nuclear factor 1, a transcription factor at the crossroads of glucose homeostasis. *J Am Soc Nephrol* 2000; 11 Suppl 16: S140-S143
- 29 Louet JF, Hayhurst G, Gonzalez FJ, Girard J, Decaux JF. The coactivator PGC-1 is involved in the regulation of the liver carnitine palmitoyltransferase I gene expression by cAMP in combination with HNF4 alpha and cAMP-response element-binding protein (CREB). *J Biol Chem* 2002; 277: 37991-38000
- 30 Shih DQ, Bussen M, Sehaye E, Ananthanarayanan M, Schneider BL, Suchy FJ, Shefer S, Bollileti JS, Gonzalez FJ, Breslow JL, Stoffel M. Hepatocyte nuclear factor-1alpha is an essential regulator of bile acid and plasma cholesterol metabolism. *Nat Genet* 2001; 27: 375-382
- 31 Miura N, Tanaka K. Analysis of the rat hepatocyte nuclear factor (HNF) 1 gene promoter: synergistic activation by HNF4 and HNF1 proteins. *Nucleic Acids Res* 1993; 21: 3731-3736
- 32 Corre S, Galibert MD. Upstream stimulating factors: highly versatile stress-responsive transcription factors. *Pigment Cell Res* 2005; 18: 337-348
- 33 Kahn A. From the glycogenic function of the liver to gene regulation by glucose. *C R Seances Soc Biol Fil* 1998; 192: 813-827
- 34 Morifuzumi S, Gourdon L, Lefrançois-Martinez AM, Kahn A, Raymondjean M. Effect of different basic helix-loop-helix leucine zipper factors on the glucose response unit of the L-type pyruvate kinase gene. *Gene Expr* 1998; 7: 103-113
- 35 Pineda Torra I, Jamshidi Y, Flavell DM, Fruchart JC, Staels B. Characterization of the human PPARalpha promoter: identification of a functional nuclear receptor response element. *Mol Endocrinol* 2002; 16: 1013-1028
- 36 Xu J, Chang V, Joseph SB, Trujillo C, Bassilian S, Saad MF, Lee WN, Kurland IJ. Peroxisomal proliferator-activated receptor alpha deficiency diminishes insulin-responsiveness of gluconeogenic/glycolytic/pentose gene expression and substrate cycle flux. *Endocrinology* 2004; 145: 1087-1095
- 37 Lee SS, Chan WY, Lo CK, Wan DC, Tsang DS, Cheung WT. Requirement of PPARalpha in maintaining phospholipid and triacylglycerol homeostasis during energy deprivation. *J Lipid Res* 2004; 45: 2025-2037
- 38 Chakravarthy MV, Pan Z, Zhu Y, Tordjman K, Schneider JC, Coleman T, Turk J, Semenkovich CF. "New" hepatic fat activates PPARalpha to maintain glucose, lipid, and cholesterol homeostasis. *Cell Metab* 2005; 1: 309-322
- 39 Wan YJ, Cai Y, Lungo W, Fu P, Locker J, French S, Sucov HM. Peroxisome proliferator-activated receptor alpha-mediated pathways are altered in hepatocyte-specific retinoid X receptor alpha-deficient mice. *J Biol Chem* 2000; 275: 28285-28290
- 40 Edwards PA, Kennedy MA, Mak PA. LXRs: oxysterol-activated nuclear receptors that regulate genes controlling lipid homeostasis. *Vascu Pharmacol* 2002; 38: 249-256
- 41 Sinal CJ, Tohkin M, Miyata M, Ward JM, Lambert G, Gonzalez FJ. Targeted disruption of the nuclear receptor FXR/BAR impairs bile acid and lipid homeostasis. *Cell* 2000; 102: 731-744
- 42 Lambert G, Amar MJ, Guo G, Brewer HB Jr, Gonzalez FJ, Sinal CJ. The farnesoid X-receptor is an essential regulator of cholesterol homeostasis. *J Biol Chem* 2003; 278: 2563-2570
- 43 Vidal H. PPAR receptors: recent data. *Ann Endocrinol (Paris)* 2005; 66: 155-159
- 44 Yoshikawa T, Shimano H, Amemiya-Kudo M, Yahagi N, Hasty AH, Matsuzaka T, Okazaki H, Tamura Y, Iizuka Y, Ohashi K, Osuga J, Harada K, Gotoda T, Kimura S, Ishibashi S, Yamada N. Identification of liver X receptor-retinoid X receptor as an activator of the sterol regulatory element-binding protein 1c gene promoter. *Mol Cell Biol* 2001; 21: 2991-3000
- 45 Kim HJ, Kim JY, Kim JY, Park SK, Seo JH, Kim JB, Lee IK, Kim KS, Choi HS. Differential regulation of human and mouse orphan nuclear receptor small heterodimer partner promoter by sterol regulatory element binding protein-1. *J Biol Chem* 2004; 279: 28122-28131
- 46 Ito S, Fujimori T, Furuya A, Satoh J, Nabeshima Y, Nabeshima Y. Impaired negative feedback suppression of bile acid synthesis in mice lacking betaKlotho. *J Clin Invest* 2005; 115: 2202-2208
- 47 Brendel C, Schoonjans K, Botrugno OA, Treuter E, Auwerx J. The small heterodimer partner interacts with the liver X receptor alpha and represses its transcriptional activity. *Mol Endocrinol* 2002; 16: 2065-2076
- 48 Cho H, Thorvaldsen JL, Chu Q, Feng F, Birnbaum MJ. Akt1/PKBalpha is required for normal growth but dispensable for maintenance of glucose homeostasis in mice. *J Biol Chem* 2001; 276: 38349-38352
- 49 Ono H, Shimano H, Katagiri H, Yahagi N, Sakoda H, Onishi Y, Anai M, Ogihara T, Fujishiro M, Viana AY, Fukushima Y, Abe M, Shojima N, Kikuchi M, Yamada N, Oka Y, Asano T. Hepatic Akt activation induces marked hypoglycemia, hepatomegaly, and hypertriglyceridemia with sterol regulatory element binding protein involvement. *Diabetes* 2003; 52: 2905-2913
- 50 Colnot S, Decaens T, Niwa-Kawakita M, Godard C, Hamard G, Kahn A, Giovannini M, Perret C. Liver-targeted disruption of Apc in mice activates beta-catenin signaling and leads to hepatocellular carcinomas. *Proc Natl Acad Sci USA* 2004; 101: 17216-17221
- 51 Murata M, Miyoshi Y, Ohsawa M, Shibata K, Ohta T, Imai Y, Nishikawa M, Iwao K, Tateishi H, Shimano T, Kobayashi T, Nakamura Y. Accumulation of beta-catenin in the cytoplasm and the nuclei during the early hepatic tumorigenesis. *Hepatology* 2001; 21: 126-135
- 52 Yamaoka H, Ohtsu K, Sueda T, Yokoyama T, Hiyama E. Diagnostic and prognostic impact of beta-catenin alterations in pediatric liver tumors. *Oncol Rep* 2006; 15: 551-556
- 53 Kolligs FT, Bonner G, Goke B. Wnt/beta-catenin/tcf signaling: a critical pathway in gastrointestinal tumorigenesis. *Digestion* 2002; 66: 131-144
- 54 Zeng X, Tamai K, Doble B, Li S, Huang H, Habas R, Okamura H, Woodgett J, He X. A dual-kinase mechanism for Wnt co-receptor phosphorylation and activation. *Nature* 2005; 438: 873-877
- 55 Taurin S, Sandbo N, Qin Y, Browning D, Dulin NO. Phosphorylation of beta-catenin by cyclic AMP-dependent protein kinase. *J Biol Chem* 2006; 281: 9971-9976
- 56 Picardi A, D'Avola D, Gentilecci LV, Galati G, Fiori E, Spataro S, Afeltra A. Diabetes in chronic liver disease: from old concepts to new evidence. *Diabetes Metab Res Rev* 2006; 22: 274-283
- 57 Gotoh J, Obata M, Yoshie M, Kasai S, Ogawa K. Cyclin D1 over-expression correlates with beta-catenin activation, but not with H-ras mutations, and phosphorylation of Akt, GSK3 beta and ERK1/2 in mouse hepatic carcinogenesis. *Carcinogenesis* 2003; 24: 435-442
- 58 Kershenevich Stalnikowitz D, Weissbrod AB. Liver fibrosis and inflammation. A review. *Ann Hepatol* 2003; 2: 159-163
- 59 Nagahara N, Katayama A. Post-translational regulation of mercaptopyruvate sulfurtransferase via a low redox potential cysteine-sulfenate in the maintenance of redox homeostasis. *J Biol Chem* 2005; 280: 34569-34576
- 60 Fitzpatrick PF. The aromatic amino acid hydroxylases. *Adv Enzymol Relat Areas Mol Biol* 2000; 74: 235-294

ically linked variable of neuraminidase inhibitors for influenza viruses.

doi:10.1016/j.antiviral.2008.01.087

74

Sub-optimal Protease Inhibition of HIV-1: Effects on Virion Morphogenesis and RNA Maturation

Michael D. Moore^{1,*}, William Fu², Roger G. Ptak², Wei-Shau Hu¹

¹ HIV Drug Resistance Program, National Cancer Institute at Frederick, Frederick, USA; ² Southern Research Institute, Frederick, USA

During or soon after release of HIV-1 from an infected cell the virion initiates the process of maturation. The viral protease becomes activated, leading to the subsequent cleavage of the viral polyproteins Gag and GagPol into their constituent parts. As a result, an internal conical core condenses surrounding the viral nucleic acid and the particle becomes infectious. Concomitant with this global alteration in virion morphogenesis is a conformational change in the viral genomic RNA from a loosely associated dimer into a more thermodynamically stable form. Protease defective viruses are capable of virus release and viral RNA encapsidation, but these particles are non-infectious and immature due to an inability to carry out proteolytic cleavage. Within these particles the viral RNA is also observed to be in an immature state, demonstrating a link between the proteinase maturation and that of the nucleic acid. We have used sub-optimal concentrations (IC50 and IC90) of two protease inhibitor drugs (Lopinavir and Atazanavir) to demonstrate their effect on the Gag polyprotein processing and RNA properties of the treated virions. The results were then correlated to their effects on virion morphogenesis as determined by EM. The results show that even with high levels of viral inhibition (IC90) most of the viral protein is processed. However, a slight but significant increase in processing intermediates was detected upon drug exposure and a small decrease (2–3 °C when 50% of dimers remained) in overall thermostability of the viral RNA dimer was also observed. These defects correlated with an increase in immature particles as observed by EM, but the numbers of immature particles did not adequately account for the level of viral inhibition. These data suggest that the presence of small quantities of residual processing intermediates, within the viral particles, is capable of disproportionately inhibiting the viral replication cycle, without having comparative effects on either RNA maturation or virion core condensation.

doi:10.1016/j.antiviral.2008.01.088

75

Significance of 3 β -dehydroxysterol-D24-reductase (DHCR24) in life cycle of Hepatitis C virus

Tomohiro Nishimura^{1,*}, Masaaki Sato¹, Makoto Saito¹, Yuri Kasama¹, Michinori Kohara², Kyoko Kohara¹

¹ Kumamoto University, Kumamoto, Japan; ² RRINSHOKEN, Tokyo, Japan

Hepatitis C virus (HCV) causes persistent infection often progressing to hepatocellular carcinoma (HCC). We previously reported that full HCV genome-expressing HepG2 cells enhanced their clonogenic capacity after 44 days of passage (M6 44 days cells). We established monoclonal antibodies (MoAbs) against surface antigens on these cells. One of the MoAbs specifically recognized the molecule which was overexpressed in the cancerous region of livers of all HCV-positive HCC patients. It was identified as 24-dehydrocholesterol reductase (DHCR24), which was reported to be involved in cholesterol biosynthesis and hydrogen peroxide-induced cytotoxicity. The full-length HCV upregulated the transcription of DHCR24 in human liver cells in the presence of p53. Expression of HCV induced the upregulation of DHCR24 and p53, and was sustained in M6 44 days cells. However, activity of p21^{WAF1/CIP1} promoter in response to hydrogen peroxide was impaired in M6 44 days cells. This might be induced by the post-translational modification of p53, which was regulated by DHCR24. Thus, DHCR24 plays a critical role in the regulation of the response to HCV and hydrogen peroxide, and this pathway is a target of HCV during its persistent expression.

doi:10.1016/j.antiviral.2008.01.089

76

Inhibition of Human T-Cell Lymphotropic Virus Type-1 Integrase by Dicafeoylquinic Acids Extracted from Coffee (*Coffea arabica*) Seeds

Ordóñez Paula^{1,*}, Panay Joel¹, Cuesta-Astroz Yesid¹, Ceballos Carolina¹, Ceron Flavio¹, Moncayo Alejandro², Martha C. Domínguez¹, Sánchez Adalberto¹, Lareo Leonardo³, García-Vallejo Felipe¹

¹ Laboratory of Molecular Biology and Pathogenesis, Universidad del Valle, Cali, Columbia; ² Department of Chemistry, Universidad del Valle, Cali, Columbia; ³ Laboratory of Computational and Structural Biochemistry and Bioinformatics, Pontificia Universidad Javeriana, Bogotá, Columbia

Human T-cell lymphotropic virus type-1 (HTLV-1) replication depends on the viral enzyme integrase (IN) that mediates integration of a DNA copy of the virus into the host cell genome. Integrase represents a novel target to which antiviral agents might be directed. The C-terminal part of the HTLV-1 pol gene is predicted to encode the HTLV-1 IN; however, this protein has not yet been detected in virions or infected cells. In order to evaluate compounds with anti-HTLV IN activity, we extracted dicafeoylquinic acids (DCQAs) from coffee (*Coffea arabica*) seeds. Using a baculovirus system we expressed a 38-kDa IN



Comparative aspects on the role of polypyrimidine tract-binding protein in internal initiation of hepatitis C virus and picornavirus RNAs

T. Nishimura^{a,e}, M. Saito^a, T. Takano^{a,b,c}, A. Nomoto^d,
M. Kohara^b, K. Tsukiyama-Kohara^{a,b,c,*}

^aDepartment of Experimental Phylaxiology, Faculty of Medical and Pharmaceutical Sciences,
Kumamoto University 1-1-1, Honjo, Kumamoto 860-8556, Japan

^bDepartment of Microbiology and Cell Biology, The Tokyo Metropolitan Institute of Medical Science,
Tokyo 113-8613, Japan

^cLaboratory Animal Research Center, Institute of Medical Science, The University of Tokyo,
Tokyo 108-8639, Japan

^dGraduate School of Medicine, The University of Tokyo, Tokyo 113-0033, Japan

^eThe Chemo-Sero-Therapeutic Research Institute, Tokyo 869-1298, Japan

Accepted 6 July 2007

Abstract

We compared the effects of polypyrimidine tract-binding protein (PTB) on hepatitis C virus (HCV genotype 1a), encephalomyocarditis virus (EMCV) and poliovirus internal ribosome entry site (IRES) activities *in vitro*. It bound strongly to EMCV IRES, but weakly to PV and HCV RNAs. PV IRES showed the strongest dependency to PTB and it showed less than one-tenth of IRES activity after the immuno-depletion of PTB from HeLa S10 lysate with pre-coated anti-PTB IgG beads, comparing to the normal IgG beads-treated S10 lysate. EMCV IRES activity was approximately 40% of that of normal control after PTB depletion.

*Corresponding author. Department of Experimental Phylaxiology, Faculty of Medical and Pharmaceutical Sciences, Kumamoto University 1-1-1, Honjo, Kumamoto 860-8556, Japan.
Tel./fax: +81 96 373 5560.

E-mail address: kkohara@kumamoto-u.ac.jp (K. Tsukiyama-Kohara).

0147-9571/\$ - see front matter © 2007 Elsevier Ltd. All rights reserved.
doi:10.1016/j.cimid.2007.07.002