

- Bartenschlager, R., Lohmann, V., 2000. Replication of hepatitis C virus. *J. Gen. Virol.* 81 (Pt 7), 1631–1648.
- Boulant, S., Targett-Adams, P., McLauchlan, J., 2007. Disrupting the association of hepatitis C virus core protein with lipid droplets correlates with a loss in production of infectious virus. *J. Gen. Virol.* 88 (Pt 8), 2204–2213.
- Brasaele, D.L., Dolios, G., Shapiro, L., Wang, R., 2004. Proteomic analysis of proteins associated with lipid droplets of basal and lipolytically stimulated 3T3-L1 adipocytes. *J. Biol. Chem.* 279 (45), 46835–46842.
- Choo, Q.L., Kuo, G., Weiner, A.J., Overby, L.R., Bradley, D.W., Houghton, M., 1989. Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 244 (4902), 359–362.
- Fukushima, M., Tanaka, Y., Sato, S., Ono, Y., Nitahara-Kasahara, Y., Suzuki, T., Miyamura, T., Hanada, K., Nishijima, M., 2006. Enhancement of de novo fatty acid biosynthesis in hepatic cell line Huh7 expressing hepatitis C virus core protein. *Biol. Pharm. Bull.* 29 (9), 1958–1961.
- Harada, T., Kim, D.W., Sagawa, K., Suzuki, T., Takahashi, K., Saito, I., Matsuura, Y., Miyamura, T., 1995. Characterization of an established human hepatoma cell line constitutively expressing non-structural proteins of hepatitis C virus by transfection of viral cDNA. *J. Gen. Virol.* 76 (Pt 5), 1215–1221.
- Hope, R.G., McLauchlan, J., 2000. Sequence motifs required for lipid droplet association and protein stability are unique to the hepatitis C virus core protein. *J. Gen. Virol.* 81 (Pt 8), 1913–1925.
- Hope, R.G., Murphy, D.J., McLauchlan, J., 2002. The domains required to direct core proteins of hepatitis C virus and GB virus-B to lipid droplets share common features with plant oleosin proteins. *J. Biol. Chem.* 277 (6), 4261–4270.
- Johnston, J.A., Ward, C.L., Kopito, R.R., 1998. Aggresomes: a cellular response to misfolded proteins. *J. Cell Biol.* 143 (7), 1883–1898.
- Kang, S.M., Shin, M.J., Kim, J.H., Oh, J.W., 2005. Proteomic profiling of cellular proteins interacting with the hepatitis C virus core protein. *Proteomics* 5 (8), 2227–2237.
- Kato, T., Date, T., Miyamoto, M., Sugiyama, M., Tanaka, Y., Orito, E., Ohno, T., Sugihara, K., Hasegawa, I., Fujiwara, K., Ito, K., Ozasa, A., Mizokami, M., Wakita, T., 2005. Detection of anti-hepatitis C virus effects of interferon and ribavirin by a sensitive replicon system. *J. Clin. Microbiol.* 43 (11), 5679–5684.
- Kim, W., Oe Uim, S., Kim, J.S., Ryu, Y.H., Byeon, J.Y., Kim, H.J., Kim, Y.L., Heo, J.S., Park, Y.M., Jung, G., 2003. Comparison of proteome between hepatitis B virus- and hepatitis C virus-associated hepatocellular carcinoma. *Clin. Cancer Res.* 9 (15), 5493–5500.
- Kuo, G., Choo, Q.L., Alter, H.J., Gitnick, G.L., Redeker, A.G., Purcell, R.H., Miyamura, T., Dienstag, J.L., Alter, M.J., Stevens, C.E., et al., 1989. An assay for circulating antibodies to a major etiologic virus of human non-A, non-B hepatitis. *Science* 244 (4902), 362–364.
- Lieber, J.C., Evans, R.M., 1996. Disruption of the vimentin intermediate filament system during adipose conversion of 3T3-L1 cells inhibits lipid droplet accumulation. *J. Cell. Sci.* 109 (Pt 13), 3047–3058.
- Matto, M., Rice, C.M., Aroeti, B., Glenn, J.S., 2004. Hepatitis C virus core protein associates with detergent-resistant membranes distinct from classical plasma membrane rafts. *J. Virol.* 78 (21), 12047–12053.
- McLauchlan, J., Lemberg, M.K., Hope, G., Martoglio, B., 2002. Intramembrane proteolysis promotes trafficking of hepatitis C virus core protein to lipid droplets. *EMBO J.* 21 (15), 3980–3988.
- Miyazari, Y., Atsuzawa, K., Usuda, N., Watashi, K., Hishiki, T., Zayas, M., Bartenschlager, R., Wakita, T., Hijikata, M., Shimotohno, K., 2007. The lipid droplet is an important organelle for hepatitis C virus production. *Nat. Cell Biol.* 9 (9), 1089–1097.
- Morishii, K., Okabayashi, T., Nakai, K., Moriya, K., Koike, K., Murata, S., Chiba, T., Tanaka, K., Suzuki, R., Suzuki, T., Miyamura, T., Matsuura, Y., 2003. Proteasome activator PA28gamma-dependent nuclear retention and degradation of hepatitis C virus core protein. *J. Virol.* 77 (19), 10237–10249.
- Morimoto, T., Fujita, M., Kawamura, T., Sunagawa, Y., Takaya, T., Wada, H., Shimatsu, A., Kita, T., Hasegawa, K., 2008. Myocardial regulation of p300 and p53 by doxorubicin involves ubiquitin pathways. *Circ. J.* 72 (9), 1506–1511.
- Moriya, K., Yotsuyanagi, H., Shintani, Y., Fujie, H., Ishibashi, K., Matsuura, Y., Miyamura, T., Koike, K., 1997. Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J. Gen. Virol.* 78 (Pt 7), 1527–1531.
- Moriya, K., Fujie, H., Shintani, Y., Yotsuyanagi, H., Tsutsumi, T., Ishibashi, K., Matsuura, Y., Kimura, S., Miyamura, T., Koike, K., 1998. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat. Med.* 4 (9), 1065–1067.
- Randall, G., Panis, M., Cooper, J.D., Tellinghuisen, T.L., Sukhodolets, K.E., Pfeffer, S., Landthaler, M., Landgraf, P., Kan, S., Lindenbach, B.D., Chien, M., Weir, D.B., Russo, J.J., Ju, J., Brownstein, M.J., Sheridan, R., Sander, C., Zavolan, M., Tuschl, T., Rice, C.M., 2007. Cellular cofactors affecting hepatitis C virus infection and replication. *Proc. Natl. Acad. Sci. U. S. A.* 104 (31), 12884–12889.
- Reed, K.E., Rice, C.M., 2000. Overview of hepatitis C virus genome structure, polyprotein processing, and protein properties. *Curr. Top. Microbiol. Immunol.* 242, 55–84.
- Ruggieri, A., Harada, T., Matsuura, Y., Miyamura, T., 1997. Sensitization to Fas-mediated apoptosis by hepatitis C virus core protein. *Virology* 229 (1), 68–76.
- Sarria, A.J., Nordeen, S.K., Evans, R.M., 1990. Regulated expression of vimentin cDNA in cells in the presence and absence of a preexisting vimentin filament network. *J. Cell Biol.* 111 (2), 553–565.
- Sato, S., Fukushima, M., Yamakawa, Y., Natsume, T., Suzuki, T., Shoji, I., Aizaki, H., Miyamura, T., Nishijima, M., 2006. Proteomic profiling of lipid droplet proteins in hepatoma cell lines expressing hepatitis C virus core protein. *J. Biochem. (Tokyo)* 139 (5), 921–930.
- Schweitzer, S.C., Evans, R.M., 1998. Vimentin and lipid metabolism. *Sub-cell. Biochem.* 31, 437–462.
- Shevchenko, A., Wilm, M., Vorm, O., Jensen, O.N., Podtelejnikov, A.V., Neubauer, G., Shevchenko, A., Mortensen, P., Mann, M., 1996. A strategy for identifying gel-separated proteins in sequence databases by MS alone. *Biochem. Soc. Trans.* 24 (3), 893–896.
- Shi, S.T., Lee, K.J., Aizaki, H., Hwang, S.B., Lai, M.M., 2003. Hepatitis C virus RNA replication occurs on a detergent-resistant membrane that cofractionates with caveolin-2. *J. Virol.* 77 (7), 4160–4168.
- Shirakura, M., Murakami, K., Ichimura, T., Suzuki, R., Shimoji, T., Fukuda, K., Abe, K., Sato, S., Fukushima, M., Yamakawa, Y., Nishijima, M., Morishii, K., Matsuura, Y., Wakita, T., Suzuki, T., Howley, P.M., Miyamura, T., Shoji, I., 2007. E6AP ubiquitin ligase mediates ubiquitination and degradation of hepatitis C virus core protein. *J. Virol.* 81 (3), 1174–1185.
- Suzuki, R., Tamura, K., Li, J., Ishii, K., Matsuura, Y., Miyamura, T., Suzuki, T., 2001. Ubiquitin-mediated degradation of hepatitis C virus core protein is regulated by processing at its carboxyl terminus. *Virology* 280 (2), 301–309.
- Suzuki, T., Ishii, K., Aizaki, H., Wakita, T., 2007. Hepatitis C viral life cycle. *Adv. Drug Deliv. Rev.* 59 (12), 1200–1212.
- Tanaka, T., Inoue, K., Hayashi, Y., Abe, A., Tsukiyama-Kohara, K., Nuriya, H., Aoki, Y., Kawaguchi, R., Kubota, K., Yoshida, M., Koike, M., Tanaka, S., Kohara, M., 2004. Virological significance of low-level hepatitis B virus infection in patients with hepatitis C virus associated liver disease. *J. Med. Virol.* 72 (2), 223–229.
- Wakita, T., Pietschmann, T., Kato, T., Date, T., Miyamoto, M., Zhao, Z., Murthy, K., Habermann, A., Krausslich, H.G., Mizokami, M., Bartenschlager, R., Liang, T.J., 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* 11 (7), 791–796.



Characterization of infectious hepatitis C virus from liver-derived cell lines

Daisuke Akazawa^{a,b}, Tomoko Date^b, Kenichi Morikawa^b, Asako Murayama^b, Noriaki Omi^{a,b}, Hitoshi Takahashi^{a,b}, Noriko Nakamura^a, Koji Ishii^b, Tetsuro Suzuki^b, Masashi Mizokami^c, Hidenori Mochizuki^a, Takaji Wakita^{b,*}

^a Pharmaceutical Research Laboratories, Toray Industries, Inc., Kanagawa, Japan

^b Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan

^c Department of Clinical Molecular Informative Medicine, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan

ARTICLE INFO

Article history:
Received 20 September 2008
Available online xxxxx

Keywords:
Cell culture
Hepatitis C virus
Infectivity
Particle
Replicon

ABSTRACT

The efficient production of infectious HCV from the JFH-1 strain is restricted to the Huh7 cell line and its derivatives. However, the factors involved in this restriction are unknown. In this study, we examined the production of infectious HCV from other liver-derived cell lines, and characterized the produced viruses. Clones of the Huh7, HepG2, and IMY-N9, harboring the JFH-1 full-genomic replicon, were obtained. The supernatant of each cell clone exhibited infectivity for naive Huh7. Each infectious supernatant was then characterized by sucrose density gradient. For all of the cell lines, the main peak of the HCV-core protein and RNA exhibited at approximately 1.15 g/mL of buoyant density. However, the supernatant from the IMY-N9 differed from that of Huh7 in the ratio of core:RNA at 1.15 g/mL and significant peaks were also observed at lower density. The virus particles produced from the different cell lines may have different characteristics.

© 2008 Elsevier Inc. All rights reserved.

Hepatitis C virus (HCV) is an enveloped virus that belongs to the *Hepacivirus* genus of the *Flaviviridae* family. HCV is a human pathogen and HCV infection is a major cause of chronic hepatitis, liver cirrhosis and hepatic carcinoma. The main therapy for HCV is treatment with pegylated-interferon and ribavirin. However, these agents show little effect for patients that have a high titer of HCV-RNA, genotype 1. Thus, it is necessary to develop new, more effective therapies and preventive treatments to counteract HCV infection. It was discovered that a genotype 2a strain of HCV, JFH-1, can efficiently replicate in the Huh7 cell line [1], and an *in vitro* culture model of infectious HCV has also been successfully developed using the JFH-1 genome [2–4]. Recently, it has become possible to produce various chimeric HCV by replacement of the JFH-1 structural protein region with that of other strains. The HCV particles produced from such chimera are expected to lead to the development of a HCV vaccine, and new anti-HCV pharmaceuticals.

The infectious HCV-derived JFH-1 genome was developed using the human hepatoma Huh7 cell line [5]. Although the sub-genomic replicon RNA of JFH-1 can autonomously replicate, not only in Huh7 cells, but in other human liver [6], non-hepatic [7], and mouse [8] cells, infectious HCV production has been restricted to Huh7-derived cells. In this study, we undertook a comparative study of infectious HCV particles produced from different cell lines including Huh7. Infectious HCV particles were successfully produced into the culture media and characterized.

Materials and methods

Cell culture. Huh7, Huh7.5.1 ([3], a generous gift from Dr. Francis V. Chisari), HepG2, and IMY-N9 cells were cultured at 37 °C in 5% CO₂. The HepG2 cells were cultured in modified Eagle's medium containing 10% fetal bovine serum. All of the other cells were cultured in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum, as described previously [6].

Plasmids. The pFGR-JFH1 and pFGR-JFH1/deltaE12 plasmids, encoding the full-genomic replicon, and envelope-deleted replicons, respectively, were generated as previously described [9].

RNA synthesis. RNA synthesis was performed as described previously [2]. Briefly, the pFGR-JFH1 plasmid was digested with XbaI and then treated with Mung Bean nuclease (New England Biolabs, Beverly, MA). The digested plasmid DNA fragment was then purified and used as a template for RNA synthesis. HCV-RNA was synthesized *in vitro* using a MEGAscript™ T7 kit (Ambion, Austin, TX). The synthesized RNA was treated with DNaseI, followed by acid phenol extraction to remove any remaining template DNA.

Establishment of replicon cells. Cell lines harboring FGR-JFH1 replicons were produced as described previously [9]. Briefly, trypsinized cells were washed with Opti-MEM™ reduced-serum medium (Invitrogen, Carlsbad, CA) and resuspended at 7.5×10^6 cells/mL with Cytomix buffer [1]. RNA (10 µg), synthesized from pFGR-JFH1, was mixed with 400 µL of cell suspension and transferred to an electroporation cuvette (Precision Universal Cuvettes, Thermo Hybrid, Middlesex, UK). The cells were then pulsed at 260V and 950 µF with the Gene Pulser II™ apparatus (Bio-Rad,

* Corresponding author. Fax: +81 3 5285 1161.
E-mail address: wakita@nih.go.jp (T. Wakita).

Hercules, CA). Transfected cells were immediately transferred to 10-cm culture dishes, each containing 8 mL of culture medium. G418 (0.8–1.0 mg/mL) (Nacalai Tesque, Kyoto, Japan) was added to the culture medium at 16–24 h after transfection. Culture medium, supplemented with G418, was replaced twice per week. Three weeks after transfection, sparsely grown G418-resistant colonies were independently isolated using a cloning cylinder (Asahi Techno Glass Co., Tokyo, Japan), and were expanded.

Preparation of supernatants from FGR-JFH1 replicon cells. Culture media was collected from Huh7, IMY-N9, and HepG2 cell lines harboring the FGR-JFH1 replicon and was passed through a 0.45- μ m filter. Filtrate culture media was then filtered and concentrated 50-fold using Amicon Ultra-15 (100,000 Molecular weight cut off; Millipore, Bedford, MA), and stored at -80°C until use.

Assay of infection of naïve Huh7 cells. Infection of naïve Huh7 cells were assayed by immunofluorescence and colony formation assays. For the immunofluorescence assay naïve Huh7.5.1 cells were seeded at 1×10^4 cells/well in an 8-well chamber slide (Becton Dickinson, Franklin Lakes, NJ), cultured overnight and then inoculated with diluted culture media containing infectious HCV particles (1×10^6 HCV-RNA copies). At 72 h after inoculation, the cells were fixed in acetone/methanol (1:1) for 10 min at -20°C , and the infected foci were visualized by immunofluorescence as follows.

An anti-core HCV protein monoclonal antibody 2H9 [2] was added to the cells at 50 $\mu\text{g}/\text{mL}$ in BlockAce (Dainippon Sumitomo Pharma, Osaka, Japan). After incubation for 1 h at room temperature, the cells were washed and incubated with a 1:400 dilution of AlexaFluor 488-conjugated anti-mouse IgG (Molecular Probes, Eugene, OR) diluted in BlockAce. The cells were then washed, treated with DAPI solution (Sigma, Saint Louis, MO) at 0.1 $\mu\text{g}/\text{mL}$ and examined by Biozero fluorescence microscopy (Keyence, Osaka, Japan).

Colony formation assays were performed as described previously [9]. Briefly, naïve Huh7 cells were inoculated with culture supernatants from replicon-expressing cell lines for 2 h, and then cultured with complete medium. Inoculated cells were cultured for 3 weeks in medium supplemented with G418 (0.3 mg/mL). Cell survival was assessed by staining with crystal violet.

Titration of infectivity. The infectivity titer of the culture supernatants was determined on Huh7.5.1 cells by end point dilution and immunofluorescence as described above. Briefly, each sample was serially diluted 10-fold in DMEM-10% FBS and 100 μL was used to inoculate Huh7.5.1 cells. Infection was examined 72 h post-inoculation by immunofluorescence using a mouse monoclonal anti-core antibody and secondary anti-mouse IgG-Alexa 488 conjugated antibodies. Infectious foci were counted and the titer was calculated and expressed as focus forming units per mL (FFU/mL).

Sucrose density gradient analysis. Concentrated cell supernatants were layered on top of a preformed continuous 10–60% sucrose gradient in TNE buffer containing 10 mM Tris, pH7.5, 150 mM NaCl, and 0.1 mM EDTA. Gradients were centrifuged in an SW41 rotor (Beckman Coulter, Fullerton, CA) at 35,000 rpm for 16 h at 4°C , and fractions (400 μL each) were collected from the bottom of the tube. The density of each fraction was estimated by weighing a 100 μL drop from each fraction following a gradient run.

Quantification of HCV-core protein and RNA. The level of the HCV-core protein in culture supernatants or sucrose density gradient fractions, was assayed using an immunoassay as described elsewhere [10]. Viral RNA was isolated from harvested culture media, or sucrose density gradient fractions, using the QiaAmp Viral RNA Extraction kit (Qiagen, Tokyo, Japan). The copy number of HCV RNA was determined by real-time detection reverse transcription-polymerase chain reaction (RTD-PCR), using an ABI Prism 7500fast sequence detector system (Applied Biosystems, Tokyo, Japan) [11].

Results

Production of infectious HCV from human liver-derived cell lines

We first determined if it was possible to produce infectious HCV from cell lines other than Huh7. We selected the HepG2 and IMY-N9 cell lines to establish human liver-derived cell lines that enable replication of the JFH-1 genome [6]. Since full-genomic JFH-1 did not transiently replicate in these cells (data not shown), we established FGR-JFH1 replicon cells that stably replicate the JFH-1 genome. In the culture media obtained from these full-genomic replicon cells, HCV-RNA titers were detected by RTD-PCR. The titer of HCV-RNA was highest in the supernatant from an IMY-N9 cell clone and lowest from a HepG2 cell clone (Table 1). When naïve Huh7.5.1 cells were inoculated with culture supernatants from the replicon cells, infected cells could be detected by immunofluorescence using an anti-HCV-core protein antibody (Fig. 1A). These data suggested that HepG2 and IMY-N9 cells are able to produce infectious HCV.

We then compared the specific infectivity of the replicon containing culture supernatants from the different cells. Specific infectivity was calculated by dividing the infectious titer, calculated by immunofluorescence of infectious foci, of the culture media by the titer obtained for HCV-RNA. Using these calculations the culture media from Huh7 and HepG2 cells showed almost the same specific infectivity whereas that from IMY-N9 cell was relatively higher (Table 1). Thus the infectious HCV in the culture media might differ according to the cell line from which it was obtained.

To clarify the differences observed in specific infectivity, we next examined the ability of the various cellular supernatants to induce colony formation. For this assay naïve Huh7 cells were inoculated with culture media of the same HCV-RNA titer as that of the FGR-JFH1 virus and were cultured in G418-containing medium. Cell survival was assayed by staining with crystal violet, and the number of colonies formed was counted. Consistent with the specific infectivity results, the supernatant of the IMY-N9 replicon cell showed higher colony formation compared with that of Huh7 and HepG2 replicon cells (Fig. 1B and C). Thus IMY-N9 cells produce infectious HCV with a relatively higher infectivity than the other cell lines suggesting that the supernatant derived from the different replicon producing cells may differ.

Characterization of the FGR-JFH1 virus from different liver-derived cells

To further characterize potential differences between the viruses produced by the different cell lines we next characterized the FGR-JFH1 virus in the media of the different cell lines by sucrose density gradient analysis. Concentrated cell supernatants were layered on top of a preformed continuous 10–60% sucrose gradient and centrifuged. Twenty-four fractions were collected and the HCV-core protein and RNA was assayed in each fraction. The peak fraction of the HCV-core protein and that of the RNA coincided at a density of 1.15 g/mL in all supernatants. However, the supernatant of the IMY-N9 cells showed different profiles for both the HCV-core protein and RNA compared to those of Huh7. Thus the IMY-N9 cells had a different ratio of

Table 1
Infectivity of the supernatant of replicon cell lines.

Producing cell	HCV-RNA (copies/mL)	Infectious titer (FFU/mL)	Specific infectivity (FFU/RNA copy)
Huh7	$1.36 \pm 0.02 \times 10^8$	$1.30 \pm 0.32 \times 10^4$	9.56×10^{-5}
IMY-N9	$2.80 \pm 0.04 \times 10^8$	$3.75 \pm 0.38 \times 10^4$	1.34×10^{-4}
HepG2	$8.80 \pm 0.75 \times 10^7$	$7.70 \pm 1.41 \times 10^3$	7.96×10^{-5}




Fig. 1. Naive Huh7 cell infection assay of JFH-1 full-genomic replicon cell culture supernatants. (A) JFH-1 full-genomic replicon (FGR-JFH1) cells were established in Huh7, HepG2, and IMY-N9 cell lines. Supernatants derived from Huh7 (left), HepG2 (middle), and IMY-N9 (right) cells (1×10^6 HCV-RNA copies) were inoculated into naive Huh7.5.1 cells (1×10^4) for 48 h, and infected cells were then detected by immunofluorescence using an anti-core antibody (clone 2H9) (green). (B) Naive Huh7 cells (5×10^5) were inoculated with mock, Huh7, HepG2, and IMY-N9-derived supernatants (10^6 HCV-RNA copies per 10-cm dish) of FGR-JFH1 cells for 2 h. Inoculated cells were cultured for 3 weeks in complete medium supplemented with G418 (0.3 mg/mL), and G418-resistant cells were stained using crystal violet. (C) The number of G418-resistant colonies obtained in (B) was calculated when 10^5 or 10^6 copies of HCV-RNA were tested. Mean values of colony number were indicated in duplicate experiment. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

HCV-core protein and RNA at a density of 1.15 g/mL (RNA/Core ratio; Huh7: 511, IMY-N9: 133 copies/fmol) and also showed a secondary peak at lower density (approximately 1.05 g/mL). For all supernatants the peak of infectivity exhibited at a density of 1.10 g/mL that was slightly lower than that of the HCV-core protein and RNA peaks. Furthermore infectivity was barely detectable in the lower density fractions (Fig. 2) suggesting that the HCV-core protein and RNA that was detected at lower density was irrelevant for infectivity of the different supernatants.

We considered the possibility that the core protein and RNA in the lighter fractions may be due to cellular debris containing a replication complex. To determine if this might be the case we therefore analyzed the supernatants from Huh7 and IMY-N9 envelope-deleted replicon cells (FGR-JFH1/deltaE12). The HCV-core protein and RNA were detected in the supernatants of these cells although the titers were very low. These supernatants were not infective for naive Huh7 cells (data not shown). Furthermore, analysis of the concentrated supernatants of these cell lines by sucrose density gradient analysis detected both the HCV-core protein and RNA, and the major peaks of HCV-RNA were detected in the lower density (approximately 1.10 g/mL) fractions (Fig. 3). However, the profiles of HCV-core protein and RNA did not coincide for either cell line.

Discussion

Infectious HCV can be produced in cell culture by using the JFH-1 genome. This system permits investigation of various aspects of the HCV life cycle such as the steps of entry into cells, replication, and secretion. Infectious HCV derived from JFH-1 is robustly produced in Huh7 cell lines [2,3], and the infectious particles have been characterized. However the difficulty in robustly producing

infectious HCV from other cell lines prevents a comparative study of HCV production among different cell lines. In this study, we compared infectious HCV production in Huh7 with that of other cell lines, and characterized the viruses produced.

First, we established Huh7, IMY-N9, and HepG2 FGR-JFH1 replicon cells. These cell lines were able to replicate the JFH-1 sub-genomic replicon [6]. The HCV-core protein and RNA were detected in all of the supernatants and all of these supernatants showed infectivity for naive Huh7. Infectivity was evaluated by transient infection and colony formation assays. These assays indicated that the infectious supernatant from IMY-N9 cell had higher infectivity than the other cell lines for naive Huh7 cells.

Next, we characterized each supernatant by sucrose density gradient analysis, which revealed both similarities and differences among the infectious supernatants. All samples showed typical peaks at 1.15 g/mL buoyant density for HCV-core protein and RNA, and infectious fractions showed an almost identical buoyant density of 1.10 g/mL. However, the supernatant from the IMY-N9 cells showed a difference in the core/RNA ratio at a density of 1.15 g/mL and higher secondary peak of HCV-core protein and RNA at a lower density (approximately 1.05 g/mL). Since the fractions at lower density did not correlate with infectivity, it is believed that the component at lower density does not contain infectious HCV particles but rather cellular debris that contains HCV proteins, RNA, and lipids [12]. HCV can associate with lipoprotein [13,14], and is secreted with VLDL [15]. Thus, the observed differences in the HCV-producing cells may derive from differences in lipoprotein synthesis. However, it is also possible that the components migrating at lower density contain virus particles. The deletion mutant of FGR-JFH1 (FGR-JFH1/deltaE12) did replicate in Huh7 and IMY-N9 cells, and these replicon cells secreted the HCV-core protein into the culture media, although at low levels. HCV-RNA was also detected in the same culture




Fig. 2. Density gradient analysis of infectious HCV derived from Huh7 and IMY-N9 cells. Concentrated supernatants of Huh7 cells (A) and IMY-N9 cells (B) were layered on top of a preformed continuous 10–60% sucrose gradient in TNE buffer. The gradients were centrifuged in a SW41 rotor at 35,000 rpm for 16 h at 4°C, and fractions (400 μ L each) were collected from the bottom of the tube. The buoyant density (closed circles), HCV-core protein (closed diamonds), HCV-RNA (open diamonds) and infectivity for naïve Huh7.5.1 cells (shown in gray) was detected in each fraction as described in Materials and methods.

medium, and the profile of this HCV-RNA differed from that of the HCV-core protein in sucrose density gradient analysis. Thus, the peak fractions containing the HCV-core protein and RNA from the supernatant of FGR-JFH1/deltaE12 cells were different from the peak fractions from that of FGR-JFH1 cells. Therefore it is possible that all of the peaks of HCV-core protein and RNA observed in the supernatant of FGR-JFH1 replicon cells may correlate to virus particles with different densities. However, the reason why they centrifuge at different densities is unclear. Interestingly, the supernatants from cells transfected with envelope-deleted replicon RNA exhibit non-identical HCV-core protein and RNA profiles on a sucrose density gradient. Envelope-deleted replicon RNA may have a decreased ability to form nucleocapsids although a detailed examination is necessary to establish this point.

We previously developed a method for infectious HCV production using the FGR-JFH1 [9], and have now succeeded in producing infectious HCV in the supernatant of cultured liver-derived cell lines harboring FGR-JFH1 RNA. Infectious HCV particles are useful for vaccine production and are considered good antigens for the generation of useful antibodies. Selection of an appropriate cell line is important for the production of HCV particles for vaccine development. The technique used in this study seemed to be appropriate for producing infectious HCV in various cell lines [8].




Fig. 3. Density gradient analysis of supernatants derived from Huh7 and IMY-N9 cells transfected with FGR-JFH1/deltaE12 RNA. Concentrated supernatants from Huh7 (A) and IMY-N9 (B) cells were analyzed by sucrose density gradient as described in the legend to Fig. 2. The buoyant density (closed circles), HCV-core protein (closed diamonds) and HCV-RNA (open diamonds) was analyzed in each fraction.

A second advantage of using HepG2 and IMY-N9 cells for the production of virus particles is that these parental cell lines, unlike the Huh7 cell line, do not express the CD81 molecule on the cell surface, however, the expression on cell clones used in this study was not confirmed. This means that the FGR-JFH1 replicon of these cell lines may have a single cycle of HCV production, encompassing replication, assembly, budding and secretion, and do not show HCV permissiveness. These cells should therefore be useful for the discovery of drugs targeted against HCV assembly and secretion.

Acknowledgments

This work was partially supported by a grant-in-aid for Scientific Research from the Japan Society for the Promotion of Science and from the Ministry of Health, Labor, and Welfare of Japan by the Research on Health Sciences Focusing on Drug Innovation from the Japan Health Sciences Foundation. Huh7 and Huh7.5.1 was a kind gift from Dr. Francis V. Chisari.

References

- [1] T. Kato, T. Date, M. Miyamoto, A. Furusaka, K. Tokushige, M. Mizokami, T. Wakita, Efficient replication of the genotype 2a hepatitis C virus subgenomic replicon, *Gastroenterology* 125 (2003) 1808–1817.

- [2] T. Wakita, T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H.G. Krausslich, M. Mizokami, R. Bartenschlager, T.J. Liang, Production of infectious hepatitis C virus in tissue culture from a cloned viral genome, *Nat. Med.* 11 (2005) 791–796.
- [3] J. Zhong, P. Gastaminza, G. Cheng, S. Kapadia, T. Kato, D.R. Burton, S.F. Wieland, S.L. Uprichard, T. Wakita, F.V. Chisari, Robust hepatitis C virus infection in vitro, *Proc. Natl. Acad. Sci. USA* 102 (2005) 9294–9299.
- [4] B.D. Lindenbach, M.J. Evans, A.J. Syder, B. Wolk, T.L. Tellinghuisen, C.C. Liu, T. Maruyama, R.O. Hynes, D.R. Burton, J.A. McKeating, C.M. Rice, Complete replication of hepatitis C virus in cell culture, *Science* 309 (2005) 623–626.
- [5] H. Nakabayashi, K. Taketa, K. Miyano, T. Yamane, J. Sato, Growth of human hepatoma cells lines with differentiated functions in chemically defined medium, *Cancer Res.* 42 (1982) 3858–3863.
- [6] T. Date, T. Kato, M. Miyamoto, Z. Zhao, K. Yasui, M. Mizokami, T. Wakita, Genotype 2a hepatitis C virus subgenomic replicon can replicate in HepG2 and IMY-N9 cells, *J. Biol. Chem.* 279 (2004) 22371–22376.
- [7] T. Kato, T. Date, M. Miyamoto, Z. Zhao, M. Mizokami, T. Wakita, Nonhepatic cell lines HeLa and 293 support efficient replication of the hepatitis C virus genotype 2a subgenomic replicon, *J. Virol.* 79 (2005) 592–596.
- [8] S.L. Uprichard, J. Chung, F.V. Chisari, T. Wakita, Replication of a hepatitis C virus replicon clone in mouse cells, *Virol. J.* 3 (2006) 89.
- [9] T. Date, M. Miyamoto, T. Kato, K. Morikawa, A. Murayama, D. Akazawa, J. Tanabe, S. Sone, M. Mizokami, T. Wakita, An infectious and selectable full-length replicon system with hepatitis C virus JFH-1 strain, *Hepatol. Res.* 37 (2007) 433–443.
- [10] K. Aoyagi, C. Ohue, K. Iida, T. Kimura, E. Tanaka, K. Kiyosawa, S. Yagi, Development of a simple and highly sensitive enzyme immunoassay for hepatitis C virus core antigen, *J. Clin. Microbiol.* 37 (1999) 1802–1808.
- [11] T. Takeuchi, A. Katsume, T. Tanaka, A. Abe, K. Inoue, K. Tsukiyama-Kohara, R. Kawaguchi, S. Tanaka, M. Kohara, Real-time detection system for quantification of hepatitis C virus genome, *Gastroenterology* 116 (1999) 636–642.
- [12] T. Pietschmann, V. Lohmann, A. Kaul, N. Krieger, G. Rinck, G. Rutter, D. Strand, R. Bartenschlager, Persistent and transient replication of full-length hepatitis C virus genomes in cell culture, *J. Virol.* 76 (2002) 4008–4021.
- [13] M. Monazahian, S. Kippenberger, A. Muller, H. Seitz, I. Bohme, S. Grethe, R. Thomssen, Binding of human lipoproteins (low, very low, high density lipoproteins) to recombinant envelope proteins of hepatitis C virus, *Med. Microbiol. Immunol.* 188 (2000) 177–184.
- [14] R. Thomssen, S. Bonk, C. Propfe, K.H. Heermann, H.G. Kochel, A. Uy, Association of hepatitis C virus in human sera with beta-lipoprotein, *Med. Microbiol. Immunol.* 181 (1992) 293–300.
- [15] S.U. Nielsen, M.F. Bassendine, A.D. Burt, C. Martin, W. Pumechockchai, G.L. Toms, Association between hepatitis C virus and very-low-density lipoprotein (VLDL)/LDL analyzed in iodixanol density gradients, *J. Virol.* 80 (2006) 2418–2428.

Interaction of Hepatitis C Virus Nonstructural Protein 5A with Core Protein Is Critical for the Production of Infectious Virus Particles[†]

Takahiro Masaki,¹ Ryosuke Suzuki,¹ Kyoko Murakami,¹ Hideki Aizaki,¹ Koji Ishii,¹ Asako Murayama,¹ Tomoko Date,¹ Yoshiharu Matsuura,² Tatsuo Miyamura,¹ Takaji Wakita,¹ and Tetsuro Suzuki^{1*}

Department of Virology II, National Institute of Infectious Diseases, Shinjuku-ku, Tokyo 162-8640, Japan,¹ and Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka University, Suita-shi, Osaka 565-0871, Japan²

Received 17 April 2008/Accepted 22 May 2008

Nonstructural protein 5A (NS5A) of the hepatitis C virus (HCV) possesses multiple and diverse functions in RNA replication, interferon resistance, and viral pathogenesis. Recent studies suggest that NS5A is involved in the assembly and maturation of infectious viral particles; however, precisely how NS5A participates in virus production has not been fully elucidated. In the present study, we demonstrate that NS5A is a prerequisite for HCV particle production as a result of its interaction with the viral capsid protein (core protein). The efficiency of virus production correlated well with the levels of interaction between NS5A and the core protein. Alanine substitutions for the C-terminal serine cluster in domain III of NS5A (amino acids 2428, 2430, and 2433) impaired NS5A basal phosphorylation, leading to a marked decrease in NS5A-core interaction, disturbance of the subcellular localization of NS5A, and disruption of virion production. Replacing the same serine cluster with glutamic acid, which mimics the presence of phosphoserines, partially preserved the NS5A-core interaction and virion production, suggesting that phosphorylation of these serine residues is important for virion production. In addition, we found that the alanine substitutions in the serine cluster suppressed the association of the core protein with viral genome RNA, possibly resulting in the inhibition of nucleocapsid assembly. These results suggest that NS5A plays a key role in regulating the early phase of HCV particle formation by interacting with core protein and that its C-terminal serine cluster is a determinant of the NS5A-core interaction.

Hepatitis C virus (HCV) infection is a major public health problem and is prevalent in about 200 million people worldwide (27, 40, 42). Current protocols for treating HCV infection fail to produce a sustained virological response in as many as half of treated individuals, and many cases progress to chronic liver disease, including chronic hepatitis, cirrhosis, and hepatocellular carcinoma (15, 31, 35, 43).

HCV is a positive-strand RNA virus classified in the *Hepacivirus* genus within the *Flaviviridae* family (55). Its approximately 9.6-kb genome is translated into a single polypeptide of about 3,000 amino acids (aa), in which the structural proteins core, E1, and E2 reside in the N-terminal region. A crucial function of core protein is assembly of the viral nucleocapsid. The amino acid sequence of this protein is well conserved among different HCV strains compared to other HCV proteins. The nonstructural (NS) proteins NS3-NS5B are considered to assemble into a membrane-associated HCV RNA replicase complex. NS3 possesses the enzymatic activities of serine protease and RNA helicase, and NS4A serves as a cofactor for NS3 protease. NS4B plays a role in the remodeling of host cell membranes, probably to generate the site for the replicase assembly. NS5B functions as the RNA-dependent RNA polymerase. NS5A is known to play an important but undefined role in viral RNA replication.

NS5A is a phosphoprotein that can be found in basally phosphorylated (56 kDa) and hyperphosphorylated (58 kDa) forms (49). Comparative sequence analyses and limited proteolysis of recombinant NS5A have demonstrated that NS5A is composed of three domains (52). Domain I is relatively conserved among HCV genotypes compared to domains II and III. Analysis of the crystal structure of the conserved domain I that immediately follows the membrane-anchoring α -helix localized at the N terminus revealed a dimeric structure (53). The interface between protein molecules is characterized by a large, basic groove, which has been proposed as a site of RNA binding. In fact, its RNA binding property has been demonstrated biochemically (17). Domains II and III of NS5A are far less understood. Domain II contains a region referred to as the interferon sensitivity determining region, and this region and its C-terminal 26 residues have been shown to be essential for interaction with the interferon-induced, double-stranded RNA-dependent protein kinase (6–10, 38, 39, 48). Domain III includes a number of potential phosphoacceptor sites and is most likely involved in basal phosphorylation. This domain tolerates insertion of large heterologous sequences such as green fluorescent protein (GFP) and is not required for function of NS5A in HCV RNA replication (1, 34). However, a study with the recently established productive HCV cell culture system using genotype 2a isolate JFH-1 (28, 56, 58) demonstrated that while insertion of GFP within the NS5A region does not affect RNA replication, it does produce marked decreases in the production of infectious virus particles (41). This suggests that the C-terminal region of NS5A may affect virus particle production independent of RNA replication. Re-

* Corresponding author. Mailing address: Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan. Phone: 81 3 5285 1111. Fax: 81 3 5285 1161. E-mail: tesuzuki@nih.go.jp.

[†] Published ahead of print on 4 June 2008.

cently, Miyazaki et al. reported that the association of core protein with the NS proteins and replication complexes around lipid droplets (LDs) is critical for producing infectious viruses (33).

In the present study, we demonstrated that NSSA is a prerequisite for HCV particle production via its interaction with core protein, and we identified serine residues in the C-terminal region of NSSA that play an important role in virion production. Substitution of the serine residues with alanine residues inhibited not only the interaction of NSSA with core protein but also HCV RNA-core association and led to a decrease in HCV particle production with no effect on RNA replication.

MATERIALS AND METHODS

DNA construction. Plasmids pJFH1, which contains the full-length JFH-1 cDNA downstream of the T7 RNA promoter sequence, and pSGR-JFH1/Luc, in which the neomycin resistance gene of pSGR-JFH1 has been replaced by the firefly luciferase reporter gene, have been previously described (24, 56). To generate the fluorochrome gene-tagged full-length JFH-1 plasmid, pJFH1/NSSA-GFP, the region encompassing the RsrII site of NSSA and the BsrGI site of NSSB was amplified by PCR, the amplification product was cloned into pGEM-T Easy vector (Promega, Madison, WI), and the resultant plasmid was designated pGEM-JFH1/RsrII-BsrGI. A GFP reporter gene was amplified by PCR from pGreen Lantern-1 (Invitrogen, Carlsbad, CA) with primers containing the XhoI sequence and inserted, after restriction digestion with XhoI, into the XhoI site of pGEM-JFH1/RsrII-BsrGI. The resulting plasmid was digested by RsrII and BsrGI and ligated into pJFH1 similarly digested by RsrII and BsrGI to produce pJFH1/NSSA-GFP. For generation of the fluorochrome gene-tagged subgenomic reporter plasmid, pJFH1/NSSA-GFP was digested by RsrII and SnaBI and ligated into pSGR-JFH1/Luc similarly digested by RsrII and SnaBI. The mutations in the NSSA gene were generated by oligonucleotide-directed mutagenesis (57). To construct plasmids expressing N-terminally FLAG-tagged HCV core protein or hemagglutinin (HA)-tagged NSSA, DNA fragments encoding core protein or NSSA (wild type or mutants) were generated from the full-length JFH-1 cDNA by PCR. The core protein coding sequence, together with a FLAG sequence linked to its N terminus, was cloned into the pCAGGS vector (37). The coding sequences of NSSA, together with an HA sequence linked to their N termini, were also cloned into pCAGGS vectors. All PCR products were confirmed by automated nucleotide sequencing with an ABI Prism 3130 Avant Genetic Analyzer (Applied Biosystems, Tokyo, Japan).

Cells and viruses. The human hepatoma cell line, Huh-7, and JFH1/4-1 cells, which are Huh-7 cells carrying a subgenomic replicon of JFH-1 (32), were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with minimal essential medium nonessential amino acids (Invitrogen), 100 units/ml of penicillin, 100 µg/ml of streptomycin, and 10% fetal bovine serum (FBS) at 37°C in a 5% CO₂ incubator. Huh-c-p7 cells, which are Huh-7 cells stably expressing the proteins core to p7 derived from the JFH-1 strain (18), were incubated in DMEM containing 300 µg/ml of zeocin (Invitrogen). HCV particles derived from JFH-1 were produced by transient transfection of Huh-7 cells with *in vitro* transcribed RNA, as described previously (56, 58). Recombinant vaccinia virus strain DIs, which expresses the bacteriophage T7 RNA polymerase under the control of the vaccinia virus early/late promoter P7.5, was generated and propagated as previously described (19).

DNA transfection, immunoprecipitation (IP), and immunoblotting. For co-expression of FLAG-tagged core protein and HA-tagged NSSA, cells were seeded onto 35-mm wells of a six-well cell culture plate and cultured overnight. Plasmid DNAs (2 µg) were transfected into cells using TransIT-LT1 transfection reagent (Mirus, Madison, WI). Cells were harvested at 48 h posttransfection, washed three times with 1 ml of ice-cold phosphate-buffered saline (PBS), and suspended in 0.25 ml lysis buffer (20 mM Tris-HCl [pH 7.4] containing 135 mM NaCl, 1% Triton X-100, 0.05% sodium dodecyl sulfate [SDS], and 10% glycerol) supplemented with 50 mM NaF, 5 mM Na₂VO₄, 1 µg/ml leupeptin, and 1 mM phenylmethylsulfonyl fluoride (PMSF). Cell lysates were sonicated at 4°C for 5 min, incubated for 30 min at 4°C, and centrifuged at 14,000 × g for 5 min at 4°C. After preclearing, the supernatant was immunoprecipitated with 10 µl of anti-FLAG M2-agarose beads (Sigma, St. Louis, MO). For expression of the full-length HCV polyprotein, Huh-7 cells transfected with 10 µg of *in vitro* transcribed RNAs by electroporation were resuspended in 20 or 30 ml of culture

medium, and 10-ml aliquots were seeded into 100-mm culture dishes. At 72 h posttransfection, the cells were incubated in 0.5 ml of lysis buffer (20 mM Tris-HCl [pH 7.4] containing 135 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, and 10% glycerol) supplemented with 50 mM NaF, 5 mM Na₂VO₄, 1 µg/ml leupeptin, and 1 mM PMSF. After preclearing, the supernatant was immunoprecipitated with 5 µg of polyclonal anti-NSSA antibody (34a) or polyclonal anti-C/EBPB antibody (Santa Cruz Biotechnology, Santa Cruz, CA), and 20 µl of protein G-agarose beads (Invitrogen). The immunocomplex was precipitated with the beads by centrifugation at 800 × g for 30 s and then washed five times with lysis buffer by centrifugation. The proteins binding to the beads were boiled in 20 µl of SDS sample buffer and then subjected to SDS-12.5% polyacrylamide gel electrophoresis (PAGE). The proteins were transferred onto a polyvinylidene difluoride membrane (Immobilion; Millipore, Bedford, MA) and then reacted with a primary antibody and a secondary horseradish peroxidase-conjugated antibody. The immunocomplexes were visualized with an ECL Plus Western Blotting Detection System (GE Healthcare, Buckinghamshire, United Kingdom) and detected using a LAS-3000 imaging analyzer (Fujifilm, Tokyo, Japan).

In vitro synthesis of HCV RNA and RNA transfection. Plasmid DNAs were digested with XbaI and treated with mung bean nuclease (New England Biolabs, Ipswich, MA) to remove the four terminal nucleotides, resulting in the correct 3' end of the HCV cDNA. Digested DNAs were purified and used as templates for RNA synthesis. HCV RNA was synthesized *in vitro* using a MEGAscript T7 kit (Ambion, Austin, TX). Synthesized RNA was treated with DNase I (Ambion), followed by acid guanidinium thiocyanate-phenol-chloroform extraction to remove any remaining template DNA. Synthesized HCV RNAs were used for electroporation. Trypsinized Huh-7 cells were washed with Opti-MEM I reduced-serum medium (Invitrogen) and resuspended at 3 × 10⁶ cells/ml with Cytomix buffer (54). RNA was mixed with 400 µl of cell suspension and transferred into an electroporation cuvette (Precision Universal Cuvettes; Thermo Hybrid, Middlesex, United Kingdom). Cells were then pulsed at 260 V and 950 µF using a Gene Pulser II unit (Bio-Rad, Hercules, CA). Transfected cells were immediately transferred onto six-well culture plates or 100-mm culture dishes.

Luciferase assay. Cells were harvested at different time points posttransfection of subgenomic reporter replicons and lysed in passive lysis buffer (Promega). The luciferase activity in cells was determined using a luciferase assay system (Promega).

Quantification of HCV core protein. HCV core protein in transfected cells or cell culture supernatants was quantified using a highly sensitive enzyme immunoassay (Ortho HCV antigen ELISA Kit; Ortho Clinical Diagnostics, Tokyo, Japan). To determine intracellular core protein amounts, cell lysates were prepared as described previously (41). To determine the efficiency of core protein release, the ratio of extracellular core protein to total core protein (the sum of intra- and extracellular core protein amounts) was calculated.

Intra- and extracellular infectivity assay. Culture supernatants were harvested 72 h posttransfection, and virus titers were determined by a 50% tissue culture infectious dose (TCID₅₀) assay as described previously (28, 46). Virus titration was performed by seeding naive Huh-7 cells in 96-well plates at a density of 1 × 10⁴ cells/well. Samples were serially diluted fivefold in complete growth medium and used to infect the seeded cells (six wells per dilution). At 72 h after infection, the inoculated cells were fixed and immunostained with a mouse monoclonal anti-core protein antibody (2H9) (56), followed by an Alexa Fluor 488-conjugated anti-mouse immunoglobulin G (IgG) (Invitrogen). Wells that showed at least one core protein-expressing cell was counted as positive. Cell-associated infectivity was determined essentially as described previously (12, 47). Briefly, cells were extensively washed with PBS, scraped, and centrifuged for 3 min at 120 × g. Cell pellets were resuspended in 1 ml of DMEM containing 10% FBS and subjected to four cycles of freezing and thawing using dry ice and a 37°C water bath. Samples were then centrifuged at 2,400 × g for 10 min at 4°C to remove cell debris, and cell-associated infectivity was determined by TCID₅₀ assay.

Expression of HCV proteins using vaccinia viruses, metabolic labeling of cells, and radioimmunoprecipitation analysis. Metabolic labeling of cells and radioimmunoprecipitation analysis were performed as described by Huang et al. (17) with some modifications. A total of 4 × 10⁵ Huh-7 cells were seeded onto each well of six-well cell culture plates and cultured overnight. A 2-µg amount of subgenomic replicon DNAs carrying defined NSSA mutations was transfected into cells using TransIT-LT1 transfection reagent, and at 12 h posttransfection the cells were then infected at a multiplicity of infection of 10 with recombinant vaccinia viruses expressing the T7 RNA polymerase. After 40 h of transfection, cells were incubated in methionine- and cysteine-deficient DMEM (Invitrogen) or phosphate-deficient DMEM (Invitrogen) for 2 h and labeled for 6 h with [³⁵S]methionine and [³⁵S]cysteine (200 µCi/well; GE Healthcare) or

[32 P]orthophosphate (250 μ Ci/well; GE Healthcare). The cells were then washed twice with cold PBS and lysed with SDS lysis buffer (50 mM Tris-HCl [pH 7.6], 0.5% SDS, 1 mM EDTA, 20 μ g/ml of PMSF). The cell lysates were passed through a 27-gauge needle several times to shear cellular DNA. After a 10-min incubation at 75°C, the lysates were clarified by centrifugation and diluted five-fold with HNAET buffer (50 mM HEPES [pH 7.5], 150 mM NaCl, 0.67% bovine serum albumin, 1 mM EDTA, 0.33% Triton X-100). After preclearing by incubation with 20 μ l of protein G-agarose beads for 1 h at 4°C, the supernatant was incubated with 2 μ g of rabbit polyclonal anti-NS5A antibody overnight at 4°C. A 20- μ l aliquot of protein G-agarose beads was further added and incubated for 2 h at 4°C. The cell pellets were washed three times with 0.5 ml of HNAETS buffer (HNAET containing 0.5% SDS), followed by washing once with 0.5 ml of HNE buffer (50 mM HEPES [pH 7.5], 150 mM NaCl and 1 mM EDTA). After treatment with or without λ protein phosphatase (New England Biolabs), the cell pellets were suspended in 20 μ l of SDS sample buffer and boiled for 10 min. The proteins were resolved on 10% SDS-polyacrylamide gels and analyzed by autoradiography.

Subcellular fractionation analysis. All steps were carried out at 4°C in the presence of a protease inhibitor cocktail (Complete; Roche, Mannheim, Germany) as described previously (20), with some modifications. Cells were suspended in four cell volumes of homogenization buffer (50 mM NaCl, 10 mM triethylamine [pH 7.4], 1 mM EDTA), snap frozen in liquid nitrogen, stored at -80°C, and thawed in a water bath at room temperature. Supernatants (0.4 ml) were layered on linear 10-ml iodixanol gradients from 2.5 to 25% and centrifuged at 37,000 rpm for 3.5 h in an SW41 rotor (Beckman, Fullerton, CA), followed by collection of 0.8-ml fractions from the top. Each fraction was concentrated by Centricon YM30 (Millipore), separated by SDS-PAGE, and immunoblotted with a rabbit polyclonal anti-calnexin antibody (Stressgen Biotechnologies, Victoria, Canada), a mouse monoclonal anti-adipose differentiation-related protein (ADRP) antibody (Progen Biotechnik, Heidelberg, Germany), or a rabbit polyclonal anti-NS5A antibody. The core protein amount in each fraction was also determined by enzyme-linked immunosorbent assay (ELISA).

IP-RT-PCR. The process of cell lysis to RNA purification was carried out essentially as described by Johnson et al. (21) with some modifications. A total of 3×10^6 Huh-7 cells were transfected with 10 μ g of *in vitro* transcribed HCV RNAs and resuspended in 20 or 30 ml of culture medium, after which 10-ml aliquots were seeded into 100-mm culture dishes. At 72 h posttransfection, the cells were scraped and incubated in 500 μ l of hypotonic buffer (10 mM HEPES [pH 7.6], 1.5 mM MgCl₂, 10 mM KCl, 0.2 mM PMSF) per dish. The cells were passed through a 20-gauge needle several times, lysed with Nonidet P-40 at a final concentration of 1%, and incubated on ice for an additional 10 min. After centrifugation at 4,000 \times g at 4°C for 15 min, glycerol was added to the supernatants at a final concentration of 5%. The cell lysates were incubated with 20 μ l of protein G-agarose beads for 30 min at room temperature. After the cell lysates were removed from protein G-agarose beads, 5 μ g of mouse monoclonal anti-core protein antibody or normal mouse IgG (Sigma) as a negative control was added, and samples were incubated for an additional 1 h at room temperature. A 20- μ l aliquot of protein G-agarose beads per sample was added to the cell lysates and incubated for 1 h. After incubation, the beads were washed three times with wash buffer (10 mM Tris-HCl [pH 7.6], 100 mM KCl, 5 mM MgCl₂, and 1 mM dithiothreitol) and eluted in 100 μ l of elution buffer (50 mM Tris-HCl [pH 8.0], 1% SDS, and 10 mM EDTA) at 65°C for 10 min. After treatment with 100 μ g of proteinase K at 37°C for 30 min, the RNAs in immunocomplexes were isolated by acid guanidinium thiocyanate-phenol-chloroform extraction. Reverse transcriptase PCR (RT-PCR) was carried out using random hexamer and Superscript II RT (Invitrogen), followed by nested PCR with LA *Taq* DNA polymerase (TaKaRa, Shiga, Japan) and primer sets amplifying the fragments of nucleotides (nt) 129 to 2367 and nt 7267 to 9463 of the JFH-1 genome. To amplify the fragment of nt 129 to 2367, the sense primer 5'-CTGTGAGGAAC TACTGTCTT-3' and the antisense primer 5'-TCACGATGTTCTGGTGAA G-3' were used for first-round PCR; the sense primer 5'-CGGGAGAGCCAT AGTGG-3' and the antisense primer 5'-CATTCCGTGGTGTAGAGTCA-3' were used for second-round PCR. To amplify the fragment of nt 7267 to 9463, the sense primer 5'-GTCCAGGGTGCCCGTTCTGGACT-3' and the antisense primer 5'-CGGGTACACGACCTTCCAC-3' were used for first-round PCR; the sense primer 5'-CACCCTGTGCTGGTGTGCT-3' and the antisense primer 5'-GTGTACCTAGTGTGTGCCCTCA-3' were used for second-round PCR.

Indirect immunofluorescence analysis. Cells incubated for 3 days after transfection with JFH-1 RNAs were seeded in an eight-well chamber slide (BD Biosciences, San Jose, CA) and cultured overnight. The adherent cells were washed twice with PBS and fixed with 4% paraformaldehyde at room temperature. After a washing step with PBS, the cells were permeabilized with PBS containing 0.3% Triton X-100 and 2% FBS for 1 h at room temperature and

stained with a rabbit polyclonal anti-NS5A antibody and a mouse monoclonal anti-core protein antibody. The fluorescent secondary antibodies were Alexa Fluor 488- or Alexa Fluor 555-conjugated anti-rabbit or anti-mouse IgG antibodies (Invitrogen). Analyses of JFH-1 were performed on a Zeiss confocal laser scanning microscope LSM 510 (Carl Zeiss, Oberkochen, Germany).

RESULTS

Mutations of serine residues at the NS5A C terminus impair basal phosphorylation but have little effect on viral RNA replication. As demonstrated in a previous study, insertion of GFP into the NS5A C terminus does not significantly affect viral RNA replication but reduces the generation of infectious HCV particles (41). The C-terminal region of NS5A contains highly conserved serine residues that are involved in basal phosphorylation (1, 23, 49). To examine the involvement of the serine clusters (cluster 3-A [CL3A] and cluster 3-B [CL3B]) in the C-terminal region of NS5A in HCV particle production, we created mutated HCV genomes as well as subgenomic replicons carrying alanine substitutions for the conserved serine residues at aa 2384, 2388, 2390, and 2391 (residues are numbered according to the positions within the original JFH-1 polyprotein) (CL3A/SA); at aa 2428, 2430, and 2433 (CL3B/SA); or an in-frame deletion spanning aa 2384 to 2433 (Δ 2384-2433) (Fig. 1). A construct with an in-frame insertion of GFP (NS5A-GFP) was also generated as described previously for the Con1 isolate (34).

First, we analyzed the effects of the NS5A mutations on HCV RNA replication using a transient RNA replication assay using subgenomic luciferase reporter replicons (Fig. 2A) and found that the serine-to-alanine substitutions (CL3A/SA and CL3B/SA) did not affect viral RNA replication. NS5A-GFP and Δ 2384-2433 slightly reduced RNA replication, indicating that the mutations of the NS5A C terminus tested in this study do not critically affect RNA replication, which is consistent with previous reports (1, 34, 51).

Next, the phosphorylation status of the mutated NS5A was analyzed as described in Materials and Methods (Fig. 2B). NS5A was isolated from radiolabeled cells by IP and analyzed either directly by SDS-PAGE or after treatment with λ protein phosphatase. Analysis of 32 P-radiolabeled proteins revealed that the CL3A/SA, CL3B/SA, and Δ 2384-2433 mutations resulted in marked reduction of basal phosphorylation (Fig. 2B, compare lane 1 with lanes 3, 5, and 7 in the top panel). All 32 P-labeled NS5A proteins were sensitive to treatment with phosphatase (lanes 2, 4, 6, and 8). The possibility that loss of signal after dephosphorylation was due to contaminating proteases present in the phosphatase preparations can be ruled out because no degradation of the 35 S-labeled proteins was observed (Fig. 2B, bottom panel). These results suggest that mutations in the C-terminal serine cluster of NS5A impair basal phosphorylation but have no significant effect on viral RNA replication.

Effect of mutations introduced into the NS5A C terminus on the production of infectious HCV particles. To analyze HCV particle production from cells transfected with the *in vitro* transcribed viral genomic RNAs, we harvested supernatants and cells at 4, 24, 48, 72, and 96 h posttransfection and measured the amounts of core protein. As shown in Fig. 3A, comparable amounts of core proteins were detected in all transfected cells 4 h after transfection, reflecting unchanged




FIG. 1. Structures of HCV constructs used in this study. Schematic diagram of the NS5A structure according to Tellinghuisen et al. (52) is shown in the top panel. The three domains are indicated by white boxes and are separated by trypsin-sensitive regions with presumably low structural complexity (low-complexity sequence [LCS]). The numbers indicate amino acid residues within the original JFH-1 polyprotein. The names listed on the left represent full-length HCV constructs, subgenomic reporter replicons, or N-terminally HA-tagged NS5A constructs used in this study. NS5A-GFP carries a GFP insertion between aa 2394 and 2395 as indicated by a shaded box. CL3A/SA and CL3B/SA carry several serine-to-alanine substitutions in the NS5A C terminus constructed as described previously (1). HCV constructs from S2428A to S2430/2433A carry single or double serine-to-alanine substitutions generated by modification of the CL3B/SA construct. The Δ2384–2433 mutant possesses an in-frame deletion in the C-terminal region of NS5A. Amino acid substitutions are marked in bold and underlined. N and C represent N terminus and C terminus, respectively.

transfection efficiencies, and the kinetics of intracellular core protein levels was similar among transfectants. By contrast, core protein released from cells transfected either with the mutated genome of CL3B/SA, Δ2384–2433, or NS5A-GFP was more than 10-fold lower than that for the wild-type JFH-1 or CL3A/SA (Fig. 3B). Figure 3C shows the efficiency of core protein release from each transfectant, which is expressed as a percentage of the extracellular core protein level relative to the amount of total core protein (the sum of intra- and extracellular core protein). Core protein release efficiency with the wild type and CL3A/SA was 2 to 13% at 48 to 96 h after transfection, while only 1% or less of core protein was released in the cases of CL3B/SA, Δ2384–2433, and NS5A-GFP strains.

To further investigate production and release of infectious virus particles, naïve Huh-7 cells were infected with culture supernatants of cells harvested 72 h posttransfection, and infectious virus titers were determined by TCID₅₀ assay at 72 h after infection. Figure 3D shows that release of infectious virus particles from cells transfected with the genome of CL3B/SA or Δ2384–2433 mutants was markedly reduced (about 10,000-fold) compared to that from wild-type- or CL3A/SA-transfected cells (white bars). To examine whether such a decrease in infectious HCV in the culture supernatants was attributable to defective virion assembly or impaired release of virions, we determined cell-associated infectivity (Fig. 3D). Production of

intracellular infectious virions in CL3B/SA- and Δ2384–2433-transfected cells was strongly impaired in comparison with that in wild-type-transfected (~1,000-fold) and CL3A/SA-transfected (~100-fold) cells. Thus, the results suggest a potential role for the serine cluster at aa 2428, 2430, and 2433 of NS5A in assembly of infectious HCV particles. Among the NS5A mutations tested, CL3B/SA is of particular interest because this mutation leads to a marked reduction in HCV production with no impact on viral RNA replication.

Serine residues at aa 2428, 2430, and 2433 are important for the interaction between NS5A and core protein. Miyanari et al. reported that the association of core protein with NS proteins is critical for infectious HCV production and that mutations of the core protein and NS5A that cause these proteins to fail to associate with each other impair the production of infectious virus (33). Based on these observations and the findings noted above, we hypothesize that NS5A plays a key role in recruiting viral RNA, which is synthesized at the viral replication complex, to nucleocapsid formation via interaction between the NS5A C-terminal region and the core protein. To prove this, we analyzed the interaction of NS5A with the core protein by coimmunoprecipitation experiments. HA-tagged NS5A constructs carrying defined mutations were generated (Fig. 1) and coexpressed with the FLAG-tagged core protein in Huh-7 cells. As shown in Fig. 4A, coimmunoprecipitation of NS5A




FIG. 2. Mutations at the C terminus of NS5A impair basal phosphorylation and have only a minor impact on RNA replication. (A) Replication of given mutants in transfected Huh-7 cells as determined by luciferase reporter assays performed at 24, 48, and 72 h posttransfection (white, gray, and black bars, respectively). Values given were normalized for transfection efficiency using the luciferase activity determined 4 h after transfection, which was set to 1. Mean values of quadruplicate measurements and the standard deviations are given. (B) Phosphorylation analysis of NS5A using the vaccinia virus T7 hybrid system. NS3-to-NS5B polyprotein fragments carrying the mutations specified above the lanes were transfected into Huh-7 cells, and proteins were radiolabeled with [³²P]orthophosphate or [³⁵S]methionine and [³⁵S]cysteine. NS5A proteins were isolated by IP and separated by SDS-PAGE (10% polyacrylamide). Mock-transfected cells served as a negative control (lanes 9 and 10). Half of the samples were treated with λ protein phosphatase (λ-PPase) (+) whereas the other half was mock treated (-) prior to SDS-PAGE. Arrows and asterisks indicate hyperphosphorylated and basally phosphorylated forms, respectively.

with the core protein was observed in cells expressing the wild-type NS5A and the CL3A/SA-mutated NS5A, but the amount of immunoprecipitated NS5A in the CL3A/SA-expressing cells was slightly lower than that in the wild-type-expressing cells. In contrast, the CL3B/SA- or the Δ2384-2433-mutated NS5A coimmunoprecipitated with the core protein only slightly or not at all.

We further examined the interaction of NS5A with core protein in cells expressing HCV genomes. At 72 h posttransfection with the wild type or CL3B/SA, cells were harvested and immunoprecipitated with an anti-NS5A antibody or an anti-C/EBPβ antibody as a negative control, followed by immunoblotting. Under these experimental conditions, the amount of extracellular core protein released from cells transfected with the CL3B/SA genome was about 10-fold lower than

that for the wild type, although comparable amounts of intracellular core protein were observed in both transfectants (Fig. 4B, left panels). As shown in the right panels of Fig. 4B, the core protein was specifically coimmunoprecipitated with NS5A in cells expressing the wild-type JFH-1 genome but not with the mutated NS5A in cells expressing the CL3B/SA genome. These results demonstrate that NS5A interacts with the core protein in cells producing infectious particles and that serine residues at aa 2428, 2430, and 2433 are important to the success of this interaction.

Two serine residues among aa 2428, 2430, and 2433 are responsible for regulating the interaction of NS5A with the core protein as well as HCV particle production. To further determine the critical residues in the C-terminal serine cluster of NS5A responsible for HCV particle production, we replaced one or two serine residues in the region with alanine (Fig. 1) and investigated which serine-to-alanine substitution influenced HCV particle production. Core protein levels in cells transfected with any construct were comparable over 4 days after transfection, indicating similar efficiencies of transfection and RNA replication from each construct (data not shown). As shown in Fig. 5A, we observed a slight delay in the kinetics of core protein release from cells transfected with the single-substitution genomes, S2428A, S2430A, and S2433A, up to 48 or 72 h posttransfection. However, core protein release from these cells reached comparable levels to that for the wild type at 96 h after transfection. In the cases of the double-substitution mutants (Fig. 5B), core protein release from cells transfected with the double-substitution genomes was markedly reduced, with 10- to 30-fold decreases compared to that for wild type observed. The kinetics of core protein release were similar to that for CL3B/SA.

Interaction of NS5A carrying single or double serine-to-alanine substitutions with the core protein was investigated by coimmunoprecipitation analysis using HA-tagged NS5A constructs. NS5A mutants carrying a single substitution were coimmunoprecipitated with the core protein (Fig. 5C), while none of the double-substitution NS5A mutants or the triple-substitution mutant, CL3B/SA, coimmunoprecipitated with the core protein (Fig. 5D). These results suggest that at least two serine residues in the C-terminal serine cluster of NS5A (aa 2428, 2430, and 2433) are necessary for the interaction between NS5A and the core protein as well as for regulation of HCV particle production and that there is positive correlation between their interaction and the amount of core protein released.

Glutamic acid partially substitutes for serine phosphorylation in the interaction of NS5A with the core protein and virus production. A consequence of phosphorylation is the addition of negative charge to a protein. In some cases, phosphoserine can be mimicked by glutamic or aspartic acid (14). To determine whether the introduction of negative charges into aa 2428, 2430, and 2433 instead of phosphoserines positively regulates the interaction of NS5A with the core protein and virus production, we replaced the serine residues with glutamic acid residues and constructed the CL3B/SE and S2428/2430E mutants (Fig. 6A). Cells transfected with the double-glutamic acid substitution, S2428/2430E, exhibited similar kinetics to the wild-type-transfected cells and released ~22-fold more core protein than S2428/2430A-transfected cells by 96 h posttransfection (Fig. 6B). In contrast,




FIG. 3. Effect of mutations introduced into the NS5A C terminus on the production of infectious HCV particles. (A) Intracellular levels of core protein measured at various time points after transfection. A total of 3×10^6 Huh-7 cells were transfected with $10 \mu\text{g}$ of in vitro-transcribed HCV RNAs specified in the inset and resuspended in 10 ml of culture medium, after which 2-ml aliquots were seeded into each well of a six-well culture plate. The cells were harvested at different time points between 4 h and 96 h posttransfection, and then $500 \mu\text{l}$ of cell lysate per well was prepared. After centrifugation, supernatants were processed for a core protein-specific ELISA. (B) Release of core protein from cells transfected with the HCV genomes specified in the inset. Cell culture supernatants harvested from cells given in panel A were analyzed by a core protein ELISA. (C) Efficiency of core protein release from cells transfected with the HCV genomes specified in the inset. The percent core protein release (vertical axis) indicates the percentage of released core protein in relation to total core protein (the sum of intra- and extracellular core protein) calculated for each time point. (D) Infectivity of virus particles contained in supernatants and cells after transfection with mutants specified below the graph. Culture supernatants and cells were harvested 72 h posttransfection, and extracellular (white bars) and intracellular infectivity (gray bars) levels were determined by TCID_{50} assay. The gray line and arrowhead represent the detection limit of the limiting dilution assay. Mean values and standard deviations for at least triplicates are shown in all panels.

the transfectant with the triple glutamic acid substitution, CL3B/SE, showed similar trends to that of CL3B/SA. In the coimmunoprecipitation experiments with FLAG-tagged core protein and HA-tagged NS5A constructs (Fig. 6C), S2428/2430E, but not S2428/2430A, restored the ability of NS5A to interact with the core protein up to a similar level to that of wild type. As expected, neither CL3B/SE nor CL3B/SA coimmunoprecipitated with the core protein. Taken together, these results indicate that negative charges at aa 2428 and 2430 preserve the ability of NS5A to interact with the core protein and positively regulate virus production. However, the data of the CL3B/SE mutant indicate that it is likely that negative charges alone are not sufficient to enhance either the interaction of NS5A with the core protein or virus production.

Subcellular localization of NS5A and core protein in Huh-7 cells expressing HCV genomes.

The coimmunoprecipitation experiments described above indicate that the wild-type NS5A but not the CL3B/SA mutant interacts with the core protein. To evaluate the NS5A-core protein interaction in intact cells, we examined the subcellular localization of NS5A with the core protein by immunofluorescence analysis. NS5A colocalized with the core protein in cells transfected with the JFH-1 wild type (Fig. 7A), whereas their colocalization was rarely observed in cells transfected with the CL3B/SA RNA (Fig. 7B).

To further analyze the subcellular compartments for the localization of NS5A and core protein in cytoplasmic membrane structures, including the endoplasmic reticulum (ER) and LDs, we performed subcellular fractionation studies as




FIG. 4. aa 2428, 2430, and 2433 are essential for the interaction between NS5A and the core protein. (A) Effect of mutations at the NS5A C terminus on the interaction of NS5A with the core protein. N-terminally FLAG-tagged core protein and N-terminally HA-tagged NS5A carrying defined mutations were coexpressed in Huh-7 cells and immunoprecipitated with anti-FLAG antibody. The resulting precipitates were examined by immunoblotting using anti-HA or FLAG antibody. One-tenth of the cell lysates used in IP is shown as the 10% input. (B) Interaction between NS5A and the core protein in HCV-replicating cells. Huh-7 cells were lysed 72 h after transfection of the *in vitro* transcript of the HCV genome (wild type or CL3B/SA) and were immunoprecipitated with anti-NS5A antibody or anti-C/EBP β antibody as a negative control. The resulting precipitates were examined by immunoblotting using anti-core protein, NS5A, or C/EBP β antibody. One-tenth of cell lysates used in IP was immunoblotted with anti-core protein antibody (10% input). Cell culture supernatants harvested from transfected cells were analyzed by a core protein ELISA in parallel. IB, immunoblotting.

described in Materials and Methods. The iodixanol gradient was collected from the top to the bottom into 12 fractions (fractions 1 to 12). As shown in Fig. 7C, an ER marker, calnexin, was found in fractions 7 to 12 and was localized primarily in fractions 11 and 12. In contrast, ADRP, a cellular marker for LDs, was mainly observed in fractions 4 to 7. These two markers were equally distributed among cells analyzed (data not shown). The distribution of the wild-type NS5A was found in fractions 4 to 7, which was parallel to the fractionation profile of ADRP. The CL3B/SA-mutated NS5A was more broadly distributed and was also observed in heavier fractions than the wild-type NS5A, which was analogous to distribution of NS5A expressed in JFH1/4-1 cells bearing subgenomic replicons. The core protein in cells expressing the JFH-1 wild type, the CL3B/SA mutant, and in Huh/c-p7 cells that express JFH-1 structural proteins was distributed in a similar fashion, indicating that the distribution of core protein is not affected by NS5A mutation. The fractionation profile of the core protein, with a peak in fraction 4 or 5, was similar to that of the wild-type

NS5A or ADRP but not to that of the CL3B/SA-mutated NS5A or calnexin, suggesting that core protein interacts with the wild-type NS5A in LD fractions, which is consistent with previous reports (33, 44, 45).

NS5A-core protein interaction is important for association of the core protein with the viral genomic RNA. To further address our hypothesis regarding involvement of NS5A in recruiting viral RNA to nucleocapsid formation, we analyzed the association of the core protein with HCV RNA in wild-type- or CL3B/SA-expressing cells by IP-RT-PCR (Fig. 8). Both cell lysates were immunoprecipitated with an anti-core protein antibody or a negative control, mouse IgG. Total RNA prepared from each immunoprecipitate was subjected to RT-PCR in order to detect HCV RNA. The amounts of immunoprecipitated core protein (Fig. 8, lower panel) as well as the expression of HCV RNA (Fig. 8, upper panels, Input) were comparable in both cells. In cells expressing the wild-type JFH-1 genome, the viral RNAs covering the 5' terminal 2.2-kb as well as the 3' terminal 2.2-kb regions were detected in immunopre-




FIG. 5. Determination of critical amino acids responsible for virus production and the interaction of NSSA with the core protein. (A and B) Effect of single or double serine-to-alanine substitutions on virus production. After transfection of in vitro transcripts of the HCV genomes specified in the inset into Huh-7 cells, the cells and culture supernatants were harvested at the time points given, and the amounts of the core protein were determined by core protein-specific ELISA. Percent core protein release (vertical axis) indicates the percentage of released core protein in relation to total core protein (the sum of intra- and extracellular core protein) calculated for each time point. Mean values and standard deviations for at least triplicate experiments are shown. (C and D) Effect of single or double serine-to-alanine substitutions on the interaction between NSSA and the core protein. N-terminally FLAG-tagged core protein and N-terminally HA-tagged NSSA carrying defined mutations were coexpressed in Huh-7 cells and immunoprecipitated with anti-FLAG antibody. The resulting precipitates were examined by immunoblotting using anti-HA or FLAG antibody. One-tenth of the cell lysates used in IP is shown as the 10% input. IB, immunoblotting.

cipitates obtained with the anti-core protein antibody but not with the mouse IgG. In contrast, in cells expressing the CL3B/SA genome, HCV RNA was not detected in the immunoprecipitates with either antibody. These results demonstrate that HCV RNA associates with the core protein in cells where NSSA interacts with core protein (JFH-1 wild type) but not in cells where their interaction is impaired (CL3B/SA).

DISCUSSION

In the present study, we demonstrated the involvement of NSSA in the production of HCV particles via the interaction of NSSA with the core protein and identified its C-terminal serine cluster 3-B (aa 2428, 2430, and 2433), which is implicated in basal phosphorylation, as a key element for the interaction of NSSA with the core protein and for infectious virus production. Serine-to-alanine substitutions at the cluster, which have no impact on viral RNA replication, inhibit the interaction between NSSA and the core protein, thereby indicating that there is a connection between NSSA-core protein association and virus production. Finally, CL3B mutation leads to impair-

ment of the association of the core protein with HCV RNA and, therefore, possibly RNA encapsidation.

Several reports have indicated that viral NS proteins are involved in the virion assembly of *Flaviviridae* viruses (25, 29, 30, 33). For instance, mutations in yellow fever virus NS2A block production of infectious virus, and this perturbation can be released by a suppressor mutation in NS3 (25), while the hydrophobic residues of Kunjin virus NS2A required for virus assembly have been mapped (26). Miyanari et al. have shown that HCV core protein recruits NS proteins to the LD-associated membranes and that the NS proteins around the LDs participate in the assembly of infectious viral particles (33). Furthermore, during preparation of the current article, two studies regarding participation of NSSA in the assembly of HCV particles were published. Appel et al. have demonstrated the essential role of domain III of NSSA in the formation of infectious particles, and deletions in this domain that disrupt colocalization of NSSA and the core protein abrogate virion production (2). Tellinghuisen et al. identified a serine residue in domain III as a key determinant for viral particle production




FIG. 6. Effect of glutamic acid substitutions for phosphoserines at aa 2428, 2430, and 2433 on virus production and the interaction of NS5A with the core protein. (A) Alanine or glutamic acid substitutions for serine residues at aa 2428, 2430, and 2433. The numbers indicate amino acid positions within the polypeptide of the JFH-1 isolate. The names shown on the left represent full-length HCV or N-terminally HA-tagged NS5A constructs used in this experiment. Amino acid substitutions are marked in bold and underlined. C represents the C terminus. (B) Effect of alanine or glutamic acid substitutions on virus production. After transfection of *in vitro* transcripts of the HCV genomes specified in the inset into Huh-7 cells, the cells and the culture supernatants were harvested at the time points given, and the amounts of core protein were determined by core protein-specific ELISA. Percent core protein release (vertical axis) indicates the percentage of released core protein in relation to total core protein (the sum of intra- and extracellular core protein) calculated for each time point. Mean values and standard deviations for at least triplicate experiments are shown. (C) Effect of alanine or glutamic acid substitutions on the interaction between NS5A and the core protein. N-terminally FLAG-tagged core protein and N-terminally HA-tagged NS5A carrying defined mutations were coexpressed in Huh-7 cells and immunoprecipitated with anti-FLAG antibody. The resulting precipitates were examined by immunoblotting (IB) using anti-HA or FLAG antibody. One-tenth of the cell lysates used in IP is as shown as the 10% input.




FIG. 7. Subcellular localization of NSSA and the core protein in HCV-replicating cells. Huh-7 cells were transfected with the *in vitro* transcript of the HCV genome, wild type (A) or CL3B/SA (B). Seventy-two hours after transfection, the cells were fixed with 4% paraformaldehyde, permeabilized with 0.3% Triton X-100, and double stained with antibodies against the core protein (green) and NSSA (red), followed by staining with an Alexa Fluor 488- or Alexa Fluor 555-conjugated antibody. High-magnification panels are enlarged images of white squares in the merge panels. (C) HCV (wild type or CL3B/SA)-replicating cells, JFH1/4-1 cells harboring a subgenomic replicon of JFH-1, or Huh/c-p7 cells stably expressing JFH-1 structural proteins were lysed by freeze-thawing, and the cell lysates were fractionated on 5 to 25% iodixanol gradients. The distributions of NSSA, calnexin (ER marker), and ADRP (LD marker) were determined by immunoblotting, and those of the core protein were examined by core protein-specific ELISA.

(50). However, the mechanism by which NS proteins participate in virus assembly or the role of the interaction between structural and NS proteins in virus life cycles has not been fully elucidated. Here, we have clearly demonstrated that HCV NSSA interacts with the core protein in coimmunoprecipitation experiments not only with coexpression of each epitope-tagged protein but also with cells expressing the viral genome; and by using immunofluorescence and subcellular fractionation analysis, we have confirmed that mutations in CL3B abolish colocalization of NSSA and the core protein, presumably around LDs. In addition, the intracellular infectivity assay and IP-RT-PCR strongly suggest that impairment of the NSSA-core protein interaction results in disruption of virus production at an early stage of virion assembly. On the basis of the present results and findings in accompanying articles, one may infer the following events: newly synthesized HCV RNAs bound to NSSA are released from the replication complex-containing membrane compartment and can be captured by the core protein via interaction with domain III of NSSA at the surface of LDs or LD-associated membranes. Consequently, the viral RNAs are encapsidated, and virion assembly proceeds in the local environment. Recruitment of newly synthesized viral RNAs to the core protein could be important for efficient nucleocapsid formation in cells, where concentrations of the viral genome and the structural proteins are typically low, and may contribute to the selection of the viral genome to be

packaged. Interaction between NSSA and the core protein has been previously reported, and the NSSA region containing an interferon sensitivity determining region and the PKR-binding sequence (aa 2212 to 2330) has been mapped to that required for binding with core protein by yeast two-hybrid and *in vitro* pull-down assays (13). However, involvement of domain III in the NSSA-core protein interaction was not analyzed in detail, and a role for the NSSA-core protein interaction in the HCV life cycle was not examined in that study.

A growing body of evidence points to phosphorylation of NSSA as being important in controlling HCV RNA replication. Although the degree and the requirement for its hyperphosphorylation diverge between different HCV isolates, mutations that are associated with increased replicative fitness of HCV replicons frequently lead to a reduced level of NSSA hyperphosphorylation (1, 5, 36). Inhibitors of serine/threonine protein kinases that block NSSA hyperphosphorylation facilitate replication of a non-culture-adapted replicon (3, 36). One model that has been proposed suggests that NSSA hyperphosphorylation negatively regulates HCV RNA replication by disrupting the interaction between NSSA and the vesicle-associated membrane protein-associated protein subtype A, a cellular factor considered necessary for efficient RNA replication (5). However, the regulatory role of the basal phosphorylation of NSSA in the viral life cycle is poorly understood. It has been reported that the C-terminal region of NSSA (aa 2350 to 2419)




FIG. 8. IP-RT-PCR of HCV-replicating cells performed to examine the association between the core protein and the HCV genome RNA. Huh-7 cells were transfected with the *in vitro* transcript of the HCV genome (wild type or CL3B/SA) and lysed in 500 μ l of hypotonic buffer at 72 h posttransfection. After IP with an anti-core protein antibody or mouse IgG, immunoprecipitates were eluted in 100 μ l of elution buffer. RNAs in immunocomplexes were isolated by acid guanidinium thiocyanate-phenol-chloroform extraction. PCR was carried out as described in Materials and Methods with primer sets amplifying the fragments of nt 129 to 2367 and nt 7267 to 9463 of the JFH-1 genome. One-tenth (10 μ l) of each eluted immunoprecipitate was used for assays of the core protein amounts to ensure IP efficiency (lower panel). RNA extracted from a small aliquot of each cell lysate used in IP-RT-PCR is shown as the input.

is involved in basal phosphorylation (23). There are highly conserved serine residues in this region, and alanine substitutions or in-frame deletion of the serine residues has been shown to impair basal phosphorylation but not to affect RNA replication in the genotype 1b isolate (1). Consistently, a metabolic ³²P labeling experiment in the present study demonstrated that NS5A mutants of the JFH-1 isolate in the region impair the basal phosphorylation. Nevertheless, Tellinghuisen et al. noted that the serine at aa 2433 of JFH-1 is involved in generating hyperphosphorylated NS5A, as shown by Western blotting (50). The basis for this difference is uncertain. To date, there is no clear evidence to determine which serine residues located in domain III are phosphoacceptor sites or whether these residues influence NS5A phosphorylation in an indirect fashion. Future study to map phosphoacceptor sites in the NS5A domain III by biochemical approaches is needed.

We found that two of the three serine residues at CL3B are responsible for regulating the interaction of NS5A with the core protein as well as for infectious virus production. To further evaluate the effect of constitutive serine phosphorylation at the cluster, we replaced the serine residues with glu-

tamic acid, which mimics the presence of phosphoserines. The S2428/2430E mutant led to restoration of the interaction of NS5A with the core protein and virus production up to levels similar to the wild type. Somewhat unexpectedly, the triple glutamic acid substitution (CL3B/SE) exhibited only a slight restoration effect or none at all. It is considered that the degree of negative charge on the glutamic acid residue is not completely equivalent to that of phosphoserine. It is likely that the range of acidity at the local environment of the NS5A domain III that will allow interaction with the core protein is rather narrow. Induction of a conformational change in NS5A by the incorporation of phosphate may also be important for its interaction with the core protein. Tellinghuisen et al. reported that a single serine-to-alanine substitution at aa 2433 blocks the production of infectious virus and that casein kinase II likely phosphorylates the residue (50). Although this seems inconsistent with our results, these investigators also showed that deletions producing a lack of all three serine residues in the cluster inhibited virus production more severely than a single mutation. We observed that a single substitution of S2428A, S2430A, or S2433A resulted in a moderate decrease

in the virus released from the transfected cells; however, more evident perturbation was obtained from double or triple substitutions (Fig. 5A and B). Tellinghuisen et al. determined the HCV production at 48 h after RNA transfection and found a marked inhibition by the single substitution S2433A. In our study, as indicated in Fig. 5A, the reduction caused by the S2433A mutant was approximately 90% at 48 h after transfection; however, the virus production from the mutant reached a similar level to that of the wild type at 96 h posttransfection.

Several previous studies have found that apolipoproteins B (apoB) and E (apoE), microsomal triglyceride transfer protein, and HCV p7 protein are key factors for production of the infectious HCV particles (4, 11, 16, 22, 47). Assembly and maturation of the viral particles appear to depend on the formation of very-low-density lipoprotein, a large particle containing apoB, apoE, and large amounts of neutral lipids in hepatic cells. p7 protein is primarily involved in a late step of virus particle production, and the findings support the idea that p7 acts as viroporin, which has the capacity to compromise cell membrane integrity and thus favors the release of viral progeny. How the early step in virion production regulated by the NS5A-core protein interaction links with the later step(s) involved in the very-low-density lipoprotein assembly or p7 function remains an interesting question to be addressed.

In summary, we demonstrated that the C-terminal serine cluster of NS5A (aa 2428, 2430, and 2433), which is involved in generating the basal phosphorylated form, is a determinant of NS5A interaction with the core protein and the subcellular localization of NS5A. Mutation of this cluster blocks the NS5A-core protein interaction, resulting in perturbation of association between the core protein and HCV RNA. It is thus tempting to consider that NS5A plays a key role in transporting the viral genome RNA synthesized by the replication complex to the surface of LDs or LD-associated membranes, where the core protein localizes, leading to facilitation of nucleocapsid formation. Structural analysis of the NS5A domain III-core protein complex should provide greater insight into the mode of interaction between these viral proteins. Identification of residues at the interface that are involved in important interactions will be of significant value in designing novel structure-based inhibitors to block the early step of HCV particle formation.

ACKNOWLEDGMENTS

We are grateful to Francis V. Chisari (The Scripps Research Institute) for providing Huh-7 cells. We thank M. Matsuda, S. Yoshizaki, T. Shimoji, M. Kaga, and M. Sasaki for technical assistance and T. Mizoguchi for secretarial work.

This work was supported by Grants-in-Aid from the Ministry of Health, Labor and Welfare; by the Program for Promotion of Fundamental Studies in Health Sciences of the Organization for Drug ADR Relief, R&D Promotion and Product Review of Japan (grant ID:01-3); by the Japan Society for the Promotion of Science; and by Research on Health Sciences focusing on Drug Innovation from the Japan Health Sciences Foundation, Japan. T.M. is the recipient of a Research Resident Fellowship from the Foundation for Promotion of Cancer Research in Japan.

REFERENCES

- Appel, N., T. Pietschmann, and R. Bartenschlager. 2005. Mutational analysis of hepatitis C virus nonstructural protein 5A: potential role of differential phosphorylation in RNA replication and identification of a genetically flexible domain. *J. Virol.* 79:3187-3194.

- Appel, N., M. Zayas, S. Miller, J. Krijnsse-Locker, T. Schaller, P. Friebe, S. Kallis, U. Engel, and R. Bartenschlager. 2008. Essential role of domain III of nonstructural protein 5A for hepatitis C virus infectious particle assembly. *PLoS Pathog.* 4:e1000035.
- Bürckstämmer, T., M. Kriegs, J. Lupberger, E. K. Pauli, S. Schmitt, and E. Hildt. 2006. Raf-1 kinase associates with hepatitis C virus NS5A and regulates viral replication. *FEBS Lett.* 580:575-580.
- Chang, K. S., J. Jiang, Z. Cai, and G. Luo. 2007. Human apolipoprotein E is required for infectivity and production of hepatitis C virus in cell culture. *J. Virol.* 81:13783-13793.
- Evans, M. J., C. M. Rice, and S. P. Goff. 2004. Phosphorylation of hepatitis C virus nonstructural protein 5A modulates its protein interactions and viral RNA replication. *Proc. Natl. Acad. Sci. USA* 101:13038-13043.
- Gale, M., Jr., C. M. Blakely, B. Kwieciszewski, S. L. Tan, M. Dossett, N. M. Tang, M. J. Kurth, S. J. Polyak, D. R. Gretch, and M. G. Katze. 1998. Control of PKR protein kinase by hepatitis C virus nonstructural 5A protein: molecular mechanisms of kinase regulation. *Mol. Cell. Biol.* 18:5208-5218.
- Gale, M., Jr., and M. G. Katze. 1998. Molecular mechanisms of interferon resistance mediated by viral-directed inhibition of PKR, the interferon-induced protein kinase. *Pharmacol. Ther.* 78:29-46.
- Gale, M., Jr., B. Kwieciszewski, M. Dossett, H. Nakao, and M. G. Katze. 1999. Antiapoptotic and oncogenic potentials of hepatitis C virus are linked to interferon resistance by viral repression of the PKR protein kinase. *J. Virol.* 73:6506-6516.
- Gale, M., Jr., M. J. Kurth, and M. G. Katze. 1998. Repression of the PKR protein kinase by the hepatitis C virus NS5A protein: a potential mechanism of interferon resistance. *Clin. Diagn. Virol.* 10:157-162.
- Gale, M., Jr., M. J. Kurth, N. M. Tang, S. L. Tan, D. A. Hopkins, T. E. Dever, S. J. Polyak, D. R. Gretch, and M. G. Katze. 1997. Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein. *Virology* 230:217-227.
- Gastaminza, P., G. Cheng, S. Wieland, J. Zhong, W. Liao, and F. V. Chisari. 2008. Cellular determinants of hepatitis C virus assembly, maturation, degradation, and secretion. *J. Virol.* 82:2120-2129.
- Gastaminza, P., S. B. Kapadia, and F. V. Chisari. 2006. Differential biological properties of infectious intracellular and secreted hepatitis C virus particles. *J. Virol.* 80:11074-11081.
- Goh, P. Y., Y. J. Tan, S. P. Lim, S. G. Lim, Y. H. Tan, and W. J. Hong. 2001. The hepatitis C virus core protein interacts with NS5A and activates its caspase-mediated proteolytic cleavage. *Virology* 290:224-236.
- Hicke, L., B. Zanolari, and H. Riezman. 1998. Cytoplasmic tail phosphorylation of the alpha-factor receptor is required for its ubiquitination and internalization. *J. Cell Biol.* 141:349-358.
- Hoofnagle, J. H. 2002. Course and outcome of hepatitis C. *Hepatology* 36:S21-S29.
- Huang, H., F. Sun, D. M. Owen, W. Li, Y. Chen, M. Gale, Jr., and J. Ye. 2007. Hepatitis C virus production by human hepatocytes dependent on assembly and secretion of very low-density lipoproteins. *Proc. Natl. Acad. Sci. USA* 104:5848-5853.
- Huang, L., J. Hwang, S. D. Sharma, M. R. Hargittai, Y. Chen, J. J. Arnold, K. D. Raney, and C. E. Cameron. 2005. Hepatitis C virus nonstructural protein 5A (NS5A) is an RNA-binding protein. *J. Biol. Chem.* 280:36417-36428.
- Ishii, K., K. Murakami, S. S. Hmw, B. Zhang, J. Li, M. Shirakura, K. Morikawa, R. Suzuki, T. Miyamura, T. Wakita, and T. Suzuki. 2008. Trans-encapsulation of hepatitis C virus subgenomic replicon RNA with viral structure proteins. *Biochem. Biophys. Res. Commun.* 371:446-450.
- Ishii, K., Y. Ueda, K. Matsuo, Y. Matsuura, T. Kitamura, K. Kato, Y. Izumi, K. Someya, T. Ohsu, M. Honda, and T. Miyamura. 2002. Structural analysis of vaccinia virus DIs strain: application as a new replication-deficient viral vector. *Virology* 302:433-444.
- Jesch, S. A., A. J. Mehta, M. Velliste, R. F. Murphy, and A. D. Linstedt. 2001. Mitotic Golgi is in a dynamic equilibrium between clustered and free vesicles independent of the ER. *Traffic* 2:873-884.
- Johnson, R. F., S. E. McCarthy, P. J. Godlewski, and R. N. Hartley. 2006. Ebola virus VP30-VP40 interaction is sufficient for packaging 3E-SSE minigenome RNA into virus-like particles. *J. Virol.* 80:5135-5144.
- Jones, C. T., C. L. Murray, D. K. Eastman, J. Tassello, and C. M. Rice. 2007. Hepatitis C virus p7 and NS2 proteins are essential for production of infectious virus. *J. Virol.* 81:8374-8383.
- Kaneko, T., Y. Tanji, S. Satoh, M. Hijikata, S. Asabe, K. Kimura, and K. Shimotohno. 1994. Production of two phosphoproteins from the NS5A region of the hepatitis C viral genome. *Biochem. Biophys. Res. Commun.* 205:320-326.
- Kato, T., T. Date, M. Miyamoto, M. Sugiyama, Y. Tanaka, E. Orito, T. Ohno, K. Sugihara, I. Hasegawa, K. Fujiwara, K. Ito, A. Ozasa, M. Mizokami, and T. Wakita. 2005. Detection of anti-hepatitis C virus effects of interferon and ribavirin by a sensitive replicon system. *J. Clin. Microbiol.* 43:5679-5684.
- Kümmerer, B. M., and C. M. Rice. 2002. Mutations in the yellow fever virus nonstructural protein NS2A selectively block production of infectious particles. *J. Virol.* 76:4773-4784.

26. Leung, J. Y., G. P. Pijlman, N. Kondratieva, J. Hyde, J. M. Mackenzie, and A. A. Khromykh. 2008. Role of nonstructural protein NS2A in flavivirus assembly. *J. Virol.* **82**:4731–4741.
27. Liang, T. J., B. Rehmann, L. B. Seeff, and J. H. Hoofnagle. 2000. Pathogenesis, natural history, treatment, and prevention of hepatitis C. *Ann. Intern. Med.* **132**:296–305.
28. Lindenbach, B. D., M. J. Evans, A. J. Syder, B. Wolk, T. L. Tellinghuisen, C. C. Rice. 2005. Complete replication of hepatitis C virus in cell culture. *Science* **309**:623–626.
29. Liu, W. J., H. B. Chen, and A. A. Khromykh. 2003. Molecular and functional analyses of Kunjin virus infectious cDNA clones demonstrate the essential roles for NS2A in virus assembly and for a nonconservative residue in NS3 in RNA replication. *J. Virol.* **77**:7804–7813.
30. Liu, W. J., P. L. Sedlak, N. Kondratieva, and A. A. Khromykh. 2002. Complementation analysis of the flavivirus Kunjin NS3 and NS5 proteins defines the minimal regions essential for formation of a replication complex and shows a requirement of NS3 in *cis* for virus assembly. *J. Virol.* **76**:10766–10775.
31. Manns, M. P., H. Wedemeyer, and M. Cornberg. 2006. Treating viral hepatitis C: efficacy, side effects, and complications. *Gut* **55**:1350–1359.
32. Miyamoto, M., T. Kato, T. Date, M. Mizokami, and T. Wakita. 2006. Comparison between subgenomic replicons of hepatitis C virus genotypes 2a (JFH-1) and 1b (Con1 NK5.1). *Intervirology* **49**:37–43.
33. Miyazaki, Y., K. Atsuzawa, N. Usuda, K. Watanashi, T. Hishiki, M. Zayas, R. Bartschlagler, T. Wakita, M. Hijikata, and K. Shimotohno. 2007. The lipid droplet is an important organelle for hepatitis C virus production. *Nat. Cell Biol.* **9**:1089–1097.
34. Moradpour, D., M. J. Evans, R. Gosert, Z. Yuan, H. E. Blum, S. P. Goff, B. D. Lindenbach, and C. M. Rice. 2004. Insertion of green fluorescent protein into nonstructural protein 5A allows direct visualization of functional hepatitis C virus replication complexes. *J. Virol.* **78**:7400–7409.
35. Murakami, K., T. Kimura, M. Osaki, K. Ishii, T. Miyamura, T. Suzuki, T. Wakita, and I. Shoji. 2008. Virological characterization of the hepatitis C virus JFH-1 strain in lymphocytic cell lines. *J. Gen. Virol.* **89**:1587–1592.
36. National Institutes of Health. 2002. NIH consensus statement on management of hepatitis C. NIH Consens. State. Sci. Statements **19**:1–46.
37. Neddermann, P., M. Quintavalle, C. Di Pietro, A. Clementi, M. Cerretani, S. Altamura, L. Bartholomew, and R. De Francesco. 2004. Reduction of hepatitis C virus NS5A hyperphosphorylation by selective inhibition of cellular kinases activates viral RNA replication in cell culture. *J. Virol.* **78**:13306–13314.
38. Niwa, H., K. Yamamura, and J. Miyazaki. 1991. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* **108**:193–199.
39. Pawlotsky, J. M. 1999. Hepatitis C virus (HCV) NS5A protein: role in HCV replication and resistance to interferon- α . *J. Viral Hepat.* **6**(Suppl. 1): 47–48.
40. Pawlotsky, J. M., G. Germanidis, A. U. Neumann, M. Pellerin, P. O. Fraiua, and D. Dhumeaux. 1998. Interferon resistance of hepatitis C virus genotype 1b: relationship to nonstructural 5A gene quasispecies mutations. *J. Virol.* **72**:2795–2805.
41. Poyard, T., M. F. Yuen, V. Ratzu, and C. L. Lai. 2003. Viral hepatitis C. *Lancet* **362**:2095–2100.
42. Schaller, T., N. Appel, G. Koutsoudakis, S. Kallis, V. Lohmann, T. Pietschmann, and R. Bartschlagler. 2007. Analysis of hepatitis C virus superinfection exclusion by using novel fluorochrome gene-tagged viral genomes. *J. Virol.* **81**:4591–4603.
43. Seeff, L. B., and J. H. Hoofnagle. 2003. Appendix: The National Institutes of Health Consensus Development Conference: management of hepatitis C 2002. *Clin. Liver Dis.* **7**:261–287.
44. Seeff, L. B., and J. H. Hoofnagle. 2002. National Institutes of Health Consensus Development Conference: management of hepatitis C. 2002. *Hepatology* **36**:S1–S2.
45. Shavinskaya, A., S. Boulant, F. Penin, J. McLauchlan, and R. Bartschlagler. 2007. The lipid droplet binding domain of hepatitis C virus core protein is a major determinant for efficient virus assembly. *J. Biol. Chem.* **282**:37158–37169.
46. Shi, S. T., S. J. Polyak, H. Tu, D. R. Taylor, D. R. Gretch, and M. M. Lai. 2002. Hepatitis C virus NS5A colocalizes with the core protein on lipid droplets and interacts with apolipoproteins. *Virology* **292**:198–210.
47. Shirakura, M., K. Murakami, T. Ichimura, R. Suzuki, T. Shimoji, K. Fukuda, K. Abe, S. Sato, M. Fukasawa, Y. Yamakawa, M. Nishijima, K. Moriishi, Y. Matsuura, T. Wakita, T. Suzuki, P. M. Howley, T. Miyamura, and I. Shoji. 2007. E6AP ubiquitin ligase mediates ubiquitination and degradation of hepatitis C virus core protein. *J. Virol.* **81**:1174–1185.
48. Steinmann, E., F. Penin, S. Kallis, A. H. Patel, R. Bartschlagler, and T. Pietschmann. 2007. Hepatitis C virus p7 protein is crucial for assembly and release of infectious virions. *PLoS Pathog.* **3**:e103.
49. Tan, S. L., and M. G. Katze. 2001. How hepatitis C virus counteracts the interferon response: the jury is still out on NS5A. *Virology* **284**:1–12.
50. Tanji, Y., T. Kaneko, S. Sato, and K. Shimotohno. 1995. Phosphorylation of hepatitis C virus-encoded nonstructural protein NS5A. *J. Virol.* **69**:3980–3986.
51. Tellinghuisen, T. L., K. L. Foss, and J. Treadaway. 2008. Regulation of hepatitis C virus production via phosphorylation of the NS5A protein. *PLoS Pathog.* **4**:e1000032.
52. Tellinghuisen, T. L., K. L. Foss, J. C. Treadaway, and C. M. Rice. 2008. Identification of residues required for RNA replication in domains II and III of the hepatitis C virus NS5A protein. *J. Virol.* **82**:1073–1083.
53. Tellinghuisen, T. L., J. Marcotrigiano, A. E. Gorbalenya, and C. M. Rice. 2004. The NS5A protein of hepatitis C virus is a zinc metalloprotein. *J. Biol. Chem.* **279**:48576–48587.
54. Tellinghuisen, T. L., J. Marcotrigiano, and C. M. Rice. 2005. Structure of the zinc-binding domain of an essential component of the hepatitis C virus replicase. *Nature* **435**:374–379.
55. van den Hoff, M. J., A. F. Moorman, and W. H. Lamers. 1992. Electroporation in "intracellular" buffer increases cell survival. *Nucleic Acids Res.* **20**:2902.
56. van Regenmortel, M. H. V., C. M. Fauquet, D. H. L. Bishop, E. B. Carstens, M. K. Estes, S. M. Lemon, J. Maniloff, M. A. Mayo, D. J. McGeoch, C. R. Pringle, and R. B. Wickner (ed.). 2000. *Virus taxonomy: classification and nomenclature of viruses*. Seventh report of the International Committee on Taxonomy of Viruses. Academic Press, San Diego, CA.
57. Wakita, T., T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H. G. Krausslich, M. Mizokami, R. Bartschlagler, and T. J. Liang. 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* **11**:791–796.
58. Watanashi, K., M. Hijikata, A. Tagawa, T. Doi, H. Marusawa, and K. Shimotohno. 2003. Modulation of retinoid signaling by a cytoplasmic viral protein via sequestration of Sp110b, a potent transcriptional corepressor of retinoic acid receptor, from the nucleus. *Mol. Cell. Biol.* **23**:7498–7509.
59. Zhong, J., P. Gastaminza, G. Cheng, S. Kapadia, T. Kato, D. R. Burton, S. F. Wieland, S. L. Uprichard, T. Wakita, and F. V. Chisari. 2005. Robust hepatitis C virus infection in vitro. *Proc. Natl. Acad. Sci. USA* **102**:9294–9299.



Trans-encapsidation of hepatitis C virus subgenomic replicon RNA with viral structure proteins

Koji Ishii*, Kyoko Murakami, Su Su Hmwe, Bin Zhang, Jin Li, Masayuki Shirakura, Kenichi Morikawa, Ryoosuke Suzuki, Tatsuo Miyamura, Takaji Wakita, Tetsuro Suzuki

Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan

ARTICLE INFO

Article history:

Received 2 April 2008

Available online 28 April 2008

Keywords:

Hepatitis C virus

Replicon

Virus-like particle

Vaccine

ABSTRACT

A trans-packaging system for hepatitis C virus (HCV) subgenomic replicon RNAs was developed. HCV subgenomic replicon was efficiently encapsidated by the HCV structural proteins that were stably expressed *in trans* under the control of a mammalian promoter. Infectious HCV-like particles (HCV-LPs), established a single-round infection, were produced and released into culture medium in titers of up to 10^3 focus forming units/ml. Expression of NS2 protein with structural proteins (core, E1, E2, and p7) was shown to be critical for the infectivity of HCV-LPs. Anti-CD81 treatment decreased the number of infected cells, suggesting that HCV-LPs infected cells in a CD81-dependent manner. The packaging cell line should be useful both for the production of single-round infectious HCV-LPs to elucidate the mechanisms of HCV assembly, particle formation and infection to host cells, and for the development of HCV replicon-based vaccines.

© 2008 Elsevier Inc. All rights reserved.

Hepatitis C virus (HCV) is a positive-strand RNA virus that belongs to the *Hepacivirus* genus in the *Flaviviridae* family. The HCV genome comprises about 9600 nucleotides that encode a single polypeptide of around 3000 amino acids [1–3], which is processed by cellular and viral encoded proteases into at least 10 different structural and nonstructural proteins [4–6]. The JFH-1 strain of HCV, classified as genotype 2a strain, is the first HCV strain that can produce HCV particles in Huh7 cells [7,8]. The synthesis of HCV-like particles (HCV-LPs) using a recombinant baculovirus containing the cDNA of HCV structural proteins has been reported [9]. HCV-LP production by mammalian expression systems using vesicular stomatitis virus [10] and semliki forest virus [11] were also reported although the amount of VLP production is not as high as that of baculovirus system.

Subgenomic replicon system is a useful tool as gene expression vectors and is desirable for the development of vaccines. In the case of flaviviruses, several systems have been described for packaging flavivirus replicons, including Kunjin virus replicons [12–14], yellow fever virus replicons [15], tick-borne encephalitis virus replicons [16], and West Nile virus replicons [17,18]. In some cases, these packaging systems have utilized cell lines expressing the flavivirus structural proteins under the control of eukaryotic promoters [16,19]. These virus-like particle (VLP)-generating systems have been useful for packaging viral genomes encoding various for-

eign genes [14,15,20,18], the study of virus tropism and various aspects of viral assembly and entry [17].

Subgenomic replicons of JFH-1 replicate efficiently in Huh7 cells and do not require cell culture-adaptive mutations [21]. The construction of a system to package HCV replicon into HCV-LPs would not only be useful to investigate as-yet unclear steps of HCV life cycles such as genome packaging and virion assembly but also offers the possibilities of a new approach for vaccine development. In this study, we constructed subgenomic replicon cell lines constitutively expressing JFH-1 structural proteins under the control of elongation factor-1 α (EF) promoter, and found stable expression of structural proteins and release of HCV-LPs from the cell line. A sucrose density gradient centrifugation of the culture medium resulted in partial purification of the HCV-LPs. Infectivity of HCV-LPs produced by this system was confirmed by colony formation assay and immunofluorescence analysis. Anti-CD81 antibody treatment decreased the infectivity of HCV-LPs, suggesting that VLPs infected to cells in CD81-dependent fashion. This is the first report that HCV structural proteins of HCV can trans-package its subgenomic replicon. The system described here should be useful to elucidate the mechanisms of HCV assembly, particle formation, and infection to host cells.

Materials and methods

Plasmid construction. Core to p7 coding region of JFH-1 was amplified using pJFH-1 [21] as a template and sense primer

* Corresponding author. Fax: +81 3 5285 1161.

E-mail address: kishii@nih.go.jp (K. Ishii).