

ing proteases and polymerases, from patients persistently infected with RNA viruses that exhibit a quasispecies nature, such as human immunodeficiency virus and HCV. Viral quasispecies are not a simple collection of diverse mutants but a group of interactive variants capable of adapting to new environments (48). Cyclosporine treatment has been shown to be effective for patients infected with HCV of genotype 1b (20) and suppresses HCV RNA replication *in vitro* (52). In addition, cyclosporine has been shown to disrupt the interaction between NS5B and cyclophilin B, which is required for an efficient RNA-binding of NS5B (53). Cyclophilins and FKBP8 are classified as immunophilins capable of binding to the immunosuppressants cyclosporine and FK506, respectively (26). The family members do not share a homologous domain other than drug-binding and enzymatically active domains, based on their amino acid sequences, substrate specificities, and inhibitor sensitivities. However, cyclosporine-resistant RNA replicon was shown to exhibit mutations not only in NS5B but also in NS5A (12, 41), suggesting that cyclosporine might affect the viral replication through the nucleotide-binding ability of NS5B, as well as the function of NS5A. Recently, geldanamycin, an inhibitor of Hsp90, was shown to drastically impair the replication of poliovirus without any emergence of escape mutants (14). Therefore, the elucidation of host proteins, including immunophilins and chaperones, participating in the HCV replication complex may lead to the development of new therapeutics for chronic hepatitis C with a broad spectrum and a low possibility of emergence of revertant viruses. In particular, disruption of the specific interaction of Val¹²¹ of NS5A with the TPR domain of FKBP8 might be an ideal target for a novel therapeutic measure.

Egger et al. reported that NS4B alters the intracellular membrane to form a membranous web structure consisting of a membrane-associated multiprotein complex localized in the cytoplasmic compartments distinct from the mitochondria *in vitro* and in the liver of an HCV-infected chimpanzee, suggesting that the membranous web forms the viral replication complex (8). An N-terminal amphipathic helix of NS4B plays an important role in the viral replication, as well as in the correct localization of other NS proteins including NS5A (9). Furthermore, VAP-B was reported to interact with Nir2 protein through the FFAT (named for two phenylalanines [i.e., FF] in the acidic tract) motif and to remodel the ER structure to form a convoluted membrane structure resembling a membranous web (3). In addition, VAP-A and B interact with not only NS5A but also NS5B (13, 16, 47), suggesting that the complex of NS5A with FKBP8 might be recruited on the membranous web by NS4B and/or VAPs and participate in the HCV replication.

FKBP8 has been shown to be localized mainly on the mitochondria and to interact with Bcl-2 to sequester Bcl-2 on the mitochondria (7, 44). However, HCV RNA was suggested to be replicated in the membranous web structure in replicon cells (8, 13, 15), and NS5A was reported to localize on the ER, Golgi apparatus (2, 6, 16), and lipid droplets (43). Figures 6C and 7A clearly indicate that the intracellular compartment including NS5A and FKBP8 is distinct from mitochondria. The HCV core protein was shown to upregulate genes related to fatty acid biosynthesis through the interaction with proteasome activator PA28 γ /REG γ in the nucleus (34) and to induce ac-

cumulation of cytoplasmic lipid droplets in the mouse liver (35). Recently, it was shown that the HCV core protein of the genotype 2a JFH1 strain recruits the replication complex to the lipid droplet-associated membranes, and HCV particles were detected in close proximity to the lipid droplets, suggesting that lipid droplets induced by core protein participate in the assembly of HCV particles (31). In addition, the lipid droplets including the core protein were surrounded by the nonstructural proteins was also detected in cells expressing the chimeric HCV genomes encoding core to a part of NS2 proteins of genotype 1b or 1a strain and the nonstructural proteins of JFH1 strain (31). In the present study, FKBP8 was shown to be colocalized with NS5A in a highly electron-dense intracellular compartment indistinguishable from the membranous web. Although the total amount of FKBP8 was not changed by the treatment of the replicon cells by IFN- α (data not shown), the membranous web structure where FKBP8 and NS5A had accumulated was removed by the treatment (Fig. 7B). These results suggest that the replication of the subgenomic HCV RNA induces the formation of a membranous web structure in which NS5A and FKBP8 are colocalized but has no effect on the expression level of FKBP8. Furthermore, we could not detect any colocalization of FKBP8 and NS5A with the lipid droplets in the replicon cells harboring a full-length genome of the genotype 1b Con1 strain (data not shown). Although the relationships between the membranous web and lipid droplets remain unknown, these discrepancies might be attributable to the difference in HCV genotypes of the nonstructural proteins that consist of the major components of the replication complex determining the efficiency of HCV replication.

In conclusion, our data indicate that NS5A directly binds to FKBP8 through the Val¹²¹ and colocalizes in the convoluted membrane structure known as the membranous web. Future studies on the role of FKBP8 in the replication of HCV might contribute to the development of a new type of anti-HCV drugs with a low frequency of emergence of drug-resistant breakthrough viruses.

ACKNOWLEDGMENTS

We thank H. Murase for secretarial work. We also thank R. Bartschlagler and C. E. Cameron for providing the plasmids. We are grateful to the staff of the Center for Research and Education for their help in using the Biacore 2000.

This work was supported in part by grants-in-aid from the Ministry of Health, Labor, and Welfare; the Ministry of Education, Culture, Sports, Science, and Technology; the 21st Century Center of Excellence Program; and the Foundation for Biomedical Research and Innovation.

REFERENCES

- Abe, T., Y. Kaname, I. Hamamoto, Y. Tsuda, X. Wen, S. Tagawa, K. Morishiki, O. Takeuchi, T. Kawai, T. Kanto, N. Hayashi, S. Akira, and Y. Matsura. 2007. Hepatitis C virus nonstructural protein 5A modulates the Toll-like receptor-MyD88-dependent signaling pathway in macrophage cell lines. *J. Virol.* 81:8953–8966.
- Aizaki, H., K. S. Choi, M. Liu, Y. J. Li, and M. M. Lai. 2006. Polypyrimidine-tract-binding protein is a component of the HCV RNA replication complex and necessary for RNA synthesis. *J. Biomed. Sci.* 13:469–480.
- Amarillo, R., S. Ramachandran, H. Sabanay, and S. Lev. 2005. Differential regulation of endoplasmic reticulum structure through VAP-Nir protein interaction. *J. Biol. Chem.* 280:5934–5944.
- Appel, N., T. Pietschmann, and R. Bartschlagler. 2005. Mutational analysis of hepatitis C virus nonstructural protein 5A: potential role of differential phosphorylation in RNA replication and identification of a genetically flexible domain. *J. Virol.* 79:3187–3194.

5. Boulant, S., P. Targett-Adams, and J. McLauchlan. 2007. Disrupting the association of hepatitis C virus core protein with lipid droplets correlates with a loss in production of infectious virus. *J. Gen. Virol.* **88**:2204–2213.
6. Brass, V., E. Bieck, R. Montserret, B. Wolk, J. A. Hellings, H. E. Blum, F. Penin, and D. Moradpour. 2002. An amino-terminal amphipathic alpha-helix mediates membrane association of the hepatitis C virus nonstructural protein 5A. *J. Biol. Chem.* **277**:8130–8139.
7. Bulgakov, O. V., J. T. Eggenschwiler, D. H. Hong, K. V. Anderson, and T. Li. 2004. FKBP8 is a negative regulator of mouse sonic hedgehog signaling in neural tissues. *Development* **131**:2149–2159.
8. Egger, D., B. Wolk, R. Gosert, L. Bianchi, H. E. Blum, D. Moradpour, and K. Bienz. 2002. Expression of hepatitis C virus proteins induces distinct membrane alterations including a candidate viral replication complex. *J. Virol.* **76**:5974–5984.
9. Elazar, M., K. H. Cheong, P. Liu, H. B. Greenberg, C. M. Rice, and J. S. Glenn. 2003. Amphipathic helix-dependent localization of NS5A mediates hepatitis C virus RNA replication. *J. Virol.* **77**:6055–6061.
10. Enomoto, N., I. Sakuma, Y. Asahina, M. Kurosaki, T. Murakami, C. Yamamoto, N. Izumi, F. Marumo, and C. Sato. 1995. Comparison of full-length sequences of interferon-sensitive and resistant hepatitis C virus 1b. Sensitivity to interferon is conferred by amino acid substitutions in the NS5A region. *J. Clin. Invest.* **96**:224–230.
11. Evans, M. J., C. M. Rice, and S. P. Goff. 2004. Phosphorylation of hepatitis C virus nonstructural protein 5A modulates its protein interactions and viral RNA replication. *Proc. Natl. Acad. Sci. USA* **101**:13038–13043.
12. Fernandes, F., D. S. Poole, S. Hoover, R. Middleton, A. C. Andrei, J. Gerstner, and R. Striker. 2007. Sensitivity of hepatitis C virus to cyclosporine A depends on nonstructural proteins NS5A and NS5B. *Hepatology* **46**:1026–1033.
13. Gao, L., H. Aizaki, J. W. He, and M. M. Lai. 2004. Interactions between viral nonstructural proteins and host protein hVAP-33 mediate the formation of hepatitis C virus RNA replication complex on lipid raft. *J. Virol.* **78**:3480–3488.
14. Geller, R., M. Vignuzzi, R. Andino, and J. Frydman. 2007. Evolutionary constraints on chaperone-mediated folding provide an antiviral approach refractory to development of drug resistance. *Genes Dev.* **21**:195–205.
15. Gosert, R., D. Egger, V. Lohmann, R. Bartenschlager, H. E. Blum, K. Bienz, and D. Moradpour. 2003. Identification of the hepatitis C virus RNA replication complex in Huh-7 cells harboring subgenomic replicons. *J. Virol.* **77**:5487–5492.
16. Hamamoto, I., Y. Nishimura, T. Okamoto, H. Aizaki, M. Liu, Y. Mori, T. Abe, T. Suzuki, M. M. Lai, T. Miyamura, K. Moriishi, and Y. Matsuura. 2005. Human VAP-B is involved in hepatitis C virus replication through interaction with NS5A and NS5B. *J. Virol.* **79**:13473–13482.
17. Ho, S. N., H. D. Hunt, R. M. Horton, J. K. Pullen, and L. R. Pease. 1989. Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene* **77**:51–59.
18. Horton, R. M., H. D. Hunt, S. N. Ho, J. K. Pullen, and L. R. Pease. 1989. Engineering hybrid genes without the use of restriction enzymes: gene splicing by overlap extension. *Gene* **77**:61–68.
19. Huang, L., E. V. Sineva, M. R. Hargittai, S. D. Sharma, M. Suthar, K. D. Raney, and C. E. Cameron. 2004. Purification and characterization of hepatitis C virus nonstructural protein 5A expressed in *Escherichia coli*. *Protein Expr. Purif.* **37**:144–153.
20. Inoue, K., K. Sekiyama, M. Yamada, T. Watanabe, H. Yasuda, and M. Yoshida. 2003. Combined interferon alpha2b and cyclosporin A in the treatment of chronic hepatitis C: controlled trial. *J. Gastroenterol.* **38**:567–572.
21. Kanda, T., A. Basu, R. Steele, T. Wakita, J. S. Rysse, R. Ray, and R. B. Ray. 2006. Generation of infectious hepatitis C virus in immortalized human hepatocytes. *J. Virol.* **80**:4633–4639.
22. Kapadia, S. B., and F. V. Chisari. 2005. Hepatitis C virus RNA replication is regulated by host geranylgeranylation and fatty acids. *Proc. Natl. Acad. Sci. USA* **102**:2561–2566.
23. Krieger, N., V. Lohmann, and R. Bartenschlager. 2001. Enhancement of hepatitis C virus RNA replication by cell culture-adaptive mutations. *J. Virol.* **75**:4614–4624.
24. Lamarre, D., P. C. Anderson, M. Bailey, P. Beaulieu, G. Bolger, P. Bonneau, M. Bos, D. R. Cameron, M. Cartier, M. G. Cordingley, A. M. Faucher, N. Goudreau, S. H. Kawai, G. Kukoj, L. Lagace, S. R. LaPlante, H. Narjes, M. A. Poupard, J. Rancourt, R. A. Sentjens, R. St. George, B. Simoneau, G. Steinmann, D. Thibault, Y. S. Tsantrizos, S. M. Weldon, C. L. Yong, and M. Llinas-Brunet. 2003. An NS3 protease inhibitor with antiviral effects in humans infected with hepatitis C virus. *Nature* **426**:186–189.
25. Lindenbach, B. D., M. J. Evans, A. J. Syder, B. Wolk, T. L. Tellinghuisen, C. C. Liu, T. Maruyama, R. O. Hynes, D. R. Burton, J. A. McKeating, and C. M. Rice. 2005. Complete replication of hepatitis C virus in cell culture. *Science* **309**:623–626.
26. Liu, J., J. D. Farmer, Jr., W. S. Lane, J. Friedman, I. Weissman, and S. L. Schreiber. 1991. Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell* **66**:807–815.
27. Lohmann, V., F. Korner, J. Koch, U. Herian, L. Theilmann, and R. Bartenschlager. 1999. Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science* **285**:110–113.
28. Macdonald, A., K. Crowder, A. Street, C. McCormick, and M. Harris. 2004. The hepatitis C virus NS5A protein binds to members of the Src family of tyrosine kinases and regulates kinase activity. *J. Gen. Virol.* **85**:721–729.
29. Macdonald, A., and M. Harris. 2004. Hepatitis C virus NS5A: tales of a promiscuous protein. *J. Gen. Virol.* **85**:2485–2502.
30. Manns, M. P., J. G. McHutchison, S. C. Gordon, V. K. Rustgi, M. Shiffman, R. Reindollar, Z. D. Goodman, K. Koury, M. Ling, and J. K. Albrecht. 2001. Peginterferon alpha-2b plus ribavirin compared with interferon alpha-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* **358**:958–965.
31. Miyanari, Y., K. Atsuzawa, N. Usuda, K. Watashi, T. Hishiki, M. Zayas, R. Bartenschlager, T. Wakita, M. Hijikata, and K. Shimotohno. 2007. The lipid droplet is an important organelle for hepatitis C virus production. *Nat. Cell Biol.* **9**:1089–1097.
32. Moradpour, D., F. Penin, and C. M. Rice. 2007. Replication of hepatitis C virus. *Nat. Rev. Microbiol.* **5**:453–463.
33. Moriishi, K., and Y. Matsuura. 2003. Mechanisms of hepatitis C virus infection. *Antivir. Chem. Chemother.* **14**:285–297.
34. Moriishi, K., R. Mochizuki, K. Moriya, H. Miyamoto, Y. Mori, T. Abe, S. Murata, K. Tanaka, T. Miyamura, T. Suzuki, K. Koike, and Y. Matsuura. 2007. Critical role of PA28gamma in hepatitis C virus-associated steatogenesis and hepatocarcinogenesis. *Proc. Natl. Acad. Sci. USA* **104**:1661–1666.
35. Moriya, K., H. Yotsuyanagi, Y. Shintani, H. Fujie, K. Ishibashi, Y. Matsuura, T. Miyamura, and K. Koike. 1997. Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J. Gen. Virol.* **78**(Pt. 7):1527–1531.
36. Niwa, H., K. Yamamura, and J. Miyazaki. 1991. Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* **108**:193–199.
37. Okamoto, T., Y. Nishimura, T. Ichimura, K. Suzuki, T. Miyamura, T. Suzuki, K. Moriishi, and Y. Matsuura. 2006. Hepatitis C virus RNA replication is regulated by FKBP8 and Hsp90. *EMBO J.* **25**:5015–5025.
38. Pawlotsky, J. M., and G. Germain. 1999. The nonstructural 5A protein of hepatitis C virus. *J. Viral Hepat.* **6**:343–356.
39. Randall, G., M. Panis, J. D. Cooper, T. L. Tellinghuisen, K. E. Sukhodolets, S. Pfeffer, M. Landthaler, P. Landgraf, S. Kan, B. D. Lindenbach, M. Chien, D. B. Weir, J. J. Russo, J. Ju, M. J. Brownstein, R. Sheridan, C. Sander, M. Zavolan, T. Tuschl, and C. M. Rice. 2007. Cellular cofactors affecting hepatitis C virus infection and replication. *Proc. Natl. Acad. Sci. USA* **104**:12884–12889.
40. Rieder, C. L., and S. S. Bowser. 1985. Correlative immunofluorescence and electron microscopy on the same section of Epon-embedded material. *J. Histochem. Cytochem.* **33**:165–171.
41. Robida, J. M., H. B. Nelson, Z. Liu, and H. Tang. 2007. Characterization of hepatitis C virus subgenomic replicon resistance to cyclosporine in vitro. *J. Virol.* **81**:5829–5840.
42. Sakamoto, H., K. Okamoto, M. Aoki, H. Kato, A. Katsume, A. Ohta, T. Tsukuda, N. Shimizu, Y. Aoki, M. Arisawa, M. Kohara, and M. Sudo. 2005. Host sphingolipid biosynthesis as a target for hepatitis C virus therapy. *Nat. Chem. Biol.* **1**:333–337.
43. Shi, S. T., S. J. Polyak, H. Tu, D. R. Taylor, D. R. Gretch, and M. M. Lai. 2002. Hepatitis C virus NS5A localizes with the core protein on lipid droplets and interacts with apolipoproteins. *Virology* **292**:198–210.
44. Shirane, M., and K. I. Nakayama. 2003. Inherent calcineurin inhibitor FKBP38 targets Bel-2 to mitochondria and inhibits apoptosis. *Nat. Cell Biol.* **5**:28–37.
45. Tellinghuisen, T. L., J. Marcotrigiano, A. E. Gorbalenya, and C. M. Rice. 2004. The NS5A protein of hepatitis C virus is a zinc metalloprotein. *J. Biol. Chem.* **279**:48576–48587.
46. Tellinghuisen, T. L., J. Marcotrigiano, and C. M. Rice. 2005. Structure of the zinc-binding domain of an essential component of the hepatitis C virus replicase. *Nature* **435**:374–379.
47. Tu, H., L. Gao, S. T. Shi, D. R. Taylor, T. Yang, A. K. Mircheff, Y. Wen, A. E. Gorbalenya, S. B. Hwang, and M. M. Lai. 1999. Hepatitis C virus RNA polymerase and NS5A complex with a SNARE-like protein. *Virology* **263**:30–41.
48. Vignuzzi, M., J. K. Stone, J. J. Arnold, C. E. Cameron, and R. Andino. 2006. Quasispecies diversity determines pathogenesis through cooperative interactions in a viral population. *Nature* **439**:344–348.
49. Wakita, T., T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H. G. Krausslich, M. Mizokami, R. Bartenschlager, and T. J. Liang. 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* **11**:791–796.
50. Wang, C., M. Gale, Jr., B. C. Keller, H. Huang, M. S. Brown, J. L. Goldstein, and J. Ye. 2005. Identification of FBL2 as a geranylgeranylated cellular protein required for hepatitis C virus RNA replication. *Mol. Cell* **18**:425–434.
51. Wasley, A., and M. J. Alter. 2000. Epidemiology of hepatitis C: geographic differences and temporal trends. *Semin. Liver Dis.* **20**:1–16.
52. Watashi, K., M. Hijikata, M. Hosaka, M. Yamaji, and K. Shimotohno. 2003.

- Cyclosporin A suppresses replication of hepatitis C virus genome in cultured hepatocytes. *Hepatology* **38**:1282-1288.
53. **Watashi, K., N. Ishii, M. Hijikata, D. Inoue, T. Murata, Y. Miyanari, and K. Shimotohno.** 2005. Cyclophilin B is a functional regulator of hepatitis C virus RNA polymerase. *Mol. Cell* **19**:111-122.
54. **Ye, J., C. Wang, R. Sumpter, Jr., M. S. Brown, J. L. Goldstein, and M. Gale, Jr.** 2003. Disruption of hepatitis C virus RNA replication through inhibition of host protein geranylgeranylation. *Proc. Natl. Acad. Sci. USA* **100**:15865-15870.
55. **Yi, M., and S. M. Lemon.** 2004. Adaptive mutations producing efficient replication of genotype 1a hepatitis C virus RNA in normal Huh7 cells. *J. Virol.* **78**:7904-7915.
56. **Yi, M., R. A. Villanueva, D. L. Thomas, T. Wakita, and S. M. Lemon.** 2006. Production of infectious genotype 1a hepatitis C virus (Hutchinson strain) in cultured human hepatoma cells. *Proc. Natl. Acad. Sci. USA* **103**:2310-2315.
57. **Zhong, J., P. Gastaminza, G. Cheng, S. Kapadia, T. Kato, D. R. Burton, S. F. Wieland, S. L. Uprichard, T. Wakita, and F. V. Chisari.** 2005. Robust hepatitis C virus infection in vitro. *Proc. Natl. Acad. Sci. USA* **102**:9294-9299.

Human Butyrate-Induced Transcript 1 Interacts with Hepatitis C Virus NS5A and Regulates Viral Replication[†]

Shuhei Tagawa,¹ Toru Okamoto,¹ Takayuki Abe,¹ Yoshio Mori,¹ Tetsuro Suzuki,²
Kohji Moriishi,¹ and Yoshiharu Matsuura^{1*}

Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka University, Osaka,¹ and
Department of Virology II, National Institute of Infectious Diseases, Tokyo,² Japan

Received 2 October 2007/Accepted 18 December 2007

Hepatitis C virus (HCV) nonstructural protein 5A (NS5A) is required for the replication of the viral genome and is involved in several host signaling pathways. To gain further insight into the functional role of NS5A in HCV replication, we screened human cDNA libraries by a yeast two-hybrid system using NS5A as the bait and identified human butyrate-induced transcript 1 (hB-ind1) as a novel NS5A-binding protein. Endogenously and exogenously expressed hB-ind1 was coimmunoprecipitated with NS5A of various genotypes through the coiled-coil domain of hB-ind1. The small interfering RNA (siRNA)-mediated knockdown of hB-ind1 in human hepatoma cell lines suppressed the replication of HCV RNA replicons and the production of infectious particles of HCV genotype 2a strain JFH1. Furthermore, these reductions were canceled by the expression of an siRNA-resistant hB-ind1 mutant. Among the NS5A-binding host proteins involved in HCV replication, hB-ind1 exhibited binding with FKBP8, and hB-ind1 interacted with Hsp90 through the FxxW motif in its N-terminal p23 homology domain. The impairment of the replication of HCV RNA replicons and of the production of infectious particles of JFH1 virus in the hB-ind1 knockdown cell lines was not reversed by the expression of an siRNA-resistant hB-ind1 mutant in which the FxxW motif was replaced by AxxA. These results suggest that hB-ind1 plays a crucial role in HCV RNA replication and the propagation of JFH1 virus through interaction with viral and host proteins.

Hepatitis C virus (HCV) infects approximately 170 million people worldwide and induces serious chronic hepatitis that results in steatosis, cirrhosis, and ultimately hepatocellular carcinoma (7, 64). More than two-thirds of the HCV-positive population in Western countries and Japan face chronic infection by genotypes 1a and 1b. The current combination therapy using pegylated alpha interferon (IFN) plus ribavirin has achieved a sustained virological response in 50% of individuals infected with HCV genotypes 1a and 1b (37, 53).

HCV belongs to the genus *Hepacivirus* of the family *Flaviviridae* and has a single-stranded, positive-sense RNA genome of approximately 9.6 kb, encoding a large polyprotein composed of approximately 3,000 amino acid residues. The polyprotein is cleaved by host and viral proteases, resulting in viral structural proteins (core, E1, and E2), a putative ion channel-forming protein (p7), and nonstructural proteins (NS2, NS3, NS4A, NS4B, NS5A, and NS5B) (40, 55). Highly structured untranslated regions are flanked at both the 5' and 3' ends of the open reading frame. The initiation of translation of the viral RNA is dependent on an internal ribosome entry site (IRES) localized in the 5' untranslated region (28, 58).

The HCV RNA is suggested to replicate in a replication complex composed of the viral nonstructural proteins and several host proteins. An HCV replicon system established as a representative functional system was composed of an antibiotic gene for selection and HCV genomic RNA for autonomous

replication in the intracellular compartments of human hepatoma cell line Huh7 without production of infectious particles (34). Recently, cell culture systems for production of an infectious HCV have been established based on HCV genotype 2a (32, 62, 74). Furthermore, a mouse model consisting of an immunodeficient mouse xenotransplanted with human liver fragments has been established for the study of in vivo replication of HCV (38). These in vitro and in vivo systems have enabled us to investigate the life cycle of HCV and to develop antiviral drugs for chronic hepatitis C.

NS5A is a phosphoprotein that possesses multiple functions in viral replication, IFN resistance, and pathogenesis (35). Adaptive mutations to increase RNA replication are frequently mapped to the coding region of NS5A, indicating that NS5A is critical for HCV replication (1, 71). NS5A has been shown to be associated with a range of cellular proteins involved in cellular signaling pathways, such as IFN-induced kinase PKR (14), growth factor receptor-binding protein 2 (Grb2) (56), p53 (36, 48), and the phosphoinositide-3-kinase p85 subunit (18), and with proteins involved in protein trafficking and membrane morphology, such as karyopherin b3 (8), apolipoprotein A1 (52), amphiphysin II (73), F-box and leucine-rich repeat protein 2 (FBL2) (26, 63, 70), and vesicle-associated membrane protein-associated protein subtype A (VAP-A) (59). We have previously reported that the host proteins VAP-B and FKBP8, a member of the FK506-binding protein (FKBP) family, interact with NS5A and that these interactions are required for efficient replication of HCV (16, 45), further supporting the hypothesis that NS5A is a pivotal component of the HCV replication complex.

To gain a better understanding of the functional role of

* Corresponding author. Mailing address: Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka University, 3-1 Yamada-oka, Suita, Osaka 565-0871, Japan. Phone: 81-6-6879-8340. Fax: 81-6-6879-8269. E-mail: matsuura@biken.osaka-u.ac.jp.

[†] Published ahead of print on 26 December 2007.

NS5A in HCV replication, we screened human libraries by employing a yeast two-hybrid system and using NS5A as the bait. We thus identified human butyrate-induced transcript 1 (hB-ind1) as an NS5A-binding protein. Murine B-ind1 has been identified as a transcript induced by treatment with sodium butyrate in BALB/c BP-A31 mouse fibroblasts (10). hB-ind1 is a multiple-membrane-spanning protein, consisting of 362 amino acids, that possesses significant homology with protein tyrosine phosphatase-like, member A (PTPLA), and co-chaperone p23 and is suggested to be involved in the Rac1 signaling pathway (10). In this study we examine the biological effects of the interaction of hB-ind1 with NS5A and other host proteins on the replication of HCV.

MATERIALS AND METHODS

Plasmids. The plasmids encoding NS5A, FKBP8, VAP-A, VAP-B, and heat shock protein 90 (Hsp90) have been described previously (45). The human FBL2 gene was amplified from the total cDNA of Huh7 by PCR. A cDNA clone containing hB-ind1 cDNA was isolated from a human fetal brain library (Clontech, Palo Alto, CA) by the advanced yeast two-hybrid system Matchmaker Two-Hybrid System 3 (Clontech) using an HCV NS5A protein as bait. Each cDNA of N-terminally FLAG-tagged hB-ind1 and its mutants was generated by cloning into pEF FlagGs pGKuro (23). pSilencer-hB-ind1, carrying a short hairpin RNA (shRNA) targeted to hB-ind1 under the control of the U6 promoter, was constructed by cloning of the oligonucleotide pair 5'-GATCCGGA AAAGCGACCACTGTTTCTCAAGAGAAAACAGTGGTCCGTTTTCCTTT TTTGGAAA-3'-5'-AGCTTTTCCAAAAGGAAAAGCGACCACTGTTT TCTCTGAGAAACAGTGGTCCGTTTTCGCG-3' between the BamHI and HindIII sites of pSilencer 2.1-U6 hygro (Ambion, Austin, TX). A plasmid encoding a mutant hB-ind1 resistant to shRNA was prepared by introduction of five silent mutations (nucleotides were changed from A to G, G to A, A to C, A to T, and C to T at positions 291, 294, 297, 300, and 301, respectively) into hB-ind1 cDNA by the method of splicing by overlap extension (19). The pSilencer negative-control plasmid (Ambion) has no homology to any human gene. The pFK-1₃₈₉ neo/NS3-3'/NK5.1 plasmid (46) was kindly provided by R. Barten Schlager, and the neomycin-resistant gene was replaced with a firefly luciferase gene. The resulting plasmid was designated pFK-1₃₈₉ FL/NS3-3'/NK5.1. The plasmids used in this study were confirmed by sequencing with ABI Prism genetic analyzer 3130 (Applied Biosystems, Tokyo, Japan).

Cells and virus infection. All cell lines were cultured at 37°C under a humidified atmosphere with 5% CO₂. Human embryo kidney 293T cells were maintained in Dulbecco's modified Eagle's medium (DMEM) (Sigma, St. Louis, MO) supplemented with 100 U/ml penicillin, 100 µg/ml streptomycin, and 10% fetal calf serum (FCS). The human hepatoma cell line Huh7.5.1 was kindly provided by F. Chisari (74). The Huh7 and Huh7.5.1 cell lines were maintained in DMEM containing nonessential amino acids (NEAA), 100 U/ml penicillin, 100 µg/ml streptomycin, and 10% FCS. The Huh9-13 cell line, an Huh7-derived cell line harboring a subgenomic HCV replicon (34), was maintained in DMEM containing 10% FCS, NEAA, and 1 mg/ml G418 (Nacalai Tesque, Kyoto, Japan). Huh7.5.1 cells were transfected with pSilencer-hB-ind1 or an empty plasmid, and drug-resistant clones were selected by treatment with hygromycin (Wako, Tokyo, Japan) at a final concentration of 10 µg/ml. Plasmids encoding a full-length or truncated (amino acid residues 101 to 277) version of hB-ind1 were transfected into Huh7.5.1 cells, and the cells surviving after selection with 0.1 µg/ml of puromycin for 1 week were used for virus infection. The viral RNA of JFH1 was introduced into Huh7.5.1 cells according to the method of Wakita et al. (62). The supernatant was collected at 7 days posttransfection and used as HCV particles that are infectious in cell culture (HCVcc).

Antibodies. A rabbit anti-hB-ind1 antibody was prepared by immunization with synthetic peptides corresponding to amino acid residues 106 to 117 of hB-ind1. A mouse monoclonal antibody to influenza virus hemagglutinin (HA) was purchased from Covance (Richmond, CA). The mouse anti-FLAG M2 antibody that was conjugated with a horseradish peroxidase and a mouse anti-β-actin monoclonal antibody were purchased from Sigma. The mouse monoclonal antibody to HCV NS5A was obtained from Austral Biologicals (San Ramon, CA).

Yeast two-hybrid assay and library screening. A human fetal brain library prepared with pAct2 was purchased from Clontech and was screened by the yeast two-hybrid system Matchmaker GAL4 Two-Hybrid System 3 (Clontech) accord-

ing to the manufacturer's protocol. The NS5A cDNA fragment encoding amino acid residues 1973 to 2419 of HCV strain Con1 was amplified by PCR and cloned into pGBKT7 (Clontech); the resulting plasmid was designated pGBKT7 HCV NS5A. The yeast *Saccharomyces cerevisiae* strain AH109, which secretes α-galactosidase under the control of the MEL1 region, was transformed with pGBKT7 HCV NS5A and grown on a medium lacking tryptophan. The clone including the bait plasmid was transformed with the library plasmids. The transformed yeast cells were grown on 2% agar plates of a dropout medium lacking tryptophan, leucine, histidine, and adenine. The resulting colonies grown on the dropout plate were inoculated again on a new dropout plate containing 20 µg/ml X-α-Gal (5-bromo-4-chloro-3-indolyl-α-D-galactopyranoside) and incubated at 30°C for 7 days. The total DNA was prepared from all blue colonies and then introduced into *Escherichia coli* strain JM109. The prey plasmids were recovered from the clones grown on LB agar plates containing 10 µg/ml ampicillin. One positive clone was isolated from among 2 million colonies of the human fetal brain library, and the nucleotide sequence of this clone includes the complete cDNA of hB-ind1 in its frame.

Transfection, immunoblotting, and immunoprecipitation. Transfection and immunoprecipitation analyses were carried out as described previously (16, 45). Immunoprecipitates boiled in loading buffer were subjected to 12.5% sodium dodecyl sulfate-polyacrylamide gel electrophoresis. The proteins were transferred to polyvinylidene difluoride membranes (Millipore, Bedford, MA) and were reacted with the appropriate antibodies. The immune complexes were visualized with SuperSignal West Femto substrate (Pierce, Rockford, IL) and detected by an LAS-3000 image analyzer system (Fujifilm, Tokyo, Japan).

Gene silencing by siRNA. The short interfering RNAs (siRNAs) Target-4 (5'-GCUGAGUGACGUACAGAAC-3') and Target-6 (5'-GGAAAAGCGAC CACUGUUU-3') were obtained for knockdown of endogenous hB-ind1 (Ambion, Austin, TX). The negative control, siCONTROL Non-Targeting siRNA 2, which exhibits no downregulation of any human genes, was purchased from Dharmacon (Buckinghamshire, United Kingdom). Huh9-13 cells harboring a subgenomic HCV replicon grown on 6-well plates were transfected with 20 nM siRNA by using siFECTOR (B-Bridge International, Sunnyvale, CA) according to the manufacturer's protocol. The transfected cells were incubated in DMEM supplemented with 10% FCS and were then harvested at 96 h posttransfection.

Real-time PCR. The HCV RNA level was estimated by the method described previously (16, 45). Total RNA was prepared from cells by using the RNeasy minikit (Qiagen, Tokyo, Japan). First-strand cDNA was synthesized using an RNA LA PCR kit (Takara Bio Inc., Shiga, Japan) and random primers. Each cDNA was estimated by Platinum Sybr green qPCR SuperMix UDG (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol. Fluorescent signals were analyzed by an ABI Prism 7000 system (Applied Biosystems). The HCV IRES, glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and hB-ind1 genes were amplified using primer pairs 5'-GAGTGTGGTCCAGCCGCA-3'-5'-CACTGCGAAGCACCCATATCA-3', 5'-GAAGGTGAAGGTCCGAGATC-3'-5'-GAAGGTGAAGGTCCGAGATC-3', and 5'-CACCTGGAGTCTTAGA CCTTGTG-3'-5'-CAGTCCGAGTTTATTAGCGCTC-3', respectively. The values for HCV genomic RNA and hB-ind1 mRNA were normalized to that for GAPDH mRNA. Each PCR product was detected as a single band of the correct size by agarose gel electrophoresis (data not shown).

In vitro transcription and RNA transfection. Plasmids pFK-1₃₈₉ neo/NS3-3'/NK5.1 and pFK-1₃₈₉ FL/NS3-3'/NK5.1 were linearized at the ScaI site and then transcribed in vitro using the MEGAScript T7 kit (Ambion) according to the manufacturer's protocol. To generate capped mRNA encoding *Renilla* luciferase, pRL-CMV was cleaved with BamHI and then transcribed using the mMESSAGE mMACHINE kit (Ambion) according to the manufacturer's protocol. These in vitro-transcribed RNAs were introduced into Huh7.5.1 cells at 4 million cells/0.4 ml by electroporation at 270 V and 960 µF using Gene Pulser (Bio-Rad, Hercules, CA).

Colony formation assay. The colony formation assay has been described previously (45). Briefly, in vitro-transcribed RNA was electroporated into Huh7 cells and plated in DMEM containing 10% FCS and NEAA. The medium was replaced with fresh DMEM containing 10% FCS, NEAA, and 1 mg/ml G418 at 24 h posttransfection. The remaining colonies were fixed with 4% paraformaldehyde and stained with crystal violet at 4 weeks after electroporation.

Luciferase assay. Transfected cells were seeded in a 12-well plate and then lysed in 200 µl of passive lysis buffer (Promega, Madison, WI) at 24 h posttransfection. Luciferase activity was measured in 20-µl aliquots of cell lysates using the Dual-Luciferase reporter assay system (Promega). Firefly luciferase activity was standardized to that of *Renilla* luciferase, and the results are expressed as the increases in relative luciferase units (RLU).




FIG. 1. Interaction of NS5A with hB-ind1 in mammalian cells. (A) HA-NS5A of strain Con1 and FLAG-tagged hB-ind1 were expressed in 293T cells and immunoprecipitated (IP) with an anti-HA or anti-FLAG antibody. Immunoprecipitates were subjected to Western blotting (IB) to detect coprecipitated counterparts. As a negative control, an empty plasmid was used instead of the plasmid encoding FLAG-hB-ind1 or HA-NS5A. Anti-FLAG and anti-HA antibodies did not recognize HA-tagged NS5A and FLAG-tagged hB-ind1, respectively. (B) HA-NS5A protein derived from genotype 1b strain Con1 or J1, genotype 1a strain H77C, or genotype 2a strain JFH1 was coexpressed with FLAG-hB-ind1 in 293T cells, immunoprecipitated with an isotype control or anti-FLAG antibody, and analyzed by Western blotting with an antibody to the FLAG or HA tag. An empty plasmid was used instead of the plasmid encoding FLAG-hB-ind1 as a negative control. (C) Endogenous hB-ind1 in Huh9-13 cells harboring subgenomic HCV replicon RNA was immunoprecipitated with normal rabbit immunoglobulin G (IgG) (lane 1) or anti-hB-ind1 rabbit IgG (lane 2), and immunoprecipitates were analyzed by Western blotting with specific antibodies.

Statistical analysis. Results are expressed as means \pm standard deviations. The significance of differences between the means was determined by Student's *t* test.

RESULTS

hB-ind1 interacts with HCV NS5A of various genotypes. NS5A derived from the genotype 1b strain Con1 was used as bait to screen the human fetal brain cDNA library by a yeast two-hybrid system, and one clone including a gene encoding the open reading frame of the hB-ind1 gene was isolated. To examine whether hB-ind1 could interact with NS5A in mammalian cells, HA-tagged NS5A (HA-NS5A) was coexpressed with FLAG-tagged hB-ind1 (FLAG-hB-ind1) in 293T cells and immunoprecipitated with an antibody to the HA or the FLAG tag. FLAG-hB-ind1 and HA-NS5A were coimmunoprecipitated by either antibody (Fig. 1A). To determine the interaction of various genotypes of NS5A with hB-ind1, HA-NS5A of the genotype 1a strain H77C, the genotype 1b strain J1, or the genotype 2a strain JFH1 was coexpressed with FLAG-hB-ind1 and immunoprecipitated with the anti-FLAG antibody. An empty plasmid was used as a negative control. FLAG-hB-ind1 was immunoprecipitated with the anti-FLAG antibody at similar levels in cells coexpressing FLAG-hB-ind1 and HA-NS5A of all genotypes. HA-NS5A of various genotypes was coprecipitated with FLAG-hB-ind1 by the anti-FLAG antibody, whereas the anti-FLAG antibody did not precipitate any HA-NS5A of the various genotypes used in this study (Fig. 1B). To further confirm the interaction between hB-ind1 and HCV NS5A in the functional setting, lysates of Huh9-13 cells harboring subgenomic HCV replicon RNA were subjected to im-

muno-precipitation analysis with a rabbit polyclonal antibody raised against hB-ind1. NS5A was coimmunoprecipitated with endogenous hB-ind1 in the lysates of replicon cells (Fig. 1C). These results indicate that hB-ind1 interacts with NS5A of various HCV genotypes in mammalian cells.

hB-ind1 interacts with NS5A through the amino acid residues from 114 to 134 including the coiled-coil domain. hB-ind1 is composed of 362 amino acid residues and has domains homologous with p23 and PTPLA in the regions from Pro⁸ to Asp¹¹² and from Gln¹⁹⁶ to Leu³⁴⁶, respectively (Fig. 2A). To determine the region responsible for the interaction with NS5A, various deletion mutants of FLAG-hB-ind1 were constructed (Fig. 2B). Each of the mutants was coexpressed with Con1 HA-NS5A in 293T cells and immunoprecipitated with an anti-HA antibody. An empty plasmid was used as a negative control in the immunoprecipitation analyses. HA-NS5A was coimmunoprecipitated with full-length hB-ind1 and with mutants possessing amino acid residues 114 to 134, corresponding to the coiled-coil domain, which generally participates in protein-protein interactions (Fig. 2B and C), whereas HA-NS5A was not coimmunoprecipitated with hB-ind1 mutants lacking the coiled-coil domain. The anti-HA antibody did not coprecipitate FLAG-hB-ind1 or its mutants. These results indicate that hB-ind1 interacts with HCV NS5A through the coiled-coil domain.

hB-ind1 participates in the replication of HCV RNA and the propagation of infectious HCV particles. To investigate the role(s) of endogenous hB-ind1 in the replication of HCV RNA, an siRNA targeted to hB-ind1 or a control siRNA was transfected into Huh9-13 cells harboring subgenomic HCV




FIG. 2. Determination of the NS5A-binding region in hB-ind1. (A) Structure and functional domains of hB-ind1. (B) Deletion mutants of hB-ind1 used in this study and the results of binding to NS5A. N-terminally FLAG-tagged hB-ind1 mutants encoding the region from residue 1 to 149, 1 to 100, 101 to 362, 150 to 362, or 101 to 277 were designated 1-149, 1-100, 101-362, 150-362, or 101-277, respectively. An N-terminally FLAG-tagged hB-ind1 mutant spanning the region from residue 101 to residue 277 but lacking residues 114 to 134 was designated 101-277 Δ 114-134. In addition, N-terminally FLAG-tagged hB-ind1 mutants lacking the region from 101 to 149 or from 114 to 134 were designated Δ 101-149 or Δ 114-134, respectively. The coiled-coil domain was located at residues 114 to 134. Each mutant gene was inserted into pEF FLAGGs pGKpuro. A summary of immunoprecipitation results is given on the right. (C) Each hB-ind1 mutant was coexpressed with Con1 HA-NS5A in 293T cells, immunoprecipitated with an anti-HA antibody, and analyzed by Western blotting with an anti-FLAG antibody. As a negative control, an empty plasmid was used instead of the plasmid encoding HA-NS5A. The anti-HA antibody did not recognize FLAG-tagged hB-ind1 or its mutants.

replicon RNA. Total RNA was extracted from the transfected cells, and levels of hB-ind1 mRNA and HCV RNA were determined by real-time PCR. At 72 h posttransfection, hB-ind1 mRNA and HCV subgenomic RNA levels in cells transfected with each of the hB-ind1 siRNAs were reduced more than 60% from the levels in cells treated with the control siRNA (Fig. 3A). The levels of expression of hB-ind1 and the HCV NS5A protein were decreased in HCV replicon cells transfected with the hB-ind1 siRNA but not in those transfected with the control siRNA (Fig. 3B).

To examine the effects of the knockdown of hB-ind1 on the replication of HCV RNA and the propagation of HCVcc, we established Huh7.5.1 cell lines stably expressing an shRNA targeted to hB-ind1. Dozens of colonies were obtained from cells transfected with a plasmid encoding the cDNA of the shRNA to hB-ind1 after selection with hygromycin. Although the levels of mRNA and expression of endogenous hB-ind1 were not changed in cells bearing a nonspecific shRNA, they were reduced in the clones bearing shRNAs targeted to hB-ind1, except for clone 1 (Fig. 3C and D). There was no significant difference in growth among the cell lines (Fig. 3E).

The replicon RNA transcribed from pFK-1₃₈₉ neo/NS3-3/

NK5.1 was transfected into the hB-ind1 knockdown cell lines Huh-si2 and Huh-si5, which were cultured for 4 weeks in the presence of G418. The numbers of colonies in the knockdown cell lines were less than one-fourth of those in the control cell line (Huh-c) (Fig. 4A). A FLAG-tagged hB-ind1 wobble mutant (FLAG-rB-ind1), which is resistant to the shRNA targeted to hB-ind1 due to the introduction of silent mutations, was capable of expressing an siRNA-resistant hB-ind1 upon introduction into cells at a level similar to that of the endogenous hB-ind1 (eB-ind1) detected in the control cell line (Fig. 4B). The reduction of colony formation by the knockdown of eB-ind1 in the hB-ind1 knockdown cell lines Huh-si2 and Huh-si5 was canceled by the expression of FLAG-rB-ind1 (Fig. 4A). To further examine the involvement of hB-ind1 in the replication of HCV, a chimeric HCV RNA encoding a firefly luciferase gene under the control of HCV IRES (Fig. 4C) was transfected into the knockdown cell lines. Knockdown of hB-ind1 reduced the RLU in Huh-si2 and Huh-si5 cells by 40% and 70%, respectively, and this reduction was also canceled by the expression of FLAG-rB-ind1. To further examine the effect of hB-ind1 knockdown on the production of HCV infectious particles, HCVcc were inoculated into the hB-ind1 knockdown




FIG. 3. Effects of hB-ind1 knockdown on HCV replication. (A) Huh9-13 cells were transfected with siRNA 4 or siRNA 6 (#4 or #6, respectively), targeted to the hB-ind1 gene, or with a nonspecific siRNA, at a final concentration of 20 nM, and were harvested at 72 h posttransfection. hB-ind1 mRNA and HCV RNA levels were determined by real-time PCR. The levels of hB-ind1 mRNA and HCV RNA were normalized to the amount of GAPDH mRNA and expressed as percentages of the control value. (B) Huh9-13 cells transfected with siRNAs were lysed at 72 h posttransfection and subjected to Western blotting (IB) with an antibody to hB-ind1, NS5A, or β -actin. (C) Establishment of hB-ind1 knockdown Huh7.5.1 cell lines. Plasmids encoding shRNAs targeted to hB-ind1 (siRNA 6) or nonspecific targets were transfected into Huh7.5.1 cells and cultivated in the presence of hygromycin. Independent clones were established by limiting dilution. The value for hB-ind1 mRNA was normalized to the amount of GAPDH mRNA and expressed as a percentage of the control value. Huh7.5.1 cell lines expressing siRNAs targeted to hB-ind1 (Huh-si1 to Huh-si5) and to a nonspecific target (Huh-c) were established. (D) Expression of hB-ind1 in knockdown cells. The knockdown cell lines were lysed and subjected to Western blotting with an antibody to hB-ind1 or β -actin. (E) Growth curves of the knockdown cell lines were determined by the method of trypan blue dye exclusion. Data in this figure are representative of three independent experiments. Error bars, standard deviations. Asterisks indicate significant differences ($P < 0.01$) from the control value.

cell lines. Both virus titers, determined by focus-forming units at 72 h postinfection in culture supernatants, and HCV RNA levels in Huh-si2 and Huh-si5 cells were significantly reduced, and these reductions were canceled by the expression of FLAG-rB-ind1 (Fig. 4D). These results suggest that hB-ind1 is involved in the replication of HCV RNA and the propagation of HCVcc.

An hB-ind1 mutant retaining the binding region to NS5A has a dominant-negative effect on the replication of HCV. To examine the involvement of hB-ind1 in the replication of HCV in greater detail, deletion mutants of hB-ind1 retaining or lacking the binding region to NS5A were expressed in Huh9-13 cells harboring subgenomic HCV replicon RNA (Fig. 5A). Although the hB-ind1 mutant possessing the NS5A binding region (101-277) and full-length hB-ind1 were detected at similar levels in replicon cells transfected with the expression plasmids (Fig. 5B), HCV RNA replication was reduced only in cells expressing the mutant retaining the binding region to NS5A, not in those expressing full-length hB-ind1 or the mutant lacking the binding region to NS5A (101-277 Δ 114-134) (Fig. 5C). However, no significant difference in NS5A expres-

sion was observed in Huh9-13 cells transfected with the expression plasmids (Fig. 5B). Production of the infectious HCV particles was also reduced in the culture supernatants of Huh7.5.1 cells expressing the hB-ind1 mutant retaining the binding region to NS5A (101-277) but not in those expressing full-length hB-ind1 or the hB-ind1 101-277 Δ 114-134 mutant (Fig. 5D). These dominant-negative effects of the hB-ind1 mutant retaining the binding region to NS5A on the replication of HCV RNA in Huh9-13 cells and on the production of infectious particles in Huh7.5.1 cells further support the notion that hB-ind1 regulates the replication of HCV RNA and the propagation of HCVcc.

hB-ind1 interacts with FKBP8 and Hsp90. Previous reports have suggested that HCV NS5A interacts with several host proteins such as FBL2 (63), VAP-A (59), VAP-B (16), and FKBP8 (45) and that these interactions participate in the replication of HCV. To determine the interplay of the NS5A-binding proteins, FLAG-tagged hB-ind1 was coexpressed with HA-tagged FBL2, VAP-A, VAP-B, or FKBP8 in 293T cells and immunoprecipitated with an anti-FLAG antibody, and FKBP8 was shown to specifically interact with hB-ind1 (Fig. 6A). We have




FIG. 4. Effects of hB-ind1 knockdown on the replication of HCV RNA and the production of infectious particles. (A) The hB-ind1 knockdown (Huh-si2 and Huh-si5) and control (Huh-c) cell lines were first transfected with either a plasmid encoding hB-ind1 resistant to siRNA by virtue of the introduction of silent mutations (FLAG-rB-ind1) or an empty vector (EV) and then further transfected with replicon RNA transcribed from pFK-1₃₈₉ neo/NS3-3'/NK5.1. (Upper panel) The cell colonies remaining after cultivation for 4 weeks in the presence of G418 were fixed with 4% paraformaldehyde and stained with crystal violet. (Lower panel) The number of colonies was standardized to the amount of transfected RNA. (B) The expression of the siRNA-resistant hB-ind1 (FLAG-rB-ind1) and the endogenous hB-ind1 (eB-ind1) in Huh-c, Huh-si2, and Huh-si5 cells transfected with either a plasmid encoding FLAG-rB-ind1 or an empty vector was analyzed by Western blotting (IB) with an antibody to hB-ind1 or β -actin. (C) HCV subgenomic replicon RNA transcribed from pFK-1₃₈₉ FL/NS3-3'/NK5.1 and capped *Renilla* luciferase RNA transcribed from pRL-CMV were cotransfected into Huh-c, Huh-si2, and Huh-si5 cells pretreated with either a plasmid encoding FLAG-rB-ind1 or an empty vector. The firefly luciferase activity was normalized to that of *Renilla* luciferase. HCV IRES-dependent translational activity was expressed as a percentage of the RLU of Huh-c cells transfected with an empty plasmid. EMCV, encephalomyocarditis virus. (D) HCVcc were inoculated into Huh-c, Huh-si2, and Huh-si5 cells pretreated with either a plasmid encoding FLAG-rB-ind1 or an empty vector. (Upper panel) The culture supernatants at 72 h postinoculation were subjected to a focus-forming assay, and virus titers are expressed as focus-forming units (FFU) per milliliter. (Lower panel) The amount of intracellular HCV RNA was measured by real-time PCR and normalized to the amount of GAPDH mRNA. The HCV RNA level is expressed as a percentage of that of Huh-c cells transfected with an empty plasmid. Data in this figure are representative of three independent experiments. Error bars, standard deviations. Asterisks indicate significant differences (**, $P < 0.01$; *, $P < 0.05$) from the control value.

previously shown that FKBP8 is capable of binding to both NS5A and Hsp90 through the tetratricopeptide repeat (TPR) domain and that the recruitment of Hsp90 to the replication complex plays a crucial role in the replication of HCV (45). Hsp90 is a molecular chaperone and requires various cochaperone proteins such as p23 for efficient chaperone activity. hB-ind1 shows homology to p23 (Fig. 2A), and the FxxW motif, essential for the binding to Hsp90, is conserved in residues Phe¹⁰⁷xTrp¹¹⁰ of hB-ind1 (11, 27, 68). To determine whether hB-ind1 interacts with Hsp90 through the FxxW motif as reported for p23, FLAG-tagged hB-ind1 or an hB-ind1 mutant in which Phe¹⁰⁷ and Trp¹¹⁰ had been replaced with Ala (FLAG-hB-ind1AxxA) was coexpressed with HA-tagged Hsp90 in 293T cells and immunoprecipitated with an anti-FLAG antibody. Hsp90 was coimmunoprecipitated with wild-type hB-ind1 but not with the

mutant hB-ind1, indicating that hB-ind1 interacts with Hsp90 through the FxxW motif (Fig. 6B).

Previously, we showed that the amino acid residues of the carboxylate clump position in the TPR domain of FKBP8 attach to the C-terminal MEEVD motif of Hsp90 (45). To examine the interaction of hB-ind1 with Hsp90 in the absence of association with FKBP8, FLAG-tagged hB-ind1 was first coexpressed with HA-tagged Hsp90 or mutant Hsp90 lacking the MEEVD motif in 293T cells and then immunoprecipitated with an anti-FLAG antibody. Similar levels of hB-ind1 were coprecipitated with Hsp90 irrespective of the deletion of the MEEVD motif of Hsp90 (Fig. 6C), suggesting that hB-ind1 alone is capable of binding to Hsp90 through the FxxW motif irrespective of the association of FKBP8. To further clarify the interplay among hB-ind1, FKBP8, and Hsp90, FLAG-tagged




FIG. 5. Dominant-negative effect of an hB-ind1 mutant on the replication of HCV. (A) Plasmids encoding full-length hB-ind1 (construct 1) or deletion mutants of hB-ind1 retaining (construct 2) or lacking (construct 3) the NS5A binding region. (B) One of the three plasmids or an empty vector (EV) was transfected into Huh9-13 cells harboring a subgenomic HCV replicon RNA and was subjected to Western blotting (IB) with specific antibodies at 72 h posttransfection. (C) The amount of intracellular HCV RNA in the Huh9-13 cells was measured at 72 h posttransfection by real-time PCR, normalized to the amount of GAPDH mRNA, and expressed as the percentage of the value for control cells transfected with an empty plasmid. (D) One of the three plasmids or an empty vector was transfected into Huh7.5.1 cells, and then HCVcc were inoculated. Virus production in the culture supernatants at 72 h postinoculation was determined by a focus-forming assay. FFU, focus-forming units. Data in this figure are representative of three independent experiments. Error bars, standard deviations. Asterisks indicate significant differences ($P < 0.01$) from the control value.

hB-ind1 was coexpressed with HA-tagged Hsp90 and/or FKBP8 and then immunoprecipitated with an anti-FLAG antibody. Coprecipitation of Hsp90 with hB-ind1 was increased by additional expression of FKBP8 (Fig. 6D). These results suggest that hB-ind1 interacts with Hsp90 through the FxxW motif and that FKBP8 also participates in the complex formation to enhance the interaction.

hB-ind1 participates in HCV propagation through the interaction with Hsp90. Next, to examine the role of the interaction of hB-ind1 with Hsp90 in the replication of HCV RNA, the replicon RNA transcribed from pFK-1₃₈₉ neo/NS3-3'/NK5.1 was transfected into hB-ind1 knockdown Huh-si5 cells expressing siRNA-resistant FLAG-rB-ind1 or FLAG-rB-ind1AxxA, in which the Hsp90 binding motif FxxW was changed to AxxA. The colony formation in Huh-si5 cells transfected with an empty plasmid was 10% of that in Huh-c cells. The expression of FLAG-rB-ind1 in Huh-si5 cells recovered the colony formation in Huh-si5 cells to 98% of that in Huh-c cells, although that of FLAG-rB-ind1 AxxA in Huh-si5 cells exhibited only 40% recovery (Fig. 7A). To further examine the role of the interaction between hB-ind1 and Hsp90 in the production of HCVcc, Huh-si5 cells expressing either FLAG-rB-ind1 or FLAG-rB-ind1AxxA were infected with HCVcc, and the virus titer in the culture supernatants and the intracellular HCV RNA level at 72 h postinfection were determined. Virus production was reduced in the culture supernatants, and viral RNA replication in the hB-ind1 knockdown cells was restored by the expression of FLAG-rB-ind1 but not by that of FLAG-rB-ind1AxxA, as seen in colony formation by the replicon

RNA (Fig. 7B). Collectively, these results suggest that the interaction of hB-ind1 with Hsp90 through the FxxW motif is required for genomic RNA replication and particle production of HCV.

DISCUSSION

In this study we have shown that hB-ind1 participates in HCV RNA replication and particle production through interaction with NS5A, FKBP8, and Hsp90. hB-ind1 was initially identified as a downstream transducer of Rac1, a member of the small GTP-binding proteins, in mouse fibroblasts treated with sodium butyrate, a multifunctional agent known to inhibit cell proliferation and to induce differentiation by modulating transcription (6, 10). Rac1 possesses diverse biological functions, including cytoskeletal dynamics, membrane ruffling, cell cycle progression, gene transcription, and cell survival (4, 31, 49). Previous studies have suggested that hB-ind1 mediates Rac1 and Jun N-terminal protein kinase-NF- κ B signaling and is involved in the regulation of gene expression (6, 10). Inhibition of Rac1 function leads to disruption of cytoskeleton dynamics, resulting in impairment of cell growth (17, 69).

Inhibition of cell growth downregulates HCV RNA replication in the replicon cell line (41, 51), and cell cycle regulation affects HCV IRES-mediated translation (20, 61). Furthermore, cytoskeletal regulation is required for HCV RNA synthesis (3). However, knockdown of hB-ind1 and expression of the deletion mutants exhibited neither morphological change nor suppression of cell growth, suggesting that the suppression




FIG. 6. Interaction of hB-ind1 with other NS5A-binding host proteins. (A) FLAG-hB-ind1 was first coexpressed with HA-tagged FBL2, VAP-A, VAP-B, or FKBP8 in 293T cells and then immunoprecipitated with an anti-FLAG or control antibody. The immunoprecipitates were detected by Western blotting (IB) with an anti-HA antibody. (B) FLAG-hB-ind1 or FLAG-hB-ind1AxxA, in which Phe¹⁰⁷ and Trp¹¹⁰ had been replaced with Ala, was coexpressed with HA-Hsp90 in 293T cells and immunoprecipitated with an anti-FLAG antibody. The immunoprecipitates were detected by Western blotting with an anti-HA or anti-FLAG antibody. (C) FLAG-hB-ind1 was coexpressed with HA-Hsp90 or mutant Hsp90 lacking the MEEVD motif (HA-Hsp90 Δ MEEVD) in 293T cells and was immunoprecipitated with an anti-FLAG antibody. The immunoprecipitates were detected by Western blotting with an anti-HA or anti-FLAG antibody. (D) HA-Hsp90, HA-FKBP8, and FLAG-hB-ind1 were coexpressed in various combinations in 293T cells and immunoprecipitated with an anti-FLAG antibody. The immunoprecipitates were detected by Western blotting with an anti-HA or anti-FLAG antibody. Data in this figure are representative of three independent experiments.

of HCV replication by dysfunction of hB-ind1 is not due to cell growth arrest or cytoskeletal disruption. Murine B-ind1 has been reported to be expressed in all mouse tissues examined, with abundant expression detected in the testis, kidney, brain, and liver (10). Significant levels of endogenous hB-ind1 expression have been detected in the human hepatic cell lines Huh7, HepG2, Hep3B, and FLC4 and in the nonhepatic human cell lines HeLa, 293T, and THP-1 (data not shown); therefore, the tissue specificity of HCV replication could not be explained by the expression of hB-ind1.

Combination therapy with IFN and cyclosporine A has been shown to be effective for patients infected with a high viral load of HCV genotype 1b (24), and cyclosporine A has been shown to suppress HCV RNA replication in vitro through deactivation of the interaction between NS5B and cyclophilin B (66). Cyclophilin and FKBP are classified as immunophilins capable of binding to immunosuppressants cyclosporine A and FK506, respectively (33). The immunophilins do not share a homologous domain with each other, based on their amino acid sequences, substrate specificities, and inhibitor sensitivities. We

have recently reported that NS5A binds specifically to FKBP8 but not to other homologous immunophilins such as FKBP52 and cyclophilin D. FKBP8 forms both a homomultimer and a heteromultimer with the chaperone protein Hsp90. Mutation analyses of FKBP8 and Hsp90 suggest that FKBP8 acts as an intermediate between NS5A and Hsp90 via the different position of the TPR domain in FKBP8 and regulates HCV genome replication (45).

The molecular chaperone Hsp90 is one of the most abundant proteins in unstressed cells and generally requires various cochaperone proteins in multiple steps to promote the folding, functional maturation, and stability of its client proteins. Newly synthesized unfolded client proteins are delivered to the Hsp70 complex via Hsp40. In most cases, Hsp70 is able to process the client proteins on its own. Certain substrates require Hsp90 for proper folding or activation. In this case, the scaffold protein Hop connects elements of the Hsp70 and Hsp90 machineries to form an intermediate complex (2, 12, 13, 47). In the late stage, the Hsp70 component dissociates, and at the same time, p23 and immunophilins enter the complex (44, 54) and the




FIG. 7. Role of the interaction of hB-ind1 with Hsp90 in the replication of HCV. (A) hB-ind1 knockdown (Huh-si5) and control (Huh-c) cell lines were transfected either with a plasmid encoding the FLAG-tagged siRNA-resistant hB-ind1 (FLAG-rB-ind1) or FLAG-rB-ind1AxxA (with substitutions in the motif required for binding to Hsp90) or with an empty vector (EV) and were then further transfected with replicon RNA transcribed from pFK-1389 neo/NS3-3'/NK5.1. (Upper panel) The cell colonies remaining after cultivation for 4 weeks in the presence of G418 were fixed with 4% paraformaldehyde and stained with crystal violet. (Lower panel) The number of colonies was standardized to the amount of transfected RNA. (B) Upper panel) Huh-si5 cells expressing either FLAG-rB-ind1 or FLAG-rB-ind1AxxA were infected with HCVcc, and virus production in the culture supernatants at 72 h postinoculation was determined by a focus-forming assay. (Lower panel) The amount of intracellular HCV RNA was measured at 72 h posttransfection by real-time PCR, normalized to the amount of GAPDH mRNA, and expressed as a percentage of the value for control cells transfected with an empty plasmid. Data in this figure are representative of three independent experiments. Error bars, standard deviations. Asterisks indicate significant differences ($P < 0.01$) from the control value.

client proteins are refolded by Hsp90 chaperone activity to achieve the mature form. After that, p23 enhances the dissociation of the mature client protein from the final complex, and the released Hsp90 enters in the next chaperone cycle (72). It has been reported that Hsp90 cochaperone frequencies differ among client proteins (50). FKBP8 interacts with the C-terminal MEEVD motif of Hsp90 through the carboxylate clump position in the TPR domain of FKBP8 (45).

The C-terminal region of hB-ind1 shares homology with PIPLA (60). Protein tyrosine phosphatases are generally involved in the signaling pathways regulating metabolism, cell growth, differentiation, and cytoskeletal dynamics through the conserved HC(x)₅R motif (57). NS5A also interacts with signal transducer and activator of transcription 1 (STAT1) and impairs IFN signaling through the suppression of STAT1 phosphorylation (30). In addition, intracellular uptake of apoptotic cells expressing NS5A by dendritic cells leads to an increase in the secretion of CXCL-8 and impairment of IFN-induced tyrosine phosphorylation of STAT1 and STAT2 (67). Although hB-ind1 lacks the conserved active motif, the interaction of NS5A with the coiled-coil domain in the central region of hB-ind1 may have an effect on the phosphorylation of host proteins involved in the replication of HCV.

Hsp90 has been shown to be involved in the enzymatic activity and intracellular localization of several viral polymerases, including those of influenza virus (39, 42), herpes simplex virus type 1 (5), and Flock house virus (25). Knockdown and treatment with an Hsp90 inhibitor have revealed that Hsp90 activity is important for the rapid growth of negative-strand RNA viruses (9). Furthermore, Hsp90 has been shown to be required for the activity of hepatitis B virus reverse

transcriptase (21, 22). Although the precise mechanisms by which Hsp90 and FKBP8 cooperate with NS5A to improve the *in vivo* replication of HCV have not been clarified yet, treatment with Hsp90 inhibitors in combination with IFN reduced HCV replication in mice xenotransplanted with human liver fragments (43).

In this study, hB-ind1 was shown to interact with Hsp90 through the FxxW motif in the N-terminal p23 homology domain, and the interaction of hB-ind1 with Hsp90 was shown to be further intensified by the expression of FKBP8, suggesting that FKBP8 and hB-ind1 cooperatively recruit Hsp90 to the HCV replication complex. Furthermore, hB-ind1 was shown to be involved in HCV genomic RNA replication and particle production through the interaction with NS5A and Hsp90. These results suggest that hB-ind1 may be involved in the Hsp90 chaperone pathway in a function similar to that of p23 in cooperation with immunophilins such as FKBP8 and that it plays a crucial role in HCV replication in terms of the correct folding of the replication complex required for efficient enzymatic activity. In addition, cyclophilin B may also participate in the translocation of NS5B, as seen in the polymerase subunits of influenza virus, to facilitate binding to the viral RNA. In contrast to cyclosporine A, FK506 *per se* exhibits no inhibition of RNA replication in HCV replicon cells (65). FKBP8 is a member of the FKBP family but lacks several amino acid residues required for peptidyl-prolyl *cis-trans* isomerase and FK506 binding activities (29). Therefore, nonimmunosuppressive FK506 derivatives that are capable of binding to FKBP8 may exhibit anti-HCV activity. Recently, geldanamycin, an inhibitor of Hsp90, was shown to drastically impair the replication of poliovirus without any escape mutant emerging (15).

Therefore, elucidation of host proteins, including immunophilins, cochaperones, and chaperones, participating in the HCV replication complex may lead to the development of new therapeutics for chronic hepatitis C with a broad spectrum and a low possibility of emergence of breakthrough viruses against antiviral drugs.

In conclusion, in this study we demonstrated that hB-ind1 is involved in HCV replication through interactions with NSSA, FKBP8, and Hsp90. Further clarification of the relationship between viral and host proteins is needed in order to understand the precise mechanism of HCV replication.

ACKNOWLEDGMENTS

We thank H. Murase for secretarial work. We also thank T. Wakita, F. Chisari, and R. Bartenschlager for providing the infectious clones of JFH1, Huh7.5.1, and replicon cell lines, respectively.

This work was supported in part by grants-in-aid from the Ministry of Health, Labor, and Welfare; the Ministry of Education, Culture, Sports, Science, and Technology; the 21st Century Center of Excellence Program; and the Foundation for Biomedical Research and Innovation.

REFERENCES

- Appel, N., T. Pietschmann, and R. Bartenschlager. 2005. Mutational analysis of hepatitis C virus nonstructural protein 5A: potential role of differential phosphorylation in RNA replication and identification of a genetically flexible domain. *J. Virol.* 79:3187–3194.
- Bohen, S. P., A. Kralli, and K. R. Yamamoto. 1995. Hold 'em and fold 'em: chaperones and signal transduction. *Science* 268:1303–1304.
- Bost, A. G., D. Venable, L. Liu, and B. A. Heinz. 2003. Cytoskeletal requirements for hepatitis C virus (HCV) RNA synthesis in the HCV replicon cell culture system. *J. Virol.* 77:4401–4408.
- Bryan, B. A., D. Li, X. Wu, and M. Liu. 2005. The Rho family of small GTPases: crucial regulators of skeletal myogenesis. *Cell. Mol. Life Sci.* 62:1547–1555.
- Burch, A. D., and S. K. Weller. 2005. Herpes simplex virus type 1 DNA polymerase requires the mammalian chaperone hsp90 for proper localization to the nucleus. *J. Virol.* 79:10740–10749.
- Burridge, K., and K. Wennerberg. 2004. Rho and Rac take center stage. *Cell* 116:167–179.
- Cerny, A., and F. V. Chisari. 1999. Pathogenesis of chronic hepatitis C: immunological features of hepatic injury and viral persistence. *Hepatology* 30:595–601.
- Chung, K. M., J. Lee, J. E. Kim, O. K. Song, S. Cho, J. Lim, M. Seedorf, B. Hahn, and S. K. Jang. 2000. Nonstructural protein 5A of hepatitis C virus inhibits the function of karyopherin $\beta 3$. *J. Virol.* 74:5233–5241.
- Connor, J. H., M. O. McKenzie, G. D. Parks, and D. S. Lyles. 2007. Antiviral activity and RNA polymerase degradation following Hsp90 inhibition in a range of negative strand viruses. *Virology* 362:109–119.
- Courilleau, D., E. Chastre, M. Sabbah, G. Redeuilh, A. Athf, and J. Mester. 2000. B-ind1, a novel mediator of Rac1 signaling cloned from sodium butyrate-treated fibroblasts. *J. Biol. Chem.* 275:17344–17348.
- Dittmar, K. D., D. R. Demady, L. F. Stancato, P. Krishna, and W. B. Pratt. 1997. Folding of the glucocorticoid receptor by the heat shock protein (hsp) 90-based chaperone machinery. The role of p23 is to stabilize receptor:hsp90 heterocomplexes formed by hsp90:p60:hsp70. *J. Biol. Chem.* 272:21213–21220.
- Dittmar, K. D., K. A. Hutchison, J. K. Owens-Grillo, and W. B. Pratt. 1996. Reconstitution of the steroid receptor: hsp90 heterocomplex assembly system of rabbit reticulocyte lysate. *J. Biol. Chem.* 271:12833–12839.
- Frydman, J., and J. Hoffeld. 1997. Chaperones get in touch: the Hip-Hop connection. *Trends Biochem. Sci.* 22:87–92.
- Gale, M. J., Jr., M. J. Korh, N. M. Tang, S. L. Tan, D. A. Hopkins, T. E. Dever, S. J. Polyak, D. R. Gretsch, and M. G. Katze. 1997. Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein. *Virology* 230:217–227.
- Geller, R., M. Vignuzzi, R. Andino, and J. Frydman. 2007. Evolutionary constraints on chaperone-mediated folding provide an antiviral approach refractory to development of drug resistance. *Genes Dev.* 21:195–205.
- Hamamoto, I., Y. Nishimura, T. Okamoto, H. Aizaki, M. Liu, Y. Mori, T. Abe, T. Suzuki, M. M. Lai, T. Miyamura, K. Moriishi, and Y. Matsuura. 2005. Human VAP-B is involved in hepatitis C virus replication through interaction with NSSA and NSSB. *J. Virol.* 79:13473–13482.
- Hardy, R. W., J. Marcotrigiano, K. J. Blight, J. E. Majors, and C. M. Rice. 2003. Hepatitis C virus RNA synthesis in a cell-free system isolated from replicon-containing hepatoma cells. *J. Virol.* 77:2029–2037.
- He, Y., H. Nakao, S. L. Tan, S. J. Polyak, P. Neddermann, S. Vijayarsi, B. L. Jacobs, and M. G. Katze. 2002. Subversion of cell signaling pathways by hepatitis C virus nonstructural 5A protein via interaction with Grb2 and P85 phosphatidylinositol 3-kinase. *J. Virol.* 76:9207–9217.
- Ho, S. N., H. D. Hunt, R. M. Horton, J. K. Pullen, and L. R. Pease. 1989. Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene* 77:51–59.
- Honda, M., S. Kaneko, E. Matsushita, K. Kobayashi, G. A. Abell, and S. M. Lemon. 2000. Cell cycle regulation of hepatitis C virus internal ribosomal entry site-directed translation. *Gastroenterology* 118:152–162.
- Hu, J., and C. Seeger. 1996. Hsp90 is required for the activity of a hepatitis B virus reverse transcriptase. *Proc. Natl. Acad. Sci. USA* 93:1060–1064.
- Hu, J., D. O. Toft, and C. Seeger. 1997. Hepadnavirus assembly and reverse transcription require a multi-component chaperone complex which is incorporated into nucleocapsids. *EMBO J.* 16:59–68.
- Huang, D. C., S. C. Cory, and A. Strasser. 1997. Bcl-2, Bcl-XL and adenovirus protein E1B19kD are functionally equivalent in their ability to inhibit cell death. *Oncogene* 14:405–414.
- Inoue, K., K. Sekiyama, M. Yamada, T. Watanabe, H. Yasuda, and M. Yoshida. 2003. Combined interferon $\alpha 2b$ and cyclosporin A in the treatment of chronic hepatitis C: controlled trial. *J. Gastroenterol.* 38:567–572.
- Kammüller, K. M., and D. J. Miller. 2005. The cellular chaperone heat shock protein 90 facilitates Flock House virus RNA replication in *Drosophila* cells. *J. Virol.* 79:6827–6837.
- Kapadia, S. B., and F. V. Chisari. 2005. Hepatitis C virus RNA replication is regulated by host geranylgeranylation and fatty acids. *Proc. Natl. Acad. Sci. USA* 102:2561–2566.
- Kosano, H., B. Stensgaard, M. C. Charlesworth, N. McMahon, and D. Toft. 1998. The assembly of progesterone receptor-hsp90 complexes using purified proteins. *J. Biol. Chem.* 273:32973–32979.
- Lai, V. C., S. Dempsey, J. Y. Lau, Z. Hong, and W. Zhong. 2003. In vitro RNA replication directed by replicase complexes isolated from the subgenomic replicon cells of hepatitis C virus. *J. Virol.* 77:2295–2300.
- Lam, E., M. Martin, and G. Wiederrecht. 1995. Isolation of a cDNA encoding a novel human FKS56-binding protein homolog containing leucine zipper and tetratricopeptide repeat motifs. *Gene* 160:297–302.
- Lan, K. H., K. L. Lan, W. P. Lee, M. L. Sheu, M. Y. Chen, Y. L. Lee, S. H. Yen, F. Y. Chang, and S. D. Lee. 2007. HCV NSSA inhibits interferon-alpha signaling through suppression of STAT1 phosphorylation in hepatocyte-derived cell lines. *J. Hepatol.* 46:759–767.
- Le, S. S., F. A. Locks, H. Udo, S. Richardson-Burns, R. A. Phelps, R. J. Bouchard, H. Barth, K. Auctories, K. L. Tyler, E. R. Kandef, K. A. Heidenreich, and D. A. Linseman. 2005. Inhibition of Rac GTPase triggers a C-Jun- and Bim-dependent mitochondrial apoptotic cascade in cerebellar granule neurons. *J. Neurochem.* 94:1025–1039.
- Lindenbach, B. D., M. J. Evans, A. J. Syder, B. Volk, T. L. Tellinghuisen, C. C. Liu, T. Maruyama, R. O. Hynes, D. R. Burton, J. A. McKeating, and C. M. Rice. 2005. Complete replication of hepatitis C virus in cell culture. *Science* 309:623–626.
- Liu, J., J. D. Farmer, Jr., W. S. Lane, J. Friedman, I. Weissman, and S. L. Schreiber. 1991. Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell* 66:807–815.
- Lohmann, V., F. Korner, J. Koch, U. Herian, L. Theilmann, and R. Bartenschlager. 1999. Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science* 285:110–113.
- Macdonald, A., K. Crowder, A. Street, C. McCormick, and M. Harris. 2004. The hepatitis C virus NSSA protein binds to members of the Src family of tyrosine kinases and regulates kinase activity. *J. Gen. Virol.* 85:721–729.
- Majumder, M., A. K. Ghosh, R. Steele, R. Ray, and R. B. Ray. 2001. Hepatitis C virus NSSA physically associates with p53 and regulates p21/waf1 gene expression in a p53-dependent manner. *J. Virol.* 75:1401–1407.
- Manns, M. P., M. Cornberg, and H. Wedemeyer. 2001. Current and future treatment of hepatitis C. *Indian J. Gastroenterol.* 20(Suppl. 1):C47–C51.
- Mercer, D. F., D. E. Schiller, J. F. Elliott, D. N. Douglas, C. Hao, A. Rinfret, W. R. Addison, K. P. Fischer, T. A. Churchill, J. R. Lakey, D. L. Tyrrell, and N. M. Kneteman. 2001. Hepatitis C virus replication in mice with chimeric human livers. *Nat. Med.* 7:927–933.
- Momose, F., T. Naito, K. Yano, S. Sugimoto, Y. Morikawa, and K. Nagata. 2002. Identification of Hsp90 as a stimulatory host factor involved in influenza virus RNA synthesis. *J. Biol. Chem.* 277:45306–45314.
- Moriishi, K., and Y. Matsuura. 2003. Mechanisms of hepatitis C virus infection. *Antivir. Chem. Chemother.* 14:285–297.
- Murata, T., T. Ohshima, M. Yamaji, M. Hosaka, Y. Miyayari, M. Hijikata, and K. Shimotohno. 2005. Suppression of hepatitis C virus replicon by TGF- β . *Virology* 331:407–417.
- Naito, T., F. Momose, A. Kawaguchi, and K. Nagata. 2007. Involvement of Hsp90 in assembly and nuclear import of influenza virus RNA polymerase subunits. *J. Virol.* 81:1339–1349.
- Nakagawa, S., T. Umehara, C. Matsuda, S. Kuge, M. Sudoh, and M. Kohara.

2007. Hsp90 inhibitors suppress HCV replication in replicon cells and humanized liver mice. *Biochem. Biophys. Res. Commun.* 353:882-888.
44. Obermann, W. M., H. Sondermann, A. A. Russo, N. P. Pavletich, and F. U. Hartl. 1998. In vivo function of Hsp90 is dependent on ATP binding and ATP hydrolysis. *J. Cell Biol.* 143:901-910.
 45. Okamoto, T., Y. Nishimura, T. Ichimura, K. Suzuki, T. Miyamura, T. Suzuki, K. Moritshi, and Y. Matsura. 2006. Hepatitis C virus RNA replication is regulated by FKBP8 and Hsp90. *EMBO J.* 25:5015-5025.
 46. Pietschmann, T., V. Lohmann, A. Kaul, N. Krieger, G. Rinck, G. Ratter, D. Strand, and R. Bartenschlager. 2002. Persistent and transient replication of full-length hepatitis C virus genomes in cell culture. *J. Virol.* 76:4008-4021.
 47. Prapapanich, V., S. Chen, E. J. Toran, R. A. Rimerman, and D. F. Smith. 1996. Mutational analysis of the hsp70-interacting protein Hip. *Mol. Cell Biol.* 16:6200-6207.
 48. Qadri, L., M. Iwahashi, and F. Simon. 2002. Hepatitis C virus NS5A protein binds TBP and p53, inhibiting their DNA binding and p53 interactions with TBP and ERCC3. *Biochim. Biophys. Acta* 1592:193-204.
 49. Ridley, A. J., H. F. Paterson, C. L. Johnston, D. Diekmann, and A. Hall. 1992. The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. *Cell* 70:401-410.
 50. Riggs, D. L., M. B. Cox, J. Cheung-Flynn, V. Prapapanich, P. E. Carrigan, and D. F. Smith. 2004. Functional specificity of co-chaperone interactions with Hsp90 client proteins. *Crit. Rev. Biochem. Mol. Biol.* 39:279-295.
 51. Scholle, F., K. Li, F. Bodola, M. Ikeda, B. A. Luxon, and S. M. Lemon. 2004. Virus-host cell interactions during hepatitis C virus RNA replication: impact of polyprotein expression on the cellular transcriptome and cell cycle association with viral RNA synthesis. *J. Virol.* 78:1513-1524.
 52. Shi, S. T., S. J. Polyak, H. Tu, D. R. Taylor, D. R. Gretch, and M. M. Lai. 2002. Hepatitis C virus NS5A colocalizes with the core protein on lipid droplets and interacts with apolipoproteins. *Virology* 292:198-210.
 53. Strader, D. B., T. Wright, D. L. Thomas, and L. B. Seef. 2004. Diagnosis, management, and treatment of hepatitis C. *Hepatology* 39:1147-1171.
 54. Sullivan, W., B. Stensgard, G. Caucutt, B. Bartha, N. McMahon, E. S. Alnemri, G. Litwack, and D. Toft. 1997. Nucleotides and two functional states of hsp90. *J. Biol. Chem.* 272:8007-8012.
 55. Takamizawa, A., C. Mori, I. Fuke, S. Manabe, S. Murakami, J. Fujita, E. Onishi, T. Andoh, I. Yoshida, and H. Okayama. 1991. Structure and organization of the hepatitis C virus genome isolated from human carriers. *J. Virol.* 65:1105-1113.
 56. Tan, S. L., H. Nakao, Y. He, S. Vijaypri, P. Neddermann, B. L. Jacobs, B. J. Mayer, and M. G. Katze. 1999. NS5A, a nonstructural protein of hepatitis C virus, binds growth factor receptor-bound protein 2 adaptor protein in a Src homology 3 domain/ligand-dependent manner and perturbs mitogenic signaling. *Proc. Natl. Acad. Sci. USA* 96:5533-5538.
 57. Tiganis, T., and A. M. Bennett. 2007. Protein tyrosine phosphatase function: the substrate perspective. *Biochem. J.* 402:1-15.
 58. Tsukiyama-Kohara, K., N. Iizuka, M. Kohara, and A. Nomoto. 1992. Internal ribosome entry site within hepatitis C virus RNA. *J. Virol.* 66:1476-1483.
 59. Tu, H., L. Gao, S. T. Shi, D. R. Taylor, T. Yang, A. K. Mircheff, Y. Wen, A. E. Gorbalenya, S. B. Hwang, and M. M. Lai. 1999. Hepatitis C virus RNA polymerase and NS5A complex with a SNARE-like protein. *Virology* 263:30-41.
 60. Uwanogho, D. A., Z. Hardcastle, P. Balogh, G. Mirza, K. L. Thornburg, J. Ragoussis, and P. T. Sharpe. 1999. Molecular cloning, chromosomal mapping, and developmental expression of a novel protein tyrosine phosphatase-like gene. *Genomics* 62:406-416.
 61. Venkatesan, A., R. Sharma, and A. Dasgupta. 2003. Cell cycle regulation of hepatitis C and encephalomyocarditis virus internal ribosome entry site-mediated translation in human embryonic kidney 293 cells. *Virus Res.* 94:85-95.
 62. Wakita, T., T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H. G. Krausslich, M. Mizokami, R. Bartenschlager, and T. J. Liang. 2005. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat. Med.* 11:791-796.
 63. Wang, C., M. Gale, Jr., B. C. Keller, H. Huang, M. S. Brown, J. L. Goldstein, and J. Ye. 2005. Identification of FBL2 as a geranylgeranylated cellular protein required for hepatitis C virus RNA replication. *Mol. Cell* 18:425-434.
 64. Wasley, A., and M. J. Alter. 2000. Epidemiology of hepatitis C: geographic differences and temporal trends. *Semin. Liver Dis.* 20:1-16.
 65. Watashi, K., M. Hijikata, M. Hosaka, M. Yamaji, and K. Shimotohno. 2003. Cyclosporin A suppresses replication of hepatitis C virus genome in cultured hepatocytes. *Hepatology* 38:1282-1288.
 66. Watashi, K., N. Ishii, M. Hijikata, D. Inoue, T. Murata, Y. Miyazaki, and K. Shimotohno. 2005. Cyclophilin B is a functional regulator of hepatitis C virus RNA polymerase. *Mol. Cell* 19:111-122.
 67. Wertheimer, A. M., S. J. Polyak, R. Leistikow, and H. R. Rosen. 2007. Engulfment of apoptotic cells expressing HCV proteins leads to differential chemokine expression and STAT signaling in human dendritic cells. *Hepatology* 45:1422-1432.
 68. Wochnik, G. M., J. C. Young, U. Schmidt, F. Holsboer, F. U. Hartl, and T. Rein. 2004. Inhibition of GR-mediated transcription by p23 requires interaction with Hsp90. *FEBS Lett.* 560:35-38.
 69. Xue, Y., F. Bi, X. Zhang, Y. Pan, N. Liu, Y. Zheng, and D. Fan. 2004. Inhibition of endothelial cell proliferation by targeting Rac1 GTPase with small interference RNA in tumor cells. *Biochem. Biophys. Res. Commun.* 320:1309-1315.
 70. Ye, J., C. Wang, R. Sumpter, Jr., M. S. Brown, J. L. Goldstein, and M. Gale, Jr. 2003. Disruption of hepatitis C virus RNA replication through inhibition of host protein geranylgeranylation. *Proc. Natl. Acad. Sci. USA* 100:15865-15870.
 71. Yi, M., and S. M. Lemon. 2004. Adaptive mutations producing efficient replication of genotype 1a hepatitis C virus RNA in normal Huh7 cells. *J. Virol.* 78:7904-7915.
 72. Young, J. C., and F. U. Hartl. 2000. Polypeptide release by Hsp90 involves ATP hydrolysis and is enhanced by the co-chaperone p23. *EMBO J.* 19:5930-5940.
 73. Zech, B., A. Kurtenbach, N. Krieger, D. Strand, S. Blencke, M. Morbitzer, K. Salasidis, M. Cotten, J. Wissing, S. Obert, R. Bartenschlager, T. Herget, and H. Daub. 2003. Identification and characterization of amphiphysin II as a novel cellular interaction partner of the hepatitis C virus NS5A protein. *J. Gen. Virol.* 84:555-560.
 74. Zhong, J., P. Gastaminza, G. Cheng, S. Kapadia, T. Kato, D. R. Burton, S. F. Wieland, S. L. Uprichard, T. Wakita, and F. V. Chisari. 2005. Robust hepatitis C virus infection in vitro. *Proc. Natl. Acad. Sci. USA* 102:9294-9299.

Interaction of Hepatitis C Virus Nonstructural Protein 5A with Core Protein Is Critical for the Production of Infectious Virus Particles[▽]

Takahiro Masaki,¹ Ryosuke Suzuki,¹ Kyoko Murakami,¹ Hideki Aizaki,¹ Koji Ishii,¹ Asako Murayama,¹ Tomoko Date,¹ Yoshiharu Matsuura,² Tatsuo Miyamura,¹ Takaji Wakita,¹ and Tetsuro Suzuki^{1*}

Department of Virology II, National Institute of Infectious Diseases, Shinjuku-ku, Tokyo 162-8640, Japan,¹ and Department of Molecular Virology, Research Institute for Microbial Diseases, Osaka University, Suita-shi, Osaka 565-0871, Japan²

Received 17 April 2008/Accepted 22 May 2008

Nonstructural protein 5A (NS5A) of the hepatitis C virus (HCV) possesses multiple and diverse functions in RNA replication, interferon resistance, and viral pathogenesis. Recent studies suggest that NS5A is involved in the assembly and maturation of infectious viral particles; however, precisely how NS5A participates in virus production has not been fully elucidated. In the present study, we demonstrate that NS5A is a prerequisite for HCV particle production as a result of its interaction with the viral capsid protein (core protein). The efficiency of virus production correlated well with the levels of interaction between NS5A and the core protein. Alanine substitutions for the C-terminal serine cluster in domain III of NS5A (amino acids 2428, 2430, and 2433) impaired NS5A basal phosphorylation, leading to a marked decrease in NS5A-core interaction, disturbance of the subcellular localization of NS5A, and disruption of virion production. Replacing the same serine cluster with glutamic acid, which mimics the presence of phosphoserines, partially preserved the NS5A-core interaction and virion production, suggesting that phosphorylation of these serine residues is important for virion production. In addition, we found that the alanine substitutions in the serine cluster suppressed the association of the core protein with viral genome RNA, possibly resulting in the inhibition of nucleocapsid assembly. These results suggest that NS5A plays a key role in regulating the early phase of HCV particle formation by interacting with core protein and that its C-terminal serine cluster is a determinant of the NS5A-core interaction.

Hepatitis C virus (HCV) infection is a major public health problem and is prevalent in about 200 million people worldwide (27, 40, 42). Current protocols for treating HCV infection fail to produce a sustained virological response in as many as half of treated individuals, and many cases progress to chronic liver disease, including chronic hepatitis, cirrhosis, and hepatocellular carcinoma (15, 31, 35, 43).

HCV is a positive-strand RNA virus classified in the *Hepacivirus* genus within the *Flaviviridae* family (55). Its approximately 9.6-kb genome is translated into a single polypeptide of about 3,000 amino acids (aa), in which the structural proteins core, E1, and E2 reside in the N-terminal region. A crucial function of core protein is assembly of the viral nucleocapsid. The amino acid sequence of this protein is well conserved among different HCV strains compared to other HCV proteins. The nonstructural (NS) proteins NS3-NS5B are considered to assemble into a membrane-associated HCV RNA replicase complex. NS3 possesses the enzymatic activities of serine protease and RNA helicase, and NS4A serves as a cofactor for NS3 protease. NS4B plays a role in the remodeling of host cell membranes, probably to generate the site for the replicase assembly. NS5B functions as the RNA-dependent RNA polymerase. NS5A is known to play an important but undefined role in viral RNA replication.

NS5A is a phosphoprotein that can be found in basally phosphorylated (56 kDa) and hyperphosphorylated (58 kDa) forms (49). Comparative sequence analyses and limited proteolysis of recombinant NS5A have demonstrated that NS5A is composed of three domains (52). Domain I is relatively conserved among HCV genotypes compared to domains II and III. Analysis of the crystal structure of the conserved domain I that immediately follows the membrane-anchoring α -helix localized at the N terminus revealed a dimeric structure (53). The interface between protein molecules is characterized by a large, basic groove, which has been proposed as a site of RNA binding. In fact, its RNA binding property has been demonstrated biochemically (17). Domains II and III of NS5A are far less understood. Domain II contains a region referred to as the interferon sensitivity determining region, and this region and its C-terminal 26 residues have been shown to be essential for interaction with the interferon-induced, double-stranded RNA-dependent protein kinase (6–10, 38, 39, 48). Domain III includes a number of potential phosphoacceptor sites and is most likely involved in basal phosphorylation. This domain tolerates insertion of large heterologous sequences such as green fluorescent protein (GFP) and is not required for function of NS5A in HCV RNA replication (1, 34). However, a study with the recently established productive HCV cell culture system using genotype 2a isolate JFH-1 (28, 56, 58) demonstrated that while insertion of GFP within the NS5A region does not affect RNA replication, it does produce marked decreases in the production of infectious virus particles (41). This suggests that the C-terminal region of NS5A may affect virus particle production independent of RNA replication. Re-

* Corresponding author. Mailing address: Department of Virology II, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan. Phone: 81 3 5285 1111. Fax: 81 3 5285 1161. E-mail: tesuzuki@nih.go.jp.

[▽] Published ahead of print on 4 June 2008.

cently, Miyazaki et al. reported that the association of core protein with the NS proteins and replication complexes around lipid droplets (LDs) is critical for producing infectious viruses (33).

In the present study, we demonstrated that NSSA is a prerequisite for HCV particle production via its interaction with core protein, and we identified serine residues in the C-terminal region of NSSA that play an important role in virion production. Substitution of the serine residues with alanine residues inhibited not only the interaction of NSSA with core protein but also HCV RNA-core association and led to a decrease in HCV particle production with no effect on RNA replication.

MATERIALS AND METHODS

DNA construction. Plasmids pJFH1, which contains the full-length JFH-1 cDNA downstream of the T7 RNA promoter sequence, and pSGR-JFH1/Luc, in which the neomycin resistance gene of pSGR-JFH1 has been replaced by the firefly luciferase reporter gene, have been previously described (24, 56). To generate the fluorochrome gene-tagged full-length JFH-1 plasmid, pJFH1/NSSA-GFP, the region encompassing the RsrII site of NSSA and the BsrGI site of NSSB was amplified by PCR, the amplification product was cloned into pGEM-T Easy vector (Promega, Madison, WI), and the resultant plasmid was designated pGEM-JFH1/RsrII-BsrGI. A GFP reporter gene was amplified by PCR from pGreen-Lantern-1 (Invitrogen, Carlsbad, CA) with primers containing the XhoI site and inserted, after restriction digestion with XhoI, into the XhoI site of pGEM-JFH1/RsrII-BsrGI. The resulting plasmid was digested by RsrII and BsrGI and ligated into pJFH1 similarly digested by RsrII and BsrGI to produce pJFH1/NSSA-GFP. For generation of the fluorochrome gene-tagged subgenomic reporter plasmid, pJFH1/NSSA-GFP was digested by RsrII and SnaBI and ligated into pSGR-JFH1/Luc similarly digested by RsrII and SnaBI. The mutations in the NSSA gene were generated by oligonucleotide-directed mutagenesis (57). To construct plasmids expressing N-terminally FLAG-tagged HCV core protein or hemagglutinin (HA)-tagged NSSA, DNA fragments encoding core protein or NSSA (wild type or mutants) were generated from the full-length JFH-1 cDNA by PCR. The core protein coding sequence, together with a FLAG sequence linked to its N terminus, was cloned into the pCAGGS vector (37). The coding sequences of NSSA, together with an HA sequence linked to their N termini, were also cloned into pCAGGS vectors. All PCR products were confirmed by automated nucleotide sequencing with an ABI Prism 3130 Avant Genetic Analyzer (Applied Biosystems, Tokyo, Japan).

Cells and viruses. The human hepatoma cell line, Huh-7, and JFH1/4-1 cells, which are Huh-7 cells carrying a subgenomic replicon of JFH-1 (32), were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with minimal essential medium nonessential amino acids (Invitrogen), 100 units/ml of penicillin, 100 µg/ml of streptomycin, and 10% fetal bovine serum (FBS) at 37°C in a 5% CO₂ incubator. Huh-c/p7 cells, which are Huh-7 cells stably expressing the proteins core to p7 derived from the JFH-1 strain (18), were incubated in DMEM containing 300 µg/ml of zeocin (Invitrogen). HCV particles derived from JFH-1 were produced by transient transfection of Huh-7 cells with in vitro transcribed RNA, as described previously (56, 58). Recombinant vaccinia virus strain DIs, which expresses the bacteriophage T7 RNA polymerase under the control of the vaccinia virus early/late promoter P7.5, was generated and propagated as previously described (19).

DNA transfection, immunoprecipitation (IP), and immunoblotting. For coexpression of FLAG-tagged core protein and HA-tagged NSSA, cells were seeded onto 35-mm wells of a six-well cell culture plate and cultured overnight. Plasmid DNAs (2 µg) were transfected into cells using TransIT-LTI transfection reagent (Mirus, Madison, WI). Cells were harvested at 48 h posttransfection, washed three times with 1 ml of ice-cold phosphate-buffered saline (PBS), and suspended in 0.25 ml lysis buffer (20 mM Tris-HCl [pH 7.4] containing 135 mM NaCl, 1% Triton X-100, 0.05% sodium dodecyl sulfate [SDS], and 10% glycerol) supplemented with 50 mM NaF, 5 mM Na₃VO₄, 1 µg/ml leupeptin, and 1 mM phenylmethylsulfonyl fluoride (PMSF). Cell lysates were sonicated at 4°C for 5 min, incubated for 30 min at 4°C, and centrifuged at 14,000 × g for 5 min at 4°C. After preclearing, the supernatant was immunoprecipitated with 10 µl of anti-FLAG M2-agarose beads (Sigma, St. Louis, MO). For expression of the full-length HCV polyprotein, Huh-7 cells transfected with 10 µg of in vitro transcribed RNAs by electroporation were resuspended in 20 or 30 ml of culture

medium, and 10-ml aliquots were seeded into 100-mm culture dishes. At 72 h posttransfection, the cells were incubated in 0.5 ml of lysis buffer (20 mM Tris-HCl [pH 7.4] containing 135 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, and 10% glycerol) supplemented with 50 mM NaF, 5 mM Na₃VO₄, 1 µg/ml leupeptin, and 1 mM PMSF. After preclearing, the supernatant was immunoprecipitated with 5 µg of polyclonal anti-NSSA antibody (34a) or polyclonal anti-C/EBPβ antibody (Santa Cruz Biotechnology, Santa Cruz, CA), and 20 µl of protein G-agarose beads (Invitrogen). The immunocomplex was precipitated with the beads by centrifugation at 800 × g for 30 s and then washed five times with lysis buffer by centrifugation. The proteins binding to the beads were boiled in 20 µl of SDS sample buffer and then subjected to SDS-12.5% polyacrylamide gel electrophoresis (PAGE). The proteins were transferred onto a polyvinylidene difluoride membrane (Immobilon; Millipore, Bedford, MA) and then reacted with a primary antibody and a secondary horseradish peroxidase-conjugated antibody. The immunocomplexes were visualized with an ECL Plus Western Blotting Detection System (GE Healthcare, Buckinghamshire, United Kingdom) and detected using a LAS-3000 imaging analyzer (FujiFilm, Tokyo, Japan).

In vitro synthesis of HCV RNA and RNA transfection. Plasmid DNAs were digested with XbaI and treated with mung bean nuclease (New England Biolabs, Ipswich, MA) to remove the four terminal nucleotides, resulting in the correct 3' end of the HCV cDNA. Digested DNAs were purified and used as templates for RNA synthesis. HCV RNA was synthesized in vitro using a MEGAscript T7 kit (Ambion, Austin, TX). Synthesized RNA was treated with DNase I (Ambion), followed by acid guanidinium thiocyanate-phenol-chloroform extraction to remove any remaining template DNA. Synthesized HCV RNAs were used for electroporation. Trypsinized Huh-7 cells were washed with Opti-MEM 1 reduced-serum medium (Invitrogen) and resuspended at 3×10^6 cells/ml with Cytochrome buffer (54). RNA was mixed with 400 µl of cell suspension and transferred into an electroporation cuvette (Precision Universal Cuvettes; Thermo Hybrid, Middlesex, United Kingdom). Cells were then pulsed at 260 V and 950 µF using a Gene Pulser II unit (Bio-Rad, Hercules, CA). Transfected cells were immediately transferred onto six-well culture plates or 100-mm culture dishes.

Luciferase assay. Cells were harvested at different time points posttransfection of subgenomic reporter replicons and lysed in passive lysis buffer (Promega). The luciferase activity in cells was determined using a luciferase assay system (Promega).

Quantification of HCV core protein. HCV core protein in transfected cells or cell culture supernatants was quantified using a highly sensitive enzyme immunoassay (Ortho HCV antigen ELISA Kit; Ortho Clinical Diagnostics, Tokyo, Japan). To determine intracellular core protein amounts, cell lysates were prepared as described previously (41). To determine the efficiency of core protein release, the ratio of extracellular core protein to total core protein (the sum of intra- and extracellular core protein amounts) was calculated.

Intra- and extracellular infectivity assay. Culture supernatants were harvested 72 h posttransfection, and virus titers were determined by a 50% tissue culture infectious dose (TCID₅₀) assay as described previously (28, 46). Virus titration was performed by seeding naive Huh-7 cells in 96-well plates at a density of 1×10^4 cells/well. Samples were serially diluted fivefold in complete growth medium and used to infect the seeded cells (six wells per dilution). At 72 h after infection, the inoculated cells were fixed and immunostained with a mouse monoclonal anti-core protein antibody (2H9) (56), followed by an Alexa Fluor 488-conjugated anti-mouse immunoglobulin G (IgG) (Invitrogen). Wells that showed at least one core protein-expressing cell was counted as positive. Cell-associated infectivity was determined essentially as described previously (12, 47). Briefly, cells were extensively washed with PBS, scraped, and centrifuged for 3 min at 120 × g. Cell pellets were resuspended in 1 ml of DMEM containing 10% FBS and subjected to four cycles of freezing and thawing using dry ice and a 37°C water bath. Samples were then centrifuged at 2,400 × g for 10 min at 4°C to remove cell debris, and cell-associated infectivity was determined by TCID₅₀ assay.

Expression of HCV proteins using vaccinia viruses, metabolic labeling of cells, and radioimmunoprecipitation analysis. Metabolic labeling of cells and radioimmunoprecipitation analysis were performed as described by Huang et al. (17) with some modifications. A total of 4×10^5 Huh-7 cells were seeded onto each well of six-well cell culture plates and cultured overnight. A 2-µg amount of subgenomic replicon DNAs carrying defined NSSA mutations was transfected into cells using TransIT-LTI transfection reagent, and at 12 h posttransfection the cells were then infected at a multiplicity of infection of 10 with recombinant vaccinia viruses expressing the T7 RNA polymerase. After 40 h of transfection, cells were incubated in methionine- and cysteine-deficient DMEM (Invitrogen) or phosphate-deficient DMEM (Invitrogen) for 2 h and labeled for 6 h with [³⁵S]methionine and [³⁵S]cysteine (200 µCi/well; GE Healthcare) or

[32 P]orthophosphate (250 μ Ci/well; GE Healthcare). The cells were then washed twice with cold PBS and lysed with SDS lysis buffer (50 mM Tris-HCl [pH 7.6], 0.5% SDS, 1 mM EDTA, 20 μ g/ml of PMSF). The cell lysates were passed through a 27-gauge needle several times to shear cellular DNA. After a 10-min incubation at 75°C, the lysates were clarified by centrifugation and diluted five-fold with HNAET buffer (50 mM HEPES [pH 7.5], 150 mM NaCl, 0.67% bovine serum albumin, 1 mM EDTA, 0.33% Triton X-100). After preclearing by incubation with 20 μ l of protein G-agarose beads for 1 h at 4°C, the supernatant was incubated with 2 μ g of rabbit polyclonal anti-NS5A antibody overnight at 4°C. A 20- μ l aliquot of protein G-agarose beads was further added and incubated for 2 h at 4°C. The cell pellets were washed three times with 0.5 ml of HNAETS buffer (HNAET containing 0.5% SDS), followed by washing once with 0.5 ml of HNE buffer (50 mM HEPES [pH 7.5], 150 mM NaCl and 1 mM EDTA). After treatment with or without λ protein phosphatase (New England Biolabs), the cell pellets were suspended in 20 μ l of SDS sample buffer and boiled for 10 min. The proteins were resolved on 10% SDS-polyacrylamide gels and analyzed by autoradiography.

Subcellular fractionation analysis. All steps were carried out at 4°C in the presence of a protease inhibitor cocktail (Complete; Roche, Mannheim, Germany) as described previously (20), with some modifications. Cells were suspended in four cell volumes of homogenization buffer (50 mM NaCl, 10 mM triethylamine [pH 7.4], 1 mM EDTA), snap frozen in liquid nitrogen, stored at -80°C, and thawed in a water bath at room temperature. Supernatants (0.4 ml) were layered on linear 10-ml iodixanol gradients from 2.5 to 25% and centrifuged at 37,000 rpm for 3.5 h in an SW41 rotor (Beckman, Fullerton, CA), followed by collection of 0.8-ml fractions from the top. Each fraction was concentrated by Centricon YM30 (Millipore), separated by SDS-PAGE, and immunoblotted with a rabbit polyclonal anti-calnexin antibody (Stressgen Biotechnologies, Victoria, Canada), a mouse monoclonal anti-adipose differentiation-related protein (ADRP) antibody (Progen Biotechnik, Heidelberg, Germany), or a rabbit polyclonal anti-NS5A antibody. The core protein amount in each fraction was also determined by enzyme-linked immunosorbent assay (ELISA).

IP-RT-PCR. The process of cell lysis to RNA purification was carried out essentially as described by Johnson et al. (21) with some modifications. A total of 3×10^6 Huh-7 cells were transfected with 10 μ g of *in vitro* transcribed HCV RNAs and resuspended in 20 or 30 ml of culture medium, after which 10-ml aliquots were seeded into 100-mm culture dishes. At 72 h posttransfection, the cells were scraped and incubated in 500 μ l of hypotonic buffer (10 mM HEPES [pH 7.6], 1.5 mM MgCl₂, 10 mM KCl, 0.2 mM PMSF) per dish. The cells were passed through a 20-gauge needle several times, lysed with Nonidet P-40 at a final concentration of 1%, and incubated on ice for an additional 10 min. After centrifugation at 4,000 \times g at 4°C for 15 min, glycerol was added to the supernatants at a final concentration of 5%. The cell lysates were incubated with 20 μ l of protein G-agarose beads for 30 min at room temperature. After the cell lysates were removed from protein G-agarose beads, 5 μ g of mouse monoclonal anti-core protein antibody or normal mouse IgG (Sigma) as a negative control was added, and samples were incubated for an additional 1 h at room temperature. A 20- μ l aliquot of protein G-agarose beads per sample was added to the cell lysates and incubated for 1 h. After incubation, the beads were washed three times with wash buffer (10 mM Tris-HCl [pH 7.6], 100 mM KCl, 5 mM MgCl₂, and 1 mM dithiothreitol) and eluted in 100 μ l of elution buffer (50 mM Tris-HCl [pH 8.0], 1% SDS, and 10 mM EDTA) at 65°C for 10 min. After treatment with 100 μ g of proteinase K at 37°C for 30 min, the RNAs in immunocomplexes were isolated by acid guanidinium thiocyanate-phenol-chloroform extraction. Reverse transcriptase PCR (RT-PCR) was carried out using random hexamer and Superscript II RT (Invitrogen), followed by nested PCR with LA *Taq* DNA polymerase (TaKaRa, Shiga, Japan) and primer sets amplifying the fragments of nucleotides (nt) 129 to 2367 and nt 7267 to 9463 of the JFH-1 genome. To amplify the fragment of nt 129 to 2367, the sense primer 5'-CTGTGAGGAAC TACTGTCTT-3' and the antisense primer 5'-TCCACGATGTTCTGGTGAA G-3' were used for first-round PCR; the sense primer 5'-CGGGAGGCCAT AGTGG-3' and the antisense primer 5'-CATTCGGTGTAGAGTGCA-3' were used for second-round PCR. To amplify the fragment of nt 7267 to 9463, the sense primer 5'-GTCCAGGGTGCCTTCTGGA-3' and the antisense primer 5'-GCGGCTCACGGACCTTTCAC-3' were used for first-round PCR; the sense primer 5'-CACCGTTGCTGGTTGTGCT-3' and the antisense primer 5'-GTGTACCTAGTGTGGCCGCTCA-3' were used for second-round PCR.

Indirect immunofluorescence analysis. Cells incubated for 3 days after transfection with JFH-1 RNAs were seeded in an eight-well chamber slide (BD Biosciences, San Jose, CA) and cultured overnight. The adherent cells were washed twice with PBS and fixed with 4% paraformaldehyde at room temperature. After a washing step with PBS, the cells were permeabilized with PBS containing 0.3% Triton X-100 and 2% FBS for 1 h at room temperature and

stained with a rabbit polyclonal anti-NS5A antibody and a mouse monoclonal anti-core protein antibody. The fluorescent secondary antibodies were Alexa Fluor 488- or Alexa Fluor 555-conjugated anti-rabbit or anti-mouse IgG antibodies (Invitrogen). Analyses of JFH-1 were performed on a Zeiss confocal laser scanning microscope LSM 510 (Carl Zeiss, Oberkochen, Germany).

RESULTS

Mutations of serine residues at the NS5A C terminus impair basal phosphorylation but have little effect on viral RNA replication. As demonstrated in a previous study, insertion of GFP into the NS5A C terminus does not significantly affect viral RNA replication but reduces the generation of infectious HCV particles (41). The C-terminal region of NS5A contains highly conserved serine residues that are involved in basal phosphorylation (1, 23, 49). To examine the involvement of the serine clusters (cluster 3-A [CL3A] and cluster 3-B [CL3B]) in the C-terminal region of NS5A in HCV particle production, we created mutated HCV genomes as well as subgenomic replicons carrying alanine substitutions for the conserved serine residues at aa 2384, 2388, 2390, and 2391 (residues are numbered according to the positions within the original JFH-1 polyprotein) (CL3A/SA); at aa 2428, 2430, and 2433 (CL3B/SA); or an in-frame deletion spanning aa 2384 to 2433 (Δ 2384-2433) (Fig. 1). A construct with an in-frame insertion of GFP (NS5A-GFP) was also generated as described previously for the Con1 isolate (34).

First, we analyzed the effects of the NS5A mutations on HCV RNA replication using a transient RNA replication assay using subgenomic luciferase reporter replicons (Fig. 2A) and found that the serine-to-alanine substitutions (CL3A/SA and CL3B/SA) did not affect viral RNA replication. NS5A-GFP and Δ 2384-2433 slightly reduced RNA replication, indicating that the mutations of the NS5A C terminus tested in this study do not critically affect RNA replication, which is consistent with previous reports (1, 34, 51).

Next, the phosphorylation status of the mutated NS5A was analyzed as described in Materials and Methods (Fig. 2B). NS5A was isolated from radiolabeled cells by IP and analyzed either directly by SDS-PAGE or after treatment with λ protein phosphatase. Analysis of 32 P-radiolabeled proteins revealed that the CL3A/SA, CL3B/SA, and Δ 2384-2433 mutations resulted in marked reduction of basal phosphorylation (Fig. 2B, compare lane 1 with lanes 3, 5, and 7 in the top panel). All 32 P-labeled NS5A proteins were sensitive to treatment with phosphatase (lanes 2, 4, 6, and 8). The possibility that loss of signal after dephosphorylation was due to contaminating proteases present in the phosphatase preparations can be ruled out because no degradation of the 35 S-labeled proteins was observed (Fig. 2B, bottom panel). These results suggest that mutations in the C-terminal serine cluster of NS5A impair basal phosphorylation but have no significant effect on viral RNA replication.

Effect of mutations introduced into the NS5A C terminus on the production of infectious HCV particles. To analyze HCV particle production from cells transfected with the *in vitro* transcribed viral genomic RNAs, we harvested supernatants and cells at 4, 24, 48, 72, and 96 h posttransfection and measured the amounts of core protein. As shown in Fig. 3A, comparable amounts of core proteins were detected in all transfected cells 4 h after transfection, reflecting unchanged




FIG. 1. Structures of HCV constructs used in this study. Schematic diagram of the NS5A structure according to Tellinghuisen et al. (52) is shown in the top panel. The three domains are indicated by white boxes and are separated by trypsin-sensitive regions with presumably low structural complexity (low-complexity sequence [LCS]). The numbers indicate amino acid residues within the original JFH-1 polyprotein. The names listed on the left represent full-length HCV constructs, subgenomic reporter replicons, or N-terminally HA-tagged NS5A constructs used in this study. NS5A-GFP carries a GFP insertion between aa 2394 and 2395 as indicated by a shaded box. CL3A/SA and CL3B/SA carry several serine-to-alanine substitutions in the NS5A C terminus constructed as described previously (1). HCV constructs from S2428A to S2430/2433A carry single or double serine-to-alanine substitutions generated by modification of the CL3B/SA construct. The Δ2384–2433 mutant possesses an in-frame deletion in the C-terminal region of NS5A. Amino acid substitutions are marked in bold and underlined. N and C represent N terminus and C terminus, respectively.

transfection efficiencies, and the kinetics of intracellular core protein levels was similar among transfectants. By contrast, core protein released from cells transfected either with the mutated genome of CL3B/SA, Δ2384–2433, or NS5A-GFP was more than 10-fold lower than that for the wild-type JFH-1 or CL3A/SA (Fig. 3B). Figure 3C shows the efficiency of core protein release from each transfectant, which is expressed as a percentage of the extracellular core protein level relative to the amount of total core protein (the sum of intra- and extracellular core protein). Core protein release efficiency with the wild type and CL3A/SA was 2 to 13% at 48 to 96 h after transfection, while only 1% or less of core protein was released in the cases of CL3B/SA, Δ2384–2433, and NS5A-GFP strains.

To further investigate production and release of infectious virus particles, naïve Huh-7 cells were infected with culture supernatants of cells harvested 72 h posttransfection, and infectious virus titers were determined by TCID₅₀ assay at 72 h after infection. Figure 3D shows that release of infectious virus particles from cells transfected with the genome of CL3B/SA or Δ2384–2433 mutants was markedly reduced (about 10,000-fold) compared to that from wild-type- or CL3A/SA-transfected cells (white bars). To examine whether such a decrease in infectious HCV in the culture supernatants was attributable to defective virion assembly or impaired release of virions, we determined cell-associated infectivity (Fig. 3D). Production of

intracellular infectious virions in CL3B/SA- and Δ2384–2433-transfected cells was strongly impaired in comparison with that in wild-type-transfected (~1,000-fold) and CL3A/SA-transfected (~100-fold) cells. Thus, the results suggest a potential role for the serine cluster at aa 2428, 2430, and 2433 of NS5A in assembly of infectious HCV particles. Among the NS5A mutations tested, CL3B/SA is of particular interest because this mutation leads to a marked reduction in HCV production with no impact on viral RNA replication.

Serine residues at aa 2428, 2430, and 2433 are important for the interaction between NS5A and core protein. Miyazari et al. reported that the association of core protein with NS proteins is critical for infectious HCV production and that mutations of the core protein and NS5A that cause these proteins to fail to associate with each other impair the production of infectious virus (33). Based on these observations and the findings noted above, we hypothesize that NS5A plays a key role in recruiting viral RNA, which is synthesized at the viral replication complex, to nucleocapsid formation via interaction between the NS5A C-terminal region and the core protein. To prove this, we analyzed the interaction of NS5A with the core protein by coimmunoprecipitation experiments. HA-tagged NS5A constructs carrying defined mutations were generated (Fig. 1) and coexpressed with the FLAG-tagged core protein in Huh-7 cells. As shown in Fig. 4A, coimmunoprecipitation of NS5A




FIG. 2. Mutations at the C terminus of NS5A impair basal phosphorylation and have only a minor impact on RNA replication. (A) Replication of given mutants in transfected Huh-7 cells as determined by luciferase reporter assays performed at 24, 48, and 72 h posttransfection (white, gray, and black bars, respectively). Values given were normalized for transfection efficiency using the luciferase activity determined 4 h after transfection, which was set to 1. Mean values of quadruplicate measurements and the standard deviations are given. (B) Phosphorylation analysis of NS5A using the vaccinia virus T7 hybrid system. NS3-to-NS5B polypeptide fragments carrying the mutations specified above the lanes were transfected into Huh-7 cells, and proteins were radiolabeled with [³²P]orthophosphate or [³⁵S]methionine and [³⁵S]cysteine. NS5A proteins were isolated by IP and separated by SDS-PAGE (10% polyacrylamide). Mock-transfected cells served as a negative control (lanes 9 and 10). Half of the samples were treated with λ protein phosphatase (λ-PPase) (+) whereas the other half was mock treated (-) prior to SDS-PAGE. Arrows and asterisks indicate hyperphosphorylated and basally phosphorylated forms, respectively.

with the core protein was observed in cells expressing the wild-type NS5A and the CL3A/SA-mutated NS5A, but the amount of immunoprecipitated NS5A in the CL3A/SA-expressing cells was slightly lower than that in the wild-type-expressing cells. In contrast, the CL3B/SA- or the Δ2384-2433-mutated NS5A coimmunoprecipitated with the core protein only slightly or not at all.

We further examined the interaction of NS5A with core protein in cells expressing HCV genomes. At 72 h posttransfection with the wild type or CL3B/SA, cells were harvested and immunoprecipitated with an anti-NS5A antibody or an anti-C/EBPβ antibody as a negative control, followed by immunoblotting. Under these experimental conditions, the amount of extracellular core protein released from cells transfected with the CL3B/SA genome was about 10-fold lower than

that for the wild type, although comparable amounts of intracellular core protein were observed in both transfectants (Fig. 4B, left panels). As shown in the right panels of Fig. 4B, the core protein was specifically coimmunoprecipitated with NS5A in cells expressing the wild-type JFH-1 genome but not with the mutated NS5A in cells expressing the CL3B/SA genome. These results demonstrate that NS5A interacts with the core protein in cells producing infectious particles and that serine residues at aa 2428, 2430, and 2433 are important to the success of this interaction.

Two serine residues among aa 2428, 2430, and 2433 are responsible for regulating the interaction of NS5A with the core protein as well as HCV particle production. To further determine the critical residues in the C-terminal serine cluster of NS5A responsible for HCV particle production, we replaced one or two serine residues in the region with alanine (Fig. 1) and investigated which serine-to-alanine substitution influenced HCV particle production. Core protein levels in cells transfected with any construct were comparable over 4 days after transfection, indicating similar efficiencies of transfection and RNA replication from each construct (data not shown). As shown in Fig. 5A, we observed a slight delay in the kinetics of core protein release from cells transfected with the single-substitution genomes, S2428A, S2430A, and S2433A, up to 48 or 72 h posttransfection. However, core protein release from these cells reached comparable levels to that for the wild type at 96 h after transfection. In the cases of the double-substitution mutants (Fig. 5B), core protein release from cells transfected with the double-substitution genomes was markedly reduced, with 10- to 30-fold decreases compared to that for wild type observed. The kinetics of core protein release were similar to that for CL3B/SA.

Interaction of NS5A carrying single or double serine-to-alanine substitutions with the core protein was investigated by coimmunoprecipitation analysis using HA-tagged NS5A constructs. NS5A mutants carrying a single substitution were coimmunoprecipitated with the core protein (Fig. 5C), while none of the double-substitution NS5A mutants or the triple-substitution mutant, CL3B/SA, coimmunoprecipitated with the core protein (Fig. 5D). These results suggest that at least two serine residues in the C-terminal serine cluster of NS5A (aa 2428, 2430, and 2433) are necessary for the interaction between NS5A and the core protein as well as for regulation of HCV particle production and that there is positive correlation between their interaction and the amount of core protein released.

Glutamic acid partially substitutes for serine phosphorylation in the interaction of NS5A with the core protein and virus production. A consequence of phosphorylation is the addition of negative charge to a protein. In some cases, phosphoserine can be mimicked by glutamic or aspartic acid (14). To determine whether the introduction of negative charges into aa 2428, 2430, and 2433 instead of phosphoserines positively regulates the interaction of NS5A with the core protein and virus production, we replaced the serine residues with glutamic acid residues and constructed the CL3B/SE and S2428/2430E mutants (Fig. 6A). Cells transfected with the double-glutamic acid substitution, S2428/2430E, exhibited similar kinetics to the wild-type-transfected cells and released ~22-fold more core protein than S2428/2430A-transfected cells by 96 h posttransfection (Fig. 6B). In contrast,




FIG. 3. Effect of mutations introduced into the NS5A C terminus on the production of infectious HCV particles. (A) Intracellular levels of core protein measured at various time points after transfection. A total of 3×10^6 Huh-7 cells were transfected with $10 \mu\text{g}$ of in vitro-transcribed HCV RNAs specified in the inset and resuspended in 10 ml of culture medium, after which 2-ml aliquots were seeded into each well of a six-well culture plate. The cells were harvested at different time points between 4 h and 96 h posttransfection, and then 500 μl of cell lysate per well was prepared. After centrifugation, supernatants were processed for a core protein-specific ELISA. (B) Release of core protein from cells transfected with the HCV genomes specified in the inset. Cell culture supernatants harvested from cells given in panel A were analyzed by a core protein ELISA. (C) Efficiency of core protein release from cells transfected with the HCV genomes specified in the inset. The percent core protein release (vertical axis) indicates the percentage of released core protein in relation to total core protein (the sum of intra- and extracellular core protein) calculated for each time point. (D) Infectivity of virus particles contained in supernatants and cells after transfection with mutants specified below the graph. Culture supernatants and cells were harvested 72 h posttransfection, and extracellular (white bars) and intracellular infectivity (gray bars) levels were determined by TCID₅₀ assay. The gray line and arrowhead represent the detection limit of the limiting dilution assay. Mean values and standard deviations for at least triplicates are shown in all panels.

the transfectant with the triple glutamic acid substitution, CL3B/SE, showed similar trends to that of CL3B/SA. In the coimmunoprecipitation experiments with FLAG-tagged core protein and HA-tagged NS5A constructs (Fig. 6C), S2428/2430E, but not S2428/2430A, restored the ability of NS5A to interact with the core protein up to a similar level to that of wild type. As expected, neither CL3B/SE nor CL3B/SA coimmunoprecipitated with the core protein. Taken together, these results indicate that negative charges at aa 2428 and 2430 preserve the ability of NS5A to interact with the core protein and positively regulate virus production. However, the data of the CL3B/SE mutant indicate that it is likely that negative charges alone are not sufficient to enhance either the interaction of NS5A with the core protein or virus production.

Subcellular localization of NS5A and core protein in Huh-7 cells expressing HCV genomes. The coimmunoprecipitation experiments described above indicate that the wild-type NS5A but not the CL3B/SA mutant interacts with the core protein. To evaluate the NS5A-core protein interaction in intact cells, we examined the subcellular localization of NS5A with the core protein by immunofluorescence analysis. NS5A colocalized with the core protein in cells transfected with the JFH-1 wild type (Fig. 7A), whereas their colocalization was rarely observed in cells transfected with the CL3B/SA RNA (Fig. 7B).

To further analyze the subcellular compartments for the localization of NS5A and core protein in cytoplasmic membrane structures, including the endoplasmic reticulum (ER) and LDs, we performed subcellular fractionation studies as