

**Table 3**  
Undetectable rate of HBV-DNA by Amplicor monitor assay in HBeAg-negative and -positive patients

HBV-DNA (log copies/ml)	Baseline	6 months	12 months	18 months	24 months
<i>HBeAg-negative</i>					
<2.6	0 (0%)	40 (75%)	50 (94%)	47 (100%)	30 (100%)
2.6–4.5	3 (6%)	13 (25%)	3 (6%)	0 (0%)	0 (0%)
≥4.5	50 (94%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Total, n (%)	53 (100%)	53 (100%)	53 (100%)	47 (100%)	30 (100%)
<i>HBeAg-positive</i>					
<2.6	0 (0%)	32 (40%)	37 (47%)	34 (54%)	34 (68%)
2.6–4.5	3 (4%)	26 (33%)	29 (37%)	21 (35%)	15 (30%)
≥4.5	76 (96%)	21 (27%)	13 (16%)	8 (13%)	1 (2%)
Total, n (%)	79 (100%)	79 (100%)	79 (100%)	63 (100%)	50 (100%)

ADV + LAM therapy, the HBV-DNA level diminished by 1.5 log copies/ml and ALT level improved to the normal range. At that time, only the mutant strain (rtA181T) was detected, suggesting that the viral reduction was due to the suppression of wild-type HBV strain. The HBV-DNA level was persistently above 5 log copies/ml even at the end of 1 year of ADV + LAM therapy. On the other hand, in the patients with rtA181T + rtM204I mutation, viral load rapidly decreased to the undetectable HBV-DNA level at the end of 6 months of ADV + LAM combination therapy (Fig. 2c).

### 3.6. Clinical course and clonal analysis in patients who developed ADV-related mutation during combination therapy

Fig. 3 shows the clinical course of patients with ADV-resistant mutants. The first ADV-resistant HBV strain was isolated from a 32-year-old Japanese man with genotype C (Fig. 3a). At 15 months after the start of LAM, viral and biochemical breakthroughs were observed. To suppress the viral HBV-DNA, ADV was added to LAM therapy. The mutant strain with rtA181T associated with ADV resistance appeared at 6 months of ADV + LAM therapy, while another rtN236T mutation appeared at 3 years of ADV therapy. Moreover, breakthrough hepatitis was observed after 3.5 years of ADV + LAM therapy (Table 4). Interestingly, the rtA181T at the end of 6 months of ADV therapy, due to single nucleotide substitution (TGG to TGA), resulted in early termination of overlapping HBs gene by creating a stop codon. On the other hand, at the end of 3 years of ADV therapy, all rtA181T mutant strains changed to double nucleotide substitutions (TGG to TTA), which induced amino acid substitutions in both polymerase (rtA181T) and HBs antigen (HBs W172L) developed.

Another mutant strain was detected in a 38-year-old Japanese man with genotype C-HBV infection (Fig. 3b). Following 46 months of ADV + LAM ther-

apy when the viral load was increased, the rtA181S mutant strain without YMDD mutation was detected; however, the viral load diminished naturally to an undetectable level in a few months.

### 3.7. Clinical events

After the addition of ADV, 4 of the 132 (3%) patients elevated in serum creatinine >0.5 mg/dl above baseline and their ADV dose was reduced to 10 mg every other day.

Eight patients developed hepatocellular carcinoma (HCC) before the addition of ADV. After the addition of ADV, four patients developed HCC. Three of the four patients (75%) had cirrhosis at the start of ADV. The median duration from the start of ADV to the development of HCC was 14 months (range, 6–26 months). At the diagnosis of HCC, 3 of the 4 patients (75%) had undetectable HBV-DNA.

Of the 41 patients with cirrhosis, 5 patients had ascites and/or pleural effusion at the start of ADV. In 4 of the 5 patients, the fluid level diminished and disappeared during combination therapy. Only one patient with HCC showed worsened liver failure and died 22 months later. All patients without HCC and decompensation at the start of ADV therapy did not develop liver decompensation during follow-up.

## 4. Discussion

The efficacy of ADV combined with LAM has been reported in some studies; however, the rate of HBV-DNA undetectability under combination therapy was found to be the same as in patients treated with ADV alone [14,19]. We investigated whether combination therapy is characterized by a low risk of ADV resistance. In this study, we studied the long-term efficacy of ADV when added to LAM in 132 patients with chronic hepatitis B who developed LAM resistance. The results demonstrated that combination therapy




Fig. 2. Clinical course of three patients who showed emergence of ADV-resistant mutants at the commencement of ADV therapy. Amino acid substitutions in the HBV polymerase are displayed in the panel above each graph. (a) Clinical course of a patient who developed the rtA181S mutant. (b) Clinical course of a patient who developed the rtA181T mutant. (c) Clinical course of a patient who developed the rtA181T with rtM204I mutant. LAM, lamivudine; ADV, adefovir; rt, reverse transcriptase; \* no data.




Fig. 3. Clinical course of two patients with LAM-resistant HBV who showed the emergence of an ADV-resistant mutant during combination therapy. Amino acid substitutions in the HBV polymerase are displayed in the panel above the graph. (a) Clinical course of a patient who developed the rtA181T + rtN236T mutant. (b) Clinical course of a patient who developed the rtA181S mutant. LAM, lamivudine; ADV, adefovir; rt, reverse transcriptase.

rapidly and consistently suppressed the HBV-DNA. Moreover, we demonstrated that the emergence of ADV-resistant mutants was rare during the combination therapy for up to 3 years. However, our virological analysis showed that substitutions at rt181, which were associated with both LAM and ADV resistance, need to be evaluated during combination therapy.

Multivariate analysis in this study revealed that the baseline HBeAg status, AST level and HBV-DNA level influenced the cumulative probability of undetectability of serum HBV-DNA. A number of previous studies also identified almost the same predictors of virological response during ADV alone or combination therapy [20,21]. In particular, the undetectable rate of HBV-

**Table 4**  
Clonal analysis of samples from the patient who developed resistance to ADV + LAM combination therapy

	Relative rate (%) of clones (No. of clones/total)				
	Wild	rtM204I/V	rtA181T(1)	rtA181T(2)	rtA181T + rtN236T
rtA181	–	–	T(HBsAgstop)*	T(sW172L)#	T(sW172L)#
rtM204	–	I/V	–	–	–
rtN236	–	–	–	–	N
(1) At the start of ADV + LAM	40 (4/10)	60 (6/10)	0	0	0
(2) 6 months after ADV + LAM	0	59 (13/22)	41 (9/22)	0	0
(3) 2 years after ADV + LAM	16 (4/25)	36 (9/25)	36 (9/25)	12 (3/25)	0
(4) 3 years after ADV + LAM	0	0	0	0	100 (20/20)

Note. rtM204I, methionine to isoleucine substitution at rt204; rtM204V, methionine to valine substitution at rt204; rtA181T, alanine to threonine substitution at rt181.

\* The single nucleotide substitution (TGG to TGA) resulted in rtA181T mutation and early termination of overlapping HBs gene by creating a stop codon.

# The double nucleotide substitution (TGG to TTA) resulted in amino acid substitutions in both polymerase (rtA181T) and HBs antigen (HBs W172L).

DNA was more frequent and faster in HBeAg-negative patients than in HBeAg-positive patients. At the end of 12 months of combination therapy, the rates of undetectability of HBV in HBeAg-negative and HBeAg-positive patients were 94% and 47%, respectively. However, in HBeAg-positive patients, the longer treatment course increased the virological response more frequently. Some patients achieved virological response after 2 more years of combination therapy (Fig. 1a). It was considered that the low risk of ADV resistance during combination therapy contributed to the longer effect of HBV suppression.

Our result is in agreement with the previous study that showed that ADV-resistant mutants are infrequent in combination therapy [14,19]; however, our study demonstrated that the ADV mutant could have emerged during combination therapy. We identified the emergence of rtA181T/S and/or rtN236T mutation in two of the 129 patients with YMDD mutant as an ADV-resistant strain during the ADV + LAM combination therapy. To our knowledge, this is the first report of emergence of ADV-resistant mutant followed by breakthrough hepatitis during combination therapy as shown in Fig. 3a. A previous open-label study in HBeAg-negative LAM-resistant patients demonstrated that combination therapy did not result in the development of resistance to ADV over a period of 3 years, in contrast to ADV monotherapy that was associated with the development of such resistance in 21% of the patients after the first year [14]. Although another recent study reported the appearance of ADV-resistant mutants in three patients during combination therapy, they were all initially switched to ADV monotherapy and later changed to the combination therapy after several months [20]. Another recent study reported an emergence rate of ADV resistance during ADV + LAM of 1% at 1 year and 4% at 3 years; however, no virological rebound was noted [21]. The patients in our study continued to show a viral load of up to 5.8 log copies/ml

and developed breakthrough hepatitis. Some studies of ADV monotherapy reported that the rise in ALT after the emergence of rtA181 and/or rtN236 mutant is mild to moderate [9,14,24]. Several *in vitro* studies including our previous study [22–25] demonstrated that the rtA181T and rtN236T mutant leads to a minor reduction in the susceptibility to both LAM and ADV. However, one study of 998 naïve patients treated with ADV showed that the rtA181V + rtN236T mutation was significantly associated with virological breakthrough [26]. In our study, a similar phenomenon emerged; patients with ADV resistance developed breakthrough hepatitis after rtN236T mutation that appeared after rtA181T mutation.

Interestingly, clonal analyses of HBV in patients with ADV-resistant mutants in this study showed that such mutants were mixed with rtA181T mutants without substitutions at rt204, and rtM204I/V mutants without substitution at rt181. Moreover, we identified two types of rtA181T mutant strains; one was a single nucleotide substitution that induced prematurely terminated HBsAg and the other was a double nucleotide substitution that induced amino acid substitutions in the HBs antigen. The rtA181T mutant with prematurely terminated HBsAg cannot replicate and spread by itself because of the lack of HBs antigen. This type of strain is thought to replicate *in vivo* supplied HBs antigen from wild-type strains as helpers. Thus, the mutants changed themselves to the HBV with mature HBsAg by additional nucleotide substitution. Our previous study identified the rtA181T with mature HBsAg first; however, the mutant emerged during LAM therapy and it did not show a stepwise process [25].

We also demonstrated that the substitution at rt181 was associated with not only ADV resistance but also LAM resistance. At the commencement of ADV + LAM combination therapy, the substitutions at rtA181 as LAM resistance were identified in three patients (2.3%), who exhibited poor viral reduction

during the combination therapy. Of note, the rtA181S mutation is a novel LAM-resistant strain that has never been reported. There are a few reports of the rtA181 mutation associated with LAM resistance. A recent study reported the presence of rtA181T mutants in 3 of 57 (5.3%) LAM-resistant patients [15] and another study showed that 6 of 145 (4%) LAM-resistant patients developed rtA181T/V mutation [21]. If ADV therapy produces insufficient reduction of LAM-resistant HBV, it is important to suspect the emergence of ADV-related mutant at the commencement of ADV therapy and plan a new treatment strategy. However, there is no consensus at present on the management of patients with ADV + LAM-resistant mutant. Entecavir was the only agent reported to be effective both *in vitro* and *in vivo*. In our study, the patient with rtA181S mutation was switched to entecavir therapy; however, this did not produce a sufficient reduction in the viral load. On the other hand, recent studies reported the efficacy of tenofovir for patients with LAM-resistant mutants [27,28]. Further studies are needed to clear this issue.

In conclusion, ADV in combination with LAM effectively suppressed viral replication and was efficacious in LAM-resistant patients with chronic HBV infection. Genotypic analysis indicated that the emergence of ADV-resistant mutants was rare in patients on ADV + LAM combination therapy at least for 2 years. However, virological analysis showed that the substitution at rt181, which was associated with both LAM and ADV resistance, was needed for careful monitoring before and during combination therapy.

#### Acknowledgement

This work was supported in part by a Grant-in-Aid from the Ministry of Health, Labor and Welfare, Japan.

#### References

- [1] Ganem D, Prince AM. Hepatitis B virus infection—natural history and clinical consequences. *N Engl J Med* 2004;350:1118–1129.
- [2] Wright TL, Lau JY. Clinical aspects of hepatitis B virus infection. *Lancet* 1993;342:1340–1344.
- [3] Lai CL, Leung N, Teo EK, Tong M, Wong F, Hann HW, et al. A 1-year trial of telbivudine, lamivudine, and the combination in patients with hepatitis B e antigen-positive chronic hepatitis B. *Gastroenterology* 2005;129:528–536.
- [4] Chan HLY, Heathcote EJ, Marcellin P, Lai CL, Cho M, Moon YM, et al. Treatment of hepatitis B e antigen-positive chronic hepatitis with telbivudine or adefovir: a randomized trial. *Ann Int Med* 2007;147:745–754.
- [5] Nevens F, Main J, Honkoop P, Tyrrell DL, Barber J, Sullivan MT, et al. Lamivudine therapy for chronic hepatitis B: a six-month randomized dose-ranging study. *Gastroenterology* 1997;113:1258–1263.
- [6] Lai CL, Chien RN, Leung NW, Chang TT, Guan R, Tai DI, et al. A one-year trial of lamivudine for chronic hepatitis B. Asia Hepatitis Lamivudine Study Group. *N Engl J Med* 1998;339:61–68.
- [7] Suzuki Y, Kumada H, Ikeda K, Chayama K, Arase Y, Saitoh S, et al. Histological changes in liver biopsies after one year of lamivudine treatment in patients with chronic hepatitis B infection. *J Hepatol* 1999;30:743–748.
- [8] Lai CL, Dienstag J, Schiff E, Leung NW, Atkins M, Hunt C, et al. Prevalence and clinical correlates of YMDD variants during lamivudine therapy for patients with chronic hepatitis B. *Clin Infect Dis* 2003;36:687–696.
- [9] Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, Chang TT, Kitis G, Rizzetto M, et al. Long-term therapy with adefovir dipivoxil for HBeAg-negative chronic hepatitis B for up to 5 years. *Gastroenterology* 2006;131:1743–1751.
- [10] Colonna RJ, Rose R, Baldick CJ, Levine S, Pokornowski K, Yu CF, et al. Entecavir resistance is rare in nucleoside naive patients with hepatitis B. *Hepatology* 2006;44:1656–1665.
- [11] Allen MI, Deslauriers M, Andrews CW, Tipples GA, Walters KA, Tyrrell DL, et al. Identification and characterization of mutations in hepatitis B virus resistant to lamivudine. Lamivudine Clinical Investigation Group. *Hepatology* 1998;27:1670–1677.
- [12] Delaney WE, Yang H, Westland CE, Das K, Arnold E, Gibbs CS, et al. The hepatitis B virus polymerase mutation rtV173L is selected during lamivudine therapy and enhances viral replication *in vitro*. *J Virol* 2003;77:11833–11841.
- [13] Sherman M, Yurdaydin C, Sollano J, Silva M, Liaw YF, Cianciara J, et al. Entecavir for treatment of lamivudine-refractory, HBeAg-positive chronic hepatitis B. *Gastroenterology* 2006;130:2039–2049.
- [14] Rapti I, Dimou E, Mitsoula P, Hadziyannis SJ. Adding-on versus switching-to adefovir therapy in lamivudine-resistant HBeAg-negative chronic hepatitis B. *Hepatology* 2007;45:307–313.
- [15] Lee YS, Suh DJ, Lim YS, Jung SW, Kim KM, Lee HC, et al. Increased risk of adefovir resistance in patients with lamivudine-resistant chronic hepatitis B after 48 weeks of adefovir dipivoxil monotherapy. *Hepatology* 2006;43:1385–1391.
- [16] Perrillo R, Hann HW, Mutimer D, Willems B, Leung N, Lee WM, et al. Adefovir dipivoxil added to ongoing lamivudine in chronic hepatitis B with YMDD mutant hepatitis B virus. *Gastroenterology* 2004;126:81–90.
- [17] Hosaka T, Suzuki F, Suzuki Y, Saitoh S, Kobayashi M, Someya T, et al. Adefovir dipivoxil for treatment of breakthrough hepatitis caused by lamivudine-resistant mutants of hepatitis B virus. *Intervirology* 2004;47:362–369.
- [18] Usuda S, Okamoto H, Iwanari H, Baba K, Tsuda F, Miyakawa Y, et al. Serological detection of hepatitis B virus genotypes by ELISA with monoclonal antibodies to type-specific epitopes in the preS2-region product. *J Virol Methods* 1999;80:97–112.
- [19] Peters MG, Hann HW, Martin P, Heathcote EJ, Buggisch P, Rubin R, et al. Adefovir dipivoxil alone or in combination with lamivudine in patients with lamivudine-resistant chronic hepatitis B. *Gastroenterology* 2004;126:91–101.
- [20] Buti M, Elefsiniotis I, Jardi R, Vargas V, Rodriguez-Frias F, Schapper M, et al. Viral genotype and baseline load predict the response to adefovir treatment in lamivudine-resistant chronic hepatitis B patients. *J Hepatol* 2007;47:366–372.
- [21] Lampertico P, Viganò M, Manenti E, Iavarone M, Sablon E, Colombo M. Low resistance to adefovir combined with lamivudine: a 3-year study of 145 lamivudine-resistant hepatitis B patients. *Gastroenterology* 2007;133:1445–1451.
- [22] Lampertico P, Viganò M, Manenti E, Iavarone M, Lunghi G, Colombo M. Adefovir rapidly suppresses hepatitis B in HBeAg-negative patients developing genotypic resistance to lamivudine. *Hepatology* 2005;42:1414–1419.
- [23] Yeh CT, Chien RN, Chu CM, Liaw YF. Clearance of the original hepatitis B virus YMDD-motif mutants with emergence of distinct lamivudine-resistant mutants during prolonged lamivudine therapy. *Hepatology* 2000;31:1318–1326.

- [24] Brunelle MN, Jacquard AC, Pichoud C, Durantel D, Carrouee-Durantel S, Villeneuve JP, et al. Susceptibility to antivirals of a human HBV strain with mutations conferring resistance to both lamivudine and adefovir. *Hepatology* 2005;41:1391–1398.
- [25] Yatsuji H, Noguchi C, Hiraga N, Mori N, Tsuge M, Imamura M, et al. Emergence of a novel lamivudine-resistant hepatitis B virus variant with a substitution outside the YMDD motif. *Antimicrob Agents Chemother* 2006;50:3867–3874.
- [26] Borroto-Esoda K, Miller MD, Arterburn S. Pooled analysis of amino acid changes in the HBV polymerase in patients from four major adefovir deipivoxil clinical trials. *J Hepatol* 2007;47:492–498.
- [27] Van Bommel F, Wunsche T, Mauss S, Reinke P, Bergk A, Schurmann D, et al. Comparison of adefovir and tenofovir in the treatment of lamivudine-resistant hepatitis B virus infection. *Hepatology* 2004;40:1421–1425.
- [28] van Bommel F, Zollner B, Sarrazin C, Spengler U, Huppe D, Moller B, et al. Tenofovir for patients with lamivudine-resistant hepatitis B virus (HBV) infection and high HBV DNA level during adefovir therapy. *Hepatology* 2006;44:318–325.

# Hepatocarcinogenesis Following HCV RNA Eradication by Interferon in Chronic Hepatitis Patients

Miharu Hirakawa, Kenji Ikeda, Yasuji Arase, Yusuke Kawamura, Hiromi Yatsuji,  
Tetsuya Hosaka, Hitomi Sezaki, Norio Akuta, Masahiro Kobayashi, Satoshi Saitoh,  
Fumitaka Suzuki, Yoshiyuki Suzuki and Hiromitsu Kumada



INTERNAL MEDICINE

*Reprinted from Internal Medicine*

Vol. 47, Pages 1637-1643

October 2008

## Hepatocarcinogenesis Following HCV RNA Eradication by Interferon in Chronic Hepatitis Patients

Miharu Hirakawa, Kenji Ikeda, Yasuji Arase, Yusuke Kawamura, Hiromi Yatsuji, Tetsuya Hosaka, Hitomi Sezaki, Norio Akuta, Masahiro Kobayashi, Satoshi Saitoh, Fumitaka Suzuki, Yoshiyuki Suzuki and Hiromitsu Kumada

### Abstract

**Objective** Interferon (IFN) therapy reduces the incidence of hepatocarcinogenesis in patients with hepatitis C viral (HCV) infection who achieve a sustained virological response (SVR). The aim of the present study was to determine the rate of hepatocarcinogenesis and the risk factor in sustained virological responders.

**Patients and Method** The study subjects were 1,193 patients with HCV-related chronic liver disease and IFN- or IFN plus ribavirin-induced SVR. The age, male/female ratio, and liver fibrosis stage [(F0-F3)/LC] were 15-83 years, 808/385, and 1106/41, respectively. Patients were followed-up for 8.3 years (range, 0 to 19.0 years) and the incidence of hepatocellular carcinoma was recorded.

**Results** Hepatocellular carcinogenesis was detected in 23 patients during the follow-up. The crude rates of hepatocarcinogenesis at 5, 10, and 15 years were 1.5%, 2.4% and 4.1%, respectively. Multivariate analysis identified cirrhosis, male sex and age older than 50 years as determinants of hepatocarcinogenesis with hazard ratios of 12.9 ( $p < 0.001$ ), 6.45 ( $p = 0.012$ ), and 20.2 ( $p = 0.004$ ), respectively.

**Conclusion** Long-term follow-up of patients with chronic HCV infection is necessary even in those who show SVR, especially in male elderly patients with cirrhosis.

**Key words:** hepatitis C virus, hepatocellular carcinoma, chronic hepatitis, sustained virological response, cox proportional hazard model

(*Inter Med* 47: 1637-1643, 2008)

(DOI: 10.2169/internalmedicine.47.1087)

### Introduction

Interferon (IFN) is effective in eliminating HCV and reducing serum alanine aminotransferase (ALT) in some patients with chronic hepatitis C viral (HCV) infection (1-3). A reduction in the incidence of hepatocellular carcinoma (HCC) in patients with HCV-associated hepatitis and cirrhosis treated with IFN has been reported by many investigators (4-14). The previous study (14) suggested that fluctuations and persistently high levels of ALT in patients with chronic HCV infection enhances the carcinogenic process. From the viewpoint of liver carcinogenesis, IFN plays a suppressive action on the development of HCC through reduction or complete remission of inflammatory activity. Multivariate

analysis has indicated that IFN lowers the carcinogenesis rate in those patients who show IFN-induced reduction in ALT levels (15). In patients with IFN-induced normalization of ALT levels, the groups at high risk for carcinogenesis were older, male, and a more advanced histologic stage (16). Patients who show elimination of HCV RNA are considered to exhibit normalization of ALT levels. Therefore, the incidence of carcinogenesis is assumed to be lower in patients with sustained virological response (SVR) than in those who show biochemical response (BR) and no response (NR) to IFN therapy. SVR was defined as persistent disappearance of HCV RNA after therapy, BR as normal ALT values without elimination of HCV RNA for at least 6 months after therapy, and NR as persistently abnormal or only transient normalization of ALT for less than 6 months. However, he-

Department of Hepatology, Toranomon Hospital, Tokyo

Received for publication March 6, 2008; Accepted for publication May 21, 2008

Correspondence to Dr. Miharu Hirakawa, ZXC00701@nifty.ne.jp

patocarcinogenesis still occurs in patients with SVR (17-31). In the studies of Toyoda et al (17) and Ikeda et al (18), the risk factors for carcinogenesis were not discussed due to few sustained virological responders with carcinogenesis. Tokita et al (19) and Kobayashi et al (20) indicated that the risk factors of hepatocarcinogenesis after elimination of HCV RNA are severe fibrosis, male sex, and regular consumption of moderate amounts of alcohol, and old age at the start of IFN treatment. Their hazard ratios could not be estimated because of the relatively small number of patients with SVR. Ikeda et al (21) indicated the hazard ratios of risk factors; older age, increased aspartate aminotransferase (AST), and decreased platelet count. However, the study population was restricted to patients who received IFN monotherapy and it did not include patients who received either pegylated interferon (PEG IFN) or combination therapy of IFN and ribavirin.

The aims of this study were to estimate the rate of hepatocellular carcinogenesis in patients with chronic HCV infection who show SVR to IFN monotherapy or combination therapy of IFN and ribavirin and to determine the risk factors that affect carcinogenesis rate in such patients using multivariate analysis.

## Patients and Methods

### Study population

In this retrospective cohort study, all patients with chronic HCV infection who started IFN therapy between February 1987 and July 2006 in the Department of Hepatology, Toranomon Hospital were analyzed in the database. Prior to IFN therapy, they were positive for anti-HCV (second- or third generation anti-HCV, enzyme-linked immunosorbent assay, Dainabot, Osaka, Japan) and HCV RNA. Anti-HCV was assayed using stored frozen sera at -80°C. HCV RNA was determined by the Amplicor method (Cobas Amplicor HCV Monitor Test, v2.0, Roche Molecular Systems, Inc., Belleville, NJ) or the branched DNA probe assay (b DNA probe assay; version 2.0; Chiron, Dai-ichi Kagaku, Tokyo, Japan). The medical records of 1,193 patients with HCV infection, who had achieved HCV RNA elimination after IFN therapy or the combination therapy of IFN and ribavirin were obtained. The sera of all patients were negative for hepatitis B surface antigen (HBsAg; radioimmunoassay, Austria, Abbott Laboratories, Detroit, MI). The study protocol was approved by the Human Ethics Review Committee of Toranomon hospital.

### Clinical background and laboratory data

The background of 1,193 patients who achieved SVR is shown in Table 1. They included 809 men and 384 women, who were 15 to 83 years old with a median age of 50 years at the commencement of therapy. HCV genotype was analyzed by the immunoserological typing method using a commercial kit (Kokusai Diagnostic Corporation, Kobe, Japan).

Table 1. Patients' Profiles, Virological, Histological Characteristics of the Patients Prior to Their Interferon (IFN) Therapy and Protocol of IFN Therapy

Number of patients	1193
Sex (M / F)	809 / 384
Age (years) *	50 (15-83)
Observation period (year) *	8.3 (0.0-19.0)
HCV genotype	
Genotype 1a, 1b	494 (41.4%)
Genotype 2a, 2b	670 (56.2%)
Genotype 1+2	5 (0.4%)
Genotype 3	1 (0.1%)
Undetermined	23 (1.9%)
Histological stage of hepatitis	
F0 (no fibrosis)	7 (0.6%)
F1 (slight fibrosis)	738 (61.9%)
F2 (moderate fibrosis)	289(24.2%)
F3 (severe fibrosis)	72 (6.0%)
F4 (cirrhosis)	41 (3.4%)
Not examined	46 (3.9%)
IFN therapy	
Monotherapy	1032 (86.5%)
Combination therapy with ribavirin	161 (13.5%)
Type of IFN	
IFN- $\alpha$	850 (71.2%)
PEG IFN- $\alpha$	46 (3.9%)
IFN- $\beta$	251 (21.0%)
IFN- $\alpha$ /PEG IFN- $\alpha$ +IFN- $\beta$	47 (3.9%)

IFN: interferon, PEG IFN: pegylated interferon

\*Data are median (minimum, maximum) values.

The HCV genotype was 1 (genotype 1a and 1b) in 494 patients, 2 (genotype 2a and 2b) in 670 patients, 1 plus 2 in 5 patients, 3 in 1 patient. Before treatment, 1,131 patients underwent liver biopsy with or without peritoneoscopy to assess the staging of liver fibrosis and the grade of inflammatory activity based on the classification of Desmet (32). Staging of liver fibrosis was defined as F0 (no fibrosis), F1 (fibrosis portal expansion), F2 (bridging fibrosis), F3 (bridging fibrosis with architectural distortion) and F4 (cirrhosis). Additionally, 16 patients were diagnosed as cirrhosis by peritoneoscopy without biopsy, laboratory values or clinical features: 1147 patients were diagnosed with chronic hepatitis (n=1,106) and cirrhosis (n=41) (F0/F1/F2/F3/F4=7/738/289/72/41).



### Treatment protocol

IFN was performed once in 973 patients and more times of therapy in 220 patients (twice/three times/four times/five times/six times=166/38/13/2/1). IFN and ribavirin combination therapy was used to eliminate HCV RNA for 161 patients, while IFN monotherapy eliminated HCV RNA for the other 1,032 patients. The type of IFN was IFN- $\alpha$  (natural or recombinant)/PEG IFN- $\alpha$  in 896 patients (75.1%); IFN- $\alpha$  in 850 patients (71.2%), PEG IFN- $\alpha$  in 46 patients (3.9%), IFN- $\beta$  (natural) in 251 patients (21.0%) and IFN- $\alpha$  or PEG IFN- $\alpha$  and IFN- $\beta$  in 47 patients (3.9%).

A total of 613 patients (51.4%) received 3 to 9 million units of IFN everyday for 8 weeks followed by twice or three times a week for 1 to 305 weeks (for 16 to 22 weeks in 75% of patients), 304 patients (25.5%) received 3 to 9 million units of IFN everyday for 1-5 weeks followed by three times a week, 5 patients (0.4%) for 12 weeks and one patient (0.1%) for 24 weeks followed by intermittent administration. A total of 124 patients (10.4%) underwent short therapy with IFN everyday for 4-8 weeks, 2 patients (0.2%) for 10-12 weeks, 18 patients (1.5%) for 18-24 weeks. 2 patients (0.2%) had a prolonged administration of IFN for 11 and 13 months. And 63 patients (5.3%) underwent intermittent administration of three times a week for 4 weeks to 70 months. This protocol is one of the low-dose intermittent IFN therapies. A total of 48 patients (4.0%) underwent 50-180  $\mu$ g of PEG IFN once a week: 8 patients for 24 weeks and 40 patients for 48 weeks.

### Follow-up and diagnosis of hepatocellular carcinoma

Almost all patients were followed-up every week or bi-weekly during IFN monotherapy. This included hematological, biochemical, and virological tests. Patients treated with pegylated IFN were also checked every week or bi-weekly. After the completion of treatment, monthly follow-up was continued until the virological response could be determined. When SVR was confirmed, imaging studies were conducted once or twice per year in the majority of patients; these included computed tomography (CT) or ultrasonography (US), except those patients who were lost to follow-up. Angiography was performed only when HCC was highly suspected on CT or US. The presence of a characteristic hypervascular nodule on angiography was considered a specific finding for HCC, and histological confirmation was usually not required in the majority of such cases. The clinical trends of tumor markers were also taken into account. When angiography could not be performed, the hepatic mass was considered HCC when CT showed a hypervascular mass and the tumor marker level was elevated. No fine needle biopsy or histopathological examination was performed before treatment.

The date of the last follow-up in this study was March 1, 2007. The median observation period of the entire group was 8.3 years with a range of 0.0 to 19.0 years. As for pa-

tients of the combination therapy, the median follow-up period was 3.2 years with a range of 0.0 to 7.5 years.

### Statistical analysis

Non-parametric procedures were employed for the analysis of background clinicopathological parameters, including Mann-Whitney U-test. The rate of hepatocarcinogenesis was calculated for the period between the end of IFN therapy and appearance of HCC, using Kaplan-Meier technique (33). Differences in carcinogenesis curves were tested using the log-rank test. Independent factors associated with the appearance of HCC were studied using stepwise Cox regression analysis (34). The following seven variables were analyzed for potential covariates for liver carcinogenesis; age, sex, fibrotic stage of hepatitis at the initiation of the IFN therapy, HCV genotype, use of ribavirin (monotherapy or combination therapy), type of IFN ( $\alpha$  or  $\beta$ ), and number of treatments. Factors found significant were entered into a multivariate Cox proportional hazard model. A P-value less than 0.05 was considered significant. Data were analyzed using the SPSS software ver. 11.0.1J (SPSS Inc., Chicago, IL).

### Crude rates of hepatocarcinogenesis

During a median observation period of 8.3 years with a range of 0.0 to 19.0 years, HCC was diagnosed in 23 (1.9%) of the 1,193 patients. The median interval between the end of therapy and detection of HCC was 3.1 years (range, 0-12.9 years).

The characteristics of HCC patients are shown in Table 2. Patients who developed HCC before the initiation of IFN therapy were excluded. Four patients (Nos. 2, 5, 15 and 22) developed HCC before the diagnosis of SVR but after the elimination of HCV RNA. The surgically resected liver tissue was also examined by the PCR method in 4 cases (Nos. 1, 6, 13 and 14), which showed no HCV RNA.

HCC patients included 21 men and 2 women; the median age at the start of IFN therapy and at diagnosis of HCC was 58 (range, 50-70) and 62 (51-76), respectively. The HCV genotype was 1 in 9 patients and 2 in 14 patients. Chronic hepatitis was diagnosed in 14 patients (F1/F2/F3=2/12/0) and cirrhosis in 9, at the time of initiation of IFN therapy. The type of therapy for hepatitis was IFN monotherapy in 22 patients and the combination therapy in 1. The number of HCC tumors was one in 18 patients, two in 3 patients and more than two in 2 patients. A typical hypervascular mass on angiography or perfusion defect on CT during arterial portography (CT-AP) was noted in 20 patients. Angiography could not be performed in the other three patients; they had a hypo-enhanced, iso enhanced, and hyper-enhanced tumor on CT, respectively. Treatment was radical in 21 patients; including hepatectomy in 17 and percutaneous locoregional therapy in 4 patients. At the time of surgical resection, the fibrosis staging was histopathologically examined in 16 patients. Twelve patients was diagnosed as hepatitis (F1/F1-2/F2/F3=2/1/6/3) and 4 as cirrhosis. In

Table 2. Carcinogenesis after HCV RNA Elimination

No	Gender	Age at the start of IFN	Age at the time of carcinogenesis	Genotype	Type of IFN	Fibrosis staging before IFN Tx	Interval between the end of IFN Tx and carcinogenesis, yr	Number of Tumor	Tumor size, mm	Treatment for HCC	Fibrosis staging at the time of carcinogenesis	Differentiation of HCC
1	M	50	51	1b	$\alpha$	F2	1.0	1	15	Hepatectomy	F3	Moderate
2	M	52	54	2a	$\alpha$	F1	0.6	1	18	Hepatectomy	F2	Well
3	F	54	59	2a	$\alpha$	F2	3.5	1	17	Hepatectomy	F1-2	Moderate
4	M	55	60	1b	$\alpha$	F2	3.7	1	16	Hepatectomy	F2	Moderate
5	M	55	56	1b	Peg $\alpha$ +Rib	F2	0.0	1	21	RFA	-	-
6	M	55	57	2a	$\alpha$ 2a	F4	1.9	1	19	Hepatectomy	F4	Moderate
7	M	57	67	2	$\alpha$	F2	8.9	1	47	Hepatectomy	F1	Moderate
8	M	55	59	2a	$\alpha$ 2a	F4	3.1	1	18	Hepatectomy	F4	Moderate
9	M	55	62	1b	$\beta$	F4	6.5	1	16	Hepatectomy	F4	Poor
10	M	57	58	1b	$\alpha$	F2	0.9	1	16	Hepatectomy	F2	Moderate
11	M	57	59	1b	$\alpha$	F2	1.2	1	20	Hepatectomy	F2	Moderate
12	M	58	66	2a	$\alpha$ 2b	F1	8.7	1	26	Hepatectomy	F1	Well
13	M	58	62	1b	$\beta$	F2	3.9	1	30	Hepatectomy	F2	poor/moderate
14	M	59	69	2b	$\alpha$	F2	9.1	1	21	Hepatectomy	F2	Moderate
15	M	59	61	1b	$\alpha$	F4	0.1	1	30	Hepatectomy	F3	poor/moderate
16	M	61	63	2	$\alpha$	F2	1.8	4+LN meta	23	Hepatectomy+MCT	F3	moderate>well
17	M	62	65	2a	$\beta$	F4	2.4	2	20,20	RFA	-	-
18	M	62	75	2a	$\alpha$ 2a	F4	12.9	1	23	Hepatectomy	F4	Moderate
19	M	63	66	2a	$\beta$	F2	3.6	Uncountable	Diffuse	No treatment	-	-
20	M	65	71	2a	$\alpha$	F2	5.0	2	12, 8	Hepatectomy	-	Necrosis
21	F	66	68	2a	$\alpha$	F4	0.9	1	13	RFA	-	-
22	M	69	72	1b	$\beta$	F4	0.4	2	13, 13	Hepatectomy	-	moderate, poor
23	M	70	76	2a	$\beta$	F4	5.4	1	10	RFA+PMCT	-	-

IFN: interferon, Tx: therapy, Peg: pegylated interferon, Rib: ribavirin, LN meta: lymph node metastasis, RFA: radiofrequency ablation, MCT: microwave coagulation therapy, PMCT: percutaneous microwave coagulation therapy.

Carcinogenesis rate




Figure 1. Rates of hepatocarcinogenesis in 41 patients with cirrhosis (F4) and 1,106 patients with liver fibrosis stage F0-F3.

comparison with the staging at the initiation of IFN therapy, 3 cases showed improvement in the fibrosis, 10 showed no change, and 3 showed progression.

The crude rates of hepatocarcinogenesis in the SVR patients were 1.5%, 2.4% and 2.7% at the end of the 5th year, 10th year and 15th year, respectively.

#### Determinants of hepatocarcinogenesis

The rate of carcinogenesis was significantly higher in 41 patients with cirrhosis (F4) than in 1,106 patients with liver fibrosis of F0-F3 ( $p < 0.0001$ , Fig. 1). The respective cumulative HCC development rates in patients with cirrhosis at 5, 10, and 15 years after SVR were 15.6%, 24.2% and 39.4%. On the other hand, the respective rates for patients with F0-F3 were 1.00%, 1.68% and 1.68% at 5, 10, and 15 years after SVR. When patients were divided into two groups with F2-4 and with F0-1, rates of the former group were 4.16%, 6.52% and 7.58%, while those of latter group were 0.13%,

Carcinogenesis rate




Figure 2. Rates of hepatocarcinogenesis in 809 male patients and 384 female patients.

0.40% and 0.40%. The incidence rates of HCC increases with the fibrotic stage; those with F1, F2 and F4 were 0.14%, 3.43% and 15.6% at 5 years, 0.40%, 5.22% and 24.2% at 10 years, and 0.40%, 5.22% and 39.4% at 15 years, respectively.

The rate of hepatocarcinogenesis among 809 male patients was significantly higher than among 384 female patients ( $p = 0.018$ , Fig. 2); the respective rates at 5, 10 and 15 years were 1.87%, 3.18% and 3.67% for males and 0.58%, 0.58% and 0.58% for females.

The rate of hepatocarcinogenesis among 570 patients aged  $> 50$  years was greater than among 623 patients aged  $< 51$  years at the start of IFN therapy ( $p < 0.0001$ , Fig. 3); the respective rates at 5, 10 and 15 years for the former group were 2.92%, 4.93% and 5.81%, compared with 0.16%, 0.16% and 0.16% for the latter.

Multivariate analysis identified three factors to be associated with the rate of development of HCC: sex, age at start IFN treatment, and fibrotic stage in the liver tissue. Multi-

ivariate analysis was performed using non-time dependent proportional hazard analysis. Fibrotic stage, sex, and age were identified as significant independent factors that influenced the rate of future hepatocarcinogenesis (Table 3). Cirrhosis (F4) was associated with a higher risk of hepatocarcinogenesis with a hazard ratio of 12.9 (95% confidence interval, 5.5-30.6,  $p < 0.001$ ) compared with F1-3 stage. Similarly, male sex (6.45,  $p = 0.012$ ) and older age than 50 years (20.2,  $p = 0.004$ ) were associated with a higher risk. Serological grouping of HCV, type of therapy (monotherapy or combination therapy), type of IFN of the final therapy, and number of therapies did not significantly influence the rate of hepatocarcinogenesis. When the patients were divided into two groups with F0-1 and with F2-4, hazard ratios of F2-4, male and older age were 13.4 (3.1-57.8,  $p < 0.0001$ ), 7.00 (1.63-29.99,  $p = 0.0009$ ) and 17.6 (2.3-131.6,  $p = 0.005$ ). When the patients were divided into two groups with F0-2 and F3-4, hazard ratios of F3-4, male and older age were 5.8 (2.5-13.8,  $p < 0.0001$ ), 6.78 (1.58-29.07,  $p = 0.01$ ) and 22.9 (3.0-172.2,  $p = 0.002$ ).

#### Carcinogenesis rate




Figure 3. Rates of hepatocarcinogenesis in 570 patients older than 50 and 623 patients younger than 51 years.

## Discussion

Epidemiological data on the rate of development of HCC in patients with chronic hepatitis (35) and those with cirrhosis (36) indicate that the life expectancy of patients with HCV-related chronic liver disease is significantly influenced by the development of HCC. Up to 75% of patients with HCV infection and cirrhosis eventually develop HCC (15). IFN can be considered to have anti-carcinogenic properties through its anti-inflammatory action, since several studies have already described that the cancer suppressive activity of IFN in those patients who show HCV RNA eradication was similar to that of patients with ALT normalization without HCV RNA elimination (BR) (15, 37-40). After excluding patients with cirrhosis, the previous report (40) showed that the rate of carcinogenesis was lower in patients with SVR than in those with BR because HCV-elimination does not result in re-elevation and exacerbation of ALT. As a follow-up to the above studies, the rate of hepatocarcinogenesis in SVR patients with either chronic hepatitis or cirrhosis was estimated in the present study.

In spite of the anti-carcinogenic effect of SVR, 23 cases developed HCC following elimination of HCV RNA among 1,193 patients. The median interval between the end of IFN therapy and carcinogenesis is 3.1 years with a range of 0.0 to 12.9 years. Among 23 cases, 22 patients had regular examinations of at least once a year, and 21 of them received radical treatment such as hepatectomy or radiofrequency ablation. The high rate of radical treatment was probably due to the preserved liver function after HCV RNA elimination.

HCCs in six cases that were detected in the year after the end of the interferon therapy could have been already present before elimination of HCV RNA. Even when we exclude these cases, multivariate analysis identified the same factors such as higher histological stage, male sex and age older age as determinants of hepatocarcinogenesis. The haz-

Table 3. Factors Associated with Hepatocarcinogenesis in Sustained Virological Responders with Chronic HCV Infection

Factors	Category	Hazard ratio	95% confidence interval	p
Fibrotic stage	1: F0-3	1		
	2: F4	12.9	(5.5-30.6)	<0.001
Gender	1: women	1		
	2: men	6.45	(1.51-27.64)	0.012
Age (years)	1: <51	1		
	2: >50	20.2	(2.7-152.9)	0.004

ard ratios of cirrhosis, male sex and age older than 50 years were 10.9 (4.0-29.8,  $p < 0.001$ ), 10.4 (1.4-78.2,  $p = 0.024$ ) and 17.0 (2.2-130.7,  $p = 0.006$ ), respectively.

The rates of hepatocarcinogenesis in patients with histological stage F0-F3 were 1.00%, 1.68%, 1.68% at 5, 10, and 15 years after SVR, respectively. These rates were about 20% less than the rates reported previously for patients with chronic hepatitis; 4.8%, 13.6%, and 26.0%, respectively (35). The rates of hepatocarcinogenesis in patients with cirrhosis were 15.6%, 24.2%, and 39.4% at 5, 10, and 15 years, respectively, which were about 65% less than the rates reported previously for patients with cirrhosis; 21.5%, 53.2% and 75.2%, respectively (36). These results indicate that IFN has a more marked effect in reducing the rate of hepatocarcinogenesis in patients with F0-F3 than in those with cirrhosis. Furthermore, the difference in the rate of hepatocarcinogenesis in patients who show SVR and those with chronic HCV-infected patients increases with time, since the likelihood of development of HCC before elimination of HCV RNA decreases as time passes after IFN therapy.

Although some studies have reported that elderly male patients with severe fibrotic stage could be at a high risk for hepatocarcinogenesis even when they show SVR, the hazard ratios in such patients have not been reported probably be-

cause of shorter follow-up period and the relatively small number of patients. In this study, the follow-up period was longer than that of previous studies, allowing meaningful multivariate analysis (e.g., Cox hazards model). The results of such analysis showed that the risk of carcinogenesis increases with the histologic stage of the liver, age and male sex. This finding was similar to that reported in a study of untreated patients (41, 42) or IFN-treated hepatitis patients with the histological stage of F0-F3 (15).

Treatment of patients with chronic HCV infection using PEG IFN- $\alpha$  and ribavirin resulted in persistently negative tests for serum HCV RNA in 40-50% of patients with HCV genotype 1 and 75-80% with HCV genotype 2 or 3. The present study also showed that neither type of IFN ( $\alpha$  or  $\beta$ ) nor the use of ribavirin altered the rate of carcinogenesis. Further studies are needed with a longer follow-up period since the follow-up period of patients treated with the combination therapy ranged from only 0 to 7.5 years (median 3.2 years) and was shorter than that of patients who received IFN monotherapy.

In conclusion, the results emphasize the importance of long-term follow-up of patients with chronic HCV infection, even those who show SVR to IFN therapy, especially male elderly patients with severe fibrosis of the liver.


## References

- Davis GL, Balart LA, Schiff ER, et al. Treatment of chronic hepatitis C with recombinant interferon alpha. A multicenter, randomized, controlled trial. *N Engl J Med* **321**: 1501-1506, 1989.
- Di Bisceglie AM, Martin P, Kassianides C, et al. Recombinant interferon alpha therapy for chronic hepatitis C: A randomized, double-blind, placebo-controlled trial. *N Engl J Med* **321**: 1506-1510, 1989.
- Causse X, Godinot H, Chevallier M, et al. Comparison of 1 or 3 MU of interferon alfa-2b and placebo in patients with chronic non-A, non-B hepatitis. *Gastroenterology* **101**: 497-502, 1991.
- Chayama K, Saitoh S, Arase Y, et al. Effect of interferon administration on serum hepatitis C virus RNA in patients with chronic hepatitis C. *Hepatology* **13**: 1040-1043, 1991.
- Nishiguchi S, Kuroki T, Nakatani S, et al. Randomized trial of effects of interferon-alpha on incidence of hepatocellular carcinoma in chronic active hepatitis C with cirrhosis. *Lancet* **346**: 1051-1055, 1995.
- Mazzella G, Accogli E, Scotti S, et al. Alpha interferon treatment may prevent hepatocellular carcinoma in HCV-related liver cirrhosis. *J Hepatol* **24**: 141-147, 1996.
- Schalm SW, Fattovich G, Brouwer JT. Therapy of hepatitis C: patients with cirrhosis. *Hepatology* **26**: 128S-132S, 1997.
- Benvegnu L, Chemello L, Noventa F, Fattovich G, Pontisso P, Alberti A. Retrospective analysis of the effect of interferon therapy on the clinical outcome of patients with viral cirrhosis. *Cancer* **83**: 901-909, 1998.
- Niederer C, Lange S, Heintges T, Erhardt A, Buschkamp M, Hurter D. Prognosis of chronic hepatitis C: results of a large, prospective cohort study. *Hepatology* **28**: 1687-1695, 1998.
- International Interferon-alpha Hepatocellular Carcinoma Study Group. Effect of interferon-alpha on progression of cirrhosis to hepatocellular carcinoma: a retrospective cohort study. *Lancet* **351**: 1535-1539, 1998.
- Kasahara A, Hayashi N, Mochizuki K, et al. Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. Osaka Liver Disease Study Group. *Hepatology* **27**: 1394-1402, 1998.
- Shindo M, Ken A, Okuno T. Varying incidence of cirrhosis and hepatocellular carcinoma in patients with chronic hepatitis C responding differently to interferon therapy. *Cancer* **85**: 1943-1950, 1999.
- Yoshida H, Shiratori Y, Moriyama M, et al. Interferon therapy reduces the risk for hepatocellular carcinoma: national surveillance program of cirrhotic and noncirrhotic patients with chronic hepatitis C in Japan IJIT Study Group. Inhibition of hepatocellular carcinogenesis by interferon therapy. *Ann Int Med* **131**: 174-181, 1999.
- Ikeda K, Saitoh S, Kobayashi M, et al. Long-term interferon therapy for 1 year or longer in reduces the hepatocellular carcinogenesis rate in patients with liver cirrhosis caused by hepatitis C virus: A pilot study. *J Gastroenterol Hepatol* **16**: 406-415, 2001.
- Ikeda K, Saitoh S, Arase Y, et al. Effect of interferon therapy on hepatocellular carcinogenesis in patients with chronic hepatitis type C: A long-term observation study of 1,643 patients using statistical bias correction with proportional hazard analysis. *Hepatology* **29**: 1124-1130, 1999.
- Makiyama A, Itoh Y, Kasahara A, et al. Characteristics of patients with chronic hepatitis C who develop hepatocellular carcinoma after a sustained response to interferon therapy. *Cancer* **101**: 1616-1622, 2004.
- Toyoda H, Kumada T, Tokuda A, et al. Long-term follow-up of sustained responders to interferon therapy in patients with chronic hepatitis C. *J Viral Hepat* **7**: 414-419, 2000.
- Ikeda M, Fujiyama S, Tanaka M, et al. Clinical features of hepatocellular carcinoma that occur after sustained virological response to interferon for chronic hepatitis C. *J Gastroenterol Hepatol* **21**:


- 122-128, 2006.
19. Tokita H, Fukui H, Tanaka A, et al. Risk factors for the development of hepatocellular carcinoma among patients with chronic hepatitis C who achieved a sustained virological response to interferon therapy. *J Gastroenterol Hepatol* **20**: 752-758, 2005.
  20. Kobayashi S, Takeda T, Enomoto M, et al. Development of hepatocellular carcinoma in patients with chronic hepatitis C who had a sustained virological response to interferon therapy: a multicenter, retrospective cohort study of 1124 patients. *Liver Intern* **27**: 186-191, 2007.
  21. Ikeda M, Fujiyama S, Tanaka M, et al. Risk factors for development of hepatocellular carcinoma in patients with chronic hepatitis C after sustained response to interferon. *J Gastroenterol* **40**: 148-156, 2005.
  22. Enokimura N, Shiraki K, Kawakita T, et al. Hepatocellular carcinoma development in sustained viral responders to interferon therapy in patients with chronic hepatitis C. *Anticancer Res* **23**: 593-596, 2003.
  23. Yamaguchi K, Omagari K, Kinoshita H, et al. Development of hepatocellular carcinoma in a patient with chronic hepatitis C after 6 years of a sustained and complete response to IFN- $\alpha$ . *J Clin Gastroenterol* **29**: 207-209, 1999.
  24. Yamada M, Ichikawa M, Matsubara A, Ishiguro Y, Yamada M, Yokoi S. Development of small hepatocellular carcinoma 80 months after clearance of hepatitis C virus with interferon therapy. *Eur J Gastroenterol Hepatol* **12**: 1029-1032, 2000.
  25. Nagano K, Fukuda Y, Nakano I, et al. A case of the development of two hepatocellular carcinomas and a cholangiocarcinoma with cirrhosis after elimination of serum hepatitis C virus RNA with interferon therapy. *Hepatogastroenterology* **47**: 1436-1438, 2000.
  26. Yamaura T, Matsumoto A, Rokuhara A, et al. Development of small hepatocellular carcinoma in a patient with chronic hepatitis C after 77 months of a sustained and complete response to interferon therapy. *J Gastroenterol Hepatol* **17**: 1229-1235, 2002.
  27. Sugiura N, Sakai Y, Ebara M, et al. Detection of hepatocellular carcinoma after interferon therapy for chronic hepatitis C: clinical study of 26 cases. *J Gastroenterol Hepatol* **11**: 535-539, 1996.
  28. Tamori A, Kuroki T, Nishiguchi S, et al. Case of small hepatocellular carcinoma in the caudate lobe detected after interferon caused disappearance of hepatitis C virus. *Hepatogastroenterology* **43**: 1079-1083, 1996.
  29. Hirashima N, Mizokami M, Orito E, et al. Case report: development of hepatocellular carcinoma in a patient with chronic hepatitis C infection after a complete and sustained response to interferon- $\alpha$ . *J Gastroenterol Hepatol* **11**: 955-958, 1996.
  30. Miyano S, Togashi H, Shinzawa H, et al. Case report: Occurrence of hepatocellular carcinoma 4.5 years after successful treatment with virus clearance for chronic hepatitis C. *J Gastroenterol Hepatol* **14**: 928-930, 1999.
  31. Inoue M, Ohhira M, Ohtake T, et al. Hepatocellular carcinoma developed in a patient with chronic hepatitis C after the disappearance of hepatitis C virus due to interferon therapy. *Hepatogastroenterology* **46**: 2554-2560, 1999.
  32. Knodell RG, Ishak KG, Black WC, et al. Formulation and application of a numerical scoring system for assessing histological activity in asymptomatic chronic active hepatitis. *Hepatology* **1**: 431-435, 1981.
  33. Kaplan EL, Meier P. Nonparametric estimation from incomplete observation. *J Am Stat Assoc* **53**: 457-481, 1958.
  34. Cox DR. Regression models and life tables. *J R Stat Soc* **34**: 248-275, 1972.
  35. Ikeda K, Saitoh S, Suzuki Y, et al. Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis—A prospective observation of 2215 patients. *J Hepatol* **28**: 930-938, 1998.
  36. Ikeda K, Saitoh S, Koida I, et al. A multivariate analysis of risk factors for hepatocellular carcinogenesis—A prospective observation of 795 cases with viral and alcoholic cirrhosis. *Hepatology* **18**: 47-53, 1993.
  37. Kasahara A, Hayashi N, Mochizuki K, et al. The Osaka Liver Disease Study Group. Clinical characteristics of patients with chronic hepatitis C showing biochemical remission, without hepatitis C virus eradication, as a result of interferon therapy. *J Viral Hepat* **7**: 343-351, 2000.
  38. Yabuuchi I, Imai Y, Kawata S, et al. Long-term responders without eradication of hepatitis C virus after interferon therapy: characterization of clinical profiles and incidence of hepatocellular carcinoma. *Liver* **20**: 290-295, 2000.
  39. Okanoue T, Itoh Y, Minami M, et al. Interferon therapy lowers the rate of progression to hepatocellular carcinoma in chronic hepatitis C but not significantly in an advanced stage: a retrospective study in 1,148 patients—Viral Hepatitis Therapy Study Group. *J Hepatol* **30**: 653-659, 1999.
  40. Ikeda K, Arase Y, Saitoh S, et al. Anti-carcinogenic impact of interferon therapy in patients with chronic hepatitis C—A large-scale long-term study in a single center. *Intervirology* **49**: 82-90, 2006.
  41. Benvegnù L, Fattovich G, Noventa F, et al. Concurrent hepatitis B and C virus infection and risk of hepatocellular carcinoma in cirrhosis. A prospective study. *Cancer* **74**: 2442-2448, 1994.
  42. Tanaka K, Sakai H, Hashizume M, Hirohata T. A long-term follow-up study on risk factors for hepatocellular carcinoma among Japanese patients with liver cirrhosis. *Jpn J Cancer Res* **89**: 1241-1250, 1998.

# *Journal of* **Gastroenterology**

- ▶ Recent advances in autoimmune pancreatitis
- ▶ CE versus DBE in obscure GI bleeding
- ▶ Efficacy of H<sub>2</sub>-receptor antagonist for NERD in Japan
- ▶ IFN plus lamivudine combination therapy



The Japanese Society  
of Gastroenterology

 Springer

# Journal of Gastroenterology

Volume 43 · Number 6 · 2008

## Special Article

M. Otsuki, J.B. Chung, K. Okazaki, et al.

Asian diagnostic criteria for autoimmune pancreatitis: consensus of the Japan-Korea Symposium on Autoimmune Pancreatitis 403

## Reviews

K. Okazaki, K. Uchida, T. Fukui  
Recent advances in autoimmune pancreatitis: concept, diagnosis, and pathogenesis 409

J.-T. Li, Z.-X. Liao, J. Ping, et al.  
Molecular mechanism of hepatic stellate cell activation and antifibrotic therapeutic strategies 419

## Alimentary Tract

Y. Usta, I.N. Saltik-Temizel, H. Demir, et al.  
Comparison of short- and long-term treatment protocols and the results of second-line quadruple therapy in children with *Helicobacter pylori* infection 429

N. Kameda, K. Higuchi, M. Shiba, et al.  
A prospective, single-blind trial comparing wireless capsule endoscopy and double-balloon enteroscopy in patients with obscure gastrointestinal bleeding 434

R. González-Segovia, J.L. Quintanar, E. Salinas, et al.  
Effect of the flavonoid quercetin on inflammation and lipid peroxidation induced by *Helicobacter pylori* in gastric mucosa of guinea pig 441

M. Hongo, Y. Kinoshita, K. Haruma  
A randomized, double-blind, placebo-controlled clinical study of the histamine H<sub>2</sub>-receptor antagonist famotidine in Japanese patients with nonerosive reflux disease 448

## Liver, Pancreas, and Biliary Tract

N. Yuki, T. Nagaoka, K. Nukui, et al.  
Adding interferon to lamivudine enhances the early virologic response and reversion of the precore mutation in difficult-to-treat HBV infection 457

A. Kasahara, K. Kita, E. Tomita, et al.  
Repeated administration of recombinant human serum albumin caused no serious allergic reactions in patients with liver cirrhosis: a multicenter clinical study 464

L. Singh, D.K. Bakshi, S. Majumdar, et al.  
Mitochondrial dysfunction and apoptosis of acinar cells in chronic pancreatitis 473

A. Fujiwara, K. Sakaguchi, S. Fujioka, et al.  
Fibrosis progression rates between chronic hepatitis B and C patients with elevated alanine aminotransferase levels 484

## Case Report

H. Hosogi, S. Nagayama, J. Kawamura, et al.  
Molecular insights into Peutz-Jeghers syndrome: two probands with a germline mutation of *LKB1* 492

## On The Cover

Original figures are shown in N. Kameda et al., pp 434-40

 Springer

Indexed in *Index Medicus*, *Current Contents*, *EMBASE*, *BIOSIS*, *CINAHL*, *Chemical Abstracts*

## Repeated administration of recombinant human serum albumin caused no serious allergic reactions in patients with liver cirrhosis: a multicenter clinical study

AKINORI KASAHARA<sup>1</sup>, KEIJI KITA<sup>2</sup>, EIICHI TOMITA<sup>3</sup>, JOUJI TOYOTA<sup>4</sup>, YASUHARU IMAI<sup>5</sup>, and HIROMITSU KUMADA<sup>6</sup>

<sup>1</sup>Department of General Medicine, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita 565-0871, Japan

<sup>2</sup>Department of Internal Medicine, Kagawa Prefectural Central Hospital, Takamatsu, Japan

<sup>3</sup>Department of Gastroenterology, Gifu Municipal Hospital, Gifu, Japan

<sup>4</sup>Department of Gastroenterology, Sapporo-Kosei General Hospital, Sapporo, Japan

<sup>5</sup>Department of Gastroenterology, Ikeda Municipal Hospital, Ikeda, Japan

<sup>6</sup>Department of Hepatology, Toranomon Hospital, Tokyo, Japan

**Background.** We carried out a multicenter study to evaluate the safety of recombinant human serum albumin (rHSA), developed using the methylotrophic yeast *Pichia pastoris*, during and after repeated administration in patients with liver cirrhosis. **Methods.** rHSA was administered to 423 cirrhosis patients with ascites or edema. rHSA was administered three times over 3 days, and each 3-day treatment course was repeated at least three times with an interval of at least 2 weeks between courses. Adverse drug reactions (ADRs) were monitored during and after repeated rHSA administration. Specific antibody titers against *Pichia* yeast components were measured before and after treatment. Efficacy was evaluated on the basis of changes in serum albumin level, colloid osmotic pressure, and body weight. **Results.** ADRs were observed in 96 of 423 patients (22.7%), with no serious allergy or difference in the incidence of ADRs observed among the first, second, and third administrations. Specific IgE and IgG antibodies were detected before treatment in 19 and 422 patients, respectively. However, allergic ADRs were observed in 14 patients in whom specific IgE antibodies were not detected. No obvious relationship between allergic ADRs and specific IgE or IgG titers was identified. Serum albumin levels and colloid osmotic pressure increased significantly ( $P < 0.0001$ ), and body weight decreased significantly ( $P < 0.0001$ ) after rHSA administration. **Conclusions.** rHSA caused no serious allergic reactions even when three treatment courses were administered at intervals of at least 2 weeks.

**Key words:** recombinant human serum albumin (rHSA), *Pichia pastoris*, liver cirrhosis, allergic reactions, antibody testing

### Introduction

Plasma-derived human serum albumin (pHSA) is used to correct circulating plasma volume and improve colloid osmotic pressure. As pHSA is manufactured from human blood, the supply is limited and the risk of infection by unknown viruses or prions cannot be completely ruled out. To alleviate these problems, recombinant human serum albumin (rHSA) has been developed from *Pichia pastoris*<sup>1</sup> without using any animal-derived materials.

rHSA and pHSA have been shown to be identical in structure and physicochemical and immunochemical properties.<sup>2–4</sup> A phase III controlled clinical study of patients in liver cirrhosis, with increased serum albumin as the primary end point, showed that the efficacy of rHSA is comparable to that of pHSA.<sup>5,6</sup> A similar efficacy has been demonstrated in patients with hemorrhagic shock, thermal burns, or nephrotic syndrome.<sup>5,6</sup> No serious allergic symptoms were observed in these studies.<sup>5,6</sup>

At present, many recombinant pharmaceuticals with established efficacy and safety are being used, and among blood products, recombinants of blood coagulation factor VIII<sup>7,8</sup> are increasingly replacing plasma-derived preparations. Despite the high purity of rHSA, the potential onset of allergic reactions triggered by *Pichia* yeast components must be investigated, because the administered doses of rHSA are higher than those of other recombinants, and because administration might be repeated. This multicenter study was therefore conducted in cirrhosis patients to evaluate the safety and efficacy of rHSA during and after repeated administration of rHSA.



## Methods

### Patients

A total of 423 patients with liver cirrhosis [etiology: hepatitis B virus (HBV) in 59 patients, hepatitis C virus (HCV) in 264, alcohol consumption in 47, primary biliary cirrhosis (PBC) in three, autoimmune hepatitis in three, non-B, non-C hepatitis in ten, HBV plus HCV in four, HCV plus alcohol consumption in seven, and other in 26] were enrolled in this study from July 2002 to January 2005. Written informed consent was obtained from all patients.

All patients meeting the following criteria were included in this study: serum albumin level  $<3.0$  g/dl; presence of cirrhotic ascites or edema (legs or back); and age, 20–75 years. Beginning with the second treatment course, patients with serum albumin level  $<3.0$  g/dl or cirrhotic ascites or edema (legs or back) were eligible.

Patients who met one or more of the following criteria were excluded from each course: prothrombin time  $<30\%$  or  $5$  s  $\geq$  reference or control time; total bilirubin  $\geq 5.0$  mg/dl; serum creatinine  $\geq 4.0$  mg/dl; presence of hepatocellular carcinoma (HCC) with tumor embolism in the portal vein (main trunk or primary/secondary branch), main trunk of the hepatic vein, or postcaval vein; New York Heart Association class III/IV; history of shock in response to any pHSA preparation component; hepatic encephalopathy of grade II or higher severity at the time of consent; and pregnancy or lactation in women. Also excluded were any patients the investigator considered inappropriate for the study.

### Protocol amendment

During the course of this study, a phase I study of rHSA was conducted in the United States in healthy volunteers with high specific IgE antibodies against *Pichia* yeast components ( $\geq 0.7$  U<sub>A</sub>/ml). In the American study, serious allergic adverse drug reactions (ADRs) were observed in two of four patients (bronchospasms in one patient and bronchospasms and generalized urticaria in one patient), leading to discontinuation of the study. Consequently, new enrollment in the present study was temporarily suspended in May 2003 while the cause was investigated. A passive cutaneous anaphylaxis (PCA) reaction test in rats showed that the products used in the American study were more antigenic than those used in the present study. Afterward, purification techniques were improved and the problem was resolved.

In November 2003, the present study was restarted with an amended protocol that included the following

additional inclusion criteria: specific IgE antibody titers against *Pichia* yeast components were required to be  $<0.35$  U<sub>A</sub>/ml before each treatment, and hospital admission is indispensable from the time of the first administration (day 1) to the day after the final administration (day 4).

### Treatment

After confirmation of a negative skin prick test against *Pichia* yeast components, 25 g/day of rHSA (25% rHSA, Bipla, Chitose, Japan) was intravenously administered over 2 h each day for 3 days. This course of treatment was repeated for three courses, and optionally for five courses. Each course was started after an interval of at least 2 weeks after the previous course.

### Concomitant medication

During the study period, concomitant use of other investigational products was not allowed. Administration of plasma protein products, blood transfusion, reinstitution of concentrated ascites, paracentesis, and invasive treatment of complications associated with the underlying disease (e.g., endoscopic variceal ligation or sclerotherapy for esophageal varices, percutaneous ethanol injection therapy, or radiofrequency ablation for HCC) were prohibited from the day the inclusion/exclusion criteria were examined, except for specific IgE testing, until day 10. Continued use of already prescribed diuretics was allowed; however, the start of diuretics was prohibited during days 1–4. Branched-chain amino acid formulae and other pharmaceuticals could be used concomitantly.

### Measurements and observation

Inclusion/exclusion criteria were examined within 3 days prior to day 1 of each treatment course, and specific IgE screening, which was added as an inclusion criterion by the protocol amendment, was performed within 14 days prior to day 1.

ADRs were monitored during the study period. Hematological tests and urinalysis were performed at baseline (within 3 days prior to day 1) and on days 4 and 10, and antibody tests were performed on day 1 (pretreatment) and days 4 and 10 in each course to evaluate safety.

Serum albumin level, colloid osmotic pressure, and body weight were measured on days 1 (pretreatment) and 4 during each course to evaluate efficacy. Serum albumin level and colloid osmotic pressure were measured at Mitsubishi Chemical Medicine Corporation (Tokyo, Japan)

**Table 1.** Descriptions of *Pichia* yeast components

PYC-a	A centrifuged supernatant of non-rHSA-producing <i>Pichia</i> yeast culture
PYC-b	A <i>Pichia</i> yeast lysate from a non-rHSA producing <i>Pichia</i> transformant
PYC-c	A <i>Pichia</i> yeast component from a non-rHSA producing <i>Pichia</i> transformant purified via passage through a streamline column

PYC, *Pichia* yeast component

#### Antibody testing

Specific IgE and IgG antibody titers against three types of *Pichia* yeast components, PYC-a, -b, and -c (Table 1), were measured by fluorescent enzyme immunoassay (ImmunoCAP;<sup>9</sup> Phadia, Uppsala, Sweden) at Mitsubishi Chemical Medicine Corporation. Intraday (ten repetitions), interday (duplicate measurements over 5 days), and interinstrument (duplicate measurements using three instruments) reproducibility was assessed, yielding a maximum coefficient of variation of 11.6% and 8.6% for specific IgE and IgG antibody titers, respectively. The quantification limit was 0.35 U<sub>A</sub>/ml and 2 mg<sub>A</sub>/l for specific IgE and IgG antibody titers, respectively.

#### Data analysis

The incidence of ADRs and abnormal laboratory changes for which a causal relationship to rHSA could not be ruled out was assessed. In addition to allergic ADRs judged by each investigator, skin symptoms, including rash, drug eruption, eczema, erythema, Henoch-Schonlein purpura, purpura, pruritus, and generalized pruritus; respiratory symptoms, including cough, wheezing, rhinorrhea, and nasal congestion; and pyrexia were selected as "allergy-related ADRs" by the sponsor (Mitsubishi Tanabe Pharma Corporation, Osaka, Japan) and analyzed.

The incidence of ADRs that occurred in patients with at least one of the three types of specific IgE antibody titers detected at one or more time points was compared with those in whom antibodies were not detected at any time points.

As the three types of specific IgG antibody titers were not normally distributed, the value of each titer was logarithm-transformed to evaluate the time course. The incidence of ADRs that occurred in patients with an increase in specific IgG antibodies of  $\geq 50\%$  from baseline of the first course or the respective course was compared with that in patients with no increase.

#### Statistics

Incidences of ADRs were compared by using Fisher's exact test. Changes in serum albumin level, colloid osmotic pressure, and body weight from day 1 (pretreat-

ment) to day 4 were evaluated using a paired *t* test. Two-sided *P* values of  $<0.05$  were considered to indicate statistical significance. These analyses were performed with SAS statistical analysis software version 8.2 (SAS Institute, Cary, NC, USA).

#### Ethics

This study was conducted in compliance with good clinical practice. The protocol was reviewed and approved by the institutional review board at each study site.

#### Results

Screenings were conducted in 472 patients. Twenty-two, three, and four patients were excluded because of the inclusion/exclusion criteria (except for the specific IgE criterion), deterioration of hepatic cirrhosis, and other reasons, respectively. Of the original 472 patients, 39 were specific IgE positive before treatment. Of these 39 patients with positive IgE, 19 were included in this study: 18 were included before the protocol amendment in November 2003, and one patient whose specific IgE was negative in the screening test but became positive at just prior to the first course treatment was included after the protocol amendment. The remaining 20 patients were excluded because of positive IgE results. Consequently, rHSA was administered to 423 patients (267 men and 156 women) in total. At baseline, their mean age was  $63.3 \pm 8.1$  years. Mean serum albumin level, total bilirubin, creatinine, and prothrombin time were  $2.5 \pm 0.3$  g/dl,  $1.9 \pm 1.0$  mg/dl,  $1.0 \pm 0.5$  mg/dl, and  $61.3 \pm 16.1\%$ , respectively. Physical findings of ascites, imaging findings of ascites, and physical findings of edema were observed in 70.2%, 84.2%, and 75.2% of patients, respectively (Table 2).

#### Adverse drug reactions

rHSA was administered to 423, 314, 219, 63, and 34 patients during the first, second, third, fourth, and fifth courses, respectively.

The overall incidence of ADRs was 22.7% (96/423 patients). Common (incidence  $\geq 1\%$ ) ADRs were pyrexia (6.9%; 29/423), hepatic encephalopathy (2.4%;

Table 2. Baseline demographics

	No. of patients or mean $\pm$ SD	%
Sex		
Male	267	63.1
Female	156	36.9
Age, years	63.3 $\pm$ 8.1	
Cause of hepatic cirrhosis		
HBV	59	13.9
HCV	264	62.4
Alcohol consumption	47	11.1
PBC	3	0.7
Autoimmune hepatitis	3	0.7
Non-B, non-C	10	2.4
HBV + HCV	4	0.9
HCV + alcohol consumption	7	1.7
Others	26	6.1
Serum albumin (g/dl)	2.5 $\pm$ 0.3	
Total bilirubin (mg/dl)	1.9 $\pm$ 1.0	
Creatinine (mg/dl)	1.0 $\pm$ 0.5	
Prothrombin time <sup>a</sup> (%)	61.3 $\pm$ 16.1	
Prothrombin time <sup>b</sup> (s)	13.6 $\pm$ 1.4	
Ascites (physical diagnosis)		
-	126	29.8
+	297	70.2
Ascites (diagnostic imaging)		
-	66	15.6
+	356	84.2
Unknown	1	0.2
Edema		
-	103	24.3
+	318	75.2
Unknown	2	0.5
Hepatocellular carcinoma		
-	276	65.2
+	147	34.8
New York Heart Association		
No heart failure	397	93.9
NYHA classification		
I	24	5.7
II	2	0.5

HBV, hepatitis B virus; HCV, hepatitis C virus; PBC, primary biliary cirrhosis; NYHA, New York Heart Association

<sup>a</sup>n = 418

<sup>b</sup>n = 5

10/423), rash (2.1%; 9/423), pruritus (1.7%; 7/423), and dehydration (1.2%; 5/423) (Table 3). Uncommon but significant ADRs were hemorrhagic shock (0.2%; 1/423), hemorrhoidal hemorrhage (0.2%; 1/423), gastric variceal hemorrhage (0.2%; 1/423), and gastrointestinal hemorrhage (0.2%; 1/423) (Table 3).

Incidences of skin symptoms, respiratory symptoms, and pyrexia, defined as allergy-related ADRs by the sponsor, were 5.4% (23/423), 1.4% (6/423), and 6.9% (29/423) respectively (Table 4). Incidences of overall and allergy-related ADRs did not increase during repeated administration (Table 4).

ADRs judged to be allergic symptoms by the investigators were observed in 14 patients: rash (eight

patients), drug eruption (two), Henoch-Schönlein purpura (two), purpura (one), and wheezing (one). None of these allergic ADRs were serious, and they regressed or improved with appropriate intervention (Table 5). In all these patients, specific IgE antibodies against *Pichia* yeast components were not detected before or after rHSA administration.

Common (incidence  $\geq$ 1%) abnormal laboratory changes for which a causal relationship with rHSA could not be ruled out were aggravated urinary occult blood (2.9%; 12/421), aggravated urine protein (2.4%; 10/421), increased blood urea nitrogen (2.4%; 10/423), decreased hemoglobin (1.9%; 8/423), increased eosinophils (1.9%; 8/422), increased neutrophils (1.9%; 8/422), decreased red blood cells (1.7%; 7/423), decreased hematocrit (1.7%; 7/423), decreased lymphocytes (1.7%; 7/422), decreased platelets (1.4%; 6/423), increased creatinine (1.4%; 6/423), increased total bilirubin (1.2%; 5/423), and aggravated urine urobilinogen (1.4%; 6/421) (Table 6).

#### Specific IgE antibodies and ADRs

In 19 of 423 patients, at least one of the three types of specific IgE antibodies against *Pichia* yeast component was detected before treatment. Of these patients, antibody titers increased after treatment in four patients, but not in the remaining 15 patients. In five of the 404 patients negative for specific IgE antibodies before treatment, at least one of the specific IgE antibodies became detectable after treatment, but not in the remaining 399 patients.

The incidences of overall and allergy-related ADRs in the 24 patients with specific IgE antibodies detected before or after treatment did not differ from those in the 399 patients in whom the antibodies were not detected (Table 7).

It should be taken into account that specific IgE <0.35 was added as an inclusion criterion during the course of this study, in November 2003. Nineteen patients positive for specific IgE were included in this study. None of these patients experienced allergic ADRs. Furthermore, in the subgroup of specific IgE-negative patients, the allergic ADR incidence in patients before and after protocol amendment was 4.4% (9/206) and 2.5% (5/198), respectively, which are statistically not different. Consequently, the protocol amendment might not have affected the incidence of allergic ADRs.

#### Specific IgG antibodies and ADRs

Specific IgG antibodies against *Pichia* yeast components were detected in 422 of 423 patients before treatment. In 88 of 423 patients, specific IgG antibody titers increased by 50% or more after rHSA administration,

**Table 3.** Adverse drug reactions

System organ class	Incidence		
	≥1%	0.5% to <1%	<0.5%
Infections and infestations			Cellulitis (0.2%) Folliculitis (0.2%)
Blood and lymphatic system disorders		Anemia (0.5%)	Thrombocytopenia (0.2%) Nephrogenic anemia (0.2%)
Metabolism and nutrition disorders	Dehydration (1.2%)	Hyperammonemia (0.9%)	Hyperuricemia (0.2%)
Psychiatric disorders			Disorientation (0.2%) Delirium (0.2%)
Nervous system disorders	Hepatic encephalopathy (2.4%)	Headache (0.7%)	Subarachnoid hemorrhage (0.2%) Iliad nerve paralysis (0.2%) Syncope (0.2%) Cerebral infarction (0.2%) Convulsion (0.2%)
Cardiac disorders			Palpitations (0.2%)
Vascular disorders			Hot flush (0.2%) Shock hemorrhagic (0.2%)
Respiratory, thoracic and mediastinal disorders		Cough (0.7%)	Pleural effusion (0.2%) Rhinorrhea (0.2%) Nasal congestion (0.2%) Wheezing (0.2%) Pulmonary edema (0.2%)
Gastrointestinal disorders		Diarrhea (0.9%) Vomiting (0.5%)	Nausea (0.2%) Stomatitis (0.2%) Ascites (0.2%) Gastric varices hemorrhage (0.2%) Gastrointestinal hemorrhage (0.2%) Hemorrhoidal hemorrhage (0.2%) Feces discolored (0.2%)
Hepatobiliary disorders		Deterioration of hepatic cirrhosis (0.5%)	
Skin and subcutaneous tissue disorders	Rash (2.1%) Pruritus (1.7%)	Henoch-Schönlein purpura (0.5%) Drug eruption (0.5%)	Erythema (0.2%) Pruritus generalized (0.2%) Purpura (0.2%) Eczema (0.2%)
Musculoskeletal and connective tissue disorders			Pain in extremity (0.2%) Buttock pain (0.2%)
General disorders and administration site conditions	Pyrexia (6.9%)	Injection site pain (0.7%)	Malaise (0.2%) Feeling hot (0.2%) Edema (0.2%)
Investigations			Platelet count decreased (0.2%)

**Table 4.** Incidences of overall and allergy-related adverse drug reactions during each course

	First course (n = 423)		Second course (n = 314)		Third course (n = 219)		Fourth course (n = 63)		Fifth course (n = 34)		Total (n = 423)	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Overall	60	14.2	28	8.9	24	11.0	5	7.9	2	5.9	96	22.7
Skin symptoms	19	4.5	2	0.6	2	0.9	0	0.0	0	0.0	23	5.5
Respiratory symptoms	1	0.2	3	1.0	3	1.4	1	1.6	1	2.9	6	1.4
Pyrexia	20	4.7	8	2.5	5	2.3	1	1.6	1	2.9	29	6.9