

control samples and family samples, respectively (MAF=2.4% and 47%), under a multiplicative model of inheritance in first-set screening samples.

Regarding interpretation of the results from this study, several limitations should be mentioned: Firstly, we did not screen the entire region of *NRG1*. On that point, Corvin et al. showed an independent 'at-risk' haplotype close to an EST cluster of unknown function (*Hs.97362*) within intron 1 of *NRG1* (Corvin et al., 2004). Secondly, our samples were not assessed with the use of the standard structured interview, and therefore have the possibility of false negatives due to misdiagnosis or sampling bias. Detailed association analyses with dense markers in the entire region of *NRG1* in well-phenotyped samples, including symptomatology, are essential in future study.

In conclusion, these results indicate that the positionally and functionally attractive regions of *NRG1* are unlikely to contribute to susceptibility to schizophrenia in the Japanese population. Moreover the nature of our results support that two-stage analysis with large sample size is appropriate to examine the susceptibility genes for common diseases; independent samples for examination of significance found in screening results should be an integral part of experimental design in genetic association analysis. Imputation methods should also be used when only haplotype association shows significance, in order to check for possible causal SNPs that can reflect the haplotype.

Role of funding source

This work was supported in part by research grants from the Ministry of Education, Culture, Sports, Science and Technology, the Ministry of Health, Labor and Welfare, and the Japan Health Sciences Foundation (Research on Health Sciences Focusing on Drug Innovation).

Contributors

MI and NT designed the study, wrote the protocol and drafted the manuscript. MI, NT, SS, BA, YW, AN, YY, TK, YK, TK, and KK performed laboratory assays and the data-analysis. RH, HU, TI, TS, and MT advised on data-analysis. NO and NI participated in the design of the study, interpretation of the data, and drafting of the manuscript. All authors contributed to and have approved the final manuscript.

Conflict of interest

None.

Acknowledgements

We thank Ms S Ishihara, Mr M Hatano and Ms M Miyata for their technical support. We also thank Dr. Michael C. O'Donovan at Cardiff University for his helpful comments.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.schres.2008.01.010.

References

- Badner, J.A., Gershon, E.S., 2002. Meta-analysis of whole-genome linkage scans of bipolar disorder and schizophrenia. *Mol. Psychiatry* 7 (4), 405–411.
- Balding, D.J., 2006. A tutorial on statistical methods for population association studies. *Nat. Rev. Genet.* 7 (10), 781–791.
- Barrett, J.C., Fry, B., Maller, J., Daly, M.J., 2005. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21 (2), 263–265.
- Corfas, G., Roy, K., Buxbaum, J.D., 2004. Neuregulin 1-erbB signaling and the molecular/cellular basis of schizophrenia. *Nat. Neurosci.* 7 (6), 575–580.
- Corvin, A.P., Morris, D.W., McGhee, K., Schwaiger, S., Scully, P., Quinn, J., Meagher, D., Clair, D.S., Waddington, J.L., Gill, M., 2004. Confirmation and refinement of an 'at-risk' haplotype for schizophrenia suggests the EST cluster, *Hs.97362*, as a potential susceptibility gene at the Neuregulin-1 locus. *Mol. Psychiatry* 9 (2), 208–213.
- de Bakker, P.I., Yelensky, R., Pe'er, I., Gabriel, S.B., Daly, M.J., Altshuler, D., 2005. Efficiency and power in genetic association studies. *Nat. Genet.* 37 (11), 1217–1223.
- Dudbridge, F., 2003. Pedigree disequilibrium tests for multilocus haplotypes. *Genet. Epidemiol.* 25 (2), 115–121.
- Fukui, N., Muratake, T., Kaneko, N., Amagane, H., Someya, T., 2006. Supportive evidence for neuregulin 1 as a susceptibility gene for schizophrenia in a Japanese population. *Neurosci. Lett.* 396 (2), 117–120.
- Harrison, P.J., Law, A.J., 2006. Neuregulin 1 and schizophrenia: genetics, gene expression, and neurobiology. *Biol. Psychiatry* 60 (2), 132–140.
- Harrison, P.J., Weinberger, D.R., 2005. Schizophrenia genes, gene expression, and neuropathology: on the matter of their convergence. *Mol. Psychiatry* 10 (1), 40–68. image 5.
- Ikeda, M., Iwata, N., Suzuki, T., Kitajima, T., Yamanouchi, Y., Kinoshita, Y., Inada, T., Ujike, H., Ozaki, N., 2005. Association analysis of chromosome 5 GABAA receptor cluster in Japanese schizophrenia patients. *Biol. Psychiatry* 58 (6), 440–445.
- Iwata, N., Suzuki, T., Ikeda, M., Kitajima, T., Yamanouchi, Y., Inada, T., Ozaki, N., 2004. No association with the neuregulin 1 haplotype to Japanese schizophrenia. *Mol. Psychiatry* 9 (2), 126–127.
- Lewis, C.M., Levinson, D.F., Wise, L.H., DeLisi, L.E., Straub, R.E., Hovatta, I., Williams, N.M., Schwab, S.G., Pulver, A.E., Faraone, S.V., Brzustowicz, L.M., Kaufmann, C.A., Garver, D.L., Gurling, H.M., Lindholm, E., Coon, H., Moises, H.W., Byerley, W., Shaw, S.H., Mesen, A., Sherrington, R., O'Neill, F.A., Walsh, D., Kendler, K.S., Ekelund, J., Paunio, T., Lonnqvist, J., Peltonen, L., O'Donovan, M.C., Owen, M.J., Wildenauer, D.B., Maier, W., Nestadt, G., Blouin, J.L., Antonarakis, S.E., Mowry, B.J., Silverman, J.M., Crowe, R.R., Cloninger, C.R., Tsuang, M.T., Malaspina, D., Harkavy-Friedman, J.M., Svrakic, D.M., Bassett, A.S., Holcomb, J., Kalsi, G., McQuillin, A., Brynjolfsson, J., Sigmundsson, T., Petursson, H., Jazin, E., Zoega, T., Helgason, T., 2003. Genome scan meta-analysis of schizophrenia and bipolar disorder, part II: schizophrenia. *Am. J. Hum. Genet.* 73 (1), 34–48.
- Marchini, J., Howie, B., Myers, S., McVean, G., Donnelly, P., 2007. A new multipoint method for genome-wide association studies by imputation of genotypes. *Nat. Genet.* 39 (7), 906–913.
- Neale, B.M., Sham, P.C., 2004. The future of association studies: gene-based analysis and replication. *Am. J. Hum. Genet.* 75 (3), 353–362.
- Pritchard, J.K., 2001. Are rare variants responsible for susceptibility to complex diseases? *Am. J. Hum. Genet.* 69 (1), 124–137.

- Purcell, S., Cherny, S.S., Sham, P.C., 2003. Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics* 19 (1), 149–150.
- Stefansson, H., Sigurdsson, E., Steinthorsdottir, V., Bjornsdottir, S., Sigmundsson, T., Ghosh, S., Brynjolfsson, J., Gunnarsdottir, S., Ivarsson, O., Chou, T.T., Hjaltason, O., Birgisdottir, B., Jonsson, H., Gudnadottir, V.G., Gudmundsdottir, E., Bjornsson, A., Ingvarsson, B., Ingason, A., Sigfusson, S., Hardardottir, H., Harvey, R.P., Lai, D., Zhou, M., Brunner, D., Mutel, V., Gonzalo, A., Lemke, G., Sainz, J., Johannesson, G., Andresson, T., Gudbjartsson, D., Manolescu, A., Frigge, M.L., Gurney, M.E., Kong, A., Gulcher, J.R., Petursson, H., Stefansson, K., 2002. Neuregulin 1 and susceptibility to schizophrenia. *Am. J. Hum. Genet.* 71 (4), 877–892.
- Stefansson, H., Sarginson, J., Kong, A., Yates, P., Steinthorsdottir, V., Gudfinnsson, E., Gunnarsdottir, S., Walker, N., Petursson, H., Crombie, C., Ingason, A., Gulcher, J.R., Stefansson, K., St Clair, D., 2003. Association of neuregulin 1 with schizophrenia confirmed in a Scottish population. *Am. J. Hum. Genet.* 72 (1), 83–87.
- Walss-Bass, C., Liu, W., Lew, D.F., Villegas, R., Montero, P., Dassori, A., Leach, R.J., Almsy, L., Escamilla, M., Raventos, H., 2006. A novel missense mutation in the transmembrane domain of neuregulin 1 is associated with schizophrenia. *Biol. Psychiatry* 60 (6), 548–553.
- Williams, N.M., Green, E.K., Macgregor, S., Dwyer, S., Norton, N., Williams, H., Raybould, R., Grozeva, D., Hamshere, M., Zammit, S., Jones, L., Cardno, A., Kirov, G., Jones, I., O'Donovan, M.C., Owen, M.J., Craddock, N., 2006. Variation at the DAOA/G30 locus influences susceptibility to major mood episodes but not psychosis in schizophrenia and bipolar disorder. *Arch. Gen. Psychiatry* 63 (4), 366–373.
- Yang, J.Z., Si, T.M., Ruan, Y., Ling, Y.S., Han, Y.H., Wang, X.L., Zhou, M., Zhang, H.Y., Kong, Q.M., Liu, C., Zhang, D.R., Yu, Y.Q., Liu, S.Z., Ju, G.Z., Shu, L., Ma, D.L., Zhang, D., 2003. Association study of neuregulin 1 gene with schizophrenia. *Mol. Psychiatry* 8 (7), 706–709.