

Fig. 2. Retinal histological section of affected Suzuki Monkey showing the accumulation of drusen between the retinal pigment epithelium and Bruch's membrane.

The eyes of monkey are structurally similar to human eyes which make them extremely valuable for AMD studies. However, there are limitations in using this species over other laboratory animals. Monkeys have a relatively longer life span, have a longer gestation period, have a lower birth numbers resulting in a slower expansion of the pedigree, more difficult to genetically manipulate, and the maintenance cost is high. In the other laboratory animals, the differences in the eye structure, lack of a fovea, and a low cone/rod ratio compared to humans have been considered to be a disadvantage for using them as AMD models. However, they are easier to manipulate genetically and easier and less expensive to maintain. This has made the development of a mouse model of AMD very attractive, and a number of mouse AMD models have been reported recently.

6 Mouse Model for AMD

The mouse model described by Ambati et al is deficient either in monocyte chemoattractant protein-1 or its cognate C-C chemokine receptor-2. These mice were found to develop the cardinal features of AMD including accumulation of lipofuscin

in drusen beneath the RPE, photoreceptor atrophy, and CNV (Ambati et al. 2003). An impairment of macrophage recruitment allowed the accumulation of C5a and IgG, which leads to the production of vascular endothelial growth factor by the RPE cells and the development of CNVs. Another mouse model that has three known AMD risk factors: age, high fat cholesterol rich diet, and expression of human apolipoprotein E (apoE2, apoE3, apoE4) has been developed (Malek, Johnson, Mace, Saloupis, Schmechel, Rickman, Toth, Sullivan, and Bowes Rickman 2005). ApoE4-deficient mice are severely affected showing diffuse subretinal pigment epithelial deposits, drusen, thickened Bruch's membrane, and atrophy, hypopigmentation, and hyperpigmentation of the RPE.

Oxidative stress has long been linked to the pathogenesis of AMD. Imamura et al reported a Cu, Zn-superoxide dismutase (SOD1)-deficient mice that had features typical of AMD in human. Senescent Sod1 (-/-) mice had drusen, thickened Bruch's membrane, and choroidal neovascularization (Imamura, Noda, Hashizume, Shinoda, Yamaguchi, Uchiyama, Shimizu, Mizushima, Shirasawa, and Tsubota 2006). The number of drusen increased with age and also after exposure of young Sod1 (-/-) mice to excess light. The retinal pigment epithelial cells of Sod1 (-/-) mice showed oxidative damage, and their beta-catenin-mediated cellular integrity was disrupted. These findings suggested that oxidative stress may affect the junctional proteins necessary for the barrier integrity of the RPE. These observations strongly suggested that oxidative stress may play a major role in AMD.

The complement components, C3a and C5a, are present in drusen, and were observed in Bruch's membrane of a laser-induced CNV mice model. Neutralization of C3a or C5a by antibody or by blockade of their receptors by a complement inhibitor significantly reduced the CNV (Nozaki, Raisler, Sakurai, Sarma, Barnum, Lambris, Chen, Zhang, Ambati, Baffi, and Ambati 2006). These observations revealed a role for immunological mechanisms for the angiogenesis and provided evidence for future therapeutic strategies for AMD. Although the pathology of AMD is pronounced in the macula area, it is not confined to this region. Characteristics of human AMD such as thickening of Bruch's membrane, accumulation of drusen, and CNV have been observed in mouse models. Nevertheless, the primate model will still be the choice for AMD studies, especially at the stage when new therapeutic methods are tested and evaluated for the first time. However, it would be wise and more productive to study both primate and mouse models in AMD research. This will be necessary to learn the mechanisms underlying the disease and to identify clinical and molecular markers for the early stages of AMD. The findings from these studies will provide critical information needed to develop therapies for AMD.

7 Acknowledgements

This work was supported by the research grant from the Ministry of Health, Labour and Welfare of Japan.

References

- Ambati, J., Anand, A., Fernandez, S., Sakurai, E., Lynn, B. C., Kuziel, W. A., Rollins, B. J., Ambati, B. K. (2003) An animal model of age-related macular degeneration in senescent Ccl-2- or Ccr-2-deficient mice. *Nat. Med.* 9, 1390-1397.
- Chader, G. J. (2002) Animal models in research on retinal degenerations: Past progress and future hope. *Vision Res* 42, 393-399.
- Dewan, A., Liu, M., Hartman, S., Zhang, S. S., Liu, D. T., Zhao, C., Tam, P. O., Chan, W. M., Lam, D. S., Snyder, M., Barnstable, C., Pang, C. P., Hoh, J. (2006) HTRA1 promoter polymorphism in wet age-related macular degeneration. *Science* 314, 989-992.
- Edwards, A. O., Ritter, R. 3rd., Abel, K. J., Manning, A., Panhuysen, C., Farrer, L. A. (2005) Complement factor H polymorphism and age-related macular degeneration. *Science* 308, 421-424.
- El-Mofty, A., Gouras, P., Eisner, G., Balazs, E. A. (1978) Macular degeneration in rhesus monkey (*Macaca mulatta*). *Exp. Eye Res.* 27, 499-502.
- Gold, B., Merriam, J. E., Zernant, J., Hancox, L. S., Taiber, A. J., Gehrs, K., Cramer, K., Neel, J., Bergeron, J., Barile, G. R., Smith, R. T. (2006) AMD Genetics Clinical Study Group; G. S. Hageman, M. Dean, R. Allikmets. Variation in factor B (BF) and complement component 2 (C2) genes is associated with age-related macular degeneration. *Nat. Genet.* 38, 458-462.
- Gotoh, N., Yamada, R., Hiratani, H., Renault, V., Kuroiwa, S., Monet, M., Toyoda, S., Chida, S., Mandai, M., Otani, A., Yoshimura, N., Matsuda, F. (2006) No association between complement factor H gene polymorphism and exudative age-related macular degeneration in Japanese. *Hum. Genet.* 120, 139-143.
- Hageman, G. S., Anderson, D. H., Johnson, L. V., Hancox, L. S., Taiber, A. J., Hardisty, L. I., Hageman, J. L., Stockman, H. A., Borchardt, J. D., Gehrs, K. M., Smith, R. J., Silvestri, G., Russell, S. R., Klaver, C. C., Barbazetto, I., Chang, S., Yannuzzi, L. A., Barile, G. R., Merriam, J. C., Smith, R. T., Olsh, A. K., Bergeron, J., Zernant, J., Merriam, J. E., Gold, B., Dean, M., Allikmets, R. (2005) A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc. Natl. Acad. Sci. U S A* 102, 7227-7232.
- Haines, J. L., Hauser, M. A., Schmidt, S., Scott, W. K., Olson, L. M., Gallins, P., Spencer, K. L., Kwan, S. Y., Noureddine, M., Gilbert, J. R., Schmetz-Boutaud, N., Agarwal, A., Postel, E. A., Pericak-Vance, M. A. (2005) Complement factor H variant increases the risk of age-related macular degeneration. *Science* 308, 419-421.
- Hammond, C. J., Webster, A. R., Snieder, H., Bird, A. C., Gilbert, C. E., Spector, T. D. (2002) Genetic influence on early age-related maculopathy: A twin study. *Ophthalmology* 109, 730-736.
- Heiba, I. M., Elston, R. C., Klein, B. E., Klein, R. (1994) Sibling correlations and segregation analysis of age-related maculopathy: The Beaver Dam Eye Study. *Genet. Epidemiol.* 11, 51-67.
- Imamura, Y., Noda, S., Hashizume, K., Shinoda, K., Yamaguchi, M., Uchiyama, S., Shimizu, T., Mizushima, Y., Shirasawa, T., Tsubota, K. (2006) Drusen, choroidal neovascularization, and retinal pigment epithelium dysfunction in SOD1-deficient mice: a model of age-related macular degeneration. *Proc Natl Acad Sci U S A* 103, 11282-11287.
- Iyengar, S. K., Song, D., Klein, B. E., Klein, R., Schick, J. H., Humphrey, J., Millard, C., Liptak, R., Russo, K., Jun, G., Lee, K. E., Fijal, B., Elston, R. C. (2004) Dissection of genomewide-scan data in extended families reveals a major locus and oligogenic susceptibility for age-related macular degeneration. *Am. J. Hum. Genet.* 74, 20-39.
- Klein, R. J., Zeiss, C., Chew, E. Y., Tsai, J. Y., Sackler, R. S., Haynes, C., Henning, A. K., SanGiovanni, J. P., Mane, S. M., Mayne, S. T., Bracken, M. B., Ferris, F. L., Ott, J., Barnstable, C., Hoh, J. (2005) Complement factor H polymorphism in age-related macular degeneration. *Science* 308, 385-389.
- Schick, J. H., Iyengar, S. K., Klein, B. E., Klein, R., Reading, K., Liptak, R., Millard, C., Lee, K. E., Tomany, S. C., Moore, E. L., Fijal, B. A., Elston, R. C. (2003) A whole-genome screen of a quantitative trait of age-related maculopathy in sibships from the Beaver Dam Eye Study. *Am. J. Hum. Genet.* 72, 1412-1424.
- Majewski, J., Schultz, D. W., Weleber, R. G., Schain, M. B., Edwards, A. O., Matisse, T. C., Acott, T. S., Ott, J., Klein, M. L. (2003) Age-related macular degeneration--a genome scan in extended families. *Am. J. Hum. Genet.* 73, 540-550.
- Malek, G., Johnson, L. V., Mace, B. E., Saloupis, P., Schmechel, D. E., Rickman, D. W., Toth, C. A., Sullivan, P. M., Bowes Rickman, C. (2005) Apolipoprotein E allele-dependent pathogenesis: a model for age-related retinal degeneration. *Proc. Natl. Acad. Sci. U S A.* 102, 11900-11905.

- Mullins, R. F., Russell, S. R., Anderson, D. H., Hageman, G. S. (2000) Drusen associated with aging and age-related macular degeneration contain proteins common to extracellular deposits associated with atherosclerosis, elastosis, amyloidosis, and dense deposit disease. *FASEB J* 14, 835-846.
- Nicolas, M. G., Fujiki, K., Murayama, K., Suzuki, M. T., Mineki, R., Hayakawa, M., Yoshikawa, Y., Cho, F., Kanai, A. (1996) Studies on the mechanism of early onset macular degeneration in cynomolgus (*Macaca fascicularis*) monkeys. I. Abnormal concentrations of two proteins in the retina. *Exp. Eye Res.* 62, 211-219.
- Nicolas, M. G., Fujiki, K., Murayama, K., Suzuki, M. T., Shindo, N., Hotta, Y., Iwata, F., Fujimura, T., Yoshikawa, Y., Cho, F., Kanai, (1996) A. Studies on the mechanism of early onset macular degeneration in cynomolgus monkeys. II. Suppression of metallothionein synthesis in the retina in oxidative stress. *Exp. Eye Res.* 62, 399-408.
- Nozaki, M., Raisler, B. J., Sakurai, E., Sarma, J. V., Barnum, S. R., Lambris, J. D., Chen, Y., Zhang, K., Ambati, B. K., Baffi, J. Z., Ambati J. (2006) Drusen complement components C3a and C5a promote choroidal neovascularization. *Proc. Natl. Acad. Sci. U S A* 103, 2328-2333.
- Okamoto, H., Umeda, S., Obazawa, M., Minami, M., Noda, T., Mizota, A., Honda, M., Tanaka, M., Koyama, R., Takagi, I., Sakamoto, Y., Saito, Y., Miyake, Y., Iwata, T. (2006) Complement factor H polymorphisms in Japanese population with age-related macular degeneration. *Mol. Vis.* 12, 156-158.
- Russell, S. R., Mullins, R. F., Schneider, B. L., Hageman, G. S. (2000) Location, substructure, and composition of basal laminar drusen compared with drusen associated with aging and age-related macular degeneration. *Am J Ophthalmol* 129, 205-214.
- Seddon, J. M., Ajani, U. A., Mitchell, B. D. (1997) Familial aggregation of age-related maculopathy. *Am. J. Ophthalmology* 123, 199-206.
- Stafford, T. J., Anness, S. H., Fine, B. S. (1984) Spontaneous degenerative maculopathy in the monkey. *Ophthalmology* 91, 513-521.
- Yang, Z., Camp, N. J., Sun, H., Tong, Z., Gibbs, D., Cameron, D. J., Chen, H., Zhao, Y., Pearson, E., Li, X., Chien, J., Dewan, A., Harmon, J., Bernstein, P. S., Shridhar, V., Zabriskie, N. A., Hoh, J., Howes, K., Zhang, K. (2006) A variant of the HTRA1 gene increases susceptibility to age-related macular degeneration. *Science* 314, 992-993.
- Umeda, S., Ayyagari, R., Allikmets, R., Suzuki, M. T., Karoukis, A. J., Ambasudhan, R., Zernant, J., Okamoto, H., Ono, F., Terao, K., Mizota, A., Yoshikawa, Y., Tanaka, Y., Iwata, T. (2005) Early-onset macular degeneration with drusen in a cynomolgus monkey (*Macaca fascicularis*) pedigree: exclusion of 13 candidate genes and loci. *Invest. Ophthalmol. Vis. Sci.* 46, 683-691.
- Umeda, S., Suzuki, M. T., Okamoto, H., Ono, F., Mizota, A., Terao, K., Yoshikawa, Y., Tanaka, Y., Iwata, T. (2005) Molecular composition of drusen and possible involvement of anti-retinal autoimmunity in two different forms of macular degeneration in cynomolgus monkey (*Macaca fascicularis*). *FASEB J.* 19, 1683-1685.

国立医療学会誌

IRYO

医療

JAPANESE JOURNAL OF NATIONAL MEDICAL SERVICES

Vol.62

No. 9

SEPTEMBER 2008

図説 感覚器疾患シリーズ No. 9

特集：視覚障害(3)

眼疾患バイオマーカーの探索

国立病院機構東京医療センター 臨床研究センター 分子細胞生物学研究部

岩田 岳

国立医療学会

JAPANESE SOCIETY OF NATIONAL MEDICAL SERVICES

国立医療学会

国立医療学会は、国立高度専門医療センター、国立ハンセン病療養所、
独立行政法人国立病院機構などの会員を
中心とした学術団体です。

いわゆる国が実施する高度専門医療、政策医療に深く関わり、
医療の進歩発展、診療、研修教育を促進し、
国民医療の向上と、会員の親睦を目指します。

国立医療学会は開かれています。上記の国立医療機関に勤務されてい
る方はどの職種からでも正会員に登録していただけます。また、上記以外の
一般の方であっても、本学会の活動趣旨に賛同していただける方は、一定の
手続きをしていただきますと、賛助会員になれます。

正会員と賛助会員には、機関誌「医療」が定期配信されます。
正会員と賛助会員は機関誌「医療」に投稿できます。
入会手続きは、裏面を参照願います。

雑誌「医療」は、

- * 国立医療学会の機関誌です。
- * 多職種の会員構成である為に、国が行うあらゆる医療分野がカバー
される総合医学誌です。
- * 国が進める高度専門医療、政策医療、あらゆる医療分野からのホット
な情報が満載されています。
- * 厚労省が進める医学研究情報、科学研究費関連の研究トピックスを
いち早く知ることが出来ます。

最近のバックナンバーは、以下の通りです。

雑誌のみの定期購読も受け付けています。各1部850円となっています
ので、興味ある方は是非ご購入の上、医療現場でお役立て下さい。

今後の筋萎縮性側索硬化症医療のあり方を考える	第60巻第10号
長寿医療の最前線	第60巻第12号
神経疾患の摂食・嚥下・栄養を考える	第61巻第2号
スタートした障害者自立支援法	第61巻第3号
曲がり角に立つ重症心身障害医療	第61巻第11号
筋ジストロフィーの歯科学的問題	第61巻第12号
ソーシャルワークの最前線	第62巻第2号
循環器病克服10年戦略	第62巻第3号
がん医療における全人的な他職種チーム医療の可能性と課題	第62巻第4号

お問い合わせ先：国立医療学会事務局
TEL/FAX:03-3410-8881



眼疾患バイオマーカーの探索

岩田 岳

IRYO Vol. 62 No. 9 (512-515) 2008

キーワード：眼、遺伝子、遺伝子多型、シークエンサー、予防医学

はじめに

近年ゲノム配列や体内のタンパク質を分析して疾患の発症を予測する研究が行われている。この背景には飛躍的な技術革新によるゲノム解析技術や質量分析を用いた微量タンパク質の網羅的解析技術などの進歩がある。これらの技術はすでに多くの疾患で成果を上げており、眼科分野でも応用されている。患者の眼球において、まだ生理機能が保持されている環境では再生医療を、そして環境が破壊している場合には人工網膜等による視力回復が検討されている。本章ではこれらの研究に加え、未発症の患者予備軍を選別し、病気の予防あるいは遅延を目的とした予防医学について、その選別部分について最近の研究を紹介したい。

感覚器症例情報収集システムの構築

眼球内部は透明な組織で構成されていることから容易に組織内部を観察できる一方、生検することがきわめて困難な組織でもある。したがって早期診断に利用できるサンプルは涙液、血液、尿に限られ、必然的に研究対象は白血球のDNAと血漿成分に絞られる。また、手術中に破棄される房水、硝子体の

一部についても倫理委員会の承認と患者の同意があれば研究材料として利用できる。われわれはこれらのDNAや体液から疾患と関連する疾患バイオマーカーの探索を行っている。平成15年、国立病院機構および厚生労働省の支援を受けて、オンラインによる症例情報の収集とDNA・血漿バンクの設立を目的とした「感覚器ネットワーク」を東京医療センター臨床研究センター（感覚器センター）に構築した（図1）。関連施設の国立病院および大学病院から白内障、緑内障、加齢黄斑変性、網膜色素変性の症例情報をウェブ画面上から登録し、同時に検体を受け取る仕組みである。症例情報には個人情報を除く患者の臨床所見が詳細に登録できるようになっており、眼底写真の添付も可能である。遺伝子やタンパク質の解析結果は症例情報と一体化され、個々の患者についてデータベースが構築されることによって、疾患バイオマーカーの探索をより容易にしている。

遺伝子解析による加齢性眼疾患の感受性遺伝子の探索

これまでの研究から加齢性眼疾患の多くは多因子疾患（遺伝、習慣、環境）であると考えられている。

国立病院機構東京医療センター 臨床研究センター 分子細胞生物学研究部
別刷請求先：岩田岳 国立病院機構東京医療センター 臨床研究センター 分子細胞生物学研究部
〒152-8902 目黒区東が丘2-5-1
(平成20年7月18日受付)

Series of Articles on Sensory Disorders No. 9

Search for Biomarker of Ocular Diseases

Takeshi Iwata

Key Words: eye, gene, single nucleotide polymorphisms, DNA sequencer, preventive medicine




図1 感覚器センターに設置された症例情報システムと各種分析装置

加齢黄斑変性の場合、加齢、遺伝、喫煙、肥満、青い光などがリスク因子として報告されている。とくに遺伝因子（感受性遺伝子）については技術的な進歩によって全ゲノムを対象とした患者と健常者の比較が可能になっている。ヒトゲノムプロジェクトによってヒトゲノムには約22,000個の遺伝子が存在し、平均で1,000塩基に1つの割合で配列が個々に異なることが明らかになった。この一塩基配列の変化である遺伝子多型：Single Nucleotide Polymorphism (SNP) を組み合わせてブロックにし、比較することによって感受性遺伝子の染色体上の位置を調べることができる。SNPの組み合わせをハプロタイプと呼ぶが、複数の国が参加して、すべてのハプロタイプを明らかにする国際ハップマッププロジェクト (International HapMap Project, <http://www.hapmap.org/>) が進行中である。

ゲノム上には1千万個のSNPが存在すると計算されるが、これだけのSNPを患者個々に解析することは困難であった。しかし、最近SNPアレーの開発が進み、50万-100万種類のSNPを同時に検出することができるようになってきている。このチップを利用しても全SNPの5-10%しかカバーできないが、感受性遺伝子の位置情報、病気との相関（頻度、オ

ッズ比）について有用な情報が得られる。われわれはこの方法によって白内障、緑内障、加齢黄斑変性、偽落屑症候群の感受性遺伝子について解析している。疾患と相関するSNPは疾患を発症する塩基配列と物理的に近い位置に存在すると考えられるが、そのものではないことが多い。これはハップマッププロジェクトが健常者を対象にして作成されたデータベースであり、患者のSNPは登録されていないからである。発症原因の塩基配列を知るためにはSNP間の塩基配列をDNAシーケンサーで読むしかない。近年この塩基配列解読についても驚異的な技術革新が進行している（図2）。1990年にアメリカエネルギー省と厚労省によって30億ドルの予算で開始されたヒトゲノムプロジェクトが10年間かけて解読した30億塩基を1日で解読するDNAシーケンサーがアメリカPacific Bioscience社を含む複数の会社から発表される予定である。アメリカではヒトゲノムを1,000ドルで完全解読する1,000ドルゲノムプロジェクトが進行しており、今後SNPアレーからゲノム配列を患者と健常者で直接比較することによってより詳細な情報が得られると考えられる。また、DNAシーケンサーの感度が上がり、必要なDNA量も年々減少している。このことから採血は不要と




図2 DNAシーケンサー1台が1日に解読できるゲノム塩基数
1人のゲノムが1日で解読できるDNAシーケンサーが発表される予定である。

なり、爪や髪の毛から抽出されるDNA量でも十分な解析が可能になると予測される。

近年の研究によってゲノム上の2%の配列が約22,000遺伝子をコードし、残り70%の配列からRNAが生成され、このRNAが遺伝子発現やタンパク質合成の制御に関係していることが明らかになってきた。このようなタンパク質を合成しないRNA(ncRNA: non-coding RNA)が眼疾患にも関わっている可能性は高く、感受性遺伝子に加え、感受性ncRNAについても積極的に研究を進めている。

眼疾患のためのプロテオミクスとバイオマーカーの探索

感受性遺伝子が明らかにされ、発症の有無が判断できるようになっても、個々の患者の発症時期を予測することは困難である。体内を循環する血液や排泄される尿の成分を分析することによって発症時期を予測する研究が行われている。近年、質量分析計が飛躍的に進歩し、コンピューターソフトによる機械の操作性も改善されたことから、専門の技官でなくても操作できるようになってきた。発見しているタンパク質の総称をプロテーム、その分析方法をプロテオミクスというが、今回われわれは白内障、緑内障、加齢黄斑変性の患者から血漿を集め、質量分

析計を用いた血漿プロテオミクスを行った。血漿にはアルブミン、グロブリンなど20種類のタンパク質が重量換算で99%を占めており、これらを除去しなければ疾患によって変動する微量タンパク質を質量分析計の検出できる感度で捕捉することが困難である。そこでわれわれは東レ株式会社との共同研究によって低分子のタンパク質だけを分離・濃縮する分画装置を開発し、患者血漿の分画を試みた(図3)。分画成分はさらに逆相クロマトグラフィー、トリブシン処理、2次元クロマトグラフィーによる分離を経て、イオントラップ型質量分析計によってタンパク質の同定が行われた。その結果、白内障、緑内障、加齢黄斑変性からそれぞれ固有に17、15、21種類のタンパク質が同定され、現在これらのタンパク質についてELISAやウエスタンブロットによって確認作業を進めている。同様な方法によって涙液、房水、硝子体液など眼由来の体液についても解析を行う予定である。また、今回同定されたタンパク質情報の確認作業と遺伝子多型情報と症例情報に結びつけるための作業を行っている。

おわりに

眼疾患のバイオマーカー探索として遺伝子解析と質量分析計を用いた眼疾患プロテオミクスについて




図3 中空糸膜を用いた低分子分画装置による微量タンパク質の濃縮
東レ株式会社が開発した分画装置によって、選択的に低分子（50kDa以下）の血漿タンパクを質量分析計で検出できる濃度に調製できるようになった。

紹介した、これまでに日本人と欧米人との間で疾患と相関する遺伝子多型が一致しないことがたびたび報告されている。しかし、日本人と体質がより近いと考えられるアジアの人々と連携しながら研究が進めば、日本人で発見された疾患バイオマーカーを利用して、広くアジアの患者の早期診断にも利用できる可能性がある。

【参考文献】

- 1) 岩田岳. 遺伝子解析の現状、またその行方は？
打倒！加齢黄斑変性，眼科インストラクションコース，メディカルビュー社；2005；p20-3.
- 2) 岩田岳. 加齢黄斑変性の遺伝子研究の最前線，特集：網膜脈絡膜変性疾患のアップデート，あたらしい眼科 2006；23：1125-31.
- 3) Okamoto H, Umeda S, Obazawa M et al. Complement Factor H Polymorphisms in Japanese Population with Age-Related Macular Degeneration. Mol Vis 2006；12：156-8.
- 4) 岩田岳. 網膜・硝子体のプロテオーム解析，日の眼科 2007；78：577-82.
- 5) Yoshida T, Wan AD, Zhang H et al. HTRA 1 Promoter Polymorphism Predisposes Japanese to AMD. Mol Vis 2007；13：545-8.

平成 年 月 日

国立医療学会理事長 殿

国立医療学会会員 新規加入申込書

下記のとおり国立医療学会への入会を申し込みます

氏 名： _____

連絡先 住所：〒 _____

TEL： _____ FAX： _____

所属施設名： _____

職 名： _____

- 正規会員（国立病院機構，国立高度専門医療センターおよび国立ハンセン病療養所等の国立医療施設に所属の方）
- 賛助会員（上記以外の施設に所属する方）

1) 入会希望年月：平成 年 月から
（年度途中からの入会の場合は会費は月割りとなります）

2) 職 種（番号に○をしてください）

- | | |
|------------|-----------------|
| 1. 医師 | 8. 言語聴覚士 |
| 2. 看護師 | 9. 臨床工学技士 |
| 3. 薬剤師 | 10. 管理栄養士 |
| 4. 臨床検査技師 | 11. 医療ソーシャルワーカー |
| 5. 診療放射線技師 | 12. 診療情報管理士 |
| 6. 理学療法士 | 13. 事務 |
| 7. 作業療法士 | 14. その他 |

※申込書はまず下記国立医療学会事務局に FAX または郵便でお申し込みください。
申し込み受付後事務局よりご連絡致します。

申込書送信先

国立医療学会事務局

〒152-0021

東京都目黒区東が丘2-5-23

FAX：03-3410-8881

雑誌「医療」バックナンバー

雑誌「医療」は、バックナンバーの購読や定期購読を受け付けています。興味のある方はぜひお買い求めになり、医療現場でお役立てください。

特集テーマ	掲載号数
頭頸部外科手術と喉頭機能外科の進歩	第60巻第4号
喉頭摘出術	第60巻第6号
正常圧水頭症（その1）	第60巻第7号
（その2）	第60巻第8号
今後の筋萎縮性側索硬化症医療のあり方を考える	第60巻第10号
長寿医療の最前線	第60巻第12号
神経疾患の摂食・嚥下・栄養を考える	第61巻第2号
スタートした障害者自立支援法	第61巻第3号
曲がり角に立つ重症心身障害医療	第61巻第11号
筋ジストロフィーの科学的問題	第61巻第12号
ソーシャルワークの最前線	第62巻第2号
循環器病克服10年戦略	第62巻第3号
がん医療における全人的な多職種チーム医療の可能性と課題	第62巻第4号

<購読のお申し込み方法>

雑誌「医療」定期購読やバックナンバー購読のお申し込みは、下の申込み用紙に必要事項をご記入の上、FAXにて国立医療学会事務局（FAX 03-3410-8881）までお申し込みください。

「医療」購読申込書

- 雑誌「医療」を最新号から（ ）冊定期購読する
- 第（ ）巻第（ ）号のバックナンバーを（ ）冊購読する

氏名			
連絡先	住所 〒		
	TEL	FAX	

LOXLI variants in elderly Japanese patients with exfoliation syndrome/glaucoma, primary open-angle glaucoma, normal tension glaucoma, and cataract

Masaki Tanito,¹ Masayoshi Minami,² Masakazu Akahori,² Sachiko Kaidzu,¹ Yasuyuki Takai,¹
 Akihiro Ohira,¹ Takeshi Iwata²

(The first three authors contributed equally to this work.)

¹Department of Ophthalmology, Shimane University Faculty of Medicine, Izumo, Shimane, Japan; ²National Institute of Sensory Organs, National Hospital Organization Tokyo Medical Center, Tokyo, Japan

Purpose: To evaluate the association of lysyl oxidase like 1 (*LOXLI*) gene variants in Japanese patients with open-angle glaucoma.

Methods: We evaluated the association of three *LOXLI* variants (rs1048661, rs3825942, and rs2165241) in 142 Japanese patients with exfoliation syndrome (EX; n=59) and exfoliation glaucoma (EG; n=83) as well as in 251 control patients aged 70 years or older with primary open-angle glaucoma (PG; n=40), normal tension glaucoma (NG; n=54), and cataract (CT; n=157).

Results: In comparison with the CT group, the single nucleotide polymorphisms (SNPs) showed significant association with EX, EG, and EX+EG. The odds ratio (OR)=19.71–28.23 and $p=1.69 \times 10^{-23}$ – 3.00×10^{-45} for allele T of rs1048661; OR=28.21–39.78 and $p=1.77 \times 10^{-4}$ – 2.42×10^{-22} for allele G of rs3825942; and OR=16.59–23.40 and $p=4.79 \times 10^{-5}$ – 1.08×10^{-9} for allele C of rs2165241. In comparison with the controls (CT+PG+NG), the haplotype rs1048661/rs3825942 (T/G) was significantly associated with EX+EG ($p=8.27 \times 10^{-44}$), and haplotype G/A had a significant protective effect ($p=2.25 \times 10^{-14}$). None of the three SNPs showed significant differences between the EX and EG groups or between the PG and NG groups.

Conclusions: These SNPs are associated with exfoliation syndrome/glaucoma in the Japanese population. The risk alleles in rs1048661 and rs2165241 are different from other populations. Additional genetic or environmental risk factors other than these *LOXLI* SNPs could be associated with the development of exfoliation syndrome as well as exfoliation glaucoma among exfoliation syndrome patients.

Exfoliation syndrome is the most common identifiable cause of open-angle glaucoma worldwide. It is an age-related, generalized disorder of the extracellular matrix that is characterized by the production and progressive accumulation of a fibrillar extracellular material in many ocular tissues [1]. A recent genome-wide association study demonstrated that one intronic single nucleotide polymorphism (SNP, rs2165241) and two exonic SNPs (rs1048661 [R141L] and rs3825942 [G153D]), which are located in the first exon of the lysyl oxidase like1 (*LOXLI*) gene on chromosome 15q24.1, are highly associated with exfoliation syndrome in Icelandic and Swedish populations while none of these SNPs were associated with primary open-angle glaucoma in either of the two populations [2].

Until now, several lines of studies have reported significant associations of these *LOXLI* SNPs with exfoliation syndrome [3-16] or the lack of associations of these SNPs with primary open-angle glaucoma [16-18] in various populations. In these studies, subjects in their 40s and 50s are sometimes recruited as controls. The prevalence of clinical exfoliation syndrome increases with age, particularly after the age of 60 years [1]. However, preclinical exfoliation syndromes in younger generations are able to distinguish from normal subjects until patients become older. Accordingly, the level of statistical significance between exfoliation syndrome and controls could be underestimated in studies that include younger patients in the control group. In this study, we evaluated the association of three *LOXLI* SNPs in Japanese patients with exfoliation syndrome/glaucoma, primary open angle glaucoma, normal tension glaucoma, and cataract. To reduce the chance of misclassifying latent syndromes within the control group, only patients aged 70 years or older were recruited as primary open-angle glaucoma, normal tension glaucoma, or cataract subjects in this study. To the best of our knowledge, this is the first study reporting an association of

Correspondence to: Takeshi Iwata, PhD, National Institute of Sensory Organs, National Hospital Organization, Tokyo Medical Center, 2-5-2 Higashigaoka, Meguro-ku, Tokyo, 152-8902, Japan; Phone: +81-3-3411-1026; FAX: +81-3-3411-1026 email: iwataakeshi@kankakuki.go.jp.

TABLE 1. SUMMARY OF STUDY POPULATIONS.

Parameter	Exfoliation syndrome or glaucoma			Primary glaucoma			CT	p value
	EX	EG	EX+EG	PG	NG	PG+NG		
No. of subjects	59	83	142	40	54	94	157	
Male:Female								
N	12:47	42:41	54:88	14:26	18:36	32:62	45:112	0.2264*
%	20.3:79.7	50.6:49.4	50.6:49.4	35.0:65.0	33.3:66.7	34.0:66.0	28.7:71.3	
Age (years)								
mean±SD	78.2±8.0	78.8±8.5	78.5±8.2	75.6±5.3	78.3±4.8	77.2±5.1	77.2±5.0	0.1271**
Range	55-95	57-95	55-95	70-87	70-91	70-91	70-90	

The asterisk and double asterisk indicate that the p values were obtained by Pearson's χ^2 test and one-way ANOVA, respectively, among three (EX+EG, PG+NG, and CT) groups. EX, exfoliation syndrome; EG, exfoliation glaucoma; PG, primary open-angle glaucoma aged 70 years or older; NG, normal tension glaucoma aged 70 years or older; CT, cataract aged 70 years or older.

LOXL1 SNPs with exfoliation syndrome/glaucoma using age-controlled subjects. This is also the first study reporting the association of *LOXL1* SNPs with normal tension glaucoma.

METHODS

Subjects: Three hundred and ninety-three unrelated Japanese subjects presenting exfoliation syndrome without glaucoma (EX), exfoliation syndrome with glaucoma (EG), primary open-angle glaucoma (PG), normal tension glaucoma (NG), and cataract (CT) were recruited at the Shimane University Hospital and Iinan Hospital in Shimane, Japan. This study adhered to the tenets of the Declaration of Helsinki. The research was reviewed and approved by the Institutional Review Boards of both hospitals. Written informed consent was obtained from all of the subjects. All of the subjects underwent a dilated pupil examination of the anterior segments, ocular media, and fundus using a slit-lamp and a funduscope. The subjects with EX and EG exhibited the typical pattern of exfoliation material on the anterior lens surface and/or pupillary margin during slit-lamp examination. The subjects with EG and PG had a history of intraocular pressure (IOP) greater than or equal to 21 mmHg, and the subjects with EX, NG, and CT had a history of IOP never exceeding 20 mmHg. The subjects with PG, NG, and EG presented a typical glaucomatous optic disc cupping or rim thinning and visual field loss. The patients who had a history of IOP greater than or equal to 21 mmHg but no glaucomatous optic disc changes nor visual field loss were assigned as EG in this study. To avoid possible misclassification of latent exfoliation syndromes as PG, NG, or CT, patients younger than 70 years old were not recruited. The number of subjects, each gender, and the mean and range of ages in each group are summarized in Table 1.

DNA genotyping: Genomic DNA was extracted from the peripheral white blood cells of each subject. Polymerase chain reaction (PCR) was performed using primers designed to amplify the genomic region containing both rs1048661 and rs3825942 (forward primer: 5'-AGG TGT ACA GCT TGC

TCA ACT C-3' and reverse primer: 5'-TAG TAC ACG AAA CCC TGG TCG T-3') or just rs2165241 (forward primer: 5'-AGA ATG CAA GAC CTC AGC ATG AG-3' and reverse primer: 5'-TAG TGG CCA GAG GTC TGC TAA G-3'). The sequence was determined based on the dideoxy terminator method using an ABI PRISM 3130xl Genetic Analyzer (Applied Biosystems, Foster City, CA) according to the manufacturer's protocol. We used SeqScape Software version 2.5 (Applied Biosystems) to analyze the sequence alignment.

Statistical analysis: The deviation of the genotype distributions from the Hardy-Weinberg equilibrium was assessed in the case and control samples using HAPLOVIEW version 4.0 [19]. Statistical analysis was performed using R version 2.6.2. Fisher's exact test was used to compare the allele or genotype frequencies of each case group with the controls. The odds ratio (OR) and 95% confidence intervals (CIs) were calculated by the logistic regression method. Individual haplotypes and their estimated population frequencies were inferred using HAPLOVIEW version 4.0 [19] with all of the parameters set at the default values.

RESULTS

The allelic and genotypic counts and frequencies of SNPs rs1048661, rs3825942, and rs2165241 within *LOXL1* are shown in Table 2. The ORs and p values for the allelic and genotypic frequencies of the three SNPs in comparison between the cases (EX, EG, and EX+EG) and controls (CT, PG, NG, and PG+NG) are shown in Table 3. In comparison with the CT group, the SNPs showed a significant association with EX, EG, and EX+EG for the T allele of rs1048661 at OR=19.71-28.23 and $p=1.69 \times 10^{-23}$ - 3.00×10^{-45} , the G allele of rs3825942 at OR=28.21-39.78 and $p=1.77 \times 10^{-8}$ - 2.42×10^{-22} , and the C allele of rs2165241 at OR=16.59-23.40 and $p=4.79 \times 10^{-5}$ - 1.08×10^{-9} (Table 3). The genotypes, TT of rs1048661 ($p=4.11 \times 10^{-25}$ - 3.78×10^{-43}), GG of rs3825942 ($p=3.53 \times 10^{-11}$ - 2.10×10^{-39}), and CC of rs2165241 ($p=1.95 \times 10^{-4}$ - 1.07×10^{-8}), also showed significant

TABLE 2. ALLELIC AND GENOTYPIC COUNTS AND FREQUENCIES OF rs1048661, rs3825942, AND rs2165241 IN POPULATION SYNDROME/GLAUCOMA (EX, EG, AND EX+EG), PRIMARY GLAUCOMA (PG, NG, PG+NG), AND CATARACT (CT).

Exfoliation syndrome or glaucoma														
Allele	EX		EG		EX+EG		PG		NG		Primary glaucoma		CT	
	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency
rs1048661														
T	111	0.941	159	0.958	270	0.951	39	0.488	50	0.463	89	0.473	140	0.446
G	7	0.059	7	0.042	14	0.049	41	0.513	58	0.537	99	0.527	174	0.554
Genotype														
TT	54	0.915	76	0.916	130	0.916	10	0.25	8	0.148	18	0.192	25	0.159
TG	3	0.051	7	0.084	10	0.070	19	0.475	34	0.630	53	0.564	90	0.573
GG	2	0.034	0	0	2	0.014	11	0.275	12	0.222	23	0.245	42	0.268
rs3825942														
G	117	0.992	165	0.994	282	0.993	64	0.800	86	0.796	150	0.798	253	0.806
A	1	0.009	1	0.006	2	0.007	16	0.200	22	0.204	38	0.202	61	0.194
Genotype														
GG	58	0.983	82	0.988	140	0.986	25	0.625	33	0.611	58	0.617	100	0.637
AG	1	0.017	1	0.012	2	0.014	14	0.350	20	0.370	34	0.362	53	0.338
AA	0	0	0	0	0	0	1	0.025	1	0.019	2	0.021	4	0.026
rs2165241														
C	117	0.992	165	0.994	282	0.993	77	0.963	93	0.861	170	0.904	275	0.876
T	1	0.009	1	0.006	2	0.007	3	0.038	15	0.139	18	0.096	39	0.124
Genotype														
CC	58	0.983	82	0.988	140	0.986	37	0.925	40	0.741	77	0.819	122	0.777
CT	1	0.017	1	0.012	2	0.014	3	0.075	13	0.241	16	0.170	31	0.198
TT	0	0	0	0	0	0	0	0	1	0.019	1	0.011	4	0.026

EX, exfoliation syndrome; EG, exfoliation glaucoma; PG, primary open angle glaucoma aged 70 years or older; NG, normal tension glaucoma aged 70 years or older; and CT, cataract aged 70 years or older.

TABLE 3. ODDS RATIOS AND P VALUES FOR ALLELIC AND GENOTYPIC FREQUENCIES OF rs1048661, rs3825942, AND rs2165241 IN COMPARISON BETWEEN CASES (EX, EG, AND EX+EG) AND CONTROLS (CT, PG, NG, AND PG+NG).

Parameter	EX			EG			EX+EG					
	Versus CT	Versus PG	Versus NG	Versus CT	Versus PG	Versus NG	Versus CT	Versus PG	Versus NG			
rs1048661												
Allele												
p value	1.69x 10 ⁻²²	3.23x 10 ⁻¹³	2.55x 10 ⁻¹⁶	5.65x 10 ⁻³³	1.92x 10 ⁻¹⁷	1.32x 10 ⁻²¹	4.71x 10 ⁻²⁶	3.00x 10 ⁻⁴⁵	2.61x 10 ⁻²⁰	4.40x 10 ⁻²⁶	1.18x 10 ⁻³¹	
OR	19.71	16.67	18.39	28.23	23.88	26.35	25.27	23.97	20.27	22.37	21.45	
95% CI	8.90-43.67	6.91-40.22	7.84-43.14	7.80-39.88	12.83-62.14	9.96-57.27	11.31-61.41	11.25-56.76	13.40-42.87	10.14-40.56	11.60-43.16	11.67-39.43
Genotype												
p value	4.11x 10 ⁻²³	1.16x 10 ⁻¹¹	1.91x 10 ⁻¹⁷	2.31x 10 ⁻¹⁹	4.80x 10 ⁻³²	2.12x 10 ⁻¹⁴	6.50x 10 ⁻²¹	3.42x 10 ⁻²⁴	3.78x 10 ⁻⁴³	1.04x 10 ⁻¹⁶	2.74x 10 ⁻²⁵	3.57x 10 ⁻³¹
rs3825942												
Allele												
p value	1.77x 10 ⁻⁴	2.10x 10 ⁻³³	2.10x 10 ⁻³³	2.10x 10 ⁻³³	2.42x 10 ⁻³²	6.48x 10 ⁻⁸	4.72x 10 ⁻⁹	1.23x 10 ⁻¹⁰	3.98x 10 ⁻¹⁶	8.68x 10 ⁻¹⁰	1.47x 10 ⁻¹¹	1.84x 10 ⁻¹⁴
OR	28.21	29.25	29.93	29.64	39.78	41.25	42.21	41.8	34	35.25	36.07	35.72
95% CI	3.86-205.97	3.79-225.66	3.96-226.36	4.01-219.07	5.46-289.78	5.36-317.49	5.59-318.47	5.67-308.20	8.23-140.45	7.91-157.16	8.31-156.49	8.50-150.11
Genotype												
p value	2.10x 10 ⁻³³	2.10x 10 ⁻³³	2.10x 10 ⁻³³	2.10x 10 ⁻³³	3.53x 10 ⁻¹¹	7.19x 10 ⁻⁸	2.88x 10 ⁻⁹	1.11x 10 ⁻¹⁰	6.37x 10 ⁻¹⁶	1.20x 10 ⁻⁹	1.06x 10 ⁻¹¹	1.17x 10 ⁻¹⁴
rs2165241												
Allele												
p value	4.79x 10 ⁻⁵	0.3051	1.03x 10 ⁻³³	1.20x 10 ⁻³	7.87x 10 ⁻⁷	0.1022	4.87x 10 ⁻⁶	8.28x 10 ⁻⁵	1.08x 10 ⁻⁹	0.0729	1.52x 10 ⁻⁷	3.02x 10 ⁻⁶
OR	16.59	4.56	18.87	12.39	23.4	6.43	26.61	17.47	20	5.49	22.74	14.93
95% CI	2.25-122.20	0.47-44.63	2.45-145.49	1.63-94.08	3.18-171.92	0.66-62.81	3.46-204.70	2.31-132.36	4.78-83.62	0.90-33.46	5.11-101.30	3.42-65.14
Genotype												
p value	1.95x 10 ⁻⁴	0.3	1.59x 10 ⁻⁴	2.82x 10 ⁻³	5.14x 10 ⁻⁶	0.1004	7.16x 10 ⁻⁶	1.23x 10 ⁻⁴	1.07x 10 ⁻⁴	0.0715	2.79x 10 ⁻⁷	4.71x 10 ⁻⁶

OR, odds ratio; CI, confidence interval; NA, not applicable; EX, exfoliation syndrome; EG, exfoliation glaucoma; PG, primary open-angle glaucoma aged 70 years or older; NG, normal tension glaucoma aged 70 years or older; and CT, cataract aged 70 years or older. p values were obtained by Fisher's exact probability test.

TABLE 4. ODDS RATIOS AND P VALUES FOR THREE SNPs IN COMPARISON BETWEEN EX AND EG.

EX versus EG				
	rs1048661		rs3825942	rs2165241
	Allele			
p value	0.5829		1	1
OR	1.43		1.41	1.41
95% CI	0.49-4.20		0.09-22.78	0.09-22.78
	Genotype			
p value	0.2005		1	1

No significant association was found between EX and EG. OR, odds ratio; CI, confidence interval; EX, exfoliation syndrome; and EG, exfoliation glaucoma. p values were obtained by Fisher's exact probability test.

associations with EX, EG, and EX+EG when they were compared to the CT group (Table 2 and Table 3). Significant associations with EX, EG, and EX+EG were detected in comparisons with the primary glaucoma groups (PG, NG, and PG+NG) for the alleles, T of rs1048661, G of rs3825942, and C of rs2165241, as well as the genotypes, TT of rs1048661, GG of rs3825942, and CC of rs2165241, with the exception of the comparisons between EX, EG, or EX+EG and the PG groups for allelic (OR=4.56-6.43 and $p=0.3051-0.0729$) and genotypic ($p=0.3000-0.0715$) frequencies in rs2165241 (Table 3).

None of the three SNPs showed a significant difference between the EX and EG groups in these allelic or genotypic frequencies (Table 4). In addition to this, none of the three SNPs showed significant differences in their allelic or genotypic frequencies in comparisons between the primary glaucoma (PG, NG, or PG+NG) and CT groups or between the PG and NG groups (data not shown), excepted for the allele C of rs2165241 ($p=0.0233$) and the genotype CC of rs2165241 ($p=0.0294$) in comparison between the PG and NG groups.

The two SNPs, rs1048661 and rs3825942, were in linkage disequilibrium ($D'=1$). In our study populations, only three of the four possible haplotypes in rs1048661/rs3825942 were detected (Table 5 and Table 6). In the comparisons between cases (EX, EG, or EX+EG) and controls (CT+PG+NG), the T and G were significantly associated with EX, EG, and EX+EG, and the G and A had a significant protective effect (Table 5 and Table 6).

DISCUSSION

Based on this study among 393 elderly Japanese patients with exfoliation syndrome, exfoliation glaucoma, primary open-angle glaucoma, and cataract, we confirmed the findings of Thorleifsson and colleagues [2] that three SNPs within *LOXL1* are strongly associated with exfoliation syndrome and glaucoma. In addition to the original study in Icelandic and Swedish populations, allele G of rs3825942 has been consistently suggested as a risk-associated allele of

exfoliation syndrome/glaucoma in five studies from the United States [3-7], two studies from Europe [8,9], one study from India [10], one study from Australia [11], and six studies including this one from Japan [12-16]. In this study, we found that allele T of rs1048661 is associated with exfoliation syndrome/glaucoma, which is consistent with other studies in the Japanese population [12-16], while allele G is reported to be risk-associated in studies from other countries [2,4,6-11]. We also found that allele C of rs2165241 is associated with exfoliation syndrome/glaucoma, which is consistent with one study from Japan [13], while allele T is risk-associated in other populations [2,4,5,7,8]. Allelic frequencies of the three SNPs reported in previous studies and the current study are summarized in Figure 1. These results suggest a possibility that the missense changes in *LOXL1* are not actually causative but mark a haplotype that carries variants that may indeed be causative.

This study revealed extremely high ORs and significant p values for the three SNPs in the comparisons between the exfoliation syndrome/glaucoma groups and cataract groups. Only patients aged 70 years or older were recruited as control subjects in this study, enabling us to reduce the chance of misclassifying latent or preclinical exfoliation syndromes into the control group. Accordingly, the inclusion criteria of the control group might contribute to the extremely significant association of the *LOXL1* SNPs with exfoliation syndrome in this study. In contrast, the significance of the association of the SNPs with exfoliation syndrome/glaucoma was relatively low when the PG group was used as a control comparison compared to when the NG or CT groups were used as control comparisons. Although the level of significance was relatively low, we found differences in frequencies between PG and NG for allele C of rs2165241 ($p=0.0233$) and the genotype CC of rs2165241 ($p=0.0294$). These may suggest the possible inclusion of late onset exfoliation syndrome in the PG group. Previously, a lack of association between *LOXL1* polymorphisms and primary open-angle glaucoma or primary angle-closure glaucoma were reported in Caucasian, African American, Ghanaian, and Indian populations [17,18]. We did

TABLE 5. COUNTS AND FREQUENCIES OF PHENOTYPE IN CASES (EX, EG, AND EX+EG) AND CONTROLS (CT, NG, PG+NG, CT+NG, AND CT+PG+NG)

Cases	EX		EG		EX+EG		CT		NG		PG+NG		CT+NG		CT+PG+NG		
	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	Count	Frequency	
TG	111	0.041	159	0.558	270	0.951											
GG	6	0.051	6	0.038	12	0.042											
GA	1	0.008	1	0.006	2	0.007											
Controls																	
TG	140	0.443	50	0.483	89	0.473											
GG	115	0.364	36	0.333	61	0.324											
GA	61	0.193	22	0.204	38	0.202											

EX, exfoliation syndrome; EG, exfoliation glaucoma; PG, primary open angle glaucoma aged 70 years or older; NG, normal tension glaucoma aged 70 years or older; and CT, cataract aged 70 years or older.

TABLE 6. P VALUES FOR HAPLOTYPE RS1048661/RS3825942 IN COMPARISONS BETWEEN CASES (EX, EG, AND EX+EG) AND CONTROLS (CT, NG, PG+NG, CT+NG, AND CT+PG+NG).

EX	Versus CT	Versus NG	Versus PG+NG	Versus CT+NG	Versus CT+PG+NG
TG	9.56x10 ⁻²¹	NA	9.56x10 ⁻²¹	6.19x10 ⁻¹⁷	1.27x10 ⁻²¹
GG	9.69x10 ⁻¹¹	NA	9.69x10 ⁻¹¹	1.76x10 ⁻¹⁰	1.43x10 ⁻¹⁰
GA	1.01x10 ⁻⁶	NA	1.01x10 ⁻⁶	5.63x10 ⁻⁷	5.63x10 ⁻⁷
EG	Versus CT	Versus NG	Versus PG+NG	Versus CT+NG	Versus CT+PG+NG
TG	1.83x10 ⁻²⁸	4.95x10 ⁻²¹	3.06x10 ⁻²³	9.77x10 ⁻³⁰	4.40x10 ⁻²⁰
GG	3.12x10 ⁻¹⁵	2.51x10 ⁻¹¹	4.82x10 ⁻¹²	2.60x10 ⁻¹⁵	3.70x10 ⁻¹⁵
GA	5.63x10 ⁻⁹	8.10x10 ⁻⁹	4.08x10 ⁻⁹	3.02x10 ⁻⁹	2.36x10 ⁻⁹
EX+EG	Versus CT	Versus NG	Versus PG+NG	Versus CT+NG	Versus CT+PG+NG
TG	1.23x10 ⁻⁴⁰	7.84x10 ⁻²⁹	1.24x10 ⁻³²	5.89x10 ⁻⁴³	8.27x10 ⁻⁴⁴
GG	5.93x10 ⁻²²	4.01x10 ⁻¹⁵	1.03x10 ⁻¹⁶	2.39x10 ⁻²²	2.83x10 ⁻²²
GA	1.17x10 ⁻¹³	3.98x10 ⁻¹³	9.33x10 ⁻¹⁴	3.68x10 ⁻¹⁴	2.25x10 ⁻¹⁴

Significant association was found between cases and controls. EX, exfoliation syndrome; EG, exfoliation glaucoma; PG, primary open angle glaucoma aged 70 years or older; NG, normal tension glaucoma aged 70 years or older; and CT, cataract aged 70 years or older. NA, not available due to a small number of samples.

not find any significant associations between any of the three SNPs with primary open-angle glaucoma, reconfirming the previous observations in our Japanese population. Most recently, one study from Japan reported a lack of association between *LOXLI* polymorphisms and primary open-angle glaucoma in two of three SNPs (e.g., rs1048661 and rs3825942) [16]. In addition to this, the current study suggests a lack of association between rs2165241 and primary open-angle glaucoma as well as a lack of association between any of the three SNPs with normal tension glaucoma.

In this study, none of the SNPs exhibited significant differences in their allelic and genotypic frequencies between exfoliation syndrome and glaucoma, suggesting that these SNPs are associated with exfoliation glaucoma through their association with exfoliation syndrome as reported previously [2]. Accordingly, our data suggest that additional genetic or environmental risk factors are associated with the development of exfoliation glaucoma among exfoliation syndrome patients. Further study is required to clarify these risks.

In summary, we have demonstrated significant associations of *LOXLI* variants with Japanese patients who have exfoliation syndrome/glaucoma. Compared to other populations, the risk alleles in rs1048661 and rs2165241 are unique in this population. The *LOXLI* association of exfoliation glaucoma is through the association of exfoliation syndrome. *LOXLI* lacked any association with primary open-angle glaucoma or normal tension glaucoma in this population. Additional genetic or environmental risk factors




Figure 1. Allelic frequencies of the three SNPs reported in previous studies and the current study. Allelic frequencies of three SNPs in cases (exfoliation syndrome and exfoliation glaucoma) and controls (cataract or normal population) that have been reported in literatures are summarized.

other than *LOXL1* are likely to be associated with an increase in exfoliation glaucoma among exfoliation syndrome patients.

ACKNOWLEDGMENTS

This study was supported in part by a grant to T.I. from the Ministry of Health, Labour and Welfare of Japan and by the National Hospital Organization of Japan.

REFERENCES

- Ritch R, Schlotzer-Schrehardt U. Exfoliation syndrome. *Surv Ophthalmol* 2001; 45:265-315. [PMID: 11166342]
- Thorleifsson G, Magnusson KP, Sulem P, Walters GB, Gudbjartsson DF, Stefansson H, Jonsson T, Jonasdottir A, Stefansson G, Masson G. Common sequence variants in the *LOXL1* gene confer susceptibility to exfoliation glaucoma. *Science* 2007; 317:1397-400. [PMID: 17690259]
- Fingert JH, Alward WL, Kwon YH, Wang K, Streb LM, Sheffield VC, Stone EM. *LOXL1* mutations are associated with exfoliation syndrome in patients from the midwestern United States. *Am J Ophthalmol* 2007; 144:974-5. [PMID: 18036875]
- Yang X, Zabriskie NA, Hau VS, Chen H, Tong Z, Gibbs D, Farhi P, Katz BJ, Luo L, Pearson E, Goldsmith J, Ma X, Kaminoh Y, Chen Y, Yu B, Zeng J, Zhang K, Yang Z. Genetic association of *LOXL1* gene variants and exfoliation glaucoma in a Utah cohort. *Cell Cycle* 2008; 7:521-4. [PMID: 18287813]
- Fan BJ, Pasquale L, Grosskreutz CL, Rhee D, Chen T, DeAngelis MM, Kim I, del Bono E, Miller JW, Li T. DNA sequence variants in the *LOXL1* gene are associated with pseudoexfoliation glaucoma in a U.S. clinic-based population with broad ethnic diversity. *BMC Med Genet* 2008; 9:5. [PMID: 18254956]
- Challa P, Schmidt S, Liu Y, Qin X, Vann RR, Gonzalez P, Allingham RR, Hauser MA. Analysis of *LOXL1* polymorphisms in a United States population with pseudoexfoliation glaucoma. *Mol Vis* 2008; 14:146-9. [PMID: 18334928]
- Aragon-Martin JA, Ritch R, Liebmann J, O'Brien C, Blaaw K, Mercieca F, Spiteri A, Cobb CJ, Damji KF, Tarkkanen A. Evaluation of *LOXL1* gene polymorphisms in exfoliation syndrome and exfoliation glaucoma. *Mol Vis* 2008; 14:533-41. [PMID: 18385788]
- Pasutto F, Krumbiegel M, Mardin CY, Paoli D, Lammer R, Weber BH, Kruse FE, Schlotzer-Schrehardt U, Reis A. Association of *LOXL1* Common Sequence Variants in German and Italian Patients with Pseudoexfoliation Syndrome and Pseudoexfoliation Glaucoma. *Invest Ophthalmol Vis Sci* 2008; 49:1459-63. [PMID: 18385063]
- Mossbock G, Renner W, Faschinger C, Schmut O, Wedrich A, Weger M. Lysyl oxidase-like protein 1 (*LOXL1*) gene polymorphisms and exfoliation glaucoma in a Central European population. *Mol Vis* 2008; 14:857-61. [PMID: 18483563]
- Ramprasad VL, George R, Soumitra N, Sharmila F, Vijaya L, Kumaramanickavel G. Association of non-synonymous single nucleotide polymorphisms in the *LOXL1* gene with pseudoexfoliation syndrome in India. *Mol Vis* 2008; 14:318-22. [PMID: 18334947]
- Hewitt AW, Sharma S, Burdon KP, Wang JJ, Baird PN, Dimasi DP, Mackey DA, Mitchell P, Craig JE. Ancestral *LOXL1* variants are associated with pseudoexfoliation in Caucasian Australians but with markedly lower penetrance than in Nordic people. *Hum Mol Genet* 2008; 17:710-6. [PMID: 18037624]
- Hayashi H, Gotoh N, Ueda Y, Nakanishi H, Yoshimura N. Lysyl oxidase-like 1 polymorphisms and exfoliation syndrome in the Japanese population. *Am J Ophthalmol* 2008; 145:582-5. [PMID: 18201684]
- Ozaki M, Lee KY, Vithana EN, Yong VH, Thalamuthu A, Mizoguchi T, Venkatraman A, Aung T. Association of *LOXL1* Gene Polymorphisms with Pseudoexfoliation in the Japanese. *Invest Ophthalmol Vis Sci* 2008; 49:3976-80. [PMID: 18450598]
- Mori K, Imai K, Matsuda A, Ikeda Y, Naruse S, Hitora-Takeshita H, Nakano M, Taniguchi T, Omi N, Tashiro K. *LOXL1* genetic polymorphisms are associated with exfoliation glaucoma in the Japanese population. *Mol Vis* 2008; 14:1037-40. [PMID: 18552979]
- Fuse N, Miyazawa A, Nakazawa T, Mengkegale M, Otomo T, Nishida K. Evaluation of *LOXL1* polymorphisms in eyes with exfoliation glaucoma in Japanese. *Mol Vis* 2008; 14:1338-43. [PMID: 18648524]
- Mabuchi F, Sakurada Y, Kashiwagi K, Yamagata Z, Iijima H, Tsukahara S. Lysyl oxidase-like 1 gene polymorphisms in Japanese patients with primary open angle glaucoma and exfoliation syndrome. *Mol Vis* 2008; 14:1303-8. [PMID: 18636115]
- Chakrabarti S, Rao KN, Kaur I, Parikh RS, Mandal AK, Chandrasekhar G, Thomas R. The *LOXL1* gene variations are not associated with primary open-angle and primary angle-closure glaucomas. *Invest Ophthalmol Vis Sci* 2008; 49:2343-7. [PMID: 18223248]
- Liu Y, Schmidt S, Qin X, Gibson JR, Hutchins K, Santiago-Turla C, Wiggs JL, Budenz DL, Akafo S, Challa P. Lack of Association between *LOXL1* variants and Primary Open-angle Glaucoma in Three Different Populations. *Invest Ophthalmol Vis Sci* 2008; 49:3465-8. [PMID: 18421074]
- Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 2005; 21:263-5. [PMID: 15297300]

The print version of this article was created on 21 October 2008. This reflects all typographical corrections and errata to the article through that date. Details of any changes may be found in the online version of the article.