

observed between hnRNPs A2/B1 and hnRNPs C1/C2.

## Discussion

Comparison of the different protein levels in the macula and peripheral retina was determined by proteomic analyses, and the results showed a number of quantitatively abundant proteins in the primate macula. One of the proteins was arrestin-C known to be specifically expressed in cone photoreceptors, a cell type highly concentrated in the macula (4). The identification of arrestin-C indicated that this proteomic approach was methodologically effective for the identification of macula enriched proteins. Another example is 3'(2'),5'-bisphosphate nucleotidase 1 previously identified by SAGE study was reported to be highly expressed in cone photoreceptors (7), and our results identified it as another macular enriched protein. Guanylate kinase (Table 2, M3), which catalyzes the phosphorylation of GMP(38) and is essential for recycling cGMP in the photoreceptors (39), was also identified as one of the macular enriched protein.

$\gamma$ -synuclein, a member of the synuclein family, was identified as a macular specific spot in the 2D-gel. Previous SAGE study also showed that  $\gamma$ -synuclein was highly expressed in the macula (6). Immunostaining showed that this protein was predominantly

located in the retinal ganglion cells and nerve fiber layer (Fig. 7D), which is also consistent with a previous study (40). These findings are in agreement with the high concentration of retinal ganglion cells in the macular area.  $\gamma$ -synuclein was initially identified as a breast cancer specific gene 1 by high throughput differential cDNA sequencing (33). This protein was expressed in peripheral nervous system, brain (41), and in ocular tissues (40).  $\gamma$ -Synuclein knockout mice are viable, fertile, with no abnormal behavior compared to wild type mice (42) with morphologically normal retina (43). The cellular function of  $\gamma$ -synuclein is still unknown, but an over-expression of synoretin, a bovine ortholog of  $\gamma$ -synuclein, activates the Elk-1 signal transduction pathway (44) and the chaperonic activity (43). Snyder et al. hypothesized that  $\gamma$ -synuclein is the regulator of 20S proteasome degradation and has a role in cellular response to oxidative stress because 20S ubiquitin-independent proteasomal pathway is responsible for the degradation of 70-80% of all mildly oxidized proteins (45). In the macula,  $\gamma$ -synuclein is likely to play an important role in the degradation of oxidized proteins.

Six macula specific mitochondria proteins were identified in this study. Mitochondria have been recognized as one of the major cellular generators of reactive oxygen species through electron transport chain reactions (55). Oxidative stress is believed to be important in the aging process and pathogenesis of diseases including AMD (56).

Photoreceptors densely localized in the macula have higher concentration of mitochondria with 2 to 3-fold greater oxygen consumption compared to the inner retina (46) and the distribution and morphological differences of mitochondria in the different regions of the retina have been reported (47). The macula may due to the higher concentration of cones exposed to more oxidative stress than peripheral retina.

Retinal abnormality of the Cu, Zn-SOD knockout mice (57) and Mn-SOD knockdown mice (58) mimic to patients with AMD. A higher demand of ATP production in cones than in rods has been suggested by the analysis of mitochondrial crista structure (59). An accumulation of 4977 bp deleted mitochondrial DNA (mtDNA<sup>4977</sup>), which is mainly caused by free radicals generated from mitochondria in the inner segments, has been shown to increase with age, and at higher concentration in the macula than in the peripheral retina (60). In our results, additional horizontal spots for Mn-SOD were observed in the 2D-western blot (Fig. 6, arrow), indicating a posttranslational modification specific to the macula. The phosphorylation of Mn-SOD by calcium ion has been demonstrated to inactivate enzyme activity (61). The activity of Mn-SOD is essential for the survival (62) and protection of the retina from oxidative stress (63). The presence of numerous phosphorylated forms of Mn-SOD in different region of the retina may explain the preferentially impairment of the macula under high oxidative stress levels.

Addition to these observation, the identified voltage-dependent anion-selective channel protein 1 (VDAC1) is located in the outer membrane of mitochondria (48, 49) and involved in cell volume regulation and apoptosis (X). Age-related maculopathy susceptibility 2 (*ARMS2/LOC387715*), a functionally unknown susceptibility gene for AMD (50-52), encodes a protein which is also localized to the outer membrane of the mitochondria (53, 54). Further functional study of VDAC1 and ARMS2 in correlation with AMD is notable.

E-FABP also plays a role as an antioxidant. 4-hydroxynonenal (4-HNE) is one of the aldehydes produced from lipid peroxidation. E-FABP is a molecular target for 4-HNE modification and functions as an antioxidant protein by scavenging reactive lipids through covalent modification of Cys-120 (34). 4-HNE modified E-FABP has been detected in the rat retina (64) and even higher expression levels in the primate macula to cope with the higher oxidative stress than the peripheral retina. In a transcriptional comparison study between macular and peripheral RPE, the expression of E-FABP was higher in the peripheral than macular RPE (9). Our immunostaining showed that E-FABP was predominantly located in the neural retina except the photoreceptors (Fig. 7E) which is consistent with the results of Kingma et al. (65). Our observation of a higher E-FABP expression in the macula is probably from the neural retina and not the RPE.

Four macula enriched hnRNPs were identified. hnRNPs are RNA-binding proteins predominantly located in the nucleus and function in splicing, transporting, and stabilizing mRNAs (66, 67). hnRNPs A2/B1 and hnRNPs C1/C2 were confirmed to be expressed at significantly higher levels in the macula by western blotting (Fig. 5G and 5H). These proteins were identified from multiple horizontal spots in the same gel indicating the possibility of phosphorylation. hnRNPs C1/C2 has been observed in several posttranslational phosphorylated forms and these forms are dependent on the duration of the oxidative stress (37). It also modulates the affinity of the mRNA binding to hnRNPs C1/C2 (68). The regional specificity of phosphorylation in this protein may affect the pathological process of macula-related diseases. For example, it has been reported that this protein stabilizes the mRNA of amyloid precursor protein (APP) and enhances translation of APP (69). APP and Amyloid beta ( $A\beta$ ) are found in drusen from AMD patients (X). Enrichment and posttranslational modification of hnRNPs C1/C2 may associate with  $A\beta$  accumulation. In addition to the hnRNPs C1/C2 phosphorylation, hnRNPs A2/B1 phosphorylation is involved in modifying its molecular function. In the central nerve system, phosphorylation of hnRNPs A2/B1 has been considered to be essential for myelination at the axon-glia contact site by mRNA transportation (70).

Tropomyosin1 a chain belongs to a multi-gene family of actin-binding proteins and 5

isoforms are listed in UniProtKB/Swiss-Prot database. In mammals, tropomyosin can give rise to many isoforms by splicing variants and specific isoforms are present in each tissue (35). Although the tropomyosin1a chain and other tropomyosins have been identified in photoreceptors (72), this is the first report to observe distribution of tropomyosin isoforms in macula and to identify brain isoform from photoreceptors. Anti-tropomyosin1a chain antibody TM311 was used for western blotting and immunohistochemistry. TM311 is a monoclonal antibody which detects broad range of alpha-, beta-, gamma- and delta- tropomyosin expressed in both muscle and non-muscle tissue except TMBR-3. By immunostaining, tropomyosin isoforms detected by TM311 were located around the choroidal vessel (Fig. 7G). MacDonald et al. demonstrated that endostatin, a cleavage fragment of collagen XVIII, functions as an anti-angiogenic factor (73) but it requires binding to tropomyosin to gain activity (74). Higher expression of tropomyosin1a chain in the macula has the potential to act as an anti-angiogenic factor for AMD by binding to endostatin.

Phosphorylation of the tropomyosin1a chain has also been described. Using human umbilical vascular endothelial cells, Houle et al. reported tropomyosin phosphorylation by oxidative stress through activation of the ERK pathway. Inhibition of the ERK pathway in the presence of oxidative stress resulted in membrane blebbing which then led to

damage of the vascular endothelium (71).

Spot number M4 corresponded to TMBr-3. TMBr-3 is expressed in all regions of the brain (75, 76) and was detected in the outer plexiform layer and photoreceptor inner segments (Fig. 7F) by double immunostaining with arrestin-C (Fig. 7B). The function of TMBr-3 in the photoreceptor inner segments is still unknown. Tropomyosin has a role in stabilizing actin filaments and regulates actin filaments in competition with other actin binding protein in non-muscle cells (77). Actin filaments located in the photoreceptor connecting cilia are considered the cytoskeletal core necessary for initiating new disk formation (78). Disk turnover is essential for maintenance of normal vision and consequently accumulation of lipofuscin in RPE cells by phagocytosis of disks has been linked to AMD (79). The function of TMBr-3 might involve disk formation by actin binding and rate of lipofuscin accumulation. Furthermore one of peptide fragment, which contains nitro tyrosine, in tropomyosin 1 alpha chain was identified from skeletal muscle (X). Nitration has been considered the potential for pathogenesis of many diseases (X). Brain type isoform also contains same amino acid sequence. In the retina, nitration of tropomyosin might be occurred and be associated with pathogenesis of AMD. Further studies are needed to understand the function of TMBr-3 in the photoreceptors.

## **Conclusions**

Proteomic analyses using 2D gel electrophoresis and LC-MS/MS were performed for comparison of regional differences in primate retina. Twenty-six proteins were identified as macula unique 2D-gel spots and 6 of these proteins were confirmed to be macula enriched by western blotting. In this study, brain type isoform of tropomyosin 1 alpha chain was first reported from retina. Furthermore, additional horizontal spots for Mn-SOD were specifically observed in the macula. Although some of these proteins are derived because of difference in major cell types between the two regions, identification of protein from macular enriched cells is important to understand the pathogenesis of AMD.

## **Acknowledgement**

This study was funded in part by a grant to TI from the Ministry of Health, Labour, and Welfare of Japan and by a grant to HO by the Ministry of Education, Sports, Science and Technology of Japan (KAKENHI 19791305).



## References

1. Provis, J. M., Penfold, P. L., Cornish, E. E., Sandercoe, T. M., and Madigan, M. C. (2005) Anatomy and development of the macula: specialisation and the vulnerability to macular degeneration. *Clin. Exp. Optom.* 88, 269-281
2. Snodderly, D. M., Brown, P. K., Delori, F. C., and Auran, J. D. (1984) The macular pigment. I. Absorbance spectra, localization, and discrimination from other yellow pigments in primate retinas. *Invest. Ophthalmol. Vis. Sci.* 25, 660-673
3. Snodderly, D. M., Auran, J. D., and Delori, F. C. (1984) The macular pigment. II. Spatial distribution in primate retinas. *Invest. Ophthalmol. Vis. Sci.* 25, 674-685
4. Curcio, C. A., Allen, K. A., Sloan, K. R., Lerea, C. L., Hurley, J. B., Klock, I. B., and Milam, A. H. (1991) Distribution and morphology of human cone photoreceptors stained with anti-blue opsin. *J. Comp. Neurol.* 312, 610-624
5. Hyman, L., and Neborsky, R. (2002) Risk factors for age-related macular degeneration: an update. *Curr. Opin. Ophthalmol.* 13, 171-175

6. Sharon, D., Blackshaw, S., Cepko, C. L., and Dryja, T. P. (2002) Profile of the genes expressed in the human peripheral retina, macula, and retinal pigment epithelium determined through serial analysis of gene expression (SAGE). *Proc. Natl. Acad. Sci. U S A* 99, 315-320
  
7. Bowes Rickman, C., Ebright, J. N., Zavodni, Z. J., Yu, L., Wang, T., Daiger, S. P., Wistow, G., Boon, K., and Hauser, M. A. (2006) Defining the human macula transcriptome and candidate retinal disease genes using EyeSAGE. *Invest. Ophthalmol. Vis. Sci.* 47, 2305-2316
  
8. Ishibashi, K., Tian, J., and Handa, J. T. (2004) Similarity of mRNA phenotypes of morphologically normal macular and peripheral retinal pigment epithelial cells in older human eyes. *Invest. Ophthalmol. Vis. Sci.* 45, 3291-3301
  
9. Radeke, M. J., Peterson, K. E., Johnson, L. V., and Anderson, D. H. (2007) Disease susceptibility of the human macula: differential gene transcription in the retinal pigmented epithelium/choroid. *Exp. Eye. Res.* 85, 366-380

10. van Soest, S. S., de Wit, G. M., Essing, A. H., ten Brink, J. B., Kamphuis, W., de Jong, P. T., and Bergen, A. A. (2007) Comparison of human retinal pigment epithelium gene expression in macula and periphery highlights potential topographic differences in Bruch's membrane. *Mol. Vis.* 13, 1608-1617

11. Nishizawa, Y., Komori, N., Usukura, J., Jackson, K. W., Tobin, S. L., and Matsumoto, H. (1999) Initiating ocular proteomics for cataloging bovine retinal proteins: microanalytical techniques permit the identification of proteins derived from a novel photoreceptor preparation. *Exp. Eye. Res.* 69, 195-212

12. Alge, C. S., Suppmann, S., Priglinger, S. G., Neubauer, A. S., May, C. A., Hauck, S., Welge-Lussen, U., Ueffing, M., and Kampik, A. (2003) Comparative proteome analysis of native differentiated and cultured dedifferentiated human RPE cells. *Invest. Ophthalmol. Vis. Sci.* 44, 3629-3641

13. Cavusoglu, N., Thierse, D., Mohand-Said, S., Chalmel, F., Poch, O., Van-Dorsseleer, A., Sahel, J. A., and Leveillard, T. (2003) Differential proteomic analysis of the mouse

retina: the induction of crystallin proteins by retinal degeneration in the rd1 mouse. *Mol.*

*Cell. Proteomics.* 2, 494-505

14. West, K. A., Yan, L., Shadrach, K., Sun, J., Hasan, A., Miyagi, M., Crabb, J. S., Hollyfield, J. G., Marmorstein, A. D., and Crabb, J. W. (2003) Protein database, human retinal pigment epithelium. *Mol. Cell. Proteomics.* 2, 37-49

15. Ethen, C. M., Reilly, C., Feng, X., Olsen, T. W., and Ferrington, D. A. (2006) The proteome of central and peripheral retina with progression of age-related macular degeneration. *Invest. Ophthalmol. Vis. Sci.* 47, 2280-2290

16. Stafford, T. J. (1974) Maculopathy in an elderly sub-human primate. *Mod. Probl. Ophthalmol.* 12, 214-219

17. El-Mofty, A., Gouras, P., Eisner, G., and Balazs, E. A. (1978) Macular degeneration in rhesus monkey (*Macaca mulatta*). *Exp. Eye. Res.* 27, 499-502

18. Stafford, T. J., Anness, S. H., and Fine, B. S. (1984) Spontaneous degenerative

maculopathy in the monkey. *Ophthalmology* 91, 513-521

19. Ishibashi, T., Sorgente, N., Patterson, R., and Ryan, S. J. (1986) Pathogenesis of drusen in the primate. *Invest. Ophthalmol. Vis. Sci.* 27, 184-193

20. Monaco, W. A., and Wormington, C. M. (1990) The rhesus monkey as an animal model for age-related maculopathy. *Optom. Vis. Sci.* 67, 532-537

21. Hope, G. M., Dawson, W. W., Engel, H. M., Ulshafer, R. J., Kessler, M. J., and Sherwood, M. B. (1992) A primate model for age related macular drusen. *Br. J. Ophthalmol.* 76, 11-16

22. Nicolas, M. G., Fujiki, K., Murayama, K., Suzuki, M. T., Mineki, R., Hayakawa, M., Yoshikawa, Y., Cho, F., and Kanai, A. (1996) Studies on the mechanism of early onset macular degeneration in cynomolgus (*Macaca fascicularis*) monkeys. I. Abnormal concentrations of two proteins in the retina. *Exp. Eye. Res.* 62, 211-219

23. Nicolas, M. G., Fujiki, K., Murayama, K., Suzuki, M. T., Shindo, N., Hotta, Y., Iwata, F.,

Fujimura, T., Yoshikawa, Y., Cho, F., and Kanai, A. (1996) Studies on the mechanism of early onset macular degeneration in cynomolgus monkeys. II. Suppression of metallothionein synthesis in the retina in oxidative stress. *Exp. Eye. Res.* 62, 399-408

24. Suzuki, M. T., Terao, K., and Yoshikawa, Y. (2003) Familial early onset macular degeneration in cynomolgus monkeys (*Macaca fascicularis*). *Primates* 44, 291-294

25. Umeda, S., Ayyagari, R., Suzuki, M. T., Ono, F., Iwata, F., Fujiki, K., Kanai, A., Takada, Y., Yoshikawa, Y., Tanaka, Y., and Iwata, T. (2003) Molecular cloning of ELOVL4 gene from cynomolgus monkey (*Macaca fascicularis*). *Exp. Anim.* 52, 129-135

26. Umeda, S., Ayyagari, R., Allikmets, R., Suzuki, M. T., Karoukis, A. J., Ambasudhan, R., Zernant, J., Okamoto, H., Ono, F., Terao, K., Mizota, A., Yoshikawa, Y., Tanaka, Y., and Iwata, T. (2005) Early-onset macular degeneration with drusen in a cynomolgus monkey (*Macaca fascicularis*) pedigree: exclusion of 13 candidate genes and loci. *Invest. Ophthalmol. Vis. Sci.* 46, 683-691

27. Umeda, S., Suzuki, M. T., Okamoto, H., Ono, F., Mizota, A., Terao, K., Yoshikawa, Y.,

Tanaka, Y., and Iwata, T. (2005) Molecular composition of drusen and possible involvement of anti-retinal autoimmunity in two different forms of macular degeneration in cynomolgus monkey (*Macaca fascicularis*). *FASEB. J.* 19, 1683-1685

28. Butt, R. H., Pfeifer, T. A., Delaney, A., Grigliatti, T. A., Tetzlaff, W. G., and Coorsen, J. R. (2007) Enabling coupled quantitative genomics and proteomics analyses from rat spinal cord samples. *Mol. Cell. Proteomics.* 6, 1574-1588

29. Hauck, S. M., Schoeffmann, S., Deeg, C. A., Gloeckner, C. J., Swiatek-de Lange, M., and Ueffing, M. (2005) Proteomic analysis of the porcine interphotoreceptor matrix. *Proteomics* 5, 3623-3636

30. Wang, Y. D., Wu, J. D., Jiang, Z. L., Wang, Y. B., Wang, X. H., Liu, C., and Tong, M. Q. (2007) Comparative proteome analysis of neural retinas from type 2 diabetic rats by two-dimensional electrophoresis. *Curr. Eye. Res.* 32, 891-901

31. Sakuma, H., Inana, G., Murakami, A., Higashide, T., and McLaren, M. J. (1996) Immunolocalization of X-arrestin in human cone photoreceptors. *FEBS. Lett.* 382,

32. Melov, S. (2000) Mitochondrial oxidative stress. Physiologic consequences and potential for a role in aging. *Ann. N. Y. Acad. Sci.* 908, 219-225

33. Ji, H., Liu, Y. E., Jia, T., Wang, M., Liu, J., Xiao, G., Joseph, B. K., Rosen, C., and Shi, Y. E. (1997) Identification of a breast cancer-specific gene, BCSG1, by direct differential cDNA sequencing. *Cancer. Res.* 57, 759-764

34. Bennaars-Eiden, A., Higgins, L., Hertzell, A. V., Kapphahn, R. J., Ferrington, D. A., and Bernlohr, D. A. (2002) Covalent modification of epithelial fatty acid-binding protein by 4-hydroxynonenal in vitro and in vivo. Evidence for a role in antioxidant biology. *J. Biol. Chem.* 277, 50693-50702

35. Schevzov, G., Vrhovski, B., Bryce, N. S., Elmir, S., Qiu, M. R., O'Neill G. M., Yang, N., Verrills, N. M., Kavallaris, M., and Gunning, P. W. (2005) Tissue-specific tropomyosin isoform composition. *J. Histochem. Cytochem.* 53, 557-570



36. Sueoka, E., Goto, Y., Sueoka, N., Kai, Y., Kozu, T., and Fujiki, H. (1999)

Heterogeneous nuclear ribonucleoprotein B1 as a new marker of early detection for human lung cancers. *Cancer Res.* 59, 1404-1407

37. Stone, J. R., and Collins, T. (2002) Rapid phosphorylation of heterogeneous nuclear ribonucleoprotein C1/C2 in response to physiologic levels of hydrogen peroxide in human endothelial cells. *J. Biol. Chem.* 277, 15621-15628

38. Hall, S. W., and Kuhn, H. (1986) Purification and properties of guanylate kinase from bovine retinas and rod outer segments. *Eur. J. Biochem.* 161, 551-556

39. Farber, D. B. (1995) From mice to men: the cyclic GMP phosphodiesterase gene in vision and disease. The Proctor Lecture. *Invest. Ophthalmol. Vis. Sci.* 36, 263-275

40. Surguchov, A., McMahan, B., Masliah, E., and Surgucheva, I. (2001) Synucleins in ocular tissues. *J. Neurosci. Res.* 65, 68-77

41. George, J. M. (2002) The synucleins. *Genome Biol.* 3, REVIEWS3002

42. Ninkina, N., Papachroni, K., Robertson, D. C., Schmidt, O., Delaney, L., O'Neill, F., Court, F., Rosenthal, A., Fleetwood-Walker, S. M., Davies, A. M., and Buchman, V. L. (2003) Neurons expressing the highest levels of gamma-synuclein are unaffected by targeted inactivation of the gene. *Mol. Cell. Biol.* 23, 8233-8245

43. Surgucheva, I., Ninkina, N., Buchman, V. L., Grasing, K., and Surguchov, A. (2005) Protein aggregation in retinal cells and approaches to cell protection. *Cell. Mol. Neurobiol.* 25, 1051-1066

44. Surguchov, A., Surgucheva, I., Solessio, E., and Baehr, W. (1999) Synoretin--A new protein belonging to the synuclein family. *Mol. Cell. Neurosci.* 13, 95-103

45. Snyder, H., Mensah, K., Hsu, C., Hashimoto, M., Surgucheva, I. G., Festoff, B., Surguchov, A., Masliah, E., Matouschek, A., and Wolozin, B. (2005) beta-Synuclein reduces proteasomal inhibition by alpha-synuclein but not gamma-synuclein. *J. Biol. Chem.* 280, 7562-7569

46. Medrano, C. J., and Fox, D. A. (1995) Oxygen consumption in the rat outer and inner retina: light- and pharmacologically-induced inhibition. *Exp. Eye. Res.* 61, 273-284
47. Hoang, Q. V., Linsenmeier, R. A., Chung, C. K., and Curcio, C. A. (2002) Photoreceptor inner segments in monkey and human retina: mitochondrial density, optics, and regional variation. *Vis. Neurosci.* 19, 395-407
48. Colombini, M. (1989) Voltage gating in the mitochondrial channel, VDAC. *J. Membr. Biol.* 111, 103-111
49. Adams, V., Griffin, L., Towbin, J., Gelb, B., Worley, K., and McCabe, E. R. (1991) Porin interaction with hexokinase and glycerol kinase: metabolic microcompartmentation at the outer mitochondrial membrane. *Biochem. Med. Metab. Biol.* 45, 271-291
50. Rivera, A., Fisher, S. A., Fritsche, L. G., Keilhauer, C. N., Lichtner, P., Meitinger, T., and Weber, B. H. (2005) Hypothetical LOC387715 is a second major susceptibility gene for age-related macular degeneration, contributing independently of complement factor H to disease risk. *Hum. Mol. Genet.* 14, 3227-3236

51. Dewan, A., Liu, M., Hartman, S., Zhang, S. S., Liu, D. T., Zhao, C., Tam, P. O., Chan, W. M., Lam, D. S., Snyder, M., Barnstable, C., Pang, C. P., and Hoh, J. (2006) HTRA1 promoter polymorphism in wet age-related macular degeneration. *Science* 314, 989-992
52. Yang, Z., Camp, N. J., Sun, H., Tong, Z., Gibbs, D., Cameron, D. J., Chen, H., Zhao, Y., Pearson, E., Li, X., Chien, J., Dewan, A., Harmon, J., Bernstein, P. S., Shridhar, V., Zabriskie, N. A., Hoh, J., Howes, K., and Zhang, K. (2006) A variant of the HTRA1 gene increases susceptibility to age-related macular degeneration. *Science* 314, 992-993
53. Kanda, A., Chen, W., Othman, M., Branham, K. E., Brooks, M., Khanna, R., He, S., Lyons, R., Abecasis, G. R., and Swaroop, A. (2007) A variant of mitochondrial protein LOC387715/ARMS2, not HTRA1, is strongly associated with age-related macular degeneration. *Proc. Natl. Acad. Sci. U S A* 104, 16227-16232
54. Fritsche, L. G., Loenhardt, T., Janssen, A., Fisher, S. A., Rivera, A., Keilhauer, C. N., and Weber, B. H. (2008) Age-related macular degeneration is associated with an unstable ARMS2 (LOC387715) mRNA. *Nat. Genet.* 40, 892-896