

Current Topics in Innate Immunity

Edited by

JOHN D. LAMBRIS

Department of Pathology & Laboratory Medicine

University of Pennsylvania

Philadelphia, Pennsylvania

 Springer

 Vegean
Conferences

Editor:

John D. Lambris, Ph.D.
University of Pennsylvania
Philadelphia, PA 19104
www.lambris.net

Library of Congress Control Number: 2007926254

Proceedings from the 4th International Conference on Innate Immunity held in Corfu, Greece,
June 4-9, 2006.

ISBN 978-0-387-71765-4

e-ISBN 978-0-387-71767-8



Printed on acid-free paper.

© 2007 Springer Science+Business Media, LLC

All rights reserved. This work may not be translated or copied in whole or in part without the written permission of the publisher (Springer Science+Business Media, LLC, 233 Spring Street, New York, NY 10013, USA), except for brief excerpts in connection with reviews or scholarly analysis. Use in connection with any form of information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed is forbidden.

The use in this publication of trade names, trademarks, service marks, and similar terms, even if they are not identified as such, is not to be taken as an expression of opinion as to whether or not they are subject to proprietary rights.

10 9 8 7 6 5 4 3 2 1

springer.com

Preface

Innate Immunity has long been regarded as the non-specific arm of immune response, acting immediately and in a generic way, to defend the host from infections. In the post genomic era, our knowledge of the innate immune system is enriched by findings on the specificity of innate immune reactions as well as to novel functions that do not strictly correlate with immunological defense and surveillance, immune modulation or inflammation. Several studies indicate that molecules involved in innate immunity exert functions that are either more complex than previously thought, or go well beyond the innate immune character of the system.

The advent of high-throughput platforms for genome and proteome-wide profiling, together with the enormous amount of raw genetic information that has accumulated in the databases, have stirred new expectations in biomedical research. They have led scientists to revisit established biological systems from a global and integrative perspective. Innate Immunity research is now faced with the challenge of trying to integrate isolated biochemical pathways into complex gene and protein regulatory circuits. In this respect, scientists from around the world convened at the 4th International Conference on Innate Immunity (June 4 - 9, 2006), in Corfu, Greece to discuss recent advances in this fast evolving field. This volume represents a collection of topics on natural killer cells, mast cells, phagocytes, toll like receptors, complement, host defense in plants and invertebrates, evasion strategies of microorganisms, pathophysiology, protein structures, design of therapeutics, and experimental approaches discussed during the conference.

I am grateful to the contributing authors for the time and effort they have devoted to writing, what I consider exceptionally informative chapters in a book that will have a significant impact on the Innate Immunity field. I am grateful to Rodanthi Lambris, for her assistance in formatting the text. I also gratefully acknowledge the generous help provided by Dimitrios Lambris in managing the organization of this meeting. Finally, I also thank Andrea Macaluso and Lisa Tenaglia of Springer Publishers for their supervision in this book's production.

John D. Lambris, Ph.D.

18. COMPLEMENT ACTIVATION OF DRUSEN IN PRIMATE MODEL (<i>MACACA FASCICULARIS</i>) FOR AGE-RELATED MACULAR DEGENERATION	251
Takeshi Iwata	
1 Introduction	251
2 Introduction of AMD	252
3 Genetics of AMD	252
4 Biochemistry of AMD	253
5 Primate Model for AMD	254
6 Mouse Model for AMD	256
7 Acknowledgements	257
References	258
19. EXPLORING THE COMPLEMENT INTERACTION NETWORK USING SURFACE PLASMON RESONANCE	260
Daniel Ricklin and John D. Lambris	
1 Introduction	260
2 Surface Plasmon Resonance – The Key to Kinetic Constants	260
3 Applications in Complement Research	264
3.1 Making connections – old and new ones	266
3.2 The good and evil side of complement: interaction with pathogens	268
3.3 The complexity of complex formation	270
3.4 Developing therapeutic interventions	272
4 Conclusions & Perspectives	273
References	274
20. GLYCOSYLATION AS A TARGET FOR RECOGNITION OF INFLUENZA VIRUSES BY THE INNATE IMMUNE SYSTEM	279
Patrick C. Reading, Michelle D. Tate, Danielle L. Pickett, and Andrew G. Brooks	
1 Introduction	279
2 Influenza Viruses	279
3 Envelope Glycoproteins of Influenza Viruses	280
4 Glycosylation of Influenza Virus HA and NA	281
4.1 Functions of glycosylation of influenza HA and NA glycoproteins	281
5 Mammalian C-Type Lectins	282

Complement Activation of Drusen in Primate Model (*Macaca fascicularis*) for Age-Related Macular Degeneration

Takeshi Iwata

Takeshi Iwata, National Institute of Sensory Organs, National Hospital Organization
Tokyo Medical Center, 2-5-1 Higashigaoka, Meguro-ku, Tokyo 152-8902 Japan

1 Introduction

Dysfunction of the visual system can alter normal human life style and lower quality of life. The most prevalent causes of visual impairment worldwide are cataracts, glaucoma, and age-related macular degeneration (AMD). These eye diseases are responsible for 69% of blindness globally. Although cataracts are the leading cause of blindness worldwide, recent advances in cataract surgery has significantly reduced the visual impairments caused by cataracts especially in developed countries. The most prevalent eye disease for elderly Europeans and Americans is AMD. This degenerative disease progresses from retinal deposits called drusen to neovascularization and retinal hemorrhages resulting in irreversible loss of central vision. In spite of the high incidence of AMD, a limited amount of information is available on the underlying pathological mechanisms causing these diseases. Obtaining tissues from the AMD donors is often difficult, and even when obtained, they are usually collected many hours or even days after death. Because of limitation for human tissue, the availability of animal models is becomes valuable because they can be used to investigate the molecular mechanisms of the disease and to test new therapeutic intervention.

The retina is composed of nine layers of neural and glial cells that are arranged concentrically at the posterior pole of the eye. Incoming light is focused on the central area of the retina called the fovea which is located in the center of the macula. In humans, the average size of the macula is only 6 mm in diameter. The outer surface of the retina is covered by a monolayer of retinal pigment epithelial (RPE) cells which forms a diffusion barrier between the neural retina and the choroidal blood supply. The RPE regulates the transport of proteins to the retina, and controls the hydration

and ionic composition of the subretinal space. The physiological condition of the RPE is closely associated with the pathogenesis of AMD.

2 Introduction of AMD

AMD is a blinding disorder characterized by a marked decrease in central vision associated with RPE atrophy with or without choroidal neovascularization (CNV). Many factors including genetic, behavioral, and environmental, are involved in this disease. AMD is characterized by the degeneration of cone photoreceptors in the foveal region of the retina resulting in a decrease of central visual acuity. The progressive impairment of the retinal pigment epithelial (RPE) cells, and damage to Bruch's membrane and choriocapillaris results in retinal atrophy and photoreceptor dysfunction. In some cases, CNV develops, and the new vessels penetrate Bruch's membrane and pass into the subretinal space.

Two types of AMD are recognized; the non-neovascular type is called the dry-type AMD and includes more than 80% of the cases, and the neovascular type is called the wet-type AMD which is progressive with a higher probability of blindness. The prevalence of AMD differs considerably among the different ethnic groups, but the incidence increases with age in all groups. A lower prevalence of AMD has been reported in individuals of African ancestry than of Anglo-Saxon ancestry. Other risk factors for AMD are cigarette smoking, obesity, hypertension, and atherosclerosis.

3 Genetics of AMD

Epidemiological studies have shown that genetic factor play critical role for AMD. Twin studies have previously shown a higher concordance for AMD in monozygotic twins than in dizygotic twins (Heiba, Elston, Klein, and Klein 1994; Seddon, Ajani, and Mitchell 1997; Hammond, Webster, Snieder, Bird, Gilbert, and Spector 2002). In addition, first degree relatives of individuals with AMD have higher incidence of AMD over individuals without a family history of AMD. Genetic segregation studies have also shown a genetic effect that accounts for approximately 60% of AMD with a single major gene accounting for about 55% of the risk of developing AMD. Previous data have suggested that the etiology of AMD has a significant genetic component. Only a small proportion of the families with AMD show Mendelian inheritance, and the majority of the individuals inherit AMD in a complex multi-gene pattern. With the help of the haplotype marker project (HapMap Project), genome wide scanning has identified at least 13 loci linked to AMD on different chromosomes (Iyengar, Song, Klein, Klein, Schick, Humphrey, Millard, Liptak, Russo, Jun, Lee, Fijal, and Elston 2004; Schick, Iyengar, Klein, Klein, Reading, Liptak, Millard, Lee, Tomany, Moore, Fijal, and Elston 2003; Majewski, Schultz, Weleber, Schain, Edwards, Matise, Acott, Ott, and Klein 2003). Recently, a polymorphism of complement factor H (CFH) gene (*Y402H*) was shown to be associated with an increased risk for AMD (Klein, Zeiss, Chew, Tsai, Sackler, Haynes, Henning, SanGiovanni, Mane, Mayne, Bracken, Ferris, Ott, Barnstable, and Hoh 2005; Edwards, Ritter, Abel, Manning, Panhuysen, and

Farrer 2005; Haines, Hauser, Schmidt, Scott, Olson, Gallins, Spencer, Kwan, Nouredine, Gilbert, Schnetz-Boutaud, Agarwal, Postel, and Pericak-Vance 2005; Hageman, Anderson, Johnson, Hancox, Taiber, Hardisty, Hageman, Stockman, Borchardt, Gehrs, Smith, Silvestri, Russell, Klaver, Barbazetto, Chang, Yannuzzi, Barile, Merriam, Smith, Olsh, Bergeron, Zernant, Merriam, Gold, Dean, and Allikmets 2005).

These results were confirmed in many of the countries with large Caucasian populations but not in Japan (Okamoto, Umeda, Obazawa, Minami, Noda, Mizota, Honda, Tanaka, Koyama, Takagi, Sakamoto, Saito, Miyake, and Iwata 2006; Gotoh, Yamada, Hiratani, Renault, Kuroiwa, Monet, Toyoda, Chida, Mandai, Otani, Yoshimura, and Matsuda 2006). This gene is located on chromosome 1q25-31 where one of the candidate loci was identified by linkage studies. Another recent study reported that a haplotype association of tandemly located complement 2 and factor B was protective for AMD (Gold, Merriam, Zernant, Hancox, Taiber, Gehrs, Cramer, Neel, Bergeron, Barile, Smith, AMD Genetics Clinical Study Group, Hageman, Dean, Allikmets 2006). HTRA1, a serine protease 11 was recently discovered to be strongly associated with AMD. Unlike the CFH, our study shows strong association with this gene for Japanese AMD patients (Yang, Camp, Sun, Tong, Gibbs, Cameron, Chen, Zhao, Pearson, Li, Chien, Dewan, Harmon, Bernstein, Shridhar, Zabriskie, Hoh, Howes, and Zhang 2006; Dewan, Liu, Hartman, Zhang, Liu, Zhao, Tam, Chan, Lam, Snyder, Barnstable, Pang, and Hoh 2006).

4 Biochemistry of AMD

The early stage of the dry type AMD is characterized by a thickening of Bruch's membrane, aggregation of pigment granules, and increasing numbers of drusen. The thickening of Bruch's membrane obstructs its function as a 'barrier' between the choroid and the RPE that protects the neural retina from the choriocapillary. Drusen are small yellowish-white deposits that are composed of lipids, proteins, glycoproteins, and glycosaminoglycans. They accumulate in the extracellular space and the inner aspects of Bruch's membrane. Drusen are not directly associated with visual loss but represent a risk factor for both the dry-type and wet-type AMD. The classification of hard and soft drusen is based on their size, shape, and color; hard drusen are yellowish with diameters $<50 \mu\text{m}$ and are found in eyes that are less likely to progress to advanced stages of the disease, while soft drusen are darker yellow and larger in size, and are found in eyes more likely to progress to more advanced stages of AMD. A small percentage of dry-type AMD patients progress to the late stage of the wet-type AMD that is characterized by geographic atrophy or detachment of RPE and the development of CNV in the macular region. The presence of a CNV is the factor that most damages the neural retina because the newly developed vessels grow from the choriocapillaris through Bruch's membrane and extend laterally through the RPE cell layer (classic CNV) or extend between the inner Bruch's membrane and RPE (occult CNV). In advanced stages of AMD, the CNV and fluid leaked into the subretinal or intraretinal regions leads to cell death and retinal detachment.

Recent analyses of the progression of drusen have provided important clues that help understand the molecular pathology of AMD. Using both immunohistochemistry

and proteomic techniques, the materials in drusen were found to be composed of molecules that mediate inflammatory and immune processes (Russell, Mullins, Schneider, and Hageman 2000; Mullins, Russell, Anderson, and Hageman 2000). These molecules include components of the complement pathway and modulators of complement activation, viz., vitronectin, clusterin, membrane cofactor protein, and complement receptor-1. In addition, molecules triggering inflammation, amyloid P component, α 1-antitrypsin, and apolipoprotein E, were identified in drusen. Cellular debris from macrophages, RPE cells, and choroidal dendritic cells has also been identified in drusen. Additional proteins such as crystallins, EEFMP1, and amyloid-beta have been also found in drusen. The presence of immunoreactive proteins and the oxidative modifications of many proteins in drusen imply that both oxidation and immune functions are involved in the pathogenesis of AMD. These findings suggest that complement activation triggers innate immune responses in the subretinal space. The co-distribution of IgG and terminal complement complexes in drusen indicate that immune responses that directly target antigens in retinal cells might also be occurring. Anti-retinal autoantibodies have been reported in a number of ocular disorders, e.g., macular degeneration in an aged monkey model.

5 Primate Model for AMD

Over the past few years, genetic engineering techniques have generated a number of animal models of AMD in mice, rats, rabbits, pigs, and dogs (Chader 2002). However in mammals, a well-defined fovea is found only in primates (humans and monkeys), and a search for a monkey line affected with macular degeneration has been persistent for a long time. A monkey with macular degeneration was first described by Stafford et al in 1974. They reported that 6.6 % of the elderly monkeys they examined showed pigmentary disorders and drusen-like spots (Stafford, Anness, and Fine 1984). El-Mofty et al reported that the incidence of maculopathy was 50% in a colony of rhesus monkeys at the Caribbean Primate Research Center of the University of Puerto Rico (El-Mofty, Gouras, Eisner, and Balazs 1978). At the Tsukuba Primate Research Center (Tsukuba City, Japan), Suzuki et al found a single cynomolgus monkey (*Macaca fascicularis*) (Suzuki Monkeys) in 1986 with a large number of small drusen around the macular region (Nicolas, Fujiki, Murayama, Suzuki, Mineki, Hayakawa, Yoshikawa, Cho, Kanai 1996; Nicolas, Fujiki, Murayama, Suzuki, Shindo, Hotta, Iwata, Fujimura, Yoshikawa, Cho, Kanai 1996; Suzuki, Terao, and Yoshikawa 2003). This single affected monkey has multiplied to a large pedigree of more than 65 affected and 210 unaffected monkeys. Drusen were observed in the macular region as early as one year after birth, and the numbers increased and spread toward the peripheral retina throughout life. No histological abnormalities have been found in the retina, retinal vessels, or choroidal vasculatures of the eyes with drusen. However, abnormality in electroretinogram (ERG) were observed in sever case showing dysfunction of the macula.

Immunohistochemical and proteomic analyses of the drusen from these monkeys showed that the drusen were very similar to those in other monkeys with aged




Fig. 1. Fundus photograph of both eyes of Suzuki Monkey showing accumulation of drusen (white spot) around the macular region.

macular degeneration sporadically found in older monkeys and also with human drusen (Umeda, Ayyagari, Allikmets, Suzuki, Karoukis, Ambasadhan, Zernant, Okamoto, Ono, Terao, Mizota, Yoshikawa, Tanaka, and Iwata 2005; Umeda, Suzuki, Okamoto, Ono, Mizota, Terao, Yoshikawa, Tanaka, and Iwata 2005; Ambati, Anand, Fernandez, Sakurai, Lynn, Kuziel, Rollins, and Ambati 2003). These observations have shown that the Suzuki Monkeys produce drusen that are biochemically similar to those in human AMD patients, but the development of the drusen occurs at an accelerated rate. More than 240 loci are being investigated to try to identify the disease causing gene and to understand the biological pathways leading to complement activation. Simultaneously, we have been studying a colony of aged monkeys which develop drusen after 15 years of birth.

Drusen components of these sporadically found affected monkeys were compared with human and Suzuki Monkeys by classical immunohistochemical techniques and by proteome analysis using mass spectrometer. Significant finding was that drusen contained protein molecules that mediate inflammatory and immune processes. These include immunoglobulins, components of complement pathway, and modulators for complement activation (e.g., vitronectin, clusterin, membrane cofactor protein, and complement receptor-1), molecules involved in the acute-phase response to inflammation (e.g., amyloid P component, α 1-antitrypsin, and apolipoprotein E), major histocompatibility complex class II antigens, and HLA-DR antigens (Umeda et al. 2005). Cellular components have also been identified in drusen, including RPE debris, lipofuscin, and melanin, as well as processes of choroidal dendritic cells, which are felt to contribute to the inflammatory response. In addition to immune components, a number of other proteins were found in drusen. These appear to be vitronectin, clusterin, TIMP-3, serum amyloid P component, apolipoprotein E, IgG, Factor X, crystallins, EEFMP1, and amyloid-beta. The presence of immunoreactive proteins and oxidative modified proteins implicate both oxidation and immune functions in the pathogenesis of AMD.




Fig. 2. Retinal histological section of affected Suzuki Monkey showing the accumulation of drusen between the retinal pigment epithelium and Bruch's membrane.

The eyes of monkey are structurally similar to human eyes which make them extremely valuable for AMD studies. However, there are limitations in using this species over other laboratory animals. Monkeys have a relatively longer life span, have a longer gestation period, have a lower birth numbers resulting in a slower expansion of the pedigree, more difficult to genetically manipulate, and the maintenance cost is high. In the other laboratory animals, the differences in the eye structure, lack of a fovea, and a low cone/rod ratio compared to humans have been considered to be a disadvantage for using them as AMD models. However, they are easier to manipulate genetically and easier and less expensive to maintain. This has made the development of a mouse model of AMD very attractive, and a number of mouse AMD models have been reported recently.

6 Mouse Model for AMD

The mouse model described by Ambati et al is deficient either in monocyte chemoattractant protein-1 or its cognate C-C chemokine receptor-2. These mice were found to develop the cardinal features of AMD including accumulation of lipofuscin

in drusen beneath the RPE, photoreceptor atrophy, and CNV (Ambati et al. 2003). An impairment of macrophage recruitment allowed the accumulation of C5a and IgG, which leads to the production of vascular endothelial growth factor by the RPE cells and the development of CNVs. Another mouse model that has three known AMD risk factors: age, high fat cholesterol rich diet, and expression of human apolipoprotein E (apoE2, apoE3, apoE4) has been developed (Malek, Johnson, Mace, Saloupis, Schmechel, Rickman, Toth, Sullivan, and Bowes Rickman 2005). ApoE4-deficient mice are severely affected showing diffuse subretinal pigment epithelial deposits, drusen, thickened Bruch's membrane, and atrophy, hypopigmentation, and hyperpigmentation of the RPE.

Oxidative stress has long been linked to the pathogenesis of AMD. Imamura et al reported a Cu, Zn-superoxide dismutase (SOD1)-deficient mice that had features typical of AMD in human. Senescent Sod1 (-/-) mice had drusen, thickened Bruch's membrane, and choroidal neovascularization (Imamura, Noda, Hashizume, Shinoda, Yamaguchi, Uchiyama, Shimizu, Mizushima, Shirasawa, and Tsubota 2006). The number of drusen increased with age and also after exposure of young Sod1 (-/-) mice to excess light. The retinal pigment epithelial cells of Sod1 (-/-) mice showed oxidative damage, and their beta-catenin-mediated cellular integrity was disrupted. These findings suggested that oxidative stress may affect the junctional proteins necessary for the barrier integrity of the RPE. These observations strongly suggested that oxidative stress may play a major role in AMD.

The complement components, C3a and C5a, are present in drusen, and were observed in Bruch's membrane of a laser-induced CNV mice model. Neutralization of C3a or C5a by antibody or by blockade of their receptors by a complement inhibitor significantly reduced the CNV (Nozaki, Raisler, Sakurai, Sarma, Barnum, Lambris, Chen, Zhang, Ambati, Baffi, and Ambati 2006). These observations revealed a role for immunological mechanisms for the angiogenesis and provided evidence for future therapeutic strategies for AMD. Although the pathology of AMD is pronounced in the macula area, it is not confined to this region. Characteristics of human AMD such as thickening of Bruch's membrane, accumulation of drusen, and CNV have been observed in mouse models. Nevertheless, the primate model will still be the choice for AMD studies, especially at the stage when new therapeutic methods are tested and evaluated for the first time. However, it would be wise and more productive to study both primate and mouse models in AMD research. This will be necessary to learn the mechanisms underlying the disease and to identify clinical and molecular markers for the early stages of AMD. The findings from these studies will provide critical information needed to develop therapies for AMD.

7 Acknowledgements

This work was supported by the research grant from the Ministry of Health, Labour and Welfare of Japan.

References

- Ambati, J., Anand, A., Fernandez, S., Sakurai, E., Lynn, B. C., Kuziel, W. A., Rollins, B. J., Ambati, B. K. (2003) An animal model of age-related macular degeneration in senescent Ccl-2- or Ccr-2-deficient mice. *Nat. Med.* 9, 1390-1397.
- Chader, G. J. (2002) Animal models in research on retinal degenerations: Past progress and future hope. *Vision Res* 42, 393-399.
- Dewan, A., Liu, M., Hartman, S., Zhang, S. S., Liu, D. T., Zhao, C., Tam, P. O., Chan, W. M., Lam, D. S., Snyder, M., Barnstable, C., Pang, C. P., Hoh, J. (2006) HTRA1 promoter polymorphism in wet age-related macular degeneration. *Science* 314, 989-992.
- Edwards, A. O., Ritter, R. 3rd., Abel, K. J., Manning, A., Panhuysen, C., Farrer, L. A. (2005) Complement factor H polymorphism and age-related macular degeneration. *Science* 308, 421-424.
- El-Mofty, A., Gouras, P., Eisner, G., Balazs, E. A. (1978) Macular degeneration in rhesus monkey (*Macaca mulatta*). *Exp. Eye Res.* 27, 499-502.
- Gold, B., Merriam, J. E., Zernant, J., Hancox, L. S., Taiber, A. J., Gehrs, K., Cramer, K., Neel, J., Bergeron, J., Barile, G. R., Smith, R. T. (2006) AMD Genetics Clinical Study Group; G. S. Hageman, M. Dean, R. Allikmets. Variation in factor B (BF) and complement component 2 (C2) genes is associated with age-related macular degeneration. *Nat. Genet.* 38, 458-462.
- Gotoh, N., Yamada, R., Hiratani, H., Renault, V., Kuroiwa, S., Monet, M., Toyoda, S., Chida, S., Mandai, M., Otani, A., Yoshimura, N., Matsuda, F. (2006) No association between complement factor H gene polymorphism and exudative age-related macular degeneration in Japanese. *Hum. Genet.* 120, 139-143.
- Hageman, G. S., Anderson, D. H., Johnson, L. V., Hancox, L. S., Taiber, A. J., Hardisty, L. I., Hageman, J. L., Stockman, H. A., Borchardt, J. D., Gehrs, K. M., Smith, R. J., Silvestri, G., Russell, S. R., Klaver, C. C., Barbazetto, I., Chang, S., Yannuzzi, L. A., Barile, G. R., Merriam, J. C., Smith, R. T., Olsh, A. K., Bergeron, J., Zernant, J., Merriam, J. E., Gold, B., Dean, M., Allikmets, R. (2005) A common haplotype in the complement regulatory gene factor H (HF1/CFH) predisposes individuals to age-related macular degeneration. *Proc. Natl. Acad. Sci. U S A* 102, 7227-7232.
- Haines, J. L., Hauser, M. A., Schmidt, S., Scott, W. K., Olson, L. M., Gallins, P., Spencer, K. L., Kwan, S. Y., Noureddine, M., Gilbert, J. R., Schetz-Boutaud, N., Agarwal, A., Postel, E. A., Pericak-Vance, M. A. (2005) Complement factor H variant increases the risk of age-related macular degeneration. *Science* 308, 419-421.
- Hammond, C. J., Webster, A. R., Snieder, H., Bird, A. C., Gilbert, C. E., Spector, T. D. (2002) Genetic influence on early age-related maculopathy: A twin study. *Ophthalmology* 109, 730-736.
- Heiba, I. M., Elston, R. C., Klein, B. E., Klein, R. (1994) Sibling correlations and segregation analysis of age-related maculopathy: The Beaver Dam Eye Study. *Genet. Epidemiol.* 11, 51-67.
- Imamura, Y., Noda, S., Hashizume, K., Shinoda, K., Yamaguchi, M., Uchiyama, S., Shimizu, T., Mizushima, Y., Shirasawa, T., Tsubota, K. (2006) Drusen, choroidal neovascularization, and retinal pigment epithelium dysfunction in SOD1-deficient mice: a model of age-related macular degeneration. *Proc Natl Acad Sci U S A* 103, 11282-11287.
- Iyengar, S. K., Song, D., Klein, B. E., Klein, R., Schick, J. H., Humphrey, J., Millard, C., Liptak, R., Russo, K., Jun, G., Lee, K. E., Fijal, B., Elston, R. C. (2004) Dissection of genomewide-scan data in extended families reveals a major locus and oligogenic susceptibility for age-related macular degeneration. *Am. J. Hum. Genet.* 74, 20-39.
- Klein, R. J., Zeiss, C., Chew, E. Y., Tsai, J. Y., Sackler, R. S., Haynes, C., Henning, A. K., SanGiovanni, J. P., Mane, S. M., Mayne, S. T., Bracken, M. B., Ferris, F. L., Ott, J., Barnstable, C., Hoh, J. (2005) Complement factor H polymorphism in age-related macular degeneration. *Science* 308, 385-389.
- Schick, J. H., Iyengar, S. K., Klein, B. E., Klein, R., Reading, K., Liptak, R., Millard, C., Lee, K. E., Tomany, S. C., Moore, E. L., Fijal, B. A., Elston, R. C. (2003) A whole-genome screen of a quantitative trait of age-related maculopathy in sibships from the Beaver Dam Eye Study. *Am. J. Hum. Genet.* 72, 1412-1424.
- Majewski, J., Schultz, D. W., Weleber, R. G., Schain, M. B., Edwards, A. O., Matise, T. C., Acott, T. S., Ott, J., Klein, M. L. (2003) Age-related macular degeneration--a genome scan in extended families. *Am. J. Hum. Genet.* 73, 540-550.
- Malek, G., Johnson, L. V., Mace, B. E., Saloupis, P., Schmechel, D. E., Rickman, D. W., Toth, C. A., Sullivan, P. M., Bowes Rickman, C. (2005) Apolipoprotein E allele-dependent pathogenesis: a model for age-related retinal degeneration. *Proc. Natl. Acad. Sci. U S A* 102, 11900-11905.

- Mullins, R. F., Russell, S. R., Anderson, D. H., Hageman, G. S. (2000) Drusen associated with aging and age-related macular degeneration contain proteins common to extracellular deposits associated with atherosclerosis, elastosis, amyloidosis, and dense deposit disease. *FASEB J* 14, 835-846.
- Nicolas, M. G., Fujiki, K., Murayama, K., Suzuki, M. T., Mineki, R., Hayakawa, M., Yoshikawa, Y., Cho, F., Kanai, A. (1996) Studies on the mechanism of early onset macular degeneration in cynomolgus (*Macaca fascicularis*) monkeys. I. Abnormal concentrations of two proteins in the retina. *Exp. Eye Res.* 62, 211-219.
- Nicolas, M. G., Fujiki, K., Murayama, K., Suzuki, M. T., Shindo, N., Hotta, Y., Iwata, F., Fujimura, T., Yoshikawa, Y., Cho, F., Kanai, (1996) A. Studies on the mechanism of early onset macular degeneration in cynomolgus monkeys. II. Suppression of metallothionein synthesis in the retina in oxidative stress. *Exp. Eye Res.* 62, 399-408.
- Nozaki, M., Raisler, B. J., Sakurai, E., Sarma, J. V., Barnum, S. R., Lambris, J. D., Chen, Y., Zhang, K., Ambati, B. K., Baffi, J. Z., Ambati J. (2006) Drusen complement components C3a and C5a promote choroidal neovascularization. *Proc. Natl. Acad. Sci. U S A* 103, 2328-2333.
- Okamoto, H., Umeda, S., Obazawa, M., Minami, M., Noda, T., Mizota, A., Honda, M., Tanaka, M., Koyama, R., Takagi, I., Sakamoto, Y., Saito, Y., Miyake, Y., Iwata, T. (2006) Complement factor H polymorphisms in Japanese population with age-related macular degeneration. *Mol. Vis.* 12, 156-158.
- Russell, S. R., Mullins, R. F., Schneider, B. L., Hageman, G. S. (2000) Location, substructure, and composition of basal laminar drusen compared with drusen associated with aging and age-related macular degeneration. *Am J Ophthalmol* 129, 205-214.
- Seddon, J. M., Ajani, U. A., Mitchell, B. D. (1997) Familial aggregation of age-related maculopathy. *Am. J. Ophthalmology* 123, 199-206.
- Stafford, T. J., Anness, S. H., Fine, B. S. (1984) Spontaneous degenerative maculopathy in the monkey. *Ophthalmology* 91, 513-521.
- Yang, Z., Camp, N. J., Sun, H., Tong, Z., Gibbs, D., Cameron, D. J., Chen, H., Zhao, Y., Pearson, E., Li, X., Chien, J., Dewan, A., Harmon, J., Bernstein, P. S., Shridhar, V., Zabriskie, N. A., Hoh, J., Howes, K., Zhang, K. (2006) A variant of the HTRA1 gene increases susceptibility to age-related macular degeneration. *Science*. 314, 992-993.
- Umeda, S., Ayyagari, R., Allikmets, R., Suzuki, M. T., Karoukis, A. J., Ambasadhan, R., Zernant, J., Okamoto, H., Ono, F., Terao, K., Mizota, A., Yoshikawa, Y., Tanaka, Y., Iwata, T. (2005) Early-onset macular degeneration with drusen in a cynomolgus monkey (*Macaca fascicularis*) pedigree: exclusion of 13 candidate genes and loci. *Invest. Ophthalmol. Vis. Sci.* 46, 683-691.
- Umeda, S., Suzuki, M. T., Okamoto, H., Ono, F., Mizota, A., Terao, K., Yoshikawa, Y., Tanaka, Y., Iwata, T. (2005) Molecular composition of drusen and possible involvement of anti-retinal autoimmunity in two different forms of macular degeneration in cynomolgus monkey (*Macaca fascicularis*). *FASEB J.* 19, 1683-1685.

Proteomic and Transcriptomic Analyses of Retinal Pigment Epithelial Cells Exposed to REF-1/TFPI-2

Masabiko Shibuya,^{1,2} Haru Okamoto,¹ Takehiro Nozawa,³ Jun Utsumi,⁴ Venkat N. Reddy,⁵ Hirotosbi Echizen,² Yasubiko Tanaka,⁶ and Takeshi Iwata¹

PURPOSE. The authors previously reported a growth-promoting factor, REF-1/TFPI-2, that is specific to retinal pigment epithelial (RPE) cells. The purpose of this study was to determine the genes and proteins of human RPE cells that are altered by exposure to TFPI-2.

METHODS. Human primary RPE cells were cultured with or without TFPI-2. Cell extracts and isolated RNA were subjected to proteomic and transcriptomic analyses, respectively. Proteins were separated by two-dimensional gel electrophoresis followed by gel staining and ion spray tandem mass spectrometry analyses. Transcriptomic analysis was performed using a DNA microarray to detect 27,868 gene expressions.

RESULTS. Proteomic analysis revealed c-Myc binding proteins and ribosomal proteins L11 preferentially induced by TFPI-2 in human RPE cells. Transcriptomic analysis detected 10,773 of 33,096 probes in the TFPI-2 treated samples, whereas only 2186 probes were detected in the nontreated samples. Among the genes up-regulated by TFPI-2 at the protein level were *c-myc*, *Mdm2*, transcription factor *E2F3*, retinoblastoma binding protein, and the *p21* gene, which is associated with the *c-myc* binding protein and ribosomal protein L11.

CONCLUSIONS. The mechanisms by which TFPI-2 promotes the proliferation of RPE cells may be associated with augmented *c-myc* synthesis and the activation of E2F in the retinoblastoma protein (Rb)/E2F pathway at the G1 phase of the RPE cells. Activation of ribosomal protein L11 and the Mdm2 complex of the p53 pathway may be counterbalanced by the hyperproliferative conditions. (*Invest Ophthalmol Vis Sci.* 2007;48:516-521) DOI:10.1167/iov.06-0434

Retinal pigment epithelial (RPE) cells play important roles in maintaining the homeostasis of the retina. RPE cells, located between the sensory retina and the choroidal blood supply, form a diffusion barrier controlling access to the subretinal space, with the RPE membrane regulating the transport

of proteins and controlling the hydration and ionic composition of the subretinal space. The sensitivity and viability of the photoreceptors thus depend on RPE-catalyzed transport activity. Proteins in the RPE cells that function in ionic, sugar, peptide, and water transport have been identified.¹ Damage to RPE cells generally leads to degeneration of the neural retina, as occurs in retinitis pigmentosa and age-related macular degeneration. Transplantation of the healthy retinal pigment cells or embryonic stem cells differentiating into RPE cells would be an ideal therapeutic approach to treat such diseases, and such attempts have been made.²

An alternative approach to treat these retinal diseases would be the use of a growth factor that promotes proliferation of the remaining RPE cells in a damaged retina or one that stimulates the regeneration of damaged RPE cells. To find such factor(s), the proteins expressed in human fibroblast cells were fractionated and assayed, leading to the isolation of RPE cell factor-1 (REF-1), which selectively promoted the proliferation of primary human RPE cells.³

Subsequently, the cDNA of REF-1 was cloned using information from the N-terminal amino acid sequences, which was identical with the tissue factor pathway inhibitor-2 (TFPI-2).³ Earlier studies have shown that TFPI-2 is a Kunitz-type serine protease inhibitor⁴⁻⁶ involved in the regulation of extrinsic blood coagulation⁷ and in the proliferation, invasion, and metastasis of various types of malignant cells.^{4,8-15} Extensive studies on the physiological roles of TEPI-2 have revealed that the ERK/MAPK pathway¹⁵ may be associated with the up-regulation of the *TFPI-2* gene and that DNA methylation^{9,10} in certain tumor cell lines may be related to the downregulation of the *TEPI-2* gene. When TFPI-2 is added to the culture medium of vascular smooth muscle cells, it promotes cell proliferation.¹⁴

Our initial finding that TFPI-2 enhanced RPE proliferation prompted us to question how this was achieved. We applied proteomic and transcriptomic analyses to screen the changes in the expression of the RNAs and proteins in RPE cells and will show that the proliferation promoting activity of TFPI-2 on RPE cells is associated with the regulation of an oncogene product, *c-myc*, and representative cancer repressor proteins retinoblastoma protein (Rb)/E2F and p53.

MATERIALS AND METHODS

TFPI-2 Treatment of Human RPE Cell Culture

Human primary RPE cells (passage 5) were seeded at a density of 2.5×10^4 cells/0.5 mL per well in 24-well plastic plates (BD Biosciences, Franklin Lakes, NJ) with Dulbecco modified MEM (DMEM; Invitrogen Japan, Tokyo, Japan) containing 15% fetal calf serum (FCS, Invitrogen). TFPI-2 was added to 20 wells with the RPE cells at 10 ng/mL concentrations and was incubated at 37°C for 24 hours for the proteomic samples, and for 6 hours, 12 hours, and 24 hours for the transcriptomic samples. An equal amount of saline was added to 20 wells containing RPE cells for controls. TFPI-2 was donated by Toray Industries, Inc., Tokyo, Japan.

From the ¹Laboratory of Cellular and Molecular Biology, National Institute of Sensory Organs, National Hospital Organization Tokyo Medical Center, Tokyo, Japan; ²Department of Pharmacotherapy, Meiji Pharmaceutical University, Tokyo, Japan; ³Analytical Instrument Division, AMR Inc., Tokyo, Japan; ⁴R&D Division, Toray Industries, Inc., Tokyo, Japan; ⁵Department of Ophthalmology, Kellogg Eye Center, University of Michigan, Ann Arbor, Michigan; and ⁶International University of Health and Welfare, Mita Hospital, Tokyo, Japan.

Supported in part by a grant-in-aid from the policy-based Medical Services Foundation.

Submitted for publication April 18, 2006; revised July 17, 2006; accepted December 4, 2006.

Disclosure: M. Shibuya, None; H. Okamoto, None; T. Nozawa, AMR Inc. (F); J. Utsumi, Toray Industries, Inc. (F); V.N. Reddy, None; H. Echizen, None; Y. Tanaka, None; T. Iwata, None

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

Corresponding author: Takeshi Iwata, Laboratory of Cellular and Molecular Biology, National Institute of Sensory Organs, National Hospital Organization Tokyo Medical Center, 2-5-1 Higashigaoka, Meguro-ku, Tokyo 152-8902, Japan; iwataakeshi@kankakuki.go.jp.

Protein Sample Preparation

To isolate whole cellular protein extracts from cultured RPE cells, the cells were rinsed 3 times with 1× PBS (pH 7.4) and were lysed in a denaturing lysis buffer containing 7 M urea, 2 M thiourea, 4% CHAPS, 40 mM Tris, 0.2% purifier (Bio-Lyte, pH range 3–10; Bio-Rad, Hercules, CA), and 50 mM dithiothreitol (DTT). The collected lysate was then centrifuged at 14,000g for 15 minutes at 4°C. Proteins in the supernatant were repeatedly concentrated and precipitated and finally desalted (ReadyPrep 2-D Cleanup kit; Bio-Rad). The protein concentration in the RPE samples was determined by a modified Lowry method adapted for use with the lysis buffer.

Two-Dimensional Electrophoresis

Protein samples were separated by a two-dimensional electrophoresis method. A 300- μ g protein sample was loaded on immobilized pH gradient (IPG) strips (pH 3–10, 7 cm; pH 4–7, 17 cm; Bio-Rad) by in-gel rehydration at 20°C overnight. For the 7-cm strip, isoelectric focusing (IEF) was used for the first dimension at an initial voltage of 250 V for 15 minutes, increased to 4000 V for 2 hours, and held until 20,000 V/h was reached. For the 17-cm strip, the initial voltage was set at 250 V, as for the 7-cm strip. Then the voltage was increased to 10,000 V for 3 hours and was held until 60,000 V/h was reached. Immediately after IEF, the IPG strips were equilibrated for 20 minutes in buffer containing 6 M urea, 2% SDS, 0.375 M Tris (pH 8.8), and 20% glycerol under a reduced condition with 2% DTT (Bio-Rad), followed by another incubation for 10 minutes in the same buffer under alkylating conditions with 2.5% iodoacetamide (Bio-Rad).¹⁵

Equilibrated IPG strips were then electrophoresed by SDS-PAGE for the second dimension. Images of the chemiluminescent signals were captured and merged with those of protein spots made visible by protein gel stain (Sypro Ruby; Bio-Rad), and the spots corresponding to the immunoreactivity were cut out. To test reproducibility, the experiment was performed twice.

Protein Identification by Mass Spectrometry

Excised gel pieces were rinsed with water and then with acetonitrile and were completely dried for the reduction-alkylation step. They were incubated with 10 mM DTT in 100 mM ammonium bicarbonate for 45 minutes at 56°C, then with 55 mM iodoacetamide in 100 mM ammonium bicarbonate for 30 minutes at room temperature in the dark. The supernatant was removed, and the washing procedure was repeated three times. Finally, the gel pieces were again completely dried before trypsin digestion and were rehydrated in a solution of trypsin (12.5 ng/ μ L; Promega, Madison, WI) in 50 mM ammonium bicarbonate. The digestion was continued for 16 hours at 37°C, and the extraction step was performed once with 25 mM ammonium bicarbonate, then twice with 5% formic acid, and finally with water. After resuspension in 40 μ L solution of aqueous 0.1% trifluoroacetic acid/2% acetonitrile, the samples were analyzed by liquid chromatography coupled to tandem mass spectrometry (LC-MS/MS). For analysis by LC-MS/MS, the tryptic digests were injected by an automatic sampler (HTS-PAL, CTC Analytics, Zwingen, Switzerland) onto a 0.2 × 50-mm capillary reversed-phase column (Magic C18, 3 μ m; Michrom BioResources, Inc., Auburn, CA) using an HPLC (Paradigm MS4; Michrom BioResources). Peptides were eluted with a gradient (95% solvent A consisting of 98% H₂O/2% acetonitrile/0.1% formic acid)/5% solvent B (10% H₂O/90% acetonitrile/0.1% formic acid) (0 minute)/35% solvent A/65% solvent B (20 minutes)/5% solvent A/95% solvent B (21 minutes)/5% solvent A/95% solvent B (23 minutes)/95% solvent A/5% solvent B (30 minutes) for 30 minutes at a flow rate of 1.5 μ L/min. Peptides were eluted directly into an ion trap mass spectrometer (ESI; Finnigan LTQ; Thermo Electron Corporation, Waltham, MA) capable of data-dependent acquisition. Each full MS scan was followed by an MS/MS scan of the most intense peak in the full MS spectrum with the dynamic exclusion enabled to allow detection of less-abundant peptide ions. Mass spectrometric scan events and HPLC solvent gradients were controlled with the use of a computer program (Paradigm Home; Michrom BioResources).

Total RNA Isolation from RPE Cells

Total RNA was isolated from the cultured RPE cells after 6 hours, 12 hours, and 24 hours with TFPI-2 using a total RNA isolation kit (RNA-Bee-RNA Isolation Reagent; Tel-Test, Friendswood, TX). Total RNA samples were treated with RNase-free DNase (Roche Diagnostics Japan) to minimize genomic DNA contamination.

DNA Microarray Analysis

DNA microarray analysis was performed (AB1700 Chemiluminescent Microarray Analyzer; Applied Biosystems, Foster City, CA). The survey array used (Human Genome Survey Array; Applied Biosystems) contained 33,096 60-mer oligonucleotide probes representing a set of 27,868 individual human genes and more than 1000 control probes. Sequences used for the microarray probe were obtained from curated transcripts (Celera Genomics Human Genome Database), RefSeq transcripts that had been structurally curated from the LocusLink public database, high-quality cDNA sequences from the Mammalian Gene Collection (MGC; <http://mgc.nci.nih.gov>), and transcripts that were experimentally validated (Applied Biosystems). The 60-mer oligo probes were synthesized using standard phosphoramidite chemistry and solid-phase synthesis and underwent quality control by mass spectrometry. The probes were deposited and covalently bound to a derivatized nylon substrate (2.5 × 3 inches) that was backed by a glass slide by contact spotting with a feature diameter of 180 μ m and more than 45 μ m between each feature. A 24-mer oligo internal control probe (ICP) was cospotted at every feature with 60-mer gene expression probe on the microarray. Digoxigenin-UTP labeled cRNA was generated and linearly amplified from 1 μ g total RNA (Chemiluminescent RT-IVT Labeling Kit, version 2.0; Applied Biosystems) according to the manufacturer's protocol. Array hybridization (two arrays per sample), chemiluminescence detection, image acquisition, and analysis were performed (Chemiluminescence Detection Kit and AB1700 Chemiluminescent Microarray Analyzer; Applied Biosystems) according to the manufacturer's protocol.

Briefly, each microarray was first prehybridized at 55°C for 1 hour in hybridization buffer with blocking reagent. Sixteen micrograms labeled cRNA targets were first fragmented into 100 to 400 bases by incubation with fragmentation buffer at 60°C for 30 minutes, mixed with internal control target (ICT; 24-mer oligo labeled with LIZR fluorescent dye), and hybridized to each prehybrid microarray in 1.5 mL vol at 55°C for 16 hours. After hybridization, the arrays were washed with hybridization wash buffer and chemiluminescence rinse buffer. Enhanced chemiluminescence signals were generated by first incubating the arrays with anti-digoxigenin alkaline phosphatase and enhanced with chemiluminescence enhancing solution and chemiluminescence substrate.

Images were collected from each microarray using the 1700 analyzer equipped with a high-resolution, large-format CCD camera, including 2 "short" chemiluminescent images (5-second exposure length each) and 2 "long" chemiluminescent images (25-second exposure length each) for gene expression analysis, two fluorescent images for feature finding and spot normalization, and two quality control images for spectrum cross-talk correction. Images were quantified, corrected for background and spot, and spatially normalized.

Data Analysis

MS data were identified with the use of a protein search program (BioWorks 3.2; Thermo Electron Corporation, Waltham, MA). For protein database searches, the same program was used to create centroid peak lists from the raw spectra. These peak lists were then submitted for database searching (BioWorks). The identity of the samples was searched from databases (nrNCBI (www.ncbi.nlm.nih.gov)) that extracted proteins and were restructured; search terms included human and *Homo sapiens*. Differentially expressed proteins were further analyzed for related genes and proteins using natural language processing software (PubGene database; PubGene Inc., Boston, MA) and data mining software of gene expression (OmniViz; OmniViz, Inc., Maynard, MA).




FIGURE 1. Two-dimensional gel electrophoresis of human RPE cells culture with (A) and without (B) TFPI-2. Spots corresponding to proteins whose expression is dependent on the presence of TFPI-2 in the culture medium are indicated by the arrows (*insets*). Proteins were detected by SYPRO Ruby staining. Spots corresponding to the differentially expressed proteins indicated by arrows (1 vs. 2 and 3 vs. 4) were subsequently subject to the LC-MS/MS analysis so that proteins could be identified.

RESULTS

Proteomic Analysis of RPE Cells Treated with TFPI-2

To determine the mechanisms responsible for the proliferation-promoting activity of TFPI-2 on RPE cells, protein synthesis and RNA expression were determined before and after TFPI-2 exposure. Differentially expressed proteins in the primary human RPE cells in response to TFPI-2 were identified by two-dimensional electrophoresis (Fig. 1). Samples were initially separated using IPG at a pH range of 3 to 10 to observe the full distribution of protein spots. The pH range was then narrowed to 4 to 7 to obtain higher resolution for spot picking. Consequently, approximately 480 spots were identified in the whole gel. We then focused on molecular weight less than 25 kDa, which is easy to check for changes. Ten spots considered differentially expressed in the two-dimensional gel were collected and subjected to LC-MS/MS analysis. Among the identified proteins, ribosomal protein L11 (RPL11; Fig. 1-1) and c-Myc binding protein (MYCBP; Fig. 1-3), known for regulating cell proliferation, were identified.¹⁶ These two proteins, identified by LC-MS/MS analysis and data analysis software (BioWorks 3.2), were consistent with those estimated from the results of two-dimensional electrophoresis (Table 1).

Transcriptomic Analysis of RPE Cells Treated with TFPI-2

The expression of 8134 genes in RPE cells was analyzed using DNA microarray with and without TFPI-2 exposure for 6 hours, 12 hours, and 24 hours. Signal normalization was performed for six independent DNA microarray chips according to the manufacturer's protocol. Genes differentially expressed by

more than threefold were considered significant and were selected for further analysis. Among the 33,096 possible probes, 10,773 probes were detected in the RPE cells incubated with TFPI-2, whereas only 2186 probes were detected without TFPI-2. Based on expression levels at the three time points (6 hours, 12 hours, and 24 hours), the time-dependent expression pattern of each gene was calculated and clustered with other genes with similar expression patterns using data mining software (OmniViz). Data analysis resulted in 38 clusters of genes that either increased or decreased their expression levels by more than twofold after TFPI-2 (Fig. 2). Nineteen genes were upregulated in 5 clusters, 108 genes in 16 clusters, and 717 genes in 22 clusters at 6 hours, 12 hours, and 24 hours, respectively. For downregulated genes, 30 genes in 16 clusters, 119 genes in 19 clusters, and 3 genes in 19 clusters were observed after 6 hours, 12 hours, and 24 hours, respectively. Transcriptomic analysis revealed significantly more genes differentially expressed at the transcriptional level than at the proteome level.

DISCUSSION

Proteins and genes whose expression was upregulated or downregulated after exposure to TFPI-2 were analyzed in human RPE cells to study the proteomic and transcriptomic changes. Protein and gene expression profiles for human RPE cells have been reported by West et al.,¹⁷ who identified 278 proteins, and Cai et al.,¹⁸ who reported 5580 ± 84 genes expressed in adult human RPE and ARPE19 cell lines using a DNA chip with 12,600 probes (Human U95Av2; Affymetrix, Santa Clara, CA). Our study showed changes in the expression of 8134 of 27,868 genes. DNA microarray analyses were simul-

TABLE 1. Two-Dimensional Gel Spots Identified by Mass Spectrometry

Protein	Number of AA	Peptide Residues	Identified Peptide from Database	MW	Score	Accession Number
c-Myc binding protein	167	108-117	TAEDAKDFFK	18642.6	10.13	1731809
Ribosomal protein L11	177	88-94	VREYELR	20125.1	20.21	14719845




FIGURE 2. Differentially expressed genes detected by DNA array are plotted as clusters. Differentially expressed genes whose expression level was increased by more than twofold (A-C) or was reduced by more than 0.5-fold (D-F) in RPE cells treated with TFPI-2 at incubation times of 6 hours, 12 hours, and 24 hours compared with the control cells are shown. Expression profile analysis revealed different gene expression patterns at each incubation time.

taneously performed at three time points (6 hours, 12 hours, and 24 hours) to monitor the course of expression of the possible 27,868 genes in human RPE cells exposed and not exposed to TFPI-2. This study was conducted at the translational and the transcriptional levels to complement the disadvantages of each method.

Raw gene expression data were further analyzed with data mining software (OmniViz) to obtain an overall picture of the transcriptional changes induced by TFPI-2 in human primary RPE cells. Genes whose expressions were changed by more than twofold were clustered into 38 groups showing a change of expression at each time point (Fig. 2). The number of genes upregulated at each time point was considerably higher than the number that was downregulated. A small number of genes was triggered by TFPI-2 treatment at 6 hours, before the major changes occurred at 24 hours. Among the initially upregulated genes were reticulon 4 interacting protein 1, phospholipase C, delta 1, granzyme M (lymphocyte met-ase 1; *GZMM*), and mitochondrial ribosomal protein L41 (*MRPL41*).

Proteomics analysis simultaneously performed at 24 hours identified two differentially expressed proteins, the *c-myc* binding protein (MYCBP) and the ribosomal protein L11 (RPL11). MYCBP and RPL11 (Fig. 3) are well known to regulate cell cycling through the Rb/E2F pathway and the p53 pathway, respectively. MYCBP stimulates *c-myc* transcription through the retinoblastoma protein (Rb)/E2F pathway (see Fig. 5). Sears et al.¹⁹ reported that activation of Myc increased the signal transduction of the cyclin D/cdk4 and cyclin E/cdk2 pathways. Activation of these pathways inactivates Rb after phosphorylation and E2F dissociation, which then promotes RPE cells to go into the S-phase of the cell cycle. The twofold transcriptional increase of *Rb* and *E2F3* in TFPI-2 exposed cells compared with control at 24 hours supports this hypothesis (Figs. 4C, 4F).

Concomitantly, the expressions of Rb and Mdm2 were upregulated twofold in growth-stimulated cells compared with control cells. Because Rb is associated with the negative regulation of the G₁-phase of the cell cycle, the enhanced expres-




FIGURE 3. Time course of gene expression for TFPI-2 (A), *c-myc* binding protein (B), and ribosomal protein L11 (C) in the cultured human RPE cells after exposure to TFPI-2.




FIGURE 4. Time courses of protein expression patterns for *c-myc* (A), *Mdm2* (B), retinoblastoma binding protein 4 (C), p-53 inducible protein (D), p21 (E), and transcription factor E2F3 in the cultured human RPE cells after exposure to TFPI-2.

sion of *Mdm2* might have been involved in the augmented degradation of Rb through the ubiquitin/proteasome-dependent pathway. Recently, Uchida et al.²⁰ suggested that *Mdm2* regulates the function of RB through the ubiquitin-dependent degradation of RB.

The *Rb* gene was the first identified tumor-suppressor gene,²¹ and it was recognized as a central component of a signaling pathway that controlled cell proliferation. Specifically, the D-type G_1 cyclins, together with their associated cyclin-dependent kinases (CKDs) Cdk4 and Cdk6, initiated the phosphorylation of Rb and Rb family members, inactivating their capacity to interact with the E2F transcription factors (Fig. 5).¹⁹ This phosphorylation leads to an accumulation of E2F1, E2F2, and E2F3a, which activate the transcription of a large number of genes essential for DNA replication and further cell cycle progression.^{22–26} Among the E2F targets are genes encoding a second class of G_1 cyclins, cyclin E, and the associated kinase Cdk2 (Fig. 5).¹⁹ The activation of cyclin

E/Cdk2 kinase activity by E2F leads to further phosphorylation and inactivation of Rb, further enhancing E2F activity and increasing the accumulation of cyclin E/Cdk2 (Fig. 5).¹⁹ This feedback loop, which leads to a continual inactivation of *Rb* independent of the action of cyclin D/Cdk4—defined as a junction in cell proliferation response when passed through the cell cycle—becomes growth factor independent.^{25,26} The activity of the G_1 Cdk is negatively regulated by a family of cyclin-dependent kinase inhibitors (CKIs), including p21^{WAF1}, p27^{KIP1}, and the p16^{INK4a} family.²⁷ The three upregulated E2Fs associate exclusively with Rb and appear to play a positive role in cell cycle progression.¹⁹

RPL11 binds the mouse double-minute 2 (*Mdm2* is the mouse homologue of *Hdm2* in humans) protein with other ribosomal proteins (L23 and L5) to form a complex to inhibit ubiquitin-dependent degradation of p53.^{28–30} The RPL11 protein is expressed in ARPE-19 cells.³¹ Inhibition of p53 degradation leads to p21 signaling, which participates in the G_1




FIGURE 5. Hypothetical network of various genes and proteins associated with the growth-promoting effect of TFPI-2 on the human RPE cells. Arrows: stimulatory signals. Straight and dotted lines: inhibitory effects.

arrest of the cell cycle but also negatively regulates cell proliferation (Fig. 5).^{30,32-34} In support of this hypothesis, *p21* transcription was increased by twofold after 24 hours by TFPI-2.

The *p53* gene mediates a major tumor-suppression pathway in mammalian cells and is frequently altered in human tumors.³⁰ Its function is kept at a low level during normal cell growth and is activated in response to various cellular stresses by acting as a sequence-specific transcription factor.³⁰ The *p53* protein induces cell cycle arrest or apoptosis.³⁰

Shinoda et al.¹⁴ reported cell growth proliferation of vascular smooth muscle endothelial cells by a purified mitogenic substance from human umbilical vein endothelial cells, later identified as TFPI-2. These authors showed the rapid activation of mitogen-activated protein kinase (MAPK) by TFPI-2 and the induced activation of proto-oncogene *c-fos* mRNA in smooth muscle cells.¹⁴ They concluded that *c-fos* activation was initiated by MAPK based on MAPK inhibitor PD098059 suppression.

In conclusion, the results of proteomic and transcriptomic analyses suggest that the proliferation of RPE cells induced by TFPI-2 is regulated through the Rb/E2F, *p53*, and Ras/Raf/MAPK pathways. We and others^{3,35} have reported a transcript of TFPI-2 in the mRNA of RPE cells. It is now reasonable to expect that RPE cells are able to self-proliferate by generating TFPI-2. Additional studies are needed to determine whether TFPI-2 can act as such an autocrine factor and can be modified for future treatment of the dry-type age-related macular degeneration and of retinitis pigmentosa.

References

- Hughes BA, Gallemore RP, Miller SS. Transport mechanisms in the retinal pigment epithelium. In: Marmor MF, Wolfensberger TJ, eds. *The Retinal Pigment Epithelium: Function and Disease*. New York: Oxford University Press; 1998:103-134.
- Haruta M. Embryonic stem cells: potential source for ocular repair. *Semin Ophthalmol*. 2005;20:17-23.
- Tanaka Y, Utsumi J, Matsui M, et al. Purification, molecular cloning, and expression of a novel growth-promoting factor for retinal pigment epithelial cells, REF-1/TFPI-2. *Invest Ophthalmol Vis Sci*. 2004;45:245-252.
- Chand HS, Foster DC, Kiesel W. Structure, function and biology of tissue factor pathway inhibitor-2. *Thromb Haemost*. 2005;94:1122-1130.
- Schmidt AE, Chand HS, Cascio D, et al. Crystal structure of Kunitz domain 1 (KD1) of tissue factor pathway inhibitor-2 in complex with trypsin: implications for KD1 specificity of inhibition. *J Biol Chem*. 2005;280:27832-27838.
- Chand HS, Schmidt AE, Bajaj SP, et al. Structure-function analysis of the reactive site in the first Kunitz-type domain of human tissue factor pathway inhibitor-2. *J Biol Chem*. 2004;279:17500-17507.
- Sprecher CA, Kiesel W, Mathewes S, et al. Molecular cloning, expression, and partial characterization of a second human tissue-factor-pathway inhibitor. *Proc Natl Acad Sci USA*. 1994;91:3353-3357.
- Yanamandra N, Kondraganti S, Gondi CS, et al. Recombinant adeno-associated virus (rAAV) expressing TFPI-2 inhibits invasion, angiogenesis and tumor growth in a human glioblastoma cell line. *Int J Cancer*. 2005;115:998-1005.
- Rollin J, Iochmann S, Blechet C, et al. Expression and methylation status of tissue factor pathway inhibitor-2 gene in non-small-cell lung cancer. *Br J Cancer*. 2005;92:775-783.
- Konduri SD, Srivenugopal KS, Yanamandra N. Promoter methylation and silencing of the tissue factor pathway inhibitor-2 (TFPI-2), a gene encoding an inhibitor of matrix metalloproteinases in human glioma cells. *Oncogene*. 2003;22:4509-4516.
- Santin AD, Zhan F, Bignotti E, et al. Gene expression profiles of primary HPV16- and HPV18-infected early stage cervical cancers and normal cervical epithelium: identification of novel candidate molecular markers for cervical cancer diagnosis and therapy. *Virology*. 2005;331:269-291.
- Sato N, Parker AR, Fukushima N, et al. Epigenetic inactivation of TFPI-2 as a common mechanism associated with growth and invasion of pancreatic ductal adenocarcinoma. *Oncogene*. 2005;24:850-858.
- Kast C, Wang M, Whiteway M. The ERK/MAPK pathway regulates the activity of the human tissue factor pathway inhibitor-2 promoter. *J Biol Chem*. 2003;278:6787-6794.
- Shinoda E, Yui Y, Hattori R, et al. Tissue factor pathway inhibitor-2 is a novel mitogen for vascular smooth muscle cells. *J Biol Chem*. 1999;274:5379-5384.
- Bahk SC, Lee SH, Jang JU, et al. Identification of crystallin family proteins in vitreous body in rat endotoxin-induced uveitis: involvement of crystallin truncation in uveitis pathogenesis. *Proteomics*. 2006;6:3436-3444.
- Taira T, Maeda J, Onishi T, et al. AMY-1, a novel C-MYC binding protein that stimulates transcription activity of C-MYC. *Genes Cells*. 1998;3:549-565.
- West KA, Yan L, Shadrach K, et al. Protein database, human retinal pigment epithelium. *Mol Cell Proteomics*. 2003;2:37-49.
- Cai H, Del Priore LV. Gene expression profile of cultured adult compared to immortalized human RPE. *Mol Vis*. 2006;12:1-14.
- Sears RC, Nevins JR. Signaling networks that link cell proliferation and cell fate. *J Biol Chem*. 2002;277:11617-11620.
- Uchida C, Miwa S, Kitagawa K, et al. Enhanced Mdm2 activity inhibits pRb function via ubiquitin-dependent degradation. *EMBO J*. 2005;24:160-169.
- Hanahan D, Weinberg RA. The hallmarks of cancer. *Cell*. 2000;100:57-70.
- Dyson N. The regulation of E2F by pRb-family proteins. *Genes Dev*. 1998;12:2245-2262.
- Nevins JR. Toward an understanding of the functional complexity of the E2F and retinoblastoma families. *Cell Growth Differ*. 1998;9:585-593.
- Harbour JW, Dean DC. Rb function in cell-cycle regulation and apoptosis. *Nat Cell Biol*. 2000;2:E65-E67.
- Dou QP, Levin AH, Zhao S, Pardee AB. Cyclin E and cyclin A as candidates for the restriction point protein. *Cancer Res*. 1993;53:1493-1497.
- Pardee AB. A restriction point for control of normal animal cell proliferation. *Proc Natl Acad Sci USA*. 1974;71:1286-1290.
- Sherr CJ, Roberts JM. CDK inhibitors: positive and negative regulators of G1-phase progression. *Genes Dev*. 1999;13:1501-1512.
- Krystof V, McNaie IW, Walkinshaw MD. Antiproliferative activity of olomoucine II, a novel 2,6,9-trisubstituted purine cyclin-dependent kinase inhibitor. *Cell Mol Life Sci*. 2005;62:1763-1771.
- Dai MS, Lu H. Inhibition of MDM2-mediated p53 ubiquitination and degradation by ribosomal protein L5. *J Biol Chem*. 2004;279:44475-44482.
- Zhang Y, Wolf GW, Bhat K, et al. Ribosomal protein L11 negatively regulates oncoprotein MDM2 and mediates a p53-dependent ribosomal-stress checkpoint pathway. *Mol Cell Biol*. 2003;23:8902-8912.
- Rao KC, Palamalai V, Dunlevy JR, et al. Peptidyl-Lys metalloendopeptidase-catalyzed ¹⁸O labeling for comparative proteomics: application to cytokine/lipopolysaccharide-treated human retinal pigment epithelium cell line. *Mol Cell Proteomics*. 2005;4:1550-1557.
- Chao C, Saito S, Kang J. p53 transcriptional activity is essential for p53-dependent apoptosis following DNA damage. *EMBO J*. 2000;19:4967-4975.
- Bai F, Matsui T, Ohtani-Fujita N, et al. Promoter activation and following induction of the p21/WAF1 gene by flavone is involved in G1 phase arrest in A549 lung adenocarcinoma cells. *FEBS Lett*. 1998;437:61-64.
- Nyunoya T, Powers LS, Yarovsky TO. Hyperoxia induces macrophage cell cycle arrest by adhesion-dependent induction of p21Cip1 and activation of the retinoblastoma protein. *J Biol Chem*. 2003;278:36099-36106.
- Ortego J, Escibano J, Coca-Prados M. Gene expression of protease and protease inhibitors in the human ciliary epithelium and ODM-2 cells. *Exp Eye Res*. 1997;65:289-299.

85. 緑内障の動物モデル (1)

—霊長類モデル, ラットモデル—

岩田 岳

独立行政法人国立病院機構東京医療センター
臨床研究センター(感覚器センター)
分子細胞生物学研究部門

緑内障研究において動物モデルの存在はきわめて重要である。現在はおもに霊長類に加えてラットやマウスなどの齧歯(げっし)類が利用されている。本セミナーでは緑内障で利用されている動物モデルについて、2回シリーズで紹介したい。

はじめに

動物モデルの利点は、隅角や視神経・網膜における変化を発症過程に沿って詳細に解析できること、さらに新薬の評価を行えることである。これまでも複数の哺乳類やその他の種で動物モデルの探索や開発が行われてきた¹⁾。利用目的に応じて1)ヒトとの視覚形態の類似性、2)発症までの時間、3)遺伝子操作の可能性、4)モデル動物作製に必要な技術、5)眼球の大きさ、6)解析に必要な技術、7)モデル動物の有効性、8)動物の維持費用などの検討が必要である。これまでも異なる種で自然発症した緑内障モデル動物が紹介されているが、一般的には手術的あるいは遺伝子改変によって作製されたモデル動物が利用されている。

●霊長類モデル

すべての動物モデルのなかで隅角や視神経乳頭の構造がヒトと最も類似する霊長類モデルが研究に適していることはいうまでもないことである。特に房水流出機構に関する研究においては貴重な存在である。しかし、1頭当たりの維持費用がマウスの約100倍かかることや、飼育・管理に高度な知識・技術が必要であることから、多くの研究では利用されていない。房水流路の遮断にはおもに線維柱帯の光凝固が利用される^{2,3)}。この手法によって手術後数日間で25~60 mmHgの眼圧上昇が期待できる。その他の手法としては前房内に赤血球⁴⁾、ラテックス⁵⁾、ポリアクリルアミドゲル⁶⁾、ステロイド⁷⁾を注入することによって眼圧上昇を促す方法が報告されているが、光凝固によって最も安定した眼圧上昇が得られている⁸⁾。霊長類における眼圧上昇は視神経乳頭、網膜神経線維、網膜神経節細胞層に障害⁹⁾をもたらす。ヒトと同様な病理学的所見が再現されることが確認されている。また、霊長類モデルを利用した、光凝固後30日に




図1 カニクイザルとマウスの視神経乳頭の比較
カニクイザル(A)の視神経乳頭の構造はヒトときわめて類似しており、マウス(B)のそれとは大きく異なる。

における網膜内の遺伝子発現の研究も報告されており¹⁰⁾、この情報は新しい治療薬の開発にも利用されている。

●ラットモデル

動物モデルを用いて薬効評価を行う場合、実験には多数の動物が必要になる。このような場合にラットは有効である。ラットは簡単に飼育でき、性質もおとなしく、眼球も手ごろな大きさであることから、市販の機器を使って麻酔なしで眼圧測定ができる¹¹⁾。ラットの眼球には緑内障に関係する部位がすべて存在する。ラットにおける眼圧上昇は強膜静脈への生理食塩水の注入¹²⁾、インドインクを使った線維柱帯の光凝固¹³⁾、線維柱帯の光凝