

- 23 Soriano SG, Cowan DB, Proctor MR, Scott RM. Levels of soluble adhesion molecules are elevated in the cerebrospinal fluid of children with moyamoya syndrome. *Neurosurgery* 2002; 50: 544–49.
- 24 Kim SK, Yoo JI, Cho BK, et al. Elevation of CRABP-I in the cerebrospinal fluid of patients with moyamoya disease. *Stroke* 2003; 34: 2835–41.
- 25 Nanba R, Kuroda S, Ishikawa T, Houkin K, Iwasaki Y. Increased expression of hepatocyte growth factor in cerebrospinal fluid and intracranial artery in moyamoya disease. *Stroke* 2004; 35: 2837–42.
- 26 Fukui M. Current state of study on moyamoya disease in Japan. *Surg Neurol* 1997; 47: 138–43.
- 27 Yamada H, Deguchi K, Tanigawara T, et al. The relationship between moyamoya disease and bacterial infection. *Clin Neurosurg* 1997; 99: S221–24.
- 28 Mineharu Y, Takenaka K, Yamakawa H, et al. Inheritance pattern of familial moyamoya disease: autosomal dominant mode and genomic imprinting. *J Neurol Neurosurg Psychiatry* 2006; 77: 1025–29.
- 29 Ikeda H, Sasaki T, Yoshimoto T, Fukui M, Arinami T. Mapping of a familial moyamoya disease gene to chromosome 3p24.2-p26. *Am J Hum Genet* 1999; 64: 533–37.
- 30 Inoue TK, Ikezaki K, Sasazuki T, Matsushima T, Fukui M. Linkage analysis of moyamoya disease on chromosome 6. *J Child Neurol* 2000; 15: 179–82.
- 31 Yamauchi T, Tada M, Houkin K, et al. Linkage of familial moyamoya disease (spontaneous occlusion of the circle of Willis) to chromosome 17q25. *Stroke* 2000; 31: 930–35.
- 32 Sakurai K, Horiochi Y, Ikeda H, et al. A novel susceptibility locus for moyamoya disease on chromosome 8q32. *J Hum Genet* 2004; 49: 278–81.
- 33 Nanba R, Tada M, Kuroda S, Houkin K, Iwasaki Y. Sequence analysis and bioinformatics analysis of chromosome 17q25 in familial moyamoya disease. *Childs Nerv Syst* 2005; 21: 62–68.
- 34 Karasawa J, Touho H, Ohnishi H, Miyamoto S, Kikuchi H. Cerebral revascularization using omental transplantation for childhood moyamoya disease. *J Neurosurg* 1993; 79: 192–96.
- 35 Miyamoto S, Kikuchi H, Karasawa J, Nagata I, Ihara I, Yamagata S. Study of the posterior circulation in moyamoya disease. Part 2: visual disturbances and surgical treatment. *J Neurosurg* 1986; 65: 454–60.
- 36 Zheng W, Wanibuchi M, Onda T, et al. A case of moyamoya disease presenting with chorea. *Childs Nerv Syst* 2006; 22: 274–78.
- 37 Bowen M, Marks MP, Steinberg GK. Neuropsychological recovery from childhood moyamoya disease. *Brain Dev* 1998; 20: 119–23.
- 38 Kuroda S, Houkin K, Ishikawa T, et al. Determinants of intellectual outcome after surgical revascularization in pediatric moyamoya disease: a multivariate analysis. *Childs Nerv Syst* 2004; 20: 302–08.
- 39 Lubman DL, Pantelis C, Desmond P, Proffitt TM, Velakoulis D. Moyamoya disease in a patient with schizophrenia. *J Int Neuropsychol Soc* 2003; 9: 806–10.
- 40 Nagata T, Harada D, Aoki K, et al. Effectiveness of carbamazepine for benzodiazepine-resistant impulsive aggression in a patient with frontal infarcts. *Psychiatry Clin Neurosci* 2007; 61: 695–97.
- 41 Kodama N, Aoki Y, Hiraga H, Wada T, Suzuki J. Electroencephalographic findings in children with moyamoya disease. *Arch Neurol* 1979; 36: 16–19.
- 42 Isobe M, Kuroda S, Kamiyama H, Abe H, Mitumori K. Cerebral blood flow reactivity to hyperventilation in children with spontaneous occlusion of the circle of Willis (moyamoya disease). *No Shinkei Geka* 1992; 20: 399–407 (in Japanese).
- 43 Kameyama M, Shirane R, Tsurumi Y, et al. Evaluation of cerebral blood flow and metabolism in childhood moyamoya disease: an investigation into "re-build-up" phenomenon on EEG by positron CT. *Childs Nerv Syst* 1986; 2: 130–33.
- 44 Kazumata K, Kuroda S, Houkin K, Abe H, Mitumori K. Regional cerebral hemodynamics during re-build-up phenomenon in childhood moyamoya disease. An analysis using ^{99m}Tc-HMPAO SPECT. *Childs Nerv Syst* 1996; 12: 161–65.
- 45 Kuroda S, Houkin K, Hoshi Y, Tamura M, Kazumata K, Abe H. Cerebral hypoxia after hyperventilation causes "re-build-up" phenomenon and TIA in childhood moyamoya disease. A near-infrared spectroscopy study. *Childs Nerv Syst* 1996; 12: 448–52.
- 46 Kuroda S, Kamiyama H, Isobe M, Houkin K, Abe H, Mitumori K. Cerebral hemodynamics and "re-build-up" phenomenon on electroencephalogram in children with moyamoya disease. *Childs Nerv Syst* 1995; 11: 214–19.
- 47 Qiao F, Kuroda S, Kamada K, Houkin K, Iwasaki Y. Source localization of the re-build up phenomenon in pediatric moyamoya disease-a dipole distribution analysis using MEG and SPECT. *Childs Nerv Syst* 2003; 19: 760–64.
- 48 Irikura K, Miyasaka Y, Kurata A, et al. A source of haemorrhage in adult patients with moyamoya disease: the significance of tributaries from the choroidal artery. *Acta Neurochir* 1996; 138: 1282–86.
- 49 Iwama T, Morimoto M, Hashimoto N, Goto Y, Todaka T, Sawada M. Mechanism of intracranial rebleeding in moyamoya disease. *Clin Neurol Neurosurg* 1997; 99: S187–90.
- 50 Kawaguchi S, Sakai T, Morimoto T, Kakizaki T, Kamada K. Characteristics of intracranial aneurysms associated with moyamoya disease. A review of 111 cases. *Acta Neurochir* 1996; 138: 1287–94.
- 51 Kuroda S, Houkin K, Kamiyama H, Abe H. Effects of surgical revascularization on peripheral artery aneurysms in moyamoya disease: report of three cases. *Neurosurgery* 2001; 49: 463–67.
- 52 Dietrichs E, Dahl A, Nyberg-Hansen R, Russell D, Rootwelt K, Vege T. Cerebral blood flow findings in moyamoya disease in adults. *Acta Neurol Scand* 1992; 85: 318–22.
- 53 Marushima A, Yanaka K, Matsuki T, Kojima H, Nose T. Subarachnoid hemorrhage not due to ruptured aneurysm in moyamoya disease. *J Clin Neurosci* 2006; 13: 146–49.
- 54 Osanai T, Kuroda S, Nakayama N, Yamauchi T, Houkin K, Iwasaki Y. Moyamoya disease presenting with subarachnoid hemorrhage localized over the frontal cortex: case report. *Surg Neurol* 2008; 69: 197–200.
- 55 Komiyama M. Moyamoya disease and pregnancy. *J Nucl Med* 1999; 40: 214–15.
- 56 Seol HJ, Wang KC, Kim SK, Hwang YS, Kim KJ, Cho BK. Headache in pediatric moyamoya disease: review of 204 consecutive cases. *J Neurosurg* 2005; 103: 439–42.
- 57 Kikuta K, Takagi Y, Arakawa Y, Miyamoto S, Hashimoto N. Absence epilepsy associated with moyamoya disease. Case report. *J Neurosurg* 2006; 104: 265–68.
- 58 Kuroda S, Ishikawa T, Houkin K, Iwasaki Y. Clinical significance of posterior cerebral artery stenosis/occlusion in moyamoya disease. *No Shinkei Geka* 2002; 30: 1295–300 (in Japanese).
- 59 Yamada I, Murata Y, Umemura I, Suzuki S, Matsushima Y. SPECT and MRI evaluations of the posterior circulation in moyamoya disease. *J Nucl Med* 1996; 37: 1613–17.
- 60 Houkin K, Kuroda S, Ishikawa T, Abe H. Neovascularization (angiogenesis) after revascularization in moyamoya disease. Which technique is most useful for moyamoya disease? *Acta Neurochir* 2000; 142: 269–76.
- 61 Houkin K, Nakayama N, Kuroda S, Ishikawa T, Nonaka T. How does angiogenesis develop in pediatric moyamoya disease after surgery? A prospective study with MR angiography. *Childs Nerv Syst* 2004; 20: 734–41.
- 62 Yamada I, Suzuki S, Matsushima Y. Moyamoya disease: diagnostic accuracy of MRI. *Neuroradiology* 1995; 37: 356–61.
- 63 Harada A, Fujii Y, Yoneoka Y, Takeuchi S, Tanaka R, Nakada T. High-field magnetic resonance imaging in patients with moyamoya disease. *J Neurosurg* 2001; 94: 233–37.
- 64 Ishikawa T, Kuroda S, Nakayama S, Terae S, Kudou K, Iwasaki Y. Prevalence of asymptomatic microbleeds in patients with moyamoya disease. *Neuroradiol* 2005; 45: 495–500.
- 65 Kikuta K, Takagi Y, Nozaki K, et al. Asymptomatic microbleeds in moyamoya disease: T2*-weighted gradient-echo magnetic resonance imaging study. *J Neurosurg* 2005; 102: 470–75.
- 66 Kikuta K, Takagi Y, Nozaki K, Okada T, Hashimoto N. Histological analysis of microbleed after surgical resection in a patient with moyamoya disease. *Neuroradiol* 2007; 47: 564–67.
- 67 Kikuta K, Takagi Y, Nozaki K, Sawamoto N, Fukuyama H, Hashimoto N. The presence of multiple microbleeds as a predictor of subsequent cerebral hemorrhage in patients with moyamoya disease. *Neurosurgery* 2008; 62: 104–11.
- 68 Houkin K, Tanaka N, Takahashi A, Kamiyama H, Abe H, Kajii N. Familial occurrence of moyamoya disease. Magnetic resonance angiography as a screening test for high-risk subjects. *Childs Nerv Syst* 1994; 10: 421–25.
- 69 Houkin K, Aoki T, Takahashi A, Abe H. Diagnosis of moyamoya disease with magnetic resonance angiography. *Stroke* 1994; 25: 2159–64.
- 70 Hasuo K, Miura F, Matsushima T. MRI and MR angiography in moyamoya disease. *J Magn Reson Imaging* 1998; 8: 762–66.

- 71 Yamada I, Suzuki S, Matsushima Y. Moyamoya disease: comparison of assessment with MR angiography and MR imaging versus conventional angiography. *Radiology* 1995; 196: 211–18.
- 72 Houkin K, Nakayama N, Kuroda S, Nonaka T, Shonai T, Yoshimoto T. Novel magnetic resonance angiography stage grading for moyamoya disease. *Cerebrovasc Dis* 2005; 20: 347–54.
- 73 Kashiwagi S, Yamashita T, Katoh S, et al. Regression of moyamoya vessels and hemodynamic changes after successful revascularization in childhood moyamoya disease. *Acta Neurol Scand Suppl* 1996; 166: 85–88.
- 74 Ogawa A, Yoshimoto T, Suzuki J, Sakurai Y. Cerebral blood flow in moyamoya disease. Part 1: correlation with age and regional distribution. *Acta Neurochir* 1990; 105: 30–34.
- 75 Kuroda S, Kamiyama H, Abe H, et al. Cerebral blood flow in children with spontaneous occlusion of the circle of Willis (moyamoya disease): comparison with healthy children and evaluation of annual changes. *Neurology Med Chir* 1993; 33: 434–38.
- 76 Kuroda S, Houkin K, Kamiyama H, Abe H, Mitsumori K. Regional cerebral hemodynamics in childhood moyamoya disease. *Childs Nerv Syst* 1995; 11: 584–90.
- 77 Ogawa A, Nakamura N, Yoshimoto T, Suzuki J. Cerebral blood flow in moyamoya disease. Part 2: Autoregulation and CO₂ response. *Acta Neurochir* 1990; 105: 107–11.
- 78 Matheja P, Weckesser M, Debus O, et al. Moyamoya syndrome: impaired hemodynamics on ECD SPECT after EEG controlled hyperventilation. *Nuklearmedizin* 2002; 41: 42–46.
- 79 Kuwabara Y, Ichiba Y, Sasaki M, et al. Cerebral hemodynamics and metabolism in moyamoya disease—a positron emission tomography study. *Clin Neurosurg* 1997; 99: S74–78.
- 80 Ikezaki K, Matsushima T, Kuwabara Y, Suzuki SO, Nomura T, Fukui M. Cerebral circulation and oxygen metabolism in childhood moyamoya disease: a perioperative positron emission tomography study. *J Neurosurg* 1994; 81: 843–50.
- 81 Kuwabara Y, Ichiba Y, Otsuka M, et al. Cerebral hemodynamic change in the child and the adult with moyamoya disease. *Stroke* 1990; 21: 272–77.
- 82 Taki W, Yonekawa Y, Kobayashi A, et al. Cerebral circulation and oxygen metabolism in moyamoya disease of ischemic type in children. *Childs Nerv Syst* 1988; 4: 259–62.
- 83 Piao R, Oku N, Kitagawa K, et al. Cerebral hemodynamics and metabolism in adult moyamoya disease: comparison of angiographic collateral circulation. *Ann Nucl Med* 2004; 18: 115–21.
- 84 Iwama T, Akiyama Y, Morimoto M, Koijima A, Hayashida K. Comparison of positron emission tomography study results of cerebral hemodynamics in patients with bleeding- and ischemic-type moyamoya disease. *Neurosurg Focus* 1998; 5: e3.
- 85 Narita T, Matsushima Y, Imai S, et al. Severe haemodynamic stress in selected subtypes of patients with moyamoya disease: a positron emission tomography study. *J Neurol Neurosurg Psychiatry* 2005; 76: 663–69.
- 86 Mesihala AH, Sviri G, Fatemi N, Britz GW, Newell DW. Long-term outcome of superficial temporal artery-middle cerebral artery bypass for patients with moyamoya disease in the US. *Neurosurg Focus* 2008; 24: E15.
- 87 Touho H, Karasawa J, Ohnishi H. Preoperative and postoperative evaluation of cerebral perfusion and vasodilatory capacity with 99mTc-HMPAO SPECT and acetazolamide in childhood moyamoya disease. *Stroke* 1996; 27: 282–89.
- 88 Iwama T, Hashimoto N, Miyake H, Yonekawa Y. Direct revascularization to the anterior cerebral artery territory in patients with moyamoya disease: report of five cases. *Neurosurgery* 1998; 42: 1157–61.
- 89 Kohno K, Oka Y, Kohno S, Ohta S, Kumon Y, Sakaki S. Cerebral blood flow measurement as an indicator for an indirect revascularization procedure for adult patients with moyamoya disease. *Neurosurgery* 1998; 42: 752–57.
- 90 Mikulis DJ, Krolczyk G, Desai H, et al. Preoperative and postoperative mapping of cerebrovascular reactivity in moyamoya disease by using blood oxygen level-dependent magnetic resonance imaging. *J Neurosurg* 2005; 103: 347–55.
- 91 Oya S, Tsutsumi K, Ueki K. Adult-onset moyamoya disease with repetitive ischemic attacks successfully treated by superficial temporal-middle cerebral artery bypass—case report. *Neurology Med Chir* 2003; 43: 138–41.
- 92 Sato H, Sato N, Tamaki N, Matsumoto S. Chronic low-perfusion state in children with moyamoya disease following revascularization. *Childs Nerv Syst* 1990; 6: 166–71.
- 93 So Y, Lee HY, Kim SK, et al. Prediction of the clinical outcome of pediatric moyamoya disease with postoperative basal/acetazolamide stress brain perfusion SPECT after revascularization surgery. *Stroke* 2005; 36: 1485–89.
- 94 Togao O, Miura F, Yoshiura T, et al. Cerebral hemodynamics in moyamoya disease: correlation between perfusion-weighted MR imaging and cerebral angiography. *AJR Am J Neuroradiol* 2006; 27: 391–97.
- 95 Tanaka Y, Narita T, Nagaoka T, et al. Quantitative evaluation of cerebral hemodynamics in patients with moyamoya disease by dynamic susceptibility contrast magnetic resonance imaging—comparison with positron emission tomography. *J Cere Blood Flow Metab* 2006; 26: 291–300.
- 96 Karasawa J, Kikuchi H, Furuse S, Kawamura J, Sakaki T. Treatment of moyamoya disease with STA-MCA anastomosis. *J Neurosurg* 1978; 49: 679–88.
- 97 Ishikawa T, Kamiyama H, Kuroda S, Yasuda H, Nakayama N, Takizawa K. Simultaneous superficial temporal artery to middle cerebral or anterior cerebral artery bypass with pan-synangiosis for moyamoya disease covering both anterior and middle cerebral artery territories. *Neurology Med Chir* 2006; 46: 462–68.
- 98 Iwama T, Hashimoto N, Tsukahara T, Miyake H. Superficial temporal artery to anterior cerebral artery direct anastomosis in patients with moyamoya disease. *Clin Neurosurg* 1997; 99: S134–36.
- 99 Golby AJ, Marks MP, Thompson RC, Steinberg GK. Direct and combined revascularization in pediatric moyamoya disease. *Neurosurgery* 1999; 45: 50–58.
- 100 Veeravagu A, Guzman R, Patil CG, Hou LC, Lee M, Steinberg GK. Moyamoya disease in pediatric patients: outcomes of neurosurgical interventions. *Neurosurg Focus* 2008; 24: E16.
- 101 Fujimura M, Kaneta T, Mugikura S, Shimizu H, Tominaga T. Temporary neurologic deterioration due to cerebral hyperperfusion after superficial temporal artery-middle cerebral artery anastomosis in patients with adult-onset moyamoya disease. *Surg Neurol* 2007; 67: 273–82.
- 102 Kuroda S, Kamiyama H, Abe H, Asaoka K, Mitsumori K. Temporary neurological deterioration caused by hyperperfusion after extracranial-intracranial bypass case report and study of cerebral hemodynamics. *Neurology Med Chir* 1994; 34: 15–19.
- 103 Nakayama N, Kuroda S, Houkin K, Takikawa S, Abe H. Intraoperative measurement of arterial blood flow using a transit time flowmeter: monitoring of hemodynamic changes during cerebrovascular surgery. *Acta Neurochir* 2001; 143: 17–24.
- 104 Fujimura M, Kaneta T, Tominaga T. Efficacy of superficial temporal artery-middle cerebral artery anastomosis with routine postoperative cerebral blood flow measurement during the acute stage in childhood moyamoya disease. *Childs Nerv Syst* 2008; 24: 827–32.
- 105 Matsushima Y, Fukui N, Tanaka K, et al. A new surgical treatment of moyamoya disease in children: a preliminary report. *Surg Neurol* 1981; 15: 313–20.
- 106 Karasawa J, Kikuchi H, Furuse S, Sakaki T, Yoshida Y. A surgical treatment of "moyamoya" disease "encephalo-myo synangiosis". *Neurology Med Chir* 1977; 17: 29–37.
- 107 Kinugasa K, Mandai S, Kamata I, Sugiu K, Ohmoto T. Surgical treatment of moyamoya disease: operative technique for encephalo-duo-arterio-myo-synangiosis, its follow-up, clinical results, and angiograms. *Neurosurgery* 1993; 32: 527–31.
- 108 Ishii R. Surgical treatment of moyamoya disease. *No Shinkei Geka* 1986; 14: 1059–68 (in Japanese).
- 109 Kawaguchi T, Fujita S, Hosoda K, et al. Multiple burr-hole operation for adult moyamoya disease. *J Neurosurg* 1996; 84: 468–76.
- 110 Ozgur BM, Arya HE, Levy ML. Indirect revascularisation for paediatric moyamoya disease: the EDAMS technique. *J Clin Neurosci* 2006; 13: 105–08.
- 111 Ishikawa T, Houkin K, Kamiyama H, Abe H. Effects of surgical revascularization on outcome of patients with pediatric moyamoya disease. *Stroke* 1997; 28: 1170–73.
- 112 Sakamoto T, Kawaguchi M, Kurehara K, Kitaguchi K, Furuya H, Karasawa J. Risk factors for neurologic deterioration after revascularization surgery in patients with moyamoya disease. *Anesth Analg* 1997; 85: 1060–65.

- 113 Kim SH, Choi JU, Yang KH, Kim TG, Kim DS. Risk factors for postoperative ischemic complications in patients with moyamoya disease. *J Neurosurg* 2005; **103**: 433–38.
- 114 Iwama T, Hashimoto N, Tsukahara T, Murai B. Peri-operative complications in adult moyamoya disease. *Acta Neurochir* 1995; **132**: 26–31.
- 115 Iwama T, Hashimoto N, Yonekawa Y. The relevance of hemodynamic factors to perioperative ischemic complications in childhood moyamoya disease. *Neurosurgery* 1996; **38**: 1120–25.
- 116 Matsushima T, Inoue T, Suzuki SO, Fujii K, Fukui M, Hasuo K. Surgical treatment of moyamoya disease in pediatric patients—comparison between the results of indirect and direct revascularization procedures. *Neurosurgery* 1992; **31**: 401–05.
- 117 Takahashi A, Kamiyama H, Houkin K, Abe H. Surgical treatment of childhood moyamoya disease—comparison of reconstructive surgery centered on the frontal region and the parietal region. *Neur Med Chir* 1995; **35**: 231–37.
- 118 Matsushima T, Inoue T, Matsuta T, et al. An indirect revascularization method in the surgical treatment of moyamoya disease—various kinds of indirect procedures and a multiple combined indirect procedure. *Neur Med Chir* 1998; **38**: 297–302.
- 119 Mizoi K, Kayama T, Yoshimoto T, Nagamini Y. Indirect revascularization for moyamoya disease: is there a beneficial effect for adult patients? *Surg Neurol* 1996; **45**: 541–48.
- 120 Houkin K, Kamiyama H, Abe H, Takahashi A, Kuroda S. Surgical therapy for adult moyamoya disease. Can surgical revascularization prevent the recurrence of intracerebral hemorrhage? *Stroke* 1996; **27**: 1342–46.
- 121 Houkin K, Ishikawa T, Yoshimoto T, Abe H. Direct and indirect revascularization for moyamoya disease: surgical techniques and peri-operative complications. *Clin Neurol Neurosurg* 1997; **99**: S142–45.
- 122 Kurokawa T, Tomita S, Ueda K, et al. Prognosis of occlusive disease of the circle of Willis (moyamoya disease) in children. *Pediatr Neurol* 1985; **1**: 274–77.
- 123 Imaizumi T, Hayashi K, Saito K, Osawa M, Fukuyama Y. Long-term outcomes of pediatric moyamoya disease monitored to adulthood. *Pediatr Neurol* 1998; **18**: 321–25.
- 124 Imaizumi C, Imaizumi T, Osawa M, Fukuyama Y, Takeshita M. Serial intelligence test scores in pediatric moyamoya disease. *Neuropediatrics* 1999; **30**: 294–99.
- 125 Karasawa J, Touho H, Ohnishi H, Miyamoto S, Kikuchi H. Long-term follow-up study after extracranial-intracranial bypass surgery for anterior circulation ischemia in childhood moyamoya disease. *J Neurosurg* 1992; **77**: 84–89.
- 126 Kuroda S, Nanba R, Ishikawa T, Houkin K, Kamiyama H, Iwasaki Y. Clinical manifestations of infantile moyamoya disease. *No Shinkei Geka* 2003; **31**: 1073–78 (in Japanese).
- 127 Kim SK, Seol HJ, Cho BK, Hwang YS, Lee DS, Wang KC. Moyamoya disease among young patients: its aggressive clinical course and the role of active surgical treatment. *Neurosurgery* 2004; **54**: 840–44.
- 128 Ezura M, Yoshimoto T, Fujiwara S, Takahashi A, Shirane R, Mizoi K. Clinical and angiographic follow-up of childhood-onset moyamoya disease. *Childs Nerv Syst* 1995; **11**: 591–94.
- 129 Hallermeier CL, Rich KM, Grubb RL Jr, et al. Clinical features and outcome in north American adults with moyamoya phenomenon. *Stroke* 2006; **37**: 1490–96.
- 130 Han DH, Nam DH, Oh CW. Moyamoya disease in adults: characteristics of clinical presentation and outcome after encephaloduro-arterio-synangiosis. *Clin Neurol Neurosurg* 1997; **99**: S151–55.
- 131 Choi JU, Kim DS, Kim EY, Lee KC. Natural history of moyamoya disease: comparison of activity of daily living in surgery and non surgery groups. *Clin Neurol Neurosurg* 1997; **99**: S11–18.
- 132 Kobayashi E, Saeki N, Oishi H, Hirai S, Yamaura A. Long-term natural history of hemorrhagic moyamoya disease in 42 patients. *J Neurosurg* 2000; **93**: 976–80.
- 133 Yoshida Y, Yoshimoto T, Shirane R, Sakurai Y. Clinical course, surgical management, and long-term outcome of moyamoya patients with rebleeding after an episode of intracerebral hemorrhage: an extensive follow-up study. *Stroke* 1999; **30**: 2272–76.
- 134 Fujii K, Ikezaki K, Irikura K, Miyasaka Y, Fukui M. The efficacy of bypass surgery for the patients with hemorrhagic moyamoya disease. *Clin Neurol Neurosurg* 1997; **99**: S194–95.
- 135 Morioka M, Hamada J, Todaka T, Yano S, Kai Y, Ushio Y. High-risk age for rebleeding in patients with hemorrhagic moyamoya disease: long-term follow-up study. *Neurosurgery* 2003; **52**: 1049–54.
- 136 Iwama T, Kotani Y, Yamakawa H, Nagata I, Hashimoto N, Sakai N. Cerebral ischemic complications following intracranial bleeding in patients with moyamoya disease—three case reports. *Neur Med Chir* 2001; **41**: 450–53.
- 137 Hirotsune N, Meguro T, Kawada S, Nakashima H, Ohmoto T. Long-term follow-up study of patients with unilateral moyamoya disease. *Clin Neurol Neurosurg* 1999; **99**: S178–81.
- 138 Kawano T, Fukui M, Hashimoto N, Yonekawa Y. Follow-up study of patients with “unilateral” moyamoya disease. *Neur Med Chir* 1994; **34**: 744–47.
- 139 Kelly ME, Bell-Stephens TE, Marks MP, Do HM, Steinberg GK. Progression of unilateral moyamoya disease: a clinical series. *Cerebrovas Dis* 2006; **22**: 109–15.
- 140 Smith ER, Scott RM. Progression of disease in unilateral moyamoya syndrome. *Neurosurg Focus* 2008; **24**: E17.
- 141 Kuroda S, Ishikawa T, Houkin K, Nanba R, Hokari M, Iwasaki Y. Incidence and clinical features of disease progression in adult moyamoya disease. *Stroke* 2005; **36**: 2148–53.
- 142 Nanba R, Kuroda S, Takeda M, et al. [Clinical features and outcomes of 10 asymptomatic adult patients with moyamoya disease]. *No Shinkei Geka* 2003; **31**: 1291–95.
- 143 Yamada M, Fujii K, Fukui M. Clinical features and outcomes in patients with asymptomatic moyamoya disease—from the results of nation-wide questionnaire survey. *No Shinkei Geka* 2005; **33**: 337–42 (in Japanese).
- 144 Miyamoto S, Akiyama M, Nagata I, et al. Long-term outcome after STA-MCA anastomosis for moyamoya disease. *Neurosurg Focus* 1998; **5**: 1–4.
- 145 Chiu D, Sheddor P, Bratina P, Grotta JC. Clinical features of moyamoya disease in the United States. *Stroke* 1998; **29**: 1347–51.
- 146 Scott RM, Smith JL, Robertson RL, Madson JR, Soriano SG, Rockoff MA. Long-term outcome in children with moyamoya syndrome after cranial revascularization by pial synangiosis. *J Neurosurg* 2004; **100**: 142–49.
- 147 Sainte-Rose C, Oliveira R, Puget S, et al. Multiple bur hole surgery for the treatment of moyamoya disease in children. *J Neurosurg* 2006; **105**: 437–43.
- 148 Kim CY, Wang KC, Kim SK, Chung YN, Kim HS, Cho BK. Encephaloduroarteriosynangiosis with bifrontal encephalogaleo(periosteal)synangiosis in the pediatric moyamoya disease: is the surgical technique and its outcomes. *Childs Nerv Syst* 2003; **19**: 316–24.
- 149 Kim SK, Wang KC, Kim IO, Lee DS, Cho BK. Combined encephaloduroarteriosynangiosis and bifrontal encephalogaleo(periosteal) synangiosis in pediatric moyamoya disease. *Neurosurgery* 2002; **50**: 88–96.
- 150 Matsushima Y, Aoyagi M, Masao H, Suzuki R, Ohno K. Mental outcome following encephaloduroarteriosynangiosis in children with moyamoya disease with the onset earlier than 5 years of age. *Childs Nerv Syst* 1990; **6**: 440–43.
- 151 Fukuyama Y, Umezawa R. Clinical and cerebral angiographic evolutions of idiopathic progressive occlusive disease of the circle of Willis (“moyamoya” disease) in children. *Brain Dev* 1985; **7**: 21–37.
- 152 Kim DS, Yoo DS, Huh PW, Kang SG, Cho KS, Kim MC. Combined direct anastomosis and encephaloduroarteriogaleosynangiosis using inverted superficial temporal artery-galeal flap and superficial temporal artery-galeal pedicle in adult moyamoya disease. *Surg Neurol* 2006; **66**: 389–94.
- 153 Okada Y, Shima T, Nishida M, Yamane K, Yamada T, Yamanaka C. Effectiveness of superficial temporal artery-middle cerebral artery anastomosis in adult moyamoya disease: cerebral hemodynamics and clinical course in ischemic and hemorrhagic varieties. *Stroke* 1998; **29**: 625–30.
- 154 Wanifuchi H, Takeshita M, Izawa M, Aoki N, Kagawa M. Management of adult moyamoya disease. *Neur Med Chir* 1993; **33**: 300–05.
- 155 Kawaguchi S, Okuno S, Sakaki T. Effect of direct arterial bypass on the prevention of future stroke in patients with the hemorrhagic variety of moyamoya disease. *J Neurosurg* 2000; **93**: 397–401.
- 156 Miyamoto S. Study design for a prospective randomized trial of extracranial-intracranial bypass surgery for adults with moyamoya disease and hemorrhagic onset—the Japan adult moyamoya trial group. *Neur Med Chir* 2004; **44**: 218–19.